

Stable Homotopy Theory and Higher Algebra

An ∞ -categorical introduction

In progress; content and numbering unstable until January 2027

Bastiaan Cnossen

Regensburg, August 14, 2026

Contents

Change log	vii
Introduction	1
Chapter 1: Introduction to ∞ -categories	10
1.1 The heuristics behind ∞ -categories	11
1.2 The language of ∞ -category theory	15
1.3 Basic constructions of ∞ -categories	19
1.4 Commutative diagrams, composition, and animaes	27
1.5 Further constructions of ∞ -categories	35
1.6 Simplicial objects and quasicategories	40
1.7 Limits and colimits	44
1.8 Foundational results in ∞ -category theory	48
Exercises	53
 I Stable homotopy theory	 59
Chapter 2: Animaes as spaces	61
2.1 Homotopy pushouts and pullbacks	63
2.2 Computing pushouts and pullbacks in localizations	66
2.3 Relative cell complexes and Serre fibrations	70
2.4 Pointed animaes, suspensions and loops	77
Exercises	84
Chapter 3: Homology and cohomology	88
3.1 Homology and cohomology theories	88
3.2 Cohomology theories and spectra	90
3.3 Ordinary homology and cohomology	92
3.4 Brown representability (online only)	99
Exercises	104
Chapter 4: Spectra and stable ∞ -categories	105
4.1 Pointed ∞ -categories	105
4.2 Stable ∞ -categories	108

4.3	Stabilization	118
4.4	The ∞ -category of spectra	129
	Exercises	139
Chapter 5: The recognition principle		141
5.1	Monoids and groups	142
5.2	The recognition principle for loop spaces	147
5.3	Commutative monoids and commutative groups	153
5.4	The recognition principle for connective spectra	165
5.5	Group completions as telescopes	168
5.6	Finite deloopings (online only)	169
	Exercises	171
Chapter 6: Homological algebra		174
6.1	Derived ∞ -categories	174
6.2	Ordinary homology and cohomology via $D(\mathbb{Z})$	183
6.3	Connective objects and t-structures	189
6.4	Projective objects and Ext-groups	199
6.5	Flat objects and Tor-groups	205
6.6	Derived functors	209
6.7	Künneth and universal coefficient theorems	220
	Exercises	225
Chapter 7: Localization and completion		228
7.1	Bousfield localizations	228
7.2	Localization at a prime	231
7.3	Completion at a prime	235
	Exercises	241
Chapter 8: Ring spectra and modules		242
8.1	Algebras and modules	243
8.2	Connective modules	247
8.3	Eilenberg–MacLane ring spectra and DGAs	251
8.4	Localizations of ring spectra	254
8.5	Projective and flat modules (online only)	264
	Exercises	271
Chapter 9: Complex K-theory		272
9.1	Vector bundles	272
9.2	Classifying vector bundles	275
9.3	Complex K-theory groups	283
9.4	Complex K-theory spectra	286
	Exercises	293
Chapter 10: Thom spectra and bordism theory		295
10.1	Thom spaces and the collapse construction	296
10.2	Thom spectra	299

10.3	Bordism and the Pontryagin–Thom theorem	307
10.4	Orientations and the Thom isomorphism	310
10.5	Coherence for one-point compactification (online only)	312
	Exercises	317
Chapter 11:	Duality phenomena	318
11.1	Symmetric monoidal duality	318
11.2	Dualizability and compactness	322
11.3	Spanier–Whitehead and Atiyah duality	323
11.4	Poincaré duality	330
	Exercises	332
II	Operads and higher algebra	333
Chapter 12:	Operads: the heuristics	335
12.1	Non-colored operads	335
12.2	Colored operads	338
12.3	Encoding the coherences: the envelope of an operad	342
12.4	Writing concatenations as products	346
	Exercises	351
Chapter 13:	Span categories	353
13.1	Constructing span categories	353
13.2	Adjunctions of span categories	362
13.3	Products and coproducts in span categories	364
	Exercises	366
Chapter 14:	Basics on ∞ -operads	368
14.1	Definitions and examples	368
14.2	Symmetric monoidal ∞ -categories and algebras over operads	378
14.3	Lax symmetric monoidal functors	382
14.4	O -monoidal ∞ -categories	389
14.5	Monoidal subcategories and monoidal Bousfield localizations	392
	Exercises	396
Chapter 15:	Cartesian and cocartesian monoidal structures	398
15.1	Unfurling	399
15.2	Cocartesian monoidal structures	406
15.3	Cartesian monoidal structures	415
	Exercises	422
Chapter 16:	Day convolution and the tensor product of spectra	423
16.1	Day convolution operads	424
16.2	Day convolution between symmetric monoidal categories	425
16.3	Examples of Day convolution	434
16.4	Rings and semirings	436

16.5 Spectra as reduced excisive functors	440
16.6 The symmetric monoidal structure on Sp	445
Exercises	449
Chapter 17: The envelope of an ∞ -operad	450
17.1 Factorization systems	451
17.2 Freely adjoining cocartesian lifts	458
17.3 Operadic envelopes	464
17.4 Comparison to Lurie’s operads	471
Exercises	476
Chapter 18: Operadic Day convolution and stabilization	477
18.1 Operadic (co)limits	478
18.2 The multiplicative Yoneda embedding	480
18.3 Day convolution into arbitrary operads	482
18.4 Stabilization of ∞ -operads	484
18.5 Modes and universal properties	490
Exercises	494
Chapter 19: Algebras and modules	495
19.1 Foundations	495
19.2 Relative tensor products and change of algebras	501
19.3 Simplicial models for algebras and modules	509
19.4 Endomorphism algebras	510
19.5 Stable and spectral consequences	513
19.6 Monads and monadicity	517
19.7 Application: connective complex K-theory	520
Exercises	525
Chapter 20: Monoidal Dwyer–Kan localizations and derived categories	526
20.1 Monoidal Dwyer–Kan localizations	526
20.2 The symmetric monoidal derived category	534
Exercises	536

III Background on ∞ -category theory 537

Chapter 21: Adjunctions	539
21.1 Definition and characterizations	539
21.2 (Co)limits and adjunctions	543
21.3 Slice categories	546
21.4 Kan extensions	549
21.5 Cofinal functors	551
21.6 Filtered and sifted ∞ -categories	552
21.7 A comparison theorem	555
21.8 Bousfield localizations	557

Exercises	560
Chapter 22: Presentable ∞ -categories	561
22.1 Presheaves and free cocompletions	561
22.2 Presentable categories and adjoint functors	562
22.3 Compact objects and Ind-categories	565
22.4 Tensor products of presentable ∞ -categories	568
22.5 Presentably symmetric monoidal categories	570
Exercises	571
Chapter 23: Cocartesian fibrations and (un)straightening	572
23.1 Cartesian and cocartesian fibrations	572
23.2 Straightening/unstraightening	579
23.3 Fiberwise criteria	583
23.4 The Hom-functor	586
23.5 Limits and colimits of ∞ -categories	587
23.6 Hom animae in functor categories	591
23.7 Descent and ∞ -topoi	594
23.8 Exponentiability of cartesian fibrations	597
Exercises	598
Chapter 24: Complete Segal animae	601
24.1 ∞ -Categories as complete Segal animae	601
24.2 Filtered colimits and geometric realizations of categories	604
24.3 Homotopy coherent nerves	606
Exercises	614
 IV Solutions and supplementary proofs	 615
Chapter 25: Solutions to Chapter 1	617
25.1 Basic constructions of ∞ -categories	617
25.2 Animae and groupoid cores	619
25.3 Commutative diagrams and composition	620
25.4 Isomorphisms and subcategories	621
25.5 Hom animae	625
 Bibliography	 627
Index of notation	633
Index	635

Change log

The following is an overview of the major updates of these notes:

- **August 2026:** Split the book into a publication version and an online version (this version); the latter uses color and contains additional material not in the published version due to space limitations. Revised Chapter 1, and added Part IV with worked solutions of several foundational exercises on ∞ -category theory. Added index and notation index.
- **July 2026:** Substantial revision. Improved exposition in Chapters 1–7, moved Chapter 3 on cohomology and Chapter 8 on ring spectra and modules, detailed revision of Part II with various reorganizations of the material.

Some new material added: Section 10.3 on classical bordism groups; proof of universal property cartesian monoidal structures in Section 15.3; corrected proof of Proposition 18.4.3; Section 18.5 on mode theory; Sections 19.6 and 21.7 on monads and monadicity; Chapter 22 on presentable ∞ -categories.
- **March 2026:** Revision of Chapter 9, expanded Section 1.1 on the heuristics behind ∞ -categories, added Section 20.1 on monoidal Dwyer-Kan localizations.
- **February 2026:** Revised Chapter 5, added discussion of descent in Section 23.7.
- **January 2026:** Added Chapter 6 on homological algebra, added Section 3.3 on ordinary and cellular homology. Restructured Chapter 19, removed chapter on deformation theory of ring spectra.
- **October 2025:** Substantial revision of Chapters 1-4, reorganized exercises.
- **August 2025:** Integrated remaining part of notes from *Introduction to Higher Algebra*, adding material on ring spectra. Rewrote Chapter 17 on operadic envelopes.
- **July 2025:** Expanded introduction. Ontological status changed from ‘lecture notes’ to ‘book project’ (to be published by Cambridge University Press in 2027).
- **June 2025:** Integrated Part II on the theory of ∞ -operads. This material was written for the lecture course *Introduction to Higher Algebra*, taught at the University of Regensburg in the summer term of 2025.
- **May 2025:** Expanded Chapter 23 on cocartesian fibrations and the straightening/unstraightening correspondence.

-
- **April 2025:** Started a change log.
 - **April 2025:** Substantial revision of the lecture notes:
 - Moved the introduction to ∞ -categories to start, making it Chapter 1;
 - Added Chapter 2 relating ∞ -categorical pushouts/pullbacks to classical homotopy pushouts/pullbacks;
 - Rewrote the chapter on Brown representability in ∞ -categorical language;
 - Added a section on derived ∞ -categories $D(\mathcal{A})$;
 - Reorganized the chapter on complex K-theory.
 - Expanded Appendix 24 on complete Segal anima; added model-independent discussion of homotopy coherent nerves;
 - Added an appendix with suggestions for further reading (which by now has been removed again).
 - **February 2025:** Added two chapters on Thom spectra and Atiyah duality, respectively;
 - **October 2024 - February 2025:** Weekly updates of the course material. We discussed, in this order, Brown representability, an introduction to ∞ -categories, stable ∞ -categories, (commutative) monoids and the recognition principle, complex K-theory, and Bousfield localizations.

Introduction

This book aims to provide an accessible introduction to the field of stable homotopy theory using the language of ∞ -categories. Our primary goal is to develop the technical foundations of the subject, including stability and the ∞ -category of spectra, the theory of ∞ -operads, and some basic theory of ring spectra.

Historical overview

To put into perspective the modern viewpoint on stable homotopy theory, let us take a step back and consider some of the historical developments leading up to it.

In 19th-century topology, mathematicians like Bernhard Riemann, Enrico Betti and Henri Poincaré worked on a classification of manifolds by assigning numerical invariants to them, like the Betti numbers and torsion coefficients. Emmy Noether observed in 1925 that these numbers correspond to the rank and torsion parts of some finitely generated abelian group. This caused a shift in perspective where the homology groups themselves were regarded as the fundamental invariant of a space, and culminated in a full axiomatization of (co)homology in 1945 by Samuel Eilenberg and Norman Steenrod.

One of the central axioms in the axiomatization of homology is the *stability axiom*, which demands that the homology of a space X agrees, up to a degree shift, with that of its suspension ΣX . The key motivation for the notion of stability was the *Freudenthal suspension theorem*, proved around the year 1937 by Hans Freudenthal [Fre37]. It states that if the first n homotopy groups of some pointed CW complex X are trivial, then the canonical comparison map $X \rightarrow \Omega \Sigma X$ induces an isomorphism on π_k for $k \leq 2n$ and a surjection for $k = 2n + 1$. Repeating this for ΣX , $\Sigma^2 X$, etcetera, one finds that the sequence of abelian groups¹

$$\pi_k(X) \rightarrow \pi_k(\Omega \Sigma X) \rightarrow \pi_k(\Omega^2 \Sigma^2 X) \rightarrow \dots$$

eventually stabilizes for every $k \geq 0$. Its colimit is called the *k -th stable homotopy group* of X , denoted $\pi_k^{\text{st}}(X)$. In contrast to their unstable analogues, the stable homotopy groups form a *generalized homology theory*, satisfying all the Eilenberg-Steenrod axioms except for the dimension axiom. Other generalized (co)homology theories discovered in the late 50's were *cobordism theory* by René Thom [Tho54], classifying manifolds with given tangential structures, and *topological K-theory* by Atiyah and Hirzebruch [AH59], classifying vector bundles over a space.

¹For $k \leq 1$, the first few terms are not yet abelian groups.

The success of homology theory in topology inspired mathematicians to develop analogous theories in other fields, leading for example to the theories of group cohomology, Lie algebra cohomology, and sheaf cohomology. An elegant cohesive framework for these developments was established by Henri Cartan and Samuel Eilenberg in 1956 in their monumental book *Homological Algebra*. Their work created a new field of mathematics, providing a powerful, abstract language that unified the various existing theories. The core technical innovation that achieved this was the definition of a *derived functor*, which in the case of the tensor product and the hom-functor results in Tor-groups and Ext-groups, respectively. Indeed, many previously ad-hoc cohomology theories could now be neatly phrased as specific instances of Ext-groups.

Independent of these advances in (co)homology theory, Spanier and Whitehead introduced the set of *stable maps* between two pointed CW-complexes X and Y :

$$[X, Y]_*^{\text{st}} := \text{colim}_n [\Sigma^n X, \Sigma^n Y]_*.$$

Finite pointed CW-complexes and stable maps assemble into a category known as the *Spanier-Whitehead category* [SW53], which they used to investigate duality phenomena of spaces. Building on their work, Spanier's student Lima [Lim59] introduced the first definition of a *spectrum*, and used it to prove a version of Spanier-Whitehead duality for a more general class of spaces. Other definitions for spectra were consequently proposed by Spanier, Kan, and Whitehead, but it was Boardman who gave the first satisfying definition of the homotopy category of spectra, with a clear published account presented by Adams [Ada74]. It is the definition still used today:²

Definition. A *spectrum* X is a sequence of pointed spaces $\{X_n\}_{n \geq 0}$ equipped with homotopy equivalences $X_n \simeq \Omega X_{n+1}$, where $\Omega(-)$ denotes the loop space; if in addition $\pi_k(X_n) = 0$ for $k < n$, we say that X is *connective*.

Some key features of the homotopy category of spectra were that it contained the Spanier-Whitehead category as a full subcategory, that it represented all cohomology theories via Brown's representability theorem, and that it admitted a commutative smash product. It allowed for calculational tools for computing stable homotopy groups, like the Adams spectral sequence [Ada58], Bousfield's localizations at homology theories [Bou79], and the development of chromatic homotopy theory [Rav84; Rav92].

The years that followed saw a surge in the study of 'brave new algebra' [May77], i.e. the importing of statements from algebra into stable homotopy theory. The key technical ingredients that made this possible were the theory of operads by May [May72] and Boardman-Vogt [BV73], and the development of point-set models for the stable homotopy category [LMS86; EKMM97; HSS00]. Most of the familiar algebraic constructions on abelian groups have direct homotopical analogues for spectra:

- There is a *direct sum* $X \oplus Y$ of two spectra, which is simultaneously a product and a coproduct;
- There is a *tensor product* $X \otimes Y$ of two spectra. This leads to the notion of a *ring spectrum*: a spectrum R equipped with a multiplication operation $m: R \otimes R \rightarrow R$ that is 'coherently' associative and unital;

²What we call a 'spectrum' was classically called an ' Ω -spectrum'.

- There is a notion of a *module* over a ring spectrum. Familiar notions for modules over (commutative) rings, like the properties of being projective, perfect, or flat, admit generalizations to the setting of ring spectra;
- Given a spectrum X and a prime p , we may form its p -*localization* $X_{(p)}$ and its p -*completion* $X_p^\wedge$, which have similar behavior as their analogues for abelian groups;
- And so on and so forth...

The main difference between these constructions and their classical counterparts is that they do not need to be derived anymore. For example, unlike the tensor product of abelian groups, the tensor product of spectra is already exact in both variables. In this way, we may think of stable homotopy theory as the ultimate realization of Cartan and Eilenberg’s vision: a setting where derived functors become primary objects of study rather than secondary constructions measuring the failure of exactness. It is for precisely this reason that the techniques and concepts from stable homotopy theory have found significant applications in numerous neighboring fields of mathematics, like (derived) algebraic geometry, algebraic K-theory, geometric topology and representation theory.

Higher category theory

To fully appreciate the pervasive role of (stable) homotopy theory in modern mathematics, we must first embrace a new way of thinking about equality:

Fundamental Principle of Homotopy Theory: When expressing that two entities are equal, we must always specify *how* they are equal by providing a homotopy/isomorphism between them.

This principle is already familiar to most mathematicians when applied to mathematical *objects*: for instance, when we want to express that two abelian groups A and B are ‘the same’, we usually mean that there is an isomorphism $A \cong B$ between them. In homotopy theory, we extend this principle to *morphisms* between objects; for example, two continuous maps $f, g: X \rightarrow Y$ are considered ‘equal’ when equipped with a homotopy $f \sim g$ between them. More broadly, in homotopy theory, the notion of ‘sameness’ is not a *property* (a binary yes/no question) but rather *structure* that must be explicitly provided (a homotopy, isomorphism, etc.). This shift in perspective originated in algebraic topology, but has by now become fundamental in many neighboring fields of mathematics.

While this principle appears straightforward, it rapidly gives rise to intricate ‘coherence problems’. Consider, for instance, the task of defining a homotopical version of an abelian group A . Following the Fundamental Principle, expressing the associativity of addition $+: A \times A \rightarrow A$ requires providing a homotopy between the two maps $A \times A \times A \rightarrow A$ given by $(a, b, c) \mapsto a + (b + c)$ and $(a, b, c) \mapsto (a + b) + c$. Once such a homotopy is given, one can construct two distinct homotopies between the maps $(a, b, c, d) \mapsto a + (b + (c + d))$ and $(a, b, c, d) \mapsto ((a + b) + c) + d$. Requiring these to coincide necessitates the introduction of *homotopies between homotopies*. This process continues indefinitely, leading to an infinite hierarchy of higher homotopies, as laid out for instance by Stasheff [Sta63].

The modern solution to such coherence problems lies in the theory of ∞ -*categories*. Conceptually, an ∞ -category behaves much like a classical 1-category: we may speak of objects and morphisms

that can be composed. But in accordance with the Fundamental Principle, it also comes equipped with a notion of homotopies between morphisms, as well as homotopies between these homotopies, and so on ad infinitum. Homotopical versions of familiar categorical constructions are conveniently formulated within this framework: for example, we may define for any ∞ -category C with finite products a new ∞ -category $\mathrm{CGrp}(C)$ of *commutative groups* in C , where all the required higher coherences are automatically built into the objects.

As we will see, there are ∞ -categories $\mathcal{S}$ and Sp whose objects are spaces³ and spectra, respectively, and whose ‘higher homotopies’ are the expected ones. The following theorem establishes a crucial relation between these two ∞ -categories:

Recognition Principle for Connective Spectra (Boardman–Vogt, May and Segal). There is a fully faithful functor of ∞ -categories

$$\mathrm{CGrp}(\mathcal{S}) \hookrightarrow \mathrm{Sp}$$

whose image consists of the connective spectra.⁴

In other words, we may think of a connective spectrum as a ‘homotopy coherent’ analogue of an abelian group. This result is part of a much broader phenomenon: many algebraic structures have natural ‘homotopy coherent’ analogues in the world of spectra (think for example of the ring spectra mentioned before).

While coherence problems in homotopy theory can be (and historically have been) handled without ∞ -categories, the ∞ -categorical framework builds the Fundamental Principle into its foundations, so that many homotopy-theoretic concepts admit native formulations. The methods from higher category theory are rapidly becoming a standard tool among researchers, and they are slowly making their way into more and more graduate courses at universities. It thus seems like the time is ripe for an accessible resource that develops the foundations of stable homotopy theory using the language of ∞ -categories. Providing such a resource is the primary objective of this book.

From abelian categories to stable ∞ -categories

With the concept of an ∞ -category at hand, we can now return to the question of what makes stable homotopy theory such a suitable framework for doing homological algebra. The short answer is that it allows for a richer notion of *exact sequences*, refining the classical notion of short exact sequences in abelian categories. To see what this means, let us start from the classical situation.

To make sense of short exact sequences in a 1-category C , some basic conditions need to be satisfied. First, C must admit a *zero object* 0 , i.e. an object with the property that for any other $X \in C$ there are unique maps $X \rightarrow 0$ and $0 \rightarrow X$. Second, C must admit kernels and cokernels. Third, consider two maps i and p in C of the form

$$A \xrightarrow{i} B \xrightarrow{p} C$$

³For reasons that will be explained, we will write $\mathcal{A}\mathfrak{n}$ rather than $\mathcal{S}$ for this ∞ -category, and refer to its objects as *animaes*.

⁴The difference between spectra and connective spectra is similar to the difference between arbitrary chain complexes and non-negatively graded chain complexes in homological algebra.

with $pi = 0$, where i is a monomorphism, and where p is an epimorphism. Then it should always be the case that A is the kernel of p if and only if C is the cokernel of i . If these equivalent conditions are satisfied, we say that i and p form a *short exact sequence*.

The story changes a little if the category C is allowed to be an ∞ -category. In this case, the equality $pi = 0$ from before should be replaced by a choice of homotopy $pi \simeq 0$. Following the Fundamental Principle, we must take this choice as part of the data of the exact sequence, and different choices will correspond to different exact sequences. Similarly, we replace the kernel of a map $g: Y \rightarrow Z$ by what is called its *fiber*: the terminal example of an object X equipped with a map $f: X \rightarrow Y$ and a nullhomotopy $gf \simeq 0$; dually, we replace cokernels by *cofibers*. These considerations motivate the following central definition in stable homotopy theory:

Definition. An ∞ -category C is called *stable* if it has a zero object, if it has fibers and cofibers, and if for every composable pair of maps $X \xrightarrow{f} Y \xrightarrow{g} Z$ equipped with a nullhomotopy $gf \simeq 0$ it holds that X is the fiber of g if and only if Z is the cofiber of f .

We refer to the sequences appearing in the definition as *exact sequences*. What makes this definition so powerful is that there are no longer any restrictions on the maps f and g in an exact sequence, in contrast to the situation for abelian categories where i needs to be a monomorphism and p needs to be an epimorphism. Indeed, while the inclusion map $\ker(p) \hookrightarrow B$ of a kernel in an abelian category is automatically a monomorphism, the structure map $\text{fib}(g) \rightarrow Y$ of a fiber in a stable ∞ -category need not be; indeed, *any* map $f: X \rightarrow Y$ arises this way. As an important consequence, it follows that any functor $F: C \rightarrow D$ between stable ∞ -categories is exact as soon as it is either left or right exact. This in particular means that structures like tensor products or internal homs in stable ∞ -categories need no longer be ‘derived’ to turn them into exact functors.

An important example of a stable ∞ -category is the ∞ -category Sp of spectra; in fact, it is in some precise sense the *universal* example. Furthermore, every abelian category $\mathcal{A}$ gives rise to a stable ∞ -category by considering its *derived* ∞ -category $D(\mathcal{A})$, which comes equipped with an inclusion $\mathcal{A} \hookrightarrow D(\mathcal{A})$ that turns short exact sequences in $\mathcal{A}$ into exact sequences in $D(\mathcal{A})$.

As we will explore during the course of this book, most of the usual constructions from homological algebra can be performed directly at the level of stable ∞ -categories, without needing to first perform them at the level of abelian categories and then passing to derived functors. The classical derived functors are then obtained as the homotopy/homology groups of these constructions. For example, every classical commutative ring R defines a commutative ring spectrum, and for two R -modules M and N the Tor- and Ext-groups between M and N may be computed as

$$\text{Tor}_n^R(M, N) \cong \pi_n(M \otimes_R N) \quad \text{and} \quad \text{Ext}_R^n(M, N) \cong \pi_{-n}(\text{hom}_R(M, N)),$$

where $-\otimes_R -$ and $\text{hom}_R(-, -)$ denote the tensor product and internal hom in the ∞ -category of R -module spectra.

The theory of ∞ -operads

The higher coherences resulting from the Fundamental Principle do, however, make it considerably more subtle to define and manipulate algebraic structures in the setting of ∞ -categories than in classical algebra. In particular, it is generally not possible to simply write down such structures ‘by

hand'; some abstract machinery is necessary to keep track of all these coherences. One of the main examples of such machinery is the formalism of ∞ -operads, developed by Lurie [Lur17] based on the classical theory of operads by May [May72] and Boardman and Vogt [BV73]. For this reason, we devote a substantial part of this book to the theory of ∞ -operads, with the aim of providing an accessible introduction to it.

Heuristically, an operad encodes all possible operations associated to a given algebraic structure. For example, there is an operad for commutative rings, which has a single n -ary operation for every $n \geq 0$, corresponding to the multiplication map $R^{\otimes n} \rightarrow R, r_1 \otimes \cdots \otimes r_n \mapsto \prod_{i=1}^n r_i$. A general operad $\mathcal{O}$ will specify for each $n \geq 0$ a set $\mathcal{O}(n)$ of possible n -ary operations. These operations should come with certain identities and compositions; for example, if we plug an n -ary operation into one input of a k -ary operation, this should yield an $(n + k - 1)$ -ary operation. We further require this composition law on operations to be associative and unital.

In the ∞ -categorical setting, the analogous concept of ∞ -operad is more subtle to define, as the unitality and associativity of the composition law are now no longer strict but are only satisfied up to coherent higher homotopies. While there are various approaches in the literature for how to deal with these coherences, we will follow Lurie by defining ∞ -operads via their 'category of operators'. In fact, we will work with a slight variant of Lurie's definition, in which the category $\mathbf{Fin}_*$ of finite pointed sets is replaced by the $(2, 1)$ -category $\mathbf{Span}(\mathbf{Fin})$ of *spans of finite sets*.

Prerequisites and how to read this book

The main prerequisites are 1-category theory and algebraic topology at the level of a first graduate course. In particular, the reader should be comfortable with limits and colimits, adjoint functors, CW complexes, homotopy groups, and singular homology and cohomology. Some familiarity with chain complexes and vector bundles is useful, but the relevant material is recalled when it first becomes essential. No prior familiarity with ∞ -categories, spectra, operads, model categories, or higher algebra is assumed.

This book is available in a published and an online version. The online version contains supplementary sections, identified by "(online only)" in their titles, and uses color in some explanatory diagrams. The supplementary sections are placed at the ends of chapters, so all material appearing in both versions has the same chapter, section, and theorem numbering. Page numbers, however, differ between the two versions.

The book is not intended to be read strictly from beginning to end. Chapter 1 provides the common foundation for Parts I and II, while Part III collects technical background that may be consulted when it is cited. The main complication is that the first two parts depend on one another. Part I constructs spectra before their coherent tensor product has been developed, so Chapter 8 states the required algebra and module theory as a black box. Part II later constructs this tensor product and proves the algebra and module package, while relying on the theory of spectra from Chapter 4 and, in its application to derived categories, on Chapter 6. The resulting dependencies are summarized in Figure 1.

Here are several possible reading routes:

- (1) For a first course in stable homotopy theory, read Chapter 1 followed by Part I in order, treating the forward references from Chapter 8 to Part II as black boxes. The final three chapters may




Figure 1: The dependency structure of the book. A solid arrow means that the source should substantially be read before the target. A dashed arrow marks a minor dependency, or material that may be black-boxed on a first reading; a double-headed dashed arrow records a mutual dependency that is resolved at the level of sections. The chapters of Part III are a reference rather than a linear block, so each of them records instead where its results are used.

be selected according to interest.

- (2) For higher algebra, begin with Chapter 1 and then read Part II in order. The construction of the tensor product of spectra in Chapter 16 uses Chapter 4; the application to connective complex K-theory at the end of Chapter 19 uses Chapter 9; and the derived-category application in Chapter 20 uses Chapter 6. These applications may be skipped if the reader is primarily interested in the abstract operadic theory. The relevant chapters of Part III may be consulted when they are first cited.
- (3) For ring spectra and derived algebra, read Part I through Chapter 7, then read Part II through Chapter 20, omitting the application to connective complex K-theory until after Chapter 9. One may then return to Chapter 8 with its higher-algebra input in place.
- (4) For the categorical foundations, read Chapter 1 followed by Part III.

Content

This book is currently still under construction, and will continue to be revised throughout 2025 and 2026; a stable version can be expected in January 2027. It is a combination of my lecture notes for the course *Introduction to stable homotopy theory*, taught in the winter term 2024/2025, with those for the course *Introduction to higher algebra*, taught in the summer term 2025.

The material is organized into a foundational chapter followed by three parts. Chapter 1 introduces ∞ -categories and anima; it precedes the parts because both mathematical cores rest on it equally. Part I then develops the foundations of stable homotopy theory: (co)homology theories, spectra and stable ∞ -categories, the recognition principle and homological algebra, followed by the principal examples, namely ring spectra, complex K-theory, Thom spectra and duality. Part II develops the theory of ∞ -operads over $\mathrm{Span}(\mathrm{Fin})$ and the higher algebra built on it, including the tensor product of spectra and the theory of algebras and modules that Part I uses. Part III gathers the ∞ -categorical background used throughout: adjunctions, presentable ∞ -categories, cocartesian fibrations and straightening, and complete Segal anima. Each part opens with an introduction describing its chapters in more detail.

Acknowledgments

This book has grown out of two graduate lecture courses on stable homotopy theory and higher algebra taught at the University of Regensburg in 2024/2025. In preparing the manuscript I have benefited from many existing sources on (stable) homotopy theory and higher algebra. I would like to acknowledge some of these influences more explicitly.

The ∞ -categorical approach to stable homotopy theory and higher algebra adopted in this book is deeply indebted to Lurie’s foundational text *Higher Algebra* [Lur17], which remains one of the central references in the field. Most of the theory of ∞ -operads developed in Part II ultimately goes back to this work, even when the order of exposition or the proofs differ from Lurie’s treatment. The exposition in Part I is strongly influenced by the lecture notes of Denis Nardin, *Introduction to stable homotopy theory* [Nar21], in particular in the choice and ordering of topics. For Part II, I have benefited from Haugseng’s introductory note on ∞ -operads [Hau23a], and from Gepner’s survey

article on higher algebra [Gep20]. The brief historical sketch of stable homotopy theory in the introduction draws on the accounts of May and Weibel [May99b; Wei99]. Classical developments of the theory of spectra that have shaped my background, and from which I have borrowed a number of examples and perspectives, may be found in the textbooks of Switzer and Adams [Swi75; Ada74].

Although no single section of the present book follows them closely, Fabian Hebestreit’s lecture course on algebraic and Hermitian K-theory [HW21] was my own first systematic introduction to ∞ -categories and modern stable homotopy theory. It has had a lasting impact on how I think about the subject and, indirectly, on the way the material is presented here.

The model-agnostic approach to ∞ -categories used throughout this book is based on joint work with Denis-Charles Cisinski, Kim Nguyen and Tashi Walde [CCNW26].

The pictorial illustrations for multimorphisms in operads were inspired by [BS22, Figure 1].

I am grateful to many colleagues and students for their help and encouragement. I would like to thank Denis-Charles Cisinski, Daniel Gratzer, Rune Haugseng, Sil Linskens, Maxime Ramzi, Stefan Schwede, Jan Steinebrunner, Tashi Walde, Ferdinand Wagner and Christoph Winges for numerous helpful conversations about the material in this book. I am especially grateful to Marc Hoyois for reading substantial parts of the manuscript and for pointing out several mathematical errors. The participants of the two Regensburg courses on which this book is based provided valuable feedback and many questions that led to clarifications and improvements in the exposition. I also thank Keima Aksaka, Karolis Dembickas, Alissa Doggwiler, Michael J. Glaeser, Tim Henke, Taiga Nakamura, Lorenzo Pascarella, Daniele Velati and Yiming Wang for sending corrections and further suggestions.

Use of generative AI

During the final preparation phase of this textbook⁵, I have made substantial use of the AI coding assistants Claude Code and Codex. This includes:

- Editorial assistance, such as typo correction and LaTeX assistance;
- Mathematical review, such as finding errors and suggesting corrections;
- Substantive mathematical contribution, such as drafting proofs, explanations, and new material.

AI involvement was particularly substantial in the following chapters and sections: Chapter 6, Sections 5.5 and 8.1, Chapter 9, Section 10.3, Sections 15.1 and 15.3, Sections 16.3 and 18.2 to 18.5, Chapter 20, and Chapter 22.

All AI-generated material incorporated into the text was reviewed and edited by me. This involved checking not only the mathematical arguments but also the compatibility with the book’s overall structure and style. I remain solely responsible for the contents of the book and for any errors that remain.

⁵This disclosure was written in July 2026. It will likely be updated before final publication.

1 Introduction to ∞ -categories

This chapter provides the ∞ -categorical foundations that underlie the rest of the book. Our goal is to develop a usable language of ∞ -categories that matches intuition from classical category theory and aligns with the way it is used in stable homotopy theory and higher algebra.

One peculiar aspect of the theory of ∞ -categories is that, while its foundations can be set up in traditional set-based mathematics, the actual practice of the theory involves some significant conceptual shifts regarding several basic mathematical notions. The premise of this book is that the most effective way to get into the field is by learning about these conceptual shifts first, and starting to do some basic mathematics with them, before discussing how these concepts can be modeled within traditional mathematics. We will achieve this by taking an axiomatic approach to ∞ -categories, focusing on just those basic properties of the theory that are needed in practice.

Section 1.1 explains the heuristics behind the theory, and Section 1.2 turns them into working syntax by setting out the axioms we assume. Section 1.3 records the constructions that build new ∞ -categories out of old ones: products and coproducts, pullbacks and pushouts, and functor categories. Section 1.4 then turns to the internal structure of a single ∞ -category: it axiomatizes which commutative diagrams are determined by their visible edges, deduces the Segal axiom and with it the composition of morphisms, and introduces anima and $\mathrm{hom\ anima}$, so that the heuristics of Section 1.1 become part of the formal setup. The section closes with the Rezk axiom. Afterwards, Section 1.5 returns to constructions with opposite categories, subcategories and localizations.

Section 1.6 discusses quasicategories, the standard set-based model. It is the one place in this chapter where we commit to a specific model, and the specifically quasicategorical parts may be skipped. Finally, Section 1.7 treats limits and colimits, and Section 1.8 collects the foundational results that later chapters use without further proof, among them the ∞ -categories $\mathbf{An}$ and $\mathbf{Cat}_\infty$ themselves and the Yoneda lemma.

The exercises in this chapter vary considerably in difficulty. Some are routine checks, while others invite the reader to work through technical consequences of the axioms in detail. The latter may instead be read together with their solutions, and are not checkpoints that must be cleared on a first reading. We use a separate *proof guide* when we state a result for later use but omit its proof from the main exposition. These guides indicate the structure of the omitted argument and may likewise be skipped on a first reading. Complete proofs and further solutions are collected in the online-only Part IV.

1.1 The heuristics behind ∞ -categories

Before setting up the axiomatics of ∞ -categories used in this book, we take a moment to explain the heuristics behind the theory, which will help motivate our choice of setup.

The fundamental principle of homotopy theory

The first concept we need to discuss is the notion of *equality*. In traditional mathematics, equality is typically viewed as a mere property: two mathematical objects are either equal or they are not. In homotopy theory and related fields, however, we encounter a more sophisticated notion of ‘sameness’ where it is not just important *that* two objects are ‘the same’, but precisely *how* they are ‘the same’. In particular, the sameness relationship between objects becomes a mathematical object in its own right: think of a homotopy between two continuous maps, or of an isomorphism between two abelian groups. This perspective is so central to understanding ∞ -categories that it deserves to be elevated to a guiding principle:

Fundamental Principle of Homotopy Theory: When expressing that two entities are equal, we must always specify *how* they are equal by providing a homotopy/isomorphism between them.

To get a feeling for this principle, let us look at three familiar settings in which it already appears:

Example 1.1.1 (Algebraic topology). Algebraic topology provides the most natural example of the Fundamental Principle:

- Two continuous maps $f, g: X \rightarrow Y$ are considered ‘equal up to homotopy’ if we provide a homotopy $H: X \times [0, 1] \rightarrow Y$ between f and g ;
- Two spaces X and Y are then considered ‘homotopically the same’ when equipped with a *homotopy equivalence*: continuous maps $f: X \rightarrow Y$ and $g: Y \rightarrow X$ together with chosen homotopies $gf \sim \text{id}_X$ and $fg \sim \text{id}_Y$.
- When studying diagrams of spaces, we similarly replace strict commutativity by homotopy commutativity. For instance, given a square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ g \downarrow & & \downarrow h \\ Z & \xrightarrow{k} & W \end{array}$$

we do not ask for a strict equality $h \circ f = k \circ g$, but instead require a specified homotopy $H: h \circ f \sim k \circ g$ witnessing how the two compositions are ‘the same’. This perspective naturally leads to *homotopy coherent diagrams*, where we must also specify compatibilities between different choices of homotopies.

Example 1.1.2 (Homological algebra). The phenomenon of ‘equality as structure’ also naturally appears in homological algebra, where it leads us to think about chain complexes not as rigid algebraic objects but as representatives of more flexible homotopical entities:

- Instead of asking whether two chain maps $f, g: C_\bullet \rightarrow D_\bullet$ between chain complexes are *equal*, we ask whether they are *chain homotopic*, meaning that there exists a family of maps $\varphi_n: C_n \rightarrow D_{n+1}$ satisfying $f_n - g_n = \partial^D \circ \varphi_n + \varphi_{n-1} \circ \partial^C$. Likewise, a chain map $C_\bullet \rightarrow D_\bullet$ is regarded as an equivalence if it admits an inverse up to chain homotopy.
- In practice, one often wants to go further and even identify two chain complexes whenever they are connected by a *quasi-isomorphism*, that is, a chain map inducing isomorphisms on all homology groups. Forcing the quasi-isomorphisms to become actual isomorphisms results in the *derived category* of chain complexes.

Example 1.1.3 (Category theory). A third example comes from category theory:

- For two objects X and Y in a 1-category C , the natural notion of equality between them is often not the strict set-theoretic equality, but rather the existence of an isomorphism $f: X \xrightarrow{\sim} Y$. Likewise, universal constructions in category theory are typically unique only ‘up to unique isomorphism’ rather than strictly equal.
- This pattern extends to functors: we wish to think of two functors $F, G: C \rightarrow D$ as ‘the same’ whenever we are given a natural isomorphism $\eta: F \xrightarrow{\sim} G$. If we think of the individual isomorphisms $\eta_X: F(X) \xrightarrow{\sim} G(X)$ as ‘equalities’, then the naturality condition becomes a statement about structural properties of these equalities.
- In monoidal categories, the associativity and unitality conditions are expressed not as strict equalities but as natural isomorphisms: for objects X, Y , and Z , we have an associator $\alpha_{X,Y,Z}: (X \otimes Y) \otimes Z \xrightarrow{\sim} X \otimes (Y \otimes Z)$ and unitors $\lambda_X: 1 \otimes X \xrightarrow{\sim} X$ and $\rho_X: X \otimes 1 \xrightarrow{\sim} X$. The coherence conditions (like the Mac Lane pentagon) then govern how these isomorphisms relate to each other.

The common pattern in all these examples is that it is not enough to know that two things are the same: one must also remember the witness, and often also the compatibilities between different such witnesses.

Animae

Once equality is no longer a mere property, the usual notion of a collection also has to change. In traditional set-based mathematics, the primitive notion of a collection is that of a *set*. The behavior of sets is governed by axioms which strongly reflect the traditional perspective on equality: given a set X , two of its elements x and y are either equal or not equal. The question of *how* x and y are equal is a meaningless one.

If we take the Fundamental Principle fully seriously, we are thus led to a different notion of collection, one in which equality between elements is a structure rather than a property. In recent years, the homotopy theory community has started to adopt the term *anima* for this concept. In terms of traditional mathematics, one may think of animae as homotopy types of topological spaces; we will

discuss this viewpoint at length in Chapter 2. But the philosophy we adopt in this book is that animae are best thought of as primitive objects in their own right, whose role is similar to that of sets in traditional mathematics. In particular, the familiar constructions on sets still have analogues for animae:

- Given animae X and Y , there is a *product* $X \times Y$, whose elements are pairs (x, y) consisting of an element x of X and an element y of Y ;
- Given animae X and Y , there is a *disjoint union* $X \sqcup Y$, an element of which is either an element of X or an element of Y ;
- There is an *empty anima* $\emptyset$, which does not have any elements;
- There is a *one-point anima* $*$, which is non-empty, and in which all elements are equal.

The main difference between animae and sets lies in what it means for two elements x and y of an anima X to be *equal*. Rather than equality being a yes/no question, an equality between x and y should be an actual mathematical object. The collection of all such equalities should then itself form an anima, which we will denote by $X(x, y)$. In turn, if α and β are two such equalities, we may form the anima $X(x, y)(\alpha, \beta)$ of equalities between α and β . This process may be repeated ad infinitum, resulting in equalities between equalities between (...) between equalities.

The following are some examples of animae:

Example 1.1.4. Every set X gives rise to an anima, which we will again denote by X . Given two elements $x, y \in X$, the anima $X(x, y)$ of equalities between x and y is either the one-point anima (if $x = y$) or the empty anima (if $x \neq y$).

Example 1.1.5. Every 1-category C gives rise to an anima $C^\simeq$, called its *groupoid core*. The elements of $C^\simeq$ are the objects of C . The anima $C^\simeq(X, Y)$ between two objects X and Y is the set of isomorphisms between X and Y (regarded as an anima via the previous example).

Example 1.1.6. Every topological space X gives rise to an anima $\Pi_\infty(X)$, called its *underlying anima* (or *fundamental ∞ -groupoid*). Every point in X defines an element of $\Pi_\infty(X)$. Given points $x, y \in X$, every path $p: [0, 1] \rightarrow X$ from x to y defines an equality in $\Pi_\infty(X)$ between x and y . Given two such paths p and q , every homotopy between p and q defines an equality between them in $\Pi_\infty(X)(x, y)$. Every homotopy between homotopies defines an equality of equalities. This pattern repeats ad infinitum. We will study the assignment $X \mapsto \Pi_\infty(X)$ in more detail in Chapter 2.

Just as sets come with maps between them, animae come with maps between them. A map of animae $f: X \rightarrow Y$ assigns to every element x of X an element $f(x)$ of Y , to every equality between x and x' an equality between $f(x)$ and $f(x')$, to every equality between two such equalities an equality between their images, and so on. The role played by bijections of sets is now played by *equivalences of animae*: those maps which admit an inverse, together with the equalities witnessing that both composites are the identity.

Uniqueness

The homotopical perspective on equality also forces us to reconsider uniqueness. Classically, saying that a set X contains a unique element with some given property means that the subset of all valid choices has exactly one element, or equivalently that the map from this subset to the one-point set $*$ is a bijection. Replacing sets by animae and bijections by equivalences, the homotopical analogue reads: the *anima* of all valid choices is *contractible*, meaning that the map from it to the one-point anima $*$ is an equivalence. In this sense, contractibility is the homotopical incarnation of ‘there is exactly one’: it says that a choice exists, that any two choices are equal, that any two equalities between choices are themselves equal, and so on.

The homotopical interpretation of uniqueness appears under various names in the classical literature. In category theory, the phrase ‘up to unique isomorphism’ is often used for objects satisfying a universal property. It expresses precisely that the anima (i.e., groupoid) of such objects, together with the isomorphisms compatible with their universal structures, is contractible. In homotopy theory, a common phrase is ‘up to contractible choice’. For example, to concatenate a finite sequence of loops in a pointed space, one may first choose a subdivision of the interval. Although no particular subdivision is preferred, the anima (i.e., space) of such choices is contractible. In this book, all uniqueness statements should be interpreted in the homotopical sense, and we simply say *unique*.

Categories

Having replaced sets by animae, we can now make the same replacement in the notion of a category. Instead of a set of objects and sets of morphisms between them, we now allow the objects to form an *anima*, and for each pair of objects the morphisms between them to form an *anima* as well. The resulting composition law is then no longer expected to be strictly associative and unital, but only associative and unital up to coherent higher isomorphism. This is the heuristic origin of the notion of an $(\infty, 1)$ -category, or just an ∞ -category for short.⁶

One should not take this heuristic too literally as a definition: replacing sets by animae suggests the right shape of the theory, but turning this into a workable notion of ∞ -category requires encoding highly non-trivial coherence data for composition. This is one of the reasons why, later in this chapter, we will take ∞ -categories as primitive and describe their behavior axiomatically rather than trying to define them directly from this heuristic.

Let us now go through some basic families of examples of ∞ -categories:

Example 1.1.7. Every classical category C defines an ∞ -category, whose anima of objects is the groupoid core $C^\simeq$ from Example 1.1.5, and whose hom animae are the hom sets of C .

Example 1.1.8. Combining the previous example with Example 1.1.6, also every topologically enriched category defines an ∞ -category, by applying Π_∞ to the hom spaces.

Example 1.1.9. Every anima X defines an ∞ -category, again denoted X , whose objects are the elements of X and whose animae of morphisms from x to y are the animae $X(x, y)$ of equalities. The

⁶The terminology ‘ ∞ -category’ is standard but in some ways unfortunate: there are no higher non-invertible morphisms, like one would have in a 2-category or a 3-category.

categorical composition corresponds to the fact that equality is reflexive and transitive. Symmetry of equality is encoded by the condition that every morphism in X admits an inverse.

In fact, in our axiomatic setup, this is how animae will be defined: as those ∞ -categories in which every morphism is invertible. The animae $X(x, y)$ from before are then nothing but the hom animae $\text{Hom}_X(x, y)$ of this ∞ -category.

While the conceptual shifts involved in ∞ -category theory may feel daunting at first, the theory broadly behaves just like classical category theory once one gets used to it. Standard constructions from category theory still make sense for ∞ -categories, such as products, coproducts, pullbacks, pushouts, functor categories, groupoid cores, subcategories, and localizations.

1.2 The language of ∞ -category theory

The previous section explained why one should expect ∞ -categories to arise once equality is treated as structure rather than property. We now turn that heuristic picture into working syntax. We will do so in two steps. First we describe the ambient language of ∞ -categories, functors, and higher isomorphisms. Then we explain how the familiar internal notions of objects, morphisms, and commutative triangles can be recovered from this language by testing against a few small 1-categories. The point of these axioms is not to formalize every coherence once and for all, but to isolate the layer of structure that we will actually use throughout the book.

1.2.1 Categories, functors and natural isomorphisms

We start with the basic objects of this language:

Axiom A.1. (0) There exist mathematical structures called ∞ -categories, denoted C, D, E , etcetera.

- (1) Given ∞ -categories C and D , we may speak of *functors* $f: C \rightarrow D$ between them. Every ∞ -category C has an *identity functor* $\text{id}_C: C \rightarrow C$. Given two functors $f: C \rightarrow D$ and $g: D \rightarrow E$, there is a *composite functor* $g \circ f: C \rightarrow E$.
- (2) Given two functors $f, f': C \rightarrow D$, we may speak of *natural isomorphisms* $\alpha: f \cong f'$. We may sometimes display natural isomorphisms diagrammatically as follows:

$$\begin{array}{ccc} & f & \\ C & \xrightarrow{\quad} & D \\ & \alpha \Downarrow \cong & \\ & f' & \end{array}$$

Every functor f has an *identity isomorphism* $\text{id}_f: f \cong f$. Every natural isomorphism α has an *inverse isomorphism* $\alpha^{-1}: f' \cong f$. Given two natural isomorphisms $\alpha: f \cong f'$ and $\beta: f' \cong f''$, there is a (*vertical*) *composite isomorphism* $\beta \circ \alpha: f \cong f''$.

To avoid cluttering of notation, we will frequently work with natural isomorphisms $f \cong f'$ that have not been given an explicit name.

- (3) Given two natural isomorphisms $\alpha, \alpha': f \cong f'$, we may speak of *isomorphisms of natural isomorphisms* $\alpha \cong \alpha'$, also called *3-isomorphisms* for short. We may again speak of identity

3-isomorphisms $\alpha \cong \alpha$, of composite 3-isomorphisms $\alpha \cong \alpha''$ and of inverse 3-isomorphisms $\alpha' \cong \alpha$.

- (4+) Proceeding in this way, we demand for every natural number n a notion of an n -isomorphism between two $(n - 1)$ -isomorphisms, and we may speak of identity n -isomorphisms, of inverse n -isomorphisms, and of composite n -isomorphisms.

Following the Fundamental Principle, we should think of a natural isomorphism $f \cong f'$ between two functors as expressing that f and f' are ‘equal’. Similarly two n -isomorphisms are regarded as ‘equal’ if we are given an $(n + 1)$ -isomorphism between them.

Axiom A.2. The composition of functors is unital and associative up to isomorphism: for functors $f: C \rightarrow D$, $g: D \rightarrow E$ and $h: E \rightarrow F$, we have natural isomorphisms⁷

$$\lambda_f: \text{id}_D \circ f \cong f, \quad \rho_f: f \circ \text{id}_C \cong f \quad \text{and} \quad \alpha_{h,g,f}: (h \circ g) \circ f \cong h \circ (g \circ f).$$

The composition of (higher) isomorphisms is similarly unital and associative up to isomorphism, and for any natural isomorphism $\alpha: f \cong f'$ there are preferred isomorphisms $\alpha^{-1} \circ \alpha \cong \text{id}_f$ and $\alpha \circ \alpha^{-1} \cong \text{id}_{f'}$.

Since we think of isomorphisms as ‘equalities’, we should make sure that all constructions we use preserve isomorphisms.

Axiom A.3. Composition of functors respects natural isomorphisms: given a natural isomorphism $\alpha: f \cong f'$ of functors $C \rightarrow D$ and a natural isomorphism $\beta: g \cong g'$ of functors $D \rightarrow E$, we obtain a new natural isomorphism $\beta * \alpha: g \circ f \cong g' \circ f'$ of functors $C \rightarrow E$, called the *horizontal composite*.

Remark 1.2.1. To be fully precise one should also write down analogous rules for horizontal composition of n -isomorphisms for all n , express their compatibility with vertical composition, and impose coherence identities for the preferred associators and unitors of functor composition, such as the triangle and pentagon identities. We will not spell out these details here and instead refer to [CCNW26, Section 1.1.2]. Throughout the book we silently assume this coherence data is available whenever we rebracket composite functors or natural isomorphisms.

Definition 1.2.2. A *commutative square of functors*

$$\begin{array}{ccc} C & \xrightarrow{f} & D \\ g \downarrow & & \downarrow h \\ C' & \xrightarrow{f'} & D' \end{array}$$

consists of four functors f, g, h and f' together with a specified natural isomorphism $\alpha: h \circ f \cong f' \circ g$.
A *commutative triangle of functors*

$$\begin{array}{ccc} & D & \\ f \nearrow & & \searrow g \\ C & \xrightarrow{h} & E \end{array}$$

⁷We have given them names to make clear that the axiom provides preferred choices for such isomorphisms, but in practice we often don’t write out the names for these isomorphisms each time.

consists of three functors f , g and h together with a specified natural isomorphism $\alpha: h \cong g \circ f$. We may similarly define a *commutative square/triangle of n -isomorphisms*.

Definition 1.2.3. A functor $f: C \rightarrow D$ is said to be an *equivalence* if there exists another functor $g: D \rightarrow C$ and natural isomorphisms $\text{id}_C \cong gf$ and $\text{id}_D \cong fg$. In this case we call g an *inverse* to f .

Exercise 1.2.4. Show that a functor f is an equivalence if and only if it admits both a section (i.e., a g such that $\text{id}_D \cong fg$) and a retraction (i.e. an h such that $\text{id}_C \cong hf$), and that in this case we have $g \cong h$.

Notation 1.2.5. It follows immediately from the previous exercise that if g and h are both inverses to f , then there is an isomorphism $g \cong h$. When an inverse to f has been chosen, we denote it by $f^{-1}: D \rightarrow C$.

Exercise 1.2.6. Show that if $f: C \rightarrow D$ and $f': D \rightarrow E$ are equivalences with inverses $g: D \rightarrow C$ and $g': E \rightarrow D$, then also the composite $f'f: C \rightarrow E$ is an equivalence with inverse $gg': E \rightarrow C$.

Lemma 1.2.7. *Equivalences satisfy the 2-out-of-3 property: given functors $f: C \rightarrow D$ and $g: D \rightarrow E$, if two of the three functors f , g and gf are equivalences, then so is the third.*

Proof. If g and f are equivalences, then so is gf by Exercise 1.2.6. If f and gf are equivalences, then we may rewrite g as the composite $g \cong g \circ \text{id}_D \cong g \circ (f \circ f^{-1}) \cong (gf) \circ f^{-1}$, so that g is the composite of two equivalences and hence itself an equivalence. A similar argument shows that f is an equivalence whenever g and gf are equivalences. $\square$

Exercise 1.2.8. Show that equivalences of ∞ -categories satisfy the 2-out-of-6 property: given functors $f: C \rightarrow D$, $g: D \rightarrow E$ and $h: E \rightarrow F$ such that gf and hg are equivalences, also the functors f , g , h and hgf are equivalences.

To make contact with familiar mathematics, we also assume that every classical 1-category may be regarded as a special case of an ∞ -category.

Axiom B. Every classical 1-category C gives rise to an ∞ -category, which we will abusively again denote by C . Given 1-categories C and D , every 1-functor $f: C \rightarrow D$ of 1-categories gives rise to a functor of ∞ -categories, and every ∞ -functor comes from a 1-functor. Every natural isomorphism $\alpha: f \xrightarrow{\cong} f'$ in the classical sense gives rise to a natural isomorphism of functors of ∞ -categories. These assignments preserve composition of functors and natural isomorphisms.

Example 1.2.9. In particular, every partially ordered set $(P, \leq)$ may be regarded as an ∞ -category by taking $\text{Hom}_P(p, q)$ to be a singleton if $p \leq q$ and empty otherwise.

1.2.2 Objects, morphisms and commutative triangles

So far, this language describes ∞ -categories from the outside, in terms of functors between them. We now explain how to recover the familiar internal notions of objects, morphisms, and commutative triangles from this same setup. Having embedded classical 1-categories into our theory, we can do this by probing an ∞ -category with a few very small ones:

Notation 1.2.10. For $n \geq 0$, we write $[n]$ for the following linearly ordered set:

$$[n] := \{0 \leq 1 \leq \dots \leq n\}.$$

For $n \geq 1$ and $0 \leq i \leq n$, we define the *face map* $d_i: [n-1] \rightarrow [n]$ as the unique injective map of posets whose image leaves out i . Similarly we define the *degeneracy map* $s_i: [n+1] \rightarrow [n]$ as the unique surjective map of posets that hits i twice.

We denote the associated three ∞ -categories again by $[0]$, $[1]$ and $[2]$. They may be displayed as follows:

$$0, \quad 0 \longrightarrow 1 \quad \text{and} \quad \begin{array}{ccc} & 1 & \\ \nearrow & & \searrow \\ 0 & \longrightarrow & 2. \end{array}$$

The five face maps $d_0, d_1: [0] \rightarrow [1]$ and $d_0, d_1, d_2: [1] \rightarrow [2]$ and the relations between them may be displayed as follows:

$$\begin{array}{ccccc} & & 0 & & \\ & d_0 \swarrow & & \searrow d_1 & \\ & 1 & \cong & 0 & \\ & \nearrow d_2 & & \nwarrow d_0 & \\ & 0 & & 1 & \\ & \nwarrow d_1 & & \nearrow d_0 & \\ & 0 & & 1 & \\ & \xrightarrow{d_1} & 0 \longrightarrow 1 & \xleftarrow{d_0} & 0. \end{array}$$

The two degeneracy maps $s_0, s_1: [2] \rightarrow [1]$ may be interpreted as the following ‘projections’:

$$\begin{array}{ccc} \begin{array}{ccc} 0 & \longrightarrow & 1 \\ & \searrow & \downarrow \\ & & 2 \end{array} & \xrightarrow{s_0} & \begin{array}{c} 0 \\ \downarrow \\ 1 \end{array} & \xleftarrow{s_1} & \begin{array}{ccc} & 0 & \\ \nearrow & & \downarrow \\ 2 & \longleftarrow & 1. \end{array} \end{array}$$

Definition 1.2.11. Let C be an ∞ -category.

- (1) An *object* of C is a functor $x: [0] \rightarrow C$;
- (2) A *morphism* of C is a functor $f: [1] \rightarrow C$. We write $f(0)$ for its *source/domain* and $f(1)$ for its *target/codomain*, which are defined as the following composites:

$$f(0): [0] \xrightarrow{d_1} [1] \xrightarrow{f} C \quad \text{and} \quad f(1): [0] \xrightarrow{d_0} [1] \xrightarrow{f} C.$$

We will also write $f: x \rightarrow y$ to indicate that f comes with isomorphisms $x \cong f(0)$ and $y \cong f(1)$.

- (3) A *commutative triangle in C* is a functor $\sigma: [2] \rightarrow C$. We will frequently denote commutative triangles by

$$\sigma = \begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{h} & z, \end{array}$$

where $f = \sigma \circ d_2$, $g = \sigma \circ d_0$ and $h = \sigma \circ d_1$. We will often leave commutative triangles unnamed, and only label their three edges.

Warning 1.2.12. While we may speak of *objects of C* , we may not speak of a *set of objects of C* ! (We will introduce the *anima of objects* in Axiom E below.)

Remark 1.2.13. We want to think of a commutative triangle σ as saying that h is a candidate composite of f and g . Note that at this stage we do not yet say anything about the existence nor the uniqueness of composites; we get back to this in Section 1.4.2.

Example 1.2.14. Given an object x of C , we define the *identity morphism* $\text{id}_x: x \rightarrow x$ as the composite $[1] \rightarrow [0] \xrightarrow{x} C$. Since the composites

$$[0] \xrightarrow{d_1} [1] \xrightarrow{s_0} [0] \quad \text{and} \quad [0] \xrightarrow{d_0} [1] \xrightarrow{s_0} [0]$$

are naturally isomorphic to the identity functor $\text{id}_{[0]}$, we see that the source and target of id_x are indeed isomorphic to x .

Exercise 1.2.15. Construct for every morphism $f: x \rightarrow y$ commutative triangles in C of the form

$$\begin{array}{ccc} & x & \\ \text{id}_x \nearrow & & \searrow f \\ x & \xrightarrow{f} & y \end{array} \quad \text{and} \quad \begin{array}{ccc} & y & \\ f \nearrow & & \searrow \text{id}_y \\ x & \xrightarrow{f} & y. \end{array}$$

Once composition of morphisms is available, this exercise will give the expected relations $f \cong f \circ \text{id}_x$ and $f \cong \text{id}_y \circ f$, see Lemma 1.4.8.

1.3 Basic constructions of ∞ -categories

We now record the ambient constructions on ∞ -categories that we will use throughout the book: terminal and initial categories, products and coproducts, pullbacks, functor categories, and pushouts. Each is introduced through its universal property, expressed only to the level of coherence that we will actually need later on.

1.3.1 Products and coproducts

We start with products and coproducts of ∞ -categories.

Axiom C.1 (Products and coproducts of ∞ -categories). Let I be a set and let C_i be an ∞ -category for every $i \in I$.

- (1) There is a *product* $\prod_{i \in I} C_i$, equipped with *projection functors* $\text{pr}_i: \prod_{i \in I} C_i \rightarrow C_i$ for every i . Given another ∞ -category T and functors $f_i: T \rightarrow C_i$ for every i , we obtain a functor $\langle f_i \rangle: T \rightarrow \prod_{i \in I} C_i$ together with natural isomorphisms $\text{pr}_i \circ \langle f_i \rangle \cong f_i$. Given two functors $g, h: T \rightarrow \prod_{i \in I} C_i$ and natural isomorphisms $\text{pr}_i \circ g \cong \text{pr}_i \circ h$, there is a natural isomorphism $g \cong h$.
- (2) There is a *coproduct* or *disjoint union* $\sqcup_{i \in I} C_i$, equipped with *inclusion functors* $\text{incl}_i: C_i \hookrightarrow \sqcup_{i \in I} C_i$ for every i . Given another ∞ -category T and functors $f_i: C_i \rightarrow T$ for every i , we obtain a functor $\langle f_i \rangle: \sqcup_{i \in I} C_i \rightarrow T$ together with natural isomorphisms $\langle f_i \rangle \circ \text{incl}_i \cong f_i$. Given two functors $g, h: \sqcup_{i \in I} C_i \rightarrow T$ and natural isomorphisms $g \circ \text{incl}_i \cong h \circ \text{incl}_i$, there is a natural isomorphism $g \cong h$.

Notation 1.3.1. For the empty-index cases, we write the resulting product and coproduct as $*$ and $\emptyset$, respectively. Thus every ∞ -category C comes equipped with canonical functors

$$C \rightarrow *, \quad \emptyset \rightarrow C.$$

Remark 1.3.2. We further assume that the canonical functor $[0] \rightarrow *$ is an equivalence. In particular, objects of an ∞ -category C may equally well be regarded as functors $[0] \rightarrow C$ or as functors $* \rightarrow C$.

In these empty-index cases we also silently assume one extra layer of coherence: any two natural isomorphisms between two functors $T \rightarrow *$ are connected by a 3-isomorphism, and similarly for functors $\emptyset \rightarrow T$. This mild strengthening is only used when comparing products with pullbacks over $*$ and coproducts with pushouts under $\emptyset$, for instance in Exercise 1.3.10.

Notation 1.3.3. When $I = \{1, 2\}$, we write $C_1 \times C_2$ and $C_1 \sqcup C_2$ for the resulting product and coproduct. Given functors $f_1: T \rightarrow C_1$ and $f_2: T \rightarrow C_2$, we write $(f_1, f_2): T \rightarrow C_1 \times C_2$ for the induced functor. Given functors $g_1: C_1 \rightarrow T$ and $g_2: C_2 \rightarrow T$, we write $\langle g_1, g_2 \rangle: C_1 \sqcup C_2 \rightarrow T$ for the induced functor.

Taking $T = *$, we see that objects of $C_1 \times C_2$ have the form (x, y) , where x is an object of C_1 and y is an object of C_2 .

Definition 1.3.4. Let C and D be ∞ -categories and let x be an object of D , regarded as a functor $x: * \rightarrow D$. We define the *constant functor* $\text{const}_x: C \rightarrow D$ as the following composite:

$$\text{const}_x: C \xrightarrow{p_C} * \xrightarrow{x} D.$$

Definition 1.3.5 (Contractible category). An ∞ -category C is called *contractible* if the functor $p_C: C \rightarrow *$ is an equivalence.

Observe that an ∞ -category C is contractible if and only if there exists an object $x: * \rightarrow C$ such that the composite $\text{const}_x: C \rightarrow C$ is naturally isomorphic to the identity $\text{id}_C: C \rightarrow C$. A natural isomorphism $H: \text{const}_x \cong \text{id}_C$ is called a *contraction of C onto x* .

The familiar symmetries of products and coproducts follow from these universal properties just as in ordinary category theory, and we will use them freely.

Exercise 1.3.6. Show that (co)products of ∞ -categories are associative, commutative and unital:

$$C \times D \xrightarrow{\sim} D \times C, \quad (C \times D) \times E \xrightarrow{\sim} C \times (D \times E), \quad C \times * \xrightarrow{\sim} C \xrightarrow{\sim} * \times C,$$

and

$$C \sqcup D \xrightarrow{\sim} D \sqcup C, \quad (C \sqcup D) \sqcup E \xrightarrow{\sim} C \sqcup (D \sqcup E), \quad \emptyset \sqcup C \xrightarrow{\sim} C \xrightarrow{\sim} C \sqcup \emptyset.$$

Construction 1.3.7. The formation of products and coproducts of ∞ -categories is functorial: given two functors $f: C \rightarrow C'$ and $g: D \rightarrow D'$, we obtain new functors

$$\begin{aligned} f \times g &:= (f \circ \text{pr}_C, g \circ \text{pr}_D): C \times D \rightarrow C' \times D' \\ f \sqcup g &:= \langle \text{incl}_{C'} \circ f, \text{incl}_{D'} \circ g \rangle: C \sqcup D \rightarrow C' \sqcup D'. \end{aligned}$$

We leave it to the reader to verify that there are natural isomorphisms

$$\begin{aligned} \text{id}_C \times \text{id}_D &\cong \text{id}_{C \times D} & \text{and} & & (f' \times g') \circ (f \times g) &\cong (f' \circ f) \times (g' \circ g), \\ \text{id}_C \sqcup \text{id}_D &\cong \text{id}_{C \sqcup D} & \text{and} & & (f' \sqcup g') \circ (f \sqcup g) &\cong (f' \circ f) \sqcup (g' \circ g) \end{aligned}$$

for functors $f: C \rightarrow C'$, $f': C' \rightarrow C''$, $g: D \rightarrow D'$ and $g': D' \rightarrow D''$.

1.3.2 Pullbacks of ∞ -categories

We will now axiomatize the *pullback* of two functors $f: C \rightarrow E$ and $g: D \rightarrow E$ of ∞ -categories. This will be similar to the axiom for the product $C \times D$, but it is more involved due to the fact that we need to record compatibilities with the structure maps to E .

Axiom C.2 (Pullbacks of ∞ -categories). Consider two functors $f: C \rightarrow E$ and $g: D \rightarrow E$. Their *pullback*, or *fiber product over E* , is an ∞ -category $C \times_E D$ equipped with functors $\text{pr}_C: C \times_E D \rightarrow C$ and $\text{pr}_D: C \times_E D \rightarrow D$ and a natural isomorphism $f \circ \text{pr}_C \cong g \circ \text{pr}_D$, or equivalently a commutative square

$$\begin{array}{ccc} C \times_E D & \xrightarrow{\text{pr}_C} & C \\ \text{pr}_D \downarrow & & \downarrow f \\ D & \xrightarrow{g} & E. \end{array}$$

Given functors $t_C: T \rightarrow C$ and $t_D: T \rightarrow D$ equipped with a natural isomorphism $\alpha: f \circ t_C \cong g \circ t_D$, we obtain a functor $t = (t_C, t_D): T \rightarrow C \times_E D$. This functor comes equipped with natural isomorphisms $\text{pr}_C \circ t \cong t_C$ and $\text{pr}_D \circ t \cong t_D$, and the composite isomorphism

$$f \circ t_C \cong f \circ \text{pr}_C \circ t \cong g \circ \text{pr}_D \circ t \cong g \circ t_D$$

is isomorphic to α . We summarize this by the following dashed-arrow diagram:⁸

$$\begin{array}{ccccc} T & & & & \\ & \searrow^{(t_C, t_D)} & & \searrow^{t_C} & \\ & C \times_E D & \xrightarrow{\text{pr}_C} & C & \\ & \text{pr}_D \downarrow & & \downarrow f & \\ & D & \xrightarrow{g} & E. & \\ & \swarrow_{t_D} & & \swarrow_{t_C} & \end{array}$$

⁸Here, and throughout the book, dashed arrows indicate morphisms that are being constructed or are otherwise under discussion. Such diagrams are understood to commute up to the evident natural isomorphisms and compatibilities.

Similarly, if $t, t' : T \rightarrow C \times_E D$ are two functors and $\alpha : \text{pr}_C \circ t \cong \text{pr}_C \circ t'$ and $\beta : \text{pr}_D \circ t \cong \text{pr}_D \circ t'$ are natural isomorphisms compatible over E , in the sense that the following square commutes,

$$\begin{array}{ccc} f \circ \text{pr}_C \circ t & \xrightarrow[\cong]{f \circ \alpha} & f \circ \text{pr}_C \circ t' \\ \cong \downarrow & & \downarrow \cong \\ g \circ \text{pr}_D \circ t & \xrightarrow[\cong]{g \circ \beta} & g \circ \text{pr}_D \circ t'. \end{array}$$

Then we obtain a natural isomorphism $(\alpha, \beta) : t \cong t'$ satisfying $\text{pr}_C \circ (\alpha, \beta) \cong \alpha$ and $\text{pr}_D \circ (\alpha, \beta) \cong \beta$, and inducing an isomorphic isomorphism of natural isomorphisms in the previous square.

Remark 1.3.8. Despite the heavier notation, Axiom C.2 has exactly the same formal shape as the product part of the previous axiom: it introduces an ∞ -category with structure, explains how compatible data on T determines a functor $T \rightarrow X$, and says that such functors are determined up to natural isomorphism by that data. For a fully precise formulation of the final compatibility condition, we refer to [CCNW26, Axiom B.5].

Exercise 1.3.9. Show that the fiber product is commutative, associative and unital: for functors $C \rightarrow E, D \rightarrow E$ and $B \rightarrow E$, there are equivalences

$$C \times_E D \xrightarrow{\sim} D \times_E C, \quad B \times_E (C \times_E D) \xrightarrow{\sim} (B \times_E C) \times_E D, \quad C \times_E E \xrightarrow{\sim} C.$$

Exercise 1.3.10. Given ∞ -categories C and D , produce an equivalence $C \times D \xrightarrow{\sim} C \times_* D$.

Exercise 1.3.11. The construction of pullbacks of ∞ -categories is functorial: given a commutative diagram

$$\begin{array}{ccccc} C & \xrightarrow{f} & E & \xleftarrow{g} & D \\ \varphi \downarrow & & \downarrow \chi & & \downarrow \psi \\ C' & \xrightarrow{f'} & E' & \xleftarrow{g'} & D' \end{array}$$

there is an induced functor

$$\varphi \times_{\chi} \psi := (\varphi \circ \text{pr}_C, \psi \circ \text{pr}_D) : C \times_E D \rightarrow C' \times_{E'} D'.$$

Show that if each of the functors φ, ψ and χ is an equivalence, then so is $\varphi \times_{\chi} \psi$. Deduce that equivalences are stable under pullback: if $g : D \rightarrow E$ is an equivalence and $f : C \rightarrow E$ is an arbitrary functor, then the projection $C \times_E D \rightarrow C$ is an equivalence.

Pullbacks of categories lead to the notion of a *fiber* of a functor:

Definition 1.3.12. Let $f : C \rightarrow D$ be a functor. For every object x of D , we define the *fiber* of f over x as the pullback

$$\begin{array}{ccc} C_x & \longrightarrow & C \\ \downarrow & \lrcorner & \downarrow f \\ * & \xrightarrow{x} & D. \end{array}$$

The definition of pullbacks of ∞ -categories naturally leads to the notion of a pullback square.

Definition 1.3.13 (Pullback square). A commutative square

$$\begin{array}{ccc} T & \xrightarrow{s} & C \\ t \downarrow & & \downarrow f \\ D & \xrightarrow{g} & E \end{array}$$

is called a *pullback square* if the induced functor $(s, t): T \rightarrow C \times_E D$ is an equivalence.

Exercise 1.3.14 (Iterated pullbacks). Consider a commutative diagram

$$\begin{array}{ccccc} C_1 & \xrightarrow{g_1} & C_2 & \xrightarrow{g_2} & C_3 \\ f_1 \downarrow & & \downarrow f_2 & & \downarrow f_3 \\ D_1 & \xrightarrow{h_1} & D_2 & \xrightarrow{h_2} & D_3. \end{array}$$

Construct an equivalence

$$D_1 \times_{D_3} C_3 \xrightarrow{\sim} D_1 \times_{D_2} (D_2 \times_{D_3} C_3).$$

Hint: define one direction by $(\text{pr}_{D_1}, (h_1 \circ \text{pr}_{D_1}, \text{pr}_{C_3}))$ and construct an inverse by projecting to D_1 and C_3 .

Lemma 1.3.15 (Pasting lemma for pullback squares). Consider a commutative diagram

$$\begin{array}{ccccc} C_1 & \xrightarrow{g_1} & C_2 & \xrightarrow{g_2} & C_3 \\ f_1 \downarrow & & \downarrow f_2 & & \downarrow f_3 \\ D_1 & \xrightarrow{h_1} & D_2 & \xrightarrow{h_2} & D_3. \end{array}$$

If the right-hand square is a pullback square, then the left-hand square is a pullback square if and only if the outer rectangle is a pullback square.

Proof. By the previous exercise there is a preferred equivalence

$$D_1 \times_{D_3} C_3 \xrightarrow{\sim} D_1 \times_{D_2} (D_2 \times_{D_3} C_3).$$

If the right-hand square is a pullback square, the functor $C_2 \rightarrow D_2 \times_{D_3} C_3$ is an equivalence. Hence, by Exercise 1.3.11, the induced functor $D_1 \times_{D_2} C_2 \rightarrow D_1 \times_{D_2} (D_2 \times_{D_3} C_3)$ is an equivalence. The claim now follows from 2-out-of-3 applied to the square

$$\begin{array}{ccc} C_1 & \xrightarrow{(f_1, g_1)} & D_1 \times_{D_2} C_2 \\ (f_1, g_2 \circ g_1) \downarrow & & \downarrow \simeq \\ D_1 \times_{D_3} C_3 & \xrightarrow{\simeq} & D_1 \times_{D_2} (D_2 \times_{D_3} C_3). \end{array}$$

□

Exercise 1.3.16. Consider a commutative square

$$\begin{array}{ccc} C & \xrightarrow{f} & D \\ g \downarrow & & \downarrow h \\ E & \xrightarrow{k} & F \end{array}$$

and assume that h is an equivalence. Show that g is an equivalence if and only if the square is a pullback square.

We record one further property of coproducts.

Axiom C.3 (Universality of coproducts). Let I be a set, and let D_i be an ∞ -category for every $i \in I$.

(1) Given functors $f_i : C_i \rightarrow D_i$ for all $i \in I$, the commutative square

$$\begin{array}{ccc} C_i & \xrightarrow{\text{incl}_i} & \bigsqcup_{i \in I} C_i \\ f_i \downarrow & & \downarrow \bigsqcup_{i \in I} f_i \\ D_i & \xrightarrow{\text{incl}_i} & \bigsqcup_{i \in I} D_i \end{array}$$

is a pullback square.

(2) For a functor $h : E \rightarrow \bigsqcup_{i \in I} D_i$, the functor

$$\langle \text{pr}_E \rangle_i : \bigsqcup_{i \in I} (E \times_{\bigsqcup_{i \in I} D_i} D_i) \rightarrow E$$

is an equivalence.

(3) The inclusions $\{i\} \hookrightarrow I$ for $i \in I$ induce an equivalence $\bigsqcup_{i \in I} \{i\} \xrightarrow{\sim} I$.

Remark 1.3.17. This is the categorical analogue of three basic facts about disjoint unions of sets: base change along an inclusion picks out the corresponding summand, every map into a disjoint union decomposes the source into the corresponding fibers, and every set is the disjoint union of its elements.

1.3.3 Functor categories

Functor categories let us package families of functors internally, and they will also allow us to define pushouts by duality in the next subsection.

Axiom C.4 (Functor category). The *functor category* between two ∞ -categories C and D is an ∞ -category denoted by $\text{Fun}(C, D)$. It comes equipped with a functor $\text{ev} : \text{Fun}(C, D) \times C \rightarrow D$, called the *evaluation functor*. If E is another ∞ -category, then every functor $f : E \times C \rightarrow D$ gives rise to a functor $f_c : E \rightarrow \text{Fun}(C, D)$ called the *currying* of f . It comes equipped with a natural isomorphism between f and the composite

$$E \times C \xrightarrow{f_c \times \text{id}_C} \text{Fun}(C, D) \times C \xrightarrow{\text{ev}} D.$$

Given two functors $g, h : E \rightarrow \text{Fun}(C, D)$ and a natural isomorphism $\alpha : \text{ev} \circ (g \times \text{id}_C) \cong \text{ev} \circ (h \times \text{id}_C)$, we obtain a natural isomorphism $\beta : g \cong h$ such that $\alpha \cong \text{ev} \star (\beta \times \text{id}_C)$.

Remark 1.3.18. Given a functor $g : E \rightarrow \text{Fun}(C, D)$, we define its *uncurrying* $g^u : E \times C \rightarrow D$ as the composite

$$E \times C \xrightarrow{g \times \text{id}_C} \text{Fun}(C, D) \times C \xrightarrow{\text{ev}} D.$$

Thus currying and uncurrying pass between functors $E \times C \rightarrow D$ and functors $E \rightarrow \text{Fun}(C, D)$, and going back and forth both ways is naturally isomorphic to the identity. In the case $E = *$, we observe that objects $* \rightarrow \text{Fun}(C, D)$ of the functor category correspond to functors $C \simeq C \times * \rightarrow D$ from C to D , justifying the terminology for $\text{Fun}(C, D)$.

Exercise 1.3.19. Show that currying and uncurrying are functorial in the parameter category E and compatible with natural isomorphisms.

The functoriality of the product naturally provides functoriality for the functor categories:

Construction 1.3.20 (Functoriality of functor categories in C and D). Given a functor $g: D \rightarrow E$, we define the functor $g \circ -: \text{Fun}(C, D) \rightarrow \text{Fun}(C, E)$ as the currying of the composite

$$\text{Fun}(C, D) \times C \xrightarrow{\text{ev}} D \xrightarrow{g} E.$$

Similarly, given a functor $f: C \rightarrow D$, we define the functor $- \circ f: \text{Fun}(D, E) \rightarrow \text{Fun}(C, E)$ as the currying of the composite

$$\text{Fun}(D, E) \times C \xrightarrow{\text{id} \times f} \text{Fun}(D, E) \times D \xrightarrow{\text{ev}} E.$$

In a completely analogous way, every natural isomorphism $\beta: g \cong g'$ of functors $D \rightarrow E$ induces a natural isomorphism $(\beta \circ -): (g \circ -) \cong (g' \circ -)$ of functors $\text{Fun}(C, D) \rightarrow \text{Fun}(C, E)$, and similarly for the construction $- \circ f$.

Alternative notations for these functors that we will frequently use are g_* and f^* :

$$g_* := g \circ -: \text{Fun}(C, D) \rightarrow \text{Fun}(C, E), \quad f^* := - \circ f: \text{Fun}(D, E) \rightarrow \text{Fun}(C, D).$$

Exercise 1.3.21. Formulate and prove that the assignments $g \mapsto (g \circ -)$ and $f \mapsto (- \circ f)$ are functorial, in the sense that they respect identity functors and composition of functors.

Exercise 1.3.22. Given ∞ -categories C, D and E , construct a composition functor

$$- \circ -: \text{Fun}(D, E) \times \text{Fun}(C, D) \rightarrow \text{Fun}(C, E).$$

Show that it reduces to the functors $g \circ -$ and $- \circ f$ when fixing one of the two variables.

Exercise 1.3.23. Let $f, g: C \rightarrow D$ be two functors of ∞ -categories, regarded as objects of $\text{Fun}(C, D)$. We may define a new ∞ -category $(f \cong g)$ via the following pullback square:

$$\begin{array}{ccc} (f \cong g) & \xrightarrow{\quad} & \text{Fun}(C, D) \\ \downarrow & \lrcorner & \downarrow \Delta \\ * & \xrightarrow{(f, g)} & \text{Fun}(C, D) \times \text{Fun}(C, D). \end{array}$$

Show that every natural isomorphism $\alpha: f \cong g$ defines an object of $(f \cong g)$. Conversely, show that every object of $(f \cong g)$ defines a natural isomorphism $\alpha: f \cong g$.

Proposition 1.3.24 (Internal universal properties). *Let T be an ∞ -category and let $\{C_i\}_{i \in I}$ be a set-indexed collection of ∞ -categories. The canonical functors below are equivalences:*

$$\begin{aligned} \text{Fun}(T, *) &\xrightarrow{\sim} *, & \text{Fun}(\emptyset, T) &\xrightarrow{\sim} *, & \text{Fun}(*, T) &\xrightarrow{\sim} T, \\ \text{Fun}\left(T, \prod_{i \in I} C_i\right) &\xrightarrow{\sim} \prod_{i \in I} \text{Fun}(T, C_i), & \text{Fun}\left(\bigsqcup_{i \in I} C_i, T\right) &\xrightarrow{\sim} \prod_{i \in I} \text{Fun}(C_i, T). \end{aligned}$$

Moreover, for functors $f: C \rightarrow E$ and $g: D \rightarrow E$ there is an equivalence

$$\mathrm{Fun}(T, C \times_E D) \xrightarrow{\sim} \mathrm{Fun}(T, C) \times_{\mathrm{Fun}(T, E)} \mathrm{Fun}(T, D),$$

and for ∞ -categories C and D there is an equivalence

$$\mathrm{Fun}(T, \mathrm{Fun}(C, D)) \xrightarrow{\sim} \mathrm{Fun}(T \times C, D).$$

Proof guide. Construct the comparison functors from the projections, inclusions, evaluation, and currying maps. To construct their inverses as actual functors, first uncurry: the product and pullback axioms assemble the resulting evaluation maps, after which one curries again. In the coproduct case, first use universality of coproducts to distribute the product with the parameter category over $\bigsqcup_i C_i$. The uniqueness clauses then give the two inverse relations. A complete proof is given in Solution 25.1.1. See also [CCNW26, Section 1.4].

1.3.4 Pushouts of ∞ -categories

Using functor categories, we may now dualize the notion of pullback square and define pushouts:

Definition 1.3.25 (Pushout square). A commutative square of ∞ -categories

$$\begin{array}{ccc} C' & \xrightarrow{u} & C \\ f' \downarrow & & \downarrow f \\ D' & \xrightarrow{v} & D \end{array}$$

is called a *pushout square* (or *cocartesian*) if, for every ∞ -category E , the induced square

$$\begin{array}{ccc} \mathrm{Fun}(D, E) & \xrightarrow{v^*} & \mathrm{Fun}(D', E) \\ f^* \downarrow & & \downarrow (f')^* \\ \mathrm{Fun}(C, E) & \xrightarrow{u^*} & \mathrm{Fun}(C', E) \end{array}$$

is a pullback square.

Axiom C.5 (Pushout of ∞ -categories). For functors $f: E \rightarrow C$ and $g: E \rightarrow D$, there exists a pushout square

$$\begin{array}{ccc} E & \xrightarrow{f} & C \\ g \downarrow & \lrcorner & \downarrow \mathrm{incl}_C \\ D & \xrightarrow{\mathrm{incl}_D} & C \sqcup_E D. \end{array}$$

We refer to the ∞ -category $C \sqcup_E D$ as the *pushout* of C and D along E .

Lemma 1.3.26 (Pasting lemma for pushout squares). Consider a commutative diagram

$$\begin{array}{ccccc} C_1 & \xrightarrow{g_1} & C_2 & \xrightarrow{g_2} & C_3 \\ f_1 \downarrow & & \downarrow f_2 & & \downarrow f_3 \\ D_1 & \xrightarrow{h_1} & D_2 & \xrightarrow{h_2} & D_3. \end{array}$$

If the left-hand square is a pushout square, then the right-hand square is a pushout square if and only if the outer rectangle is a pushout square.

Proof. This follows immediately from Lemma 1.3.15. □

1.4 Commutative diagrams, composition, and animae

After studying constructions that build new ∞ -categories out of old ones, we now return to the internal geometry of a single ∞ -category. The first step is to enlarge the class of diagrams we can speak about and to formalize the common practice of displaying only the visible edges of such diagrams. This leads to the Segal axiom, which makes composition of morphisms precise. Once composition is in place, we can define isomorphisms and recover, in a formal way, the notion of anima introduced heuristically in Section 1.1.

1.4.1 Commutative diagrams

Besides commutative triangles, several other simple diagram shapes occur constantly in practice: commutative squares, spans, cospans, and infinite strings of composable morphisms. The next axiom formalizes the idea that for these diagram shapes, the higher data is determined by the lower-dimensional data we choose to display.

Definition 1.4.1 (Diagram shapes). A *commutative square* in an ∞ -category C is a functor $[1] \times [1] \rightarrow C$. We will often display such a square as

$$\begin{array}{ccc} x & \xrightarrow{f} & y \\ g \downarrow & & \downarrow h \\ z & \xrightarrow{k} & w. \end{array}$$

Throughout, we read the first coordinate of $[1] \times [1]$ as the row and the second as the column, so that the four objects $(0, 0)$, $(0, 1)$, $(1, 0)$ and $(1, 1)$ are displayed as x , y , z and w , respectively. In particular $\{0\} \times [1]$ and $\{1\} \times [1]$ are the top and bottom rows, while $[1] \times \{0\}$ and $[1] \times \{1\}$ are the left and right columns.

Let $\sqsubset \subseteq [1] \times [1]$ be the subposet spanned by the three objects $(0, 0)$, $(0, 1)$ and $(1, 0)$, and let $\sqsupset \subseteq [1] \times [1]$ be the subposet spanned by the three objects $(1, 0)$, $(0, 1)$ and $(1, 1)$. A *span* in C is a functor $\sqsubset \rightarrow C$, and a *cospan* in C is a functor $\sqsupset \rightarrow C$. We display them as

$$\begin{array}{ccc} x & \xrightarrow{f} & y \\ g \downarrow & & \\ z. & & \end{array} \quad \text{and} \quad \begin{array}{ccc} & & y \\ & & \downarrow h \\ z & \xrightarrow{k} & w. \end{array}$$

Every commutative square gives rise to both a span and a cospan via restriction along the two inclusions $\sqsubset \hookrightarrow [1] \times [1]$ and $\sqsupset \hookrightarrow [1] \times [1]$.

Finally, let $[\omega] := \{0 \leq 1 \leq 2 \leq 3 \leq \dots\}$ be the poset of natural numbers. An *infinite sequence of morphisms in C* is a functor $x_\bullet: [\omega] \rightarrow C$, which we display as

$$x_0 \xrightarrow{f_0} x_1 \xrightarrow{f_1} x_2 \xrightarrow{f_2} \dots,$$

where x_n is the evaluation of $x_\bullet$ at $n \in [\omega]$ and f_n is obtained from the inclusion $[1] \cong \{n \leq n+1\} \hookrightarrow [\omega]$.

These pictures record only part of the underlying functorial data: they display the visible edges, while suppressing composites and higher cells. We axiomatize that this omitted data is recoverable.

Axiom D (Commutative diagram axiom). The following commutative squares are pushouts of ∞ -categories:

$$\begin{array}{ccc}
[0] & \xrightarrow{0} & [1] \\
n-1 \downarrow & & \downarrow n-1 \rightarrow n \\
[n-1] & \hookrightarrow & [n],
\end{array}
\qquad
\begin{array}{ccc}
[1] & \xrightarrow{d_1} & [2] \\
d_1 \downarrow & & \downarrow (s_1, s_0) \\
[2] & \xrightarrow{(s_0, s_1)} & [1] \times [1],
\end{array}$$

$$\begin{array}{ccc}
[0] & \xrightarrow{d_1} & [1] \\
d_1 \downarrow & & \downarrow (0,0) \rightarrow (1,0) \\
[1] & \xrightarrow{(0,0) \rightarrow (0,1)} & \lrcorner,
\end{array}
\qquad
\begin{array}{ccc}
[0] & \xrightarrow{d_0} & [1] \\
d_0 \downarrow & & \downarrow (1,0) \rightarrow (1,1) \\
[1] & \xrightarrow{(0,1) \rightarrow (1,1)} & \lrcorner,
\end{array}$$

$$\begin{array}{ccc}
\sqcup_{l \in \mathbb{N}} \{l+1\} & \longrightarrow & \sqcup_{k \in \mathbb{N}} \{2k \leq 2k+1\} \\
\downarrow & & \downarrow \\
\sqcup_{k \in \mathbb{N}} \{2k+1 \leq 2k+2\} & \longrightarrow & [\omega].
\end{array}$$

Here the first square is required to be a pushout for every $n \geq 1$. In the last square, the right and bottom maps are the inclusions, while the top and left maps send the vertex $\{l+1\}$ to the corresponding vertex with the same label.

Exercise 1.4.2. Unwind the definitions, and show that these five pushout conditions express that finite and infinite strings of composable morphisms, commutative squares, spans, and cospans are determined by the lower-dimensional data displayed above.

We will use two maps that turn commutative triangles into commutative squares with an identity edge.

Construction 1.4.3. There are monotone maps $p_0, p_2: [1] \times [1] \rightarrow [2]$, pictorially represented by

$$p_0 = \begin{array}{ccc} 0 & \longrightarrow & 1 \\ \parallel & \searrow & \downarrow \\ 0 & \longrightarrow & 2, \end{array}
\qquad
p_2 = \begin{array}{ccc} 0 & \longrightarrow & 2 \\ \downarrow & \searrow & \parallel \\ 1 & \longrightarrow & 2. \end{array}$$

Writing $j_0 := (s_0, s_1)$ and $j_2 := (s_1, s_0)$ for the two functors $[2] \rightarrow [1] \times [1]$ appearing in Axiom D, there are natural isomorphisms $p_0 \circ j_0 \cong \text{id}_{[2]}$ and $p_2 \circ j_2 \cong \text{id}_{[2]}$.

Precomposition with p_0 or p_2 displays a commutative triangle as a commutative square whose left or right column, respectively, is an identity. The following proposition makes this precise.

Proposition 1.4.4 (Triangles as squares with an identity edge). *For every ∞ -category C , the following two commutative squares are pullback squares:*

$$\begin{array}{ccc}
\text{Fun}([2], C) & \xrightarrow{p_0^*} & \text{Fun}([1] \times [1], C) \\
\text{ev}_0 \downarrow & & \downarrow \text{res} \\
C & \xrightarrow{(p_{[1]})^*} & \text{Fun}([1] \times \{0\}, C)
\end{array}
\qquad
\text{and}
\qquad
\begin{array}{ccc}
\text{Fun}([2], C) & \xrightarrow{p_2^*} & \text{Fun}([1] \times [1], C) \\
\text{ev}_2 \downarrow & & \downarrow \text{res} \\
C & \xrightarrow{(p_{[1]})^*} & \text{Fun}([1] \times \{1\}, C).
\end{array}$$

Here the functors labeled ‘res’ are given by precomposition with the inclusions of the left and right columns into $[1] \times [1]$.

Proof guide. Factor each displayed square into two squares. The commutative-square case and the case $n = 2$ of Axiom D identify the resulting squares as pullbacks, and Lemma 1.3.15 finishes the argument. Details are given in Solution 25.3.1.

1.4.2 Composition of morphisms

Given a commutative triangle $\sigma: [2] \rightarrow C$ in an ∞ -category C , we may extract two morphisms $f := \sigma \circ d_2$ and $g := \sigma \circ d_0$:

$$\sigma = \begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{\quad} & z. \end{array}$$

By the case $n = 2$ of Axiom D, the pair (f, g) completely determines the commutative triangle. We refer to this special case as the *Segal Axiom*:

Proposition 1.4.5 (Segal Axiom). *For an ∞ -category C , the restriction functor*

$$(d_0^*, d_2^*): \text{Fun}([2], C) \rightarrow \text{Fun}([1], C) \times_{\text{ev}_0, C, \text{ev}_1} \text{Fun}([1], C), \quad \sigma \mapsto (g, f)$$

is an equivalence. □

In particular, for morphisms $f: x \rightarrow y$ and $g: y \rightarrow z$ we obtain a unique commutative triangle $\sigma: [2] \rightarrow C$ satisfying $d_2^*(\sigma) \cong f$ and $d_0^*(\sigma) \cong g$. The resulting morphism $g \circ f := d_1^*(\sigma): [1] \rightarrow C$ is called the *composite of f and g* .

Definition 1.4.6 (Arrow category). Let C be an ∞ -category. We write

$$\text{Ar}(C) := \text{Fun}([1], C)$$

and refer to it as the *arrow category* of C . Its objects are precisely the morphisms of C . Restriction along the two maps $[0] \rightarrow [1]$ defines the *source* and *target* functors

$$s := d_1^* = \text{ev}_0: \text{Ar}(C) \rightarrow C, \quad t := d_0^* = \text{ev}_1: \text{Ar}(C) \rightarrow C.$$

The assignment $(g, f) \mapsto g \circ f$ is functorial:

Construction 1.4.7 (Composition functor). Consider the zig-zag

$$\text{Ar}(C) \times_{s, C, t} \text{Ar}(C) \xleftarrow[\sim]{(d_0^*, d_2^*)} \text{Fun}([2], C) \xrightarrow{d_1^*} \text{Ar}(C).$$

The first map is an equivalence, and hence it admits an inverse, providing a composite functor

$$- \circ -: \text{Ar}(C) \times_{s, C, t} \text{Ar}(C) \xrightarrow{\sim} \text{Fun}([2], C) \xrightarrow{d_1^*} \text{Ar}(C).$$

We refer to this as the *composition functor*.

We will now show that composition in an ∞ -category is unital and associative.

Lemma 1.4.8 (Unitality). *For every morphism $f: x \rightarrow y$ in an ∞ -category C , there are natural isomorphisms*

$$\mathrm{id}_y \circ f \cong f \cong f \circ \mathrm{id}_x$$

in $\mathrm{Ar}(C)$. Moreover, these isomorphisms are natural in f , in the sense that they form natural isomorphisms of functors $\mathrm{Ar}(C) \rightarrow \mathrm{Ar}(C)$.

Proof. For a fixed morphism f , the isomorphisms are an immediate consequence of Exercise 1.2.15, using the degenerate commutative triangles $s_1^*(f) := f \circ s_1$ and $s_0^*(f) := f \circ s_0$. For the naturality of the relation $f \circ \mathrm{id}_x \cong f$ we have to show that the composite

$$\mathrm{Ar}(C) \xrightarrow{\sim} \mathrm{Ar}(C) \times_{s,C,\mathrm{id}} C \xrightarrow{1 \times p_{[1]}^*} \mathrm{Ar}(C) \times_{s,C,t} \mathrm{Ar}(C) \xrightarrow{- \circ -} \mathrm{Ar}(C)$$

is isomorphic to the identity functor. This follows from the following commutative diagram:

$$\begin{array}{ccccc} \mathrm{Ar}(C) & \xrightarrow{s_0^*} & \mathrm{Fun}([2], C) & \xrightarrow{d_1^*} & \mathrm{Ar}(C) \\ \downarrow \simeq & & \downarrow \simeq & & \parallel \\ \mathrm{Ar}(C) \times_{s,C,\mathrm{id}} C & \xrightarrow{1 \times p_{[1]}^*} & \mathrm{Ar}(C) \times_{s,C,t} \mathrm{Ar}(C) & \xrightarrow{- \circ -} & \mathrm{Ar}(C). \end{array}$$

A similar discussion applies to the relation $f \cong \mathrm{id}_y \circ f$. □

Proposition 1.4.9 (Associativity). *For composable morphisms $f: x \rightarrow y$, $g: y \rightarrow z$ and $h: z \rightarrow w$ in an ∞ -category C , there is a natural isomorphism*

$$h \circ (g \circ f) \cong (h \circ g) \circ f$$

in $\mathrm{Ar}(C)$.

Proof. Throughout this proof we display a morphism in $\mathrm{Ar}(C)$ as a commutative square whose two *vertical* arrows are its source and its target, so that the horizontal direction records the morphism in $\mathrm{Ar}(C)$; the identification $\mathrm{Fun}([1], \mathrm{Ar}(C)) \simeq \mathrm{Fun}([1] \times [1], C)$ we use is the one that swaps the two factors of $[1] \times [1]$ relative to the convention of Definition 1.4.1. With this reading, consider the following two morphisms in $\mathrm{Ar}(C)$:

$$\begin{array}{ccc} x & \xrightarrow{f} & y \\ \mathrm{id}_x \downarrow & \searrow g \circ f & \downarrow g \\ x & \xrightarrow{g \circ f} & z \end{array} \quad \text{and} \quad \begin{array}{ccc} y & \xrightarrow{h \circ g} & w \\ g \downarrow & \searrow h \circ g & \downarrow \mathrm{id}_w \\ z & \xrightarrow{h} & w. \end{array}$$

These morphisms are natural in f , g and h : we have functors

$$\mathrm{Ar}(C) \times_{s,C,t} \mathrm{Ar}(C) \xrightarrow{\simeq} \mathrm{Fun}([2], C) \xrightarrow{p_0^*} \mathrm{Fun}([1] \times [1], C),$$

$$\begin{array}{ccccc}
x & \xrightarrow{f} & y & & \\
& & \downarrow g & \mapsto & \\
& & z & &
\end{array}
\quad \mapsto \quad
\begin{array}{ccc}
x & \xrightarrow{f} & y \\
& \searrow g \circ f & \downarrow g \\
& & z
\end{array}
\quad \mapsto \quad
\begin{array}{ccccc}
x & \xrightarrow{f} & y & & \\
& \searrow g \circ f & \downarrow g & & \\
\text{id}_x \downarrow & & x & \xrightarrow{g \circ f} & z,
\end{array}$$

and similarly for the other diagram. Forming the composite of these two morphisms in $\text{Ar}(C)$, we thus obtain a commutative square in C of the form

$$\begin{array}{ccc}
x & \xrightarrow{(h \circ g) \circ f} & w \\
\text{id}_x \downarrow & \searrow & \downarrow \text{id}_w \\
x & \xrightarrow{h \circ (g \circ f)} & w,
\end{array}$$

natural in f , g and h . Using Lemma 1.4.8, the diagonal of this square is then naturally isomorphic to both $(h \circ g) \circ f$ and $h \circ (g \circ f)$, finishing the proof. $\square$

1.4.3 Isomorphisms, anima and groupoid cores

With composition available, we can now speak about invertibility of morphisms.

Definition 1.4.10 (Isomorphisms). Consider a morphism $f: x \rightarrow y$ in an ∞ -category C . We say that f is *invertible*, or that it is an *isomorphism*, if there exists a morphism $f^{-1}: y \rightarrow x$ together with commutative triangles in C of the form

$$\begin{array}{ccc}
y & \xrightarrow{f^{-1}} & x \\
& \searrow & \downarrow f \\
& & y
\end{array}
\quad
\begin{array}{ccc}
x & \xrightarrow{f} & y \\
& \searrow & \downarrow f^{-1} \\
& & x.
\end{array}$$

We may now introduce the formal version of the notion of anima from Section 1.1.

Definition 1.4.11 (Anima). An ∞ -category C is called an *anima*, or an ∞ -*groupoid*, if every morphism in C is invertible.

Recall from Example 1.1.5 that every 1-category C has an anima of objects $C^\simeq$, called its *groupoid core*: the subgroupoid of C which contains all objects but only the invertible morphisms. The groupoid $C^\simeq$ is uniquely characterized by the property that the inclusion $C^\simeq \hookrightarrow C$ is terminal among functors from a groupoid into C . We use this universal property to axiomatize the ∞ -categorical version of the core:

Axiom E (Groupoid core axiom). For every ∞ -category C , there is an anima $C^\simeq$ called the (*groupoid*) *core* of C , which comes equipped with a functor $\gamma_C: C^\simeq \rightarrow C$. Every functor $F: X \rightarrow C$ from an anima X factors through γ_C , and for functors $G, H: X \rightarrow C^\simeq$ every natural isomorphism $\gamma_C \circ G \cong \gamma_C \circ H$ may be lifted to a natural isomorphism $G \cong H$.

Furthermore, the map $\langle 0, 1 \rangle: * \sqcup * \rightarrow [1]$ factors through an equivalence $* \sqcup * \xrightarrow{\sim} [1]^\simeq$.

Definition 1.4.12. For ∞ -categories C and D , we write $\text{Map}(C, D)$ for the anima $\text{Fun}(C, D)^\simeq$, and refer to it as the *mapping anima* between C and D .

Exercise 1.4.13. Show that for an anima X and an ∞ -category C the core functor $\gamma_C: C^\simeq \rightarrow C$ induces an equivalence

$$\mathrm{Map}(X, C^\simeq) \xrightarrow{\sim} \mathrm{Map}(X, C).$$

Hint: Apply the groupoid core axiom to the evaluation functor $\mathrm{Map}(X, C) \times X \rightarrow C$, and then curry and pass to the core once more.

Exercise 1.4.14. Show that the groupoid-core construction preserves pullbacks: for functors $C \rightarrow E \leftarrow D$, construct a preferred equivalence

$$(C \times_E D)^\simeq \simeq C^\simeq \times_{E^\simeq} D^\simeq.$$

Hint: Construct both directions from the groupoid-core and pullback universal properties, and compare the composites after projecting to the factors.

Exercise 1.4.15. Let $F: C \rightarrow D$ be a functor. Show that the following conditions are equivalent:

- (1) The functor F is an equivalence;
- (2) For every ∞ -category E the functor $F \circ -: \mathrm{Map}(E, C) \rightarrow \mathrm{Map}(E, D)$ is an equivalence;
- (3) For every ∞ -category E the functor $- \circ F: \mathrm{Map}(D, E) \rightarrow \mathrm{Map}(C, E)$ is an equivalence.

Exercise 1.4.16. Show that a morphism $f: X \rightarrow Y$ of anima is an equivalence if and only if the induced map $f^*: \mathrm{Map}(Y, Z) \rightarrow \mathrm{Map}(X, Z)$ is an equivalence for every anima Z .

1.4.4 Hom anima

In Section 1.1, we heuristically described the morphisms between two objects of an ∞ -category as forming an anima. We encode this object within our axiomatic setup by the following pullback.

Definition 1.4.17 (Hom anima). Let C be any ∞ -category. Given objects x and y of C , we define the *hom anima* $\mathrm{Hom}_C(x, y)$ via the following pullback square:

$$\begin{array}{ccc} \mathrm{Hom}_C(x, y) & \longrightarrow & \mathrm{Ar}(C) \\ \downarrow & \lrcorner & \downarrow (s, t) \\ * & \xrightarrow{(x, y)} & C \times C. \end{array}$$

Observe that the objects of $\mathrm{Hom}_C(x, y)$ are triples (f, α, β) , where $f: x' \rightarrow y'$ is a morphism in C and $\alpha: x \cong x'$ and $\beta: y \cong y'$ are isomorphisms in C . We will often abuse notation and pretend that $x = x'$ and $y = y'$.

Remark 1.4.18. At this point, the pullback only defines $\mathrm{Hom}_C(x, y)$ as an ∞ -category. The terminology *hom anima* will be justified after the Rezk axiom: by Proposition 1.4.25, $\mathrm{Hom}_C(x, y)$ is indeed an anima.

Exercise 1.4.19. Let x, y and z be objects of an ∞ -category C . Construct a composition functor

$$- \circ -: \mathrm{Hom}_C(y, z) \times \mathrm{Hom}_C(x, y) \rightarrow \mathrm{Hom}_C(x, z), \quad (g, f) \mapsto g \circ f.$$

Hint: Restrict the composition functor $\mathrm{Ar}(C) \times_{s, C, t} \mathrm{Ar}(C) \rightarrow \mathrm{Ar}(C)$ along the pullbacks defining the three hom anima.

Definition 1.4.20 (Natural transformations). If C and D are ∞ -categories, we define a *natural transformation* of functors $C \rightarrow D$ to be a morphism in $\text{Fun}(C, D)$. By (un)currying, this may equivalently be encoded as a functor $[1] \times C \rightarrow D$. We denote the hom anima in $\text{Fun}(C, D)$ by $\text{Nat}(f, g)$:

$$\text{Nat}(f, g) := \text{Hom}_{\text{Fun}(C, D)}(f, g).$$

1.4.5 The Rezk axiom

In a 1-category, every invertible morphism is isomorphic in the arrow category to an identity morphism. Thus, if $\text{Iso}(C) \subseteq \text{Ar}(C)$ denotes the full subcategory spanned by the invertible morphisms, then the functor

$$C \rightarrow \text{Iso}(C), \quad x \mapsto \text{id}_x,$$

is an equivalence. The Rezk axiom is the ∞ -categorical analogue of this statement. Since we do not yet have subcategories at our disposal, we first construct $\text{Iso}(C)$ indirectly:

Definition 1.4.21. Given an ∞ -category C , we define the ∞ -category $\text{Iso}(C)$ via the following pullback diagram:

$$\begin{array}{ccc} \text{Iso}(C) & \longrightarrow & \text{Ar}(C) \times_{t, C, s} \text{Ar}(C) \times_{t, C, s} \text{Ar}(C) \\ \downarrow & \lrcorner & \downarrow (g, f, h) \mapsto (f \circ g, h \circ f) \\ C \times C & \xrightarrow{(x, y) \mapsto (\text{id}_y, \text{id}_x)} & \text{Ar}(C) \times \text{Ar}(C). \end{array}$$

We define $\pi_{\text{Iso}}: \text{Iso}(C) \rightarrow \text{Ar}(C)$ to be the composite functor

$$\text{Iso}(C) \rightarrow \text{Ar}(C) \times_{t, C, s} \text{Ar}(C) \times_{t, C, s} \text{Ar}(C) \xrightarrow{\text{pr}_2} \text{Ar}(C).$$

An object of $\text{Iso}(C)$ is a triple of morphisms $(g: y' \rightarrow x, f: x \rightarrow y, h: y \rightarrow x')$ together with isomorphisms $f \circ g \cong \text{id}_y$ and $h \circ f \cong \text{id}_x$. Its image under π_{Iso} is the morphism f .

Exercise 1.4.22. Give a precise construction of the right vertical functor in the pullback defining $\text{Iso}(C)$. *Hint:* Project to the composable pairs (f, g) and (h, f) , apply the composition functor to each pair, and then take their product.

Remark 1.4.23. Note that $\text{Iso}(C)$ encodes separate left and right inverses g and h of f , rather than a single two-sided inverse f^{-1} . The reason for this is to guarantee that the fibers of $\pi_{\text{Iso}}: \text{Iso}(C) \rightarrow \text{Ar}(C)$ are either empty or contractible, so that being an isomorphism is really a *property* of a morphism rather than additional structure. This is made precise by Proposition 1.5.6.

As one expects, the identity map $\text{id}_x: x \rightarrow x$ is invertible in C for every object x of C :

Construction 1.4.24. The identity morphisms define a canonical functor

$$i: C \rightarrow \text{Iso}(C).$$

Indeed, restriction along $p_{[1]}: [1] \rightarrow [0]$ gives a functor

$$p_{[1]}^*: C \rightarrow \text{Ar}(C), \quad x \mapsto \text{id}_x,$$

whose composites with the source and target functors $s, t: \text{Ar}(C) \rightarrow C$ are both isomorphic to id_C . The triple $(p_{[1]}^*, p_{[1]}^*, p_{[1]}^*)$ therefore lifts to a functor

$$C \rightarrow \text{Ar}(C) \times_{t, C, s} \text{Ar}(C) \times_{t, C, s} \text{Ar}(C), \quad x \mapsto (\text{id}_x, \text{id}_x, \text{id}_x).$$

By Lemma 1.4.8 there is a natural isomorphism $\text{id}_x \circ \text{id}_x \cong \text{id}_x$, so the composite of this functor with $(g, f, h) \mapsto (f \circ g, h \circ f)$ is naturally isomorphic to the composite of the diagonal $\Delta: C \rightarrow C \times C$ with $(x, y) \mapsto (\text{id}_y, \text{id}_x)$. The universal property of the pullback defining $\text{Iso}(C)$ thus provides the desired functor i , and it satisfies $\pi_{\text{Iso}} \circ i \cong p_{[1]}^*$.

Axiom F (Rezk axiom). For every ∞ -category C the functor $i: C \rightarrow \text{Iso}(C)$ is an equivalence.

The Rezk axiom now implies that the hom animae constructed above really are animae.

Proposition 1.4.25 (Hom animae are animae). *Let C be an ∞ -category and let x and y be objects in C . Then the hom anima $\text{Hom}_C(x, y)$ is indeed an anima.*

A proof is given in Solution 25.5.1.

Remark 1.4.26. The condition that the functor $i: C \rightarrow \text{Iso}(C)$ is an equivalence is known as the *completeness condition*. It was used by Rezk [Rez01] to provide a model of ∞ -categories, called *complete Segal spaces*. A detailed discussion of the equivalence between ∞ -categories and complete Segal animae will be given in Chapter 24.

The Rezk axiom implies that two objects x and y in C may be identified with each other as soon as we find an invertible morphism $f: x \rightarrow y$ between them:

Corollary 1.4.27. *Let x and y be objects of C and assume there exists an invertible morphism $f: x \rightarrow y$. Then there exists a natural isomorphism $x \cong y$ as functors $* \rightarrow C$.*

Proof. The inverse and the two triangles for f determine an object $\tilde{f}$ of $\text{Iso}(C)$ whose image under π_{Iso} is f . Since the functor $i: C \rightarrow \text{Iso}(C)$ is an equivalence, there exists an object z of C together with an isomorphism $i(z) \cong \tilde{f}$ in $\text{Iso}(C)$. Applying the source and target functors $s, t: \text{Ar}(C) \rightarrow C$ to the image of this isomorphism under π_{Iso} produces isomorphisms $z \cong x$ and $z \cong y$ in C . By inversion we obtain $x \cong z$ and by composition we obtain $x \cong y$, as desired. $\square$

The previous corollary is the object-level shadow of a more general fact. For functors $f, g: C \rightarrow D$, we now have two a priori different ways to say that f and g are the same: on the one hand, we have the primitive notion of a natural isomorphism $\alpha: f \cong g$ from Axiom A.1; on the other hand, we may consider an invertible natural transformation $\alpha: f \xrightarrow{\sim} g$, that is, an invertible morphism in the functor category $\text{Fun}(C, D)$. The Rezk axiom shows that these are equivalent notions.

Corollary 1.4.28. *Let $f, g: C \rightarrow D$ be functors. Then the ∞ -category $(f \cong g)$ encoding primitive natural isomorphisms is equivalent to the fiber over (f, g) of*

$$(s, t): \text{Iso}(\text{Fun}(C, D)) \rightarrow \text{Fun}(C, D) \times \text{Fun}(C, D).$$

In particular, primitive natural isomorphisms $f \cong g$ correspond to invertible natural transformations $f \xrightarrow{\sim} g$.

Proof. Recall from Exercise 1.3.23 that there is an ∞ -category $(f \cong g)$ encoding the primitive natural isomorphisms between f and g , defined as the fiber over (f, g) of the diagonal functor

$$\mathrm{Fun}(C, D) \rightarrow \mathrm{Fun}(C, D) \times \mathrm{Fun}(C, D).$$

Now note that this diagonal functor factors as

$$\mathrm{Fun}(C, D) \xrightarrow{i} \mathrm{Iso}(\mathrm{Fun}(C, D)) \xrightarrow{(s, t)} \mathrm{Fun}(C, D) \times \mathrm{Fun}(C, D).$$

Since the first functor is an equivalence by the Rezk axiom, the fiber $(f \cong g)$ may equivalently be computed as the fiber of the functor (s, t) over (f, g) : there is a pullback square

$$\begin{array}{ccc} (f \cong g) & \xrightarrow{\quad} & \mathrm{Iso}(\mathrm{Fun}(C, D)) \\ \downarrow & \lrcorner & \downarrow (s, t) \\ * & \xrightarrow{(f, g)} & \mathrm{Fun}(C, D) \times \mathrm{Fun}(C, D). \end{array}$$

An object of this fiber consists of a natural transformation $f \rightarrow g$ together with left and right inverse data. Hence it encodes an invertible natural transformation $f \xrightarrow{\sim} g$, as claimed. $\square$

1.5 Further constructions of ∞ -categories

We will now introduce further constructions of ∞ -categories: opposite categories, subcategories and localizations.

1.5.1 Opposite categories

Axiom G. For every ∞ -category C there is an *opposite category* C^{op} . Similarly every functor $F: C \rightarrow D$ induces an opposite functor $F^{\mathrm{op}}: C^{\mathrm{op}} \rightarrow D^{\mathrm{op}}$, and this process respects composition of functors. We have equivalences $(C^{\mathrm{op}})^{\mathrm{op}} \simeq C$ and $(F^{\mathrm{op}})^{\mathrm{op}} \simeq F$.

If C and D come from classical 1-categories then C^{op} and F^{op} agree with the usual constructions of opposite categories.

If X is an anima then we have an equivalence $X^{\mathrm{op}} \simeq X$.

Note that functors $[1] \rightarrow C^{\mathrm{op}}$ correspond to functors $[1] \simeq [1]^{\mathrm{op}} \rightarrow (C^{\mathrm{op}})^{\mathrm{op}} \simeq C$, so that morphisms in C^{op} are just morphisms in C but with source and target flipped. Also since $[2] \simeq [2]^{\mathrm{op}}$, this correspondence between morphisms in C and morphisms in C^{op} preserves composition. In particular C is an anima if and only if C^{op} is an anima.

Proposition 1.5.1. *The construction $C \mapsto C^{\mathrm{op}}$ is compatible with all constructions of ∞ -categories:*

- (1) *There are equivalences $*^{\mathrm{op}} \simeq *$ and $\emptyset^{\mathrm{op}} \simeq \emptyset$.*
- (2) *For ∞ -categories C and D there are equivalences*

$$\begin{aligned} (C \times D)^{\mathrm{op}} &\simeq C^{\mathrm{op}} \times D^{\mathrm{op}}, \\ C^{\mathrm{op}} \sqcup D^{\mathrm{op}} &\simeq (C \sqcup D)^{\mathrm{op}}, \\ \mathrm{Fun}(C^{\mathrm{op}}, D^{\mathrm{op}}) &\simeq \mathrm{Fun}(C, D)^{\mathrm{op}}. \end{aligned}$$

(3) For functors $F: C \rightarrow E$ and $G: D \rightarrow E$ there is an equivalence

$$(C \times_E D)^{\text{op}} \simeq C^{\text{op}} \times_{E^{\text{op}}} D^{\text{op}}.$$

(4) For an ∞ -category C there is an equivalence $(C^{\text{op}})^{\simeq} \simeq (C^{\simeq})^{\text{op}}$.

Proof. All of these properties follow readily from the universal properties of these constructions. For illustration, let us merely sketch the proof that $(-)^{\text{op}}$ preserves products: functors $T \rightarrow (C \times D)^{\text{op}}$ correspond to functors $T^{\text{op}} \rightarrow C \times D$, which in turn are pairs of functors $(T^{\text{op}} \rightarrow C, T^{\text{op}} \rightarrow D)$, which in turn translate back into pairs of functors $(T \rightarrow C^{\text{op}}, T \rightarrow D^{\text{op}})$, i.e. functors $T \rightarrow C^{\text{op}} \times D^{\text{op}}$. $\square$

Exercise 1.5.2. Let x and y be objects of an ∞ -category C . Show that there is an equivalence

$$\text{Hom}_{C^{\text{op}}}(x, y) \simeq \text{Hom}_C(y, x).$$

Hint: apply $(-)^{\text{op}}$ to the pullback square defining $\text{Hom}_C(y, x)$ and use that $\text{Hom}_C(y, x)$ is an anima.

1.5.2 Subcategories

Next, we would like to define the notion of *subcategories*. Just like in classical category theory, a subcategory of an ∞ -category C is determined by choosing some objects of C and some morphisms between them, subject to closure under identities and composition. However, while the subcategory can be constructed ‘by hand’ in the classical situation, we must now make sure that all the higher coherence data from C is inherited by the subcategory. In order to formulate this, we start by isolating the homotopical analogue of a subcategory inclusion.

Definition 1.5.3. A functor $F: C \rightarrow D$ is called a *monomorphism* if the commutative square

$$\begin{array}{ccc} C & \xlongequal{\quad} & C \\ \parallel & & \downarrow F \\ C & \xrightarrow{F} & D \end{array}$$

is a pullback square, i.e. the functor $\Delta_F := (\text{id}_C, \text{id}_C): C \rightarrow C \times_D C$ is an equivalence of ∞ -categories.

Remark 1.5.4. Note that this means that two functors $G, H: T \rightarrow C$ are naturally isomorphic whenever the functors $F \circ G$ and $F \circ H$ are naturally isomorphic, reflecting the usual categorical notion of monomorphism.

We may sometimes write $F: C \hookrightarrow D$ if F is a monomorphism. If y is an object of D , we say that y is *contained in* C , sometimes written as $y \in C$, if there exists some object x of C satisfying $F(x) \cong y$.

Exercise 1.5.5. Let $F: C \hookrightarrow D$ be a monomorphism, and let y be an object of D which is contained in C . Show that the fiber C_y of F at y is contractible.

Hint: Show that any functor $x: * \rightarrow C_y$ is an equivalence by exhibiting it as a pullback of the equivalence $\Delta_F: C \rightarrow C \times_D C$.

The projection from the ∞ -category of isomorphisms introduced in Definition 1.4.21 is the principal example needed below.

Proposition 1.5.6 (Isomorphisms form a subcategory of arrows). *For every ∞ -category C , the functor*

$$\pi_{\text{Iso}}: \text{Iso}(C) \rightarrow \text{Ar}(C)$$

is a monomorphism.

A proof is given in Solution 25.4.1.

We postpone the remaining routine properties of monomorphisms to the exercise section at the end of the chapter.

Axiom H.1 (Monomorphisms define subcategories). Let $i: A \rightarrow C$ be a monomorphism of ∞ -categories. Then for every ∞ -category T , the commutative square

$$\begin{array}{ccc} \text{Map}(T, A) & \xrightarrow{\quad\quad\quad} & \text{Map}(T, C) \\ \downarrow & & \downarrow \\ \text{Map}(\text{Map}([1], T), \text{Map}([1], A)) & \hookrightarrow & \text{Map}(\text{Map}([1], T), \text{Map}([1], C)) \end{array}$$

is a pullback square. The vertical functors record the action of a functor on morphisms.

Exercise 1.5.7. Construct the two vertical functors and the natural isomorphism that makes the square in Axiom H.1 commute. *Hint:* For $D = A, C$, restrict the composition functor $\text{Fun}(T, D) \times \text{Fun}([1], T) \rightarrow \text{Fun}([1], D)$ to the relevant cores, factor through the appropriate groupoid cores, and curry.

Informally, the axiom says that in order for a functor $F: T \rightarrow C$ to factor through A , it suffices to check that F maps morphisms of T to morphisms of C that lie in A . Accordingly, we will use the terms *monomorphism* and *subcategory inclusion* interchangeably.

We will now introduce a method for constructing subcategories of C by specifying a collection of morphisms closed under composition.

Definition 1.5.8. Let C be an ∞ -category. A *collection of morphisms in C* is a monomorphism $M \hookrightarrow \text{Map}([1], C)$. We say it is *closed under composition* if the following two conditions are satisfied:

- For a morphism $f: x \rightarrow y$ in C , if $f \in M$, then also $\text{id}_x \in M$ and $\text{id}_y \in M$;
- For morphisms $f: x \rightarrow y$ and $g: y \rightarrow z$ in C , if $g, f \in M$, then also $g \circ f \in M$.

Axiom H.2 (Subcategory axiom). Consider an ∞ -category C equipped with a collection of morphisms $m: M \hookrightarrow \text{Map}([1], C)$ of C which is closed under composition. Then there exists a monomorphism $i_M: \langle M \rangle_C \rightarrow C$ such that the induced map on morphisms factors through an equivalence $\text{Map}([1], \langle M \rangle_C) \xrightarrow{\sim} M$.

By Axiom H.1, the monomorphism i_M has the expected universal property of a subcategory. We refer to the ∞ -category $\langle M \rangle_C$ as the *(non-full) subcategory spanned by the morphisms in M* .

Definition 1.5.9 (Full subcategory). A functor $i: A \rightarrow C$ is called a *full subcategory inclusion* if $i^\simeq: A^\simeq \rightarrow C^\simeq$ is a monomorphism, and for every ∞ -category T the commutative square

$$\begin{array}{ccc} \mathrm{Map}(T, A) & \longrightarrow & \mathrm{Map}(T, C) \\ \downarrow & & \downarrow \\ \mathrm{Map}(T^\simeq, A^\simeq) & \hookrightarrow & \mathrm{Map}(T^\simeq, C^\simeq) \end{array}$$

is a pullback square. The vertical functors are constructed by restricting evaluation to $\mathrm{Map}(T, A) \times T^\simeq$ and $\mathrm{Map}(T, C) \times T^\simeq$, factoring through the corresponding groupoid cores, and currying.

Definition 1.5.10. A *collection of objects* of an ∞ -category C is a monomorphism $\Gamma \hookrightarrow C^\simeq$. The endpoint functors on $\mathrm{Ar}(C)$ restrict along the groupoid cores to functors $s, t: \mathrm{Map}([1], C) \rightarrow C^\simeq$, for which we use the same notation. Given a collection of objects Γ , we define M_Γ as the following pullback:

$$\begin{array}{ccc} M_\Gamma & \hookrightarrow & \mathrm{Map}([1], C) \\ \downarrow & \lrcorner & \downarrow (s, t) \\ \Gamma \times \Gamma & \hookrightarrow & C^\simeq \times C^\simeq. \end{array}$$

The collection M_Γ is closed under composition, since identities and composites have endpoints in Γ whenever the original morphisms do. Hence Axiom H.2 determines a monomorphism $\langle M_\Gamma \rangle_C \hookrightarrow C$. We refer to the subcategory $\langle M_\Gamma \rangle_C$ of C as the *full subcategory spanned by the objects in Γ* .

Proposition 1.5.11 (Full subcategories from collections of objects). *The functor $\langle M_\Gamma \rangle_C \hookrightarrow C$ is a full subcategory inclusion, and there is an equivalence*

$$\langle M_\Gamma \rangle_C^\simeq \simeq \Gamma$$

over $C^\simeq$.

Proof guide. Combine the pullback in Axiom H.1, first for $T = *$ and then for general T , with the pullback defining M_Γ . For $T = *$, pullback pasting identifies the core of $\langle M_\Gamma \rangle_C$ with $\Gamma \times_{C^\simeq} \Gamma$, which is equivalent to Γ because $\Gamma \rightarrow C^\simeq$ is a monomorphism. For general T , the same comparison says that a functor $T \rightarrow C$ factors through $\langle M_\Gamma \rangle_C$ precisely when its restriction to $T^\simeq$ factors through Γ . Details are given in Solution 25.4.3.

Recognizing equivalences

Recall that a functor between classical 1-categories is an equivalence if and only if it is both fully faithful and essentially surjective. The same statement is true for ∞ -categories. Since this is such a fundamental result, we will take it as one of our axioms.

Definition 1.5.12. A functor $F: C \rightarrow D$ is called *fully faithful* if for every pair of objects x and y of C , the induced map of animae

$$F: \mathrm{Hom}_C(x, y) \rightarrow \mathrm{Hom}_D(Fx, Fy)$$

is an equivalence.

Definition 1.5.13. A functor $F: C \rightarrow D$ is called *essentially surjective* if for every object y of D there exists an object x of C together with an isomorphism $Fx \cong y$.

Axiom I (Joyal). A functor $F: C \rightarrow D$ is an equivalence if and only if it is fully faithful and essentially surjective.

Exercise 1.5.14. Show that a functor $F: C \rightarrow D$ is an equivalence if and only if the induced functor

$$F_*: \text{Map}([1], C) \rightarrow \text{Map}([1], D)$$

is an equivalence.

Exercise 1.5.15. Show that a functor $F: C \rightarrow D$ is fully faithful if and only if it is a full subcategory inclusion (Definition 1.5.9).

Another statement familiar from classical category theory is that natural isomorphisms can be checked pointwise. We will also assume this result as a black-box.

Axiom J (Joyal). Let $F, G: C \rightarrow D$ be two functors and let $\alpha: F \Rightarrow G$ be a natural transformation. Then α is a natural isomorphism if and only if for every object x of C the morphism $\alpha(x): F(x) \rightarrow G(x)$ is an isomorphism in D .

Remark 1.5.16. It is possible to formulate axioms for ∞ -categories that would allow one to deduce Axioms I and J.

1.5.3 Localizations

Another important ∞ -categorical construction is that of *localizations*, where we formally invert a collection of morphisms in an ∞ -category.

Definition 1.5.17. Let C be an ∞ -category, and let $W \hookrightarrow \text{Map}([1], C)$ be a collection of morphisms in C . We say that a functor $F: C \rightarrow D$ *inverts the morphisms in W* if for every morphism $f: x \rightarrow y$ in C contained in $W \subseteq \text{Map}([1], C)$, the image $F(f): F(x) \rightarrow F(y)$ is invertible in D . We let $\text{Fun}^W(C, D)$ be the full subcategory of $\text{Fun}(C, D)$ spanned by those functors that invert the morphisms in W .

Exercise 1.5.18. Make the definition of $\text{Fun}^W(C, D)$ precise by constructing the corresponding collection of objects of $\text{Fun}(C, D)$. *Hint:* Pull back the collection of isomorphisms in D along the functor which records the action of a functor $C \rightarrow D$ on the morphisms in W .

Definition 1.5.19 (Localization). Let C be an ∞ -category, and let $W \hookrightarrow \text{Map}([1], C)$ be a collection of morphisms in C . We say that a functor $l: C \rightarrow L$ is a *localization of C at W* if l inverts the morphisms in W , and if for every ∞ -category D , the functor

$$l^*: \text{Fun}(L, D) \rightarrow \text{Fun}(C, D)$$

is a full subcategory inclusion which identifies $\text{Fun}(L, D)$ with the full subcategory $\text{Fun}^W(C, D)$.

Note that the universal property of localizations uniquely determines the localization. We will generally denote such a localization by $C[W^{-1}]$.

Axiom K (Existence of localizations). Let C be an ∞ -category, and let $W \hookrightarrow \text{Map}([1], C)$ be a collection of morphisms in C . Then there exists a localization $l: C \rightarrow C[W^{-1}]$ of C at W .

Remark 1.5.20 (See Exercise 1.13). One can show that the functor $p_{[1]}: [1] \rightarrow *$ exhibits $*$ as the localization of $[1]$ at the collection $W = \text{Map}([1], [1])$ of all its morphisms.

Exercise 1.5.21. Let $l: C \rightarrow C[W^{-1}]$ be a localization of C at a collection of morphisms $W \hookrightarrow \text{Map}([1], C)$. Show that l can be computed by the following pushout square of ∞ -categories:

$$\begin{array}{ccccc} [1] \times W & \hookrightarrow & [1] \times \text{Ar}(C) & \xrightarrow{\text{ev}} & C \\ \text{pr}_2 \downarrow & & & & \downarrow l \\ W & \xrightarrow{\quad \quad \quad} & & \xrightarrow{\quad \quad \quad} & C[W^{-1}]. \end{array}$$

Hint: Show that if i is a full subcategory inclusion, then so is $\text{Fun}(W, i)$, and then use Exercises 1.10 and 1.13.

Definition 1.5.22. We define the *geometric realization* $\|C\|$ of an ∞ -category C as the localization $C[W^{-1}]$, where $W := \text{Map}([1], C)$ is the collection of *all* morphisms in C .

Exercise 1.5.23 (See Exercise 1.20 for hints). Show that $\|C\|$ is an anima.

Remark 1.5.24. The terminology ‘geometric realization’ will be justified in Lemma 24.2.5, where we will show that $\|C\|$ is equivalent to the geometric realization (i.e. colimit) of the nerve $N(C): \Delta^{\text{op}} \rightarrow \text{An}$ of C .

Exercise 1.5.25. Let C and D be ∞ -categories with collections of morphisms W and W' , respectively, and assume that both collections contain all identity morphisms. Use the universal property to show that

$$(C \times D)[(W \times W')^{-1}] \xrightarrow{\sim} C[W^{-1}] \times D[W'^{-1}].$$

Deduce that $\|C \times D\| \xrightarrow{\sim} \|C\| \times \|D\|$.

1.6 Simplicial objects and quasicategories

So far, we have been treating ∞ -category theory very axiomatically, focusing on the essential features of the theory itself and ignoring how precisely one might model these objects within traditional set-based mathematics. In this section, we briefly discuss the most common traditional model for ∞ -categories, namely *quasicategories*. Along the way, we will also introduce some simplicial constructions and notation that will be used later independently of quasicategories themselves. Readers who are only interested in the axiomatic development may therefore treat the specifically quasicategorical parts of this section as optional.

Before we start, some words on history are in order. Quasicategories have played a pivotal role in the acceptance of ∞ -category theory as a legitimate part of mathematics. The conceptual idea of ‘doing category theory up to homotopy’ was around for a while, but a major difficulty was how to make this idea rigorous within ordinary mathematics. While the definition of a quasicategory already appeared in the work of Boardman and Vogt [BV73] under the name ‘weak Kan complexes’, it was through deep insights and important foundational work by André Joyal [Joy08] and subsequently

Jacob Lurie [Lur09b; Lur17] that the theory obtained a rigorous, well-developed and practical form that could immediately be used by researchers.

We begin with some simplicial preliminaries that will be useful later in any case.

Definition 1.6.1. The *simplex category* Δ is the 1-category of non-empty finite linearly ordered sets and order-preserving maps between them.

Observe that any object in Δ is isomorphic to the linear order $[n] = \{0 \leq 1 \leq \dots \leq n\}$ for some n , and that the isomorphism is unique if it exists. We will therefore frequently denote generic objects of Δ simply by $[n]$.

Observation 1.6.2. Every morphism in Δ can be written as a finite composite of face maps $d_i: [n-1] \rightarrow [n]$ and degeneracy maps $s_i: [n+1] \rightarrow [n]$ (see Notation 1.2.10). The relations these maps satisfy are known as the *simplicial identities*.

Definition 1.6.3. Let C be an ∞ -category. A *simplicial object in C* is defined to be a functor $\Delta^{\text{op}} \rightarrow C$. We will denote the ∞ -category of simplicial objects by

$$\text{s}C := \text{Fun}(\Delta^{\text{op}}, C).$$

In light of Observation 1.6.2, we may informally think of a simplicial object in C as a collection of objects X_n of C for every natural number n together with face maps $d_i: X_n \rightarrow X_{n-1}$ and degeneracy maps $s_i: X_n \rightarrow X_{n+1}$ satisfying the simplicial identities. We frequently draw simplicial objects as follows:

$$\dots \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} X_2 \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} X_1 \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} X_0.$$

Since this looks somewhat chaotic, it is common to leave the degeneracy maps implicit and only draw the face maps:

$$\dots \begin{array}{c} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} X_2 \begin{array}{c} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} X_1 \begin{array}{c} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} X_0.$$

We will now specialize to $C = \text{Set}$. For a simplicial set $X \in \text{sSet}$, we refer to the set X_n as its *set of n -simplices*.

Definition 1.6.4 (Simplices). For every natural number n , we define the *n -simplex* Δ^n as the simplicial set represented by $[n]$:

$$\Delta^n := \text{Hom}_{\Delta}(-, [n]): \Delta^{\text{op}} \rightarrow \text{Set}.$$

By the Yoneda lemma this defines a fully faithful functor

$$\Delta^{\bullet}: \Delta \hookrightarrow \text{sSet}.$$


Again by the Yoneda lemma there is a natural bijection $X_n \cong \text{Hom}_{\text{sSet}}(\Delta^n, X)$, so that we may think of an n -simplex in X as a morphism $\sigma: \Delta^n \rightarrow X$ of simplicial sets.

Definition 1.6.5 (Horns). For $n \geq 1$ and $0 \leq k \leq n$, we define the *k -horn* Λ_k^n as the subsimplicial set

$$\Lambda_k^n \subseteq \Delta^n$$

whose m -simplices are those maps $\varphi: [m] \rightarrow [n]$ in Δ such that there is some element $i \in [n] \setminus \{k\}$ which is not in the image of φ . Equivalently, it is the union inside Δ^n of all of its faces except for the k -th face.

Example 1.6.6. For $n = 2$ the three horns Λ_0^2 , Λ_1^2 and Λ_2^2 may be displayed as follows:



Definition 1.6.7. Let X be a simplicial set.

- (1) We say that X is a *Kan complex* if for every $n \geq 1$ and $0 \leq k \leq n$, every morphism of simplicial sets $\Lambda_k^n \rightarrow X$ extends to a morphism $\Delta^n \rightarrow X$:

$$\begin{array}{ccc} \Lambda_k^n & \xrightarrow{\quad} & X \\ \downarrow & \nearrow \text{dashed} & \\ \Delta^n & & \end{array}$$

- (2) We say that X is a *quasicategory* if the above condition holds for all $n \geq 1$ but only for $0 < k < n$.

We denote by

$$\mathbf{Kan} \subseteq \mathbf{qCat} \subseteq \mathbf{sSet}$$

the full subcategories on the Kan complexes and the quasicategories, respectively.

We now discuss three basic sources of quasicategories, of which the first two will be most relevant for us:

Example 1.6.8 (Singular simplicial complex). Let X be a topological space. We will construct a simplicial set $\text{Sing}(X) \in \mathbf{sSet}$ called the *singular simplicial complex of X* .

- For every $n \geq 0$, let $|\Delta^n|$ denote the topological n -simplex:

$$|\Delta^n| := \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0 + \dots + x_n = 1, x_i \geq 0 \text{ for } i = 0, \dots, n\}.$$

- For every order-preserving map $\varphi: [n] \rightarrow [m]$ we get a continuous map

$$|\Delta^\varphi|: |\Delta^n| \rightarrow |\Delta^m|, \quad (x_0, \dots, x_n) \mapsto (y_0, \dots, y_m), \quad y_i := \sum_{j \in \varphi^{-1}(i)} x_j.$$

This defines a functor

$$|\Delta^\bullet|: \mathbf{\Delta} \rightarrow \mathbf{Top}, \quad [n] \mapsto |\Delta^n|, \quad \varphi \mapsto |\Delta^\varphi|.$$

- We now define the simplicial set $\text{Sing}(X)$ as

$$\text{Sing}(X)_n := \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X),$$

i.e. it is the composite $\mathbf{\Delta}^{\text{op}} \xrightarrow{|\Delta^\bullet|} \mathbf{Top}^{\text{op}} \xrightarrow{\text{Hom}_{\mathbf{Top}}(-, X)} \mathbf{Set}$.

The simplicial set $\text{Sing}(X)$ is a familiar object from algebraic topology, and is used for example in the definition of the singular (co)homology of X . It is not difficult to see that $\text{Sing}(X)$ is functorial in X , so that it defines a functor

$$\text{Sing}: \text{Top} \rightarrow \text{sSet}.$$

Definition 1.6.9 (Geometric realization). Let $K: \Delta^{\text{op}} \rightarrow \text{Set}$ be a simplicial set. Its *geometric realization* is the quotient space

$$|K| := \left(\bigsqcup_{n \geq 0} K_n \times |\Delta^n| \right) / \sim,$$

where each set K_n carries the discrete topology and the equivalence relation is generated by

$$(\varphi^*(x), y) \sim (x, |\Delta^\varphi|(y)),$$

for all maps $\varphi: [n] \rightarrow [m]$ in Δ , elements $x \in K_m$, and points $y \in |\Delta^n|$. This construction defines a functor

$$|-|: \text{sSet} \rightarrow \text{Top}$$

which is left adjoint to the singular complex functor $\text{Sing}: \text{Top} \rightarrow \text{sSet}$.

Exercise 1.6.10. Show that the singular simplicial complex $\text{Sing}(X)$ of a topological space X is a Kan complex.

Example 1.6.11 (Nerve of a 1-category). Let C be a (small, classical) 1-category. We define the *nerve of C* as the simplicial set $N(C) \in \text{sSet}$ given by

$$N(C)_n := \text{Hom}_{\text{Cat}^{(1)}}([n], C),$$

i.e. it is the composite $\Delta^{\text{op}} \hookrightarrow (\text{Cat}_1^{(1)})^{\text{op}} \xrightarrow{\text{Hom}_{\text{Cat}^{(1)}}(-, C)} \text{Set}$. Here $\text{Cat}_1^{(1)}$ denotes the 1-category of small 1-categories.⁹ Again this construction is functorial in C , resulting in a functor

$$N: \text{Cat}_1^{(1)} \rightarrow \text{sSet}.$$

Exercise 1.6.12. Show that the nerve $N(C)$ of a classical 1-category C is a quasicategory. Provide an example for which it is not a Kan complex. Give a complete characterization of those 1-categories C for which $N(C)$ is a Kan complex.

Exercise 1.6.13. Show that the functor $N: \text{Cat}_1^{(1)} \rightarrow \text{sSet}$ is fully faithful, and that its essential image consists precisely of the simplicial sets in which every inner horn has a unique filler.

Example 1.6.14 (Homotopy coherent nerve). A further family of quasicategories is given by the homotopy coherent nerve $N^{\text{top}}(D)$ of a topologically enriched category D (see Definition 24.3.1). Since this construction will not be used in the present book, we do not discuss it here in detail; see [Lur09b, Definition 1.1.5.5]. In Section 24.3, we will instead construct the homotopy coherent nerve directly as an ∞ -category, without passing through quasicategories.

We now relate the theory of quasicategories to our axioms of ∞ -category theory:

⁹We use this notation to distinguish it from the (2,1)-category Cat_1 considered in Example 1.8.14(2).

Proposition 1.6.15. *Quasicategories satisfy axioms A–K.*

Proof sketch. This is a standard collection of results in the theory of quasicategories. Axioms A and B come from the basic definitions and the nerve construction, although spelling out the full tower of higher isomorphisms in Axiom A, together with its preferred associators and unitors, takes some bookkeeping. The remaining axioms are provided by the usual formalism of quasicategories: functor categories, opposite categories, groupoid cores, subcategories determined by 1-simplices, and localizations all exist and satisfy the expected universal properties. The verifications of Axioms I and J on recognizing equivalences are non-trivial theorems in the quasicategory model, due to Joyal [Joy08]; see also [Lan21, Theorems 2.2.1 and 2.3.20]. Since none of these model-specific facts will be used later in the book, we do not verify them here. $\square$

Warning 1.6.16. Readers who prefer a concrete model may henceforth think of ∞ -categories as quasicategories. This should only be used as a conceptual aid: statements that refer to the underlying simplicial set, such as equality of vertices or horn fillers, are statements about the model rather than about the abstract ∞ -category. In the rest of the book we therefore continue to formulate everything model-independently.

1.7 Limits and colimits

Limits and colimits are among the most important universal constructions in category theory. In the ∞ -categorical setting, their definitions look formally almost identical to the classical ones, provided one replaces sets of morphisms by hom animaes.

Definition 1.7.1 (Limit/colimit). Let I and C be ∞ -categories and let $F: I \rightarrow C$ be a functor.

- (1) A *cone* on F is an object X in C together with a natural transformation $\varepsilon: \text{const}_X \rightarrow F$ of functors $I \rightarrow C$.
- (2) A cone (X, ε) is called a *limit cone* if for every other object Y of C the composite

$$\text{Hom}_C(Y, X) \xrightarrow{\text{const}} \text{Nat}(\text{const}_Y, \text{const}_X) \xrightarrow{\varepsilon \circ -} \text{Nat}(\text{const}_Y, F)$$

is an equivalence.

In this case, the object X is called the *limit* of F . It is unique, and will be denoted either by $\lim_I F$ or $\lim_{i \in I} F(i)$.

Dually, one defines a *cocone* to be a natural transformation $\eta: F \rightarrow \text{const}_X$. It is a *colimit cocone* if for every Y the composite

$$\text{Hom}_C(X, Y) \xrightarrow{\text{const}} \text{Nat}(\text{const}_X, \text{const}_Y) \xrightarrow{- \circ \eta} \text{Nat}(F, \text{const}_Y)$$

is an equivalence. The object X is then called the *colimit* of F .

Observation 1.7.2. Note that a functor $F: I \rightarrow C$ admits a limit in C if and only if the opposite functor $F^{\text{op}}: I^{\text{op}} \rightarrow C^{\text{op}}$ admits a colimit in C^{op} .

Remark 1.7.3. If I and C are 1-categories, then so is $\text{Fun}(I, C)$, and hence all animae occurring in Definition 1.7.1 are sets. The condition on a cone (X, ε) then says precisely that for every object Y the map of sets

$$\text{Hom}_C(Y, X) \rightarrow \text{Nat}(\text{const}_Y, F), \quad u \mapsto \varepsilon \circ \text{const}_u,$$

is a bijection, that is, that every cone on F with tip Y factors uniquely through (X, ε) . So Definition 1.7.1 recovers the classical notion of a limit, and dually of a colimit, in a 1-category.

Definition 1.7.4. We say an ∞ -category C *admits all I -indexed (co)limits* if for every functor $F: I \rightarrow C$ there exists a (co)limit (co)cone for F . If $G: C \rightarrow D$ is a functor between ∞ -categories that admit I -indexed (co)limits, then we say that G *preserves I -indexed (co)limits* if for every functor $F: I \rightarrow C$, the functor G sends (co)limit (co)cones of F in C to (co)limit (co)cones of GF in D .

For later diagrammatic arguments, it is convenient to encode cones and cocones as functors out of auxiliary indexing categories:

Definition 1.7.5 (Cocone ∞ -categories). For an ∞ -category I , we define the ∞ -category $I^\flat$, called the *cocone of I* , via the following pushout square of ∞ -categories:

$$\begin{array}{ccc} I \times \{1\} & \hookrightarrow & I \times [1] \\ \downarrow & & \downarrow \\ * & \xrightarrow{\quad} & I^\flat. \end{array}$$

We will often denote the object classified by the bottom morphism as ∞ . Note that a functor $\overline{F}: I^\flat \rightarrow C$ consists of a functor $I \times [1] \rightarrow C$ whose restriction to $I \times \{1\}$ is constant. Equivalently, it consists of a functor $F := \overline{F}|_{I \times \{0\}}: I \rightarrow C$ and an object $W := \overline{F}(\infty)$ in C together with a cocone $F \rightarrow \text{const}_W$.

We say that a functor $\overline{F}: I^\flat \rightarrow C$ is a *colimit diagram* if the associated cocone $F \rightarrow \text{const}_W$ is a colimit cocone.

We dually define $I^\sharp$, called the *cone of I* , as the following pushout of ∞ -categories:

$$\begin{array}{ccc} I \times \{0\} & \hookrightarrow & I \times [1] \\ \downarrow & & \downarrow \\ * & \xrightarrow{\quad} & I^\sharp. \end{array}$$

A functor $\overline{F}: I^\sharp \rightarrow C$ consists of a functor $F: I \rightarrow C$ together with a cone $\text{const}_W \rightarrow F$. We say that $\overline{F}$ is a *limit diagram* if this cone is a limit cone.

The notions of limits and colimits are closely related to that of *adjunctions*, introduced in Chapter 21. We discuss the relations between (co)limits and adjunctions in Section 21.2.

Let us discuss some special cases of limits/colimits that will be relevant.

Products and coproducts

Definition 1.7.6. Let I be a set, regarded as a discrete ∞ -category. Given a collection of objects x_i of C , we refer to a colimit of the corresponding functor $I \rightarrow C$ as a *coproduct*, and denote it by $\coprod_{i \in I} x_i$ if it exists. We refer to a limit of $I \rightarrow C$ as a *product* and denote it by $\prod_{i \in I} x_i$.

Definition 1.7.7 (Initial/terminal objects). An object x of an ∞ -category C is called *terminal* if for every other object y the hom anima $\mathrm{Hom}_C(y, x)$ is contractible. We say that x is *initial* if for every other y the hom anima $\mathrm{Hom}_C(x, y)$ is contractible.

Exercise 1.7.8. Let I be a set, and let $\{x_i\}$ be a collection of objects of an ∞ -category C .

- (1) Show that an object x in C is a product of the objects x_i if and only if it comes equipped with morphisms $\mathrm{pr}_i: x \rightarrow x_i$ such that for any other object y of C the map

$$(\mathrm{pr}_i \circ -)_i: \mathrm{Hom}_C(y, x) \rightarrow \prod_{i \in I} \mathrm{Hom}_C(y, x_i)$$

is an equivalence of anima.

- (2) Formulate and prove the dual statement for coproducts.

Hint: show that there is an equivalence $\mathrm{Fun}(I, C) \simeq \prod_{i \in I} C$.

Exercise 1.7.9. Show that the terminal objects of C are precisely the limits of the unique functor $F: \emptyset \rightarrow C$, and that the initial objects are the colimits of $\emptyset \rightarrow C$.

Example 1.7.10. The empty anima $\emptyset$ is initial in $\mathbf{An}$ and the one-point anima $*$ is terminal in $\mathbf{An}$. Indeed, by Axiom L below the hom anima $\mathrm{Hom}_{\mathbf{An}}(X, Y)$ may be identified with the mapping anima $\mathrm{Map}(X, Y) = \mathrm{Fun}(X, Y)^\simeq$, and both $\mathrm{Fun}(\emptyset, Y)$ and $\mathrm{Fun}(X, *)$ are contractible.

Pullbacks and pushouts

Recall from Definition 1.4.1 the posets $\sqcap$ and $\sqcup$:

$$\sqcap = \begin{array}{ccc} (0, 0) & \longrightarrow & (0, 1) \\ \downarrow & & \\ (1, 0) & & \end{array} \quad \text{and} \quad \sqcup = \begin{array}{ccc} & & (0, 1) \\ & & \downarrow \\ (1, 0) & \longrightarrow & (1, 1). \end{array}$$

By Axiom D, restricting along the two edges induces equivalences

$$\begin{aligned} \mathrm{Fun}(\sqcap, C) &\xrightarrow{\sim} \mathrm{Fun}([1], C) \times_{d_1^*, C, d_1^*} \mathrm{Fun}([1], C), \\ \mathrm{Fun}(\sqcup, C) &\xrightarrow{\sim} \mathrm{Fun}([1], C) \times_{d_0^*, C, d_0^*} \mathrm{Fun}([1], C) \end{aligned}$$

for every ∞ -category C . In particular, a span $\sqcap \rightarrow C$ is equivalently encoded by a pair of morphisms $(f: x \rightarrow y, g: x \rightarrow z)$ with the same source, while a cospan $\sqcup \rightarrow C$ is equivalently encoded by a pair of morphisms $(h: y \rightarrow w, k: z \rightarrow w)$ with the same target.

Definition 1.7.11. Let C be an ∞ -category, and let $f: x \rightarrow y$ and $g: x \rightarrow z$ be two morphisms in C , corresponding to a functor $\sqcap \rightarrow C$. We refer to a colimit of this functor as a *pushout of y and z along x* . We will denote a pushout by $y \sqcup_x z$, if it exists.

Dually, given morphisms $h: y \rightarrow w$ and $k: z \rightarrow w$, we refer to a limit of the corresponding functor $\sqcup \rightarrow C$ as a *pullback of y and z over w* , and denote it by $y \times_w z$.

Remark 1.7.12. For $C = \mathbf{An}$, these notions recover the classical homotopy pushouts and homotopy pullbacks of algebraic topology. We will show in Section 2.3 that the functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ sends homotopy pullbacks of spaces to pullbacks of animae, and homotopy pushouts along relative cell complexes to pushouts of animae, see Propositions 2.3.10 and 2.3.22. This is one of the main ways in which the abstract theory of this chapter connects to classical homotopy theory.

Recall that cocones on $\ulcorner$ -indexed diagrams are encoded by the cocone category $\ulcorner^\triangleright$ from Definition 1.7.5. We will now see that this category is equivalent to the indexing diagram encoding commutative squares.

Lemma 1.7.13. *The functor $\ulcorner \times [1] \rightarrow [1] \times [1]$ described informally by the diagram*

$$\begin{array}{ccccc} (1, 0) & \longleftarrow & (0, 0) & \longrightarrow & (0, 1) \\ \downarrow & & \downarrow & & \downarrow \\ (1, 1) & \equiv & (1, 1) & \equiv & (1, 1) \end{array}$$

induces an equivalence $\ulcorner^\triangleright \xrightarrow{\sim} [1] \times [1]$.

Proof. Since $\ulcorner \times [1]$ is a pushout of two copies of $[1] \times [1]$ and $[1] \times [1]$ is a pushout of two copies of $[2]$, we may construct such a functor by gluing two copies of the functor $p_2: [1] \times [1] \rightarrow [2]$ from Construction 1.4.3. By construction, the restriction of this functor to $\ulcorner \times \{1\}$ is the identity on $(1, 1)$, hence it factors through a map $\ulcorner^\triangleright \rightarrow [1] \times [1]$. To show that it is an equivalence, we must show that for any ∞ -category C the induced commutative square

$$\begin{array}{ccc} \mathrm{Fun}([1] \times [1], C) & \longrightarrow & \mathrm{Fun}(\ulcorner \times [1], C) \\ \mathrm{ev}_{(1,1)} \downarrow & & \downarrow \mathrm{res} \\ C & \longrightarrow & \mathrm{Fun}(\ulcorner \times \{1\}, C) \end{array}$$

is a pullback square. Again using the pushout descriptions of $\ulcorner \times [1]$, $[1] \times [1]$ and $\ulcorner \times \{1\}$, this square may be identified with

$$\begin{array}{ccc} \mathrm{Fun}([2], C) \times_{d_0^*, \mathrm{Fun}([1], C), d_2^*} \mathrm{Fun}([2], C) & \longrightarrow & \mathrm{Fun}([1] \times [1], C) \times_{\mathrm{res}, \mathrm{Fun}([1], C), \mathrm{res}} \mathrm{Fun}([1] \times [1], C) \\ \downarrow & & \downarrow \mathrm{res} \\ C \times_C C & \longrightarrow & \mathrm{Fun}([1], C) \times_{d_1^*, C, d_1^*} \mathrm{Fun}([1], C). \end{array}$$

The claim now follows from the right pullback square in Proposition 1.4.4. $\square$

Exercise 1.7.14. Construct an analogous equivalence $\lrcorner^{\triangleleft} \xrightarrow{\sim} [1] \times [1]$.

Convention 1.7.15. From now on, given a diagram $\ulcorner \rightarrow C$ encoded by two morphisms $f: x \rightarrow y$ and $g: x \rightarrow z$, we will always use the lemma to identify cocones over this diagram with commutative squares in C of the form

$$\begin{array}{ccc} x & \xrightarrow{f} & y \\ g \downarrow & & \downarrow h \\ z & \xrightarrow{k} & w. \end{array} \tag{1.1}$$

Similarly, given morphisms $h: y \rightarrow w$ and $k: z \rightarrow w$, we will identify cones over the corresponding diagram $\lrcorner \rightarrow C$ with commutative squares of the above form.

Definition 1.7.16. We say a commutative square in C of the form (1.1) is a *pushout square* if the corresponding cocone in C is a colimit cocone, in which case we have $w \simeq y \sqcup_x z$. Dually, we say the square is a *pullback square* if the corresponding cone in C is a limit cone, in which case we have $x \simeq y \times_w z$.

Exercise 1.7.17 (Pasting lemma for pullback squares). Consider a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{g_1} & Y & \xrightarrow{g_2} & Z \\ f_1 \downarrow & & \downarrow f_2 & & \downarrow f_3 \\ X' & \xrightarrow{h_1} & Y' & \xrightarrow{h_2} & Z' \end{array}$$

If the right-hand square is a pullback square, then the left-hand square is a pullback square if and only if the outer rectangle is a pullback square. *Hint: compare with Lemma 1.3.15.*

Exercise 1.7.18. Formulate and prove the dual pasting lemma for pushout squares.

Geometric realizations and totalizations

Definition 1.7.19. Let C be an ∞ -category, and let $X: \Delta^{\text{op}} \rightarrow C$ be a simplicial object. The colimit of X , if it exists, is called the *geometric realization of X* , and is denoted by

$$|X| := \operatorname{colim}_{[n] \in \Delta^{\text{op}}} X_n.$$

If C admits all Δ^{op} -indexed colimits, we denote the resulting colimit functor by $|-|: sC \rightarrow C$.

Remark 1.7.20. The terminology is inspired by the classical geometric realization functor from Definition 1.6.9, which glues topological simplices according to the structure maps of a simplicial set.

Definition 1.7.21. Let C be an ∞ -category and let $X: \Delta \rightarrow C$ be a cosimplicial object. The limit of X , if it exists, is called the *totalization of X* .

1.8 Foundational results in ∞ -category theory

Having established the basic language and constructions of ∞ -categories, we now collect a small amount of foundational infrastructure that will be used repeatedly later without further proof. Since our ultimate interest in this book is stable homotopy theory and higher algebra, we treat the results below as black boxes. Table 1.1 records additional background material developed later in the book.

Background material	Discussed in
Adjunctions and Kan extensions	Chapter 21
Presheaves and free cocompletions	Chapter 22
The Hom functor	Section 23.4

Table 1.1: Background material on ∞ -category theory that is used as a black box here and developed later in the book.

The ∞ -categories $\mathbf{Cat}_\infty$ and $\mathbf{An}$

To formulate Yoneda-style constructions and presheaf categories, we need ambient ∞ -categories of small animae and small ∞ -categories. The following axiom isolates their key features that are used throughout this book.

Axiom L. There exist ∞ -categories $\mathbf{An}$ and $\mathbf{Cat}_\infty$, together with a fully faithful functor $\mathbf{An} \hookrightarrow \mathbf{Cat}_\infty$. They satisfy the following properties:

- There is a functor $\pi_{\text{univ}} : (\mathbf{Cat}_\infty)_\bullet \rightarrow \mathbf{Cat}_\infty$ called the *universal cocartesian fibration*.¹⁰ Every object $c \in \mathbf{Cat}_\infty$ defines an ∞ -category C via the following pullback square:

$$\begin{array}{ccc} C & \longrightarrow & (\mathbf{Cat}_\infty)_\bullet \\ \downarrow & \lrcorner & \downarrow \pi_{\text{univ}} \\ * & \xrightarrow{c} & \mathbf{Cat}_\infty \end{array}$$

An ∞ -category C arising this way is called a *small ∞ -category*. If c is contained in the full subcategory $\mathbf{An}$, then C is an anima; animae arising this way are called *small animae*.

- For objects $c, d \in \mathbf{Cat}_\infty$ corresponding to ∞ -categories C and D , there is an equivalence of animae

$$\mathrm{Hom}_{\mathbf{Cat}_\infty}(c, d) \xrightarrow{\sim} \mathrm{Map}(C, D),$$

and in particular morphisms in $\mathbf{Cat}_\infty$ correspond to functors of ∞ -categories. Composition in $\mathbf{Cat}_\infty$ corresponds to composition of functors.

- The ∞ -categories $\mathbf{An}$ and $\mathbf{Cat}_\infty$ admit all small limits and colimits; here the adjective ‘small’ means that the indexing ∞ -category is required to be small. The inclusion $\mathbf{An} \hookrightarrow \mathbf{Cat}_\infty$ preserves small limits and colimits.
- The initial and terminal ∞ -categories are small. Small ∞ -categories are closed under products, coproducts, pullbacks, pushouts, functor categories, subcategories, and localizations.
- There is a fully faithful functor

$$\mathbf{\Delta} \hookrightarrow \mathbf{Cat}_\infty$$

given on objects by sending a linearly ordered set P to its associated ∞ -category. In particular, the ∞ -categories $[n]$ are small for all $n \geq 0$.

- There is a fully faithful functor

$$\mathbf{Set} \hookrightarrow \mathbf{An}$$

given on objects by sending a set to its associated anima.

Remark 1.8.1 (Large ∞ -categories). The ∞ -categories $\mathbf{An}$ and $\mathbf{Cat}_\infty$ are sometimes referred to as the *universe of animae* and the *universe of ∞ -categories*. It will sometimes be convenient to assume there are two more such universes $\widehat{\mathbf{An}}$ and $\widehat{\mathbf{Cat}}_\infty$ that satisfy all the same conditions as $\mathbf{An}$ and $\mathbf{Cat}_\infty$, with the following two further conditions:

¹⁰This terminology will be explained in Chapter 23.

- There is a fully faithful inclusion $\text{Cat}_\infty \hookrightarrow \widehat{\text{Cat}}_\infty$ which on small animae restricts to a functor $\text{An} \hookrightarrow \widehat{\text{An}}$;
- The ∞ -categories An and Cat_∞ are small with respect to $\widehat{\text{Cat}}_\infty$.

We will refer to ∞ -categories that are small with respect to $\widehat{\text{Cat}}_\infty$ as *large ∞ -categories*. The two points may thus be summarized by saying that every small ∞ -category is also large, and that the ∞ -categories An and Cat_∞ are large.

All categorical constructions are understood relative to fixed ambient universes, which we silently enlarge when necessary.

Remark 1.8.2 (Two meanings of ‘pullback square’). The ambient category Cat_∞ lets us reconcile two pieces of terminology that we have so far used in parallel. In Section 1.3 we defined products, coproducts, pullbacks and pushouts *of* ∞ -categories, together with the corresponding notions of pullback and pushout square, purely in terms of functors and natural isomorphisms. In Section 1.7 we defined limits and colimits *inside* an ∞ -category, and hence in particular inside Cat_∞ . For small ∞ -categories the two notions agree:

- A commutative square of functors in the sense of Definition 1.2.2 between small ∞ -categories is the same thing as a commutative square in Cat_∞ in the sense of Definition 1.4.1. Indeed, by Axiom L the hom animae of Cat_∞ are the mapping animae $\text{Map}(C, D)$, so filling a square of functors amounts to giving a natural isomorphism between the two composites.
- Such a square is a pullback square of ∞ -categories in the sense of Section 1.3 if and only if it is a pullback square in Cat_∞ in the sense of Section 1.7, and similarly for pushouts, products and coproducts. In both cases the condition says that for every ∞ -category T the induced map on mapping animae out of, respectively into, T is an equivalence.

The one point to keep in mind is a matter of universes: the definitions in Section 1.3 quantify over *all* ∞ -categories T , whereas a (co)limit in Cat_∞ only tests against small ones. By Remark 1.8.1 this is harmless, since the constructions of Section 1.3 are compatible with enlarging the ambient universe. From now on we use the two readings interchangeably.

Remark 1.8.3. In the quasicategorical model for ∞ -categories, the statements in Axiom L are proved by [Lur09b]. The quasicategory modeling Cat_∞ is taken to be the homotopy coherent nerve of the Kan-enriched category of quasicategories, while the quasicategory modeling An is the homotopy coherent nerve of the Kan-enriched category of Kan complexes.

Presheaves and free cocompletions

With these ambient categories in place, we can record the presheaf formalism that will be used repeatedly later.

Let C be an ∞ -category. Then the hom animae $\text{Hom}_C(x, y)$ for $x, y \in C$ assemble into a functor

$$\text{Hom}_C : C^{\text{op}} \times C \rightarrow \text{An}$$

called the *Hom functor*. Formally constructing this functor turns out to be surprisingly involved: it relies on the advanced concept of straightening/unstraightening from Chapter 23. We refer to Section 23.4 for a construction of the Hom functor.

Definition 1.8.4. A *presheaf* on an ∞ -category C is a functor $F : C^{\text{op}} \rightarrow \mathbf{An}$. We define the *presheaf category* of C as

$$\mathbf{PSh}(C) := \mathbf{Fun}(C^{\text{op}}, \mathbf{An}).$$

By currying, the Hom functor defines a functor $Y : C \rightarrow \mathbf{PSh}(C)$ called the *Yoneda embedding*; for $x \in C$ we refer to the presheaf $Y(x) := \text{Hom}_C(-, x)$ as the *presheaf represented by x* . A presheaf is called *representable* if it is isomorphic to $Y(x)$ for some x .

The ∞ -categorical analogue of the Yoneda lemma is even more important than its classical analogue:

Theorem 1.8.5 (Yoneda lemma, [Lan21, Propositions 4.2.10 and 4.2.11]). *For a presheaf F on C and an object $x \in C$, evaluation at $\text{id}_x \in \text{Hom}_C(x, x) = Y(x)(x)$ induces an equivalence*

$$\text{Nat}(Y(x), F) \xrightarrow{\sim} F(x), \quad (\varphi : Y(x) \rightarrow F) \mapsto \varphi(\text{id}_x).$$

In particular the Yoneda embedding $Y : C \hookrightarrow \mathbf{PSh}(C)$ is fully faithful.

The following two results will be proved in Corollary 23.6.7 and Corollary 21.2.9, respectively.

Proposition 1.8.6. *The functor $\text{Hom}_C(-, -) : C^{\text{op}} \times C \rightarrow \mathbf{An}$ preserves limits in both variables separately.*

Proposition 1.8.7. *The presheaf category $\mathbf{PSh}(C)$ admits all small limits and colimits. The evaluation functors $\text{ev}_x : \mathbf{PSh}(C) \rightarrow \mathbf{An}$ preserve all small limits and colimits.*

Exercise 1.8.8. Let C be an ∞ -category. Deduce that the Yoneda embedding $Y : C \hookrightarrow \mathbf{PSh}(C)$ preserves all limits that exist in C .

While the inclusion of C into $\mathbf{PSh}(C)$ preserves all limits, it does not in general preserve colimits. Instead, it *freely adds colimits* to C , in the following sense:

Theorem 1.8.9 (Lurie [Lur09b, Theorem 5.1.5.6]). *Let C be a small ∞ -category and let D be a large ∞ -category admitting all small colimits. Then restriction along the Yoneda embedding $Y : C \hookrightarrow \mathbf{PSh}(C)$ induces an equivalence*

$$Y^* : \mathbf{Fun}^{\text{colim}}(\mathbf{PSh}(C), D) \xrightarrow{\sim} \mathbf{Fun}(C, D),$$

where the left-hand side denotes the full subcategory of colimit-preserving functors $\mathbf{PSh}(C) \rightarrow D$.

More informally, this says that every functor $u : C \rightarrow D$ admits a unique colimit-preserving extension $Y_!(u) : \mathbf{PSh}(C) \rightarrow D$. This functor can be described directly in terms of the concept of *left Kan extension*, developed systematically in Section 21.4. Informally, a left Kan extension $i_!(u) : E \rightarrow D$ of a functor $u : C \rightarrow D$ along a functor $i : C \rightarrow E$ is the best approximation to an extension of u to E : its value at an object e of E is assembled as a colimit of the values $u(c)$ over all maps $i(c) \rightarrow e$. In the situation $i = Y$ of the theorem, the full faithfulness of the Yoneda embedding guarantees that for $e = Y(c)$, this Kan extension recovers $u(c)$.

The special case $C = *$ is the form that we will use most often in Part I. Since $\mathbf{PSh}(*) \simeq \mathbf{An}$, it gives the following universal property.

Corollary 1.8.10 (Universal property of animae). *Let D be a large ∞ -category admitting all small colimits. Evaluation at $*$ $\in \mathbf{An}$ induces an equivalence*

$$\mathrm{ev}_*: \mathrm{Fun}^{\mathrm{colim}}(\mathbf{An}, D) \xrightarrow{\sim} D.$$

Given an object $d \in D$, the inverse equivalence sends d to a functor of the form

$$\mathbf{An} \rightarrow D, \quad X \mapsto \mathrm{colim}_X d,$$

where the right-hand side denotes the colimit of the constant diagram with value d , indexed by the anima X .

Returning now to the simplicial language from Section 1.6, we may use this theorem to assign an ∞ -category to every simplicial anima:

Construction 1.8.11 (Associated ∞ -category). Every simplicial anima $X: \Delta^{\mathrm{op}} \rightarrow \mathbf{An}$ has an associated ∞ -category $\mathrm{ac}(X) \in \mathrm{Cat}_\infty$. To construct it, consider the inclusion $\Delta \hookrightarrow \mathrm{Cat}_\infty$ from Axiom L. By Theorem 1.8.9, this uniquely extends to a colimit-preserving functor

$$\mathrm{ac}: \mathbf{sAn} = \mathrm{PSh}(\Delta) \rightarrow \mathrm{Cat}_\infty.$$

In particular, every simplicial set X (hence every quasicategory) defines an ∞ -category $\mathrm{ac}(X)$ by regarding it as a simplicial anima via the inclusion $\mathbf{sSet} \hookrightarrow \mathbf{sAn}$.

Remark 1.8.12. Note that the composite

$$\mathrm{Cat}_1^{(1)} \xrightarrow{N} \mathbf{sSet} \hookrightarrow \mathbf{sAn} \xrightarrow{\mathrm{ac}} \mathrm{Cat}_\infty$$

provides a way of associating an ∞ -category to a classical 1-category. From this point forward, we stop working with Axiom B and use this refined assignment. The two assignments agree on the indexing categories used so far in this chapter: on the linear orders $[n]$ this is the fully faithful functor $\Delta \hookrightarrow \mathrm{Cat}_\infty$ from Axiom L, and the categories $[1] \times [1]$, $\lceil$, $\rfloor$ and $[\omega]$ are obtained from the $[n]$ by the coproducts and pushouts of Axiom D, which both assignments preserve.

Adjunctions

The last piece of black-box infrastructure we record here is the notion of an *adjunction*, which interacts closely with the limit and colimit constructions introduced above:

Definition 1.8.13. Let $F: C \rightarrow D$ and $G: D \rightarrow C$ be two functors. An *adjunction* between C and D consists of a natural isomorphism

$$\mathrm{Hom}_D(F(-), -) \cong \mathrm{Hom}_C(-, G(-))$$

of functors $C^{\mathrm{op}} \times D \rightarrow \mathbf{An}$. We often write $F \dashv G$ to indicate that such an isomorphism has been given.

As in classical category theory, adjunctions may equivalently be characterized in terms of a unit $\eta: \mathrm{id}_C \rightarrow GF$ and a counit $\varepsilon: FG \rightarrow \mathrm{id}_D$ satisfying the triangle identities $(\varepsilon F)(F\eta) \cong \mathrm{id}_F$ and $(G\varepsilon)(\eta G) \cong \mathrm{id}_G$. An important feature of adjunctions is that they can be characterized *pointwise*:

given a functor $F: C \rightarrow D$, if one can provide for every object Y of D a suitable object $G(Y)$ of C equipped with a map $\varepsilon_Y: FG(Y) \rightarrow Y$ satisfying a suitable adjointness condition, then these objects automatically assemble into a right adjoint $G: D \rightarrow C$, and the maps ε_Y assemble into a counit ε for the adjunction.

The theory of adjunctions is very rich and there is a lot to say about it. Details on the pointwise characterization of adjunctions, along with related concepts such as Kan extensions, (co)final functors, and Bousfield localizations, can be found in Chapter 21.

Example 1.8.14. Let us state as black boxes two fundamental examples of adjunctions we will use:

- (1) (Path components) The fully faithful inclusion $\text{Set} \hookrightarrow \text{An}$ admits a left adjoint $\pi_0: \text{An} \rightarrow \text{Set}$. This functor preserves arbitrary products.

We call $\pi_0(X)$ the *set of path components* of the anima X . We often abuse terminology and refer to an anima as a *set* if it is in the essential image of the inclusion $\text{Set} \hookrightarrow \text{An}$ (cf. Example 1.1.4).

- (2) (Homotopy category) We say that an ∞ -category C is a *1-category* if the hom anima $\text{Hom}_C(X, Y)$ is a set for all $X, Y \in C$. We denote by

$$\text{Cat}_1 \hookrightarrow \text{Cat}_\infty$$

the full subcategory spanned by the 1-categories. Then this inclusion admits a left adjoint

$$\text{Ho}: \text{Cat}_\infty \rightarrow \text{Cat}_1.$$

Given an ∞ -category C , we refer to the 1-category $\text{Ho}(C)$ as its *homotopy category*; it has the same set of isomorphism classes of objects as C , but has hom sets given by the set of path components of the hom anima of C :

$$\text{Hom}_{\text{Ho}(C)}(X, Y) \cong \pi_0(\text{Hom}_C(X, Y)).$$

Remark 1.8.15. The functor $\text{Cat}_1^{(1)} \rightarrow \text{Cat}_\infty$ from Remark 1.8.12 lands in the full subcategory Cat_1 of 1-categories. In particular, every classical 1-category defines a 1-category in the above sense. Conversely, every 1-category arises this way. We will use the phrase ‘classical 1-category’ if we wish to refer to the classical definition of categories in terms of sets of objects and morphisms.

Exercises

Exercise 1.1. Show that coproducts of ∞ -categories are *disjoint*: given ∞ -categories C and D , the commutative square

$$\begin{array}{ccc} \emptyset & \longrightarrow & C \\ \downarrow & & \downarrow \text{incl}_C \\ D & \xrightarrow{\text{incl}_D} & C \sqcup D \end{array}$$

is a pullback square. In a similar spirit, show that for every functor $f: C \rightarrow D$ the commutative square

$$\begin{array}{ccc} \emptyset & \xrightarrow{\text{id}_\emptyset} & \emptyset \\ \downarrow & & \downarrow \\ C & \xrightarrow{f} & D \end{array}$$

is a pullback square. In both cases, explain also which commutative square is meant, that is, which natural isomorphism fills it. *Hint:* use Axiom C.3.

Exercise 1.2. Consider two functors $f: C \rightarrow E$ and $g: D \rightarrow E$. Show that the commutative square

$$\begin{array}{ccc} C \times_E D & \xrightarrow{f \circ \text{pr}_C} & E \\ (\text{pr}_C, \text{pr}_D) \downarrow & & \downarrow (\text{id}_E, \text{id}_E) \\ C \times D & \xrightarrow{f \times g} & E \times E \end{array}$$

is a pullback square. In other words, construct an equivalence

$$C \times_E D \simeq (C \times D) \times_{E \times E} E.$$

Hint: Consider the commutative diagram

$$\begin{array}{ccccc} C \times_E D & \xrightarrow{\text{pr}_D} & D & \xrightarrow{g} & E \\ (\text{pr}_C, \text{pr}_D) \downarrow & & \downarrow (g, \text{id}_D) & & \downarrow (\text{id}_E, \text{id}_E) \\ C \times D & \xrightarrow{f \times \text{id}_D} & E \times D & \xrightarrow{\text{id}_E \times g} & E \times E \\ \text{pr}_C \downarrow & & \downarrow \text{pr}_E & & \\ C & \xrightarrow{f} & E & & \end{array}$$

The vertical composite of the two left-hand squares is the square defining $C \times_E D$. Use Exercises 1.3.10 and 1.3.16 and Lemma 1.3.15 to show that the bottom-left and top-right squares are pullback squares, and then apply pullback pasting twice.

Exercise 1.3. Use Axiom D to construct functors

$$\min: [1] \times [1] \rightarrow [1] \quad \text{and} \quad \max: [1] \times [1] \rightarrow [1]$$

that behave like the minimum and the maximum on the linear order $\{0 \leq 1\}$, in the sense that

$$\begin{aligned} \max(x, 0) &\cong x \cong \max(0, x), & \max(x, 1) &\cong 1 \cong \max(1, x), \\ \min(x, 0) &\cong 0 \cong \min(0, x), & \min(x, 1) &\cong x \cong \min(1, x). \end{aligned}$$

Spelled out in terms of Definition 1.4.1, this says that the restrictions of $\max$ to the left column $[1] \times \{0\}$ and to the top row $\{0\} \times [1]$ are $\text{id}_{[1]}$, while its restrictions to the right column $[1] \times \{1\}$ and to the bottom row $\{1\} \times [1]$ are const_1 ; for $\min$ the restrictions to the left column and the top row are const_0 , and those to the right column and the bottom row are $\text{id}_{[1]}$. Also show that the functors you construct do satisfy these conditions.

Exercise 1.4. Let C and D be ∞ -categories and let $x, x' \in C$ and $y, y' \in D$ be objects. Produce an equivalence

$$\mathrm{Hom}_{C \times D}((x, y), (x', y')) \simeq \mathrm{Hom}_C(x, x') \times \mathrm{Hom}_D(y, y').$$

Show that composition in $C \times D$ corresponds to composition in C and D under these equivalences.

Exercise 1.5. Let $f: C \rightarrow E$ and $g: D \rightarrow E$ be functors, and let $x, x' \in C \times_E D$ be objects. Show that there is a pullback square

$$\begin{array}{ccc} \mathrm{Hom}_{C \times_E D}(x, x') & \xrightarrow{\quad} & \mathrm{Hom}_C(\mathrm{pr}_C(x), \mathrm{pr}_C(x')) \\ \downarrow & \lrcorner & \downarrow \\ \mathrm{Hom}_D(\mathrm{pr}_D(x), \mathrm{pr}_D(x')) & \xrightarrow{\quad} & \mathrm{Hom}_E(f \mathrm{pr}_C(x), f \mathrm{pr}_C(x')). \end{array}$$

Here the lower horizontal functor is induced by g , followed by transport along the structural isomorphisms $g \mathrm{pr}_D(x) \cong f \mathrm{pr}_C(x)$ and $g \mathrm{pr}_D(x') \cong f \mathrm{pr}_C(x')$ carried by the two objects of the pullback.

Exercise 1.6. Show that a functor $F: C \rightarrow D$ is a monomorphism if and only if the projection $\mathrm{pr}_1: C \times_D C \rightarrow C$ is an equivalence.

Exercise 1.7. Consider a pullback square

$$\begin{array}{ccc} C' & \xrightarrow{\quad} & C \\ F' \downarrow & \lrcorner & \downarrow F \\ D' & \xrightarrow{\quad} & D. \end{array}$$

Show that if F is a monomorphism, then so is F' .

Exercise 1.8. Show that monomorphisms are closed under retracts. Conclude that if $i: A \rightarrow C$ induces a monomorphism $\mathrm{Map}([1], A) \hookrightarrow \mathrm{Map}([1], C)$, then it also induces a monomorphism $A^\simeq \hookrightarrow C^\simeq$.

Exercise 1.9. Let $i: A \rightarrow C$ be a monomorphism. Show that $i^\simeq: A^\simeq \rightarrow C^\simeq$ is a monomorphism. Show that for every ∞ -category T , the functor $i_*: \mathrm{Fun}(T, A) \rightarrow \mathrm{Fun}(T, C)$ is a monomorphism.

Exercise 1.10. Consider a pullback square of ∞ -categories

$$\begin{array}{ccc} A' & \xrightarrow{\quad} & A \\ i' \downarrow & \lrcorner & \downarrow i \\ C' & \xrightarrow{\quad} & C. \end{array}$$

Show that if i is a monomorphism, then so is i' . Show moreover that if i is a full subcategory inclusion, then so is i' .

Exercise 1.11. Construct the commutative square appearing in the definition of full subcategory inclusion (Definition 1.5.9).

Exercise 1.12. The goal of this exercise is to show that the anima $\mathrm{Map}([1], [1])$ is equivalent to a disjoint union of three points.

- (1) Construct a retraction r of the evaluation functor

$$e := (\text{ev}_0, \text{ev}_1): \text{Map}([1], [1]) \rightarrow [1]^\simeq \times [1]^\simeq,$$

that is, a map $r: [1]^\simeq \times [1]^\simeq \rightarrow \text{Map}([1], [1])$ together with an equivalence $r \circ e \simeq \text{id}$.

Hint: using Exercise 1.3, consider the functor

$$m: [1] \times [1] \times [1] \rightarrow [1], \quad (x, y, z) \mapsto \min(\max(x, y), \min(\max(x, z), \max(y, z))).$$

- (2) Deduce from (1) that $\text{Map}([1], [1])$ is a set. Under the equivalence $[1]^\simeq \times [1]^\simeq \simeq \{(0, 0), (0, 1), (1, 0), (1, 1)\}$, show that the fiber of e over $(1, 0)$ is empty, while each of the other three fibers is inhabited.

- (3) Conclude that the canonical map

$$* \sqcup * \sqcup * \rightarrow \text{Map}([1], [1])$$

selecting the three functors const_0 , $\text{id}_{[1]}$, and const_1 is an equivalence.

Exercise 1.13. Show that the functor $p_{[1]}: [1] \rightarrow *$ exhibits $*$ as the localization of $[1]$ at the collection $W = \text{Map}([1], [1])$ of all morphisms.

- (1) Show that $p_{[1]}$ inverts all morphisms in $\text{Map}([1], [1])$.
- (2) Let D be an ∞ -category. Use Exercise 1.12 to rewrite the pullback square defining $\text{Map}^W([1], D)$, and deduce that $\text{Map}^W([1], D)$ is equivalent to $\text{Iso}(D)^\simeq$. Use the Rezk axiom to conclude that the functor

$$p_{[1]}^*: D^\simeq \rightarrow \text{Map}^W([1], D)$$

is an equivalence.

- (3) By Axiom K, let $l: [1] \rightarrow L$ be a localization of $[1]$ at W . Use the universal property of l together with (2) to show that for every ∞ -category D , the unique functor $L \rightarrow *$ induces an equivalence

$$\text{Map}(L, D) \rightarrow \text{Map}(*, D),$$

and thus is an equivalence.

Exercise 1.14. Let $l: C \rightarrow L$ be a functor, and let $W \hookrightarrow \text{Map}([1], C)$ be the collection defined by the pullback square

$$\begin{array}{ccc} W & \xrightarrow{\quad} & \text{Map}([1], C) \\ \downarrow & \lrcorner & \downarrow l_* \\ \text{Iso}(L)^\simeq & \xrightarrow{\pi_{\text{Iso}}} & \text{Map}([1], L). \end{array}$$

Show that l is a localization of C at *some* collection of morphisms if and only if it is a localization of C at this class W .

Exercise 1.15. Show that any localization $l: C \rightarrow L$ is essentially surjective: for any object $x \in L$, there exists some $y \in C$ with $x \cong l(y)$. You may use without proof that every map of animae admits an image factorization. *Hint:* Apply this fact to $C^\simeq \rightarrow L^\simeq$, let $L' \subseteq L$ be the full subcategory on the resulting image collection, and apply the universal property of l to the map $C \rightarrow L'$.

Exercise 1.16. Show that localizations are closed under pushouts: for a pushout square of ∞ -categories

$$\begin{array}{ccc} C & \longrightarrow & C' \\ \downarrow l & \lrcorner & \downarrow l' \\ L & \longrightarrow & L', \end{array}$$

if l is a localization, then so is l' .

Exercise 1.17 (cf. Proposition 1.4.4). Use the previous exercise to show that the two functors

$$p_0: [1] \times [1] \rightarrow [2] \quad \text{and} \quad p_2: [1] \times [1] \rightarrow [2]$$

are localizations of $[1] \times [1]$ at the morphism $(0, 0) \rightarrow (1, 0)$ and $(0, 1) \rightarrow (1, 1)$, respectively; that is, at the left and the right column of the square. Here p_0 and p_2 are the functors constructed in Construction 1.4.3, pictorially represented by the following two diagrams:

$$p_0 = \begin{array}{ccc} 0 & \longrightarrow & 1 \\ \parallel & & \downarrow \\ 0 & \longrightarrow & 2, \end{array} \quad p_2 = \begin{array}{ccc} 0 & \longrightarrow & 2 \\ \downarrow & & \parallel \\ 1 & \longrightarrow & 2. \end{array}$$

Exercise 1.18. Let C be an ∞ -category. Show that the following statements are equivalent:

- (1) The ∞ -category C is an anima;
- (2) The functor $p_{[1]}^*: C \rightarrow \text{Ar}(C)$ is an equivalence;
- (3) The functor $p_{[1]}^*: C^\simeq \rightarrow \text{Map}([1], C)$ is an equivalence;
- (4) The functor $\pi_{\text{Iso}}: \text{Iso}(C) \rightarrow \text{Ar}(C)$ is an equivalence.

Hint: Use Exercise 1.13 and the Rezk axiom to show that $p_{[1]}^*: C \rightarrow \text{Ar}(C)$ identifies C with $\text{Iso}(C)$.

Exercise 1.19. Let C be an ∞ -category. The goal of this exercise is to identify the groupoid core $C^\simeq$ of C with the subcategory $\langle M \rangle_C$ of C spanned by the isomorphisms $M := \text{Iso}(C)^\simeq \hookrightarrow \text{Map}([1], C)$.

- (1) Show that M is closed under composition, so that the subcategory $\langle M \rangle_C$ is defined.
- (2) Show that the map $p_{[1]}^*: \langle M \rangle_C^\simeq \hookrightarrow \text{Map}([1], \langle M \rangle_C)$ is an equivalence, and deduce from Exercise 1.18 that $\langle M \rangle_C$ is an anima.
- (3) Show that the core functor $\gamma_C: C^\simeq \rightarrow C$ sends all morphisms to isomorphisms, hence induces a functor $C^\simeq \rightarrow \langle M \rangle_C$.
- (4) Show that this functor induces an equivalence on groupoid cores. Use (2) to conclude that it is an equivalence.

Exercise 1.20. Let C be an ∞ -category. The goal of this exercise is to show that its geometric realization $\|C\|$ is an anima.

- (1) Use Exercise 1.19 to show that the localization functor $C \rightarrow \|C\|$ factors through a functor $C \rightarrow \|C\|^\simeq$.
- (2) Argue that this functor inverts all morphisms, hence induces a functor $\|C\| \rightarrow \|C\|^\simeq$.
- (3) Show that this provides an inverse to the inclusion $\|C\|^\simeq \rightarrow \|C\|$ and conclude that $\|C\|$ is an anima.

Exercise 1.21. This exercise uses the theory developed in Chapter 21. Use the pointwise criterion for adjunctions (Lemma 21.1.4) to show that the inclusion $\mathbf{An} \hookrightarrow \mathbf{Cat}_\infty$ admits both a left adjoint $\|- \|: \mathbf{Cat}_\infty \rightarrow \mathbf{An}$ as well as a right adjoint $(-)^\simeq: \mathbf{Cat}_\infty \rightarrow \mathbf{An}$.

Part I

Stable homotopy theory

Introduction to Part I

This first part develops the foundations of stable homotopy theory in the language of ∞ -categories. Its central object is the ∞ -category Sp of spectra: a setting in which every generalized (co)homology theory is representable, in which the stability axiom becomes a structural property rather than an axiom, and in which the constructions of homological algebra may be performed without first deriving them.

Throughout we build on Chapter 1, which precedes both mathematical parts and supplies the model-independent language of ∞ -categories and of *animaes*, the ∞ -categories in which every morphism is invertible. We begin here with the topological preliminaries. Chapter 2 identifies animaes with spaces up to weak homotopy equivalence, and transports the classical homotopy pushouts and pullbacks, suspensions and loop spaces into this setting. Chapter 3 treats (co)homology theories axiomatically and introduces spectra together with Brown’s representability theorem; the online version also develops cellular homology from the axioms and proves Brown representability.

Spectra are then studied in their own right. Chapter 4 develops fiber and cofiber sequences, the notion of a stable ∞ -category, and the stabilization $\mathrm{Sp}(C)$. Chapter 5 proves the recognition principle, identifying connective spectra with commutative groups in An , so that a connective spectrum becomes a coherent analogue of an abelian group. Chapter 6 develops homological algebra from this viewpoint: derived ∞ -categories, t-structures and their hearts, Ext- and Tor-groups, and derived functors. Chapter 7 treats localization and completion of spectra at a prime.

The remaining chapters treat the principal examples. Chapter 8 develops ring spectra and their modules, taking the coherent algebra of Part II as a black box. Chapter 9 constructs complex K-theory, Chapter 10 the Thom spectra and their bordism theories, and Chapter 11 treats Spanier–Whitehead, Atiyah and Poincaré duality.

2 Animaes as spaces

In Chapter 1, we introduced the notion of an *anima*: the homotopical analogue of a set, formally defined as an ∞ -category in which all morphisms are invertible. At first glance, this abstract concept may seem disconnected from algebraic topology and classical homotopy theory. The goal of this chapter is to explain that there is in fact a close relationship: animaes may be thought of as *spaces up to (weak) homotopy equivalence*. In particular, we will see that the classical constructions of cones, suspensions, loop spaces, homotopy pushouts, homotopy pullbacks and wedges have analogues phrased purely in terms of the ∞ -category $\mathbf{An}$.

The connection between spaces and animaes is the content of Grothendieck’s *homotopy hypothesis*. To make it precise, we first need a way to pass from spaces to animaes, that is, a functor $\mathbf{Top} \rightarrow \mathbf{An}$. A topological space already behaves informally like an ∞ -groupoid: its points are objects, paths are morphisms, homotopies between paths are higher morphisms, and this hierarchy continues indefinitely. This structure on X is neatly encoded by its singular complex: Recall from Example 1.6.8 that a topological space X has a simplicial set

$$\mathrm{Sing}(X)_n := \mathrm{Hom}_{\mathbf{Top}}(|\Delta^n|, X).$$

Thus its 0-simplices are the points of X , its 1-simplices are paths, and its higher simplices encode homotopies and their higher coherences.

We may regard $\mathrm{Sing}(X)$ as a simplicial anima via the inclusion $\mathbf{Set} \hookrightarrow \mathbf{An}$ from Axiom L. We then define the *underlying anima*, or *fundamental ∞ -groupoid*, of X by

$$\Pi_\infty(X) := \mathrm{colim}_{[n] \in \Delta^{\mathrm{op}}} \mathrm{Sing}(X)_n \in \mathbf{An}.$$

This colimit glues together the singular simplices of X . Although each individual topological simplex is contractible, the way these simplices meet along their faces retains the homotopy type of X . Here it is crucial that the colimit is taken in $\mathbf{An}$: it remembers the gluing homotopy coherently, rather than merely identifying points as an ordinary colimit in $\mathbf{Set}$ would do. The construction is functorial in X , and thus defines a functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$.

How faithfully does Π_∞ reflect the homotopy theory of spaces? As a first hint, one can show with a little effort that it sends every homotopy equivalence of topological spaces to an isomorphism of animaes; in short, this holds because it preserves finite products and sends the interval to a contractible anima. In particular, the underlying anima of a topological space is a homotopy invariant of the space.

With some more effort, one can show that $\Pi_\infty(X)$ is in fact a *weak homotopy invariant* of the space X . Recall that algebraic topology studies spaces by assigning algebraic invariants to them. Among these invariants are the *homotopy groups*: generalizing the definition of the fundamental group as

homotopy classes of loops in a space, one defines for each $n \geq 1$ the n -th homotopy group of a space X at a basepoint x as the set

$$\pi_n(X, x) := [S^n, X]_*$$

of pointed homotopy classes of pointed continuous maps from the topological n -sphere $S^n \subseteq \mathbb{R}^{n+1}$ into X , with group structure defined via some form of concatenation. A continuous map $f: X \rightarrow Y$ is called a *weak homotopy equivalence* if it induces a bijection on sets of path components and isomorphisms $\pi_n(X, x) \xrightarrow{\cong} \pi_n(Y, f(x))$ on homotopy groups for all $n \geq 1$ and all choices of basepoint $x \in X$. An algebraic invariant of topological spaces is called a *weak homotopy invariant* if it sends every weak homotopy equivalence to an isomorphism in its target category; examples are the singular homology and cohomology of a space. A large part of algebraic topology is devoted to studying spaces up to weak homotopy equivalence.

If we allow our invariants of spaces to take values in arbitrary ∞ -categories, then the functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ constructed before turns out to be a weak homotopy invariant. Proving this requires classical results in simplicial homotopy theory that go beyond the scope of this book, and we refer to [Lur26, Tag 00U2 and Tag 013P] for an exposition of this theory. The key fact relevant here is that a continuous map $X \rightarrow Y$ of topological spaces is a weak homotopy equivalence if and only if the induced map of Kan complexes $\mathrm{Sing}(X) \rightarrow \mathrm{Sing}(Y)$ is a simplicial homotopy equivalence. Since the geometric realization $|\Delta^1|$ is contractible, it follows that $|-|: \mathbf{sSet} \rightarrow \mathbf{An}$ inverts simplicial homotopy equivalences, and we conclude that Π_∞ is a weak homotopy invariant.

Grothendieck's homotopy hypothesis says that Π_∞ is not just any weak homotopy invariant; it is, in the strongest possible sense, the *universal* one. We enforce this behavior in our synthetic setup via the following axiom:

Axiom M (Grothendieck's homotopy hypothesis). The morphisms inverted by the functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ are precisely the weak homotopy equivalences. Moreover, Π_∞ exhibits $\mathbf{An}$ as a localization of $\mathbf{Top}$ at the weak homotopy equivalences, that is, it induces an equivalence of ∞ -categories

$$\mathbf{Top}[\{\text{weak homotopy equivalences}\}^{-1}] \xrightarrow{\sim} \mathbf{An}.$$

In light of the homotopy hypothesis, we may think of $\mathbf{An}$ as the universal home for studying topological spaces up to weak homotopy equivalence. The rest of the chapter adds substance to this claim by transporting familiar constructions from algebraic topology into the world of animae. The main focus will be the classical theory of *homotopy pushouts and pullbacks*, which we recall in Section 2.1. In Section 2.2 we introduce a general framework for computing pushouts and pullbacks in localizations due to Cisinski [Cis19, Chapter 7], which we apply in Section 2.3 to show that Π_∞ sends homotopy pushouts and pullbacks of spaces to ∞ -categorical pushouts and pullbacks of animae. This yields the basic constructions on pointed animae, such as fibers, cofibers, suspensions, and loop spaces, developed in Section 2.4, which will become crucial in the remainder of the book.

Remark. As justification for Axiom M, we note it is satisfied in the quasicategorical model, where $\mathbf{An}$ is the quasicategory $\mathcal{S}$ of spaces. There Lurie identifies $\mathcal{S}$ with the localization of the 1-category $\mathbf{Kan}$ at the homotopy equivalences of Kan complexes [Lur17, Proposition 1.3.4.7], and by classical results from model category theory [Qui67] this agrees with the localization of $\mathbf{Top}$ at the weak homotopy equivalences.

2.1 Homotopy pushouts and pullbacks

Ordinary pushouts and pullbacks are among the most basic constructions one performs on spaces: think of gluing two spaces along a common subspace, or of the fiber of a fiber bundle over a point. For homotopy theory, however, they have a fundamental defect: they are not automatically homotopy invariant, in that two diagrams of spaces related by homotopy equivalences may have pushouts, or pullbacks, that are not homotopy equivalent.

Example 2.1.1. Given two maps $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ of topological spaces, recall that the pushout in $\mathbf{Top}$ is given by the quotient space

$$X \sqcup_Z Y := (X \sqcup Y)/\sim, \quad \text{where} \quad f(z) \sim g(z) \text{ for all } z \in Z.$$

If we let D^n denote the n -dimensional disk, and let $S^{n-1} \hookrightarrow D^n$ denote its boundary, then we have the following two pushout squares in $\mathbf{Top}$:

$$\begin{array}{ccc} S^{n-1} & \hookrightarrow & D^n \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & S^n \end{array} \quad \text{and} \quad \begin{array}{ccc} S^{n-1} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & *. \end{array}$$

These two diagrams are homotopy equivalent, by contractibility of the disk D^n , yet their pushouts S^n and $*$ are not homotopy equivalent for $n \geq 1$.

Example 2.1.2. Given two continuous maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$, their pullback $X \times_Z Y$ is given by the subspace

$$X \times_Z Y := \{(x, y) \in X \times Y \mid f(x) = g(y)\} \subseteq X \times Y.$$

To see that pullbacks are not homotopy invariant, suppose that Z is a topological space and $p: [0, 1] \rightarrow Z$ a path whose endpoints $z_0 := p(0)$ and $z_1 := p(1)$ differ, and consider the following two pullback squares in $\mathbf{Top}$:

$$\begin{array}{ccc} * & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow z_0 \\ * & \xrightarrow{z_0} & Z \end{array} \quad \text{and} \quad \begin{array}{ccc} \emptyset & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow z_0 \\ * & \xrightarrow{z_1} & Z. \end{array}$$

These two diagrams are homotopy equivalent, since the maps $z_0, z_1: * \rightarrow Z$ are homotopic via p , yet their pullbacks $*$ and $\emptyset$ are not homotopy equivalent (there is not even a map $* \rightarrow \emptyset$).

We may get around the lack of homotopy invariance of pushouts and pullbacks by replacing them by their homotopy invariant analogues, called *homotopy pushouts* and *homotopy pullbacks*. For the convenience of the reader we recall their definitions.

Definition 2.1.3. Consider maps $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ of topological spaces. Their *homotopy pushout* $X \sqcup_Z^h Y$ (or *double mapping cylinder*) is defined as

$$X \sqcup_Z^h Y := X \sqcup_Z (Z \times [0, 1]) \sqcup_Z Y,$$

where the right-hand side is the quotient of $X \sqcup (Z \times [0, 1]) \sqcup Y$ by the equivalence relation generated by $(z, 0) \sim f(z)$ and $(z, 1) \sim g(z)$ for all $z \in Z$. We denote by $\iota_X: X \hookrightarrow X \sqcup_Z^h Y$ and $\iota_Y: Y \hookrightarrow X \sqcup_Z^h Y$ the canonical inclusions. There is also a canonical homotopy

$$H_{\text{can}}: \iota_X \circ f \sim \iota_Y \circ g,$$

represented by the map $Z \times [0, 1] \rightarrow X \sqcup_Z^h Y$ sending (z, t) to $[(z, t)]$.

This construction aligns with the Fundamental Principle: rather than forcing a strict equality $f(z) = g(z)$ for all $z \in Z$, like in the strict pushout, it instead introduces a path $f(z) \rightsquigarrow g(z)$ witnessing their identification up to homotopy. As a result, $X \sqcup_Z^h Y$ classifies *homotopy commutative squares*: continuous maps from $X \sqcup_Z^h Y$ into some other space T are given by two maps $t: X \rightarrow T$ and $s: Y \rightarrow T$ together with a homotopy $H: s \circ f \sim t \circ g$:

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ f \downarrow & H & \downarrow s \\ X & \xrightarrow{t} & T. \end{array}$$

The homotopy pushout $X \sqcup_Z^h Y$ is illustrated in Figure 2.1.




Figure 2.1: The homotopy pushout $X \sqcup_Z^h Y$, obtained by gluing the cylinder $Z \times [0, 1]$ to X at one end and to Y at the other.

Exercise 2.1.4. Show that the homotopy type of $X \sqcup_Z^h Y$ depends only on the homotopy types of X, Y, Z and the homotopy classes of f, g . In more detail, consider a homotopy commutative diagram of topological spaces and continuous maps of the form

$$\begin{array}{ccccc} X & \xleftarrow{f} & Z & \xrightarrow{g} & Y \\ \alpha \downarrow & H & \downarrow \gamma & K & \downarrow \beta \\ X' & \xleftarrow{f'} & Z' & \xrightarrow{g'} & Y', \end{array}$$

where $H: \alpha \circ f \sim f' \circ \gamma$ and $K: \beta \circ g \sim g' \circ \gamma$ are homotopies. Construct a map $\alpha \sqcup_Y^h \beta: X \sqcup_Z^h Y \rightarrow X' \sqcup_{Z'}^h Y'$, and show it is a homotopy equivalence whenever α, β and γ are homotopy equivalences.

Let us mention the two most important examples of homotopy pushouts, whose ∞ -categorical incarnations will recur throughout the entire book:

- For a continuous map $f: X \rightarrow Y$, we define its (unreduced) *homotopy cofiber* $C(f)$ as the homotopy pushout of f with $X \rightarrow *$. As a special case, we have the *cone* $C(X) := C(\text{id}_X)$ of a topological space X .
- For a topological space X , we define its (unreduced) *suspension* SX as the homotopy pushout of $X \rightarrow *$ with $X \rightarrow *$.

Both are illustrated in Figure 2.2.




Figure 2.2: The homotopy cofiber $C(f) = Y \sqcup_X^h *$ of a map $f: X \rightarrow Y$ (left), and the suspension $SX = * \sqcup_X^h *$ of a space X (right).

Remark 2.1.5. In practice, we will most be interested in the *reduced* variants of each of these constructions: if X , Y and Z are pointed topological spaces and f and g preserve the basepoints, then the reduced homotopy pushout is the quotient of $X \sqcup_Z^h Y$ in which we demand that $(z_0, t) \simeq (z_0, 0)$ for all $t \in [0, 1]$, where $z_0 \in Z$ is the basepoint of Z . The reduced form of the homotopy cofiber and suspension are denoted by $\tilde{C}(f)$ and ΣX , respectively.

The definition of homotopy pullbacks is dual.

Definition 2.1.6. Consider maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ of topological spaces. We define their *homotopy pullback* $X \times_Z^h Y$ as the subspace of the product $X \times Z^{[0,1]} \times Y$ given as

$$X \times_Z^h Y := \{(x, p, y) \in X \times Z^{[0,1]} \times Y \mid p(0) = f(x), p(1) = g(y)\}.$$

Here $Z^{[0,1]} = \text{Map}([0, 1], Z)$ is the space of paths in Z with the compact-open topology, which in this case is completely determined by the requirement that a map $T \rightarrow Z^{[0,1]}$ is continuous if and only if the curried map $T \times [0, 1] \rightarrow Z$ is continuous.

Again, note how the homotopy pullback adheres to the spirit of the Fundamental Principle: for a point in the homotopy pullback, we do not require $f(x)$ and $g(y)$ to be strictly equal; instead, we require the additional data of a path p from $f(x)$ to $g(y)$ that expresses that these points are ‘equal up to homotopy’. As a result, a map $T \rightarrow X \times_Z^h Y$ into the homotopy pullback corresponds to a pair of maps $T \rightarrow X$ and $T \rightarrow Y$ together with a homotopy between the two composites $T \rightarrow Z$:

$$\begin{array}{ccc} T & \xrightarrow{t} & Y \\ s \downarrow & H & \downarrow g \\ X & \xrightarrow{f} & Z. \end{array}$$

Exercise 2.1.7. Formulate and prove the homotopy invariance of homotopy pullbacks, dual to Exercise 2.1.4.

Two important examples of homotopy pullbacks:

- Given a map $f: X \rightarrow Y$, the *homotopy fiber* of f at a point $y \in Y$ is the homotopy pullback of f along the map $y: * \rightarrow Y$. Its elements consist of a point $x \in X$ equipped with a path $p: f(x) \rightsquigarrow y$.
- The *loop space* ΩX of a pointed space (X, x) is the homotopy pullback of the map $x: * \rightarrow X$ along itself. Its elements are x -centered loops $p: x \rightsquigarrow x$.

While the homotopy invariance of homotopy pushouts and pullbacks ensures that the problems illustrated by Example 2.1.1 and Example 2.1.2 do not occur, it is not immediately clear that the localization functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ turns homotopy pushouts and pullbacks into ∞ -categorical pushouts and pullbacks. In the next section, we provide the tools to tackle this problem.

2.2 Computing pushouts and pullbacks in localizations

The problem of computing pushouts and pullbacks of anima in terms of pushouts and pullbacks of topological spaces is a special case of a problem that appears in other situations as well: given an ∞ -category C equipped with a collection of morphisms W designated as weak equivalences, how can pushouts and pullbacks in the localization $C[W^{-1}]$ be computed in terms of pushouts and pullbacks in the original ∞ -category C ? The goal of this section is to introduce a general framework for addressing this question, due to Cisinski [Cis19, Chapter 7] and closely related to Quillen’s framework of *model categories* [Qui67]. In the case of pullbacks, the idea is to find a suitable class of morphisms in C called *fibrations* with the property that the localization functor $C \rightarrow C[W^{-1}]$ preserves pullbacks along fibrations. In other words: strict pullbacks along fibrations in C are already the ‘homotopically correct’ pullbacks. Dually, pushouts in $C[W^{-1}]$ may be computed by finding a class of *cofibrations* in C and forming strict pushouts along cofibrations.

Remark 2.2.1. Model category theory has played an important role in the development of homotopy theory, being one of the main theoretical frameworks researchers used for studying homotopy theories before the widespread adoption of ∞ -category theory. A model category is a 1-category equipped with weak equivalences, fibrations, and cofibrations satisfying various axioms that ensure these classes interact appropriately. From an ∞ -categorical perspective, we may think of a model category C as providing a ‘model’ for its associated ∞ -categorical localization $C[W^{-1}]$. In particular, the fibrations and cofibrations in a model category do not affect the ‘underlying homotopy theory’ presented by the model category, and should rather be thought of as tools for gaining control over categorical constructions like pushouts and pullbacks in this localization.

Remark 2.2.2. This section can be skipped on a first reading. The only results we will use are the specific consequences for the localization functor $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ stated in Theorems 2.3.8 and 2.3.18; readers willing to accept those as black boxes may safely move to Section 2.3.

Definition 2.2.3 (Class of fibrations). Let C be an ∞ -category with a terminal object $*$. A collection of morphisms $F \subseteq \mathbf{Map}([1], C)$ is called a *class of fibrations* if the following conditions are satisfied:

- (1) It contains all the isomorphisms;
- (2) It is closed under composition;
- (3) It is closed under base change: for all morphisms $f: X \rightarrow Y$ in F and $v: Y' \rightarrow Y$ in C , the pullback $Y' \times_Y X$ exists, and the projection $Y' \times_Y X \rightarrow Y'$ is again in F .

A morphism in F is called a *fibration*. An object X in C is called *fibrant* if the unique map $X \rightarrow *$ is a fibration. Dually, a collection of morphisms I is called a *class of cofibrations* if the corresponding class I^{op} in C^{op} defines a class of fibrations in C^{op} . A morphism in I is a *cofibration*, and an object X is *cofibrant* if $\emptyset \rightarrow X$ is a cofibration.

Definition 2.2.4 (∞ -category with weak equivalences and fibrations). An ∞ -category with *weak equivalences and fibrations* is a triple (C, W, F) , where C is an ∞ -category with a terminal object, and W and F are collections of morphisms satisfying the following conditions:

- (1) (2-out-of-3) For composable morphisms f and g , if two of f, g, gf are in W , then so is the third.
- (2) (Fibrations) The class F is a class of fibrations.
- (3) (Pullback axiom) For any pullback square

$$\begin{array}{ccc} X' & \xrightarrow{u} & X \\ p' \downarrow & \lrcorner & \downarrow p \\ Y' & \xrightarrow{v} & Y \end{array}$$

in which p is a fibration between fibrant objects, and Y' is fibrant, if p belongs to W then so does p' .

- (4) (Factorization axiom) For any morphism $f: X \rightarrow Y$ in C with fibrant codomain, there exists a morphism $w: X \rightarrow X'$ in W and a fibration $p: X' \rightarrow Y$ such that $f = p \circ w$.

The morphisms in W are called *weak equivalences*. Morphisms in $W \cap F$ are *trivial fibrations*. We similarly define an ∞ -category with *weak equivalences and cofibrations* by dualizing (replacing F with a class of cofibrations I and appropriately modifying axioms (3) and (4) to involve pushouts and cofibrant objects).

Example 2.2.5. Let C be an ∞ -category that admits a terminal object and admits all pullbacks. Then we may turn C into an ∞ -category with weak equivalences and fibrations by taking W to consist only of the isomorphisms and F to consist of all morphisms in C .

Example 2.2.6 (Structures on Top). As will be discussed below, the category Top can be equipped with such structures relevant to $\text{An} \simeq \text{Top}[W^{-1}]$ where W is the class of weak homotopy equivalences:

- (Top, W, I = relative cell complexes) is an ∞ -category with weak equivalences and cofibrations (see Theorem 2.3.8). The cofibrant objects are the cell complexes.
- (Top, W, F = Serre fibrations) is an ∞ -category with weak equivalences and fibrations (see Theorem 2.3.18). All objects in Top are fibrant in this structure.

The following definition captures those functors that preserve the relevant structure:

Definition 2.2.7 (Left exact functor). Let (C, W_C, F_C) and (D, W_D, F_D) be ∞ -categories with weak equivalences and fibrations. A functor $F: C \rightarrow D$ is called *left exact* if:

- (1) F preserves terminal objects.
- (2) F sends fibrations between fibrant objects in C to fibrations in D , and trivial fibrations between fibrant objects in C to trivial fibrations in D .
- (3) For any pullback square in C

$$\begin{array}{ccc} X' & \xrightarrow{u} & X \\ p' \downarrow & \lrcorner & \downarrow p \\ Y' & \xrightarrow{v} & Y \end{array}$$

in which $p \in F_C$ and Y, Y' are fibrant in C , the induced square

$$\begin{array}{ccc} F(X') & \xrightarrow{F(u)} & F(X) \\ F(p') \downarrow & \lrcorner & \downarrow F(p) \\ F(Y') & \xrightarrow{F(v)} & F(Y) \end{array}$$

is a pullback square in D .

If C, D have weak equivalences and cofibrations, we say that $F: C \rightarrow D$ is *right exact* if $F^{\text{op}}: C^{\text{op}} \rightarrow D^{\text{op}}$ is left exact (relative to the dual fibration structures). In particular, F preserves initial objects and pushouts along cofibrations between cofibrant objects, provided that the other object in the span is also cofibrant.

We now come to the main result of this section, which we will state as a black box:

Theorem 2.2.8 (Cisinski [Cis19, Proposition 7.5.6, Theorem 7.5.18]). *Let (C, W, F) be an ∞ -category with weak equivalences and fibrations, let $C_f \subseteq C$ be the full subcategory of fibrant objects, and write W_f for the weak equivalences between fibrant objects.*

- (1) *The inclusion $\iota: C_f \hookrightarrow C$ induces an equivalence*

$$\bar{\iota}: C_f[W_f^{-1}] \xrightarrow{\sim} C[W^{-1}].$$

- (2) *The localization $C[W^{-1}]$ has finite limits (i.e., a terminal object and pullbacks), and the localization functor $\gamma: C \rightarrow C[W^{-1}]$ is left exact (when $C[W^{-1}]$ is given the trivial structure as in Example 2.2.5).*

- (3) *For any ∞ -category D with finite limits, composition with γ induces an equivalence*

$$\text{Fun}^{\text{lex}}(C[W^{-1}], D) \xrightarrow{\sim} \text{Fun}_W^{\text{lex}}(C, D)$$

between left exact functors $C[W^{-1}] \rightarrow D$ and left exact functors $C \rightarrow D$ that invert the morphisms in W . In particular, every left exact functor $C \rightarrow D$ which inverts the morphisms W induces a left exact functor $C[W^{-1}] \rightarrow D$.

Corollary 2.2.9. *Let (C, W, I) be an ∞ -category with weak equivalences and cofibrations, let $C_c \subseteq C$ be the full subcategory of cofibrant objects, and write W_c for the weak equivalences between cofibrant objects.*

(1) *The inclusion $\iota: C_c \hookrightarrow C$ induces an equivalence*

$$\bar{\iota}: C_c[W_c^{-1}] \xrightarrow{\cong} C[W^{-1}].$$

(2) *The localization $C[W^{-1}]$ has finite colimits (i.e., an initial object and pushouts), and the localization functor $\gamma: C \rightarrow C[W^{-1}]$ is right exact.*

(3) *For any ∞ -category D with finite colimits, composition with γ induces an equivalence*

$$\mathrm{Fun}^{\mathrm{rex}}(C[W^{-1}], D) \xrightarrow{\sim} \mathrm{Fun}_W^{\mathrm{rex}}(C, D).$$

Proof. Apply Theorem 2.2.8 to the opposite category C^{op} . □

Thus the localization of an ∞ -category with weak equivalences and fibrations (respectively, cofibrations) may be computed using only its fibrant (respectively, cofibrant) objects, without choosing replacements functorially.

The exactness results above cover the interaction of localizations with pushouts and pullbacks. We will now extend this discussion to arbitrary limits and colimits, following Cisinski [Cis19, Section 7.7]

Definition 2.2.10. Let C be an ∞ -category with weak equivalences and fibrations. We say that C is *homotopy complete* if the following two conditions are satisfied:

- (1) The product $\prod_{i \in I} X_i$ of any small collection of fibrant objects $(X_i)_{i \in I}$ exists in C and is again fibrant;
- (2) Given a small collection $(f_i: X_i \rightarrow Y_i)$ of morphisms between fibrant objects, if each f_i is a fibration (resp. trivial fibration), then also $\prod_i f_i$ is a fibration (resp. trivial fibration).

A functor between homotopy complete ∞ -categories with weak equivalences and fibrations is called *homotopy continuous* if it is left exact and preserves small products of fibrant objects.

Dually, we say an ∞ -category with weak equivalences and cofibrations C is *homotopy cocomplete* if C^{op} is homotopy complete; we obtain the analogous notion of a *homotopy cocontinuous* functor.

Theorem 2.2.11 (Cisinski [Cis19, Proposition 7.7.4, Theorem 7.7.6]). *Let C be a homotopy complete ∞ -category with weak equivalences and fibrations, and write $L(C) := C[W^{-1}]$ for its localization.*

- (1) *The localization $L(C)$ has small limits, and the localization functor $\gamma: C \rightarrow L(C)$ is homotopy continuous;*
- (2) *If all objects of C are fibrant, then $L(C)$ is in fact universal with this property: for any ∞ -category E with small limits, the functor γ induces an equivalence of ∞ -categories*

$$\gamma^*: \mathrm{Fun}^{\mathrm{lim}}(L(C), E) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{hcont}}(C, E),$$

where the right-hand side denotes the full subcategory spanned by the homotopy continuous functors.

- (3) Given a homotopy continuous functor $F: C \rightarrow D$, the induced functor $L(F): L(C) \rightarrow L(D)$ preserves small limits;

The dual result holds for homotopy cocomplete ∞ -categories with weak equivalences and cofibrations.

2.3 Relative cell complexes and Serre fibrations

Having established a general toolkit for understanding limits and colimits in localizations, we return our attention to the localization functor $\Pi_\infty: \text{Top} \rightarrow \text{An}$. In order to apply the theorems from the previous section, we need to equip Top with suitable notions of fibrations and cofibrations. The standard choice, originating from Quillen's work [Qui67], uses Serre fibrations and relative cell complexes. We will recall the relevant definitions and statements.

2.3.1 Relative cell complexes

We start with the cofibrations.

Definition 2.3.1. A continuous map $i: A \rightarrow X$ is called a *relative cell complex* if it is isomorphic under A to a map obtained by iteratively attaching cells: there exists an ordinal λ , a sequence $(X_\alpha)_{\alpha \leq \lambda}$ and a homeomorphism $X_\lambda \cong X$ under A such that

- $X_0 = A$;
- For $\alpha < \lambda$, $X_{\alpha+1}$ is formed from X_α via a pushout square

$$\begin{array}{ccc} \bigsqcup_{j \in I_\alpha} S^{n_j-1} & \longrightarrow & X_\alpha \\ \downarrow & & \downarrow \\ \bigsqcup_{j \in I_\alpha} D^{n_j} & \longrightarrow & X_{\alpha+1}, \end{array}$$

for some index set I_α and dimensions $n_j \geq 0$.

- For limit ordinals $\beta \leq \lambda$, $X_\beta = \text{colim}_{\alpha < \beta} X_\alpha$.

A topological space X is a *cell complex* if the map $\emptyset \rightarrow X$ is a relative cell complex. We denote by $\text{Cell} \subseteq \text{Top}$ the full subcategory spanned by the cell complexes.

Remark 2.3.2. While the definition involves ordinals to handle potentially transfinite constructions, for many common examples the process involves only a finite or countable number of steps. For example, every cell complex is homotopy equivalent to a *CW-complex*, which is a cell complex where $\lambda = \omega$ is the set of natural numbers, and where the dimensions of the cells attached at step $n \in \omega$ is n .

Cell complexes play an important role in homotopy theory, due to the following two important theorems:

Theorem 2.3.3 (Whitehead's theorem, [Hat02, Theorem 4.5]). *Let $f: X \rightarrow Y$ be a weak homotopy equivalence between spaces having the homotopy type of cell complexes. Then f is a homotopy equivalence.*

Proof. Choose homotopy equivalences $u: X' \rightarrow X$ and $v: Y' \rightarrow Y$ with X' and Y' cell complexes, and let $w: Y \rightarrow Y'$ be a homotopy inverse to v . The composite $wfu: X' \rightarrow Y'$ is a weak homotopy equivalence, hence a homotopy equivalence by the usual Whitehead theorem. Since u and w are homotopy equivalences, so is f . $\square$

Theorem 2.3.4 (CW-approximation, [Hir15, Theorem 2.1]). *For every topological space X , there exists a cell complex Z equipped with a weak homotopy equivalence $q: Z \xrightarrow{\sim} X$. Moreover, Z and q can be taken to be functorial in X .*

Remark 2.3.5. An explicit construction of such a functorial CW-approximation may be given as follows. Recall from Definition 1.6.9 that the singular complex functor $\text{Sing}: \text{Top} \rightarrow \text{sSet}$ admits the geometric realization functor $|-|: \text{sSet} \rightarrow \text{Top}$ as a left adjoint. One can prove that the geometric realization $|K|$ is a CW-complex for every simplicial set K . Furthermore, for any topological space X the counit $|\text{Sing}(X)| \rightarrow X$ of the adjunction is a weak homotopy equivalence.

As a consequence of these two theorems, animae are cell complexes up to homotopy equivalence:

Proposition 2.3.6. *The functor $\Pi_\infty: \text{Cell} \rightarrow \text{An}$ induces an equivalence*

$$\text{Cell}[\{\text{homotopy equivalences}\}^{-1}] \xrightarrow{\sim} \text{An}.$$

Proof. In light of the homotopy hypothesis, Axiom M, it suffices to show that the functor

$$\text{Cell}[\{\text{homotopy equivalences}\}^{-1}] \rightarrow \text{Top}[\{\text{weak homotopy equivalences}\}^{-1}]$$

induced by the inclusion is an equivalence of ∞ -categories. By Theorem 2.3.4 there exists a functor $Z: \text{Top} \rightarrow \text{Cell}$ equipped with a natural transformation $q: Z \Rightarrow \text{id}_{\text{Top}}$ such that the map $q_X: Z(X) \rightarrow X$ is a weak homotopy equivalence for all X . Given a weak homotopy equivalence $f: X \rightarrow Y$, the 2-out-of-3 property applied to the diagram

$$\begin{array}{ccc} Z(X) & \xrightarrow{q_X} & X \\ Z(f) \downarrow & & \downarrow f \\ Z(Y) & \xrightarrow{q_Y} & Y \end{array}$$

shows that the map $Z(f)$ is again a weak homotopy equivalence, and by Whitehead's theorem it is in fact a homotopy equivalence. It follows that Z induces a functor $\text{Top}[\{\text{weak homotopy equivalences}\}^{-1}] \rightarrow \text{Cell}[\{\text{homotopy equivalences}\}^{-1}]$, and the natural transformation $q: Z \rightarrow \text{id}$ exhibits this as the requisite inverse functor. $\square$

Corollary 2.3.7. *The homotopy category $\text{Ho}(\text{An})$ is equivalent to the category hCell whose objects are the cell complexes and whose morphism set between X and Y is the set $[X, Y]$ of homotopy classes of continuous maps $X \rightarrow Y$.*

Proof. It suffices to observe that a functor $F: \text{Cell} \rightarrow C$ into a 1-category inverts homotopy equivalences if and only if it identifies homotopic maps. One implication follows by applying F to homotopy inverses. Conversely, if $H: f \sim g$ is a homotopy, then

$$F(f) = F(H)F(i_0) = F(H)F(\text{pr})^{-1} = F(H)F(i_1) = F(g),$$

where $i_0, i_1: X \rightarrow X \times [0, 1]$ are the endpoint inclusions and $\text{pr}: X \times [0, 1] \rightarrow X$ is the projection. Thus hCell has the same universal property as the homotopy category of the localization in Proposition 2.3.6. $\square$

We are now ready to establish the claimed structure on Top :

Theorem 2.3.8 (cf. Hovey [Hov99, Theorem 2.4.19]). *Let W_{whe} be the collection of weak homotopy equivalences in Top , and let I_{cell} be the collection of relative cell complexes. Then $(\text{Top}, W_{\text{whe}}, I_{\text{cell}})$ is an ∞ -category with weak equivalences and cofibrations. Furthermore, it admits functorial factorizations and is homotopy cocomplete.*

Proof sketch. We need to verify the axioms from Definition 2.2.4 (dualized for cofibrations):

- (1) (2-out-of-3): It is clear from the definition that the class W_{whe} satisfies the 2-out-of-3 property.
- (2) (Class of cofibrations): The class I_{cell} contains all isomorphisms (by attaching no cells) and is closed under composition (attaching cells iteratively). It is also closed under pushouts: if $i: A \rightarrow X$ is a relative cell complex and $f: A \rightarrow A'$ is any map, consider the pushout square

$$\begin{array}{ccc} A & \xrightarrow{f} & A' \\ i \downarrow & \lrcorner & \downarrow i' \\ X & \xrightarrow{f'} & X'. \end{array}$$

Then i' is also a relative cell complex, where the filtration is inductively defined by $X'_{\lambda+1} := X_{\lambda+1} \sqcup_{X_\lambda} X'_\lambda$.

- (3) (Pushout axiom): We need to show that if $i: A \rightarrow X$ is a trivial cofibration (i.e., a relative cell complex and a weak homotopy equivalence) between cofibrant objects (cell complexes), then the pushout $i': A' \rightarrow X'$ is also a weak homotopy equivalence. This is a non-trivial fact in model category theory. One first shows that every trivial cofibration has the left lifting property with respect to the Serre fibrations (see Definition 2.3.12 below), then observes that this property is preserved by pushouts, and finally deduces that every map with this lifting property is a weak homotopy equivalence. See Hovey [Hov99, Section 2.4].
- (4) (Factorization axiom): The small object argument functorially factors every map $f: X \rightarrow Y$ as $X \xrightarrow{i} Z \xrightarrow{p} Y$, where i is a relative cell complex and p is a trivial Serre fibration, hence in particular a weak homotopy equivalence. See Hovey [Hov99, Sections 2.1 and 2.4].

Since all axioms are satisfied, $(\text{Top}, W_{\text{whe}}, I_{\text{cell}})$ is an ∞ -category with weak equivalences and cofibrations (with functorial factorizations). It is straightforward to see that a small coproduct of (trivial) cofibrations is again a (trivial) cofibration, so that Top is in fact homotopy cocomplete. $\square$

Proposition 2.3.9. *The functor $\Pi_\infty: \text{Top} \rightarrow \text{An}$ sends every pushout square in Top whose vertices are cell complexes and one of whose maps is a relative cell complex to a pushout square in An .*

Proof. This follows directly from Theorem 2.3.8 and the right exactness of localization functors stated in Corollary 2.2.9. $\square$

We now arrive at the main result of this section.

Proposition 2.3.10. *Let $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ be continuous maps, where all three spaces have the homotopy type of a cell complex. The canonical homotopy H_{can} from Definition 2.1.3 determines a commutative square of animae*

$$\begin{array}{ccc} \Pi_\infty(Z) & \xrightarrow{\Pi_\infty(g)} & \Pi_\infty(Y) \\ \Pi_\infty(f) \downarrow & & \downarrow \Pi_\infty(\iota_Y) \\ \Pi_\infty(X) & \xrightarrow{\Pi_\infty(\iota_X)} & \Pi_\infty(X \sqcup_Z^h Y). \end{array}$$

and this square is a pushout.

Proof. Consider the mapping cylinder of the map f :

$$M(f) := (X \sqcup (Z \times [0, 1])) / (z, 0) \sim f(z).$$

The map f factors as $Z \xrightarrow{i} M(f) \xrightarrow{p} X$, where $i(z) := [(z, 1)]$, where $p|_X = \text{id}_X$ and where $p|_{Z \times [0, 1]}(z, t) := f(z)$. Observe that the map $p: M(f) \rightarrow X$ is a homotopy equivalence, with homotopy inverse sending x to $[x]$. In particular, it induces an equivalence $\Pi_\infty(M(f)) \xrightarrow{\sim} \Pi_\infty(X)$ of animae. Further observe that the homotopy pushout $X \sqcup_Z^h Y$ can be identified with the strict pushout $M(f) \sqcup_Z Y$ via the maps g and the projection $M(f) \rightarrow X$:

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ i \downarrow & \lrcorner & \downarrow \\ M(f) & \longrightarrow & X \sqcup_Z^h Y. \end{array} \quad (2.1)$$

We will show that the functor $\Pi_\infty: \text{Top} \rightarrow \text{An}$ preserves this pushout square.

Case 1: Assume first that the spaces X, Y and Z are cell complexes and that the map f is a relative cell complex. In this case, the map $i: Z \rightarrow M(f)$ is also a relative cell complex: it may be written as the composite of $Z \times \{1\} \hookrightarrow Z \times [0, 1]$, which is a relative cell complex by the product cell filtration, and the map $Z \times [0, 1] \rightarrow M(f)$, which is a pushout of the map $Z \rightarrow X$ and hence also a relative cell complex. The fact that $\Pi_\infty(-)$ preserves the pushout square in Equation (2.1) is now an instance of Proposition 2.3.9.

Case 2: We will now prove the general case. By Theorem 2.3.4, there exists a cell complex Z' and a weak homotopy equivalence $Z' \xrightarrow{\sim} Z$. By factorizing the composite maps $Z' \rightarrow X$ and $Z' \rightarrow Y$ into a relative cell complex followed by a weak equivalence, we obtain a commutative diagram

$$\begin{array}{ccccc} X' & \xleftarrow{f'} & Z' & \xrightarrow{g'} & Y' \\ \sim \downarrow & & \downarrow \sim & & \downarrow \sim \\ X & \xleftarrow{f} & Z & \xrightarrow{g} & Y, \end{array}$$

where X' , Y' and Z' are cell complexes, the vertical maps are weak homotopy equivalences, and f' and g' are relative cell complexes. Since all six spaces have the homotopy type of a cell complex, it follows from Whitehead's theorem that the vertical maps are even homotopy equivalences. By homotopy invariance of homotopy pushouts, Exercise 2.1.4, the induced map $X' \sqcup_{Z'}^h Y' \rightarrow X \sqcup_Z^h Y$ is again a homotopy equivalence, and in particular it becomes an equivalence after applying $\Pi_\infty(-)$. This means that we have reduced the problem to Case 1, thus finishing the proof. $\square$

We next record when an actual quotient computes a homotopy cofiber. Recall that a pair (Y, A) consisting of a closed subspace $A \subseteq Y$ is called a *neighborhood deformation retract pair*, or *NDR-pair*, if there are a continuous map $u: Y \rightarrow [0, 1]$ and a homotopy $h: Y \times [0, 1] \rightarrow Y$ such that

$$u^{-1}(0) = A, \quad h(y, 0) = y, \quad h(a, t) = a,$$

and $h(y, 1) \in A$ whenever $u(y) < 1$. An inclusion $A \hookrightarrow Y$ is called *collared* if there is an open neighborhood U of A in Y and a homeomorphism

$$A \times [0, 1) \xrightarrow{\cong} U$$

whose restriction to $A \times \{0\}$ is the given inclusion. Every collared inclusion is an NDR-pair. More generally, the NDR condition is equivalent to the homotopy extension property; see May [May99a, Chapter 6, Section 4].

Corollary 2.3.11. *Let (Y, A) be an NDR-pair with A nonempty, and assume that A and Y have the homotopy type of cell complexes. Then the quotient map $Y \rightarrow Y/A$ induces an isomorphism*

$$\Pi_\infty(Y) \sqcup_{\Pi_\infty(A)} * \cong \Pi_\infty(Y/A)$$

in $\mathbf{An}_*$. In particular, this applies whenever the inclusion $A \hookrightarrow Y$ is collared.

Proof. The NDR condition implies that the canonical map from the homotopy cofiber of $A \hookrightarrow Y$ to the actual quotient Y/A is a homotopy equivalence; see May [May99a, Chapter 8, Section 4]. The claim now follows by applying Proposition 2.3.10 to the diagram $Y \leftarrow A \rightarrow *$. $\square$

2.3.2 Serre fibrations

We now turn to the fibration structure on $\mathbf{Top}$.

Definition 2.3.12. Let $p: E \rightarrow B$ be a continuous map of topological spaces. We say that p is a *Serre fibration* if it satisfies the homotopy lifting property with respect to all disks: for every $n \geq 0$ and every solid commutative diagram of the form

$$\begin{array}{ccc} D^n \times \{0\} & \xrightarrow{f} & E \\ i \downarrow & \nearrow \bar{H} & \downarrow p \\ D^n \times [0, 1] & \xrightarrow{H} & B, \end{array}$$

there exists a dashed morphism making the diagram commute.

We state without proof some standard facts about Serre fibrations:

Theorem 2.3.13 (Locality for Serre fibrations, tom Dieck [tDie08, Theorem 6.3.3]). *Let $p: E \rightarrow B$ be a continuous map, and assume that there exists an open cover $\{U_i\}_{i \in I}$ such that the restriction $p|_{U_i}: p^{-1}(U_i) \rightarrow U_i$ is a Serre fibration for each $i \in I$. Then p is itself a Serre fibration.*

In particular, locally trivial fiber bundles are Serre fibrations.

Theorem 2.3.14 (Long exact sequence of homotopy groups, Switzer [Swi75, Theorem 2.59]). *Let $p: E \rightarrow B$ be a Serre fibration and consider a point $e \in E$ with image $b := p(e)$. Define the fiber F_b as the following (strict) pullback of topological spaces:*

$$\begin{array}{ccc} F_b & \xhookrightarrow{i} & E \\ \downarrow & \lrcorner & \downarrow p \\ \text{pt} & \xhookrightarrow{b} & B. \end{array}$$

Then there is a long exact sequence of homotopy groups of the form

$$\cdots \rightarrow \pi_{n+1}(B, b) \rightarrow \pi_n(F_b, e) \xrightarrow{i_*} \pi_n(E, e) \xrightarrow{p_*} \pi_n(B, b) \rightarrow \pi_{n-1}(F_b, e) \rightarrow \cdots$$

Corollary 2.3.15. *A Serre fibration $p: E \rightarrow B$ is a weak homotopy equivalence if and only if the fiber F_b is weakly contractible for every $b \in B$.*

Proof. By working separately over each path component of B , we may assume that B is path-connected. Picking a preferred basepoint b_0 of B , it then suffices to show that p is a weak homotopy equivalence if and only if the fiber $F := F_{b_0}$ is weakly contractible. This follows immediately from the long exact sequence from Theorem 2.3.14. $\square$

A general source of Serre fibrations is the path space construction.

Construction 2.3.16. We define the *mapping path space* $P(f)$ of a continuous map $f: E \rightarrow B$ as

$$P(f) := \{(e, \gamma) \in E \times B^{[0,1]} \mid f(e) = \gamma(0)\}.$$

Lemma 2.3.17. *The projection $p: P(f) \rightarrow B$ given by $(e, \gamma) \mapsto \gamma(1)$ is a Serre fibration.*

Proof. Consider a solid commutative diagram of the form

$$\begin{array}{ccc} D^n \times \{0\} & \xrightarrow{g} & P(f) \\ i \downarrow & \nearrow \overline{H} & \downarrow p \\ D^n \times [0, 1] & \xrightarrow{H} & B. \end{array}$$

We may define the dashed map $\overline{H}$ as $\overline{H}(x, s) := (e(x), K(x, s, -))$, where

- The map $e: D^n \rightarrow E$ is the first component of g ;
- The map $K: D^n \times [0, 1] \times [0, 1] \rightarrow B$ is a continuous map satisfying the following three constraints:
 - To guarantee that $\overline{H}(x, s) \in P(f)$ we need $K(x, s, 0) = f(e(x))$;

- To guarantee commutativity of the top triangle, we need $K(x, 0, t) = \gamma(x)(t)$, where $\gamma(x)$ is the second component of $g(x)$;
- To guarantee commutativity of the bottom triangle, we need $K(x, s, 1) = H(x, s)$.

Such a continuous map exists, since the inclusion $([0, 1] \times \{0, 1\}) \cup (\{0\} \times [0, 1]) \hookrightarrow [0, 1] \times [0, 1]$ has a continuous retraction. $\square$

Theorem 2.3.18 (cf. Hovey [Hov99, Theorem 2.4.19]). *Consider the category Top of topological spaces, let F_{Serre} denote the collection of Serre fibrations and let W_{whe} denote the collection of weak homotopy equivalences. Then the triple $(\text{Top}, W_{\text{whe}}, F_{\text{Serre}})$ is an ∞ -category with weak equivalences and fibrations. It has functorial factorizations, and is homotopy complete.*

Proof. It is clear that the weak homotopy equivalences satisfy 2-out-of-3, and it is also straightforward to check that Serre fibrations are closed under pullbacks and composition and contain all homeomorphisms. This gives conditions (1) and (2) from Definition 2.2.4.

To see condition (3), consider a pullback square

$$\begin{array}{ccc} E' & \xrightarrow{f} & E \\ p' \downarrow & \lrcorner & \downarrow p \\ B' & \xrightarrow{g} & B \end{array}$$

of topological spaces, where p is both a Serre fibration and a weak homotopy equivalence. We need to show that also p' is a weak homotopy equivalence. By Corollary 2.3.15, it suffices to show that for every point $b' \in B'$ the fiber $F'_{b'} := p'^{-1}(b')$ is weakly contractible. But since p' is a pullback of p , this fiber is homeomorphic to the fiber $F_b = p^{-1}(b)$, which in turn is weakly contractible by Corollary 2.3.15.

Finally, for condition (4), note that a continuous map $f: E \rightarrow B$ may be factored as

$$E \xrightarrow{i} P(f) \xrightarrow{p} B,$$

where p is the path space fibration from Lemma 2.3.17, and where i maps e to $(e, \text{const}_{f(e)})$. The first map is a homotopy equivalence, hence in particular a weak homotopy equivalence, and the second map is a Serre fibration by Lemma 2.3.17. Since the definition of $P(f)$ is functorial in f , this finishes the proof that $(\text{Top}, W_{\text{whe}}, F_{\text{Serre}})$ is a category with weak equivalences and fibrations, with functorial factorizations. It is also clear that a product of Serre fibrations is again a Serre fibration, finishing the proof. $\square$

Proposition 2.3.19. *The localization functor $\Pi_{\infty}(-): \text{Top} \rightarrow \text{An}$ preserves all pullbacks along Serre fibrations.*

Proof. In light of Theorem 2.3.18 this is an instance of Theorem 2.2.8. $\square$

Corollary 2.3.20. *The functor $\Pi_{\infty}: \text{Top} \rightarrow \text{An}$ preserves finite products:*

$$\Pi_{\infty}(X \times Y) \xrightarrow{\sim} \Pi_{\infty}(X) \times \Pi_{\infty}(Y).$$

$\square$

Exercise 2.3.21. Prove the statement of Corollary 2.3.20 directly from the definition of Π_{∞} . *Hint:* Use Lemma 21.6.12.

We are now ready to prove our main application concerning homotopy pullbacks of topological spaces:

Proposition 2.3.22. *The localization functor $\Pi_\infty(-): \mathbf{Top} \rightarrow \mathbf{An}$ sends homotopy pullbacks of topological spaces to pullbacks of anima.*

Proof. Consider continuous maps $f: X \rightarrow Z$ and $g: Y \rightarrow Z$, and consider the pullback square

$$\begin{array}{ccc} X \times_Z^h Y & \longrightarrow & P(f) \\ \downarrow & \lrcorner & \downarrow p \\ Y & \xrightarrow{g} & Z \end{array}$$

defining $X \times_Z^h Y$. The path fibration p is a Serre fibration by Lemma 2.3.17, hence this square is sent to a pullback square in $\mathbf{An}$ by Proposition 2.3.19. The canonical inclusion $i: X \hookrightarrow P(f)$ is a homotopy equivalence, giving an equivalence $\Pi_\infty(P(f)) \simeq \Pi_\infty(X)$. This finishes the proof. $\square$

2.4 Pointed anima, suspensions and loops

The previous sections show that homotopy pushouts and pullbacks of spaces are carried by Π_∞ to pushouts and pullbacks in $\mathbf{An}$. We can therefore define suspensions, cofibers, loop spaces, and fibers directly in the ∞ -category $\mathbf{An}_*$ by their universal properties, and compare them with the classical constructions from algebraic topology. We write $\mathbf{Top}_* := \mathbf{Top}_{*/}$ for the category of pointed topological spaces and basepoint-preserving continuous maps.

Definition 2.4.1 (Pointed anima). A *pointed anima* is a pair (X, x) consisting of an anima X and a distinguished point $x \in X$. A morphism of pointed anima $f: (X, x) \rightarrow (Y, y)$ consists of a map $f: X \rightarrow Y$ in $\mathbf{An}$ together with an isomorphism $f(x) \cong y$.

We denote by $\mathbf{An}_*$ the ∞ -category of pointed anima and basepoint-preserving maps, defined as the slice category $\mathbf{An}_{*/}$ from Definition 21.3.1.

Remark 2.4.2. The forgetful functor $\mathbf{An}_* \rightarrow \mathbf{An}$ preserves pullbacks and pushouts.

Notation 2.4.3. The one-point anima $*$ is canonically pointed; we denote the corresponding pointed anima by $0 \in \mathbf{An}_*$ and call it the *zero object*. One can show it is both initial and terminal in $\mathbf{An}_*$: given pointed anima $X, Y \in \mathbf{An}_*$, there are unique morphisms $X \rightarrow 0$ and $0 \rightarrow Y$. (See Lemma 4.1.9 for more details.) We denote their composite by

$$0: X \rightarrow Y$$

and call it the *null map*.

Definition 2.4.4 (Fiber and cofiber sequences in $\mathbf{An}_*$). A *nullsequence* in $\mathbf{An}_*$ is a sequence of morphisms

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

equipped with a specified *nullhomotopy* $g \circ f \cong 0$. Equivalently, it is a commutative square in $\mathbf{An}_*$ of the form

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow g \\ 0 & \longrightarrow & Z. \end{array}$$

Such a nullsequence is called:

- a *fiber sequence* if this square is a pullback square in $\mathbf{An}_*$ (or, equivalently, in $\mathbf{An}$). In this case we call X the *fiber of g* and write $X \simeq \text{fib}(g)$;
- a *cofiber sequence* if the same square is a pushout square in $\mathbf{An}_*$ (or, equivalently, in $\mathbf{An}$). In this case we call Z the *cofiber of f* and write $Z \simeq \text{cofib}(f)$.

For pointed animae $U, V \in \mathbf{An}_*$, we write

$$[U, V]_* := \pi_0 \text{Hom}_{\mathbf{An}_*}(U, V)$$

for the pointed set of homotopy classes of pointed maps $U \rightarrow V$, whose basepoint is the null map. Recall that a sequence $A \rightarrow B \rightarrow C$ of pointed sets is *exact at B* if the image of $A \rightarrow B$ is the preimage of the basepoint under $B \rightarrow C$.

Lemma 2.4.5. *Let $W \in \mathbf{An}_*$ be a pointed anima.*

(1) *Every cofiber sequence $X \rightarrow Y \rightarrow Z$ induces a fiber sequence of pointed mapping animae*

$$\text{Hom}_{\mathbf{An}_*}(Z, W) \longrightarrow \text{Hom}_{\mathbf{An}_*}(Y, W) \longrightarrow \text{Hom}_{\mathbf{An}_*}(X, W),$$

and hence an exact sequence of pointed sets

$$[Z, W]_* \longrightarrow [Y, W]_* \longrightarrow [X, W]_*.$$

(2) *Every fiber sequence $X \rightarrow Y \rightarrow Z$ induces a fiber sequence of pointed mapping animae*

$$\text{Hom}_{\mathbf{An}_*}(W, X) \longrightarrow \text{Hom}_{\mathbf{An}_*}(W, Y) \longrightarrow \text{Hom}_{\mathbf{An}_*}(W, Z),$$

and hence an exact sequence of pointed sets

$$[W, X]_* \longrightarrow [W, Y]_* \longrightarrow [W, Z]_*.$$

Proof. The functor $\text{Hom}_{\mathbf{An}_*}(-, W): \mathbf{An}_*^{\text{op}} \rightarrow \mathbf{An}$ sends pushout squares to pullback squares. Since $\text{Hom}_{\mathbf{An}_*}(0, W)$ is terminal, applying this functor to the pushout square defining a cofiber sequence shows that the first sequence of mapping animae is a fiber sequence. Dually, the functor $\text{Hom}_{\mathbf{An}_*}(W, -): \mathbf{An}_* \rightarrow \mathbf{An}$ preserves pullbacks and sends 0 to a terminal anima, which proves the corresponding claim for fiber sequences.

It remains to pass to pointed homotopy classes. For any fiber sequence $F \rightarrow E \rightarrow B$ of pointed animae, a component of E lies in the image of $\pi_0(F) \rightarrow \pi_0(E)$ precisely when its image in B lies in the component of the basepoint. Thus $\pi_0(F) \rightarrow \pi_0(E) \rightarrow \pi_0(B)$ is exact as a sequence of pointed sets, proving both claims. $\square$

Using fiber and cofiber sequences, we can define suspension and loop functors on $\mathbf{An}_*$ in a purely ∞ -categorical way.

Definition 2.4.6 (Loop and suspension in $\mathbf{An}_*$). Consider a pointed anima $X \in \mathbf{An}_*$. Its *loop anima* ΩX is defined as the fiber of the unique map $0 \rightarrow X$:

$$\begin{array}{ccc} \Omega X & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & X. \end{array}$$

Its *suspension* ΣX is defined as the cofiber of the unique map $X \rightarrow 0$:

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & \Sigma X. \end{array}$$

Exercise 2.4.7. Show that the assignments $X \mapsto \Omega X$ and $X \mapsto \Sigma X$ define endofunctors

$$\Omega, \Sigma: \mathbf{An}_* \rightarrow \mathbf{An}_*.$$

Hint: Use the pushout description of the diagram $\lrcorner$ from Axiom D to construct a functor $C \rightarrow \mathbf{Fun}(\lrcorner, C)$ given on objects by $X \mapsto (0 \rightarrow X \leftarrow 0)$.

Lemma 2.4.8. The functor $\Sigma: \mathbf{An}_* \rightarrow \mathbf{An}_*$ is left adjoint to $\Omega: \mathbf{An}_* \rightarrow \mathbf{An}_*$: for all $X, Y \in \mathbf{An}_*$ there is a natural equivalence

$$\mathrm{Hom}_{\mathbf{An}_*}(\Sigma X, Y) \simeq \mathrm{Hom}_{\mathbf{An}_*}(X, \Omega Y).$$

Proof. By the definition of ΩY as a pullback, the right-hand side is equivalent to the anima

$$\mathrm{Hom}_{\mathbf{Fun}(\lrcorner, \mathbf{An}_*)}(\mathrm{const}_X, 0 \rightarrow Y \leftarrow 0).$$

As a consequence of Lemma 1.7.13, we see that this is naturally isomorphic to the anima of commutative squares of the form

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & Y, \end{array}$$

i.e. the fiber of the evaluation map $\mathrm{Fun}([1] \times [1], \mathbf{An}_*) \rightarrow (\mathbf{An}_*)^{\times 4}$ at $(X, 0, 0, Y)$. By dualizing this argument, we see that also the left-hand side is naturally isomorphic to this anima, finishing the proof. $\square$

Definition 2.4.9. Let $(X_i)_{i \in I}$ be a collection of pointed animae. Their coproduct in $\mathbf{An}_*$ is called the *wedge* and is denoted by $\bigvee_{i \in I} X_i$. It may be computed by the following pushout square in $\mathbf{An}$:

$$\begin{array}{ccc} \bigsqcup_{i \in I} * & \longrightarrow & \bigsqcup_{i \in I} X_i \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \bigvee_{i \in I} X_i. \end{array}$$

Definition 2.4.10 (Smash product). Let X and Y be pointed anima. Their *smash product* is the pointed anima

$$X \wedge Y := \text{cofib}(X \vee Y \longrightarrow X \times Y),$$

where the map is induced by $(\text{id}_X, 0): X \rightarrow X \times Y$ and $(0, \text{id}_Y): Y \rightarrow X \times Y$. Interchanging the two factors induces a natural *swap map*

$$\sigma_{X,Y}: X \wedge Y \xrightarrow{\cong} Y \wedge X.$$

We now relate these ∞ -categorical constructions in An_* to the classical constructions for pointed topological spaces. Applying Π_∞ to the basepoint defines a functor $\Pi_\infty: \text{Top}_* \rightarrow \text{An}_*$. It follows from Propositions 2.3.10 and 2.3.22 that this functor sends homotopy pullbacks to pullbacks and suitable homotopy pushouts to pushouts.

We will also use the following direct consequence of the construction of Π_∞ .

Lemma 2.4.11. *The functor $\Pi_\infty: \text{Top} \rightarrow \text{An}$ preserves small coproducts.*

Proof. Every continuous map from a topological simplex into a coproduct of spaces lands in a unique summand, since the simplex is connected. Hence the singular complex functor preserves coproducts. The claim follows because $\Pi_\infty(X)$ is the colimit in An of the simplicial set $\text{Sing}(X)$ and colimits commute with colimits. $\square$

In particular:

Proposition 2.4.12. *The functor $\Pi_\infty: \text{Top}_* \rightarrow \text{An}_*$ has the following properties:*

- (1) *Given a pointed map $f: X \rightarrow Y$ between spaces having the homotopy type of cell complexes, the unreduced homotopy cofiber $C(f)$, pointed by its cone point, satisfies $\Pi_\infty(C(f)) \cong \text{cofib}(\Pi_\infty(f))$.*
- (2) *If X has the homotopy type of a cell complex, the unreduced suspension SX , pointed by either cone point, satisfies $\Pi_\infty(SX) \cong \Sigma \Pi_\infty(X)$.*
- (3) *For a collection $(X_i)_{i \in I}$ of pointed spaces having the homotopy type of cell complexes, there is an isomorphism $\Pi_\infty(\bigvee_{i \in I}^h X_i) \cong \bigvee_{i \in I} \Pi_\infty(X_i)$, where the homotopy wedge $\bigvee_i^h X_i$ is the homotopy pushout of the span $\bigsqcup_i X_i \leftarrow \bigsqcup_i * \rightarrow *$.*
- (4) *Given a pointed map $f: X \rightarrow Y$ with homotopy fiber F_f over the basepoint, there is an isomorphism $\Pi_\infty(F_f) \cong \text{fib}(\Pi_\infty(f))$.*
- (5) *The loop space of a pointed space X satisfies $\Pi_\infty(\Omega X) \cong \Omega \Pi_\infty(X)$.*

Proof. Parts (1) and (2) are direct consequences of Proposition 2.3.10, while parts (4) and (5) follow from Proposition 2.3.22. For part (3), combine Proposition 2.3.10 with Lemma 2.4.11 and the pushout description of wedges in Definition 2.4.9. $\square$

Remark 2.4.13. A pointed topological space (X, x_0) is called *well-pointed* if $(X, \{x_0\})$ is an NDR-pair. Under this hypothesis, the familiar reduced models compute the constructions in the preceding proposition. More precisely, for every pointed map $f: X \rightarrow Y$, the collapse maps $C(f) \rightarrow \tilde{C}(f)$

and $SX \rightarrow \Sigma X$ are homotopy equivalences. Likewise, if every X_i is well-pointed, then the strict wedge $\bigvee_i X_i$ computes the homotopy wedge $\bigvee_i^h X_i$. Thus the conclusions of the proposition apply in particular to reduced cofibers, reduced suspensions and strict wedges of pointed cell complexes whose basepoints are 0-cells.

This result motivates the following homotopical definition of spheres in $\mathbf{An}_*$:

Definition 2.4.14. We define the *homotopical n -sphere* $S^n \in \mathbf{An}_*$ inductively by setting $S^0 := * \sqcup *$ (with the first point as basepoint, say), and taking $S^{n+1} := \Sigma(S^n)$.

Corollary 2.4.15. *The functor $\Pi_\infty(-) : \mathbf{Top}_* \rightarrow \mathbf{An}_*$ sends the topological n -sphere $S^n \subseteq \mathbb{R}^{n+1}$ to the homotopical n -sphere S^n .*

Proof. Since Π_∞ preserves coproducts, we have $\Pi_\infty(S^0) \simeq S^0$. The claim now follows inductively from part (2) of Proposition 2.4.12. $\square$

Homotopy groups of animae

We now introduce homotopy groups of animae and prove some basic facts about them, like a version of the Whitehead theorem for animae and the long exact sequence of homotopy groups for a fiber sequence.

Definition 2.4.16 (Path components of an anima). Recall from Example 1.8.14(1) that the inclusion $\mathbf{Set} \hookrightarrow \mathbf{An}$ admits a left adjoint $\pi_0 : \mathbf{An} \rightarrow \mathbf{Set}$, sending an anima X to its *set of path components* $\pi_0(X)$. The unit of the adjunction gives a map of animae $X \rightarrow \pi_0(X)$, and for an element $[x] \in \pi_0(X)$ we define the corresponding path component of X as the fiber of this map over $[x]$. We say that X is *connected* if $\pi_0(X)$ is a singleton.

Definition 2.4.17 (Homotopy groups of an anima). Let $X \in \mathbf{An}_*$ be a pointed anima and $n \geq 0$. The *n -th homotopy group* of X is defined as

$$\pi_n(X) := [S^n, X]_*.$$

We also write $\pi_n(X, x)$ if we wish to make explicit the basepoint x of X .

Since $S^n = \Sigma^n(S^0)$, the adjunction Lemma 2.4.8 implies that

$$\pi_n(X) \cong \pi_0(\Omega^n X),$$

where the functor Ω^n is defined inductively as the n -fold iteration of Ω .

Remark 2.4.18. Let (Y, y_0) be any pointed topological space, and let $X := \Pi_\infty(Y, y_0) \in \mathbf{An}_*$. Then for every $n \geq 0$ there is a natural isomorphism

$$\pi_n(X) = [S^n, X]_* \cong [S^n, Y]_*$$

with the classical n -th homotopy group of (Y, y_0) . Indeed, Proposition 2.4.12 gives natural isomorphisms

$$\Omega^n X \cong \Pi_\infty(\Omega^n Y).$$

The path components of the underlying anima of a topological space agree with its usual path components: after choosing a cell-complex approximation, this follows from Corollary 2.3.7 by mapping out of a point. It follows that

$$\pi_n(X) \cong \pi_0 \Pi_\infty(\Omega^n Y) \cong \pi_0(\Omega^n Y) = [S^n, Y]_*.$$

Remark 2.4.19 (Degree). For every $n \geq 1$, the degree of a self-map of the n -sphere induces an isomorphism

$$\deg: \pi_n(S^n) \xrightarrow{\cong} \mathbb{Z}.$$

Under these isomorphisms, suspension preserves degree. In particular, a pointed self-map of S^n is pointed homotopic to the identity if and only if it has degree 1.

Remark 2.4.20 (The suspension as a cogroup). For every pointed anima X , the suspension ΣX is a *cogroup object* in the homotopy category $\mathrm{Ho}(\mathrm{An}_*)$: there are pointed maps

$$\Sigma X \rightarrow *, \quad \text{pinch}: \Sigma X \rightarrow \Sigma X \vee \Sigma X, \quad i: \Sigma X \rightarrow \Sigma X,$$

satisfying counitality, coassociativity, and the coinverse axiom in $\mathrm{Ho}(\mathrm{An}_*)$. To see this, use the homotopy hypothesis to choose a topological space Y whose underlying anima is isomorphic to the underlying anima of X , and choose a point of Y in the component corresponding to the basepoint of X . Applying the factorization from Theorem 2.3.8 to this basepoint map, we may further assume that Y is a cell complex whose basepoint is a 0-cell. Then $\Sigma X \simeq \Pi_\infty(\Sigma Y)$ by Proposition 2.4.12 and Remark 2.4.13, and the maps above are the images under Π_∞ of the classical pinch comultiplication on the topological suspension ΣY . The classical pointed homotopies expressing the cogroup axioms are carried by Π_∞ to homotopies in An_* . The double suspension $\Sigma^2 X$ carries comultiplications in two a priori distinct ways, but by the Eckmann–Hilton argument they agree and the resulting comultiplication is cocommutative.

Consequently, for every pointed anima Y the representable functor $[-, Y]_*$ sends wedges to products, so the cogroup structure on ΣX makes $[\Sigma X, Y]_*$ into a group; this group is abelian when the source is a double suspension, as for $[\Sigma^2 X, Y]_*$.

Proposition 2.4.21 (Whitehead theorem for animae). *A morphism $f: X \rightarrow Y$ of animae is an isomorphism if and only if it induces a bijection $\pi_0(X) \xrightarrow{\cong} \pi_0(Y)$ on path components and for every $x \in X$ it induces an isomorphism $\pi_n(X, x) \xrightarrow{\cong} \pi_n(Y, f(x))$ on higher homotopy groups for all $n \geq 1$.*

Proof. By Corollary 2.3.7, the map f is the image under $\Pi_\infty: \mathrm{Cell} \rightarrow \mathrm{An}$ of a continuous map between cell complexes. In light of Remark 2.4.18, the claim is thus an instance of the usual Whitehead theorem for cell complexes, recorded in Theorem 2.3.3. $\square$

Corollary 2.4.22. *Let $f: X \rightarrow Y$ be a map of pointed animae with X and Y connected. Then f is an isomorphism if and only if the induced map*

$$\Omega f: \Omega X \rightarrow \Omega Y$$

is an isomorphism in An .

Proof. If f is an isomorphism, then clearly so is Ωf . Conversely, if Ωf is an isomorphism, then it in particular induces isomorphisms on all homotopy groups, and using the identification $\pi_n(X) \cong \pi_{n-1}(\Omega X)$ for $n \geq 1$ it follows that the map $\pi_n(f): \pi_n(X) \rightarrow \pi_n(Y)$ is an isomorphism for all $n \geq 1$. Since X and Y are connected, $\pi_0(X) \rightarrow \pi_0(Y)$ is automatically a bijection, so f is an isomorphism by Proposition 2.4.21. $\square$

We now briefly recall that the homotopy groups carry natural group structures.

Construction 2.4.23 (Loop anima as a group object). Given a pointed anima $X \in \mathbf{An}_*$, we may consider the maps

$$\begin{aligned} e: * &\xrightarrow{(\text{id}, \text{id})} * \times_X * = \Omega X, \\ M: \Omega X \times \Omega X &\cong * \times_X * \times_X * \xrightarrow{(\text{pr}_1, \text{pr}_3)} * \times_X * = \Omega X, \\ i: \Omega X &= * \times_X * \xrightarrow{(\text{pr}_2, \text{pr}_1)} * \times_X * = \Omega X. \end{aligned}$$

These maps satisfy the usual group relations up to homotopy, so that ΩX acquires the structure of a group object in $\mathbf{Ho}(\mathbf{An}_*)$. We leave the details to the reader.

Corollary 2.4.24. *For every pointed anima X and $n \geq 1$, the set $\pi_n(X)$ carries a natural group structure, and for $n \geq 2$ this group is abelian.*

Proof. The group structure on $\pi_1(X) \cong \pi_0(\Omega X)$ comes from the group object structure of ΩX in Construction 2.4.23. For $n \geq 2$, $\pi_n(X) \cong \pi_{n-2}(\Omega^2 X)$ acquires two group structures which commute with each other, and it follows from the Eckmann–Hilton argument that they agree and are abelian. $\square$

By Lemma 2.4.5, applying pointed homotopy classes to a fiber sequence produces an exact sequence. We now extend such a sequence indefinitely by repeatedly passing to fibers.

Recall that the loop object $\Omega X = 0 \times_X 0$ admits a canonical *inversion map*

$$\iota_X: \Omega X \rightarrow \Omega X,$$

induced by interchanging the two copies of 0 in the pullback. Given a pointed map $f: Y \rightarrow \Omega X$, we write

$$-f := \iota_X \circ f: Y \rightarrow \Omega X.$$

Proposition 2.4.25 (Puppe sequence). *Given a fiber sequence $F \xrightarrow{i} E \xrightarrow{p} B$ in $\mathbf{An}_*$, there is a sequence*

$$\dots \xrightarrow{\Omega^2 p} \Omega^2 B \xrightarrow{-\Omega \delta} \Omega F \xrightarrow{-\Omega i} \Omega E \xrightarrow{-\Omega p} \Omega B \xrightarrow{\delta} F \xrightarrow{i} E \xrightarrow{p} B,$$

where each consecutive triple forms a fiber sequence.

Proof. To see that ΩB is indeed the fiber of i , consider the following commutative diagram:

$$\begin{array}{ccccc} \Omega B & \overset{\delta}{\dashrightarrow} & F & \xrightarrow{i} & 0 \\ \downarrow & & \downarrow i & \lrcorner & \downarrow b \\ 0 & \xrightarrow{e} & E & \xrightarrow{p} & B. \end{array}$$

Here the outer square is the one defining ΩB , and the right square is obtained by flipping the given fiber sequence. The map δ is induced by the universal property of the pullback F , and it follows from the pasting law of pullback squares that also the left square is a pullback square, thus exhibiting ΩB as the fiber of i .

The rest of the Puppe sequence is obtained by iterating this argument indefinitely. The signs show up because in the iteration procedure we flip the homotopies in the squares; under the pullback description of loop objects, this amounts to composing with the inversion map defined above. For example, to determine the fiber of δ , we consider the following commutative diagram:

$$\begin{array}{ccccc}
 \Omega E & \overset{\curvearrowright}{\dashrightarrow} & \Omega B & \xrightarrow{\quad} & 0 \\
 \downarrow & \lrcorner & \downarrow \delta & \lrcorner & \downarrow e \\
 0 & \xrightarrow{f} & F & \xrightarrow{i} & E.
 \end{array}
 \quad \square$$

Corollary 2.4.26 (Long exact sequence on homotopy groups). *A fiber sequence $F \xrightarrow{i} E \xrightarrow{p} B$ of pointed animae induces a long exact sequence*

$$\cdots \rightarrow \pi_2(B) \xrightarrow{\delta_*} \pi_1(F) \xrightarrow{i_*} \pi_1(E) \xrightarrow{p_*} \pi_1(B) \xrightarrow{\delta_*} \pi_0(F) \rightarrow \pi_0(E) \rightarrow \pi_0(B).$$

Proof. Since each consecutive triple in the Puppe sequence is a fiber sequence, Lemma 2.4.5, applied with $W = S^0$, shows that applying $\pi_0(-) = [S^0, -]_*$ produces an exact sequence

$$\cdots \rightarrow \pi_0(\Omega^2 B) \rightarrow \pi_0(\Omega F) \rightarrow \pi_0(\Omega E) \rightarrow \pi_0(\Omega B) \rightarrow \pi_0(F) \rightarrow \pi_0(E) \rightarrow \pi_0(B).$$

In light of the natural isomorphisms $\pi_0(\Omega^n X) \cong \pi_n(X)$ we are done. $\square$

Remark 2.4.27. Forming cofibers instead of fibers similarly gives a dual Puppe sequence $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{\delta} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y \xrightarrow{-\Sigma g} \Sigma Z \xrightarrow{-\Sigma \delta} \Sigma^2 X \rightarrow \cdots$, see Exercise 3.1.3.

Exercise 2.4.28 (Sequential colimits as pushouts). Let $Z_0 \xrightarrow{f_0} Z_1 \xrightarrow{f_1} Z_2 \xrightarrow{f_2} \cdots$ be a sequence of pointed animae. Show that its colimit may be computed as the pushout

$$\begin{array}{ccc}
 \bigvee_{n \geq 0} Z_n & \xrightarrow{a} & \bigvee_{k \geq 0} Z_{2k} \\
 b \downarrow & & \downarrow \\
 \bigvee_{k \geq 0} Z_{2k+1} & \longrightarrow & \operatorname{colim}_n Z_n,
 \end{array}$$

where on the summand Z_n the maps a and b are defined as follows. If $n = 2k$ is even, then a is the inclusion $\operatorname{id}: Z_{2k} \rightarrow Z_{2k}$ of a wedge summand and b is the map $f_{2k}: Z_{2k} \rightarrow Z_{2k+1}$; if $n = 2k + 1$ is odd, then a is the map $f_{2k+1}: Z_{2k+1} \rightarrow Z_{2k+2}$ and b is the inclusion $\operatorname{id}: Z_{2k+1} \rightarrow Z_{2k+1}$.

Hint: Recall the pushout description of $[\omega]$ in Axiom D.

Exercises

Exercise 2.1 (Basic underlying animae). Compute the underlying anima $\Pi_\infty(X)$ for each of the following topological spaces:

$$X = \emptyset, \quad X = *, \quad X = S$$

where S is a discrete set, and

$$X = [0, 1], \quad X = S^0, \quad X = \bigsqcup_{i=1}^n *.$$

Then check directly in these examples how Π_∞ behaves with respect to finite products and finite coproducts.

Exercise 2.2 (The pushout counterexample, repaired). For $n \geq 1$, recall the two strict pushout squares from Example 2.1.1:

$$\begin{array}{ccc} S^{n-1} & \hookrightarrow & D^n \\ \downarrow & & \downarrow \\ * & \longrightarrow & S^n \end{array} \quad \begin{array}{ccc} S^{n-1} & \longrightarrow & * \\ \downarrow & & \downarrow \\ * & \longrightarrow & *. \end{array}$$

Show that the corresponding homotopy pushouts are homotopy equivalent. Identify both of them with S^n .

Exercise 2.3 (Maps out of a homotopy pushout). Let $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ be continuous maps, and let T be another topological space. Show that continuous maps

$$X \sqcup_Z^h Y \rightarrow T$$

are naturally the same as triples (t_X, t_Y, H) , where $t_X: X \rightarrow T$ and $t_Y: Y \rightarrow T$ are continuous maps and $H: Z \times [0, 1] \rightarrow T$ is a homotopy

$$H: t_X \circ f \sim t_Y \circ g.$$

In other words, maps out of the homotopy pushout are homotopy coherent cocones under the span $X \leftarrow Z \rightarrow Y$.

Exercise 2.4 (Maps into a homotopy pullback). Let $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ be continuous maps, and let T be another topological space. Show that continuous maps

$$T \rightarrow X \times_Z^h Y$$

are naturally the same as triples (t_X, t_Y, H) , where $t_X: T \rightarrow X$ and $t_Y: T \rightarrow Y$ are continuous maps and $H: T \times [0, 1] \rightarrow Z$ is a homotopy

$$H: f \circ t_X \sim g \circ t_Y.$$

Exercise 2.5 (Homotopy fibers in examples). Compute the homotopy fiber over the basepoint in each of the following cases:

- (1) the inclusion of the basepoint $* \rightarrow S^1$;
- (2) the covering map $\mathbb{R} \rightarrow S^1, t \mapsto e^{2\pi it}$;
- (3) for $n \geq 1$, the degree- n map $S^1 \rightarrow S^1, z \mapsto z^n$.

For the first example, show that the homotopy fiber is homotopy equivalent to a discrete copy of $\mathbb{Z}$. For the last two examples, compare the homotopy fiber with the ordinary fiber.

Exercise 2.6 (Loop spaces as homotopy fibers). Let (X, x) be a pointed topological space. Use the definition of homotopy pullback to show that the homotopy pullback of

$$* \xrightarrow{x} X \xleftarrow{x} *$$

is the usual based loop space of X at x .

Exercise 2.7 (Mapping path factorization). Let $f: E \rightarrow B$ be a continuous map, and let

$$P(f) := \{(e, \gamma) \in E \times B^{[0,1]} \mid f(e) = \gamma(0)\}$$

be its mapping path space. Let $i: E \rightarrow P(f)$ be the map $e \mapsto (e, \text{const}_{f(e)})$, and let $p: P(f) \rightarrow B$ be the map $(e, \gamma) \mapsto \gamma(1)$.

- (1) Show that $p \circ i = f$.
- (2) Write down an explicit homotopy inverse to i and show that i is a homotopy equivalence.
- (3) Identify $P(f)$ in the special cases $f = \text{id}_B$ and $f: E \rightarrow *$.
- (4) Explain geometrically why p behaves like a fibration: a point in $P(f)$ remembers not only a point of E , but also a path in the base.

Exercise 2.8 (Cell attachments and cofibers). Let $n \geq 1$ and consider the inclusion $S^{n-1} \hookrightarrow D^n$.

- (1) Compute the reduced homotopy cofiber of this inclusion as a pointed topological space.
- (2) Show that it is homotopy equivalent to S^n .
- (3) Translate the same computation into An_* using Proposition 2.4.12.

Exercise 2.9 (Wedges and pointed coproducts). Use the pushout formula from Definition 2.4.9 to compute the following wedges in An_* :

$$S^0 \vee S^0, \quad S^1 \vee S^1, \quad \bigvee_{i=1}^n S^0.$$

Compare your answers with the corresponding topological wedge sums.

Exercise 2.10 (Suspension examples). Use the intrinsic definition of suspension in An_* .

- (1) Show that $\Sigma S^n \cong S^{n+1}$ for all $n \geq 0$.
- (2) Compute $\Sigma(*)$ and $\Sigma(S^0)$.
- (3) Show that $\Sigma(X \vee Y) \cong \Sigma X \vee \Sigma Y$ for pointed anima X and Y .

Exercise 2.11 (Smash products). Use the pushout descriptions of wedges and cofibers to establish natural isomorphisms $S^0 \wedge X \cong X$ and $X_+ \wedge Y_+ \cong (X \times Y)_+$. Show that $(-) \wedge X$ preserves pushouts, and deduce that

$$S^1 \wedge X \cong \Sigma X \quad \text{and} \quad S^m \wedge S^n \cong S^{m+n}$$

for all $m, n \geq 0$.

Exercise 2.12. Show by counterexample that the conclusion of Corollary 2.4.22 fails if X and Y are not assumed connective.

Exercise 2.13 (The coPuppe sequence for a cell attachment). Apply the cofiber version of the Puppe sequence, described in the remark after Proposition 2.4.25, to the cofiber sequence

$$S^{n-1} \rightarrow D^n \rightarrow S^n$$

in pointed anima.

- (1) Write out the first six terms of the resulting sequence.
- (2) Use the contractibility of D^n to simplify the sequence.
- (3) Identify the connecting map $S^n \rightarrow \Sigma S^{n-1}$ up to isomorphism.

3 Homology and cohomology

A central theme in algebraic topology is the assignment of algebraic invariants to topological spaces. Among the most important are the *homology groups* $H_n(X; \mathbb{Z})$ and *cohomology groups* $H^n(X; \mathbb{Z})$. They are computable, satisfy convenient formal properties, and distinguish many spaces that are otherwise hard to tell apart.

More generally, a *generalized (co)homology theory* assigns graded abelian groups to spaces in a weakly homotopy invariant way and interacts predictably with wedges, cofiber sequences, and suspensions. We formulate these axioms directly for pointed animae, where homotopy invariance is automatic. We then explain the close relation with *spectra*, viewed for now as sequences of pointed animae $E_0, E_1, E_2, \dots$ equipped with isomorphisms $E_n \simeq \Omega E_{n+1}$. Every spectrum represents a cohomology theory, and Brown representability gives the converse.

Ordinary homology and cohomology are singled out among generalized theories by the *dimension axiom*. We take their existence for granted here and construct them intrinsically in Chapter 6. The online version also develops their cellular computation and gives a constructive proof of Brown representability. This chapter thereby connects classical algebraic topology with the theory of spectra and stable ∞ -categories developed in Chapter 4. Readers already comfortable with (co)homology theories may begin with Section 3.2.

3.1 Homology and cohomology theories

We start with the axiomatic approach to homology and cohomology theories.

Convention 3.1.1 (Graded abelian groups). A *graded abelian group* is a sequence $A_* = (A_n)_{n \in \mathbb{Z}}$ of abelian groups indexed by the integers. These form a 1-category $\mathbf{Ab}^{\mathbb{Z}}$ whose morphisms $f_*: A_* \rightarrow B_*$ consist of homomorphisms $f_n: A_n \rightarrow B_n$ for each $n \in \mathbb{Z}$.

Definition 3.1.2 (Homology and cohomology theories). A *reduced homology theory* is a pair $(\tilde{E}_*, \sigma)$ consisting of a functor

$$\tilde{E}_*: \mathbf{An}_* \rightarrow \mathbf{Ab}^{\mathbb{Z}}$$

and a natural isomorphism

$$\sigma: \tilde{E}_*(-) \xrightarrow{\sim} \tilde{E}_{*+1}(\Sigma(-)),$$

called the *suspension isomorphism*, satisfying:

(1) (*Wedge axiom*) For every collection $(X_i)_{i \in I}$ of pointed animae, the canonical map

$$\bigoplus_{i \in I} \tilde{E}_*(X_i) \rightarrow \tilde{E}_*\left(\bigvee_{i \in I} X_i\right)$$

is an isomorphism.

(2) (*Exactness*) For every cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in $\mathbf{An}_*$, the sequence of abelian groups

$$\widetilde{E}_*(X) \xrightarrow{f_*} \widetilde{E}_*(Y) \xrightarrow{g_*} \widetilde{E}_*(Z)$$

is exact in each degree.

Dually, a *reduced cohomology theory* is a pair $(\widetilde{E}^*, \sigma)$ consisting of a functor $\widetilde{E}^*: \mathbf{An}_*^{\text{op}} \rightarrow \mathbf{Ab}^{\mathbb{Z}}$ and a natural isomorphism $\sigma: \widetilde{E}^*(-) \xrightarrow{\sim} \widetilde{E}^{*+1}(\Sigma(-))$ satisfying:

(1) (*Wedge axiom*) For every collection $(X_i)_{i \in I}$ of pointed animae, the canonical map $\widetilde{E}^*\left(\bigvee_{i \in I} X_i\right) \rightarrow \prod_{i \in I} \widetilde{E}^*(X_i)$ is an isomorphism.

(2) (*Exactness*) For every cofiber sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in $\mathbf{An}_*$, the sequence $\widetilde{E}^*(Z) \xrightarrow{g^*} \widetilde{E}^*(Y) \xrightarrow{f^*} \widetilde{E}^*(X)$ is exact in each degree.

The suspension isomorphism extends exactness to the familiar long exact sequences. Indeed, applying exactness to the coPuppe sequence of a cofiber sequence $X \rightarrow Y \rightarrow Z$ produces boundary maps

$$\partial: \widetilde{E}_n(Z) \rightarrow \widetilde{E}_{n-1}(X) \quad \text{and} \quad \partial: \widetilde{E}^n(X) \rightarrow \widetilde{E}^{n+1}(Z).$$

We record this as an exercise because the construction is a useful way to become comfortable with cofiber sequences.

Exercise 3.1.3 (The coPuppe sequence and long exact sequences). Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be a cofiber sequence in $\mathbf{An}_*$. By dualizing the construction of Proposition 2.4.25, construct the *coPuppe sequence*: an infinite sequence

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y \xrightarrow{-\Sigma g} \Sigma Z \xrightarrow{-\Sigma h} \Sigma^2 X \xrightarrow{\Sigma^2 f} \dots$$

in which every consecutive triple forms a cofiber sequence. Deduce that any homology theory gives rise to a long exact sequence

$$\dots \rightarrow \widetilde{E}_{n+1}(Z) \xrightarrow{\partial} \widetilde{E}_n(X) \xrightarrow{f_*} \widetilde{E}_n(Y) \xrightarrow{g_*} \widetilde{E}_n(Z) \xrightarrow{\partial} \widetilde{E}_{n-1}(X) \rightarrow \dots$$

Formulate and prove the analogous statement for cohomology theories.

Reduced theories give unreduced theories by adding a disjoint basepoint. For an unpointed anima X , write $X_+ := X \sqcup \text{pt}$, pointed by the added point. We then set

$$E_n(X) := \widetilde{E}_n(X_+) \quad \text{and} \quad E^n(X) := \widetilde{E}^n(X_+).$$

For a map $A \rightarrow X$ of unpointed animae, the relative groups are defined by

$$E_n(X, A) := \widetilde{E}_n(X/A) \quad \text{and} \quad E^n(X, A) := \widetilde{E}^n(X/A),$$

where X/A denotes the cofiber of $A_+ \rightarrow X_+$.

Exercise 3.1.4 (Long exact sequence of a pair). Let $i: A \rightarrow X$ be a map of animae, and let $j: X_+ \rightarrow X/A$ denote the cofiber map. Show that every homology theory gives rise to a natural long exact sequence

$$\cdots \rightarrow E_n(A) \xrightarrow{i_*} E_n(X) \xrightarrow{j_*} E_n(X, A) \xrightarrow{\partial} E_{n-1}(A) \xrightarrow{i_*} E_{n-1}(X) \rightarrow \cdots .$$

Formulate and prove the analogous long exact sequence for cohomology:

$$\cdots \rightarrow E^n(X, A) \xrightarrow{j^*} E^n(X) \xrightarrow{i^*} E^n(A) \xrightarrow{\partial} E^{n+1}(X, A) \xrightarrow{j^*} E^{n+1}(X) \rightarrow \cdots .$$

Exercise 3.1.5. Show that precomposition with $\Pi_\infty: \text{Top}_* \rightarrow \text{An}_*$ establishes a bijection between reduced cohomology theories in the sense of Definition 3.1.2 and pairs (E^*, σ) consisting of a functor $E^*: \text{Top}_*^{\text{op}} \rightarrow \text{Ab}^{\mathbb{Z}}$ and a natural isomorphism $\sigma: E^*(-) \xrightarrow{\sim} E^{*+1}(\Sigma(-))$ satisfying:

- (1) Weak homotopy equivalences induce isomorphisms on E^* .
- (2) The wedge axiom holds for wedges of pointed CW-complexes whose basepoints are 0-cells.
- (3) For every relative CW-pair (X, A) , the sequence $E^*(X/A) \rightarrow E^*(X) \rightarrow E^*(A)$ is exact.

Formulate the analogous characterization of homology theories.

Hint: Use Theorem 2.3.4 to reduce to CW-complexes, and Proposition 2.4.12 to transport the relevant constructions.

3.2 Cohomology theories and spectra

We now relate cohomology theories to spectra. We first explain how every spectrum determines a reduced cohomology theory. Brown representability supplies the converse, showing that every reduced cohomology theory arises in this way.

Definition 3.2.1 (Spectrum). A *spectrum* E is a sequence of pointed animae $(E_m)_{m \geq 0}$ together with isomorphisms

$$\sigma_m: E_m \xrightarrow{\sim} \Omega E_{m+1}$$

for all $m \geq 0$. We call the maps σ_m the *structure maps* of E .

For convenience, we extend the sequence to all integer indices by setting

$$E_k := \Omega^{-k} E_0 \quad (k < 0).$$

With this convention, the structure maps extend to all $k \in \mathbb{Z}$: for $k < 0$, we take σ_k to be the identity map under the identification

$$E_k = \Omega^{-k} E_0 = \Omega(\Omega^{-k-1} E_0) = \Omega E_{k+1}.$$

Definition 3.2.2 (Morphism of spectra). Let $E = (E_m, \sigma_m)$ and $F = (F_m, \tau_m)$ be spectra. A *morphism of spectra* $f: E \rightarrow F$ consists of pointed maps

$$f_m: E_m \rightarrow F_m \quad (m \geq 0)$$

together with chosen homotopies making the following squares commute for every $m \geq 0$:

$$\begin{array}{ccc} E_m & \xrightarrow[\sim]{\sigma_m} & \Omega E_{m+1} \\ f_m \downarrow & & \downarrow \Omega f_{m+1} \\ F_m & \xrightarrow[\sim]{\tau_m} & \Omega F_{m+1}. \end{array}$$

For $k < 0$, we extend f to the negative levels by setting $f_k := \Omega^{-k} f_0: E_k \rightarrow F_k$.

Definition 3.2.3 (*E-cohomology*). Let E be a spectrum and let $X \in \mathbf{An}_*$ be a pointed anima. For $k \in \mathbb{Z}$, we define the *reduced E-cohomology* of X by

$$\tilde{E}^k(X) := [X, E_k]_*.$$

The isomorphism $E_k \xrightarrow{\sim} \Omega^2 E_{k+2}$ identifies this set with the second homotopy group of the pointed mapping anima $\mathrm{Hom}_{\mathbf{An}_*}(X, E_{k+2})$, and therefore equips it with a natural abelian group structure. The structure isomorphisms of E induce natural isomorphisms

$$\tilde{E}^k(X) = [X, E_k]_* \xrightarrow{\cong} [X, \Omega E_{k+1}]_* \xrightarrow{\cong} [\Sigma X, E_{k+1}]_* = \tilde{E}^{k+1}(\Sigma X),$$

which define the suspension isomorphism for $\tilde{E}^*$. Both maps are group homomorphisms: the first is induced by an isomorphism of pointed animae, while the second comes from the natural equivalence $\mathrm{Hom}_{\mathbf{An}_*}(\Sigma X, Y) \simeq \mathrm{Hom}_{\mathbf{An}_*}(X, \Omega Y)$ of Lemma 2.4.8, which induces isomorphisms on the homotopy groups of mapping animae.

A morphism of spectra $f: E \rightarrow F$ induces a natural transformation $\tilde{E}^* \rightarrow \tilde{F}^*$ by postcomposition with the maps $f_k: E_k \rightarrow F_k$. These maps are homomorphisms for the abelian group structures above, and the chosen homotopies in Definition 3.2.2 ensure that the transformation commutes with the suspension isomorphisms.

Proposition 3.2.4. *For every spectrum E , the pair $(\tilde{E}^*, \sigma)$ is a reduced cohomology theory.*

Proof. The suspension isomorphism was constructed above. For a collection $(X_i)_{i \in I}$ of pointed animae, the universal property of the wedge gives an isomorphism

$$\tilde{E}^k\left(\bigvee_{i \in I} X_i\right) = \left[\bigvee_{i \in I} X_i, E_k\right]_* \cong \prod_{i \in I} [X_i, E_k]_* = \prod_{i \in I} \tilde{E}^k(X_i),$$

which is the wedge axiom. Finally, Lemma 2.4.5 shows that every cofiber sequence $X \rightarrow Y \rightarrow Z$ induces an exact sequence

$$[Z, E_k]_* \longrightarrow [Y, E_k]_* \longrightarrow [X, E_k]_*$$

of pointed sets. Since these maps are homomorphisms for the abelian group structures constructed above, this proves the exactness axiom. $\square$

Remark 3.2.5. After constructing the tensor product of spectra, we will associate to E a reduced homology theory by the formula

$$\tilde{E}_k(X) := \pi_k(\Sigma^\infty X \otimes E);$$

we refer to Section 4.4.5 for details. The idea of using spectra to define homology and cohomology theories, and with it the notion of a generalized homology theory, goes back to Whitehead [Whi62].

Theorem 3.2.6 (Brown representability, Brown [Bro62], Switzer [Swi75]). *For every reduced cohomology theory (h^*, σ) , there is a spectrum E and a natural isomorphism $h^* \xrightarrow{\sim} \tilde{E}^*$ compatible with the suspension isomorphisms. Moreover, every natural transformation $\tilde{E}^* \rightarrow \tilde{F}^*$ of reduced cohomology theories that commutes with the suspension isomorphisms is induced by a morphism of spectra $E \rightarrow F$.*

The first assertion is a standard consequence of Brown's representability theorem. After translating Brown's result to the language of animae, apply it degreewise to the restriction of h^k to connected pointed animae, and let Y_k be a connected representing anima. Since ΣX is connected for every pointed anima X , the suspension isomorphism gives natural isomorphisms

$$h^k(X) \cong h^{k+1}(\Sigma X) \cong [\Sigma X, Y_{k+1}]_* \cong [X, \Omega Y_{k+1}]_*.$$

Thus we may take $E_k := \Omega Y_{k+1}$. The suspension isomorphisms induce maps $Y_{k+1} \rightarrow \Omega Y_{k+2}$ that are isomorphisms on all positive homotopy groups; looping them gives the required isomorphisms $E_k \xrightarrow{\sim} \Omega E_{k+1}$. The final assertion follows degreewise from the Yoneda lemma, as in Exercise 3.1. A complete proof is given in Section 3.4.

3.3 Ordinary homology and cohomology

Ordinary homology and cohomology are the generalized theories singled out by the dimension axiom. In this chapter we use them axiomatically, postponing their intrinsic construction to Section 6.2. The online version continues with the cellular computation of ordinary homology from a CW-structure.

Theorem 3.3.1 (Existence of ordinary homology and cohomology). *For every abelian group A , there exist reduced homology and cohomology theories*

$$\tilde{H}_*(-; A) : \mathbf{An}_* \rightarrow \mathbf{Ab}^{\mathbb{Z}} \quad \text{and} \quad \tilde{H}^*(-; A) : \mathbf{An}_*^{\text{op}} \rightarrow \mathbf{Ab}^{\mathbb{Z}}$$

satisfying the dimension axiom

$$\tilde{H}_n(S^0; A) \cong \begin{cases} A & \text{if } n = 0, \\ 0 & \text{if } n \neq 0, \end{cases} \quad \text{and} \quad \tilde{H}^n(S^0; A) \cong \begin{cases} A & \text{if } n = 0, \\ 0 & \text{if } n \neq 0. \end{cases}$$

The associated unreduced theories are denoted $H_(-; A)$ and $H^*(-; A)$. When $A = \mathbb{Z}$, we usually omit it from the notation.*

Remark 3.3.2 (Singular (co)homology). Given a topological space X , one can write down an explicit chain complex of abelian groups $C_*^{\text{sing}}(X; \mathbb{Z})$ called the *singular chain complex of X* . The n -th term of this chain complex is given by

$$C_n^{\text{sing}}(X; \mathbb{Z}) := \mathbb{Z}[\text{Sing}_n(X)],$$

the free abelian group generated by the set $\text{Sing}_n(X) = \text{Hom}_{\text{Top}}(|\Delta^n|, X)$ of continuous maps from the geometric n -simplex to X . The boundary map $d_n : C_n^{\text{sing}}(X; \mathbb{Z}) \rightarrow C_{n-1}^{\text{sing}}(X; \mathbb{Z})$ sends a generating n -simplex σ to the alternating sum $\sum_{i=0}^n (-1)^i d_i(\sigma)$ of its faces.

For an abelian group A , tensoring with A and taking hom into A produces chain and cochain complexes

$$C_*^{\text{sing}}(X; A) := C_*^{\text{sing}}(X; \mathbb{Z}) \otimes A \quad \text{and} \quad C_{\text{sing}}^*(X; A) := \text{Hom}_{\text{Ab}}(C_*^{\text{sing}}(X; \mathbb{Z}), A).$$

Their homology groups are the classical singular homology and cohomology groups of X with coefficients in A . This is the traditional construction of ordinary (co)homology. The fact that this satisfies the axioms is non-trivial.

In this book, we will not take singular chains as the definition, because it depends on choosing a topological presentation of an anima. Instead, in Section 6.2 we will construct ordinary homology and cohomology intrinsically using the derived category $D(\mathbb{Z})$. The comparison there shows that, for a topological space X , the intrinsic ordinary (co)homology of the underlying anima $\Pi_\infty(X)$ agrees with the singular (co)homology of X .

Remark 3.3.3. The dimension axiom implies the usual values on spheres. Namely, since $S^n \simeq \Sigma^n S^0$ as pointed anima, the suspension isomorphism gives

$$\widetilde{H}_k(S^n; A) \cong \begin{cases} A & \text{if } k = n, \\ 0 & \text{if } k \neq n, \end{cases}$$

and similarly for cohomology.

The next chapter organizes spectra and their morphisms into an ∞ -category. Before turning to it, the online version continues with an optional cellular computation of ordinary homology, followed by a proof of Brown representability. Neither is needed later.

3.3.1 Cellular homology (online only)

Convention 3.3.4. Throughout this section, we write D^n for the n -disk and S^{n-1} for its boundary sphere, viewed as anima. While contractibility of the disk implies that $D^n \simeq \text{pt}$, this notation emphasizes the analogy to cell attachments in CW-complexes. Given a map $f: A \rightarrow X$ of anima, we write X/A for the cofiber of $A \rightarrow X$.

Definition 3.3.5 (CW-structure). A *CW-structure* on an anima X consists of a sequence of anima

$$\emptyset = X^{-2} = X^{-1} \rightarrow X^0 \rightarrow X^1 \rightarrow X^2 \rightarrow \dots$$

called the *skeleta* of X , together with an equivalence $X \simeq \text{colim}_n X^n$, satisfying:

- (1) The anima X^0 is a set;
- (2) For each $n \geq 1$, there exists a set I_n of n -cells and a pushout square of anima

$$\begin{array}{ccc} \coprod_{\alpha \in I_n} S^{n-1} & \xrightarrow{\varphi} & X^{n-1} \\ \downarrow & & \downarrow \\ \coprod_{\alpha \in I_n} D^n & \xrightarrow{\quad} & X^n. \end{array}$$

The map φ is called the *attaching map* for the n -cells.

We say that X is *finite-dimensional* if there are no n -cells for sufficiently large n , and *finite* if it has finitely many cells in total.

Remark 3.3.6 (Cellular approximation). We will freely use the *cellular approximation theorem*: every pointed map between pointed cell complexes is homotopic to a cellular one. Together with the comparison of classical and intrinsic homotopy groups in Remark 2.4.18, it has the following consequence for an anima X with a CW-structure, which is all we will need. The inclusion of the n -skeleton induces a map

$$[S^k, X^n]_* \rightarrow [S^k, X]_*$$

that is a bijection for $k < n$ and a surjection for $k = n$; the same holds for the inclusion $X^n \rightarrow X^m$ of skeleta with $m > n$. More generally, if W is obtained from an anima Z by attaching cells of dimension $\geq m$, then $[S^k, Z]_* \rightarrow [S^k, W]_*$ is a bijection for $k < m - 1$ and a surjection for $k = m - 1$. Equivalently, every pointed map $S^k \rightarrow W$ is homotopic to one factoring through Z once $k \leq m - 1$, and two such maps that become homotopic in W already become homotopic after attaching the cells of dimension m .

Example 3.3.7. The results from Chapter 2 show: for a topological CW-complex X with skeleta X^n , its underlying anima $\Pi_\infty(X)$ obtains a CW-structure with skeleta $\Pi_\infty(X^n)$.

Example 3.3.8. The n -sphere S^n admits a CW-structure with one 0-cell and one n -cell: we have a pushout square

$$\begin{array}{ccc} S^{n-1} & \longrightarrow & \text{pt} \\ \downarrow & \lrcorner & \downarrow \\ D^n & \longrightarrow & S^n. \end{array}$$

The key to cellular homology is understanding how homology changes as we pass from one skeleton to the next. Recall from Section 3.1 that the relative groups are defined by $H_k(X, A) := \tilde{H}_k(X/A)$ and fit into the long exact sequence of Exercise 3.1.4.

Lemma 3.3.9. *Let X be an anima with a CW-structure. For all $k, n \in \mathbb{Z}$ with $n \geq 0$:*

- (a) *The relative homology $H_k(X^n, X^{n-1})$ is zero for $k \neq n$ and is free abelian for $k = n$, with one generator for each n -cell of X .*
- (b) *$H_k(X^n) = 0$ for $k > n$.*
- (c) *The map $H_k(X^n) \rightarrow H_k(X)$ induced by the structure map $X^n \rightarrow X$ is an isomorphism for $k < n$ and surjective for $k = n$.*

Proof. For part (a), the case $n = 0$ follows because X^0 is a disjoint union of points. For $n \geq 1$, the pushout square defining X^n from X^{n-1} shows that

$$X^n/X^{n-1} \simeq \bigvee_{\alpha \in I_n} D^n/S^{n-1} \simeq \bigvee_{\alpha \in I_n} S^n,$$

a wedge of n -spheres indexed by the n -cells. Therefore

$$H_k(X^n, X^{n-1}) = \tilde{H}_k\left(\bigvee_{\alpha \in I_n} S^n\right) \cong \bigoplus_{\alpha \in I_n} \tilde{H}_k(S^n) \cong \begin{cases} \bigoplus_{\alpha \in I_n} \mathbb{Z} & \text{if } k = n, \\ 0 & \text{if } k \neq n. \end{cases}$$

For part (b), the claim is clear for $n = 0$: we have $H_k(X^0) = 0$ for $k > 0$ as X^0 is a disjoint union of points, which have trivial homology. Consider now the long exact sequence of the pair (X^n, X^{n-1}) :

$$H_{k+1}(X^n, X^{n-1}) \rightarrow H_k(X^{n-1}) \rightarrow H_k(X^n) \rightarrow H_k(X^n, X^{n-1}).$$

By part (a), the first term vanishes when $k \neq n - 1$ and the last term vanishes when $k \neq n$. In particular, for $k > n$ the map $H_k(X^{n-1}) \rightarrow H_k(X^n)$ is an isomorphism and the claim follows by induction. For part (c), we may similarly use the long exact sequence of the pair (X^{n+1}, X^n) to conclude that the map $H_k(X^n) \rightarrow H_k(X^{n+1})$ is surjective for $k \leq n$ and injective (hence an isomorphism) for $k < n$.

It remains to pass from the skeleta to their colimit. Adding a disjoint basepoint commutes with colimits. Apply the pushout presentation of Exercise 2.4.28 to the sequence $(X^r)_+$. Since the resulting square is a pushout, the cofiber of its top horizontal map is isomorphic to the cofiber of its bottom horizontal map. Comparing the corresponding long exact sequences gives the Mayer–Vietoris sequence of the pushout. By the wedge axiom, and after multiplying the odd source summands by -1 , the relevant portion is

$$\bigoplus_{r \geq 0} H_k(X^r) \xrightarrow{\delta_k} \bigoplus_{r \geq 0} H_k(X^r) \longrightarrow H_k(X) \longrightarrow \bigoplus_{r \geq 0} H_{k-1}(X^r) \xrightarrow{\delta_{k-1}} \bigoplus_{r \geq 0} H_{k-1}(X^r).$$

Writing ι_r for the inclusion of the r -th summand, the first map is given by

$$\delta_k(\iota_r(x)) = \iota_r(x) - \iota_{r+1}((X^r \rightarrow X^{r+1})_*(x)).$$

The map δ_k is injective: if an element of the direct sum lies in its kernel, its components vanish successively, starting with the zeroth component. Its cokernel is the algebraic colimit $\text{colim}_r H_k(X^r)$. The same injectivity statement for δ_{k-1} therefore identifies the middle map with an isomorphism $\text{colim}_r H_k(X^r) \xrightarrow{\sim} H_k(X)$. The claim now follows from the stabilization properties established above. $\square$

We now construct the cellular chain complex.

Construction 3.3.10 (Cellular chain complex). Let X be an anima with a CW-structure. Using Lemma 3.3.9, portions of the long exact sequences for the pairs (X^{n+1}, X^n) , (X^n, X^{n-1}) , and (X^{n-1}, X^{n-2}) fit into the following commutative diagram:

$$\begin{array}{ccccccc}
 & & & & & & 0 \\
 & & & & & \nearrow & \\
 & & & & H_n(X^{n+1}) & & \\
 & & & \nearrow & & & \\
 0 & \searrow & & & & & \\
 & & H_n(X^n) & \nearrow & & & \\
 & \nearrow \partial_{n+1} & & \searrow j_n & & & \\
 \dots \rightarrow & H_{n+1}(X^{n+1}, X^n) & \xrightarrow{d_{n+1}} & H_n(X^n, X^{n-1}) & \xrightarrow{d_n} & H_{n-1}(X^{n-1}, X^{n-2}) \rightarrow \dots \\
 & & & \searrow \partial_n & \nearrow j_{n-1} & & \\
 & & & & H_{n-1}(X^{n-1}) & & \\
 & & & \nearrow & & & \\
 & & & 0 & & &
 \end{array}$$

Here the maps j_n and ∂_n come from the long exact sequence of the pair (X^n, X^{n-1}) . Since $H_n(X^{n-1}) = 0$ by Lemma 3.3.9(b), the map j_n is injective for every n . We define the *cellular boundary maps* as the compositions

$$d_n := j_{n-1} \circ \partial_n: H_n(X^n, X^{n-1}) \rightarrow H_{n-1}(X^{n-1}, X^{n-2}).$$

Since $\partial_n \circ j_n = 0$ (as these are consecutive maps in an exact sequence), we have $d_n \circ d_{n+1} = 0$. Thus the groups

$$C_n^{\text{CW}}(X) := H_n(X^n, X^{n-1})$$

with differentials d_n form a chain complex, called the *cellular chain complex* of X .

By Lemma 3.3.9(a), the cellular chain group $C_n^{\text{CW}}(X)$ is a free abelian group with one generator for each n -cell of X . We write $e_\alpha^n \in C_n^{\text{CW}}(X)$ for the generator corresponding to the n -cell indexed by $\alpha \in I_n$.

Definition 3.3.11 (Cellular homology). The *cellular homology groups* of an anima X with CW-structure are the homology groups of the cellular chain complex:

$$H_n^{\text{CW}}(X) := H_n(C_*^{\text{CW}}(X)) = \ker(d_n)/\text{im}(d_{n+1}).$$

The main theorem of this section is that cellular homology agrees with ordinary homology:

Theorem 3.3.12. *For any anima X with a CW-structure and any $n \geq 0$, there is an isomorphism*

$$H_n^{\text{CW}}(X) \cong H_n(X).$$

Proof. By Lemma 3.3.9(c), the map $H_n(X^{n+1}) \rightarrow H_n(X)$ is an isomorphism. Combining this with the previous commutative diagram, we conclude that $H_n(X)$ is isomorphic to the cokernel of the map $\partial_{n+1}: H_{n+1}(X^{n+1}, X^n) \rightarrow H_n(X^n)$. By injectivity of j_n , the target $H_n(X^n)$ of this map may be identified with the kernel of $\partial_n: H_n(X^n, X^{n-1}) \rightarrow H_{n-1}(X^{n-1})$, which in turn may be identified with $\ker(d_n)$ because of the injectivity of j_{n-1} . Under the resulting isomorphism $j_n: H_n(X^n) \xrightarrow{\cong} \ker(d_n)$, the image of ∂_{n+1} gets mapped to the image of $d_{n+1} = j_n \partial_{n+1}$, and we conclude that

$$H_n(X) \cong \ker(d_n)/\text{im}(d_{n+1}) = H_n^{\text{CW}}(X). \quad \square$$

We leave the following easy consequences to the reader:

Exercise 3.3.13. Let X be an anima with a CW-structure.

- (1) If X has no n -cells, then $H_n(X) = 0$.
- (2) If X has k cells in dimension n , then $H_n(X)$ is generated by at most k elements.
- (3) If X has no two cells in adjacent dimensions, then $H_n(X)$ is free abelian with rank equal to the number of n -cells.

Example 3.3.14. Complex projective space $\mathbb{CP}^n$ has a filtration whose successive strata are the affine cells $\mathbb{C}^0, \mathbb{C}^1, \dots, \mathbb{C}^n$, and hence a CW-structure with one cell in each even dimension $0, 2, 4, \dots, 2n$. Since no two cells are in adjacent dimensions, Exercise 3.3.13(3) gives

$$H_k(\mathbb{CP}^n) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0, 2, 4, \dots, 2n, \\ 0 & \text{otherwise.} \end{cases}$$

Example 3.3.15. For $n, m \geq 1$ with $n \neq m$, the product $S^n \times S^m$ has a CW-structure with one 0-cell, one n -cell, one m -cell, and one $(n + m)$ -cell. If $n, m \geq 2$ and $|n - m| > 1$, then no two cells are in adjacent dimensions, so

$$H_k(S^n \times S^m) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0, n, m, \text{ or } n + m, \\ 0 & \text{otherwise.} \end{cases}$$

In the remaining cases, some cells lie in adjacent dimensions, but the corresponding cellular boundary maps are zero, giving the same answer.

To perform explicit computations, we need a formula for the cellular boundary maps in terms of the attaching maps of the cells. The formula involves the degree of maps between spheres.

Remark 3.3.16 (Degree). For every $n \geq 1$, a map $f: S^n \rightarrow S^n$ induces multiplication by a unique integer on $H_n(S^n) \cong \mathbb{Z}$. This integer is called the *degree* of f . For pointed maps, this agrees with the degree recorded in Remark 2.4.19.

Proposition 3.3.17 (Cellular boundary formula). *Let X be an anima with a CW-structure, and let $n \geq 2$. Then the cellular boundary map is given at the generator e_α^n by*

$$d_n(e_\alpha^n) = \sum_{\beta \in I_{n-1}} d_{\alpha\beta} \cdot e_\beta^{n-1},$$

where the sum is over the set $(n - 1)$ -cells, and $d_{\alpha\beta}$ is the degree of the composite map

$$\Delta_{\alpha\beta}: S^{n-1} \xrightarrow{\varphi_\alpha} X^{n-1} \xrightarrow{q} X^{n-1}/X^{n-2} \simeq \bigvee_{\gamma \in I_{n-1}} S_\gamma^{n-1} \xrightarrow{p_\beta} S_\beta^{n-1}.$$

Here q is the quotient map and p_β is the projection onto the β -th wedge summand.

Note that $d_{\alpha\beta}$ is non-zero only for finitely many β , since by compactness of S^{n-1} the map from S^{n-1} to the wedge $\bigvee_\gamma S_\gamma^{n-1}$ only hits finitely many wedge summands.

Proof. Recall that d_n is the composite

$$\tilde{H}_n(X^n/X^{n-1}) \xrightarrow{\partial_n} H_{n-1}(X^{n-1}) \xrightarrow{j_{n-1}} \tilde{H}_{n-1}(X^{n-1}/X^{n-2}),$$

where ∂_n is the connecting homomorphism and $j_{n-1} = q_*$ is induced by the quotient map q . Let

$$\iota_\alpha: S^n \hookrightarrow \bigvee_{\gamma \in I_n} S_\gamma^n \cong X^n/X^{n-1}$$

be the inclusion of the α -th wedge summand. The generator $e_\alpha^n \in \tilde{H}_n(X^n/X^{n-1})$ is the image of the generator $[\iota_n] \in \tilde{H}_n(S^n)$ under the map on homology induced by ι_α . The map ι_α may alternatively be obtained from the attaching map $\varphi_\alpha: S^{n-1} \rightarrow X^{n-1}$ by passing to cofibers in the following diagram:

$$\begin{array}{ccccc} S^{n-1} & \longrightarrow & D^n & \longrightarrow & S^n \\ \varphi_\alpha \downarrow & & \downarrow & & \downarrow \iota_\alpha \\ X^{n-1} & \longrightarrow & X^n & \longrightarrow & X^n/X^{n-1}. \end{array}$$

By the naturality of the boundary map, it follows that the top square in the following diagram commutes:

$$\begin{array}{ccccc}
\tilde{H}_n(S^n) & \xrightarrow[\cong]{\partial_n} & \tilde{H}_{n-1}(S^{n-1}) & & \\
(\iota_\alpha)_* \downarrow & & \downarrow (\varphi_\alpha)_* & \searrow (\Delta_{\alpha\beta})_* & \\
\tilde{H}_n(X^n/X^{n-1}) & \xrightarrow{\partial_n} & H_{n-1}(X^{n-1}) & & \\
& \searrow d_n & \downarrow q_* & & \\
& & \tilde{H}_{n-1}(X^{n-1}/X^{n-2}) & \xrightarrow{(p_\beta)_*} & \tilde{H}_{n-1}(S_\beta^{n-1}).
\end{array}$$

It follows that the image under d_n of the generator e_α^n is the same as the image of $[\iota_{n-1}] \in \tilde{H}_{n-1}(S^{n-1})$ of the middle vertical map. Under the isomorphism $\tilde{H}_{n-1}(X^{n-1}/X^{n-2}) \cong \bigoplus_{\gamma \in I_{n-1}} \tilde{H}_{n-1}(S_\gamma^{n-1})$, the β -th component is obtained by applying the projection $(p_\beta)_*$. We conclude that $d_{\alpha\beta}$ is the coefficient of the map $(p_\beta \circ q \circ \varphi_\alpha)_* = (\Delta_{\alpha\beta})_*: H_{n-1}(S^{n-1}) \rightarrow H_{n-1}(S^{n-1})$. By Remark 3.3.16, this is $\deg(\Delta_{\alpha\beta})$. $\square$

For $n = 1$, the boundary of a 1-cell is the difference of the classes of its two endpoints. This follows directly from the connecting homomorphism for the pair (X^1, X^0) . In particular, this boundary vanishes when X^0 consists of a single point.

Example 3.3.18 (Orientable surfaces). Let M_g be the closed orientable surface of genus g . It admits a CW-structure with one 0-cell, $2g$ one-cells $a_1, b_1, \dots, a_g, b_g$, and one 2-cell attached along the word $[a_1, b_1] \cdots [a_g, b_g] = a_1 b_1 a_1^{-1} b_1^{-1} \cdots a_g b_g a_g^{-1} b_g^{-1}$. The cellular chain complex is

$$0 \rightarrow \mathbb{Z} \xrightarrow{d_2} \mathbb{Z}^{2g} \xrightarrow{d_1} \mathbb{Z} \rightarrow 0.$$

The boundary d_1 is zero because each one-cell starts and ends at the unique zero-cell. For d_2 , each generator a_i appears in the attaching word with total exponent $1 + (-1) = 0$, and similarly for b_i . Thus each map $\Delta_{\alpha\beta}$ has degree zero, giving $d_2 = 0$. Since both boundary maps vanish, we obtain $H_0(M_g) \cong \mathbb{Z}$, $H_1(M_g) \cong \mathbb{Z}^{2g}$, and $H_2(M_g) \cong \mathbb{Z}$.

Example 3.3.19 (Non-orientable surfaces). Let N_g be the closed non-orientable surface of genus g (the connected sum of g copies of $\mathbb{RP}^2$). It admits a CW-structure with one 0-cell, g one-cells $a_1, \dots, a_g$, and one 2-cell attached along the word $a_1^2 a_2^2 \cdots a_g^2$. The cellular chain complex is

$$0 \rightarrow \mathbb{Z} \xrightarrow{d_2} \mathbb{Z}^g \xrightarrow{d_1} \mathbb{Z} \rightarrow 0.$$

Again $d_1 = 0$, since each one-cell starts and ends at the unique zero-cell. For d_2 , each a_i appears with total exponent 2, so the degree of each $\Delta_{\alpha\beta}$ equals 2. Thus $d_2(1) = (2, 2, \dots, 2) \in \mathbb{Z}^g$. Since d_2 is injective, $H_2(N_g) = 0$. To compute $H_1(N_g) = \ker(d_1)/\text{im}(d_2) = \mathbb{Z}^g / \langle (2, \dots, 2) \rangle$, we change basis: replacing the last standard basis vector $(0, \dots, 0, 1)$ by $(1, \dots, 1)$, we see that $H_1(N_g) \cong \mathbb{Z}^{g-1} \oplus \mathbb{Z}/2\mathbb{Z}$.

Example 3.3.20 (Real projective space). Real projective space $\mathbb{RP}^n$ has a CW-structure with one cell in each dimension $0, 1, \dots, n$. The attaching map $\varphi_k: S^{k-1} \rightarrow \mathbb{RP}^{k-1}$ for the k -cell is the quotient map identifying antipodal points. One may compute using geometric methods that the composite $\Delta: S^{k-1} \xrightarrow{\varphi_k} \mathbb{RP}^{k-1} \rightarrow \mathbb{RP}^{k-1}/\mathbb{RP}^{k-2} \simeq S^{k-1}$ has degree $1 + (-1)^k$: the two preimages of a generic

point contribute with the same sign when k is even and opposite signs when k is odd. Thus the cellular boundary maps are

$$d_k = \begin{cases} 0 & \text{if } k \text{ is odd,} \\ 2 & \text{if } k \text{ is even.} \end{cases}$$

This gives

$$H_k(\mathbb{R}P^n) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0, \\ \mathbb{Z}/2\mathbb{Z} & \text{if } 0 < k < n \text{ and } k \text{ is odd,} \\ \mathbb{Z} & \text{if } k = n \text{ and } n \text{ is odd,} \\ 0 & \text{otherwise.} \end{cases}$$

3.4 Brown representability (online only)

We now prove the Brown representability theorem stated in Theorem 3.2.6. The proof has two parts. We first prove Brown's general criterion for a set-valued functor on connected pointed animae. We then apply it degreewise to a reduced cohomology theory and assemble the representing animae into a spectrum.

3.4.1 The representability theorem

Write $\text{An}_{*, \geq 1}$ for the full subcategory of An_* spanned by the connected pointed animae. Since π_0 preserves colimits, this subcategory is closed under wedges and pushouts.

Definition 3.4.1 (Brown's conditions). Let

$$F: \text{Ho}(\text{An}_{*, \geq 1})^{\text{op}} \longrightarrow \text{Set}$$

be a functor. We say that:

- (1) The functor F is *representable* if it is naturally isomorphic to $[-, Z]_*$ for some $Z \in \text{An}_{*, \geq 1}$;
- (2) The functor F satisfies the *wedge axiom* if, for every small collection $(X_i)_{i \in I}$ of connected pointed animae, the canonical map

$$F\left(\bigvee_{i \in I} X_i\right) \longrightarrow \prod_{i \in I} F(X_i)$$

is a bijection;

- (3) The functor F satisfies the *Mayer–Vietoris property* if every pushout square

$$\begin{array}{ccc} C & \xrightarrow{k} & A \\ l \downarrow & & \downarrow i \\ B & \xrightarrow{j} & X \end{array}$$

in $\text{An}_{*, \geq 1}$ induces a surjection

$$(i^*, j^*): F(X) \longrightarrow F(A) \times_{F(C)} F(B).$$

The wedge axiom for the empty collection implies that $F(\text{pt})$ is a singleton. It also gives the values of F on suspensions an algebraic structure which will be used in the proof.

Remark 3.4.2. Suppose that F satisfies the wedge axiom. The cogroup structure on ΣX from Remark 2.4.20 makes $F(\Sigma X)$ a group: the pinch map induces its multiplication, while the collapse and reflection maps induce its unit and inverses. The two cogroup structures on $\Sigma^2 X$ agree by the Eckmann–Hilton argument, so $F(\Sigma^2 X)$ is an abelian group. A natural transformation between two functors satisfying the wedge axiom preserves these group structures.

Theorem 3.4.3 (Brown representability, Brown [Bro62]). *A functor $F: \text{Ho}(\text{An}_{*, \geq 1})^{\text{op}} \rightarrow \text{Set}$ is representable if and only if it satisfies the wedge axiom and the Mayer–Vietoris property.*

For a connected pointed anima Z and an element $\xi \in F(Z)$, the Yoneda lemma associates to ξ a natural transformation

$$T_\xi: [X, Z]_* \longrightarrow F(X), \quad [f] \longmapsto f^*(\xi).$$

Brown’s construction produces a representing object by successively improving the behavior of this transformation on spheres.

Definition 3.4.4 (Universal element). Let $n \geq 1$. An element $\xi \in F(Z)$ is *n-universal* if

$$T_\xi: [S^k, Z]_* \longrightarrow F(S^k)$$

is a bijection for $1 \leq k < n$ and a surjection for $k = n$. It is *universal* if this map is a bijection for every $k \geq 1$.

Proposition 3.4.5 (Universal approximation). *Let F satisfy the wedge axiom and the Mayer–Vietoris property. For every connected pointed anima X and every $\eta \in F(X)$, there are a connected pointed anima Z_X , a pointed map $f_X: X \rightarrow Z_X$, and a universal element $\xi_X \in F(Z_X)$ such that $f_X^*(\xi_X) = \eta$.*

Proof. Step 1: Constructing increasingly universal elements. We construct a sequence

$$X =: Z_0 \xrightarrow{f_0} Z_1 \xrightarrow{f_1} Z_2 \longrightarrow \cdots$$

and elements $\xi_n \in F(Z_n)$ such that $f_n^*(\xi_{n+1}) = \xi_n$, the element ξ_n is n -universal for $n \geq 1$, and T_{ξ_n} is surjective on S^k for every $k \geq 1$.

Set $\xi_0 := \eta$ and define

$$Z_1 := X \vee \bigvee_{m \geq 1} \bigvee_{\gamma \in F(S^m)} S^m.$$

By the wedge axiom, there is a unique element $\xi_1 \in F(Z_1)$ whose restriction to X is η and whose restriction to the sphere indexed by $\gamma \in F(S^m)$ is γ . The inclusion of that sphere shows that T_{ξ_1} is surjective on S^m for every $m \geq 1$. In particular, ξ_1 is 1-universal.

Suppose now that $n \geq 1$ and that (Z_n, ξ_n) has been constructed. By Remark 3.4.2, the map

$$T_{\xi_n}: [S^n, Z_n]_* \longrightarrow F(S^n)$$

is a group homomorphism. Let K_n denote its kernel. Choose a representative $g: S^n \rightarrow Z_n$ of every class $[g] \in K_n$, and form the pushout

$$\begin{array}{ccc} \bigvee_{[g] \in K_n} S^n & \xrightarrow{\vee g} & Z_n \\ \downarrow & & \downarrow f_n \\ \text{pt} & \longrightarrow & Z_{n+1}. \end{array}$$

The restrictions of ξ_n to all the attaching spheres are zero. The Mayer–Vietoris property therefore supplies an element $\xi_{n+1} \in F(Z_{n+1})$ satisfying $f_n^*(\xi_{n+1}) = \xi_n$.

The map $T_{\xi_{n+1}}$ remains surjective on every sphere, since T_{ξ_n} is surjective and $\xi_n = f_n^* \xi_{n+1}$. Attaching $(n+1)$ -cells induces bijections

$$[S^k, Z_n]_* \xrightarrow{\cong} [S^k, Z_{n+1}]_*$$

for $k < n$, so ξ_{n+1} remains n -universal. Finally, every map $S^n \rightarrow Z_{n+1}$ factors up to homotopy through Z_n . If its class lies in the kernel of $T_{\xi_{n+1}}$, a lift to Z_n lies in K_n and hence becomes null-homotopic after the corresponding cell is attached. Thus $T_{\xi_{n+1}}$ is injective on S^n , proving that ξ_{n+1} is $(n+1)$ -universal.

Step 2: Passing to the limit. Set

$$Z_X := \text{colim}_n Z_n.$$

The mapping-telescope description from Exercise 2.4.28 gives a pushout square

$$\begin{array}{ccc} \bigvee_{n \geq 0} Z_n & \longrightarrow & \bigvee_{r \geq 0} Z_{2r} \\ \downarrow & & \downarrow \\ \bigvee_{r \geq 0} Z_{2r+1} & \longrightarrow & Z_X. \end{array}$$

On the summand Z_n , the two maps are the identity and the transition map to the adjacent stage, assigned to the even and odd target wedges according to the parity of n . The relations $f_n^*(\xi_{n+1}) = \xi_n$, together with the wedge axiom and the Mayer–Vietoris property, therefore produce an element $\xi_X \in F(Z_X)$ restricting to ξ_n on every stage. In particular, the composite $f_X: X = Z_0 \rightarrow Z_X$ satisfies $f_X^*(\xi_X) = \eta$.

For a fixed $k \geq 1$, the map $Z_n \rightarrow Z_X$ is obtained by attaching cells of dimension at least $n+1$. Hence Remark 3.3.6 shows that

$$[S^k, Z_n]_* \longrightarrow [S^k, Z_X]_*$$

is a bijection once $n > k$. Since ξ_n is n -universal, T_{ξ_X} is consequently a bijection on S^k . Thus ξ_X is universal. $\square$

Proof of Theorem 3.4.3. If $F \cong [-, Z]_*$ is representable, the wedge axiom follows from the universal property of coproducts. Mapping a pushout into Z gives a pullback of mapping animae. A pair of homotopy classes whose restrictions to the intersection agree can be represented together with a homotopy between those restrictions, and therefore lifts to a point of this pullback. This proves the Mayer–Vietoris property.

Conversely, suppose that F satisfies Brown's conditions. Apply Proposition 3.4.5 to $X = \text{pt}$ and the unique element of $F(\text{pt})$. We obtain a connected pointed anima Z and a universal element $\xi \in F(Z)$. We claim that

$$T_\xi : [X, Z]_* \longrightarrow F(X)$$

is a bijection for every connected pointed anima X .

For surjectivity, let $\eta \in F(X)$ and let $\tilde{\eta} \in F(X \vee Z)$ correspond under the wedge axiom to (η, ξ) . Applying Proposition 3.4.5 to $(X \vee Z, \tilde{\eta})$ produces a universal element $\tilde{\xi} \in F(\tilde{Z})$ and a map $X \vee Z \rightarrow \tilde{Z}$ carrying $\tilde{\xi}$ to $\tilde{\eta}$. Its restriction $g : Z \rightarrow \tilde{Z}$ induces isomorphisms on all positive homotopy groups because both ξ and $\tilde{\xi}$ are universal. Since both anima are connected, g is an isomorphism by Proposition 2.4.21. Composing the restriction $X \rightarrow \tilde{Z}$ with an inverse of g gives a class in $[X, Z]_*$ which maps to η .

For injectivity, suppose that $f, g : X \rightarrow Z$ satisfy $f^*\xi = g^*\xi =: \eta$. Form the pushout

$$\begin{array}{ccc} X \vee X & \xrightarrow{(f,g)} & Z \\ \nabla \downarrow & & \downarrow \\ X & \longrightarrow & \tilde{X}, \end{array}$$

where ∇ is the fold map. The Mayer–Vietoris property gives an element $\tilde{\eta} \in F(\tilde{X})$ restricting to ξ on Z and to η on X . Apply Proposition 3.4.5 to $(\tilde{X}, \tilde{\eta})$. As in the surjectivity argument, the resulting composite $Z \rightarrow \tilde{X} \rightarrow \tilde{Z}$ is an isomorphism because it carries one universal element to another. The pushout square then shows, after composing with its inverse, that both f and g are homotopic to the same map $X \rightarrow Z$. Hence $[f] = [g]$. $\square$

3.4.2 From representing anima to spectra

We now apply the abstract theorem to reduced cohomology theories. The only condition which does not follow immediately from the axioms is the Mayer–Vietoris property.

Lemma 3.4.6. *Let (h^*, σ) be a reduced cohomology theory. For each $n \in \mathbb{Z}$, the restriction*

$$h^n : \text{Ho}(\text{An}_{*, \geq 1})^{\text{op}} \longrightarrow \text{Ab} \longrightarrow \text{Set}$$

satisfies the Mayer–Vietoris property.

Proof. Consider a pushout square of connected pointed anima

$$\begin{array}{ccc} C & \xrightarrow{k} & A \\ l \downarrow & & \downarrow i \\ B & \xrightarrow{j} & X. \end{array}$$

The pushout property gives an isomorphism $\text{cofib}(k) \xrightarrow{\sim} \text{cofib}(j)$. The long exact cohomology sequences of Exercise 3.1.3 therefore fit into a commutative diagram

$$\begin{array}{ccccccc} h^n(\text{cofib}(j)) & \longrightarrow & h^n(X) & \xrightarrow{j^*} & h^n(B) & \longrightarrow & h^{n+1}(\text{cofib}(j)) \\ \cong \downarrow & & \downarrow i^* & & \downarrow l^* & & \downarrow \cong \\ h^n(\text{cofib}(k)) & \longrightarrow & h^n(A) & \xrightarrow{k^*} & h^n(C) & \longrightarrow & h^{n+1}(\text{cofib}(k)). \end{array}$$

Suppose that $\alpha \in h^n(A)$ and $\beta \in h^n(B)$ satisfy $k^*\alpha = l^*\beta$. The boundary of β vanishes, since its image under the right-hand isomorphism is the boundary of $l^*\beta = k^*\alpha$. Hence β lifts to some $\gamma_0 \in h^n(X)$. The element $\alpha - i^*\gamma_0$ lies in the kernel of k^* , so it comes from $h^n(\text{cofib}(k))$. Transporting a preimage across the left-hand isomorphism and then mapping it to $h^n(X)$ gives an element γ_1 whose restriction to B vanishes and whose restriction to A is $\alpha - i^*\gamma_0$. Thus $\gamma_0 + \gamma_1$ restricts to (α, β) , proving the required surjectivity. $\square$

Proof of Theorem 3.2.6. Let (h^*, σ) be a reduced cohomology theory. For each $m \geq 0$, the restriction of h^m to connected pointed animae satisfies the wedge axiom by definition and the Mayer–Vietoris property by Lemma 3.4.6. The abstract Brown representability theorem, Theorem 3.4.3, gives a connected pointed anima Y_m and a natural isomorphism

$$h^m(X) \cong [X, Y_m]_*$$

on connected pointed animae.

Set

$$E_m := \Omega Y_{m+1} \quad (m \geq 0).$$

Since ΣX is connected for every pointed anima X , the suspension isomorphism gives natural isomorphisms

$$h^m(X) \cong h^{m+1}(\Sigma X) \cong [\Sigma X, Y_{m+1}]_* \cong [X, E_m]_*$$

for every pointed anima X . Thus E_m represents h^m on all of An_* . The same calculation shows that ΩE_{m+1} also represents h^m . The Yoneda lemma therefore supplies an isomorphism

$$E_m \xrightarrow{\sim} \Omega E_{m+1}$$

compatible with the given suspension isomorphism. These isomorphisms make $(E_m)_{m \geq 0}$ into a spectrum E . The representing isomorphisms identify h^k with $\tilde{E}^k$ for $k \geq 0$, and for $k < 0$ we obtain

$$\tilde{E}^k(X) = [X, \Omega^{-k} E_0]_* \cong h^0(\Sigma^{-k} X) \cong h^k(X).$$

This proves the first assertion.

Now let $\alpha: \tilde{E}^* \rightarrow \tilde{F}^*$ be a natural transformation commuting with suspension. For each $m \geq 0$, the degree- m component is a natural transformation

$$[-, E_m]_* \longrightarrow [-, F_m]_*,$$

so Exercise 3.1 identifies it with postcomposition by a unique homotopy class of pointed maps $f_m: E_m \rightarrow F_m$. Compatibility with suspension says precisely that the squares

$$\begin{array}{ccc} E_m & \xrightarrow{\sim} & \Omega E_{m+1} \\ f_m \downarrow & & \downarrow \Omega f_{m+1} \\ F_m & \xrightarrow{\sim} & \Omega F_{m+1} \end{array}$$

commute in $\text{Ho}(\text{An}_*)$. Choosing representatives and the corresponding homotopies gives a morphism of spectra $f: E \rightarrow F$ in the sense of Definition 3.2.2. Its induced natural transformation is α in nonnegative degrees, and hence in every degree by compatibility with suspension. $\square$

Exercise 3.4.7 (Mayer–Vietoris sequence). Let $F: \text{Ho}(\text{An}_{*, \geq 1})^{\text{op}} \rightarrow \text{Ab}$ satisfy the wedge axiom and the Mayer–Vietoris property, and consider a pushout square of connected pointed animae

$$\begin{array}{ccc} C & \xrightarrow{k} & A \\ \downarrow i & & \downarrow i \\ B & \xrightarrow{j} & X. \end{array}$$

Show that the cofiber of $(i, j): A \vee B \rightarrow X$ is isomorphic to ΣC , and use the coPuppe sequence to construct a long exact Mayer–Vietoris sequence

$$\cdots \rightarrow F(\Sigma X) \rightarrow F(\Sigma A) \times F(\Sigma B) \rightarrow F(\Sigma C) \rightarrow F(X) \rightarrow F(A) \times F(B) \rightarrow F(C).$$

Determine the maps and signs explicitly.

Exercises

Exercise 3.1. Use the Yoneda lemma to show that for two pointed animae X and Y , any natural transformation $[-, X]_* \rightarrow [-, Y]_*$ of functors $\text{Ho}(\text{An}_*)^{\text{op}} \rightarrow \text{Set}$ is given by postcomposition with a unique morphism $f: X \rightarrow Y$ in $\text{Ho}(\text{An}_*)$.

Exercise 3.2. Let C be an ∞ -category with a terminal object and pushouts. Consider a commutative square

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow \\ C & \xrightarrow{g} & D, \end{array}$$

and let $\text{cofib}(f) \rightarrow \text{cofib}(g)$ denote the induced map on cofibers. Show that this map on cofibers is an isomorphism whenever the square is a pushout square in C .

4 Spectra and stable ∞ -categories

In the previous chapter, we encountered spectra as objects representing generalized cohomology theories. We now shift perspective, and study the ∞ -category of spectra in its own right. Its formal properties are reminiscent of homological algebra, and lead to a notion of *stable ∞ -categories*.

As we will see, there are two equivalent perspectives on stable ∞ -categories. The first emphasizes the relation to the stability property of cohomology theories by considering the suspension and loop functors and asking them to be invertible. The second emphasizes the analogy with exact sequences in an abelian category by considering fibers and cofibers, the homotopical analogues of kernels and cokernels, and demanding that every nullsequence

$$X \rightarrow Y \rightarrow Z$$

is a fiber sequence if and only if it is a cofiber sequence. It is this second perspective that connects stable ∞ -categories to homological algebra.

The ∞ -category of spectra admits a universal property: it is the terminal stable ∞ -category equipped with a left exact functor to $\mathbf{An}$. This is a special case of the *stabilization* $\mathrm{Sp}(C)$ of an ∞ -category C with finite limits. We may think of $\mathrm{Sp}(C)$ as obtained from C by formally making the loops operation invertible.

Section 4.1 develops fibers, cofibers, suspensions and loops in pointed ∞ -categories. Section 4.2 then studies stable ∞ -categories themselves, and Section 4.3 constructs $\mathrm{Sp}(C)$ by means of prespectra and spectrification. Finally, Section 4.4 specializes to Sp and treats suspension spectra, mapping spectra, tensor products, homotopy groups, and the homology theories represented by spectra.

4.1 Pointed ∞ -categories

In Section 2.4 we introduced the basic constructions of fibers, cofibers, suspensions and loops in the ∞ -category $\mathbf{An}_*$ of pointed anima. In this section we develop the corresponding formalism for an arbitrary pointed ∞ -category C . This will provide the language needed to formulate the definition of a stable ∞ -category in Section 4.2, and will also be used repeatedly in later chapters.

Definition 4.1.1 (Pointed ∞ -category). We call an ∞ -category C *pointed* if it has a *zero object*, i.e. an object 0 that is both initial and terminal (Definition 1.7.7). Given objects $X, Y \in C$, we define the *null map* $0: X \rightarrow Y$ as the composite of the unique maps $X \rightarrow 0$ and $0 \rightarrow Y$.

A functor $F: C \rightarrow D$ between pointed ∞ -categories is called *pointed* if it preserves zero objects.

Example 4.1.2. The category $\mathbf{Ab}$ of abelian groups is pointed: the zero object is the trivial group 0 . More generally, any additive category is pointed.

Example 4.1.3. The category Top_* of pointed topological spaces is pointed, with the one-point space functioning as zero object.

Example 4.1.4. The ∞ -category An_* of pointed anima (see Section 2.4) is pointed, with the one-point anima as zero object.

Definition 4.1.5 (Fiber and cofiber sequences). Let C be a pointed ∞ -category. A *nullsequence* in C is a sequence of morphisms

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

equipped with a specified *nullhomotopy* $g \circ f \cong 0$. Equivalently, a nullsequence is a commutative square in C of the form

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow g \\ 0 & \longrightarrow & Z. \end{array}$$

If this square is a pullback square in C , we say it is a *fiber sequence*, and refer to X as the *fiber of g* , written $X \simeq \text{fib}(g)$. If the square is a pushout square, we say it is a *cofiber sequence*, and refer to Z as the *cofiber of f* , written $Z \simeq \text{cofib}(f)$.

We say that C *admits fibers* if such a fiber sequence exists for every morphism $g: Y \rightarrow Z$. Dually, we say that C *admits cofibers* if such a cofiber sequence exists for every morphism $f: X \rightarrow Y$.

Definition 4.1.6 (Loop and suspension). Let C be a pointed ∞ -category. If C admits fibers, we define the *loop object* of $X \in C$ by

$$\Omega X := \text{fib}(0 \rightarrow X).$$

If C admits cofibers, we define its *suspension* by

$$\Sigma X := \text{cofib}(X \rightarrow 0).$$

Just as in Exercise 2.4.7, the loop and suspension constructions define endofunctors $\Omega, \Sigma: C \rightarrow C$.

Lemma 4.1.7. *Let C be a pointed ∞ -category that admits fibers and cofibers. Then the functor $\Sigma: C \rightarrow C$ is left adjoint to $\Omega: C \rightarrow C$: for all $X, Y \in C$ there is a natural equivalence*

$$\text{Hom}_C(\Sigma X, Y) \simeq \text{Hom}_C(X, \Omega Y).$$

Proof. The proof is identical to that of Lemma 2.4.8. □

Pointed objects

The pointed ∞ -categories Top_* and An_* are a special case of a general construction for turning an ∞ -category C with terminal object into a pointed ∞ -category.

Definition 4.1.8 (Pointed objects). If C is an ∞ -category with a terminal object $*$, we define its *∞ -category of pointed objects* C_* as the slice category of C over $*$, i.e. as the following pullback square:

$$\begin{array}{ccc} C_* & \longrightarrow & \text{Ar}(C) \\ \downarrow & \lrcorner & \downarrow s \\ * & \xrightarrow{*} & C. \end{array}$$

Objects of C_* are objects X of C equipped with a map $x: * \rightarrow X$ from the terminal object, thought of as a chosen ‘basepoint’ of X . We will frequently denote such objects by (X, x) , or sometimes abusively as X .

A simple computation shows that the hom anima between pointed objects (X, x) and (Y, y) in C is computed as the following fiber:

$$\mathrm{Hom}_{C_*}((X, x), (Y, y)) \simeq \mathrm{fib}_y \left(\mathrm{Hom}_C(X, Y) \xrightarrow{- \circ x} \mathrm{Hom}_C(*, Y) \right).$$

(See Lemma 21.3.2 for a proof of the more general computation of hom animae in slice categories.) In other words, a morphism in C_* from (X, x) to (Y, y) consists of a morphism $f: X \rightarrow Y$ in C together with an isomorphism $f(x) \cong y$ of maps $* \rightarrow Y$.

Lemma 4.1.9. *Let C be an ∞ -category with terminal object $*$. Then:*

- (1) *The ∞ -category C_* is pointed, with zero object given by $0 := (*, \mathrm{id}_*: * \rightarrow *)$.*
- (2) *The ∞ -category C is pointed if and only if the forgetful functor*

$$\mathrm{fgt}: C_* = C_{*/} \longrightarrow C$$

is an equivalence.

Proof. (1) For initiality of 0 , note that the map $\mathrm{Hom}_C(*, Y) \rightarrow \mathrm{Hom}_C(*, Y)$ given by precomposing with the identity is an equivalence, hence its fiber $\mathrm{Hom}_{C_*}(0, Y)$ is contractible for every $Y \in C_*$. For terminality, we observe that

$$\mathrm{Hom}_{C_*}(X, 0) \simeq \mathrm{fib} \left(\mathrm{Hom}_C(X, *) \rightarrow \mathrm{Hom}_C(*, *) \right)$$

is the fiber of a map between two contractible animae, which thus is itself contractible.

(2) The ‘only if’ part is clear from (1). Conversely, assume that C is pointed. Then the object $*$ is also an initial object in C , and hence the target functor $C_* = C_{*/} \rightarrow C$ is an equivalence by Lemma 21.3.4. $\square$

Exercise 4.1.10. Show that C_* admits pullbacks whenever C admits pullbacks, and that the functor $C_* \rightarrow C$ preserves pullbacks in this case. Similarly, show that C_* admits pushouts whenever C admits pushouts, and that $C_* \rightarrow C$ preserves pushouts.

The pointed ∞ -category C_* is in a precise sense the ‘universal approximation’ of C by a pointed ∞ -category. We will end this section by recording a precise formulation of this universal property, which will be used later.

Notation 4.1.11. Let $\mathrm{Cat}_\infty^{\mathrm{term}} \subseteq \mathrm{Cat}_\infty$ denote the subcategory of ∞ -categories with a terminal object, and functors preserving the terminal object. Let $\mathrm{Cat}_\infty^{\mathrm{pt}} \subseteq \mathrm{Cat}_\infty^{\mathrm{term}}$ denote the full subcategory of pointed ∞ -categories. Given ∞ -categories C and D with terminal objects, we denote by

$$\mathrm{Fun}_*(C, D) \subseteq \mathrm{Fun}(C, D)$$

the full subcategory spanned by functors preserving the terminal object.

The constructions of C_* and the forgetful functor $\text{fgt}: C_* \rightarrow C$ are functorial in C , giving rise to a functor

$$(-)_*: \text{Cat}_\infty^{\text{term}} \rightarrow \text{Cat}_\infty^{\text{pt}}$$

and a natural transformation $\text{fgt}: (-)_* \rightarrow \text{id}$.

Corollary 4.1.12. *Let D be a pointed ∞ -category and let C be an ∞ -category with a terminal object. Then the ∞ -category $\text{Fun}_*(D, C)$ is pointed.*

Proof. The constant functor $\text{const}_*: D \rightarrow C$ is terminal in $\text{Fun}(D, C)$, hence also in the full subcategory $\text{Fun}_*(D, C)$. We claim that it is also initial. Let $0 \in D$ denote a zero object, and let $F: D \rightarrow C$ be a functor preserving terminal objects. Since 0 is also initial in D , limits over D are given by evaluation at 0 by Lemma 21.2.4. We therefore get

$$\text{Hom}_{\text{Fun}_*(D, C)}(\text{const}_*, F) \simeq \text{Hom}_C(*, \lim_{d \in D} F(d)) \simeq \text{Hom}_C(*, F(0)).$$

Since F preserves terminal objects and 0 is terminal in D , the object $F(0)$ is terminal in C . The last hom anima is therefore contractible, proving that const_* is initial. $\square$

Lemma 4.1.13. *Let D be pointed and let C have a terminal object. Then the forgetful functor $\text{fgt}: C_* \rightarrow C$ induces an equivalence*

$$\text{fgt}_*: \text{Fun}_*(D, C_*) \xrightarrow{\sim} \text{Fun}_*(D, C).$$

In particular, the functor $(-)_: \text{Cat}_\infty^{\text{term}} \rightarrow \text{Cat}_\infty^{\text{pt}}$ is right adjoint to the inclusion $\text{Cat}_\infty^{\text{pt}} \hookrightarrow \text{Cat}_\infty^{\text{term}}$.*

Proof. Since $C_* = C_{*/}$, there is an equivalence

$$\text{Fun}_*(D, C_*) \simeq \text{Fun}_*(D, C)_*.$$

Indeed, a natural transformation $\text{const}_* \rightarrow F$ is necessarily an isomorphism at the zero object as soon as F preserves terminal objects. By Corollary 4.1.12 and Lemma 4.1.9, the forgetful functor from the right-hand side to $\text{Fun}_*(D, C)$ is an equivalence. Passing to underlying anima exhibits $(-)_*$ as a right adjoint to the inclusion. $\square$

4.2 Stable ∞ -categories

We now come to the central definition of stable homotopy theory: that of a stable ∞ -category. To motivate the definition, let us take a look at some of the features of the category Ab of abelian groups which make it suitable for doing homotopical algebra:

- (1) The category Ab is *additive*: the direct sum $A \oplus B$ of two abelian groups is both a product and a coproduct in Ab , and the hom sets $\text{Hom}_{\text{Ab}}(A, B)$ inherit the structure of abelian groups;
- (2) The category Ab admits *kernels* and *cokernels*;

(3) There is a notion of *short exact sequence*: given a sequence of abelian groups

$$0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0$$

satisfying $pi = 0$ with i a monomorphism (i.e. an injective map) and p an epimorphism (i.e. a surjective map), the map i exhibits A as a kernel of p if and only if the map p exhibits C as a cokernel of A .

Categories with properties (1)-(3) are called *abelian categories*; we will study them more closely in Section 6.1.1. The definition of stable ∞ -categories closely mirrors that of abelian categories:

Definition 4.2.1 (Stable ∞ -category). An ∞ -category C is called *stable* if:

- (1) The ∞ -category C is pointed;
- (2) The ∞ -category C admits fibers and cofibers;
- (3) A nullsequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in C is a fiber sequence if and only if it is a cofiber sequence.

We refer to the nullsequences in (3) as *exact sequences*.¹¹ A functor $F: C \rightarrow D$ between two stable ∞ -categories is called *exact* if it preserves the zero object and sends exact sequences to exact sequences. Given two stable ∞ -categories C and D , we denote by

$$\mathrm{Fun}^{\mathrm{ex}}(C, D) \subseteq \mathrm{Fun}(C, D)$$

the full subcategory spanned by the exact functors.

Note that this definition differs in two crucial ways from the definition of an abelian category. First, in condition (3) there is no requirement for f and g to be monomorphisms or epimorphisms; they are allowed to be arbitrary morphisms in C . Second, in condition (1) we are merely assuming that C is pointed, and are not a priori requiring any additive structure. In some sense, condition (3) has absorbed the requirement of additivity: as we will show in Lemma 4.2.17, stable ∞ -categories are automatically additive.

While the definition of stable ∞ -categories we gave is the one that most closely resembles the definition of abelian categories, there are various equivalent characterizations of stable ∞ -categories that we will frequently use.

Theorem 4.2.2. *For a pointed ∞ -category C , the following conditions are equivalent:*

- (1) *The ∞ -category C admits pushouts and pullbacks, and a commutative square*

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ h \downarrow & & \downarrow g \\ W & \xrightarrow{k} & Z \end{array}$$

in C is a pushout square if and only if it is a pullback square.

¹¹Alternative terminology for ‘exact sequence’ includes *bifiber sequence*, *exact triangle* and *distinguished triangle*.

- (2) The ∞ -category C is stable;
- (3) The ∞ -category C admits fibers and the loop functor $\Omega: C \rightarrow C$ is an equivalence;
- (4) The ∞ -category C admits cofibers and the suspension functor $\Sigma: C \rightarrow C$ is an equivalence.

Proof. It is clear that (1) implies (2) by taking $W = 0$. If we assume (2), then the fiber sequence

$$\begin{array}{ccc} \Omega X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & X \end{array}$$

is by assumption also a pushout square and thus the counit map $\Sigma\Omega X \rightarrow X$ is an equivalence. Similarly the unit map $X \rightarrow \Omega\Sigma X$ is an equivalence. It follows that Σ and Ω are inverse equivalences, so that (3) and (4) hold.

We now show that (3) implies (1); applying the argument to C^{op} will then show that also (4) implies (1), finishing the argument. We will first show that C admits pushouts and that pushout squares agree with pullback squares. To this end, consider the full subcategory

$$P \subseteq \text{Fun}([1] \times [1], C)$$

spanned by the pullback squares. We then have the following claim:

Claim: The restriction functor $P \rightarrow \text{Fun}(\sqcap, C)$ is an equivalence.

Proof of claim: To construct an inverse, consider a diagram $z \leftarrow x \rightarrow y$ in C , and consider the following pullback diagram (ignoring the dashed morphisms):

$$\begin{array}{ccccccc} \Omega x & \longrightarrow & \Omega z & \longrightarrow & 0 & & \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & & \\ \Omega y & \longrightarrow & a & \longrightarrow & b & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow & \lrcorner & \\ 0 & \longrightarrow & c & \longrightarrow & x & \longrightarrow & y \\ & & \downarrow & \lrcorner & \downarrow & & \downarrow \\ & & 0 & \longrightarrow & z & \dashrightarrow & \Omega^{-1}a. \end{array}$$

Note that each of the pullbacks that appear here may be obtained as a fiber, hence is assumed to exist in C . More explicitly, starting from the span $z \leftarrow x \rightarrow y$, set

$$b := \text{fib}(x \rightarrow y), \quad c := \text{fib}(x \rightarrow z).$$

Then define a as the fiber of the composite $b \rightarrow x \rightarrow z$. The nullhomotopy of $a \rightarrow b \rightarrow x \rightarrow z$ induces a map $a \rightarrow c$, and the square

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \downarrow \\ c & \longrightarrow & x \end{array}$$

is a pullback by the pasting law. The other displayed pullback squares are obtained in the same way from the fiber sequences $b \rightarrow x \rightarrow y$, $c \rightarrow x \rightarrow z$, and $a \rightarrow b \rightarrow z$. The construction is functorial in the input span because each step is obtained by applying the fiber construction to a functorially specified morphism, together with the universal maps in its defining pullback square. Thus a morphism of spans induces compatible morphisms of all the objects in the displayed diagram. We now claim that an inverse to the restriction functor is given by

$$\left(\begin{array}{ccc} x & \longrightarrow & y \\ \downarrow & & \\ z & & \end{array} \right) \mapsto \Omega^{-1} \left(\begin{array}{ccc} \Omega x & \longrightarrow & \Omega y \\ \downarrow \lrcorner & & \downarrow \\ \Omega z & \longrightarrow & a \end{array} \right).$$

Note that this functor does indeed land in P since the equivalence $\Omega^{-1}: C \xrightarrow{\sim} C$ preserves pullback squares. Furthermore, it provides an extension of the original diagram, hence defines a right inverse. Finally, if we start with any pullback square, we may rewrite it in the form

$$\begin{array}{ccc} x & \longrightarrow & y \\ \downarrow \lrcorner & & \downarrow \\ z & \longrightarrow & \Omega^{-1}a, \end{array}$$

and we see that the above procedure recovers the original pullback square, showing that it is also a left inverse. This finishes the proof of the claim. $\square$

We will now use the claim to produce pushouts in C . Consider again a diagram $z \leftarrow x \rightarrow y$ in C , and use the claim to extend it uniquely to a pullback diagram:

$$\begin{array}{ccc} x & \longrightarrow & y \\ \downarrow \lrcorner & & \downarrow \\ z & \longrightarrow & w. \end{array}$$

We claim that this square is also a pushout square in C . Indeed, for every other object a in C we have natural equivalences

$$\text{Nat} \left(\begin{array}{ccc} x & \longrightarrow & y \\ \downarrow & & \\ z & & \end{array}, \begin{array}{ccc} a & \xlongequal{\quad} & a \\ \parallel & & \\ a & \xlongequal{\quad} & a \end{array} \right) \simeq \text{Nat} \left(\begin{array}{ccc} x & \longrightarrow & y \\ \downarrow \lrcorner & & \downarrow \\ z & \longrightarrow & w \end{array}, \begin{array}{ccc} a & \xlongequal{\quad} & a \\ \parallel \lrcorner & & \parallel \\ a & \xlongequal{\quad} & a \end{array} \right) \simeq \text{Hom}_C(w, a),$$

where the first equivalence is induced from the equivalence $P \simeq \text{Fun}(\lrcorner, C)$, whereas the second equivalence holds because $[1] \times [1]$ has a terminal object $(1, 1)$, so that the functor $\text{ev}_{(1,1)}: \text{Fun}([1] \times [1], C) \rightarrow C$ is left adjoint to the constant diagram functor $C \rightarrow \text{Fun}([1] \times [1], C)$. In particular, we see that C admits pushouts, and that every pullback square is a pushout square.

Note that in particular C admits cofibers and that $\Sigma: C \rightarrow C$ is an equivalence, so applying the above logic to C^{op} we see that C also admits all pullbacks, and that every pushout square is a pullback square. We conclude that condition (1) is satisfied, finishing the proof. $\square$

Terminology 4.2.3. Let C be a stable ∞ -category. We say that a commutative square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ h \downarrow & & \downarrow g \\ W & \xrightarrow{k} & Z \end{array}$$

is an *exact square* if it is a pullback square, or equivalently a pushout square.

Warning 4.2.4. Despite the similarity in their definitions, abelian categories are *never* stable ∞ -categories: the loop functor $\Omega: \mathcal{A} \rightarrow \mathcal{A}$ in a pointed 1-category $\mathcal{A}$ is necessarily the zero-functor, hence is only an equivalence if $\mathcal{A}$ is the trivial category.

Nevertheless, every abelian category $\mathcal{A}$ embeds fully faithfully into a stable ∞ -category $D(\mathcal{A})$, called its *derived ∞ -category*, to be introduced in Section 6.1 below. The inclusion $\mathcal{A} \hookrightarrow D(\mathcal{A})$ sends short exact sequences in $\mathcal{A}$ to exact sequences in $D(\mathcal{A})$.

Proposition 4.2.5. *Let $F: C \rightarrow D$ be a functor between stable ∞ -categories. Then the following are equivalent:*

- (1) *The functor F is exact;*
- (2) *The functor F is right exact, i.e. preserves initial objects and pushouts;*
- (3) *The functor F is left exact, i.e. preserves terminal objects and pullbacks.*

Proof. Since C and D are pointed, it is clear that F is pointed in each of the three cases. The argument from Theorem 4.2.2 shows that a pointed functor preserves exact sequences if and only if it preserves pushout squares if and only if it preserves pullback squares. $\square$

Remark 4.2.6. Conditions (1)-(3) on F are also equivalent to F preserving loop spaces, or equivalently suspensions; see Exercise 16.5.6.

Since exact functors are clearly closed under composition, the following definition makes sense:

Definition 4.2.7. We define the ∞ -category of *small stable ∞ -categories* as the (non-full) subcategory

$$\mathrm{Cat}_{\infty}^{\mathrm{st}} \subseteq \mathrm{Cat}_{\infty}$$

whose objects are the stable ∞ -categories and whose morphisms are the exact functors between them.

Lemma 4.2.8. *The ∞ -category $\mathrm{Cat}_{\infty}^{\mathrm{st}}$ admits small limits, and the inclusion $\mathrm{Cat}_{\infty}^{\mathrm{st}} \hookrightarrow \mathrm{Cat}_{\infty}$ preserves small limits.*

Proof. Let $C_{\bullet}: K \rightarrow \mathrm{Cat}_{\infty}^{\mathrm{st}}$ be a diagram of stable ∞ -categories and exact functors. By Corollary 23.5.5, the limit $\lim_{k \in K} C_k$ admits an initial and terminal object, and admits pushouts and pullbacks. Furthermore, each of the forgetful functors $\lim_{k \in K} C_k \rightarrow C_k$ preserve these (co)limits. Stability of $\lim_{k \in K} C_k$ now follows directly from the stability of the ∞ -categories C_k for all $k \in K$. $\square$

Exercise 4.2.9. Let I and C be ∞ -categories and assume C is stable. Show that also $\mathrm{Fun}(I, C)$ is stable. *Hint:* Use Lemma 21.2.8.

4.2.1 Shift functors

Let C be a stable ∞ -category. Since $\Sigma: C \rightarrow C$ and $\Omega: C \rightarrow C$ are inverse equivalences, we can define shift functors for any integer n .

Definition 4.2.10 (Shift functors). For $n \in \mathbb{Z}$, we define the n -th shift functor $[n]: C \rightarrow C$ by

$$[n] := \begin{cases} \Sigma^n & \text{if } n \geq 0 \\ \Omega^{-n} & \text{if } n \leq 0 \end{cases}.$$

The two formulas agree for $n = 0$.

Warning 4.2.11. While the shift notation is convenient, it should be used with care to avoid sign issues.

For integers n and m , one may identify the composite $[n] \circ [m]$ with $[n + m]$ by explicitly checking each of the four cases. In particular, this shows that $[n] \circ [m]$ is equivalent to $[m] \circ [n]$: both are shifting by the sum $m + n$.

The subtlety is that there are several inequivalent ways to exhibit this identification. For example, since $[n]: C \rightarrow C$ is exact, it commutes with $[m]$, giving another identification $[n] \circ [m] \simeq [m] \circ [n]$. To see that this identification differs from the previous one, let us consider the case $n = m = 1$, so that both shift functors are simply the suspension functor. The first identification is then nothing but the identity on $\Sigma^2: C \rightarrow C$. However, the second identification $\Sigma^2 \simeq \Sigma^2$ is the one coming from commutation of Σ with itself. To see that these are not the same, consider the case $C = \text{An}_*$, for which the commutation map is obtained by forming smash products with the swap map $S^1 \wedge S^1 \xrightarrow{\sim} S^1 \wedge S^1$ from Definition 2.4.10. Since this map has degree -1 , it is not homotopic to the identity.

We will now discuss various basic facts about stable ∞ -categories that illustrate their general behavior.

Lemma 4.2.12. If $X \xrightarrow{f} Y \xrightarrow{g} Z$ is an exact sequence in a stable ∞ -category C , then for any integer n , the sequence

$$X[n] \xrightarrow{f[n]} Y[n] \xrightarrow{g[n]} Z[n]$$

is also exact. Furthermore, the sequences

$$Y \xrightarrow{g} Z \rightarrow X[1] \quad \text{and} \quad Z[-1] \rightarrow X \xrightarrow{f} Y$$

are exact sequences.

Proof. Since $[n]$ is an equivalence, it preserves finite limits and colimits, hence exact sequences. The second statement follows from the following pushout/pullback diagrams:

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \longrightarrow & 0 \\ \downarrow & & \downarrow g & \lrcorner & \downarrow \\ 0 & \longrightarrow & Z & \longrightarrow & X[1] \end{array} \quad \text{and} \quad \begin{array}{ccccc} Z[-1] & \longrightarrow & X & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow f & \lrcorner & \downarrow \\ 0 & \longrightarrow & Y & \xrightarrow{g} & Z. \end{array}$$

Here we used the pasting law for pushouts/pullbacks to identify the objects $X[1]$ and $Z[-1]$. $\square$

Lemma 4.2.13. *Let $f: X \rightarrow Y$ be a morphism in a stable ∞ -category. Then there is an isomorphism $\mathrm{cofib}(f) \cong \mathrm{fib}(f)[1]$.*

Proof. Consider the following diagram:

$$\begin{array}{ccccc} \mathrm{fib}(f) & \longrightarrow & X & \longrightarrow & 0 \\ \downarrow & \lrcorner & \downarrow f & & \downarrow \\ 0 & \longrightarrow & Y & \longrightarrow & \mathrm{cofib}(f). \end{array}$$

The left square is a pullback square, and hence by stability also a pushout square. The right square is by definition a pushout square. Hence the outer rectangle is a pushout square, giving the desired isomorphism $\mathrm{cofib}(f) \simeq \Sigma(\mathrm{fib}(f)) = \mathrm{fib}(f)[1]$. $\square$

Lemma 4.2.14. *Let $f: X \rightarrow Y$ be a morphism in a stable ∞ -category C . The following are equivalent:*

- (1) *The morphism f is an isomorphism;*
- (2) *We have $\mathrm{cofib}(f) \simeq 0$;*
- (3) *We have $\mathrm{fib}(f) \simeq 0$.*

Proof. We show that (1) is equivalent to (2); the equivalence between (1) and (3) is dual. Consider the following commutative square:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & \lrcorner & \downarrow \\ 0 & \longrightarrow & \mathrm{cofib}(f). \end{array}$$

Since this square is both a pushout square as well as a pullback square and isomorphisms are closed under pushouts and pullbacks, we see that f is an isomorphism if and only if the map $0 \rightarrow \mathrm{cofib}(f)$ is an isomorphism, proving the claim. $\square$

4.2.2 The additive structure of stable ∞ -categories

Stable ∞ -categories possess a strong additive structure.

Definition 4.2.15 ((Semi)additive ∞ -category). An ∞ -category C is *semiadditive* if it has a zero object, finite products, finite coproducts, and the canonical map

$$\begin{pmatrix} \mathrm{id}_X & 0 \\ 0 & \mathrm{id}_Y \end{pmatrix}: X \sqcup Y \rightarrow X \times Y$$

is an isomorphism. We write $X \oplus Y$ for the resulting biproducts. We say that C is *additive* if, in addition, the shear map

$$\begin{pmatrix} \mathrm{id}_X & \mathrm{id}_X \\ 0 & \mathrm{id}_X \end{pmatrix}: X \oplus X \rightarrow X \oplus X$$

is an equivalence.

Semiadditivity is sometimes also called *preadditivity*.

Remark 4.2.16. For objects X, Y in a semiadditive ∞ -category C , we may equip $\pi_0 \operatorname{Hom}_C(X, Y)$ with the structure of an abelian monoid: given morphisms $f, g: X \rightarrow Y$, we define their sum $f + g$ as the following composite:

$$X \xrightarrow{(\operatorname{id}, \operatorname{id})} X \oplus X \xrightarrow{f \oplus g} Y \oplus Y \xrightarrow{(\operatorname{id})} Y.$$

We leave it to the reader to check that this is indeed unital, associative and commutative. Then C is *additive* if and only if $\pi_0 \operatorname{Hom}_C(X, Y)$ is in fact a *group*. In particular, for every morphism $f: X \rightarrow Y$ we may form its negative $-f: X \rightarrow Y$ which satisfies $f + (-f) = 0$.

Lemma 4.2.17. *Every stable ∞ -category C is additive.*

Proof. Stable ∞ -categories have zero objects and pullbacks/pushouts, and therefore finite products ($X \times Y = X \times_0 Y$) and coproducts ($X \sqcup Y = X \sqcup_0 Y$). Additivity of C then amounts to showing that for all objects X and Y in C , the two commutative squares

$$\begin{array}{ccc} 0 & \longrightarrow & X \\ \downarrow & & \downarrow (\operatorname{id}, 0) \\ Y & \xrightarrow{(0, \operatorname{id})} & X \times Y \end{array} \quad \text{and} \quad \begin{array}{ccc} 0 & \longrightarrow & X \\ \downarrow & & \downarrow (\operatorname{id}, \operatorname{id}) \\ X & \xrightarrow{(0, \operatorname{id})} & X \times X \end{array}$$

are exact. To see this, we consider the following two diagrams:

$$\begin{array}{ccccc} 0 & \longrightarrow & X & & \\ \downarrow & & \downarrow (\operatorname{id}, 0) & & \\ Y & \xrightarrow{(0, \operatorname{id})} & X \times Y & \xrightarrow{\operatorname{pr}_2} & Y \\ \downarrow & & \downarrow \operatorname{pr}_1 & & \downarrow \\ 0 & \longrightarrow & X & \longrightarrow & 0 \end{array} \quad \text{and} \quad \begin{array}{ccccc} 0 & \longrightarrow & X & & \\ \downarrow & & \downarrow (\operatorname{id}, \operatorname{id}) & & \\ X & \xrightarrow{(0, \operatorname{id})} & X \times X & \xrightarrow{\operatorname{pr}_2} & X \\ \downarrow & & \downarrow \operatorname{pr}_1 & & \downarrow \\ 0 & \longrightarrow & X & \longrightarrow & 0. \end{array}$$

In both cases, the right bottom square, the bottom rectangle and the left rectangle are pullback squares, and hence by the pasting law for pullback diagrams also the upper left square is exact. This finishes the proof. $\square$

Recall that the *splitting lemma* provides equivalent conditions for a short exact sequence in an abelian category to be a *split short exact sequence*. The following is an analogue of the splitting lemma for stable ∞ -categories:

Exercise 4.2.18 (Splitting lemma). Let $X \xrightarrow{i} Y \xrightarrow{p} Z$ be an exact sequence in a stable ∞ -category C . The following are equivalent:

- (1) The map i admits a retraction $r: Y \rightarrow X$ (i.e., $ri \simeq \operatorname{id}_X$);
- (2) The map p admits a section $s: Z \rightarrow Y$ (i.e., $ps \simeq \operatorname{id}_Z$);
- (3) There is an isomorphism $Y \cong X \oplus Z$ making the following diagram commute:

$$\begin{array}{ccccc} X & \xrightarrow{i} & Y & \xrightarrow{p} & Z \\ & \searrow & \downarrow \cong & \nearrow & \\ & & X \oplus Z & & \end{array}$$

(The arrows from X to $X \oplus Z$ and from Z to $X \oplus Z$ are labeled $(\operatorname{id}_X, 0)$ and pr_Z respectively.)

Hint. Given a retraction of i , paste the two resulting pushout squares to identify Y with $X \oplus Z$. The argument starting from a section of p is dual.

Exercise 4.2.19 (Recognizing exact squares). Let C be a stable ∞ -category and consider a commutative square in C :

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ u \downarrow & & \downarrow v \\ Z & \xrightarrow{g} & W. \end{array}$$

The following conditions are equivalent:

- (1) The square is exact;
- (2) The sequence $X \xrightarrow{(f,u)} Y \oplus Z \xrightarrow{\begin{pmatrix} v \\ -g \end{pmatrix}} W$ is an exact sequence.

Hint. Compare the original square with the exactness of the displayed sequence by means of the diagonal $W \rightarrow W \oplus W$ and the difference map $W \oplus W \rightarrow W$. The necessary comparison square is exact because its associated shear map is an isomorphism.

4.2.3 Finite limits and colimits

From the fact that stable ∞ -categories admit zero objects, pushouts and pullbacks, we may deduce that they in fact admit *all* finite limits and colimits.

Definition 4.2.20 (Finite ∞ -categories). We denote by

$$\mathrm{Cat}_{\infty}^{\mathrm{fin}} \subseteq \mathrm{Cat}_{\infty}$$

the smallest full subcategory which contains the ∞ -categories $\emptyset$, $*$ and $[1]$, and which is closed under pushouts of ∞ -categories. A small ∞ -category I is called *finite* if it is contained in $\mathrm{Cat}_{\infty}^{\mathrm{fin}}$.

Remark 4.2.21. It is proved by [Vol25, Proposition 2.4] that this model-independent definition of finite ∞ -categories is compatible with the definition used in quasicategory theory: an ∞ -category is finite if and only if it is the image of a finite simplicial set under the associated ∞ -category functor $\mathrm{ac}: \mathrm{sSet} \rightarrow \mathrm{Cat}_{\infty}$ from Construction 1.8.11.

Example 4.2.22. The following ∞ -categories are finite:

- The ∞ -categories $\emptyset$, $*$ and $[1]$;
- The ∞ -category $* \sqcup *$;
- The ∞ -categories $\sqcup$ and $\sqcap$;
- The ∞ -categories $[n]$.

Remark 4.2.23. There is a functor $(-)^{\mathrm{op}}: \mathrm{Cat}_{\infty} \rightarrow \mathrm{Cat}_{\infty}$ implementing the assignment $C \mapsto C^{\mathrm{op}}$. This functor is an equivalence, hence in particular preserves pushouts and full subcategories. Since each of the ∞ -categories $\emptyset$, $*$ and $[1]$ are equivalent to their opposites, it follows that an ∞ -category C is finite if and only if its opposite C^{op} is finite.

Definition 4.2.24 (Left exact ∞ -categories). An ∞ -category C is called *left exact*, or said to *admit finite limits*, if it admits all I -indexed limits for all finite ∞ -categories I . Similarly, a functor $F: C \rightarrow D$ between left exact ∞ -categories is called *left exact* if it preserves I -indexed limits for all $I \in \text{Cat}_{\infty}^{\text{fin}}$. We denote by

$$\text{Cat}_{\infty}^{\text{lex}} \subseteq \text{Cat}_{\infty}$$

the subcategory consisting of the left exact ∞ -categories and left exact functors. Given two left exact ∞ -categories C and D , we denote by

$$\text{Fun}^{\text{lex}}(C, D) \subseteq \text{Fun}(C, D)$$

the full subcategory spanned by the left exact functors.

One dually obtains a notion of *right exactness* in terms of finite colimits.

Theorem 4.2.25 (Lurie [Lur09b, Corollaries 4.4.2.4 and 4.4.2.5]). (1) *An ∞ -category C admits finite limits if and only if it has a terminal object and admits pullbacks.*

(2) *A functor $F: C \rightarrow D$ is left exact if and only if it preserves terminal objects and pullbacks.*

Dually C has finite colimits if and only if it has an initial object and admits pushouts, and F preserves finite colimits if and only if it preserves initial objects and pushouts.

Proof sketch. One direction is clear. The other direction needs the non-trivial fact that when the ∞ -category I can be written as a colimit $I = \text{colim}_j I_j$, we get an alternative description of $\lim_{i \in I} F(i)$ as $\lim_{j \in J^{\text{op}}} \lim_{i \in I_j} F(i)$. (See Theorem 21.2.11 for a more precise formulation). $\square$

Corollary 4.2.26. *Every stable ∞ -category admits finite limits and colimits.* $\square$

Corollary 4.2.27. *A functor $F: C \rightarrow D$ between stable ∞ -categories is exact if and only if it is left exact if and only if it is right exact: we have equalities*

$$\text{Fun}^{\text{ex}}(C, D) = \text{Fun}^{\text{lex}}(C, D) = \text{Fun}^{\text{rex}}(C, D)$$

as full subcategories of $\text{Fun}(C, D)$.

We record an alternative characterization of stable ∞ -categories, due to Moritz Groth [Gro16]:

Proposition 4.2.28. *An ∞ -category C is stable if and only if it has all finite limits and colimits and finite limits commute with finite colimits.*

Proof. Let C be a stable ∞ -category. If I is a finite ∞ -category, then C has I -indexed colimits by the previous corollary, hence there is a functor $\text{colim}: \text{Fun}(I, C) \rightarrow C$. Its domain is also stable and colim preserves finite colimits, because colimits commute with colimits. So the functor is exact and hence preserves finite limits as well.

Conversely, assume that C is finitely complete and cocomplete and that finite limits commute with finite colimits in C . Considering the empty category, the fact that empty limits commute with empty

colimits expresses that C is pointed. To show that C is stable, consider for every $X \in C$ the following diagram:

$$\begin{array}{ccccc}
 X & \longleftarrow & X & \longrightarrow & 0 \\
 \downarrow & & \downarrow & & \downarrow \\
 0 & \longleftarrow & X & \longrightarrow & 0 \\
 \uparrow & & \uparrow & & \uparrow \\
 0 & \longleftarrow & X & \longrightarrow & X.
 \end{array}$$

Taking pushouts of the rows we get the diagram $0 \rightarrow \Sigma X \leftarrow 0$ whose pullback is $\Omega \Sigma X$. If instead we take pullbacks of the columns, we get $X \leftarrow X \rightarrow X$, whose pushout is X . Pullbacks commuting with pushouts tell us then the canonical map $X \rightarrow \Omega \Sigma X$ is an equivalence. A similar argument shows that $\Sigma \Omega X \rightarrow X$ is an equivalence, so that C is stable by Theorem 4.2.2. $\square$

4.3 Stabilization

If C is an ∞ -category with finite limits, there is a stable ∞ -category $\mathrm{Sp}(C)$ that universally ‘approximates C from the left’ by a stable ∞ -category, known as the *stabilization of C* . The stabilization comes equipped with a left exact functor $\mathrm{Sp}(C) \rightarrow C$ satisfying the property that any other left exact functor $D \rightarrow C$ from a stable ∞ -category D uniquely factors through some left exact functor $D \rightarrow \mathrm{Sp}(C)$. The following definition captures this more precisely:

Definition 4.3.1. A *stabilization* of an ∞ -category C with finite limits is a stable ∞ -category $\mathrm{Sp}(C)$ equipped with a left exact functor $\Omega^\infty: \mathrm{Sp}(C) \rightarrow C$ such that for every stable ∞ -category D the composition functor

$$\Omega^\infty \circ -: \mathrm{Fun}^{\mathrm{ex}}(D, \mathrm{Sp}(C)) = \mathrm{Fun}^{\mathrm{lex}}(D, \mathrm{Sp}(C)) \rightarrow \mathrm{Fun}^{\mathrm{lex}}(D, C)$$

is an equivalence, where the first equality holds by Corollary 4.2.27.

Note that this universal property uniquely determines the stabilization whenever it exists; in fact, it determines a right adjoint $\mathrm{Sp}: \mathrm{Cat}_\infty^{\mathrm{lex}} \rightarrow \mathrm{Cat}_\infty^{\mathrm{st}}$ to the inclusion functor, cf. Corollary 4.3.17 below.

Example 4.3.2. The prototypical example is $C = \mathrm{An}$, whose stabilization $\mathrm{Sp}(\mathrm{An})$ is the ∞ -category of spectra Sp that we will study in Section 4.4.

We prove the existence of $\mathrm{Sp}(C)$ by providing an explicit construction: first we define prespectra as structured sequences in C_* , and then we define spectra as those prespectra whose structure maps are isomorphisms. We will prove the following properties of $\mathrm{Sp}(C)$ along the way:

- If C admits limits, then so does $\mathrm{Sp}(C)$ (Lemma 4.3.18);
- If C admits colimits and the loop functor $\Omega: C_* \rightarrow C_*$ preserves sequential colimits, then $\mathrm{Sp}(C)$ admits colimits (Lemma 4.3.20);
- In the latter case, the functor $\Omega^\infty: \mathrm{Sp}(C) \rightarrow C$ admits a left adjoint $\Sigma^\infty: C \rightarrow \mathrm{Sp}(C)$ (Lemma 4.3.22 and Corollary 4.3.23).

4.3.1 Prespectra and spectrum objects

The construction of (pre)spectra we use closely matches the classical definition of a spectrum as a sequence of pointed spaces $\{X_n\}$ equipped with maps $X_n \rightarrow \Omega X_{n+1}$. We first allow these maps to be arbitrary; the stabilization is obtained by imposing that they be isomorphisms.

Definition 4.3.3 (Prespectra). Let C be an ∞ -category admitting finite limits. Write

$$C_*^{\mathbb{N}} := \prod_{n \geq 0} C_*$$

for the ∞ -category of sequences of pointed objects of C . We define the shift functor and the levelwise loop functor by

$$\text{sh}: C_*^{\mathbb{N}} \rightarrow C_*^{\mathbb{N}}, \quad (X_n)_{n \geq 0} \mapsto (X_{n+1})_{n \geq 0},$$

and

$$\Omega: C_*^{\mathbb{N}} \rightarrow C_*^{\mathbb{N}}, \quad (X_n)_{n \geq 0} \mapsto (\Omega X_n)_{n \geq 0}.$$

The ∞ -category of *prespectra* in C is defined as

$$\text{PSp}(C) := C_*^{\mathbb{N}} \times_{C_*^{\mathbb{N}} \times C_*^{\mathbb{N}}} \text{Ar}(C_*^{\mathbb{N}}),$$

where the map $C_*^{\mathbb{N}} \rightarrow C_*^{\mathbb{N}} \times C_*^{\mathbb{N}}$ is $(\text{id}, \Omega \text{sh})$ and the map $\text{Ar}(C_*^{\mathbb{N}}) \rightarrow C_*^{\mathbb{N}} \times C_*^{\mathbb{N}}$ is the source-target functor (s, t) .

Unwinding definitions, a prespectrum X is a sequence $(X_n)_{n \geq 0}$ of pointed objects of C , together with structure maps

$$\sigma_n^X: X_n \rightarrow \Omega X_{n+1}.$$

A morphism $f: X \rightarrow Y$ of prespectra is a sequence of pointed maps $f_n: X_n \rightarrow Y_n$ together with homotopies filling the squares

$$\begin{array}{ccc} X_n & \xrightarrow{f_n} & Y_n \\ \sigma_n^X \downarrow & & \downarrow \sigma_n^Y \\ \Omega X_{n+1} & \xrightarrow{\Omega f_{n+1}} & \Omega Y_{n+1}. \end{array}$$

Definition 4.3.4 (Spectrum objects). Let C be an ∞ -category admitting finite limits. We define its ∞ -category of *spectrum objects*, or *stabilization*, as the full subcategory

$$\text{Sp}(C) \subseteq \text{PSp}(C)$$

spanned by those prespectra X for which every structure map $\sigma_n^X: X_n \rightarrow \Omega X_{n+1}$ is an isomorphism. We denote the evaluation functors by

$$(-)_n: \text{Sp}(C) \rightarrow C_*.$$

The functor $(-)_0: \text{Sp}(C) \rightarrow C_*$ will also be denoted by Ω^∞ ; when the target is written as C , we implicitly compose with the forgetful functor $C_* \rightarrow C$.

Remark 4.3.5. Assume C is a small ∞ -category. Then the structure maps $\sigma_n: (-)_n \xrightarrow{\sim} \Omega(-)_{n+1}$ turn the functors $(-)_n: \text{Sp}(C) \rightarrow C_*$ into a cone in Cat_∞ , resulting in a comparison functor

$$\text{Sp}(C) \rightarrow \lim(\cdots \rightarrow C_* \xrightarrow{\Omega} C_* \xrightarrow{\Omega} C_*).$$

This functor is an equivalence of ∞ -categories.

Lemma 4.3.6. *The functors $(-)_n: \mathbf{PSp}(C) \rightarrow C_*$ are jointly conservative. In particular, a morphism $f: X \rightarrow Y$ of spectra in C is an isomorphism if and only if each induced map $f_n: X_n \rightarrow Y_n$ is an isomorphism.*

Proof. The projection $\mathbf{PSp}(C) \rightarrow C_*^{\mathbb{N}}$ is conservative because a morphism in the arrow category is an isomorphism precisely when its source and target maps are isomorphisms. The projections $C_*^{\mathbb{N}} \rightarrow C_*$ are jointly conservative by definition of the product. $\square$

Lemma 4.3.7. *Let X and Y be prespectra in C . Then the hom anima $\mathrm{Hom}_{\mathbf{PSp}(C)}(X, Y)$ is the equalizer of the two maps*

$$\mathrm{Hom}_{C_*^{\mathbb{N}}}(X, Y) \rightrightarrows \mathrm{Hom}_{C_*^{\mathbb{N}}}(X, \Omega \mathrm{sh} Y).$$

The first map sends $f: X \rightarrow Y$ to the composite

$$X \xrightarrow{f} Y \xrightarrow{\sigma^Y} \Omega \mathrm{sh} Y,$$

and the second sends it to the composite

$$X \xrightarrow{\sigma^X} \Omega \mathrm{sh} X \xrightarrow{\Omega \mathrm{sh}(f)} \Omega \mathrm{sh} Y.$$

Proof. This is just the definition of $\mathbf{PSp}(C)$ as the lax equalizer of id and $\Omega \mathrm{sh}$. Indeed, a morphism $X \rightarrow Y$ in $\mathbf{PSp}(C)$ is a map of the underlying sequences together with a homotopy filling the square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \sigma^X \downarrow & & \downarrow \sigma^Y \\ \Omega \mathrm{sh} X & \xrightarrow{\Omega \mathrm{sh}(f)} & \Omega \mathrm{sh} Y. \end{array}$$

Equivalently, it is a point of the stated equalizer. $\square$

Lemma 4.3.8. *Let I be an ∞ -category.*

- (1) *If C admits I -indexed limits, then $\mathbf{PSp}(C)$ admits I -indexed limits and the projection $\mathbf{PSp}(C) \rightarrow C_*^{\mathbb{N}}$ creates them.*
- (2) *If C_* admits I -indexed colimits, then $\mathbf{PSp}(C)$ admits I -indexed colimits and the projection $\mathbf{PSp}(C) \rightarrow C_*^{\mathbb{N}}$ creates them.*

Proof. For limits, let $X: I \rightarrow \mathbf{PSp}(C)$ be a diagram and write $\overline{X}: I \rightarrow C_*^{\mathbb{N}}$ for the underlying diagram of sequences. Since C_* has I -indexed limits and Ω preserves limits, the endofunctor $\Omega \mathrm{sh}$ preserves I -indexed limits. Hence the limit $L := \lim_i \overline{X}_i$ in $C_*^{\mathbb{N}}$ inherits a structure map

$$L \simeq \lim_i \overline{X}_i \rightarrow \lim_i \Omega \mathrm{sh}(\overline{X}_i) \simeq \Omega \mathrm{sh}(L),$$

which makes it the limit in $\mathbf{PSp}(C)$.

For colimits, let $D := \mathrm{colim}_i \overline{X}_i$ in $C_*^{\mathbb{N}}$. For every i , the composite

$$\overline{X}_i \xrightarrow{\sigma^{X_i}} \Omega \mathrm{sh}(\overline{X}_i) \rightarrow \Omega \mathrm{sh}(D)$$

is compatible with the diagram, and therefore induces a structure map $D \rightarrow \Omega \text{sh}(D)$. We claim that the resulting prespectrum is the colimit. Indeed, for any prespectrum Y the hom anima $\text{Hom}_{\text{PSp}(C)}(D, Y)$ is the equalizer of the two maps

$$\text{Hom}_{C_*^{\mathbb{N}}}(D, Y) \rightrightarrows \text{Hom}_{C_*^{\mathbb{N}}}(D, \Omega \text{sh} Y)$$

given by postcomposition with σ^Y and precomposition with the structure map of D , by Lemma 4.3.7. Since D is the colimit of the $\overline{X}_i$ in $C_*^{\mathbb{N}}$, and since limits commute with limits in An , this equalizer identifies with

$$\lim_i \text{Hom}_{\text{PSp}(C)}(X_i, Y),$$

which is the desired universal property. $\square$

We now turn to the main point: the loop functor on $\text{Sp}(C)$ is an equivalence. The naive argument is to shift levels, since $X_n \simeq \Omega X_{n+1}$ by definition. The subtlety is that comparing the structure maps of X with those of $\Omega(\text{shift } X)$ requires identifying $\Omega^2 \Omega$ with $\Omega \Omega^2$, and there are two ways to do so. The next results handle this coherence issue by passing first to an even-level model and then showing that the resulting automorphism of Ω^3 is trivial.

Lemma 4.3.9. *Let $\text{PSp}^{(2)}(C)$ be the lax equalizer*

$$\text{PSp}^{(2)}(C) := C_*^{\mathbb{N}} \times_{C_*^{\mathbb{N}} \times C_*^{\mathbb{N}}} \text{Ar}(C_*^{\mathbb{N}}),$$

where the map $C_*^{\mathbb{N}} \rightarrow C_*^{\mathbb{N}} \times C_*^{\mathbb{N}}$ is $(\text{id}, \Omega^2 \text{sh})$ and the map from the arrow category is the source-target functor. Let $\text{Sp}^{(2)}(C) \subseteq \text{PSp}^{(2)}(C)$ be the full subcategory spanned by those objects whose structure maps are isomorphisms. Then the even-level functor

$$E : \text{Sp}(C) \rightarrow \text{Sp}^{(2)}(C), \quad E(X)_k := X_{2k},$$

with structure maps

$$\tilde{\sigma}_k^X : X_{2k} \xrightarrow{\sigma_{2k}^X} \Omega X_{2k+1} \xrightarrow{\Omega(\sigma_{2k+1}^X)} \Omega^2 X_{2k+2},$$

is an equivalence.

Proof. We construct an inverse. Given $Y \in \text{Sp}^{(2)}(C)$ with structure maps $\tau_k : Y_k \xrightarrow{\sim} \Omega^2 Y_{k+1}$, define a spectrum QY by

$$(QY)_{2k} := Y_k \quad \text{and} \quad (QY)_{2k+1} := \Omega Y_{k+1}.$$

The structure maps of QY are $\tau_k : Y_k \rightarrow \Omega^2 Y_{k+1} = \Omega(QY)_{2k+1}$ in even degrees and the identity map $\Omega Y_{k+1} \rightarrow \Omega Y_{k+1} = \Omega(QY)_{2k+2}$ in odd degrees.

It is immediate that $EQ = \text{id}$. Conversely, for $X \in \text{Sp}(C)$ there is a natural isomorphism $X \rightarrow QEX$ which is the identity in even degrees and the structure map $\sigma_{2k+1}^X : X_{2k+1} \rightarrow \Omega X_{2k+2}$ in odd degrees. The compatibility with the structure maps follows directly from the definition of $\tilde{\sigma}_k^X$. $\square$

Remark 4.3.10. When C is small, the limit description of $\text{Sp}(C)$ from Remark 4.3.5 identifies $\text{Sp}^{(2)}(C)$ with the analogous limit over the subdiagram indexed by the even natural numbers. Since these are cofinal, the result also follows formally from Theorem 21.5.1.

Next up, we need to understand the interaction of the double loop functor Ω^2 with the loop functor Ω . If D is a pointed ∞ -category with finite limits, the functor $\Omega^2: D \rightarrow D$ commutes with limits, and in particular commutes with the formation of loop objects. This provides a natural isomorphism

$$\Omega^2(\Omega X) \xrightarrow{\cong} \Omega(\Omega^2 X).$$

The following proposition records the specific case we need.

Proposition 4.3.11 (Universal property of finite animae). *Let $\mathbf{An}^{\text{fin}}$ denote the smallest full subcategory of $\mathbf{An}$ which contains the point and is closed under finite colimits. For every ∞ -category D with finite limits, evaluation at the point induces an equivalence*

$$\text{ev}_*: \text{Fun}^{\text{lex}}((\mathbf{An}^{\text{fin}})^{\text{op}}, D) \xrightarrow{\sim} D.$$

Proof. This is the opposite of the universal property of the free finite cocompletion of the point, which is a special case of Proposition 22.1.2. $\square$

Lemma 4.3.12 (Universal property of finite pointed animae). *Let $\mathbf{An}_*^{\text{fin}}$ be the small full subcategory of $\mathbf{An}_*$ generated under finite colimits by S^0 . For every pointed ∞ -category D with finite limits, evaluation at S^0 induces an equivalence*

$$\text{ev}_{S^0}: \text{Fun}^{\text{lex}}((\mathbf{An}_*^{\text{fin}})^{\text{op}}, D) \xrightarrow{\sim} D.$$

Proof. Precomposition with $(-)_+: \mathbf{An}^{\text{fin}} \rightarrow \mathbf{An}_*^{\text{fin}}$ defines a functor

$$\text{Fun}^{\text{lex}}((\mathbf{An}_*^{\text{fin}})^{\text{op}}, D) \longrightarrow \text{Fun}^{\text{lex}}((\mathbf{An}^{\text{fin}})^{\text{op}}, D).$$

An inverse sends F to the functor

$$(Y, y) \longmapsto \text{fib}(F(Y) \rightarrow F(*)).$$

This is pointed and left exact. The two inverse identities follow from $Y_+ = Y \sqcup *$ and from the pushout $Y_+ \sqcup_{S^0} * \simeq (Y, y)$. The result now follows from Proposition 4.3.11. $\square$

Proposition 4.3.13. *For any pointed ∞ -category D with finite limits, the composite map*

$$\Omega^3 X \simeq \Omega^2(\Omega X) \xrightarrow{\cong} \Omega(\Omega^2 X) \simeq \Omega^3 X$$

is homotopic to the identity map on $\Omega^3 X$, naturally in $X \in D$.

Proof. Given $X \in D$, Lemma 4.3.12 gives a pointed left exact functor

$$X^{(-)}: (\mathbf{An}_*^{\text{fin}})^{\text{op}} \rightarrow D$$

satisfying $X^{S^0} \simeq X$. Since $X^{(-)}$ sends pushouts in $\mathbf{An}_*^{\text{fin}}$ to pullbacks in D , we have

$$X^{S^{n+1}} \simeq X^{\Sigma S^n} \simeq \Omega X^{S^n}$$

for every $n \geq 0$. Thus the comparison automorphism of $\Omega^3 X$ is induced by the analogous automorphism of S^3 in $\mathbf{An}_*^{\text{fin}}$.

It remains to prove the claim in An_*^{fin} for the generator $X = S^0$. We must show that the composite

$$\tau: S^3 \simeq \Sigma(\Sigma^2 S^0) \xrightarrow{\cong} \Sigma^2(\Sigma S^0) \simeq S^3$$

is pointed homotopic to the identity. By Remark 2.4.19, it suffices to show that τ has degree 1. Using the swap map $\sigma_{X,Y}$ from Definition 2.4.10, we may identify τ with σ_{S^1, S^2} . The isomorphism $S^2 \cong S^1 \wedge S^1$ from Exercise 2.11 then gives a decomposition

$$S^3 \simeq S^1 \wedge S^1 \wedge S^1 \xrightarrow{\text{id} \wedge \sigma_{S^1, S^1}} S^1 \wedge S^1 \wedge S^1 \xrightarrow{\sigma_{S^1, S^1} \wedge \text{id}} S^1 \wedge S^1 \wedge S^1 \simeq S^3.$$

Both maps in this decomposition are suspensions of σ_{S^1, S^1} , up to the natural identifications of smash products with suspensions. Their degrees are therefore equal to that of σ_{S^1, S^1} by Remark 2.4.19. Since this swap map is an isomorphism, its degree is ± 1 . We conclude that

$$\deg(\tau) = \deg(\text{id} \wedge \sigma_{S^1, S^1}) \cdot \deg(\sigma_{S^1, S^1} \wedge \text{id}) = \deg(\sigma_{S^1, S^1})^2 = 1,$$

as desired. The construction is natural in X , since the comparison is induced by the automorphism τ of S^3 under the equivalence of Lemma 4.3.12. $\square$

Theorem 4.3.14. *For every ∞ -category C with finite limits, the ∞ -category $\text{Sp}(C)$ is stable.*

Proof. By Lemma 4.3.8, finite limits in $\text{PSp}(C)$ are computed pointwise. Since Ω preserves finite limits, the full subcategory $\text{Sp}(C) \subseteq \text{PSp}(C)$ is closed under finite limits. It is also pointed: the constant zero prespectrum is both initial and terminal, since the zero object of C_* is both initial and terminal.

It remains, by Theorem 4.2.2, to show that the loop functor on $\text{Sp}(C)$ is an equivalence. Define the shift functor

$$\text{shift}: \text{Sp}(C) \rightarrow \text{Sp}(C), \quad \text{shift}(X)_n := X_{n+1},$$

with structure maps inherited from X . This functor is an equivalence: an inverse sends X to the spectrum $\text{shift}^{-1} X$ with

$$(\text{shift}^{-1} X)_0 := \Omega X_0 \quad \text{and} \quad (\text{shift}^{-1} X)_n := X_{n-1} \quad (n \geq 1),$$

whose structure map in degree 0 is the identity of ΩX_0 , and whose structure map in degree $n \geq 1$ is σ_{n-1}^X .

We claim that there is a natural isomorphism

$$\alpha: X \xrightarrow{\cong} \Omega(\text{shift } X).$$

By Lemma 4.3.9, it suffices to construct this map after passing to the even-level model. There we set

$$\alpha_{2k} := \sigma_{2k}^X: X_{2k} \xrightarrow{\cong} \Omega X_{2k+1} = \Omega(\text{shift } X)_{2k}.$$

We must check compatibility with the two-step structure maps. For X , the relevant structure map is

$$\tilde{\sigma}_k^X = \left(X_{2k} \xrightarrow{\sigma_{2k}^X} \Omega X_{2k+1} \xrightarrow{\Omega(\sigma_{2k+1}^X)} \Omega^2 X_{2k+2} \right).$$

For $\Omega(\text{shift } X)$, finite limits are computed pointwise, but its two-step structure map is not obtained by merely writing down the same composite with an extra Ω in front. After precomposing with $\alpha_{2k} = \sigma_{2k}^X$, the first composite in the compatibility square is

$$X_{2k} \xrightarrow{\sigma_{2k}^X} \Omega X_{2k+1} \xrightarrow{\Omega(\sigma_{2k+1}^X)} \Omega^2 X_{2k+2} \xrightarrow{\Omega(\Omega(\sigma_{2k+2}^X))} \Omega(\Omega^2 X_{2k+3}) \xrightarrow{\cong} \Omega^2(\Omega X_{2k+3}),$$

where the final arrow is the canonical comparison coming from the fact that Ω^2 preserves loop objects. The other composite in the compatibility square is

$$X_{2k} \xrightarrow{\sigma_{2k}^X} \Omega X_{2k+1} \xrightarrow{\Omega(\sigma_{2k+1}^X)} \Omega^2 X_{2k+2} \xrightarrow{\Omega^2(\sigma_{2k+2}^X)} \Omega^2(\Omega X_{2k+3}).$$

Thus the two composites whose equality is required differ only by the resulting automorphism of $\Omega^3 X_{2k+3}$. This automorphism is the identity by Proposition 4.3.13. Hence α is an isomorphism of spectra.

Consequently $\text{id}_{\text{Sp}(C)} \simeq \Omega \circ \text{shift}$. Since shift is an equivalence, Ω is an equivalence as well. $\square$

Corollary 4.3.15. *The suspension functor on $\text{Sp}(C)$ is naturally isomorphic to the shift functor. In particular, for every spectrum X and all $m, n \geq 0$, there is a natural isomorphism*

$$(X[n])_m \cong X_{m+n}.$$

Proof. The proof of Theorem 4.3.14 gives a natural isomorphism $\text{id}_{\text{Sp}(C)} \cong \Omega \circ \text{shift}$. Since Σ is inverse to Ω , this identifies Σ with shift . $\square$

Proposition 4.3.16. *For an ∞ -category C with finite limits, the functor $\Omega^\infty: \text{Sp}(C) \rightarrow C$ exhibits $\text{Sp}(C)$ as a stabilization of C .*

Proof. Let D be a stable ∞ -category. We need to show that postcomposition with Ω^∞ induces an equivalence

$$\text{Fun}^{\text{lex}}(D, \text{Sp}(C)) \xrightarrow{\sim} \text{Fun}^{\text{lex}}(D, C).$$

We construct an inverse. Let $F: D \rightarrow C$ be left exact. Since D is pointed, the functor F has a canonical lift to C_* : the object $F(d)$ is pointed by the map $* \simeq F(0) \rightarrow F(d)$ induced by $0 \rightarrow d$. For every $n \geq 0$, define a functor

$$\tilde{F}_n: D \rightarrow C_*, \quad \tilde{F}_n(d) := F(d[n]).$$

The unit equivalences $d[n] \rightarrow \Omega_D d[n+1]$, together with left exactness of F , provide natural isomorphisms

$$\tilde{F}_n \xrightarrow{\sim} \Omega \tilde{F}_{n+1}.$$

Explicitly, these are obtained by applying F to the unit equivalences and then using the natural isomorphisms $F(\Omega_D d[n+1]) \xrightarrow{\sim} \Omega F(d[n+1])$ provided by left exactness. Thus the functors $\tilde{F}_n$ and these structure isomorphisms assemble into a functor

$$\tilde{F}: D \rightarrow \text{Sp}(C).$$

This construction is natural in F , and hence defines a functor $\text{Fun}^{\text{lex}}(D, C) \rightarrow \text{Fun}^{\text{lex}}(D, \text{Sp}(C))$. Since limits in $\text{Sp}(C)$ are computed levelwise, and since each shift $d \mapsto d[n]$ is an equivalence of D , the functor $\tilde{F}$ is left exact.

The composite $\Omega^\infty \widetilde{F}$ is F , since evaluation at level zero gives $\widetilde{F}(d)_0 = F(d)$. Conversely, let $G: D \rightarrow \mathrm{Sp}(C)$ be left exact. Since D and $\mathrm{Sp}(C)$ are stable, G is exact by Corollary 4.2.27. Hence

$$G(d[n]) \simeq G(d)[n].$$

By Corollary 4.3.15, evaluation at level zero gives natural isomorphisms

$$(\Omega^\infty G)(d[n]) = G(d[n])_0 \simeq G(d)_n.$$

These isomorphisms are compatible with the structure maps, by naturality of the unit $d[n] \rightarrow \Omega_D d[n+1]$ and of the equivalence between suspension and shift in $\mathrm{Sp}(C)$. Thus $\widetilde{\Omega^\infty G} \simeq G$, naturally in G . $\square$

Corollary 4.3.17. *The fully faithful inclusion functor $\mathrm{Cat}_\infty^{\mathrm{st}} \hookrightarrow \mathrm{Cat}_\infty^{\mathrm{lex}}$ admits a right adjoint*

$$\mathrm{Sp}: \mathrm{Cat}_\infty^{\mathrm{lex}} \rightarrow \mathrm{Cat}_\infty^{\mathrm{st}}$$

given on objects by sending C to its stabilization $\mathrm{Sp}(C)$. The counit $\mathrm{Sp}(C) \rightarrow C$ of the adjunction is given by Ω^∞ .

Proof. By the pointwise criterion for adjunctions from Lemma 21.1.4, it suffices to show that for every stable ∞ -category D , composition with $\Omega^\infty: \mathrm{Sp}(C) \rightarrow C$ induces an equivalence

$$\Omega^\infty \circ -: \mathrm{Hom}_{\mathrm{Cat}_\infty^{\mathrm{st}}}(D, \mathrm{Sp}(C)) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Cat}_\infty^{\mathrm{lex}}}(D, C).$$

But since a functor $D \rightarrow \mathrm{Sp}(C)$ is exact if and only if it is left exact, this map is obtained from the defining equivalence $\mathrm{Fun}^{\mathrm{lex}}(D, \mathrm{Sp}(C)) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{lex}}(D, C)$ by passing to groupoid cores. $\square$

4.3.2 Spectrification and suspension spectra

We now explain how to compute colimits of spectra. The point is that colimits are easy in prespectra, and a prespectrum can be turned into a spectrum by repeatedly applying the shift-loop operation.

Lemma 4.3.18 (Limits of spectra). *Let C be an ∞ -category with finite limits. If C admits I -indexed limits, then $\mathrm{Sp}(C)$ admits I -indexed limits, and the evaluation functors $(-)_n: \mathrm{Sp}(C) \rightarrow C_*$ preserve them.*

Proof. By Lemma 4.3.8, the limit of a diagram of spectra can first be computed in $\mathrm{PSp}(C)$. Since Ω preserves limits, the structure maps of this pointwise limit are again equivalences. Thus the limit prespectrum is a spectrum. $\square$

Proposition 4.3.19 (Spectrification). *Let C be an ∞ -category with finite limits and sequential colimits. Assume that $\Omega: C_* \rightarrow C_*$ preserves sequential colimits. Then the inclusion $\mathrm{Sp}(C) \hookrightarrow \mathrm{PSp}(C)$ admits a left adjoint*

$$(-)^{\mathrm{sp}}: \mathrm{PSp}(C) \rightarrow \mathrm{Sp}(C), \quad X \mapsto X^{\mathrm{sp}},$$

called the spectrification functor.

Proof. The endofunctor $\Omega \text{ sh}$ of $C_*^{\mathbb{N}}$ induces an endofunctor, still denoted $\Omega \text{ sh}$, of $\text{PSp}(C)$ by

$$(\Omega \text{ sh } X)_n := \Omega X_{n+1},$$

with structure maps $\Omega(\sigma_{n+1}^X): \Omega X_{n+1} \rightarrow \Omega^2 X_{n+2}$. The structure maps of X define a natural map $\eta_X: X \rightarrow \Omega \text{ sh } X$. Since sh preserves colimits and Ω preserves sequential colimits, the endofunctor $\Omega \text{ sh}$ preserves sequential colimits of prespectra.

For a prespectrum X , define

$$X^{\text{sp}} := \text{colim} \left(X \xrightarrow{\eta_X} \Omega \text{ sh } X \xrightarrow{\Omega \text{ sh}(\eta_X)} (\Omega \text{ sh})^2 X \rightarrow \dots \right)$$

in $\text{PSp}(C)$. Then

$$\Omega \text{ sh}(X^{\text{sp}}) \simeq \text{colim}_{k \geq 0} (\Omega \text{ sh})^{k+1} X \simeq \text{colim}_{k \geq 0} (\Omega \text{ sh})^k X = X^{\text{sp}},$$

where the second isomorphism uses that the successor map $k \mapsto k + 1$ is cofinal in $\mathbb{N}$. Under this isomorphism, the structure map of X^{sp} is an isomorphism, so X^{sp} is a spectrum.

It remains to prove the universal property. Let Y be a spectrum. Since $\eta_Y: Y \rightarrow \Omega \text{ sh } Y$ is an isomorphism, composition with η_X induces an equivalence

$$\text{Hom}_{\text{PSp}(C)}(\Omega \text{ sh } X, Y) \xrightarrow{\sim} \text{Hom}_{\text{PSp}(C)}(X, Y);$$

an inverse sends a map $f: X \rightarrow Y$ to the composite $\Omega \text{ sh } X \xrightarrow{\Omega \text{ sh}(f)} \Omega \text{ sh } Y \xrightarrow{\eta_Y^{-1}} Y$. Therefore

$$\text{Hom}_{\text{Sp}(C)}(X^{\text{sp}}, Y) \simeq \lim_k \text{Hom}_{\text{PSp}(C)}((\Omega \text{ sh})^k X, Y) \simeq \text{Hom}_{\text{PSp}(C)}(X, Y),$$

which is the desired adjunction. $\square$

Lemma 4.3.20 (Colimits of spectra). *Let C be an ∞ -category with finite limits and sequential colimits, and assume that $\Omega: C_* \rightarrow C_*$ preserves sequential colimits. If C_* admits I -indexed colimits for some ∞ -category I , then $\text{Sp}(C)$ admits I -indexed colimits.*

Proof. Let $X: I \rightarrow \text{Sp}(C)$ be a diagram. Compute its colimit in $\text{PSp}(C)$, where colimits are pointwise by Lemma 4.3.8, and then apply spectrification. Since spectrification is left adjoint to the inclusion $\text{Sp}(C) \hookrightarrow \text{PSp}(C)$, the resulting spectrum has the universal property of the colimit in $\text{Sp}(C)$. $\square$

Lemma 4.3.21. *Filtered colimits in Sp are computed levelwise. In particular, the functor $\Omega^\infty: \text{Sp} \rightarrow \text{An}$ preserves filtered colimits.*

Proof. Filtered colimits in An_* are created in An , and the loop functor $\Omega: \text{An}_* \rightarrow \text{An}_*$ preserves them. Hence the levelwise colimit in $\text{PSp}(\text{An})$ of a filtered diagram of spectra again has isomorphisms as its structure maps, so it is already a spectrum. The final claim follows by evaluating at level zero. $\square$

Lemma 4.3.22 (Suspension spectra). *Let C be an ∞ -category admitting finite limits, finite colimits, and sequential colimits, and assume that $\Omega: C_* \rightarrow C_*$ preserves sequential colimits. Then the functor $\Omega^\infty: \text{Sp}(C) \rightarrow C_*$ admits a left adjoint*

$$\Sigma^\infty: C_* \rightarrow \text{Sp}(C).$$

Proof. First define the free prespectrum functor

$$\Sigma^{\infty, \text{pre}} : C_* \rightarrow \text{PSp}(C)$$

by

$$\Sigma^{\infty, \text{pre}}(X)_n := \Sigma^n X,$$

with structure maps $\Sigma^n X \rightarrow \Omega \Sigma^{n+1} X$ given by the unit of the adjunction $\Sigma \dashv \Omega$ in C_* . We claim that $\Sigma^{\infty, \text{pre}}$ is left adjoint to evaluation at level zero $\text{ev}_0 : \text{PSp}(C) \rightarrow C_*$. Indeed, a map $\Sigma^{\infty, \text{pre}}(X) \rightarrow Y$ is determined by its level-zero component $X \rightarrow Y_0$: once $f_n : \Sigma^n X \rightarrow Y_n$ is known, compatibility with the structure maps forces $f_{n+1} : \Sigma^{n+1} X \rightarrow Y_{n+1}$ to be adjoint to the composite

$$\Sigma^n X \xrightarrow{f_n} Y_n \xrightarrow{\sigma_n^Y} \Omega Y_{n+1}.$$

Conversely, this recursive construction produces a compatible map of prespectra. The construction is natural in X and Y and applies equally to parametrized families of maps, so it gives an isomorphism of hom animae

$$\text{Hom}_{\text{PSp}(C)}(\Sigma^{\infty, \text{pre}} X, Y) \simeq \text{Hom}_{C_*}(X, Y_0).$$

Now set

$$\Sigma^{\infty}(X) := (\Sigma^{\infty, \text{pre}} X)^{\text{sp}}.$$

For $Y \in \text{Sp}(C)$, spectrification gives natural equivalences

$$\text{Hom}_{\text{Sp}(C)}(\Sigma^{\infty} X, Y) \simeq \text{Hom}_{\text{PSp}(C)}(\Sigma^{\infty, \text{pre}} X, Y) \simeq \text{Hom}_{C_*}(X, \Omega^{\infty} Y),$$

which proves the adjunction. $\square$

Corollary 4.3.23. *In the situation of Lemma 4.3.22, the functor $\Omega^{\infty} : \text{Sp}(C) \rightarrow C$ admits a left adjoint*

$$\Sigma_+^{\infty} : C \rightarrow \text{Sp}(C)$$

given by $\Sigma_+^{\infty}(X) := \Sigma^{\infty}(X_+)$, where $X_+ := X \sqcup *$.

Proof. Since adjunctions compose, this is immediate from the previous lemma and the fact that a left adjoint to the forgetful functor $C_* \rightarrow C$ is given by $(-)_+ : C \rightarrow C_*$, $X \mapsto X_+ := X \sqcup *$. $\square$

4.3.3 Standard presentations

Spectrification also gives a canonical way to build a spectrum from its levels.

Definition 4.3.24. Let C be an ∞ -category with finite limits. For $n \geq 0$, define

$$\Omega^{\infty-n} : \text{Sp}(C) \rightarrow C_*, \quad \Omega^{\infty-n}(X) := \Omega^{\infty}(X[n]),$$

where $X[n]$ denotes the n -fold suspension of X in the stable ∞ -category $\text{Sp}(C)$. By Corollary 4.3.15, this functor evaluates at the n -th level in the prespectrum model: $\Omega^{\infty-n}(X) \simeq X_n$.

Lemma 4.3.25. *Let C be an ∞ -category with finite limits. For $n \geq 0$ there are natural isomorphisms*

$$\Omega^n \Omega^{\infty-n} \cong \Omega^{\infty} \quad \text{and} \quad \Omega \Omega^{\infty-(n+1)} \cong \Omega^{\infty-n}.$$

Proof. This follows from stability of $\mathrm{Sp}(C)$ and the fact that $\Omega^\infty: \mathrm{Sp}(C) \rightarrow C_*$ preserves finite limits. $\square$

Lemma 4.3.26. *In the situation of Lemma 4.3.22, the functor $\Omega^{\infty-n}: \mathrm{Sp}(C) \rightarrow C_*$ admits a left adjoint*

$$\Sigma^{\infty-n}: C_* \rightarrow \mathrm{Sp}(C)$$

given by $\Sigma^{\infty-n}(X) := (\Sigma^\infty(X))[-n]$. It satisfies the relations

$$\Sigma^{\infty-n}\Sigma^n \cong \Sigma^\infty \quad \text{and} \quad \Sigma^{\infty-(n+1)}\Sigma \simeq \Sigma^{\infty-n}.$$

Proof. Since adjunctions compose, the functor $(\Sigma^\infty(-))[-n]$ is left adjoint to $\Omega^\infty((-)[n]) = \Omega^{\infty-n}$. The stated relations follow by passing to left adjoints in the previous lemma. $\square$

Remark 4.3.27. In the situation of Lemma 4.3.22, the suspension spectrum $\Sigma^\infty(X)$ of a pointed object $X \in C_*$ is the spectrification of the suspension prespectrum

$$(X, \Sigma X, \Sigma^2 X, \dots).$$

Equivalently, its n -th level is given by

$$\Sigma^\infty(X)_n \simeq \mathrm{colim}_{k \geq 0} \Omega^k \Sigma^{k+n} X.$$

The structure maps are the inverses of the natural isomorphisms

$$\Omega \Sigma^\infty(X)_{n+1} \simeq \Omega(\mathrm{colim}_{k \geq 0} \Omega^k \Sigma^{k+n+1} X) \simeq \mathrm{colim}_{k \geq 0} \Omega^{k+1} \Sigma^{k+n+1} X \simeq \Sigma^\infty(X)_n,$$

where we use that $\Omega: C_* \rightarrow C_*$ preserves sequential colimits.

Exercise 4.3.28. In the situation of Lemma 4.3.22, use the prespectrum model to verify directly that for any spectrum $Y \in \mathrm{Sp}(C)$ and any pointed object $X \in C_*$ there is a natural isomorphism

$$\mathrm{Hom}_{\mathrm{Sp}(C)}(\Sigma^\infty X, Y) \xrightarrow{\sim} \mathrm{Hom}_{C_*}(X, \Omega^\infty Y).$$

Proposition 4.3.29 (Associated spectrum). *In the situation of Lemma 4.3.22, let X be a prespectrum in C . Its associated spectrum X^{sp} is naturally isomorphic to the following colimit in $\mathrm{Sp}(C)$:*

$$X^{\mathrm{sp}} \simeq \mathrm{colim}(\Sigma^\infty X_0 \rightarrow \Sigma^{\infty-1} X_1 \rightarrow \Sigma^{\infty-2} X_2 \rightarrow \Sigma^{\infty-3} X_3 \rightarrow \dots).$$

Here the n -th transition map is given by the composite

$$\Sigma^{\infty-n} X_n \xrightarrow{\cong} \Sigma^{\infty-(n+1)} \Sigma X_n \xrightarrow{\Sigma \sigma_n^X} \Sigma^{\infty-(n+1)} \Sigma \Omega X_{n+1} \xrightarrow{\varepsilon} \Sigma^{\infty-(n+1)} X_{n+1},$$

where the first isomorphism is from Lemma 4.3.26, and the last map is induced by the counit $\varepsilon: \Sigma \Omega \rightarrow \mathrm{id}$ of the adjunction $\Sigma \dashv \Omega$ on C_ .*

Proof. Let Y be a spectrum in $\mathrm{Sp}(C)$. Mapping out of the displayed colimit gives

$$\lim_n \mathrm{Hom}_{\mathrm{Sp}(C)}(\Sigma^{\infty-n} X_n, Y) \simeq \lim_n \mathrm{Hom}_{C_*}(X_n, \Omega^{\infty-n} Y).$$

The transition maps in this limit are precisely the compatibility conditions for a map of prespectra $X \rightarrow Y$. Hence the right-hand side identifies with $\mathrm{Hom}_{\mathrm{Sp}(C)}(X, Y)$. By the spectrification adjunction from Proposition 4.3.19, this is naturally equivalent to $\mathrm{Hom}_{\mathrm{Sp}(C)}(X^{\mathrm{sp}}, Y)$. The claim follows from Yoneda. $\square$

Corollary 4.3.30 (Standard presentation). *In the situation of Lemma 4.3.22, every spectrum $X \in \mathbf{Sp}(C)$ is naturally isomorphic to the spectrum associated to its underlying prespectrum:*

$$\operatorname{colim}_{n \geq 0} \Sigma^{\infty - n} X_n \xrightarrow{\cong} X,$$

where $X_n = \Omega^{\infty - n} X$.

Proof. Apply the previous proposition to the underlying prespectrum of X . Since X is already a spectrum, spectrification fixes it. $\square$

4.4 The ∞ -category of spectra

We now specialize the preceding construction to anima. After recording the basic properties of the resulting ∞ -category of spectra, we construct hom spectra and tensoring by spectra, and use these to describe colimit-preserving functors out of $\mathbf{Sp}$.

Definition 4.4.1. The ∞ -category of spectra $\mathbf{Sp}$ is defined as the stabilization of the ∞ -category of anima, $\mathbf{An}$,

$$\mathbf{Sp} := \mathbf{Sp}(\mathbf{An}).$$

Objects of $\mathbf{Sp}$ are called *spectra*. Concretely, a spectrum is a sequence of pointed anima X_n equipped with isomorphisms

$$\sigma_n : X_n \xrightarrow{\sim} \Omega(X_{n+1}).$$

By Lemma 4.3.7, a morphism in $\mathbf{Sp} = \mathbf{Sp}(\mathbf{An})$ is precisely a sequence of levelwise pointed maps together with homotopies compatible with the structure maps. Thus these are exactly the morphisms introduced in Definition 3.2.2. The following are the central properties governing the behavior of $\mathbf{Sp}$:

- (1) The ∞ -category $\mathbf{Sp}$ comes equipped with a left exact functor $\Omega^\infty : \mathbf{Sp} \rightarrow \mathbf{An}$. Given a spectrum X , we refer to $\Omega^\infty(X) = X_0$ as its *underlying anima*.
- (2) Given a stable ∞ -category D , any left exact functor $D \rightarrow \mathbf{An}$ uniquely lifts to an exact functor $D \rightarrow \mathbf{Sp}$: composition with Ω^∞ induces an equivalence:

$$\Omega^\infty \circ - : \operatorname{Fun}^{\operatorname{ex}}(D, \mathbf{Sp}) \xrightarrow{\sim} \operatorname{Fun}^{\operatorname{lex}}(D, \mathbf{An}).$$

- (3) The ∞ -category $\mathbf{Sp}$ admits small limits (Lemma 4.3.18) and colimits (Lemma 4.3.20).
- (4) The functors $\Omega^\infty : \mathbf{Sp} \rightarrow \mathbf{An}$ and $\Omega^\infty : \mathbf{Sp} \rightarrow \mathbf{An}_*$ admit left adjoints

$$\Sigma^\infty : \mathbf{An}_* \rightarrow \mathbf{Sp} \quad \text{and} \quad \Sigma_+^\infty : \mathbf{An} \rightarrow \mathbf{Sp}$$

satisfying $\Sigma_+^\infty(X) \cong \Sigma^\infty(X_+)$ (Lemma 4.3.22 and Corollary 4.3.23).

Given a pointed anima X , we refer to the spectrum $\Sigma^\infty(X)$ as the *reduced suspension spectrum* of X . For unpointed X , we call $\Sigma_+^\infty(X)$ the *unreduced suspension spectrum* of X .

Definition 4.4.2 (Sphere spectrum). We define the *sphere spectrum* as $\mathbb{S} := \Sigma^\infty(S^0) \simeq \Sigma_+^\infty(*) \in \mathbf{Sp}$. For an anima X , we will frequently use the alternative notation

$$\mathbb{S}[X] := \Sigma_+^\infty(X)$$

for the unreduced suspension spectrum. We wish to think of the spectrum $\mathbb{S}[X]$ as the ‘free spectrum generated by X ’. This notation mimics the notation for the free abelian group $\mathbb{Z}[S]$ generated by a set S , characterized by the property that group homomorphisms $\mathbb{Z}[S] \rightarrow A$ correspond to maps of sets $S \rightarrow A$.

Table 4.1 collects the resulting analogies between ordinary algebra and ‘higher algebra’.

Ordinary algebra	Higher algebra
Set	Anima
Abelian group	Spectrum
Integers $\mathbb{Z}$	Sphere spectrum $\mathbb{S}$
Underlying set	Underlying anima $\Omega^\infty(X)$
Free abelian group $\mathbb{Z}[S]$	Unreduced suspension spectrum $\mathbb{S}[X]$

Table 4.1: The dictionary between ordinary algebra and ‘higher algebra’.

Remark 4.4.3. Given a pointed anima $(X, x) \in \mathbf{An}_*$, note that we may write X as the cofiber of the basepoint inclusion $*_+ \rightarrow X_+$. Since $\Sigma^\infty(-)$ preserves cofibers, we may write the reduced suspension spectrum of X as

$$\Sigma^\infty(X, x) \simeq \mathrm{cofib}(\mathbb{S} \xrightarrow{x} \mathbb{S}[X]).$$

4.4.1 Hom spectra

Given an abelian category $\mathcal{A}$, the hom sets $\mathrm{Hom}_{\mathcal{A}}(A, B)$ can be equipped with a canonical abelian group structure, where the sum $f + g$ of two morphisms $f, g: A \rightarrow B$ is given by the composite

$$A \xrightarrow{\Delta} A \oplus A \xrightarrow{f \oplus g} B \oplus B \xrightarrow{\nabla} B.$$

In an analogous way, the hom animae $\mathrm{Hom}_C(X, Y)$ in a stable ∞ -category C can be canonically lifted to spectra:

Proposition 4.4.4. *For a stable ∞ -category C , there exists a unique functor*

$$\mathrm{hom}_C(-, -): C^{\mathrm{op}} \times C \rightarrow \mathbf{Sp}$$

that is exact in both variables and fits in a commutative triangle as follows:

$$\begin{array}{ccc} & & \mathbf{Sp} \\ & \nearrow \mathrm{hom}_C & \downarrow \Omega^\infty \\ C^{\mathrm{op}} \times C & \xrightarrow{\mathrm{Hom}_C} & \mathbf{An}. \end{array}$$

Proof. Consider the Yoneda embedding

$$C \hookrightarrow \text{Fun}(C^{\text{op}}, \text{An}), \quad Y \mapsto \text{Hom}_C(-, Y).$$

This lands in left exact functors, so by the equivalence $\text{Fun}^{\text{ex}}(C^{\text{op}}, \text{Sp}) \xrightarrow{\sim} \text{Fun}^{\text{lex}}(C^{\text{op}}, \text{An})$ we may uniquely lift it to a (fully faithful) functor

$$C \hookrightarrow \text{Fun}^{\text{ex}}(C^{\text{op}}, \text{Sp}), \quad Y \mapsto \text{hom}_C(-, Y).$$

The resulting functor $\text{hom}_C: C^{\text{op}} \times C \rightarrow \text{Sp}$ is exact in the first variable by construction. It remains to show it is also exact in the second variable: for fixed X , we need to show that the functor $\text{hom}_C(X, -): C \rightarrow \text{Sp}$ is exact. It suffices to check this after composing with the functors $(-)_n: \text{Sp} \rightarrow \text{An}$, which are left exact and jointly conservative. Since we have $\text{hom}_C(X, Y)_n \simeq \text{hom}_C(X[-n], Y)_0 \simeq \text{Hom}_C(X[-n], Y)$, this follows from the left exactness of $\text{Hom}_C(X[-n], -): \text{Sp} \rightarrow \text{An}$. $\square$

Terminology 4.4.5. We refer to the spectra $\text{hom}_C(X, Y)$ as the *hom spectra* or *mapping spectra* of C .

We explicitly record the following observation made in the proof of the proposition:

Corollary 4.4.6 (Stable Yoneda lemma). *For a stable ∞ -category C , the functor*

$$C \rightarrow \text{Fun}(C^{\text{op}}, \text{Sp}), \quad Y \mapsto \text{hom}_C(-, Y)$$

is fully faithful. $\square$

Exercise 4.4.7. Show that for $C = \text{Sp}$, the Hom spectrum $\text{hom}(X, Y) := \text{hom}_{\text{Sp}}(X, Y)$ may be explicitly described at level n by

$$\text{hom}(X, Y)_n \simeq \text{Hom}_{\text{Sp}}(X, Y[n]).$$

Lemma 4.4.8. *The functor $\text{hom}(\mathbb{S}, -): \text{Sp} \rightarrow \text{Sp}$ is equivalent to the identity.*

Proof. It suffices to show that these two functors agree after applying Ω^∞ , and indeed we have

$$\Omega^\infty \text{hom}(\mathbb{S}, -) \cong \text{Hom}_{\text{Sp}}(\mathbb{S}, -) \cong \text{Hom}_{\text{Sp}}(\mathbb{S}[*], -) \cong \text{Hom}_{\text{An}}(*, \Omega^\infty(-)) \cong \Omega^\infty(-).$$

Here we use that $\text{Hom}_{\text{An}}(*, -): \text{An} \rightarrow \text{An}$ is equivalent to the identity, since its left adjoint $- \times *: \text{An} \rightarrow \text{An}$ is equivalent to the identity. $\square$

We will use the standard presentation from Corollary 4.3.30 in the following form: if X is a spectrum with levels $X_n = \Omega^{\infty-n} X$, then

$$X \simeq \text{colim}_{n \geq 0} \Sigma^{\infty-n} X_n.$$

Together with Remark 4.4.3, this says that every spectrum can be built from unreduced suspension spectra using shifts, cofibers and colimits.

4.4.2 The universal property of spectra

The ∞ -category of spectra is defined via a universal property for left exact functors into it. We will now prove another universal property in terms of functors *out of it*: colimit-preserving functors $\mathrm{Sp} \rightarrow C$ into a cocomplete stable ∞ -category C are uniquely determined by their value on the sphere spectrum.

Proposition 4.4.9 (Tensoring by spectra). *Let C be a stable ∞ -category with small colimits. Then C is tensored over Sp : there exists a unique functor*

$$- \otimes -: \mathrm{Sp} \times C \rightarrow C$$

equipped with a natural equivalence of spectra

$$\mathrm{hom}_C(X \otimes Y, Z) \cong \mathrm{hom}_{\mathrm{Sp}}(X, \mathrm{hom}_C(Y, Z))$$

for $X \in \mathrm{Sp}$ and $Y, Z \in C$.

Proof. The uniqueness of $X \otimes Y$ is clear from the stable Yoneda lemma, Corollary 4.4.6. For existence, we need to show that for every $Y \in C$ the functor

$$\Phi: \mathrm{Sp}^{\mathrm{op}} \rightarrow \mathrm{Fun}(C, \mathrm{Sp}), \quad \Phi(X) := \mathrm{hom}_{\mathrm{Sp}}(X, \mathrm{hom}_C(Y, -))$$

lands in the full subcategory $C^{\mathrm{op}} \hookrightarrow \mathrm{Fun}(C, \mathrm{Sp})$ of corepresentable functors, i.e. functors of the form $\mathrm{hom}_C(W, -)$ for some $W \in C$.

Step 1: We first show that $\Phi(X)$ is corepresentable whenever $X = \mathbb{S}[X']$ is an unreduced suspension spectrum for some anima $X' \in \mathrm{An}$. To this end, recall from Theorem 1.8.9 that evaluation at $*$ induces an equivalence

$$\mathrm{Fun}^{\mathrm{colim}}(\mathrm{An}, C) \xrightarrow{\sim} C.$$

We let $- \otimes Y: \mathrm{An} \rightarrow C$ denote the unique colimit-preserving functor satisfying $* \otimes Y \cong Y$. We then claim that there exists a natural isomorphism

$$\Phi(\mathbb{S}[X']) \cong \mathrm{hom}_C(X' \otimes Y, -)$$

of functors $C \rightarrow \mathrm{Sp}$ for every $X' \in \mathrm{An}$. For this, note that both sides define colimit-preserving functors $\mathrm{An} \rightarrow \mathrm{Fun}(C, \mathrm{Sp})^{\mathrm{op}}$ by varying X' . Indeed, on either side a colimit in X' becomes a limit in $\mathrm{Fun}(C, \mathrm{Sp})$, because the relevant mapping-spectrum functor is contravariant in the variable determined by X' ; this is precisely a colimit after passing to $\mathrm{Fun}(C, \mathrm{Sp})^{\mathrm{op}}$. It therefore suffices to produce an isomorphism for $X' = *$. And indeed, in that case both sides evaluate to $\mathrm{hom}_C(Y, -)$ by Lemma 4.4.8 and the given isomorphism $* \otimes Y \cong Y$.

Step 2: We now show the claim for an arbitrary spectrum X . Observe that the functor Φ is exact and sends colimits of spectra to limits in the functor category. Also observe that the corepresentable functors in $\mathrm{Fun}(C, \mathrm{Sp})$ are closed under limits and shifts. It thus remains to show that every spectrum X can be built from unreduced suspension spectra using shifts, cofibers and colimits. By Corollary 4.3.30, every spectrum is a colimit of shifts of reduced suspension spectra. Moreover, Remark 4.4.3 writes every reduced suspension spectrum $\Sigma^\infty(Z, z)$ as the cofiber of a map $\mathbb{S} \rightarrow \mathbb{S}[Z]$ between unreduced suspension spectra. $\square$

Corollary 4.4.10. *Let C be a stable ∞ -category with small colimits. For every object $Y \in C$, the functor $\mathrm{hom}_C(Y, -): C \rightarrow \mathrm{Sp}$ admits a left adjoint*

$$- \otimes Y: \mathrm{Sp} \rightarrow C.$$

Proof. Applying Ω^∞ to the natural isomorphism from Proposition 4.4.9 provides the adjunction equivalence

$$\mathrm{Hom}_C(X \otimes Y, Z) \cong \mathrm{Hom}_{\mathrm{Sp}}(X, \mathrm{hom}_C(Y, Z))$$

for $X \in \mathrm{Sp}$ and $Z \in C$. $\square$

Fixing some stable ∞ -category C with small colimits, let us now establish various basic properties of tensoring on C .

Lemma 4.4.11. *The functor $- \otimes -: \mathrm{Sp} \times C \rightarrow C$ preserves colimits in both variables.*

Proof. By the Yoneda lemma, it suffices to show that $\mathrm{Hom}_C(- \otimes -, Z): \mathrm{Sp}^{\mathrm{op}} \times C^{\mathrm{op}} \rightarrow \mathrm{An}$ preserves limits in both variables, for all $Z \in C$. This functor is naturally isomorphic to $\mathrm{Hom}_{\mathrm{Sp}}(-, \mathrm{hom}_C(-, Z))$. This clearly preserves limits in the first variable. For the second variable, this follows from the fact that $\Omega^\infty \mathrm{hom}_C(-, Z) = \mathrm{Hom}_C(-, Z): C^{\mathrm{op}} \rightarrow \mathrm{An}$ preserves limits. $\square$

Lemma 4.4.12. *The functor $\mathbb{S} \otimes -: C \rightarrow C$ is equivalent to the identity.*

Proof. For objects $Y, Z \in C$, there are natural isomorphisms

$$\mathrm{hom}_C(Y, Z) \xrightarrow{\text{Lemma 4.4.8}} \mathrm{hom}_{\mathrm{Sp}}(\mathbb{S}, \mathrm{hom}_C(Y, Z)) \cong \mathrm{hom}_C(\mathbb{S} \otimes Y, Z),$$

hence by the stable Yoneda lemma we obtain a natural isomorphism $\mathbb{S} \otimes Y \cong Y$. $\square$

Lemma 4.4.13. *For a colimit-preserving functor $F: \mathrm{Sp} \rightarrow C$, there exists a natural isomorphism*

$$F(-) \cong - \otimes F(\mathbb{S}).$$

Moreover, this natural isomorphism can be chosen naturally in $F \in \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, C)$.

Proof. We first produce a natural isomorphism $F(\mathbb{S}[-]) \cong \mathbb{S}[-] \otimes F(\mathbb{S})$ of functors $\mathrm{An} \rightarrow C$. In light of the equivalence $\mathrm{ev}_*: \mathrm{Fun}^{\mathrm{colim}}(\mathrm{An}, C) \xrightarrow{\sim} C$, it suffices to produce an isomorphism $F(\mathbb{S}) \cong \mathbb{S} \otimes F(\mathbb{S})$, for which we take the one from Lemma 4.4.12.

Now, for computing F on an arbitrary spectrum X , we use the standard presentation from Corollary 4.3.30. Using that F preserves colimits and shifts, we then get

$$F(X) \simeq F(\mathrm{colim}_n \Sigma^{\infty-n} X_n) \simeq \mathrm{colim}_n \Omega^n F(\Sigma^\infty X_n) \simeq \mathrm{colim}_n \Omega^n \mathrm{cofib}(F(\mathbb{S}) \rightarrow F(\mathbb{S}[X_n])),$$

where the last equivalence uses $\Sigma^\infty X_n \simeq \mathrm{cofib}(\mathbb{S} \rightarrow \mathbb{S}[X_n])$. Applying the same reasoning to $F' = - \otimes F(\mathbb{S})$, we similarly obtain

$$X \otimes F(\mathbb{S}) \simeq \mathrm{colim}_n \Omega^n \mathrm{cofib}(F(\mathbb{S}) \rightarrow \mathbb{S}[X_n] \otimes F(\mathbb{S})).$$

The claim now follows from the identification $F(\mathbb{S}[X_n]) \cong \mathbb{S}[X_n] \otimes F(\mathbb{S})$ established before. These equivalences are natural in both F and X : the standard presentation is natural in X , while the initial identification of colimit-preserving functors $\mathrm{An} \rightarrow C$ is natural in F . We therefore obtain the claimed natural isomorphism in $\mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, C)$. $\square$

Theorem 4.4.14 (Universality of Sp). *Let C be a stable ∞ -category with small colimits. Then evaluation at the sphere spectrum induces an equivalence*

$$\mathrm{ev}_{\mathbb{S}}: \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, C) \xrightarrow{\sim} C.$$

Proof. We claim that an inverse is given by the tensor product functor

$$C \rightarrow \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, C), \quad X \mapsto - \otimes X.$$

A natural isomorphism $\mathbb{S} \otimes X \cong X$ is provided by Lemma 4.4.12. A natural isomorphism $F \cong - \otimes F(\mathbb{S})$ is provided by Lemma 4.4.13. $\square$

We may summarize this result by saying that Sp is the *free cocomplete stable ∞ -category generated by $\mathbb{S}$* .

Corollary 4.4.15. *Let C be a stable ∞ -category with small colimits. Then every colimit-preserving functor $F: \mathrm{Sp} \rightarrow C$ admits a right adjoint $F^R: C \rightarrow \mathrm{Sp}$.*

Proof. By Theorem 4.4.14, F is of the form $- \otimes X$ for some $X \in C$, hence has right adjoint given by the mapping spectrum functor $\mathrm{hom}_C(X, -): C \rightarrow \mathrm{Sp}$. $\square$

4.4.3 Tensor products of spectra

Given two abelian groups A and B , we may form their *tensor product* $A \otimes B$, which is characterized by the property that group homomorphisms $A \otimes B \rightarrow C$ correspond to bilinear maps $A \times B \rightarrow C$, or equivalently to morphisms of abelian groups $A \rightarrow \mathrm{Hom}_{\mathrm{Ab}}(B, C)$. We will now construct a tensor product of spectra with analogous properties.

Proposition 4.4.16. *There exists a unique tensor product functor*

$$- \otimes -: \mathrm{Sp} \times \mathrm{Sp} \rightarrow \mathrm{Sp}$$

which preserves colimits in both variables and satisfies $\mathbb{S} \otimes \mathbb{S} \cong \mathbb{S}$.

Proof. Currying and two applications of Theorem 4.4.14 give equivalences

$$\mathrm{Fun}^{\mathrm{colim}, \mathrm{colim}}(\mathrm{Sp} \times \mathrm{Sp}, \mathrm{Sp}) \simeq \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, \mathrm{Sp})) \simeq \mathrm{Fun}^{\mathrm{colim}}(\mathrm{Sp}, \mathrm{Sp}) \simeq \mathrm{Sp},$$

where the intermediate functor category is stable and admits small colimits computed pointwise. Under this equivalence, a functor is sent to its value on $(\mathbb{S}, \mathbb{S})$. The sphere spectrum therefore determines the required functor uniquely. $\square$

Let us now establish various basic properties of the tensor product.

Lemma 4.4.17. *For spectra $X, Y, Z \in \mathrm{Sp}$, there are natural isomorphisms of spectra*

$$\mathbb{S} \otimes X \cong X \cong X \otimes \mathbb{S}, \quad X \otimes Y \cong Y \otimes X \quad (X \otimes Y) \otimes Z \cong X \otimes (Y \otimes Z)$$

and

$$\mathrm{hom}(Y, \mathrm{hom}(X, Z)) \cong \mathrm{hom}(X \otimes Y, Z) \cong \mathrm{hom}(X, \mathrm{hom}(Y, Z)).$$

Proof. For the first row, each of these expressions define functors that are colimit-preserving in each of their arguments. By Theorem 4.4.14, it thus suffices to produce such isomorphisms after evaluation at the sphere spectrum, where they are immediate consequences of the isomorphism $\mathbb{S} \otimes \mathbb{S} \cong \mathbb{S}$.

For the second row, we may fix Z and regard these expressions as functors $\mathrm{Sp}^{\mathrm{op}} \times \mathrm{Sp}^{\mathrm{op}} \rightarrow \mathrm{Sp}$ that preserve limits in both variables, or equivalently as functors $\mathrm{Sp} \times \mathrm{Sp} \rightarrow \mathrm{Sp}^{\mathrm{op}}$ preserving colimits in both variables. It then again suffices to prove they are isomorphic after setting $X = Y = \mathbb{S}$, which follows from the natural isomorphism $\mathrm{hom}(\mathbb{S}, Z) \cong Z$ from Lemma 4.4.8. $\square$

Lemma 4.4.18. *There exists a natural isomorphism*

$$\mathbb{S}[X] \otimes \mathbb{S}[Y] \cong \mathbb{S}[X \times Y]$$

for all $X, Y \in \mathrm{An}$.

Proof. Both of the functors $\mathbb{S}[-] \otimes \mathbb{S}[-], \mathbb{S}[- \times -] : \mathrm{An} \times \mathrm{An} \rightarrow \mathrm{Sp}$ preserve colimits in both variables, hence they are naturally isomorphic if and only if they agree on $(*, *)$, see Corollary 1.8.10. But on $(*, *)$ both sides evaluate to $\mathbb{S} \otimes \mathbb{S} \simeq \mathbb{S}$. $\square$

Lemma 4.4.19. *There exists a natural isomorphism*

$$\Sigma^\infty(X) \otimes \Sigma^\infty(Y) \cong \Sigma^\infty(X \wedge Y)$$

for all $X, Y \in \mathrm{An}_*$.

Proof. Since $- \otimes -$ preserves cofibers in each variable, this follows by combining the previous lemma with the expression $\Sigma^\infty(Z) \simeq \mathrm{cofib}(\mathbb{S} \rightarrow \mathbb{S}[Z])$ from Remark 4.4.3 and the isomorphism $X_+ \wedge Y_+ \cong (X \times Y)_+$ from Exercise 2.11. $\square$

Remark 4.4.20. Let X and Y be spectra. The standard presentations of X and Y provide the following description of their tensor product:

$$X \otimes Y \cong \mathrm{colim}_n \mathrm{colim}_m (\Sigma^{\infty-n} X_n \otimes \Sigma^{\infty-m} Y_m) \simeq \mathrm{colim}_n \mathrm{colim}_m \Sigma^{\infty-n-m} (X_n \wedge Y_m),$$

where the last equivalence holds by the previous lemma. However, this formula should be used with care: it makes implicit use of the additivity of the shift functors $X[-n][-m] \cong X[-(n+m)]$ from Warning 4.2.11. To avoid introducing sign errors, we will not work with this formula for the tensor product.

4.4.4 Homotopy groups

Important invariants of spectra are their *homotopy groups*:

Definition 4.4.21. For a spectrum X and an integer k , we define its k -th homotopy group $\pi_k(X) \in \mathrm{Ab}$ as

$$\pi_k(X) := [\mathbb{S}[k], X] := \pi_0 \mathrm{Hom}_{\mathrm{Sp}}(\mathbb{S}[k], X).$$

The assignment $X \mapsto \pi_k(X)$ defines a functor $\pi_* : \mathrm{Sp} \rightarrow \mathrm{Ab}$.

Exercise 4.4.22. Show that we have $\pi_k(X[n]) \simeq \pi_{k-n}(X)$ for all $k, n \in \mathbb{Z}$.

Exercise 4.4.23. Show that for $n \geq 0$ and $k + n \geq 0$ we have $\pi_k(X) \cong \pi_{n+k}(X_n)$. In particular we have $\pi_k(X) = \pi_k(\Omega^\infty X)$ for $k \geq 0$.

Example 4.4.24. For a pointed anima X , the negative homotopy groups of its suspension spectrum $\Sigma^\infty X$ vanish. For $k \geq 0$, Remark 4.3.27 gives

$$\pi_k(\Sigma^\infty(X)) \cong \pi_k(\Omega^\infty \Sigma^\infty(X)) \cong \operatorname{colim}_{n \geq 0} \pi_k(\Omega^n \Sigma^n X) = \operatorname{colim}_n \pi_{n+k}(\Sigma^n X).$$

These groups are known as the *stable homotopy groups* of X . By the classical Freudenthal suspension theorem [Fre37] the maps $\pi_{n+k}(\Sigma^n X) \rightarrow \pi_{n+k+1}(\Sigma^{n+1} X)$ are isomorphisms for $n \geq k + 2$.

Example 4.4.25. By Remark 2.4.19, there is an isomorphism $\deg: \pi_n(S^n) \xrightarrow{\cong} \mathbb{Z}$ for every $n \geq 1$, and the suspension maps correspond to the identity of $\mathbb{Z}$. It follows that there is an isomorphism $\pi_0(\mathbb{S}) \cong \operatorname{colim}_n \pi_n(S^n) \cong \mathbb{Z}$.

Corollary 4.4.26. *The functors $\pi_k: \mathbf{Sp} \rightarrow \mathbf{Ab}$ for $k \in \mathbb{Z}$ are jointly conservative: a morphism of spectra $f: X \rightarrow Y$ is an isomorphism if and only if the induced map $\pi_k(f): \pi_k(X) \rightarrow \pi_k(Y)$ is an isomorphism.*

Proof. By Lemma 4.3.6, f is an isomorphism if and only if each map $f_n: X_n \rightarrow Y_n$ of anima is an isomorphism. By the Whitehead Theorem for anima, Proposition 2.4.21, this is the case if f_n induces isomorphisms on all homotopy groups. By Exercise 4.4.23 this condition is equivalent to the condition that f induces isomorphisms on all homotopy groups. $\square$

Lemma 4.4.27. *For every $k \in \mathbb{Z}$, the functor $\pi_k: \mathbf{Sp} \rightarrow \mathbf{Ab}$ preserves filtered colimits.*

Proof. By Lemma 4.3.21, a filtered colimit of spectra is computed levelwise. The claim then follows from Exercise 4.4.23 together with the fact that the homotopy groups of pointed anima preserve filtered colimits. $\square$

Lemma 4.4.28. *For every $k \in \mathbb{Z}$, the functor $\pi_k: \mathbf{Sp} \rightarrow \mathbf{Ab}$ preserves arbitrary coproducts.*

Proof. Finite coproducts in $\mathbf{Sp}$ agree with finite products, and π_k preserves both of these. Since an arbitrary coproduct is the filtered colimit of its finite subcoproducts, the claim follows from Lemma 4.4.27. $\square$

Proposition 4.4.29 (Long exact sequence of homotopy groups). *Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be an exact sequence of spectra. Then there is a long exact sequence of homotopy groups of the form*

$$\dots \rightarrow \pi_1(Y) \xrightarrow{g_*} \pi_1(Z) \rightarrow \pi_0(X) \xrightarrow{f_*} \pi_0(Y) \xrightarrow{g_*} \pi_0(Z) \rightarrow \pi_{-1}(X) \xrightarrow{f_*} \pi_{-1}(Y) \rightarrow \dots$$

Proof. Since every exact sequence of spectra $X \xrightarrow{f} Y \xrightarrow{g} Z$ leads to two new exact sequences

$$Z[-1] \rightarrow X \xrightarrow{f} Y \quad \text{and} \quad Y \xrightarrow{g} Z \rightarrow X[1],$$

it inductively suffices to prove that the sequence $\pi_0(X) \xrightarrow{f_*} \pi_0(Y) \xrightarrow{g_*} \pi_0(Z)$ is exact. It is clear that the composite is the zero map. Conversely, consider a class $[\sigma] \in \pi_0(Y)$, represented by a morphism of spectra $\sigma: \mathbb{S} \rightarrow Y$, and assume that the class $[g \circ \sigma] \in \pi_0(Z)$ is zero. Then $\sigma: \mathbb{S} \rightarrow Y$ factors through the fiber X of $g: Y \rightarrow Z$, and hence is of the form $f \circ \tau$ for some $\tau: \mathbb{S} \rightarrow X$. It follows that $[\sigma]$ is in the image of $f_*: \pi_0(X) \rightarrow \pi_0(Y)$, as desired. $\square$

4.4.5 Homology theories represented by spectra

In Chapter 3, a spectrum E directly gave a cohomology theory by mapping into its individual pointed anima E_k . We now have the additional structure needed to associate a homology theory to E as well.

Definition 4.4.30 (E -homology). Let E be a spectrum and let $X \in \mathbf{An}_*$ be a pointed anima. For $k \in \mathbb{Z}$, we define the *reduced E -homology* of X by

$$\tilde{E}_k(X) := \pi_k(\Sigma^\infty X \otimes E).$$

Proposition 4.4.31. *For every spectrum E , the functors $\tilde{E}_*$ admit natural suspension isomorphisms that make them into a reduced homology theory in the sense of Definition 3.1.2.*

Proof. The natural isomorphism $\Sigma^\infty(\Sigma X) \cong (\Sigma^\infty X)[1]$ gives suspension isomorphisms

$$\tilde{E}_{k+1}(\Sigma X) = \pi_{k+1}((\Sigma^\infty X \otimes E)[1]) \cong \pi_k(\Sigma^\infty X \otimes E) = \tilde{E}_k(X).$$

Both Σ^∞ and $- \otimes E$ preserve cofiber sequences. Exactness therefore follows from the long exact sequence of homotopy groups in Proposition 4.4.29.

It remains to verify the wedge axiom. The functor $\Sigma^\infty(-) \otimes E$ preserves coproducts, and each $\pi_k : \mathbf{Sp} \rightarrow \mathbf{Ab}$ preserves coproducts by Lemma 4.4.28. Thus

$$\bigoplus_{i \in I} \tilde{E}_k(X_i) \xrightarrow{\cong} \tilde{E}_k\left(\bigvee_{i \in I} X_i\right),$$

as required. □

Remark 4.4.32 (Comparison with the earlier definition). The cohomology theory associated to E in Definition 3.2.3 can now be expressed in terms of the mapping spectrum. Indeed, the adjunction $\Sigma^\infty \dashv \Omega^\infty$ gives

$$\begin{aligned} \pi_{-k} \operatorname{hom}(\Sigma^\infty X, E) &\cong [\Sigma^\infty X[-k], E] \\ &\cong [\Sigma^\infty X, E[k]] \\ &\cong [X, \Omega^\infty(E[k])]_* \\ &\cong [X, E_k]_* = \tilde{E}^k(X). \end{aligned}$$

Here the last isomorphism uses the spectrum structure maps for $k \geq 0$ and the convention $E_k = \Omega^{-k} E_0$ for $k < 0$. All displayed isomorphisms are isomorphisms of abelian groups: under $\Omega^\infty(E[k]) \simeq E_k$, the canonical infinite-loop group structure agrees with the double-loop group structure used in Definition 3.2.3.

4.4.6 Mapping telescopes and the Milnor sequence

We record a general formula for sequential colimits in stable ∞ -categories, an abstract form of the classical *mapping telescope* which will let us later resolve a spectrum by direct sums of spheres. By duality this provides a description of an inverse limit, giving us the Milnor sequence.

Lemma 4.4.33 (Mapping telescope). *Let C be a stable ∞ -category with countable coproducts, and let $Y_0 \xrightarrow{f_0} Y_1 \xrightarrow{f_1} Y_2 \rightarrow \dots$ be a sequential diagram in C . Write $\iota_k: Y_k \rightarrow \bigoplus_{j \geq 0} Y_j$ for the inclusion of the k -th summand. Then there is an exact sequence*

$$\bigoplus_{k \geq 0} Y_k \xrightarrow{\text{id} - f} \bigoplus_{k \geq 0} Y_k \longrightarrow \text{colim}_k Y_k,$$

where $\text{id} - f$ restricts on the k -th summand to the difference $\iota_k - \iota_{k+1} \circ f_k$.

Proof. By the pushout description of $[\omega]$ in Axiom D, the same argument as in Exercise 2.4.28 expresses $\text{colim}_k Y_k$ as a pushout of the two maps from $\bigoplus_k Y_k$ to the coproducts of the even and odd terms. In a stable ∞ -category, the pushout of a span $B \xleftarrow{a} A \xrightarrow{b} C$ is the cofiber of

$$(a, -b): A \longrightarrow B \oplus C.$$

After identifying the coproduct of the even and odd terms with $\bigoplus_k Y_k$ and changing the signs on the odd source summands, this map is precisely $\text{id} - f$. $\square$

Corollary 4.4.34 (Dual mapping telescope). *Let C be a stable ∞ -category with countable products. For every tower*

$$\dots \xrightarrow{f_2} Y_2 \xrightarrow{f_1} Y_1 \xrightarrow{f_0} Y_0$$

in C , there is an exact sequence

$$\lim_k Y_k \longrightarrow \prod_{k \geq 0} Y_k \xrightarrow{\text{id} - f} \prod_{k \geq 0} Y_k,$$

where the k -th component of $(\text{id} - f)(y)$ is $y_k - f_k(y_{k+1})$.

Proof. Apply Lemma 4.4.33 in the opposite stable ∞ -category C^{op} . $\square$

To state the resulting short exact sequence, we first introduce the one derived-limit construction that we need.

Definition 4.4.35 (First derived limit). Let $\dots \xrightarrow{f_2} A_2 \xrightarrow{f_1} A_1 \xrightarrow{f_0} A_0$ be a tower of abelian groups. Define a homomorphism

$$d: \prod_{k \geq 0} A_k \longrightarrow \prod_{k \geq 0} A_k, \quad d(a)_k := a_k - f_k(a_{k+1}).$$

The *first derived limit* of the tower is the abelian group

$$\lim_k^1 A_k := \text{coker}(d).$$

Notice that the ordinary inverse limit $\lim_k A_k$ is the kernel of d .

Corollary 4.4.36 (Milnor sequence). *For every tower $\dots \rightarrow Y_2 \rightarrow Y_1 \rightarrow Y_0$ of spectra and every $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \longrightarrow \lim_k^1 \pi_{n+1}(Y_k) \longrightarrow \pi_n(\lim_k Y_k) \longrightarrow \lim_k \pi_n(Y_k) \longrightarrow 0.$$

Proof. Homotopy groups commute with products of spectra: indeed, the mapping anima functor $\mathrm{Hom}_{\mathrm{Sp}}(\mathbb{S}[n], -)$ preserves limits, and π_0 preserves products of anima. Consequently, applying the long exact sequence of Proposition 4.4.29 to the fiber sequence of Corollary 4.4.34 gives an exact sequence

$$\prod_k \pi_{n+1}(Y_k) \xrightarrow{d} \prod_k \pi_{n+1}(Y_k) \longrightarrow \pi_n(\lim_k Y_k) \longrightarrow \prod_k \pi_n(Y_k) \xrightarrow{d} \prod_k \pi_n(Y_k).$$

The claim follows by identifying the cokernel of the first map and the kernel of the last map using Definition 4.4.35. $\square$

Exercises

Exercise 4.1. Let C and D be ∞ -categories with finite limits. Show that there is an equivalence

$$\mathrm{Sp}(C \times D) \xrightarrow{\sim} \mathrm{Sp}(C) \times \mathrm{Sp}(D).$$

Exercise 4.2. Let C be a semiadditive ∞ -category.

- (1) Show that for all objects X and Y in C the set $[X, Y] := \pi_0 \mathrm{Hom}_C(X, Y)$ of homotopy classes of morphisms from X to Y admits a canonical structure of an abelian monoid, where addition of $f, g \in [X, Y]$ is given by the composite

$$f + g: X \xrightarrow{\Delta} X \oplus X \xrightarrow{f \oplus g} Y \oplus Y \xrightarrow{\nabla} Y.$$

- (2) Show that for all morphisms $f: X' \rightarrow X$ and $g: Y \rightarrow Y'$ in C the composition maps

$$- \circ f: [X, Y] \rightarrow [X', Y] \quad \text{and} \quad g \circ -: [X, Y] \rightarrow [X, Y']$$

are homomorphisms of abelian monoids.

- (3) Let M be an abelian monoid. Show that M is an abelian group if and only if the map

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}: M \oplus M \rightarrow M \oplus M, \quad (x, y) \mapsto (x, x + y)$$

is a bijection of sets.

- (4) Conclude that the semiadditive ∞ -category C is in fact *additive* if and only if the abelian monoid $[X, Y]$ is an abelian group for all X and Y .

Exercise 4.3. Let C be a pointed ∞ -category with finite limits.

- (1) Show that for every object X , the loop space ΩX admits the structure of a group object in the homotopy category $\mathrm{Ho}(C)$: there are maps

$$e: * \rightarrow \Omega X, \quad m: \Omega X \times \Omega X \rightarrow \Omega X, \quad \text{and} \quad i: \Omega X \rightarrow \Omega X$$

that satisfy the usual group relations up to homotopy.

- (2) Show that for objects X and Y in C , the set $[X, \Omega Y]$ is canonically a group. Show that $[X, \Omega^2 Y]$ is an abelian group.
- (3) Deduce that when C is stable, the Hom-set $[X, Y]$ in $\text{Ho}(C)$ is an abelian group for all X and Y , and that composition in C defines group homomorphisms.
- (4) Show that the resulting abelian group structure on $[X, Y]$ agrees with the one defined in Exercise 4.2.

Remark 4.4.37. Applying the previous exercise to C^{op} , we deduce that for every X the suspension ΣX is a *cogroup object* in $\text{Ho}(C)$, and hence that $[\Sigma X, Y]$ admits an abelian group structure for all Y .

Remark 4.4.38. In Chapter 5, we will upgrade the group structure of ΩX in $\text{Ho}(C)$ to a group structure in C .

Exercise 4.4. Let X be a pointed anima. Consider the prespectrum $\Sigma^{\infty, \text{pre}}(X) := (\Sigma^n X, \sigma_n)$, where the structure maps are given by the unit maps

$$\Sigma^n X \rightarrow \Omega \Sigma(\Sigma^n X) = \Omega \Sigma^{n+1} X$$

of the adjunction $\Sigma \dashv \Omega$. Show that the spectrum associated to the prespectrum $\Sigma^{\infty, \text{pre}}(X)$ is $\Sigma^{\infty}(X)$.

Hint: show that all the maps

$$\Sigma^{\infty-n}(\Sigma^n X) \rightarrow \Sigma^{\infty-(n+1)}(\Sigma^{n+1} X)$$

appearing in the definition of the associated spectrum are isomorphisms.

Exercise 4.5. Show that the functor $\Sigma^{\infty}: \text{An}_* \rightarrow \text{Sp}$ sends smash products of pointed animae to tensor products of spectra: produce a natural isomorphism

$$\Sigma^{\infty}(X) \otimes \Sigma^{\infty}(Y) \cong \Sigma^{\infty}(X \wedge Y)$$

for $X, Y \in \text{An}_*$.

Exercise 4.6. Let $\text{An}^{\text{fin}} \subseteq \text{An}$ consist of those animae which are finite as an ∞ -category. Show that An^{fin} is the smallest subcategory closed under finite colimits that contains the point.

Exercise 4.7. Let X be a finite cell complex (Definition 2.3.1). Show that $\Pi_{\infty}(X)$ is a finite anima.

Exercise 4.8. Show that for every spectrum X , the functor $\text{hom}(-, X): \text{Sp}^{\text{op}} \rightarrow \text{Sp}$ admits a left adjoint, which is given by $\text{hom}(-, X)^{\text{op}}: \text{Sp} \rightarrow \text{Sp}^{\text{op}}$.

5 The recognition principle

In the previous chapter, we saw that there is a strong analogy between spectra and abelian groups. The goal of this chapter is to provide more substance to this analogy: we define an ∞ -category $\mathbf{CGrp}(\mathbf{An})$ of commutative groups in $\mathbf{An}$, also known as *E_∞ -groups*, and construct a fully faithful functor

$$\mathbf{B}^\infty : \mathbf{CGrp}(\mathbf{An}) \hookrightarrow \mathbf{Sp}$$

whose image is the full subcategory $\mathbf{Sp}_{\geq 0}$ of connective spectra. This result is classically known as the *recognition principle for infinite loop spaces*.

The ∞ -categorical definition of commutative groups is more subtle than the one from classical algebra. Homotopical analogues of familiar algebraic structures come with infinite hierarchies of coherence conditions, making it impossible to define them by simply writing down a list of relations. Following ideas of Segal, we take a direct approach¹²: we define (commutative) monoids and groups as functors out of suitable indexing ∞ -categories, namely the opposite simplex category Δ^{op} and the span category of finite sets $\mathbf{Span}(\mathbf{Fin})$. This functor-based perspective elegantly encodes all required coherences at once.

The equivalence between commutative groups in $\mathbf{An}$ and connective spectra is a consequence of another result called the *recognition principle for loop spaces*. For every pointed anima X the loop space ΩX inherits the structure of a group object in $\mathbf{An}_*$. The recognition principle says that, conversely, every group object G arises this way: there exists a pointed anima X with $G \simeq \Omega X$, and if we require X to be connected then it is unique. This X is called the *delooping* of G and is denoted $\mathbf{B}G$.

If G is in fact a *commutative* group object, the delooping $\mathbf{B}G$ is again a commutative group, so the construction can be iterated: $\mathbf{B}^2 G := \mathbf{B}(\mathbf{B}G)$, and so on. The resulting sequence of deloopings assembles into a connective spectrum $\mathbf{B}^\infty G$. Conversely, the underlying anima of a spectrum comes with a preferred commutative group structure, which for $\mathbf{B}^\infty G$ recovers G .

This chapter is organized as follows. We introduce monoids and groups in Section 5.1, and treat the recognition principle for loop spaces in Section 5.2. We then introduce commutative monoids and groups in Section 5.3, and formulate and prove the recognition principle for infinite loop spaces in Section 5.4. Finally, Section 5.5 gives an explicit telescope model for group completions under a useful hypothesis.

¹²A more general approach to homotopy coherent algebraic structures is given in Part II via the theory of ∞ -operads.

5.1 Monoids and groups

Recall that a monoid is a set M equipped with a unit element $1 \in M$ and a product $x \cdot y \in M$ for all $x, y \in M$, satisfying associativity and unitality. Given such a monoid, we obtain for every $n \geq 0$ an n -fold multiplication map $M^n \rightarrow M$ sending $(x_1, \dots, x_n)$ to $x_1 \cdots x_n$. In the ∞ -categorical setting, we make all these n -fold multiplication maps part of the data, together with all the coherences between them. These coherences are encoded by the simplex category Δ .

Definition 5.1.1 (Monoid). Let C be an ∞ -category that admits finite products. We define a *monoid object in C* (or *E_1 -monoid*) to be a simplicial object

$$M: \Delta^{\text{op}} \rightarrow C, \quad [n] \mapsto M_n := M([n]),$$

satisfying the *Segal condition*: for every $[n] \in \Delta$ the *Segal map*

$$(e_i^*)_{i=1}^n: M_n \rightarrow \prod_{i=1}^n M_1$$

induced by the n inclusion maps $e_i: [1] \cong \{i-1 \leq i\} \hookrightarrow [n]$ is an isomorphism. In particular, the condition for $n = 0$ requires M_0 to be a terminal object of C . We write

$$\text{Mon}(C) \subseteq \text{Fun}(\Delta^{\text{op}}, C)$$

for the full subcategory spanned by the monoids in C . We refer to M_1 as the *underlying object* of M .

Convention 5.1.2. Given a monoid M , we will often denote its underlying object in C again by M . We will also often identify M_n with the n -fold product $M^n := \prod_{i=1}^n M$ via the Segal map.

The unique map $s^0: [1] \rightarrow [0]$ and the inclusion $d^1: [1] \cong \{0 \leq 2\} \hookrightarrow [2]$ in Δ induce maps $M_0 \rightarrow M_1$ and $M_2 \rightarrow M_1$, which under our convention take the form

$$e: * \rightarrow M \quad \text{and} \quad m: M \times M \rightarrow M.$$

We refer to e as the *unit* or *neutral element* of M , and refer to m as the *multiplication* of M . The unitality and associativity of multiplication are encoded by the following commutative diagrams in Δ , where we denote an order-preserving map $[m] \rightarrow [n]$ by listing the images $(a_0, \dots, a_m)$ of the elements $0, \dots, m$:

$$\begin{array}{ccc} & [1] & \\ \swarrow & \downarrow (0,2) & \searrow \\ [1] & [2] & [1] \\ \swarrow (0,0,1) & \searrow (0,1,1) & \end{array} \quad \text{and} \quad \begin{array}{ccc} [1] & \xrightarrow{(0,2)} & [2] \\ (0,2) \downarrow & & \downarrow (0,2,3) \\ [2] & \xrightarrow{(0,1,3)} & [3]. \end{array}$$

Similarly, all the expected ‘higher coherences’ relating the various ways of multiplying four, five, or more elements in M follow from suitable higher dimensional diagrams in Δ . More explicitly, for a map $\varphi: [m] \rightarrow [n]$ in Δ we should think of the resulting map $\varphi^*: M^n \rightarrow M^m$ as follows:

$$\varphi^*(x_1, \dots, x_n) := (y_1, \dots, y_m), \quad y_i := \prod_{\varphi(i-1) < j \leq \varphi(i)} x_j.$$

This formula becomes transparent once we think of the elements $x_1, \dots, x_n$ as labels on the n inequalities of $[n]$, and likewise of $y_1, \dots, y_m$ as labels on the m inequalities of $[m]$. Drawing the map φ then cuts the strip between $[m]$ and $[n]$ into m blocks, one for each inequality of $[m]$, and the label of an inequality of $[m]$ is the product of the labels it sees in its block. For $\varphi: [4] \rightarrow [5]$ with $\varphi(0) = 0$, $\varphi(1) = \varphi(2) = 2$, $\varphi(3) = 4$ and $\varphi(4) = 5$ this looks as follows:

$$\begin{array}{ccccccccccc}
 [4] = \{ & 0 & \leq & 1 & \leq & 2 & \leq & 3 & \leq & 4 & \} \\
 \downarrow \varphi & \downarrow & & \swarrow & \searrow & & \downarrow & \downarrow & & \downarrow & \\
 & & y_1 = x_1 x_2 & & y_2 = e & & y_3 = x_3 x_4 & & y_4 = x_5 & & \\
 & & \swarrow & \searrow & & \swarrow & \searrow & & \swarrow & \searrow & \\
 [5] = \{ & 0 & \leq & 1 & \leq & 2 & \leq & 3 & \leq & 4 & \leq & 5 & \}
 \end{array} \tag{5.1}$$

The i -th block is bounded by the arrows starting at $i - 1$ and at i , so the inequalities of $[5]$ it contains are exactly those indexed by $\varphi(i - 1) < j \leq \varphi(i)$; multiplying their labels gives y_i . Note that the strip is read against the direction of φ : the labels travel from $[5]$ up to $[4]$, which is precisely the contravariance of M . Since $\varphi(1) = \varphi(2)$, the two arrows bounding the second block meet at 2, so this block contains no inequality of $[5]$ at all; its label is therefore the empty product, that is, the unit e . In this example we thus obtain

$$\varphi^*(x_1, \dots, x_5) = (x_1 x_2, e, x_3 x_4, x_5).$$

Exercise 5.1.3. Let $F: C \rightarrow D$ be a functor preserving finite products. Show that F induces a functor $\text{Mon}(C) \rightarrow \text{Mon}(D)$.

Example 5.1.4. By Exercise 5.1, a topological monoid is a monoid object in Top . Combining the preceding exercise with the fact that $\Pi_\infty: \text{Top} \rightarrow \text{An}$ preserves finite products (see Corollary 2.3.20), we see that every topological monoid M defines a monoid object $\Pi_\infty(M)$ in An .

Definition 5.1.5. We say that a monoid $M \in \text{Mon}(C)$ is *grouplike*, or a *group in C* , if the so-called *shear map*

$$(\text{pr}_1, m): M \times M \rightarrow M \times M, \quad (x, y) \mapsto (x, xy)$$

is an isomorphism in C . We denote by $\text{Grp}(C) \subseteq \text{Mon}(C)$ the full subcategory spanned by the groups in C .

Exercise 5.1.6. Show that a monoid $M \in \text{Mon}(\text{Set})$ is a group in the sense of Definition 5.1.5 if and only if every element x admits an inverse x^{-1} .

Exercise 5.1.7. For a group object $G \in \text{Grp}(C)$, construct an *inverse map* $(-)^{-1}: G \rightarrow G$ in C .

5.1.1 Group structures on loop spaces

We will now introduce a general class of examples of group objects in ∞ -categories: the loop objects ΩX on pointed objects X . If C is an ∞ -category with finite limits, then we may apply Exercise 4.3 in the pointed ∞ -category C_* to obtain a group structure on ΩX in $\text{Ho}(C_*)$, and hence in $\text{Ho}(C)$. We will now explain how to lift this structure to a group structure in C .

Let us first explain informally where the multiplication on ΩX comes from. The loop object is defined as the pullback $\Omega X = * \times_X *$; for $C = \text{An}$, its points are the loops in X at the basepoint x .

Multiplication should be given by composing loops, and it is the universal property of pullbacks that produces this map for us: consider the iterated pullback $* \times_X * \times_X *$, which comes with three natural projections to $* \times_X *$. Projecting away the third or the first factor yields two maps which exhibit the iterated pullback as the product $\Omega X \times \Omega X$, as verified in Lemma 5.1.13. Projecting away the *middle* factor instead yields a map

$$m: \Omega X \times \Omega X \rightarrow \Omega X,$$

and this is composition of loops. Similarly, the universal property of the pullback $\Omega X = * \times_X *$ turns the constant homotopy at the basepoint into a unit map $e: * \rightarrow \Omega X$, the constant loop.

The higher coherences arise from the same principle. The $(n+1)$ -fold iterated pullback $* \times_X \cdots \times_X *$ is the object of n -tuples of composable loops, and projecting away some of the inner factors realizes all the ways of composing consecutive loops in such a tuple. For example, associativity compares the two ways of composing three loops, and both are obtained from projections $* \times_X * \times_X * \times_X * \rightarrow * \times_X *$ which forget the two inner factors one at a time; the universal property identifies both composites with the projection onto the two outer factors. Rather than listing such coherences one at a time, we should remember the entire family of iterated pullbacks together with all projections between them, as well as the maps which repeat a factor and correspond to inserting a constant loop. This is precisely the structure of a simplicial object satisfying the Segal condition: face maps compose loops, and degeneracy maps insert constant loops.

It remains to construct this simplicial object, with all its coherences, functorially from the basepoint map $x: * \rightarrow X$. This will be achieved by a right Kan extension, indexed by the following enlargement of the simplex category.

Definition 5.1.8. We define the *augmented simplex category* as $\Delta_+ := \Delta^{\natural}$, the cone of the simplex category. We denote the cone point by $[-1]$. A functor $\Delta_+^{\text{op}} \rightarrow C$ is called an *augmented simplicial object*. It corresponds to a simplicial object $X_\bullet: \Delta^{\text{op}} \rightarrow C$ equipped with a cocone $X_\bullet \Rightarrow \text{const}_{X_{-1}}$.

Note that we may identify Δ_+ with the full subcategory of the category Poset of posets spanned by the linearly ordered sets $[n] = \{0 \leq \cdots \leq n\}$, where we set $[-1] = \emptyset$. There are fully faithful inclusions

$$i: \Delta \hookrightarrow \Delta_+ \quad \text{and} \quad j: [1] \hookrightarrow \Delta_+, \quad j(0) = [-1], \quad j(1) = [0].$$

Definition 5.1.9. Let C be an ∞ -category with pullbacks. We define the *Čech nerve functor* $\check{C}: \text{Ar}(C) \rightarrow \text{Fun}(\Delta^{\text{op}}, C)$ as the composite

$$\check{C}_\bullet: \text{Ar}(C) \simeq \text{Fun}([1]^{\text{op}}, C) \xrightarrow{j_*} \text{Fun}(\Delta_+^{\text{op}}, C) \xrightarrow{i^*} \text{Fun}(\Delta^{\text{op}}, C),$$

where j_* is the right Kan extension functor along the inclusion $j^{\text{op}}: [1]^{\text{op}} \hookrightarrow \Delta_+^{\text{op}}$, and where i^* is restriction along the inclusion i^{op} . Given a morphism $f: X \rightarrow Y$ in C , we refer to the simplicial object $\check{C}_\bullet(f)$ as the *Čech nerve of f* .

Here the equivalence $\text{Ar}(C) \simeq \text{Fun}([1]^{\text{op}}, C)$ uses the order-reversing isomorphism $[1] \cong [1]^{\text{op}}$: a morphism $f: X \rightarrow Y$ corresponds to the functor with value X at 1 and Y at 0. In the pointwise calculation below, we use the same order reversal to write it in the usual form $f: [1] \rightarrow C$, with $f(0) = X$ and $f(1) = Y$.

For a discussion of the theory of Kan extensions, we refer the reader to Section 21.4.

$$I_n := \begin{array}{c} 0 \\ 1 \\ \vdots \\ n-1 \\ n \end{array} \rightarrow \infty$$
$$f_n = \begin{array}{ccc} & X & \\ & \searrow f & \\ X & & \\ & \nearrow & \\ \vdots & & \\ X & \nearrow f & \\ & \nwarrow & \\ & X & \end{array} Y.$$
$$\check{C}_n(f) \simeq X^{\times_Y^{n+1}} = X \times_Y X \times_Y \cdots \times_Y X.$$
$$\Omega(X, x)_1 \cong \Omega(X, x),$$
$$(e_i^*)_{i=1}^n : (\Omega X)_n \rightarrow \prod_{i=1}^n \Omega X$$

is an isomorphism in C . For $n = 0$, observe that I_0 is isomorphic to the walking morphism [1], consisting of objects 0 and ∞ , and the functor $f_0: I_0 \rightarrow C$ is the diagram $* \rightarrow X$. Since [1] has 0 as an initial object, this limit is simply $*$; see Lemma 21.2.4. The case $n = 1$ is the identification from the preceding remark. For $n \geq 1$, note that we may write I_{n+1} as a pushout of I_n and I_1 along I_0 :

$$\begin{array}{ccc} I_0 & \longrightarrow & I_n \\ \downarrow & \lrcorner & \downarrow \\ I_1 & \xrightarrow{I_{e_{n+1}}} & I_{n+1} \end{array}$$

By applying Theorem 21.2.11(2) with $J = \lrcorner$, we thus obtain a pullback square of the form

$$\begin{array}{ccc} \lim(f_{n+1}: I_{n+1} \rightarrow C) & \longrightarrow & \lim(f_n: I_n \rightarrow C) \\ \downarrow & \lrcorner & \downarrow \\ \lim(f_1: I_1 \rightarrow C) & \longrightarrow & \lim(f_0: I_0 \rightarrow C), \end{array}$$

where we take f to be the basepoint map $x: * \rightarrow X$. By induction, this pullback square is isomorphic to the commutative square

$$\begin{array}{ccc} (\Omega X)_{n+1} & \xrightarrow{(e_i^*)_{i=1}^n} & \prod_{i=1}^n \Omega X \\ e_{n+1}^* \downarrow & & \downarrow \\ \Omega X & \longrightarrow & *, \end{array}$$

and so this being a pullback square means that the map $(\Omega X)_{n+1} \xrightarrow{(e_i^*)_{i=1}^{n+1}} \prod_{i=1}^{n+1} \Omega X$ is an isomorphism, finishing the induction step. This shows that ΩX is a monoid in C .

To see that ΩX is grouplike, write $p_{ij}: * \times_X * \times_X * \rightarrow * \times_X *$ for the projection retaining the i -th and j -th factors. Under the Segal isomorphism

$$(p_{01}, p_{12}): * \times_X * \times_X * \xrightarrow{\sim} \Omega X \times \Omega X,$$

the shear map is identified with (p_{01}, p_{02}) . But associativity of pullbacks gives another isomorphism

$$(p_{01}, p_{02}): * \times_X * \times_X * \xrightarrow{\sim} (* \times_X *) \times_* (* \times_X *) \cong \Omega X \times \Omega X.$$

Thus the shear map is an isomorphism. □

5.1.2 Classifying animae

The loop group functor admits a left adjoint:

Construction 5.1.14 (Classifying anima of a monoid). Let C be an ∞ -category admitting finite products and geometric realizations. Given a monoid M , we define its *classifying anima* $\mathbf{B}M$ as

$$\mathbf{B}M := |M| = \operatorname{colim}_{[n] \in \Delta^{\text{op}}} M_n \in C.$$

We may turn this into a pointed object in C via the map $* = M_0 \rightarrow \operatorname{colim}_{[n] \in \Delta^{\text{op}}} M_n$. This construction is functorial in M , resulting in a functor $\mathbf{B}: \operatorname{Mon}(C) \rightarrow C_*$.

Proposition 5.1.15. *Assume that C admits both finite limits and geometric realizations. Then the functors $\mathbf{B}$ and Ω define adjunctions*

$$\mathbf{B} : \text{Mon}(C) \rightleftarrows C_* : \Omega \quad \text{and} \quad \mathbf{B} : \text{Grp}(C) \rightleftarrows C_* : \Omega.$$

Proof. Consider the following two adjunctions:

$$\text{Fun}(\Delta^{\text{op}}, C) \begin{array}{c} \xrightarrow{i_!} \\ \perp \\ \xleftarrow{i^*} \end{array} \text{Fun}((\Delta_+)^{\text{op}}, C) \begin{array}{c} \xrightarrow{j^*} \\ \perp \\ \xleftarrow{j_*} \end{array} \text{Fun}([1]^{\text{op}}, C).$$

The bottom composite i^*j_* is precisely the Čech nerve functor $\check{C}_\bullet$. The functor $i_!$ is given by left Kan extension along i . From the pointwise formula for left Kan extensions, we see that the top composite $j^*i_!$ sends a simplicial object X in C to the map $X_0 \rightarrow |X| = \text{colim}_n X_n$.

Now, note that the top composite $j^*i_!$ sends the full subcategory $\text{Mon}(C) \subseteq \text{Fun}(\Delta^{\text{op}}, C)$ into the full subcategory $C_* \subseteq \text{Ar}(C)$, and that the resulting functor $\text{Mon}(C) \rightarrow C_*$ is precisely $\mathbf{B}$. Similarly, the bottom composite i^*j_* restricts to $\Omega : C_* \rightarrow \text{Mon}(C)$. It follows that the adjunction $j^*i_! \dashv i^*j_*$ restricts to an adjunction $\mathbf{B} \dashv \Omega$ between $\text{Mon}(C)$ and C_* . Since Ω takes values in $\text{Grp}(C) \subseteq \text{Mon}(C)$, this adjunction further restricts to an adjunction between $\text{Grp}(C)$ and C_* . $\square$

For later use, we record the following observation:

Lemma 5.1.16. *The classifying anima functor $\mathbf{B} : \text{Mon}(\text{An}) \rightarrow \text{An}$ preserves finite products.*

Proof. This functor factors as

$$\text{Mon}(\text{An}) \hookrightarrow \text{sAn} \xrightarrow{|\cdot|} \text{An}.$$

The geometric realization functor $|\cdot|$ preserves finite products by Lemma 21.6.12, and the inclusion preserves finite products since monoids are closed under products in sAn . $\square$

5.2 The recognition principle for loop spaces

The goal of this section is to prove the following theorem:

Theorem 5.2.1 (Recognition principle for loop spaces, cf. Stasheff [Sta63], May [May72], Boardman and Vogt [BV73], Lurie [Lur09b, Theorem 7.2.2.11]). *Let M be a monoid in An and let X be a pointed anima.*

- (1) *The unit map $M \rightarrow \Omega \mathbf{B} M$ is an isomorphism if and only if M is a group in An ;*
- (2) *The pointed anima $\mathbf{B} M$ is connected;*
- (3) *The counit $\mathbf{B} \Omega X \rightarrow X$ is an inclusion of the connected component of the basepoint of X .*

The classical formulation of this result is in terms of the operadic language of A_∞ -spaces or E_1 -spaces: it says that any grouplike E_1 -space is equivalent to a loop space. The ∞ -categorical formulation in terms of the adjunction $\mathbf{B} \dashv \Omega$ has the advantage that it immediately provides us with an equivalence of ∞ -categories:

Corollary 5.2.2. *The adjunction $\mathbf{B} \dashv \mathbf{\Omega}$ restricts to an equivalence of ∞ -categories*

$$\mathbf{B}: \mathrm{Grp}(\mathrm{An}) \xrightleftharpoons{\cong} \mathrm{An}_{*, \geq 1} : \mathbf{\Omega}$$

between group objects in An and connected pointed animae.

Proof. The adjunction from Proposition 5.1.15 restricts to groups and pointed connected animae since $\mathbf{\Omega}(X)$ is always a group and $\mathbf{B}M$ is always connected. The unit $M \rightarrow \mathbf{\Omega} \mathbf{B}M$ is always an isomorphism by part (1) of the theorem, while the counit $\mathbf{B} \mathbf{\Omega} X \rightarrow X$ is always an isomorphism by part (3) of the theorem. This shows that the two functors are in fact inverse to one another. $\square$

5.2.1 Proof of the recognition principle

The proof of Theorem 5.2.1 will occupy the remainder of this section; a reader willing to take the theorem on faith may safely continue with Section 5.3. Let us first explain the two main ingredients of the proof. To identify a grouplike monoid M with $\mathbf{\Omega} \mathbf{B}M$, we must compute the pullback square defining the loop anima of $\mathbf{B}M$ and identify it with the underlying anima of M . We will obtain this from a descent property of the ∞ -category An , by comparing the simplicial object M with its *décalage* PM , obtained by shifting every simplicial degree up by one. The shifted object admits extra degeneracies, which force its geometric realization to be contractible, while grouplikeness of M ensures that the comparison map $PM \rightarrow M$ is cartesian. Descent then turns this levelwise statement into the desired pullback square after geometric realization.

Extra degeneracies

We record a useful technique for computing geometric realizations.

Definition 5.2.3 (Extra degeneracies). We define the subcategory $\Delta^{\mathrm{deg}} \subseteq \Delta$ as the subcategory consisting of all objects $[n]$, but with only those morphisms $\varphi: [n] \rightarrow [m]$ satisfying $\varphi(n) = m$. An *augmented simplicial object with extra degeneracies* in an ∞ -category C is a functor $Y: (\Delta^{\mathrm{deg}})^{\mathrm{op}} \rightarrow C$.

There exists a functor $(+1): \Delta_+ \hookrightarrow \Delta^{\mathrm{deg}}$ given on objects by $[n] \mapsto [n+1]$, and on morphisms by sending $\varphi: [n] \rightarrow [m]$ to the map

$$[n+1] \rightarrow [m+1], \quad i \mapsto \begin{cases} \varphi(i) & 0 \leq i \leq n \\ m+1 & i = n+1. \end{cases}$$

Given an augmented simplicial object with extra degeneracies $Y: (\Delta^{\mathrm{deg}})^{\mathrm{op}} \rightarrow C$, precomposition with $(+1)$ yields the *underlying augmented simplicial object* of Y . Conversely, we say that an augmented simplicial object $X: \Delta_+^{\mathrm{op}} \rightarrow C$ *admits extra degeneracies* if there exists an augmented simplicial object with extra degeneracies Y such that X is isomorphic to the underlying augmented simplicial object of Y .

To understand the structure of Δ^{deg} , note that all the face maps that are allowed in Δ^{deg} are actually already in the image of the functor $(+1): \Delta_+ \hookrightarrow \Delta^{\mathrm{deg}}$. Among the degeneracy maps, the map $s_n: [n+1] \rightarrow [n]$ in Δ^{deg} is the only one which does not lie in this image. Hence an augmented simplicial

object with extra degeneracies can be pictured as follows, with non-dashed arrows encoding the underlying augmented simplicial object and with dashed arrows encoding the ‘extra degeneracies’:

$$\dots \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} Y_2 \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} Y_1 \xrightarrow{\quad} Y_0.$$

The utility of extra degeneracies lies in the following easy computation of geometric realizations:

Lemma 5.2.4 (Extra degeneracy argument). *Let $X: \Delta_+^{\text{op}} \rightarrow C$ be an augmented simplicial object that admits extra degeneracies. Then X is a colimit diagram, in the sense that the corresponding cocone $X|_{\Delta^{\text{op}}} \Rightarrow \text{const}_{X_{-1}}$ is a colimit cocone:*

$$\text{colim}_{[n] \in \Delta^{\text{op}}} X_n \xrightarrow{\sim} X_{-1}.$$

Proof. By assumption, there exists an augmented simplicial object with extra degeneracies $Y: (\Delta^{\text{deg}})^{\text{op}} \rightarrow C$ such that $X_n = Y_{n+1}$. We need to show that the preferred map from $\text{colim}_{[n] \in \Delta^{\text{op}}} X_n$ to X_{-1} induced by the cocone is an isomorphism.

Observe that the inclusion functor $(+1): \Delta \hookrightarrow \Delta^{\text{deg}}$ admits a right adjoint given by the inclusion $\Delta^{\text{deg}} \hookrightarrow \Delta$: for natural numbers $n, m \geq 0$, providing a map $\varphi: [n] \rightarrow [m]$ of partially ordered sets is the same as providing a map $\bar{\varphi}: [n+1] \rightarrow [m]$ of posets satisfying the condition that $\bar{\varphi}(n+1) = m$. It follows from Corollary 21.5.6 that the functor $(+1): \Delta \hookrightarrow \Delta^{\text{deg}}$ is an initial functor, and hence its opposite $(+1)^{\text{op}}: \Delta^{\text{op}} \hookrightarrow (\Delta^{\text{deg}})^{\text{op}}$ is a final functor. By Theorem 21.5.1, this means that the geometric realization of X may be computed as

$$\text{colim}_{[n] \in \Delta^{\text{op}}} X_n = \text{colim}_{[n] \in \Delta^{\text{op}}} Y_{n+1} \simeq \text{colim}_{[m] \in (\Delta^{\text{deg}})^{\text{op}}} Y_m.$$

But since Δ^{deg} has an initial object given by $[0]$, $(\Delta^{\text{deg}})^{\text{op}}$ has a terminal object, and so it follows from Lemma 21.2.4 that $\text{colim}_{[m] \in (\Delta^{\text{deg}})^{\text{op}}} Y_m$ is isomorphic to $Y_0 = X_{-1}$ as desired. $\square$

Descent

The ∞ -category An has a special feature that distinguishes it from other ∞ -categories: it satisfies *descent*. In the following statement, we use the notation $I^{\flat} := (I \times [1]) \times_{I \times \{1\}} *$ from Definition 1.7.5.

Theorem 5.2.5 (Descent for colimits). *Let I be a small ∞ -category, let $\bar{F}, \bar{G}: I^{\flat} \rightarrow \text{An}$ be functors and let $\bar{\alpha}: \bar{F} \Rightarrow \bar{G}$ be a natural transformation such that the restriction $\alpha := \bar{\alpha}|_I: F \Rightarrow G$ is a cartesian transformation, in the sense that for every morphism $i \rightarrow j$ in I the commutative square*

$$\begin{array}{ccc} F(i) & \longrightarrow & F(j) \\ \alpha(i) \downarrow & & \downarrow \alpha(j) \\ G(i) & \longrightarrow & G(j) \end{array} \quad (5.2)$$

is a pullback square. Assume that $\bar{G}$ is a colimit diagram. Then $\bar{F}$ is a colimit diagram if and only if $\bar{\alpha}$ is a cartesian transformation, i.e. also all the squares

$$\begin{array}{ccc} F(i) & \longrightarrow & \bar{F}(\infty) \\ \alpha(i) \downarrow & & \downarrow \bar{\alpha}(\infty) \\ G(i) & \longrightarrow & \bar{G}(\infty) = \text{colim}_I G. \end{array}$$

are pullback squares.

Proof. See Section 23.7; see also [Lur09b, Theorem 6.1.3.9]. $\square$

The theorem has two complementary uses. If both $\overline{F}$ and $\overline{G}$ are colimit diagrams and α is cartesian, then every $F(i)$ can be recovered by pulling back the map $\text{colim}_I F \rightarrow \text{colim}_I G$ along $G(i) \rightarrow \text{colim}_I G$. Conversely, if only $\overline{G}$ is known to be a colimit diagram, cartesianness of the extended transformation $\overline{\alpha}$ forces $\overline{F}$ to be a colimit diagram as well. This second form is the one we will use below. See Section 23.7 for the conceptual interpretation in terms of slice categories.

Example 5.2.6 (Universality of coproducts). Consider $I = * \sqcup *$, so that $I^\triangleright \simeq \sqcup$. The transformation $\overline{\alpha}$ will thus be a diagram of the form

$$\begin{array}{ccccc} X_0 & \longrightarrow & X_2 & \longleftarrow & X_1 \\ \downarrow \alpha_0 & & \downarrow \alpha_2 & & \downarrow \alpha_1 \\ Y_0 & \longrightarrow & Y_0 \sqcup Y_1 & \longleftarrow & Y_1, \end{array}$$

where in the bottom we get a coproduct, reflecting the condition that $\overline{G}$ is a colimit diagram. The theorem then expresses that the following two conditions are equivalent:

- The two squares in the diagram are both pullback squares;
- The functor $X_0 \sqcup X_1 \rightarrow X_2$ is an isomorphism.

The fact that these two conditions are equivalent is precisely an instance of the universality of coproducts from Axiom C.3, with the two parts of the axiom corresponding to the two implications.

The décalage argument

We are now ready to prove the recognition principle.

Proof of Theorem 5.2.1. We start with (2): the fact that $\mathbf{B}M$ is connected for any monoid object M in $\mathbf{An}$. By definition, $\mathbf{B}M$ is connected if and only if its set of path components $\pi_0(\mathbf{B}M)$ is a single point. Since $\pi_0: \mathbf{An} \rightarrow \mathbf{Set}$ is a left adjoint, it preserves colimits (see Lemma 21.2.6), and thus $\pi_0(\mathbf{B}M)$ is a colimit in sets of the following simplicial diagram:

$$\dots \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \pi_0(M_2) \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \pi_0(M_1) \begin{array}{c} \rightrightarrows \\ \rightrightarrows \\ \rightrightarrows \end{array} \pi_0(M_0).$$

The colimit of a simplicial object in sets is the coequalizer of its two face maps in degrees 1 and 0. Hence the canonical map $\pi_0(M_0) \rightarrow \text{colim}_{[n]} \pi_0(M_n)$ is surjective. Since $M_0 \cong *$, it follows that $\pi_0(\mathbf{B}M)$ has a single element.

Let us assume (1) for a moment and use it to deduce (3), the fact that the counit $\varepsilon_X: \mathbf{B}\Omega X \rightarrow X$ is the inclusion of the path component X_0 of the basepoint of X . Since $\mathbf{B}\Omega X$ is connected, the counit map certainly lands in X_0 and we need to show that the resulting map $\mathbf{B}\Omega X \rightarrow X_0$ is an isomorphism of animae. Since both sides are connected pointed animae, it suffices by Corollary 2.4.22 to show that the map $\Omega(\varepsilon_X): \Omega(\mathbf{B}\Omega X) \rightarrow \Omega(X_0) \simeq \Omega(X)$ is an isomorphism. By the triangle identities for

the adjunction, there is a commutative triangle of the form

$$\begin{array}{ccc} & \Omega(\mathbf{B}\Omega X) & \\ \eta_{\Omega X} \nearrow & & \searrow \Omega(\varepsilon_X) \\ \Omega X & \xlongequal{\quad} & \Omega X, \end{array}$$

and so $\Omega(\varepsilon_X)$ is an isomorphism if and only if $\eta_{\Omega}: \Omega X \rightarrow \Omega(\mathbf{B}\Omega X)$ is an isomorphism. But since ΩX is a group object, this is an instance of part (1).

It remains to prove part (1), i.e. that M is a group if and only if the map $\eta_M: M \rightarrow \Omega \mathbf{B}M$ is an isomorphism of monoids. One direction is clear: $\Omega \mathbf{B}M$ is a group, so if this map is an isomorphism then also M is a group. For the other direction, assume that M is a group. To show that $\eta_M: M \rightarrow \Omega \mathbf{B}M$ is an isomorphism, it suffices to show that the map $(\eta_M)_1: M_1 \rightarrow (\Omega \mathbf{B}M)_1 = \Omega \mathbf{B}M$ on underlying animae is an isomorphism: since $M_n \simeq M_1^{\times n}$ and $(\Omega \mathbf{B}M)_n \simeq ((\Omega \mathbf{B}M)_1)^{\times n}$ we then get that it is an isomorphism at every level. The map $M_1 \rightarrow \Omega \mathbf{B}M$ is induced by the commutative square

$$\begin{array}{ccc} M_1 & \xrightarrow{d_0} & M_0 \simeq * \\ d_1 \downarrow & & \downarrow \\ M_0 \simeq * & \longrightarrow & \mathbf{B}M \end{array}$$

and so is an isomorphism if and only if this is a pullback square of animae. We will deduce this as an instance of Theorem 5.2.5. To this end, consider the diagram $\overline{M}: (\Delta_+)^{\text{op}} \rightarrow \mathbf{An}$ that extends M with its colimit $\overline{M}_{-1} := \mathbf{B}M = \text{colim}_{[n] \in \Delta^{\text{op}}} M_n$. Also consider the functor

$$\overline{PM}: (\Delta_+)^{\text{op}} \rightarrow \mathbf{An}, \quad [n] \mapsto \overline{M}_{n+1},$$

and consider the natural transformation $\overline{\alpha}: \overline{PM} \rightarrow \overline{M}$ whose components are the maps $d_{n+1}: M_{n+1} \rightarrow M_n$. Let us spell out why these maps assemble functorially. The successor functor $P: \Delta_+ \rightarrow \Delta_+$, obtained by adjoining a new final element, admits a natural transformation $\text{id} \rightarrow P$ whose component at $[n]$ is the final coface $d^{n+1}: [n] \rightarrow [n+1]$. Precomposition with $\overline{M}: (\Delta_+)^{\text{op}} \rightarrow \mathbf{An}$ therefore gives $\overline{\alpha}: P^* \overline{M} = \overline{PM} \rightarrow \overline{M}$; at $[-1]$ its component is the augmentation $M_0 \rightarrow \mathbf{B}M$. This is the usual décalage construction. The successor functor is the composite $\Delta_+ \xrightarrow{(+1)} \Delta^{\text{deg}} \hookrightarrow \Delta_+$ from Definition 5.2.3. Thus $\overline{PM}$ is the underlying augmented simplicial object of $\overline{M}|_{(\Delta^{\text{deg}})^{\text{op}}}$ and admits extra degeneracies. It is therefore a colimit diagram by Lemma 5.2.4, so that $\text{colim}_{[n] \in \Delta^{\text{op}}} \overline{PM}_n \simeq \overline{PM}_{-1} = M_0$. The pullback square we wish to prove now takes the form

$$\begin{array}{ccc} PM_0 & \longrightarrow & \overline{PM}_{-1} \\ \downarrow \alpha_0 & & \downarrow \overline{\alpha}_{-1} \\ M_0 & \longrightarrow & \overline{M}_{-1}. \end{array}$$

By Theorem 5.2.5, it thus remains to show that α is a cartesian natural transformation. In other words, we need to show that for every morphism $\varphi: [n] \rightarrow [m]$ the square

$$\begin{array}{ccc} PM_m & \xrightarrow{\varphi^*} & PM_n \\ \alpha_m \downarrow & & \downarrow \alpha_n \\ M_m & \xrightarrow{\varphi^*} & M_n \end{array} \tag{5.3}$$

is a pullback square. We will prove this claim in increasingly general cases:

Step 1: We start with the case of the map $\varphi = d^1: [0] \rightarrow [1]$. In this case, the square (5.3) reduces to the following:

$$\begin{array}{ccc} M_1^2 & \xrightarrow{m} & M_1 \\ \text{pr}_1 \downarrow & & \downarrow \\ M_1 & \longrightarrow & *. \end{array} \quad (5.4)$$

This square is a pullback square if and only if the shear map $(\text{pr}_1, m): M_1 \times M_1 \rightarrow M_1 \times M_1$ is an isomorphism, which holds by assumption since M is a group.

Step 2: We will now prove the claim for $\varphi = d^i: [n] \rightarrow [n+1]$, where $n \geq 0$ and $0 \leq i \leq n+1$. We use the identifications $M_k \cong M_1^k$ coming from the Segal condition. For $i = 0$, the square (5.3) is the square which drops the first coordinate horizontally and the last coordinate vertically. It is a pullback because a pair of $(n+1)$ -tuples with the same middle n coordinates glues uniquely to an $(n+2)$ -tuple. For $0 < i < n+1$, the horizontal maps multiply the i -th and $(i+1)$ -st coordinates, while the vertical maps drop the last coordinate. Since the multiplication does not involve the last coordinate, this square is the product with M_1 of a degenerate pullback square, and is therefore again a pullback.

It remains to consider the final coface $i = n+1$. Here the multiplication involves the coordinate which is forgotten by the vertical maps, and the square takes the form

$$\begin{array}{ccc} M_1^{n+2} & \xrightarrow{(m_1, \dots, m_{n+2}) \mapsto (m_1, \dots, m_n, m_{n+1}m_{n+2})} & M_1^{n+1} \\ (m_1, \dots, m_{n+2}) \mapsto (m_1, \dots, m_{n+1}) \downarrow & & \downarrow (x_1, \dots, x_{n+1}) \mapsto (x_1, \dots, x_n) \\ M_1^{n+1} & \xrightarrow{(m_1, \dots, m_{n+1}) \mapsto (m_1, \dots, m_n)} & M_1^n. \end{array}$$

None of the maps affect the first n coordinates, and on the last two coordinates this is precisely the square (5.4). Hence it is the product of a degenerate pullback square with the pullback square from Step 1. This proves the claim for all coface maps d^i .

Step 3: Next, assume that φ is a codegeneracy map of the form $s^i: [n+1] \rightarrow [n]$. Since s^i has a section of the form $d^i: [n] \rightarrow [n+1]$, it follows from the pasting law of pullback squares that the square (5.3) is a pullback square:

$$\begin{array}{ccccc} PM_n & \xrightarrow{s_i^*} & PM_{n+1} & \xrightarrow{d_i^*} & PM_n \\ \alpha_n \downarrow & & \alpha_{n+1} \downarrow & \lrcorner & \downarrow \alpha_n \\ M_n & \xrightarrow{s_i^*} & M_{n+1} & \xrightarrow{d_i^*} & M_n. \end{array}$$

Step 4: We have now shown the claim when φ is either a coface map d^i or a codegeneracy map s^i . Since every map $\varphi: [n] \rightarrow [m]$ is an iterated composite of coface maps and codegeneracy maps, we then get the claim for general φ by using the pasting law of pullback squares again. This finishes the proof of the theorem. $\square$

5.3 Commutative monoids and commutative groups

We now move to *commutative* monoids in an ∞ -category. Just as for monoids, these will be defined as functors out of a suitable indexing category that encodes all the operations we have in a commutative monoid.

For a commutative monoid M , given a tuple of elements $(x_s)_{s \in S}$ indexed by a finite set S , we can form the sum $\sum_{s \in S} x_s$. Commutativity says precisely that this sum is independent of any chosen ordering of S . Writing $M^S := \prod_{s \in S} M$ for the S -fold product, we thus have addition maps $M^S \rightarrow M$ for every finite set S .

How do these addition maps behave with respect to maps of finite sets? Let $\mathbf{Fin}$ denote the category of finite sets and all maps between them. For a map $g: S \rightarrow T$ of finite sets, there is an induced map $g_\oplus: M^S \rightarrow M^T$ given by summing over fibers: the t -th component of $g_\oplus((x_s)_{s \in S})$ is $\sum_{s \in g^{-1}(\{t\})} x_s$. This makes the assignment $S \mapsto M^S$ into a functor $\mathbf{Fin} \rightarrow \mathbf{Set}$.

Just as for monoids, we should not look at arbitrary functors $\mathbf{Fin} \rightarrow C$, but only at those satisfying a Segal-type condition of the form $M(S) \cong M(*)^S$. To phrase this condition precisely, we need to specify projections $M(S) \rightarrow M(*)$ for each $s \in S$, that is, we need an *a priori* way to compare $M(S)$ with the product $\prod_{s \in S} M(*)$. Observe that the S -fold product M^S is not only covariantly functorial in S via addition, but also *contravariantly* functorial via restriction: a map $f: S \rightarrow T$ induces a restriction map $f^*: M^T \rightarrow M^S$ given by $f^*(x_t)_{t \in T} := (x_{f(s)})_{s \in S}$.

To encode commutative monoids, we need both functorialities, addition and restriction, packaged into a single category. The objects of this category are finite sets. What should the morphisms from S to T be? These should encode the operation of ‘first restricting along f , then adding along g ’: given maps $f: U \rightarrow S$ and $g: U \rightarrow T$, we obtain a composite

$$M^S \xrightarrow{f^*} M^U \xrightarrow{g_\oplus} M^T,$$

where the component of the output at $t \in T$ is given by summing $x_{f(u)}$ over all $u \in g^{-1}(\{t\})$. Thus, a morphism from S to T should be a pair of maps $S \leftarrow U \rightarrow T$, that is, a *span* from S to T .

What about composition? Suppose we first apply $S \xleftarrow{f} U \xrightarrow{g} T$ and then $T \xleftarrow{h} V \xrightarrow{k} R$. The first operation sends $(x_s)_{s \in S}$ to $(y_t)_{t \in T}$ where $y_t = \sum_{u \in g^{-1}(\{t\})} x_{f(u)}$. Applying the second operation, the component at $r \in R$ is

$$\sum_{v \in k^{-1}(\{r\})} y_{h(v)} = \sum_{v \in k^{-1}(\{r\})} \sum_{u \in g^{-1}(\{h(v)\})} x_{f(u)}.$$

This double sum ranges over pairs (u, v) with $g(u) = h(v)$ and $k(v) = r$. Setting $P := \{(u, v) \in U \times V \mid g(u) = h(v)\} = U \times_T V$, with its projections $h': P \rightarrow U$ and $g': P \rightarrow V$, we see that:

- the fiber $g'^{-1}(\{h(v)\})$ is in bijection with the fiber $(g')^{-1}(\{v\})$, and
- the element $f(u)$ can be written as $(f \circ h')(u, v)$.

Hence the double sum equals $\sum_{p \in (k \circ g')^{-1}(\{r\})} x_{(f \circ h')(p)}$, which is exactly the result of applying the span $S \xleftarrow{f \circ h'} P \xrightarrow{k \circ g'} R$. In other words, composing ‘first add along g , then restrict along h ’ gives the same result as ‘first restrict along h' , then add along g' ’. Spans thus compose by taking pullbacks, leading to the *span category of finite sets*.

Construction 5.3.1 (Span category). There is an ∞ -category $\text{Span}(\text{Fin})$, called the *span category of finite sets*. Its objects are finite sets, and for finite sets S and T its hom animae are given by the groupoids

$$\text{Hom}_{\text{Span}(\text{Fin})}(S, T) := \left\{ \begin{array}{ccc} & U & \\ f \swarrow & \downarrow \cong & \searrow g \\ S & & T \\ f' \swarrow & \downarrow & \searrow g' \\ & U' & \end{array} \right\}$$

of *spans from S to T* ; so the objects of this groupoid are spans $S \leftarrow U \rightarrow T$ and the morphisms are isomorphisms of spans. The identity span is given by $S \xleftarrow{\text{id}_S} S \xrightarrow{\text{id}_S} S$, and composition of spans is defined by taking pullbacks:

$$\begin{array}{ccccc} & & U \times_T V & & \\ & \swarrow h' & & \searrow g' & \\ & U & & V & \\ f \swarrow & & g & & \searrow h \\ S & & T & & R. \end{array}$$

Remark 5.3.2. Since the hom animae of $\text{Span}(\text{Fin})$ are groupoids rather than sets, $\text{Span}(\text{Fin})$ is a $(2, 1)$ -category rather than a 1-category. The rigorous construction of this ∞ -category is deferred to Chapter 13; for now, it suffices to know that $\text{Span}(\text{Fin})$ exists and behaves as described above.

Lemma 5.3.3. *The ∞ -category $\text{Span}(\text{Fin})$ is semiadditive, with biproducts given by disjoint unions of finite sets.*

Proof. Consider finite sets S_i for $i = 1, \dots, n$. We have spans of the form

$$\bigsqcup_{i=1}^n S_i \hookrightarrow S_i \xrightarrow{\text{id}_{S_i}} S_i$$

We claim that these morphisms exhibit $\bigsqcup_{i=1}^n S_i$ as a product of the sets S_i in $\text{Span}(\text{Fin})$. Indeed, note that for every map $U \rightarrow \bigsqcup_{i=1}^n S_i$ of finite sets we may write $U = \bigsqcup_{i=1}^n U_i$, where U_i is the preimage of S_i . This gives the desired equivalence of groupoids

$$\text{Hom}_{\text{Span}(\text{Fin})}(T, \bigsqcup_{i=1}^n S_i) = \left\{ \begin{array}{ccc} & \bigsqcup_{i=1}^n U_i & \\ f \swarrow & \downarrow \cong & \searrow \bigsqcup_{i=1}^n g_i \\ T & & \bigsqcup_{i=1}^n S_i \\ f' \swarrow & \downarrow & \searrow \bigsqcup_{i=1}^n g'_i \\ & \bigsqcup_{i=1}^n U'_i & \end{array} \right\} \simeq \prod_{i=1}^n \text{Hom}_{\text{Span}(\text{Fin})}(T, S_i).$$

By the dual argument (reversing a span gives a bijection between spans from S to T and spans from T to S) we see that similarly the spans

$$S_i \xleftarrow{\text{id}_{S_i}} S_i \hookrightarrow \bigsqcup_{i=1}^n S_i.$$

exhibit $\bigsqcup_{i=1}^n S_i$ also as the *coproduct* of the S_i in $\text{Span}(\text{Fin})$. The composite from S_i into this coproduct and then to the j -th product factor is represented by the pullback $S_i \times_{\bigsqcup_k S_k} S_j$. It is the

identity span when $i = j$ and the zero span otherwise. Hence the canonical comparison from the coproduct to the product is an isomorphism. For $n = 0$, the same argument says that the empty set is a zero object. Thus $\text{Span}(\text{Fin})$ is semiadditive. $\square$

Definition 5.3.4 (Commutative monoid). Let C be an ∞ -category with finite products. We define a *commutative monoid* in C to be a product-preserving functor

$$M: \text{Span}(\text{Fin}) \rightarrow C.$$

We denote by

$$\text{CMon}(C) \subseteq \text{Fun}(\text{Span}(\text{Fin}), C)$$

the full subcategory spanned by the commutative monoids. We refer to the evaluation $M(*)$ at the one-point set as the *underlying object* of M . We will sometimes abuse notation and simply refer to $M(*)$ as M .

Remark 5.3.5 (Restriction and addition maps). Let $M: \text{Span}(\text{Fin}) \rightarrow C$ be a commutative monoid. Every map $f: S \rightarrow T$ of finite sets determines two ‘half-degenerate’ spans

$$f^* = (T \xleftarrow{f} S \xrightarrow{\text{id}_S} S) \quad \text{and} \quad f_{\oplus} = (S \xleftarrow{\text{id}_S} S \xrightarrow{f} T),$$

called the *restriction* and the *addition* spans, respectively. Since products in $\text{Span}(\text{Fin})$ are computed as disjoint unions by Lemma 5.3.3, the condition that M preserves finite products means that for every finite set S the map

$$(e_s^*)_{s \in S}: M(S) \rightarrow M^S := \prod_{s \in S} M(*)$$

is an isomorphism, where $e_s: \{s\} \hookrightarrow S$ is the inclusion. We will henceforth identify $M(S)$ with M^S . It follows that every map $f: S \rightarrow T$ defines both a restriction map $f^*: M^T \rightarrow M^S$ as well as an addition map $f_{\oplus}: M^S \rightarrow M^T$, functorially in f . As a special case, the maps $\emptyset \rightarrow *$ and $* \sqcup * \rightarrow *$ in Fin then yield the *unit* and *addition*:

$$0: * \rightarrow M \quad \text{and} \quad +: M \times M \rightarrow M.$$

Definition 5.3.6 (Commutative group). A commutative monoid M is called a *commutative group* if the *shear map* $(\text{pr}_1, +): M \times M \rightarrow M \times M$ is an isomorphism. We denote by

$$\text{CGrp}(C) \subseteq \text{CMon}(C)$$

the full subcategory spanned by the commutative groups in C .

5.3.1 Alternative definition

We now give an alternative characterization of commutative monoids that is commonly used in the literature and goes back to Segal [Seg68]. This characterization uses the category of finite pointed sets rather than the full span category.

Definition 5.3.7 (Segal condition). Let C be an ∞ -category with finite products. Denote by Fin_* the category of *finite pointed sets*, whose objects are pairs $(S, *)$ consisting of a finite set S and a

distinguished basepoint $*$ $\in S$, and whose morphisms are basepoint-preserving maps. For a finite set S , we write $S_+ := S \sqcup \{*\}$ for S with an added disjoint basepoint.

A functor $M: \mathbf{Fin}_* \rightarrow C$ is said to satisfy the *Segal condition* if for every finite set S the canonical map

$$M(S_+) \rightarrow \prod_{s \in S} M(\{s\}_+)$$

induced by the pointed maps $S_+ \rightarrow \{s\}_+$ is an isomorphism, where the pointed map sends everything except for $s \in S$ to the base point of $\{s\}_+$. We denote by

$$\mathbf{Fun}^{\text{Segal}}(\mathbf{Fin}_*, C) \subseteq \mathbf{Fun}(\mathbf{Fin}_*, C)$$

the full subcategory spanned by functors satisfying the Segal condition.

Lemma 5.3.8. *Consider the wide subcategory $\text{Span}(\mathbf{Fin}, \text{inj}, \text{all})$ of $\text{Span}(\mathbf{Fin})$ whose morphisms are those spans $S \hookleftarrow U \rightarrow T$ for which the left-pointing map is an injection. This is a 1-category, and there is an equivalence of 1-categories $\mathbf{Fin}_* \simeq \text{Span}(\mathbf{Fin}, \text{inj}, \text{all})$.*

Proof. It is clear that $\text{Span}(\mathbf{Fin}, \text{inj}, \text{all})$ is a 1-category: given two injections $U \hookleftarrow S$ and $U' \hookleftarrow S$, if there exists a bijection $U \xrightarrow{\sim} U'$ over S , then it is unique.

We now construct mutually inverse functors to $\mathbf{Fin}_*$. Define a functor

$$\Phi: \text{Span}(\mathbf{Fin}, \text{inj}, \text{all}) \rightarrow \mathbf{Fin}_*$$

as follows: on objects, Φ sends a finite set S to $S_+ = S \sqcup \{*\}$. On morphisms, Φ sends a span $S \hookleftarrow U \rightarrow T$ (where the left-pointing map is an injection) to the pointed map $S_+ \rightarrow T_+$ that agrees with $U \rightarrow T$ on U and sends the complement $S \setminus U$ (together with the basepoint of S_+) to the basepoint of T_+ .

Conversely, define a functor

$$\Psi: \mathbf{Fin}_* \rightarrow \text{Span}(\mathbf{Fin}, \text{inj}, \text{all})$$

as follows: on objects, Ψ sends a finite pointed set $(S, *)$ to $S \setminus \{*\}$. On morphisms, Ψ sends a pointed map $f: (S, *) \rightarrow (T, *)$ to the span

$$S \setminus \{*\} \hookleftarrow f^{-1}(T \setminus \{*\}) \xrightarrow{f} T \setminus \{*\},$$

where the left-pointing map is the inclusion. One readily verifies that Φ and Ψ are inverse equivalences. $\square$

Proposition 5.3.9. *Let C be an ∞ -category with finite products. Restriction along the inclusion $i: \mathbf{Fin}_* \simeq \text{Span}(\mathbf{Fin}, \text{inj}, \text{all}) \hookrightarrow \text{Span}(\mathbf{Fin})$ induces an equivalence of ∞ -categories*

$$i^*: \mathbf{CMon}(C) \xrightarrow{\sim} \mathbf{Fun}^{\text{Segal}}(\mathbf{Fin}_*, C).$$

The inverse is given by right Kan extension along i .

Proof. It is clear from the definitions that restriction along i sends product-preserving functors to functors satisfying the Segal condition.

The functor i^* is conservative: if $\alpha: M \rightarrow N$ is a natural transformation between commutative monoids such that $i^*\alpha$ is an isomorphism, then in particular $\alpha_*: M(*) \rightarrow N(*)$ is an isomorphism, and since both M and N preserve products we have that the map $\alpha_S: M(S) \cong M(*)^S \rightarrow N(*)^S \cong N(S)$ is an isomorphism for all finite sets S .

It thus suffices to show that for every functor $M: \text{Fin}_* \rightarrow C$ satisfying the Segal condition, the right Kan extension $i_*(M)$ exists and the counit map $i^*i_*(M) \rightarrow M$ is an isomorphism. By the pointwise formula for right Kan extensions, we must show that for every finite set X the canonical map

$$\lim_{(X \rightarrow S) \in i_{X/}} M(S) \longrightarrow M(X)$$

is an isomorphism, where the limit is taken over the relative slice category $i_{X/} = \text{Span}(\text{Fin}, \text{inj}, \text{all}) \times_{\text{Span}(\text{Fin})} \text{Span}(\text{Fin})_{X/}$. Consider the embedding

$$\varphi: (\text{Fin}^{\text{op}})_{X/} \times_{\text{Fin}^{\text{op}}} \text{Fin}_{\text{inj}}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin}, \text{inj}, \text{all}) \times_{\text{Span}(\text{Fin})} \text{Span}(\text{Fin})_{X/}$$

that sends a map $f: S \rightarrow X$ to the span $X \xleftarrow{f} S \xrightarrow{\text{id}} S$. This functor φ has a right adjoint, given by sending a span $X \leftarrow U \rightarrow S$ to the left-pointing map $U \rightarrow X$. By Corollary 21.5.6, the functor φ is therefore initial, so the limit may be computed over the left-hand category.

On this category, the restriction of M sends a map $f: S \rightarrow X$ to $M(S) \cong M(*)^S = \lim_{s \in S} M(*)$ (using the Segal condition). By the formula for iterated limits from Theorem 21.2.11, we may thus write the left-hand side as the limit of $M(*)$ over the set $\text{colim}_{(X \rightarrow S)} S$. It thus suffices to show that X is the colimit of the forgetful functor

$$\text{Fin}_{/X} \times_{\text{Fin}} \text{Fin}_{\text{inj}} \longrightarrow \text{Fin}.$$

This functor is left Kan extended from the full subcategory spanned by the inclusions $\{x\} \hookrightarrow X$ for $x \in X$, and the colimit over this subcategory is precisely X . Moreover, the product projections in $\text{Span}(\text{Fin})$ have injective left legs and therefore belong to the wide subcategory on which i is defined. The product comparison maps for i_*M consequently agree, under the counit, with the Segal maps of M and are isomorphisms. Thus i_*M preserves finite products. $\square$

Remark 5.3.10. The ∞ -category $\text{CMon}(C)$ is usually defined as $\text{Fun}^{\text{Segal}}(\text{Fin}_*, C)$ in the literature; this was also the original approach taken by Segal [Seg68] after which the Segal condition is named. Proposition 5.3.9 shows that our definition agrees with this classical one.

5.3.2 The underlying monoid of a commutative monoid

As the name suggests, every commutative monoid has an underlying monoid.

Construction 5.3.11. We will construct a functor $\text{Cut}: \Delta^{\text{op}} \rightarrow \text{Span}(\text{Fin}, \text{inj}, \text{all}) \subseteq \text{Span}(\text{Fin})$:

- On objects, it sends $[n]$ to the set $\langle n \rangle := \{1, \dots, n\}$.
- On morphisms, it sends $\varphi: [m] \rightarrow [n]$ to the span

$$\langle n \rangle \longleftarrow \{\varphi(0) + 1, \varphi(0) + 2, \dots, \varphi(m) - 1, \varphi(m)\} \longrightarrow \langle m \rangle,$$

where the left-pointing map is the inclusion and the right-pointing map sends i to the unique $1 \leq j \leq m$ such that $\varphi(j - 1) < i \leq \varphi(j)$.

More conceptually, we may think of the set $\text{Cut}(P)$ for a poset P as the set of ‘Dedekind cuts’ of P , defined as pairs (P_0, P_1) of non-empty subposets $P_0, P_1 \subseteq P$ satisfying $P = P_0 \cup P_1$ and $p_0 < p_1$ for all $p_0 \in P_0$ and $p_1 \in P_1$. We want to think of the element $k \in \{1, \dots, n\} = \text{Cut}([n])$ as the following partition of $[n]$:

$$k \quad \leftrightarrow \quad (\{0, \dots, k-1\}, \{k, \dots, n\}).$$

On morphisms, $\text{Cut}(\varphi)$ is then simply given by taking preimages of the two subsets P_0 and P_1 , which is only defined if both of these preimages are non-empty.

Alternatively, we may think of the set $\text{Cut}([n])$ as labeling the n inequalities in the partially ordered set $[n]$ that we can use to cut $[n]$ into two pieces:

$$[n] \quad = \quad \{0 \stackrel{1}{\leq} 1 \stackrel{2}{\leq} 2 \stackrel{3}{\leq} \dots \stackrel{n}{\leq} n\}.$$

This is the bookkeeping already used in Equation (5.1): the middle term of the span $\text{Cut}(\varphi)$ consists of those inequalities of $[n]$ which lie in one of the blocks cut out by φ , the right-pointing map records for each of them the inequality of $[m]$ whose block contains it, and the left-pointing inclusion discards the inequalities of $[n]$ lying in no block at all.

Exercise 5.3.12. Let $M: \text{Span}(\text{Fin}) \rightarrow C$ be a commutative monoid. Show that $M \circ \text{Cut}: \Delta^{\text{op}} \rightarrow C$ defines a monoid in C , whose multiplication agrees with the addition map of M . Deduce that $M \circ \text{Cut}$ is a group if and only if M is a commutative group, and that there is a pullback square

$$\begin{array}{ccc} \text{CGrp}(C) & \longrightarrow & \text{CMon}(C) \\ \downarrow & & \downarrow \\ \text{Grp}(C) & \longrightarrow & \text{Mon}(C). \end{array}$$

In particular, we obtain forgetful functors

$$U := \text{Cut}^*: \text{CMon}(C) \rightarrow \text{Mon}(C) \quad \text{and} \quad U := \text{Cut}^*: \text{CGrp}(C) \rightarrow \text{Grp}(C).$$

Warning 5.3.13. For a classical 1-category C , the forgetful functor $\text{CMon}(C) \rightarrow \text{Mon}(C)$ is fully faithful by Exercise 5.6. For a general ∞ -category, this is *not* true: commutativity is no longer a *property* of a monoid, but additional *structure* one needs to provide.

5.3.3 Monoids and commutative monoids in semiadditive ∞ -categories

For a semiadditive ∞ -category C , it turns out that every object admits a unique structure of a (commutative) monoid. The idea is simple: in a semiadditive category, every object M comes equipped with a canonical addition map

$$+: M \times M \xrightarrow{\sim} M \oplus M \xrightarrow{\nabla} M,$$

given by the fold map of the biproduct, and a zero map $0: * \rightarrow M$, the unique map from the initial object. It readily follows from the universal property of coproducts that this operation is unital, associative and commutative, resulting in a commutative monoid object in the homotopy category $\text{Ho}(C)$. To promote M to a genuine commutative monoid object in C , we need these identities

to hold coherently. While this is again a consequence of the universal property of coproducts, the required bookkeeping is more difficult as we need to handle all coherences simultaneously. The goal of this subsection is to carry out this bookkeeping. Only the statements of the results below will be used later, and the proofs all follow the same pattern of restriction and Kan extension along suitable subcategories, so they may be skimmed on a first reading. Let us start with the non-commutative case.

Proposition 5.3.14. *Let C be a semiadditive ∞ -category. Then the forgetful functor $\text{Mon}(C) \rightarrow C$ is an equivalence. If C is additive, then also the forgetful functor $\text{Grp}(C) \rightarrow C$ is an equivalence.*

Proof. Let $\Delta_{\text{int}}^{\leq 1} \subseteq \Delta$ be the subcategory with objects $[0]$ and $[1]$ and whose only non-identity morphisms are the two face maps $d^0, d^1: [0] \rightarrow [1]$. We will show that restriction along the inclusions

$$j: \{[1]\} \hookrightarrow (\Delta_{\text{int}}^{\leq 1})^{\text{op}} \quad \text{and} \quad i: (\Delta_{\text{int}}^{\leq 1})^{\text{op}} \hookrightarrow \Delta^{\text{op}}$$

define equivalences

$$\text{Mon}(C) \xrightarrow{\sim} \text{Fun}^*((\Delta_{\text{int}}^{\leq 1})^{\text{op}}, C) \xrightarrow{\sim} C,$$

where Fun^* denotes the full subcategory of those functors whose value at $[0]$ is terminal.

For the first equivalence, we use left Kan extension along i . We apply the pointwise formula (Theorem 21.4.3) to compute the left Kan extension. For this, we need to understand the relative slice $i_{/[n]}$ for each $[n] \in \Delta$. Objects of this category are pairs (x, φ) where $x \in \{[0], [1]\}$ and $\varphi: x \rightarrow [n]$ is a morphism in Δ^{op} , i.e. a morphism $[n] \rightarrow x$ in Δ . There are $n+2$ order-preserving maps $[n] \rightarrow [1]$, which we may label by the sequence of values they take: there are the two *constant* maps $\underline{0}, \underline{1}: [n] \rightarrow [1]$, and for each $1 \leq k \leq n$ a map φ_k sending $0, 1, \dots, k-1$ to 0 and $k, \dots, n$ to 1 . There is also a unique map $\star: [n] \rightarrow [0]$. For morphisms in $i_{/[n]}$, we observe that the only non-identity morphisms in $(\Delta_{\text{int}}^{\leq 1})^{\text{op}}$ are the maps $[1] \rightarrow [0]$ induced by the face maps $d^0, d^1: [0] \rightarrow [1]$. By checking the commutative triangles, one sees that the only non-identity morphisms in $i_{/[n]}$ are the maps $\underline{0} \rightarrow \star$ and $\underline{1} \rightarrow \star$. It follows that the category $i_{/[n]}$ has precisely $n+1$ connected components: there is a ‘span’ component $\underline{0} \rightarrow \star \leftarrow \underline{1}$, and there are n isolated objects $\varphi_1, \dots, \varphi_n$.

Now let $F: (\Delta_{\text{int}}^{\leq 1})^{\text{op}} \rightarrow C$ be a functor with $F([0]) = *$, and write $X := F([1])$. The left Kan extension $i_!F$ evaluated at $[n]$ is the colimit over $i_{/[n]}$ of the composite $i_{/[n]} \rightarrow (\Delta_{\text{int}}^{\leq 1})^{\text{op}} \xrightarrow{F} C$. This colimit decomposes as a coproduct over the connected components. The span component contributes the colimit $\text{colim}(X \rightarrow * \leftarrow X) = *$, the terminal object, which is also initial in the pointed category C . Each isolated object φ_k contributes a copy of X . Hence $(i_!F)([n]) \simeq X^{\sqcup n}$, the n -fold coproduct. Under these identifications, the Segal map $(i_!F)([n]) \rightarrow (i_!F)([1])^n$ is the canonical comparison from the n -fold coproduct of X to its n -fold product, hence is an isomorphism by semiadditivity. Thus $i_!F$ is a monoid.

Conversely, let $M: \Delta^{\text{op}} \rightarrow C$ be a monoid. There is a canonical comparison map from the left Kan extension of the restriction, $i_!i^*M \rightarrow M$. To show this is an isomorphism, it suffices to check this at the underlying object $[1]$. There it is clear: as we just computed, the value at $[1]$ of both sides is simply $M([1])$.

For the second equivalence, we use right Kan extension along j . Since j is fully faithful, right Kan extension j_* is fully faithful whenever it exists. The relative slice $j_{x/}$ over $x \in \{[0], [1]\}$ is as follows: for $x = [1]$, the identity $\text{id}_{[1]}$ is an initial object, so the limit evaluates to the original value.

For $x = [0]$, the relative slice is empty, so the limit is the terminal object. This shows that right Kan extension along j always exists and produces a functor in $\text{Fun}^*((\Delta_{\text{int}}^{\leq 1})^{\text{op}}, C)$. Conversely, any functor with value $*$ at $[0]$ is right Kan extended from $\{[1]\}$.

For the claim about groups, note that if C is additive and M is a monoid in C with underlying object X , then the shear map is given by $\begin{pmatrix} \text{id}_X & \text{id}_X \\ 0 & \text{id}_X \end{pmatrix}: X^2 \rightarrow X^2$. Its inverse is $\begin{pmatrix} \text{id}_X & -\text{id}_X \\ 0 & \text{id}_X \end{pmatrix}$, so every monoid is automatically a group. $\square$

Remark 5.3.15. Once we know that $\text{CMon}(C)$ is semiadditive (which will be proved in general in Proposition 5.3.19), the present proposition applied to $\text{CMon}(C)$ gives an equivalence $\text{Mon}(\text{CMon}(C)) \simeq \text{CMon}(C)$, under which the canonical monoid structure on an object $M \in \text{CMon}(C)$ has multiplication given by the fold map $M \oplus M \rightarrow M$, which agrees with the algebraic addition $+: M \times M \rightarrow M$ of the commutative monoid structure. As a consequence, the monoid shear map (pr_1, m) from Definition 5.1.5 coincides with the algebraic shear map $(\text{pr}_1, +)$ from Definition 5.3.6, and there are canonical identifications

$$\text{CMon}(\text{Grp}(C)) \simeq \text{Grp}(\text{CMon}(C)) \simeq \text{CGrp}(C).$$

The first equivalence is an instance of currying functors $\text{Span}(\text{Fin}) \times \Delta^{\text{op}} \rightarrow C$: being a commutative monoid in the first coordinate commutes with being a group in the second. The second equivalence uses that both sides are the full subcategory of $\text{Mon}(\text{CMon}(C)) \simeq \text{CMon}(C)$ spanned by those objects whose shear map is invertible.

Having established the case of monoids and groups, we now turn to commutative monoids and commutative groups.

Proposition 5.3.16. *Let C be a semiadditive ∞ -category. Then the forgetful functors*

$$\text{CMon}(C) \rightarrow \text{Mon}(C) \rightarrow C$$

are equivalences. If C is additive, then also the forgetful functors

$$\text{CGrp}(C) \rightarrow \text{Grp}(C) \rightarrow C$$

are equivalences.

Proof. The equivalences $\text{Mon}(C) \xrightarrow{\sim} C$ and $\text{Grp}(C) \xrightarrow{\sim} C$ are Proposition 5.3.14. By Exercise 5.3.12, we have $\text{CGrp}(C) \simeq \text{CMon}(C) \times_{\text{Mon}(C)} \text{Grp}(C)$. The case for commutative groups therefore follows from that of commutative monoids, so it remains to show that the forgetful functor $\text{CMon}(C) \rightarrow C$ is an equivalence.

The proof strategy is similar to, but more involved than, that of Proposition 5.3.14. Consider the following subcategories of $\text{Span}(\text{Fin})$:

$$* \hookrightarrow \text{Fin}^{\leq 1} \hookrightarrow \text{Span}(\text{Fin}^{\leq 1}) \hookrightarrow \text{Span}(\text{Fin}, \text{inj}, \text{all}) \hookrightarrow \text{Span}(\text{Fin}).$$

Here $\text{Fin}^{\leq 1} \subseteq \text{Fin}$ is the full subcategory on sets of cardinality at most 1, i.e. it has objects $\emptyset$ and $*$ and a single non-identity morphism $\emptyset \rightarrow *$. We claim that restriction along each of these inclusions defines equivalences

$$\begin{aligned} \text{CMon}(C) &= \text{Fun}^{\times}(\text{Span}(\text{Fin}), C) \xrightarrow{\sim} \text{Fun}^{\text{Segal}}(\text{Span}(\text{Fin}, \text{inj}, \text{all}), C) \\ &\xrightarrow{\sim} \text{Fun}^*(\text{Span}(\text{Fin}^{\leq 1}), C) \xrightarrow{\sim} \text{Fun}^*(\text{Fin}^{\leq 1}, C) \xrightarrow{\sim} C, \end{aligned}$$

with inverses given by right, left, right and left Kan extension, respectively. Here $\text{Fun}^*(-, -)$ denotes those functors that send the empty set to the terminal object of C . For the first equivalence, this was proved in Proposition 5.3.9. The proof strategy for the other three equivalences is similar. We will spell out the details for the second equivalence, and leave the easier third and fourth equivalences to the reader.

Denoting the relevant inclusion by $i: \text{Span}(\text{Fin}^{\leq 1}) \hookrightarrow \text{Span}(\text{Fin}, \text{inj}, \text{all})$, we need to show that the restriction functor

$$i^*: \text{Fun}^{\text{Segal}}(\text{Span}(\text{Fin}, \text{inj}, \text{all}), C) \rightarrow \text{Fun}^*(\text{Span}(\text{Fin}^{\leq 1}), C)$$

is an equivalence. Note that by semiadditivity of C , a functor $F: \text{Span}(\text{Fin}, \text{inj}, \text{all}) \rightarrow C$ satisfies the Segal condition if and only if its restriction $F|_{\text{Fin}}: \text{Fin} \rightarrow C$ is left Kan extended from the point. Similarly, a functor $G: \text{Span}(\text{Fin}^{\leq 1}) \rightarrow C$ satisfies $G(\emptyset) = *$ if and only if its restriction $G|_{\text{Fin}^{\leq 1}}: \text{Fin}^{\leq 1} \rightarrow C$ is left Kan extended from the point. We now claim that left Kan extension along i restricts to a functor

$$i_!: \text{Fun}^*(\text{Span}(\text{Fin}^{\leq 1}), C) \rightarrow \text{Fun}^{\text{Segal}}(\text{Span}(\text{Fin}, \text{inj}, \text{all}), C).$$

To see this, let $G: \text{Span}(\text{Fin}^{\leq 1}) \rightarrow C$ be a functor satisfying $G(\emptyset) = *$. It will suffice to show that the restriction $i_!(G)|_{\text{Fin}}$ of the left Kan extension of G is itself the left Kan extension of $G|_{\text{Fin}^{\leq 1}}$, since then it must be Kan extended from the point. If we let $i': \text{Fin}^{\leq 1} \rightarrow \text{Fin}$ denote the inclusion, there is a canonical map

$$i'_!(G|_{\text{Fin}^{\leq 1}}) \rightarrow i_!(G)|_{\text{Fin}}.$$

Using the pointwise formula for Kan extensions, Theorem 21.4.3, we may compute both sides as colimits of G over the relative slices of i' and i , and the previous comparison map is induced by the inclusion map $\text{Fin}^{\leq 1}_{/S} \rightarrow \text{Span}(\text{Fin}^{\leq 1})_{/S}$ of relative slices, for $S \in \text{Fin}$. Since this inclusion admits a left adjoint (given by taking the forward map of a span into S , see Exercise 5.7), this functor is final by Corollary 21.5.6, and hence restriction along it does not change the value of the colimit.

Finally we observe that the Kan extension $i_!G$ has the same value on the point as G itself. From this one concludes that the unit $G \rightarrow i^*i_!G$ and counit $i_!i^*F \rightarrow F$ of the adjunction are both natural isomorphisms, and hence these two functors are inverse equivalences. $\square$

5.3.4 Semiadditivity

One of the archetypical examples of a semiadditive category is the 1-category of commutative monoids: the product $A \times B$ of two commutative monoids is also the coproduct. In this section, we will generalize this observation: given an arbitrary ∞ -category C with finite products, the ∞ -category $\text{CMon}(C)$ is semiadditive. Combining this fact with the results from the previous section, we may conclude that the forgetful functor $\text{CMon}(C) \rightarrow C$ is in fact *universal* among semiadditive ∞ -categories equipped with a finite-product-preserving functor to C . Similarly, the forgetful functor $\text{CGrp}(C) \rightarrow C$ is universal among finite-product-preserving functors $D \rightarrow C$ from an *additive* ∞ -category.

We start with two preliminary lemmas:

Lemma 5.3.17. *Let C be an ∞ -category with finite products. Then the ∞ -category $\text{CMon}(C)$ has finite products and the evaluation map $\text{ev}_T: \text{CMon}(C) \rightarrow C$ preserves finite products for each $T \in \text{Fin}$.*

Proof. The functor category $\text{Fun}(\text{Span}(\text{Fin}), C)$ admits finite products (Lemma 21.2.8) which are computed pointwise, and the subcategory $\text{CMon}(C)$ is closed under finite products. $\square$

Lemma 5.3.18. *Let C be an ∞ -category with finite products. Then there exists a cotensoring functor*

$$(-)^{(-)} : \text{Span}(\text{Fin}) \times \text{CMon}(C) \rightarrow \text{CMon}(C), \quad (S, X) \mapsto X^S,$$

defined by $X^S(T) := X(S \times T)$, which preserves finite products in both variables and whose restriction to $\{\} \times \text{CMon}(C)$ is the identity functor.*

Proof. For a finite set S and a commutative monoid X we define the cotensoring by $X^S(T) := X(S \times T)$ for $T \in \text{Fin}$. The functoriality in S and T comes from the observation that the product functor $- \times - : \text{Fin} \times \text{Fin} \rightarrow \text{Fin}$ induces a functor on span categories by taking products of spans:

$$- \times - : \text{Span}(\text{Fin}) \times \text{Span}(\text{Fin}) \rightarrow \text{Span}(\text{Fin}), \quad (S, T) \mapsto S \times T.$$

(This is no longer the categorical product in $\text{Span}(\text{Fin})$!)

- Since we have a natural bijection $* \times T \cong T$, it is clear that the cotensoring is the identity on $\{*\} \times \text{CMon}(C)$.
- Fixing $S \in \text{Fin}$, the functor $(-)^S : \text{CMon}(C) \rightarrow \text{CMon}(C)$ preserves finite products: we may check this pointwise for every $T \in \text{Fin}$, where it is by definition given by the evaluation $X \mapsto X(S \times T)$, which preserves finite products.
- Fixing $X \in \text{CMon}(C)$, the functor $X^{(-)} : \text{Span}(\text{Fin}) \rightarrow \text{CMon}(C)$ preserves finite products: again we may check this pointwise for every $T \in \text{Fin}$, where we may write it as the composite

$$\text{Span}(\text{Fin}) \xrightarrow{- \times T} \text{Span}(\text{Fin}) \xrightarrow{X} C.$$

But X preserves finite products by assumption and $- \times T$ preserves finite products since the functor $- \times T : \text{Fin} \rightarrow \text{Fin}$ preserves finite coproducts: $\bigsqcup_{i=1}^n (S_i \times T) \xrightarrow{\sim} (\bigsqcup_{i=1}^n S_i) \times T$.

In particular we see that X^S is the S -fold product of X in $\text{CMon}(C)$, explaining the notation. $\square$

Proposition 5.3.19. *Let C be an ∞ -category with finite products. Then the ∞ -category $\text{CMon}(C)$ is semiadditive. The subcategory $\text{CGrp}(C)$ is even additive.*

Proof. We first show that $\text{CMon}(C)$ is semiadditive. This amounts to two claims: first, $\text{CMon}(C)$ has a zero object, and second, finite products are also coproducts.

For the first claim, note that $\text{Span}(\text{Fin})$ is pointed, with zero object given by the empty set. By Corollary 4.1.12, the ∞ -category $\text{Fun}_*(\text{Span}(\text{Fin}), C)$ is pointed, with zero object given by the constant functor $\text{const}_* : \text{Span}(\text{Fin}) \rightarrow C$. Since $\text{CMon}(C) \subseteq \text{Fun}_*(\text{Span}(\text{Fin}), C)$ is the full subcategory spanned by the finite-product-preserving functors and contains const_* , this same object is also a zero object of $\text{CMon}(C)$.

For the second claim, we show that for commutative monoids X and Y , the product $X \times Y$ equipped with the maps $(\text{id}_X, 0) : X \rightarrow X \times Y$ and $(0, \text{id}_Y) : Y \rightarrow X \times Y$ is also a coproduct. In other words, we need to show that the corresponding natural transformation

$$((\text{id}_X, 0), (0, \text{id}_Y)) : (X, Y) \rightarrow (X \times Y, X \times Y) = \Delta(X \times Y)$$

is the unit of an adjunction between the product functor and the diagonal functor $\Delta: \mathbf{CMon}(C) \rightarrow \mathbf{CMon}(C) \times \mathbf{CMon}(C)$ (with the product being the *left* adjoint!). By Proposition 21.1.2, it suffices to supplement this unit by a counit satisfying the triangle identities. We therefore have to produce for every third commutative monoid Z a compatible counit map

$$m_Z: Z \times Z \rightarrow Z$$

satisfying the triangle identities; this is the map that will a posteriori correspond to the fold map $Z \oplus Z \rightarrow Z$. Using the cotensoring from Lemma 5.3.18, we define m_Z as the composite

$$m_Z: Z \times Z \cong Z^{*\sqcup*} \xrightarrow{Z^\nabla} Z^* \cong Z,$$

where $\nabla: * \sqcup * \rightarrow *$ is the fold map in $\mathbf{Fin}$, viewed as a span $(* \sqcup * \xleftarrow{\text{id}} * \sqcup * \xrightarrow{\nabla} *)$. The triangle identities take the following form:

$$\begin{array}{ccc} & (Z \times Z, Z \times Z) & \\ ((\text{id}_Z, 0), (0, \text{id}_Z)) \nearrow & & \searrow (m_Z, m_Z) \\ (Z, Z) & \xlongequal{\quad} & (Z, Z) \end{array} \quad \text{and} \quad \begin{array}{ccc} & (X \times Y) \times (X \times Y) & \\ (\text{id}_X, 0) \times (0, \text{id}_Y) \nearrow & & \searrow m_{X \times Y} \\ X \times Y & \xlongequal{\quad} & X \times Y. \end{array}$$

For the right-hand triangle, we use that the cotensoring preserves products, so that the map $m_{X \times Y}$ may be rewritten as the product map

$$m_X \times m_Y: (X \times X) \times (Y \times Y) \rightarrow X \times Y.$$

We then see that to produce both triangles, it suffices to produce for every $X \in \mathbf{CMon}(C)$ two natural homotopies between the composites

$$X \xrightarrow{(\text{id}_X, 0)} X \times X \xrightarrow{m_X} X, \quad X \xrightarrow{(0, \text{id}_X)} X \times X \xrightarrow{m_X} X$$

and the identity on X . For this, observe that we may write the zero map $0: \text{const}_* \rightarrow X$ as the map $X^\emptyset \rightarrow X^*$ induced on cotensorings by the inclusion $\emptyset \hookrightarrow *$ in $\mathbf{Fin}$. Consequently, under the isomorphism $X^* \times X^* \cong X^{*\sqcup*}$, the maps $(\text{id}_X, 0)$ and $(0, \text{id}_X)$ are induced by the two inclusions $\iota_1, \iota_2: * \hookrightarrow * \sqcup *$. The triangle identities then reduce to the observation that the two composites in $\mathbf{Fin}$

$$* \xrightarrow{\iota_1} * \sqcup * \xrightarrow{\nabla} * \quad \text{and} \quad * \xrightarrow{\iota_2} * \sqcup * \xrightarrow{\nabla} *$$

are both the identity, which is immediate. This proves that $\mathbf{CMon}(C)$ is semiadditive.

Finally, $\mathbf{CGrp}(C) \subseteq \mathbf{CMon}(C)$ is closed under finite products and contains the zero object, so the biproducts constructed above restrict to $\mathbf{CGrp}(C)$. Their fold map is the algebraic addition m_M constructed above. Hence the additive shear map agrees with the algebraic shear map of Definition 5.3.6, which is invertible by definition. Thus $\mathbf{CGrp}(C)$ is additive. $\square$

Remark 5.3.20. Observe that there are equivalences $\mathbf{CMon}(C_*) \simeq \mathbf{CMon}(C)_*$ and $\mathbf{CGrp}(C_*) \simeq \mathbf{CGrp}(C)_*$. Since $\mathbf{CMon}(C)$ and $\mathbf{CGrp}(C)$ are pointed, it follows from Lemma 4.1.9 that the forgetful functor $C_* \rightarrow C$ induces equivalences

$$\mathbf{CMon}(C_*) \xrightarrow{\sim} \mathbf{CMon}(C) \quad \text{and} \quad \mathbf{CGrp}(C_*) \xrightarrow{\sim} \mathbf{CGrp}(C).$$

Using a similar argument we find equivalences

$$\mathbf{Mon}(C_*) \xrightarrow{\sim} \mathbf{Mon}(C) \quad \text{and} \quad \mathbf{Grp}(C_*) \xrightarrow{\sim} \mathbf{Grp}(C).$$

With this at hand, we can now prove universal properties for the assignments $C \mapsto \mathbf{CMon}(C)$ and $C \mapsto \mathbf{CGrp}(C)$.

Notation 5.3.21. Let $\mathbf{Cat}_\infty^{\text{prod}}$ be the subcategory of $\mathbf{Cat}_\infty$ spanned by those ∞ -categories that have finite products and the finite-product-preserving functors. We further denote by

$$\mathbf{Cat}_\infty^{\text{sadd}} \subseteq \mathbf{Cat}_\infty^{\text{prod}} \quad \text{and} \quad \mathbf{Cat}_\infty^{\text{add}} \subseteq \mathbf{Cat}_\infty^{\text{prod}}$$

the full subcategories spanned by the semiadditive and additive ∞ -categories, respectively.

Proposition 5.3.22. *The two functors*

$$\mathbf{CMon}(-) : \mathbf{Cat}_\infty^{\text{prod}} \rightarrow \mathbf{Cat}_\infty^{\text{sadd}} \quad \text{and} \quad \mathbf{CGrp}(-) : \mathbf{Cat}_\infty^{\text{prod}} \rightarrow \mathbf{Cat}_\infty^{\text{add}}$$

are right adjoint to the respective inclusion functors.

Proof. We will check the (dual version of the) criterion from Proposition 21.8.9. We will treat the case for $\mathbf{CMon}(-)$; the case for $\mathbf{CGrp}(-)$ is identical. The forgetful functor defines a natural transformation $\varepsilon : \mathbf{CMon}(-) \rightarrow \text{id}_{\mathbf{Cat}_\infty^{\text{prod}}}$, and we must check that for every ∞ -category C with finite products the two forgetful functors

$$\varepsilon_{\mathbf{CMon}(C)} : \mathbf{CMon}(\mathbf{CMon}(C)) \rightarrow \mathbf{CMon}(C) \quad \text{and} \quad \mathbf{CMon}(\varepsilon_C) : \mathbf{CMon}(\mathbf{CMon}(C)) \rightarrow \mathbf{CMon}(C)$$

are equivalences. Since $\mathbf{CMon}(C)$ is semiadditive by Proposition 5.3.19, the claim for $\varepsilon_{\mathbf{CMon}(C)}$ is an instance of Proposition 5.3.16. But now the case for $\mathbf{CMon}(\varepsilon_C)$ follows immediately: the swap map $\text{Span}(\text{Fin}) \times \text{Span}(\text{Fin}) \xrightarrow{\sim} \text{Span}(\text{Fin}) \times \text{Span}(\text{Fin})$ determines an equivalence

$$\mathbf{CMon}(\mathbf{CMon}(C)) \xrightarrow{\sim} \mathbf{CMon}(\mathbf{CMon}(C)), \quad (M_n)_m \mapsto (M_m)_n,$$

and under this equivalence the forgetful functor $\varepsilon_{\mathbf{CMon}(C)}((M_n)_m) = (M_n)_1$ corresponds to the forgetful functor $\mathbf{CMon}(\varepsilon_C)((M_m)_n) = (M_1)_n$. $\square$

Corollary 5.3.23. *Let A be a semiadditive ∞ -category and let C be an ∞ -category with finite products. The forgetful functor $\mathbf{CMon}(C) \rightarrow C$ induces an equivalence*

$$\text{Fun}^\times(A, \mathbf{CMon}(C)) \xrightarrow{\sim} \text{Fun}^\times(A, C).$$

Proof. By Yoneda, it suffices to show that this induces an equivalence on mapping animae out of every ∞ -category E . This follows from the natural equivalences

$$\begin{aligned} \text{Hom}_{\mathbf{Cat}_\infty}(E, \text{Fun}^\times(A, \mathbf{CMon}(C))) &\simeq \text{Hom}_{\mathbf{Cat}_\infty^{\text{prod}}}(A, \text{Fun}(E, \mathbf{CMon}(C))) \\ &\simeq \text{Hom}_{\mathbf{Cat}_\infty^{\text{prod}}}(A, \mathbf{CMon}(\text{Fun}(E, C))) \\ &\simeq \text{Hom}_{\mathbf{Cat}_\infty^{\text{prod}}}(A, \text{Fun}(E, C)) \\ &\simeq \text{Hom}_{\mathbf{Cat}_\infty}(E, \text{Fun}^\times(A, C)), \end{aligned}$$

where the second equivalence uses pointwise finite products and the third is the adjunction from Proposition 5.3.22. $\square$

As a useful consequence, we see that the hom animae of any semiadditive ∞ -category admit canonical refinements to commutative monoids.

Proposition 5.3.24. *For a semiadditive ∞ -category C , the hom functor $\mathrm{Hom}_C : C^{\mathrm{op}} \times C \rightarrow \mathbf{An}$ lifts uniquely to a functor*

$$\mathrm{Hom}_C : C^{\mathrm{op}} \times C \rightarrow \mathbf{CMon}(\mathbf{An})$$

which preserves direct sums in both variables. If C is additive, this lands in $\mathbf{CGrp}(\mathbf{An})$.

Proof. This follows directly from the universal property of $\mathbf{CMon}(\mathbf{An})$ established in Proposition 5.3.22; the proof is completely analogous to that of Proposition 4.4.4 and is omitted. $\square$

5.4 The recognition principle for connective spectra

In Section 5.2 we proved the recognition principle for loop spaces: the loop functor

$$\Omega : \mathbf{An}_{*, \geq 1} \xrightarrow{\sim} \mathbf{Grp}(\mathbf{An})$$

is an equivalence, with inverse given by the classifying anima construction $\mathbf{B} : \mathbf{Grp}(\mathbf{An}) \rightarrow \mathbf{An}_{*, \geq 1}$. In this section, we will show that this equivalence can be enhanced to an equivalence

$$\Omega^\infty(-) : \mathbf{Sp}_{\geq 0} \xrightarrow{\sim} \mathbf{CGrp}(\mathbf{An})$$

between the ∞ -category of connective spectra and the ∞ -category of commutative groups in $\mathbf{An}$. Let us start by recalling the definition of $\mathbf{Sp}_{\geq 0}$:

Definition 5.4.1. An anima X is called *0-connected* if it is connected, i.e. if the set $\pi_0(X)$ has a single element. It is called *n -connected* for $n \geq 1$ if it is connected and the homotopy groups $\pi_k(X)$ are trivial whenever $k \leq n$. It is *(-1) -connected* whenever it is non-empty. We write

$$\mathbf{An}_{\geq n} \subseteq \mathbf{An}$$

for the full subcategory spanned by the $(n-1)$ -connected anima. Thus $\mathbf{An}_{\geq 0}$ is the full subcategory of non-empty anima. Note that X is n -connected if and only if it is connected and ΩX is $(n-1)$ -connected.

A spectrum X is called *connective* if, for every $n \geq 1$, its n -th space $X_n = \Omega^{\infty-n} X$ is $(n-1)$ -connected. We write

$$\mathbf{Sp}_{\geq 0} \subseteq \mathbf{Sp}$$

for the full subcategory spanned by the connective spectra. Connective spectra are also known as *infinite loop spaces*.

Remark 5.4.2. Since we have $\pi_i(X) \cong \pi_{n+i}(X_n)$ whenever $n+i \geq 0$, we see that a spectrum X is connective if and only if $\pi_i(X) = 0$ for $i < 0$.

The point of commutativity is that the delooping process can be repeated indefinitely: a commutative group G has a delooping $\mathbf{B}G$ which is again a commutative group, then a second delooping $\mathbf{B}^2G$, and so on, and these deloopings assemble into a spectrum whose zeroth anima is G . Conversely, the underlying anima of every spectrum carries a canonical commutative group structure. We now make this precise, beginning with the delooping of commutative groups.

Proposition 5.4.3. *Applying $\mathbf{CMon}(-)$ to the adjunction $\mathbf{B} \dashv \mathbf{\Omega}$ produces an adjunction*

$$\mathbf{B}: \mathbf{CMon}(\mathbf{An}) \simeq \mathbf{CMon}(\mathbf{Mon}(\mathbf{An})) \rightleftarrows \mathbf{CMon}(\mathbf{An}_*) \xrightarrow[\simeq]{\text{Remark 5.3.20}} \mathbf{CMon}(\mathbf{An}) : \mathbf{\Omega}.$$

Moreover, both functors actually land in $\mathbf{CGrp}(\mathbf{An})$ and induce for all $n \geq 0$ an equivalence

$$\mathbf{B}: \mathbf{CGrp}(\mathbf{An}_{\geq n}) \xrightleftharpoons{\simeq} \mathbf{CGrp}(\mathbf{An}_{\geq n+1}) : \mathbf{\Omega}.$$

Proof. Both $\mathbf{B}$ and $\mathbf{\Omega}$ preserve finite products: for $\mathbf{B}$ this is Lemma 5.1.16, while $\mathbf{\Omega}$ preserves them because it is a right adjoint. We may therefore apply $\mathbf{CMon}(-)$ to the adjunction.

For the first claim, it remains to argue that the forgetful functor $\mathbf{Mon}(\mathbf{An}) \rightarrow \mathbf{An}$ induces an equivalence

$$\mathbf{CMon}(\mathbf{Mon}(\mathbf{An})) \xrightarrow{\sim} \mathbf{CMon}(\mathbf{An}).$$

Since both $\mathbf{CMon}(-)$ and $\mathbf{Mon}(-)$ are defined as certain limit-preserving functors, there is an obvious equivalence $\mathbf{CMon}(\mathbf{Mon}(\mathbf{An})) \simeq \mathbf{Mon}(\mathbf{CMon}(\mathbf{An}))$, and under this equivalence the previous functor corresponds to the forgetful functor

$$\mathbf{Mon}(\mathbf{CMon}(\mathbf{An})) \rightarrow \mathbf{CMon}(\mathbf{An}).$$

But since $\mathbf{CMon}(\mathbf{An})$ is semiadditive by Proposition 5.3.19, this forgetful functor is an equivalence by Proposition 5.3.16.

Combining Exercise 5.8 with Remark 5.3.15, a commutative monoid $X \in \mathbf{CMon}(\mathbf{An})$ is a commutative group if and only if its monoid of path components $\pi_0(X)$ is a group. We check that both functors land in $\mathbf{CGrp}(\mathbf{An})$: the anima $\mathbf{BM}$ is connected, so $\pi_0(\mathbf{BM}) = *$ is trivially a group; and for $X \in \mathbf{CMon}(\mathbf{An}_*)$ we have $\pi_0(\mathbf{\Omega}X) = \pi_1(X)$, which is a group because loop spaces are grouplike. Hence the adjunction restricts to commutative groups, and by Theorem 5.2.1 we get that it restricts to an equivalence

$$\mathbf{B}: \mathbf{CGrp}(\mathbf{An}) \xrightarrow[\simeq]{\text{Remark 5.3.15}} \mathbf{CMon}(\mathbf{Grp}(\mathbf{An})) \xrightleftharpoons{\simeq} \mathbf{CMon}(\mathbf{An}_{\geq 1}) \simeq \mathbf{CGrp}(\mathbf{An}_{\geq 1}) : \mathbf{\Omega}.$$

Here we use Remark 5.3.20 to pass between commutative monoids in pointed anima and pointed commutative monoids in anima. The claim for arbitrary n follows because $\mathbf{BG}$ is n -connected if and only if $G \simeq \mathbf{\Omega BG}$ is $(n-1)$ -connected. $\square$

Corollary 5.4.4 (Group completion). *The inclusion $\mathbf{CGrp}(\mathbf{An}) \hookrightarrow \mathbf{CMon}(\mathbf{An})$ admits a left adjoint*

$$(-)^{\text{grp}}: \mathbf{CMon}(\mathbf{An}) \rightarrow \mathbf{CGrp}(\mathbf{An})$$

called group completion. It is given on objects by $M \mapsto \mathbf{\Omega BM}$.

Proof. It follows immediately from the proposition that for every commutative group $G \in \mathbf{CGrp}(\mathbf{An})$, precomposition with the unit $M \rightarrow \mathbf{\Omega BM}$ defines an equivalence

$$\text{Hom}_{\mathbf{CGrp}(\mathbf{An})}(\mathbf{\Omega BM}, G) \xrightarrow{\sim} \text{Hom}_{\mathbf{CMon}(\mathbf{An})}(M, G),$$

since both sides are equivalent to $\text{Hom}_{\mathbf{CMon}(\mathbf{An}_*)}(\mathbf{BM}, \mathbf{BG})$. $\square$

After all this hard work, we are now able to prove the claim that connective spectra are commutative groups in $\mathbf{An}$:

Theorem 5.4.5 (Recognition principle for connective spectra, cf. May [May72], Boardman and Vogt [BV73]). *The functor $\Omega^\infty : \mathbf{Sp} \rightarrow \mathbf{An}$ uniquely lifts to a functor*

$$\mathbf{\Omega}^\infty : \mathbf{Sp} \rightarrow \mathbf{CGrp}(\mathbf{An}).$$

This functor admits a fully faithful left adjoint

$$\mathbf{B}^\infty : \mathbf{CGrp}(\mathbf{An}) \hookrightarrow \mathbf{Sp}$$

whose image is precisely the subcategory $\mathbf{Sp}_{\geq 0}$ of connective spectra.

Proof. Since Ω^∞ preserves finite products and $\mathbf{Sp}$ is additive by Lemma 4.2.17, the fact that Ω^∞ uniquely lifts to $\mathbf{CGrp}(\mathbf{An})$ is an immediate consequence of the fact that $\mathbf{CGrp}(-) : \mathbf{Cat}_\infty^{\text{prod}} \rightarrow \mathbf{Cat}_\infty^{\text{add}}$ is right adjoint to the inclusion functor, see Proposition 5.3.22.

For the left adjoint, we use the equivalence $\mathbf{B}^n : \mathbf{CGrp}(\mathbf{An}) \xrightarrow{\sim} \mathbf{CGrp}(\mathbf{An}_{\geq n})$ from Proposition 5.4.3, with inverse given by the iterated loop space functor Ω^n . For a commutative group G , we may now define the spectrum $\mathbf{B}^\infty G$ as follows:

$$\mathbf{B}^\infty G := (\mathbf{B}^n G, \sigma_n : \mathbf{B}^n G \xrightarrow{\sim} \Omega \mathbf{B}^{n+1} G),$$

where the structure maps σ_n are the units of the adjunction $\mathbf{B} \dashv \Omega$. To see that this is a left adjoint to $\Omega^\infty : \mathbf{Sp} \rightarrow \mathbf{CGrp}(\mathbf{An})$, note that by additivity of $\mathbf{Sp}$ we get an equivalence $\mathbf{Sp} \simeq \mathbf{CGrp}(\mathbf{Sp}) \simeq \mathbf{Sp}(\mathbf{CGrp}(\mathbf{An}))$, so that we may write $\mathbf{Sp}$ as the following limit:

$$\mathbf{Sp} \simeq \lim(\cdots \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An}) \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An}) \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An})).$$

We may then compute

$$\begin{aligned} \text{Hom}_{\mathbf{Sp}}(\mathbf{B}^\infty G, X) &\simeq \lim_n \text{Hom}_{\mathbf{CGrp}(\mathbf{An})}(\mathbf{B}^n G, X_n) \\ &\simeq \lim_n \text{Hom}_{\mathbf{CGrp}(\mathbf{An})}(G, \Omega^n X_n) \\ &\simeq \lim_n \text{Hom}_{\mathbf{CGrp}(\mathbf{An})}(G, X_0) \\ &\simeq \text{Hom}_{\mathbf{CGrp}(\mathbf{An})}(G, X_0). \end{aligned}$$

Indeed, the structure isomorphisms $\sigma_n : X_n \xrightarrow{\sim} \Omega X_{n+1}$ induce compatible isomorphisms $\Omega^n X_n \xrightarrow{\sim} \Omega^{n+1} X_{n+1}$, identifying the resulting diagram constantly with X_0 . Since all these equivalences are natural in G and X , this provides the adjunction. From this computation it also follows immediately that the unit $G \rightarrow \Omega^\infty \mathbf{B}^\infty G$ of the adjunction is simply the identity map of G , hence is an isomorphism, showing that the functor $\mathbf{B}^\infty : \mathbf{CGrp}(\mathbf{An}) \rightarrow \mathbf{Sp}$ is fully faithful. The essential image is the subcategory of the above limit given by

$$\mathbf{CGrp}(\mathbf{An}) \simeq \lim(\cdots \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An}_{\geq 2}) \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An}_{\geq 1}) \xrightarrow{\Omega} \mathbf{CGrp}(\mathbf{An})),$$

which are precisely those spectra X such that X_n is $(n-1)$ -connected for all $n \geq 1$, i.e. the connective spectra. $\square$

Corollary 5.4.6. *The inclusion $\mathrm{Sp}_{\geq 0} \hookrightarrow \mathrm{Sp}$ admits a right adjoint $\tau_{\geq 0}: \mathrm{Sp} \rightarrow \mathrm{Sp}_{\geq 0}$ given by $\tau_{\geq 0}(X) = \mathbf{B}^\infty \Omega^\infty X$. $\square$*

Corollary 5.4.7. *There is a fully faithful functor*

$$\mathbf{B}^\infty: \mathrm{Ab} = \mathrm{CGrp}(\mathrm{Set}) \hookrightarrow \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\mathbf{B}^\infty} \mathrm{Sp}$$

from the category of abelian groups into the ∞ -category of spectra. Here the first functor is induced by the fully faithful finite-product-preserving inclusion $\mathrm{Set} \hookrightarrow \mathrm{An}$. $\square$

Definition 5.4.8. The spectrum $\mathbf{B}^\infty A$ associated to an abelian group is called the *Eilenberg–MacLane spectrum* associated to A , and is classically denoted by HA . Because the functor $\mathbf{B}^\infty$ is fully faithful, we will frequently treat it as an inclusion and abusively denote the Eilenberg–MacLane spectrum again by A .

Remark 5.4.9. The agreement with the construction of Eilenberg–MacLane spectra from derived categories is explained in Remark 6.2.4.

5.5 Group completions as telescopes

By Corollary 5.4.4, every commutative monoid $M \in \mathrm{CMon}(\mathrm{An})$ admits a group completion

$$M \longrightarrow M^{\mathrm{grp}} = \Omega \mathbf{B} M.$$

This is a universal construction, but in examples we often want a more explicit model for the underlying anima of M^{grp} .

For an ordinary commutative monoid A generated by an element x , the group A^{grp} admits a familiar description: it is obtained by formally inverting x . Equivalently, one may take the sequential colimit

$$A \xrightarrow{x+(-)} A \xrightarrow{x+(-)} A \xrightarrow{x+(-)} \dots$$

The result is $A[x^{-1}]$, which is already a group because x generates A .

In anima the same construction is subtler. The path components behave as expected, but the higher homotopy can remember the symmetry coherences of the commutative monoid. The telescope theorem below gives a useful condition under which the same formal-inversion telescope computes the group completion.

Definition 5.5.1 (Telescope). Let $M \in \mathrm{CMon}(\mathrm{An})$ and let $x: * \rightarrow M$ be a point. We define the *telescope of M at x* as the sequential colimit

$$T(M, x) := \mathrm{colim}(M \xrightarrow{x+(-)} M \xrightarrow{x+(-)} M \xrightarrow{x+(-)} \dots) \in \mathrm{An}.$$

Since the image of x is invertible in M^{grp} , the canonical map $M \rightarrow M^{\mathrm{grp}}$ induces a comparison map

$$\gamma_{M,x}: T(M, x) \rightarrow M^{\mathrm{grp}}.$$

We use the following form of Nikolaus [Nik17, Proposition 6]. Nikolaus gives several equivalent hypotheses; the abelian fundamental-group condition used here is the version needed for complex K-theory.

Theorem 5.5.2 (Group completion theorem, telescope form). *Let $M \in \mathbf{CMon}(\mathbf{An})$ and let $x: * \rightarrow M$ be a point. Assume that:*

- (1) *The class $[x]$ generates the commutative monoid $\pi_0(M)$;*
- (2) *Each path component of $T(M, x)$ has abelian fundamental group.*

Then the comparison map

$$\gamma_{M,x}: T(M, x) \rightarrow M^{\text{grp}}$$

is an isomorphism of anima.

Proof. Under assumption (1), the singleton $\{[x]\}$ is a generating set for $\pi_0(M)$, and $T(M, x)$ is the object denoted M_∞ by Nikolaus. The claim is therefore the implication (4) $\Rightarrow$ (3) of Nikolaus [Nik17, Proposition 6]. $\square$

Remark 5.5.3. The abelian fundamental-group hypothesis is not merely cosmetic. Nikolaus explains that for the commutative monoid $\bigsqcup_{n \geq 0} \mathbf{B}\Sigma_n$, the telescope gives $\mathbb{Z} \times \mathbf{B}\Sigma_\infty$, while the group completion is obtained from this by plus construction. The full Proposition 6 in Nikolaus [Nik17] gives equivalent criteria in terms of a cyclic-permutation obstruction, and also a useful hypoabelian variant. We will only need the abelian case above.

Remark 5.5.4. Nikolaus also proves a homological group-completion theorem: after applying a multiplicative generalized homology theory, the map $M \rightarrow \Omega \mathbf{B}M$ becomes algebraic localization under suitable Ore-type hypotheses. Its proof uses localization of ring spectra and their module categories. We will not develop that result in this book; the telescope criterion above is the part needed for complex K-theory.

5.6 Finite deloopings (online only)

For completeness, we record a finite-stage version of the recognition principle. It is not needed for Theorem 5.4.5, but it explains how increasing coherence of the multiplication corresponds to the ability to deloop more often.

As an immediate corollary of the recognition principle for loop spaces we obtain a recognition principle for n -fold loop spaces.

Definition 5.6.1. Let C be an ∞ -category with finite products. For $n \geq 0$ we iteratively define the ∞ -category of n -fold monoids in C :

- For $n = 0$ we define $\text{Mon}^{(0)}(C) := C_*$;
- For $n \geq 1$ we define $\text{Mon}^{(n)}(C) := \text{Mon}(\text{Mon}^{(n-1)}(C))$.

We similarly define the ∞ -category of n -fold groups in C by $\text{Grp}^{(0)}(C) = C_*$ and $\text{Grp}^{(n)}(C) := \text{Grp}(\text{Grp}^{(n-1)}(C))$.

Remark 5.6.2. For $n = 1$ note that we get $\text{Mon}^{(1)}(C) \xrightarrow{\sim} \text{Mon}(C)$ and $\text{Grp}^{(1)}(C) \xrightarrow{\sim} \text{Grp}(C)$ by Remark 5.3.20.

Remark 5.6.3. The n -fold monoids defined above are usually called $\mathcal{E}_n$ -monoids in the literature. Indeed, if C is equipped with its cartesian symmetric monoidal structure, then $\text{Mon}^{(n)}(C)$ can be described as the ∞ -category of algebras over the little n -cubes ∞ -operad $\mathcal{E}_n$; see [Lur17, Theorem 5.1.2.2 and Proposition 2.4.2.5]. We will not use this description here.

For general n , an n -fold monoid has n different multiplication operations that all commute with each other. In light of the Eckmann–Hilton argument these operations are all identified in the homotopy category $\text{Ho}(C)$ and are homotopy commutative for $n \geq 2$. We should think of higher n as ‘more commutativity’.

Proposition 5.6.4 (Recognition principle for iterated loop spaces). *For every $n \geq 0$ there is an adjunction*

$$\mathbf{B}^n : \text{Mon}^{(n)}(\text{An}) \rightleftarrows \text{An}_* : \Omega^n$$

which restricts to an equivalence

$$\mathbf{B}^n : \text{Grp}^{(n)}(\text{An}) \xrightarrow{\simeq} \text{An}_{*, \geq n} : \Omega^n.$$

Proof. Consider the adjunction $\mathbf{B} : \text{Mon}(\text{An}) \rightleftarrows \text{An}_* : \Omega$. Since both functors preserve finite products (see Lemma 5.1.16), we may apply the monoid construction repeatedly and paste the resulting adjunctions. Using the equivalences $\text{Mon}(C_*) \simeq \text{Mon}(C)$ from Remark 5.3.20 at the intermediate stages gives the displayed adjunction $\mathbf{B}^n \dashv \Omega^n$.

The recognition principle for loop spaces, Theorem 5.2.1, says that at each stage this adjunction restricts to an equivalence between group objects and connected pointed objects. Iterating raises connectivity by one at every stage, and therefore identifies $\text{Grp}^{(n)}(\text{An})$ with $\text{An}_{*, \geq n}$. $\square$

Corollary 5.6.5. *The ∞ -category $\text{Sp}_{\geq 0}$ is equivalent to the limit of the following diagram of ∞ -categories:*

$$\cdots \rightarrow \text{Grp}^{(n)}(\text{An}) \rightarrow \cdots \rightarrow \text{Grp}^{(2)}(\text{An}) \rightarrow \text{Grp}(\text{An}) \rightarrow \text{An}_*. \quad \square$$

Proof. By definition, $\text{Sp}_{\geq 0}$ is the full subcategory of Sp spanned by those spectra (X_n) such that the pointed anima X_n is $(n-1)$ -connected for all $n \geq 1$, hence is given by the limit of the following diagram of ∞ -categories:

$$\cdots \xrightarrow{\Omega} \text{An}_{*, \geq n} \xrightarrow{\Omega} \cdots \xrightarrow{\Omega} \text{An}_{*, \geq 2} \xrightarrow{\Omega} \text{An}_{*, \geq 1} \xrightarrow{\Omega} \text{An}_*.$$

The claim thus follows from the fact that these two diagrams are equivalent to each other, in light of the following commutative squares:

$$\begin{array}{ccc} \text{An}_{*, \geq n} & \xrightarrow{\Omega} & \text{An}_{*, \geq n-1} \\ \Omega^n \downarrow \simeq & & \simeq \downarrow \Omega^{n-1} \\ \text{Grp}^{(n)}(\text{An}) & \xrightarrow{\text{fgt}} & \text{Grp}^{(n-1)}(\text{An}). \end{array} \quad \square$$

Exercises

Exercise 5.1. Let C be a classical 1-category. Show that the category $\text{Mon}(C)$ is equivalent to the usual category of monoids in C :

- (1) Given $X \in \text{Mon}(C)$, show that the maps $e : * \rightarrow X_1$ and $m : X_1 \times X_1 \rightarrow X_1$ do indeed give X_1 the structure of a monoid;
- (2) Given a monoid M in C , show that we obtain an object $X \in \text{Mon}(C)$ by setting $X_n := M^n$, and by defining for $\varphi : [m] \rightarrow [n]$ in Δ the map $\varphi^* : M^n \rightarrow M^m$ by

$$\varphi^*(x_1, \dots, x_n) := (y_1, \dots, y_m), \quad y_i := \prod_{\varphi(i-1) < j \leq \varphi(i)} x_j.$$

- (3) Show that these two constructions define the desired equivalence of categories.

Exercise 5.2. Show that the inclusion functor

$$\text{Grp}(\text{An}) \hookrightarrow \text{Mon}(\text{An})$$

admits a left adjoint

$$(-)^{\text{grp}} : \text{Mon}(\text{An}) \rightarrow \text{Grp}(\text{An})$$

which on objects is given by $M \mapsto \mathbf{OB}M$. (This is called the *group completion functor*.)

Exercise 5.3. Show that the forgetful functor $\text{Grp}(\text{An}) \rightarrow \text{An}_*$ admits a left adjoint

$$F^{\text{Grp}}(-) : \text{An}_* \rightarrow \text{Grp}(\text{An})$$

which is given on objects by $X \mapsto \mathbf{O}\Sigma X$. (This is called the *free group functor*.)

Exercise 5.4. Let x be an object of an ∞ -category C . Denote by $\text{Aut}_C(x)$ the connected component of $\text{End}_C(x) := \text{Hom}_C(x, x)$ spanned by those endomorphisms $x \rightarrow x$ that are invertible. Show that composition of morphisms in C can be enhanced to a group structure on $\text{Aut}_C(x)$.

Remark. One can also show that $\text{End}_C(x)$ itself has the structure of a monoid via composition of morphisms, but this requires some theory we haven't introduced.

Exercise 5.5. The goal of this exercise is to show that the forgetful functor $\text{Mon}(\text{An}) \rightarrow \text{An}$ admits a left adjoint

$$F^{\text{Mon}}(-) : \text{An} \rightarrow \text{Mon}(\text{An})$$

given by the ‘anima of words in X of arbitrary length’:

$$F^{\text{Mon}}(X)_1 \simeq \bigsqcup_{n \geq 0} X^n.$$

- (1) Let $\Delta_{\text{int}} \subseteq \Delta$ denote the subcategory which has all objects, but which has only those morphisms $[n] \rightarrow [m]$ that are inclusions of an interval $\{i \leq i+1 \leq \dots \leq j-1 \leq j\}$ into $[m]$ for some $0 \leq i \leq j \leq m$ (these are called the *inert* morphisms). Show that evaluation at $[1] \in \Delta_{\text{int}}$ defines an equivalence

$$\text{Fun}^{\text{Segal}}(\Delta_{\text{int}}^{\text{op}}, \text{An}) \xrightarrow{\sim} \text{An},$$

where the left-hand side denotes those functors $X: \Delta_{\text{int}}^{\text{op}} \rightarrow \mathbf{An}$ satisfying the *Segal condition*, i.e. the map

$$(e_i^*)_{i=1}^n: X_n \rightarrow X_1^n$$

is an isomorphism for all $n \geq 0$.

Hint: the inverse is given by right Kan extension.

- (2) Show that the left Kan extension functor

$$i_! : \text{Fun}(\Delta_{\text{int}}^{\text{op}}, \mathbf{An}) \rightarrow \text{Fun}(\Delta^{\text{op}}, \mathbf{An})$$

preserves Segal objects.

- (3) Show that the resulting composite

$$F^{\text{Mon}}(-): \mathbf{An} \simeq \text{Fun}^{\text{Segal}}(\Delta_{\text{int}}^{\text{op}}, \mathbf{An}) \xrightarrow{i_!} \text{Fun}^{\text{Segal}}(\Delta^{\text{op}}, \mathbf{An}) = \mathbf{Mon}(\mathbf{An})$$

is left adjoint to the forgetful functor, and use the pointwise formula for Kan extensions to show that the formula

$$F^{\text{Mon}}(X)_1 \simeq \bigsqcup_{n \geq 0} X^n$$

is satisfied.

Exercise 5.6. Assume that C is a classical 1-category with finite products.

- (1) Show that the category $\mathbf{CMon}(C)$ is equivalent to the category of commutative monoids in C defined in the usual way: an object M of C equipped with maps $+: M \times M \rightarrow M$ and $0: * \rightarrow M$ such that the following three diagrams commute:

$$\begin{array}{ccc} M \times * & \xrightarrow{\text{id}_M \times 0} & M \times M & \xleftarrow{0 \times \text{id}_M} & M \\ & \searrow \simeq & \downarrow + & & \swarrow \simeq \\ & & M, & & \end{array} \quad \begin{array}{ccc} M \times M \times M & \xrightarrow{\text{id}_M \times +} & M \times M \\ \downarrow + \times \text{id}_M & & \downarrow + \\ M \times M & \xrightarrow{+} & M, \end{array} \quad \begin{array}{ccc} M \times M & \xrightarrow[\simeq]{\tau} & M \times M \\ & \searrow + & \swarrow + \\ & & M. \end{array}$$

- (2) Combining part (1) with Exercise 5.1, deduce that in this case the forgetful functor $U: \mathbf{CMon}(C) \hookrightarrow \mathbf{Mon}(C)$ is fully faithful.

Exercise 5.7. For a finite set S , the inclusion $\mathbf{Fin} \hookrightarrow \mathbf{Span}(\mathbf{Fin})$ of the subcategory containing all objects but only the forward-pointing maps induces a functor on slice categories of the form

$$\mathbf{Fin}_{/S} \rightarrow \mathbf{Span}(\mathbf{Fin})_{/S}.$$

Show that this functor admits a left adjoint, which is given on objects by sending a span

$$T \xleftarrow{f} U \xrightarrow{g} S$$

to the map $g: U \rightarrow S$.

Exercise 5.8. Let $M \in \mathbf{Mon}(\mathbf{An})$ be a monoid in *animae*. Show that M is a group in $\mathbf{An}$ if and only if its monoid of path components $\pi_0(M)$ is a group in the ordinary sense.

Conclude that the commutative monoid $\mathbf{BM}$ is a commutative group for every commutative monoid M .

Exercise 5.9. Show that for two ∞ -categories C and D , the projection maps $C \times D \rightarrow C$ and $C \times D \rightarrow D$ induce an equivalence

$$|C \times D| \xrightarrow{\sim} |C| \times |D|$$

on geometric realizations. (Recall that $|C|$ is obtained from C by inverting all morphisms: it is the localization $C[W^{-1}]$ with $W = \text{Map}([1], C)$.)

6 Homological algebra

Homological algebra is the study of chain complexes and their derived categories. While its origins lie in classical algebra and algebraic topology, it is deeply intertwined with stable homotopy theory: the derived category of an abelian category is a stable ∞ -category, and many of its key structures, like exact sequences and Ext-groups, are instances of general phenomena in stable ∞ -categories.

In this chapter, we develop some of the core aspects of homological algebra from the ∞ -categorical perspective. We begin in Section 6.1 by constructing the derived ∞ -category $D(\mathcal{A})$ of an abelian category $\mathcal{A}$ as a localization of the category of chain complexes at quasi-isomorphisms. Using the general framework from Section 2.2, we show that $D(\mathcal{A})$ is a stable ∞ -category. We then specialize to $D(\mathbb{Z})$ in Section 6.2, where Eilenberg–MacLane spectra provide the intrinsic construction of ordinary homology and cohomology promised in Chapter 3.

In Section 6.3, we introduce *t-structures*, an axiomatic framework that captures the interplay between stable homotopy theory and classical algebra. A t-structure comes with classes of *connective* and *coconnective* objects, which behave like connective and coconnective spectra. Their intersection turns out to be a 1-category, and in fact an abelian category, called the *heart* of the t-structure. The derived category $D(\mathcal{A})$ carries a canonical t-structure whose heart recovers the original abelian category $\mathcal{A}$.

The remaining sections treat projective and flat objects, Ext- and Tor-groups, and derived functors. In Section 6.4, we define Ext-groups as homotopy groups of mapping spectra and show that they can be computed by projective resolutions. In Section 6.5, we develop the parallel theory of flat objects and Tor-groups. In Section 6.6, we define derived functors, compute them by projective replacement, and construct the derived tensor product and derived internal hom. Finally, Section 6.7 proves the Künneth formula and the universal coefficient theorems, first algebraically for complexes over a principal ideal domain and then for the ordinary homology and cohomology of anima.

6.1 Derived ∞ -categories

One of the foundational definitions of homological algebra is that of a *chain complex*: a sequence $\{C_n\}_{n \in \mathbb{Z}}$ of abelian groups, equipped with boundary maps $d_n: C_n \rightarrow C_{n-1}$ satisfying $d_{n-1} \circ d_n = 0$. For every chain complex $C_\bullet$, we may define its *n-th homology group* as the quotient of the kernel of $d_n: C_n \rightarrow C_{n-1}$ by the image of $d_{n+1}: C_{n+1} \rightarrow C_n$. One of the main examples of chain complexes in algebraic topology is the ‘singular chain complex’ $C_\bullet^{\text{sing}}(X; \mathbb{Z})$ of a topological space X , whose homology groups are precisely the ordinary homology groups of X studied axiomatically in Section 3.3 (cf. Remark 3.3.2).

While chain complexes define a 1-category $\text{Ch}(\mathbb{Z})$, this category is in some sense ‘too rigid’ from the perspective of homotopy theory. For example, while the homology groups $H_*(X; \mathbb{Z})$ of a topological space are homotopy invariants, the singular chain complex $C_\bullet^{\text{sing}}(X; \mathbb{Z})$ itself is not homotopy invariant: the singular chain complexes of two homotopy equivalent (but non-homeomorphic) spaces are hardly ever isomorphic to each other. To resolve this issue, we may pass to the derived category $D(\mathbb{Z})$, which is obtained from $\text{Ch}(\mathbb{Z})$ by inverting all *quasi-isomorphisms*: those morphisms of chain complexes that induce isomorphisms on homology groups.

In classical treatments, $D(\mathbb{Z})$ is constructed as a triangulated category: a 1-category equipped with additional structure encoding exact sequences. However, this structure is somewhat unwieldy, and many natural constructions (such as limits and colimits, or functor categories) are difficult to perform in the triangulated setting. We will instead define $D(\mathcal{A})$ as the ∞ -categorical localization

$$D(\mathcal{A}) := \text{Ch}(\mathcal{A})[\{\text{quasi-isomorphisms}\}^{-1}]$$

in the sense of Definition 1.5.19. The resulting ∞ -category is stable, and the triangulated structure studied classically emerges as a shadow of the exact sequences coming from stability.

We now develop this theory systematically. After recalling the necessary background on abelian categories and their chain complexes in Section 6.1.1, we construct the derived ∞ -category $D(\mathcal{A})$ of an abelian category $\mathcal{A}$ in Section 6.1.2 and establish its stability in Section 6.1.3.

6.1.1 Chain complexes in abelian categories

Notation 6.1.1. Let $\mathcal{A}$ be an additive 1-category. We refer to fibers and cofibers in $\mathcal{A}$ as *kernels* and *cokernels*, respectively:

$$\begin{array}{ccc} \ker(f) & \hookrightarrow & A \\ \downarrow & \lrcorner & \downarrow f \\ 0 & \hookrightarrow & B \end{array} \quad \text{and} \quad \begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow \\ 0 & \twoheadrightarrow & \text{coker}(f) \end{array}$$

If f is a monomorphism, we will use B/A as alternative notation for the cokernel.

Note that the map $\ker(f) \rightarrow A$ is automatically a monomorphism, while $B \twoheadrightarrow \text{coker}(f)$ is automatically an epimorphism.

Definition 6.1.2. A 1-category $\mathcal{A}$ is called an *abelian category* if it is additive, has kernels and cokernels, and a nullsequence $A \xrightarrow{i} B \xrightarrow{p} C$ with i a monomorphism and p an epimorphism is a fiber sequence (i.e. $A \xrightarrow{\sim} \ker(p)$) if and only if it is a cofiber sequence (i.e. $\text{coker}(i) \xrightarrow{\sim} C$). We will display such nullsequences as

$$0 \rightarrow A \xrightarrow{i} B \xrightarrow{p} C \rightarrow 0,$$

and refer to them as *short exact sequences*.

Example 6.1.3. The main example of an abelian category is the category Ab of abelian groups. More generally the category $\text{LMod}_R(\text{Ab})$ of left modules over an associative ring R is abelian.

Example 6.1.4. The opposite $\mathcal{A}^{\text{op}}$ of an abelian category is again abelian.

The following are the basic properties of abelian categories we will need:

Proposition 6.1.5. *Let $\mathcal{A}$ be an abelian category.*

- (1) *Every monomorphism $i: A \hookrightarrow B$ is the kernel of its cokernel, and every epimorphism $p: B \twoheadrightarrow C$ is the cokernel of its kernel.*
- (2) *A morphism f is an isomorphism if and only if it is both a monomorphism and an epimorphism.*
- (3) *The category $\mathcal{A}$ admits all pullbacks and pushouts.*
- (4) *Monomorphisms are closed under pushouts, and any pushout square along a monomorphism is also a pullback square.*
- (5) *Epimorphisms are closed under pullbacks and any pullback square along an epimorphism is also a pushout square.*
- (6) *Every morphism $f: A \rightarrow B$ factors uniquely¹³ as an epimorphism followed by a monomorphism.*

The factorization from (6) is denoted by $A \twoheadrightarrow \operatorname{im}(f) \hookrightarrow B$, and we refer to the object $\operatorname{im}(f)$ as the *image* of f .

Proof. These are the standard exactness properties of an abelian category; see [Wei94, Section 1.2]. We only recall that the image factorization is obtained by setting

$$\operatorname{im}(f) := \ker(B \twoheadrightarrow \operatorname{coker}(f)),$$

and that a pushout of $f: A \rightarrow B$ along $g: A \rightarrow C$ is computed as the cokernel of $(f, -g): A \rightarrow B \oplus C$. These descriptions give the remaining assertions and their duals. $\square$

Definition 6.1.6. A nullsequence $A \xrightarrow{f} B \xrightarrow{g} C$ in $\mathcal{A}$ is called *exact* if the map $\operatorname{im}(f) \rightarrow \ker(g)$ is an isomorphism.

Definition 6.1.7. A *chain complex* in $\mathcal{A}$ is a pair $(C_\bullet, d_\bullet)$ consisting of a $\mathbb{Z}$ -graded object $C_\bullet = (C_n)_{n \in \mathbb{Z}}$, $C_n \in \mathcal{A}$, and a collection of ‘boundary maps’ $d_\bullet = (d_n: C_n \rightarrow C_{n-1})_{n \in \mathbb{Z}}$ satisfying the property that $d_{n-1} \circ d_n = 0$ for all n . We will often denote the chain complex simply by $C_\bullet$ or C .

A *chain map* (or *morphism of chain complexes*) $f_\bullet: C_\bullet \rightarrow D_\bullet$ consists of a collection of morphisms $f_n: C_n \rightarrow D_n$ that commute with the boundary maps, in the sense that $d_n^D \circ f_n = f_{n-1} \circ d_n^C$ for all n . There is a clear way to compose these, resulting in a 1-category $\operatorname{Ch}(\mathcal{A})$ of chain complexes in $\mathcal{A}$.

Remark 6.1.8 (Cohomological grading of complexes). We grade chain complexes *homologically*, so that the differential $d_n: C_n \rightarrow C_{n-1}$ *lowers* degree. Some references grade *cohomologically* instead, writing a complex as $(C^n)_{n \in \mathbb{Z}}$ with differential $d^n: C^n \rightarrow C^{n+1}$ *raising* degree. The two conventions record the same data, related by negating the index: a homologically graded complex $(C_\bullet, d_\bullet)$ becomes the cohomologically graded complex with $C^n := C_{-n}$ and $d^n := d_{-n}$, and conversely. We use homological grading throughout.

¹³This is classically formulated as ‘unique up to unique isomorphism’.

Example 6.1.9. Given an object $A \in \mathcal{A}$ and $k \in \mathbb{Z}$, we let $A[k]$ denote the chain complex consisting of A concentrated in degree k , meaning that $A[k]_k := A$ while $A[k]_n = 0$ for all $n \neq k$; all boundary maps of $A[k]$ are necessarily zero. This construction defines a fully faithful functor $(-)[k]: \mathcal{A} \hookrightarrow \text{Ch}(\mathcal{A})$ for all k .

Example 6.1.10. Let $C_\bullet$ be a chain complex. Given $k \in \mathbb{Z}$, we define the *shift* $C[k]_\bullet$ of C by $C[k]_n := C_{n-k}$ and $d_n^{C[k]} := (-1)^k d_{n-k}^C$.

Remark 6.1.11. The sign $(-1)^k$ in the differential of $C[k]$ is a deliberate choice. It is slightly unusual: many references define the shift without any sign, setting $d_n^{C[k]} := d_{n-k}^C$. Including the sign has the advantage that the shift agrees on the nose with the suspension of the derived ∞ -category (see Theorem 6.1.29) rather than merely up to a natural isomorphism, and it is the convention that fits best with the shift functors of stable homotopy theory. As the two conventions produce isomorphic chain complexes, the choice affects none of the homological invariants.

Definition 6.1.12. Given a chain complex $C_\bullet$ and an integer n , the *n -th homology* of $C_\bullet$ is the object

$$H_n(C) := \ker(d_n) / \text{im}(d_{n+1}) \in \mathcal{A}.$$

Any chain map $f_\bullet: C_\bullet \rightarrow D_\bullet$ induces a map $H_n(f): H_n(C) \rightarrow H_n(D)$ on n -th homology, resulting in a functor $H_n(-): \text{Ch}(\mathcal{A}) \rightarrow \mathcal{A}$.

Definition 6.1.13. A chain map $f_\bullet$ is called a *quasi-isomorphism* if the induced map $H_n(f): H_n(C) \rightarrow H_n(D)$ is an isomorphism for all n .

Given two chain maps $f, g: C_\bullet \rightarrow D_\bullet$, a *chain homotopy* between f and g is a family of maps $(H_n: C_n \rightarrow D_{n+1})_{n \in \mathbb{Z}}$ satisfying $f_n - g_n = \partial^D \circ H_n + H_{n-1} \circ \partial^C$. We say f is a *chain homotopy equivalence* if there exists a chain map $g: D_\bullet \rightarrow C_\bullet$ and chain homotopies $g \circ f \sim \text{id}_{C_\bullet}$ and $f \circ g \sim \text{id}_{D_\bullet}$.

Definition 6.1.14 (Mapping cone). The *mapping cone* of a chain map $f: C_\bullet \rightarrow D_\bullet$ is the chain complex $\text{Cone}(f)$ with

$$\text{Cone}(f)_n := D_n \oplus C_{n-1}, \quad d^{\text{Cone}(f)}(y, x) := (d^D(y) + f(x), -d^C(x)).$$

It fits into a natural short exact sequence $0 \rightarrow D_\bullet \rightarrow \text{Cone}(f) \rightarrow C_\bullet[1] \rightarrow 0$.

Exercise 6.1.15. Show that every chain homotopy equivalence is a quasi-isomorphism.

An important fact from homological algebra we will use is that short exact sequences of chain complexes result in long exact sequences of homology groups. We will state it as a black box; a proof may be found in [Wei94, Theorem 1.3.1].

Proposition 6.1.16. Let $0 \rightarrow C_\bullet \xrightarrow{f} D_\bullet \xrightarrow{g} E_\bullet \rightarrow 0$ be a short exact sequence in $\text{Ch}(\mathcal{A})$. Then there exist maps $\partial_n: H_n(E) \rightarrow H_{n-1}(C)$ for all n , and the sequence

$$\dots \xrightarrow{g} H_{n+1}(E) \xrightarrow{\partial_{n+1}} H_n(C) \xrightarrow{f} H_n(D) \xrightarrow{g} H_n(E) \xrightarrow{\partial_n} H_{n-1}(C) \xrightarrow{f} \dots$$

in $\mathcal{A}$ is exact. □

When $\mathcal{A}$ is modules over a ring, the boundary map has the familiar elementwise description. Represent a class in $H_n(E)$ by a cycle $x \in E_n$, lift it to $y \in D_n$, and write $d(y) = f(z)$ for some $z \in C_{n-1}$. Then z is a cycle and $\partial_n([x]) = [z]$. The standard diagram chase shows that this is independent of all choices and proves exactness; see again [Wei94, Theorem 1.3.1].

Corollary 6.1.17. *Given a short exact sequence $0 \rightarrow C_\bullet \xrightarrow{f} D_\bullet \xrightarrow{g} E_\bullet \rightarrow 0$, the map f is a quasi-isomorphism if and only if $E_\bullet$ has trivial homology.* $\square$

6.1.2 Derived ∞ -categories

Fix an abelian category $\mathcal{A}$ for the remainder of this subsection.

Definition 6.1.18. We define the *derived ∞ -category of $\mathcal{A}$* as the (∞ -categorical) localization of the 1-category $\text{Ch}(\mathcal{A})$ of chain complexes at the class of quasi-isomorphisms:

$$D(\mathcal{A}) := \text{Ch}(\mathcal{A})[\{\text{quasi-isomorphisms}\}^{-1}].$$

We will sometimes refer to objects of $D(\mathcal{A})$ as *complexes*. This is slightly abusive terminology: objects of $D(\mathcal{A})$ should best be thought of as ‘chain complexes up to quasi-isomorphism’ and in particular do not determine a well-defined chain complex in $\mathcal{A}$.

Example 6.1.19. Let R be an associative ring, and consider the abelian category $\mathcal{A} = \text{RMod}_R(\text{Ab})$ of right R -modules in the category of abelian groups. We denote the resulting derived ∞ -category by $D(R) := D(\text{RMod}_R(\text{Ab}))$. In particular $D(\mathbb{Z}) = D(\text{Ab})$.

Recall from Example 6.1.9 that $\text{Ch}(\mathcal{A})$ contains $\mathcal{A}$ as a full subcategory, given by those chain complexes concentrated in degree 0. We will now show that the resulting functor $\mathcal{A} \rightarrow D(\mathcal{A})$, $A \mapsto A[0]$ is still fully faithful. Note that this is not immediately clear, since the localization functor $\text{Ch}(\mathcal{A}) \rightarrow D(\mathcal{A})$ is far from fully faithful: for objects $A, B \in \mathcal{A}$ there are no nonzero chain maps $A[0] \rightarrow B[n]$ when $n > 0$ (the nonzero terms sit in different degrees), whereas the corresponding morphisms in $D(\mathcal{A})$ compute the Ext-groups $\text{Ext}_{\mathcal{A}}^n(A, B)$ (see Section 6.4), which are frequently nonzero.

We will proceed by showing that complexes can be ‘truncated’ both from below and from above: given a complex C , there is a universal approximation of C from the left by a complex concentrated in non-negative degrees, and from the right by a complex concentrated in non-positive degrees. By truncating from both directions, we can force the complex to be concentrated in degree 0. These truncation operations will descend to the level of derived ∞ -categories, and will induce an equivalence between $\mathcal{A}$ and the full subcategory of $D(\mathcal{A})$ on those complexes whose homology is concentrated in degree zero.

Notation 6.1.20. Given $k \in \mathbb{Z}$, we denote by $\text{Ch}(\mathcal{A})_{\geq k} \subseteq \text{Ch}(\mathcal{A})$ the full subcategory spanned by those chain complexes $C_\bullet$ satisfying $C_n = 0$ for $n < k$. Similarly, we write $\text{Ch}(\mathcal{A})_{\leq k}$ for the full subcategory on those chain complexes satisfying $C_n = 0$ for $n > k$. We denote by

$$D(\mathcal{A})_{\geq k} := \text{Ch}(\mathcal{A})_{\geq k}[\{\text{quasi-isos}\}^{-1}], \quad D(\mathcal{A})_{\leq k} := \text{Ch}(\mathcal{A})_{\leq k}[\{\text{quasi-isos}\}^{-1}]$$

the localizations of these subcategories at the quasi-isomorphisms.

Proposition 6.1.21. *For an integer k , the following hold:*

- (1) *The inclusion $\text{Ch}(\mathcal{A})_{\geq k} \hookrightarrow \text{Ch}(\mathcal{A})$ admits a right adjoint $\tau_{\geq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})_{\geq k}$.*
- (2) *The induced functor $\text{D}(\mathcal{A})_{\geq k} \rightarrow \text{D}(\mathcal{A})$ is fully faithful and admits a right adjoint $\tau_{\geq k}: \text{D}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})_{\geq k}$.*
- (3) *Dually, the inclusion $\text{Ch}(\mathcal{A})_{\leq k} \hookrightarrow \text{Ch}(\mathcal{A})$ admits a left adjoint $\tau_{\leq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})_{\leq k}$.*
- (4) *The induced functor $\text{D}(\mathcal{A})_{\leq k} \rightarrow \text{D}(\mathcal{A})$ is fully faithful and admits a left adjoint $\tau_{\leq k}: \text{D}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})_{\leq k}$.*

Proof. (1) Given a chain complex C and an integer $k \in \mathbb{Z}$, we construct a new chain complex $\tau_{\geq k}(C)$ as follows:

$$\tau_{\geq k}(C)_n := \begin{cases} C_n & n > k \\ \ker(d_k: C_k \rightarrow C_{k-1}) & n = k \\ 0 & n < k. \end{cases}$$

The structure maps $d_i: \tau_{\geq k}(C)_n \rightarrow \tau_{\geq k}(C)_{n-1}$ are the ones from $C_\bullet$ when $n > k$, are zero when $n < k$, and for $n = k$ is the canonical map $C_{k+1} \rightarrow \ker(d_k)$ induced by $d_{k+1}: C_{k+1} \rightarrow C_k$. This construction is functorial in C , defining a functor

$$\tau_{\geq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})_{\geq k}.$$

There is a canonical chain map $\varepsilon: \tau_{\geq k}(C) \rightarrow C$ given by the identity in degree $n > k$, by the zero map in degree $n < k$, and in degree $n = k$ by the inclusion $\ker(d_k) \hookrightarrow C_k$. It is clear that this chain map is a monomorphism. Furthermore, if D is a chain complex such that D_n vanishes for $n < k$, any chain map $f: D \rightarrow C$ uniquely factors through $\tau_{\geq k}(C)$: the commutative square

$$\begin{array}{ccc} D_k & \xrightarrow{f_k} & C_k \\ d_k \downarrow & & \downarrow d_k \\ 0 & \xrightarrow{f_{k-1}} & C_{k-1} \end{array}$$

guarantees that f_k factors through $\ker(d_k) \hookrightarrow C_k$. This shows that the transformation $\varepsilon: \tau_{\geq k} \rightarrow \text{id}$ exhibits the functor $\tau_{\geq k}: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})_{\geq k}$ as right adjoint to the inclusion.

(2) Observe that the counit $\varepsilon: \tau_{\geq k}(C) \rightarrow C$ of the adjunction from part (1) induces isomorphisms $H_n(\tau_{\geq k}(C)) \xrightarrow{\cong} H_n(C)$ for $n \geq k$. Since we have $H_n(\tau_{\geq k}(C)) = 0$ for $n < k$, it follows that the functor $\tau_{\geq k}$ sends quasi-isomorphisms to quasi-isomorphisms, hence induces a functor

$$\tau_{\geq k}: \text{D}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})_{\geq k}.$$

In a similar way, the unit and counit of the adjunction from part (1) induce natural transformations at the level of derived ∞ -categories, exhibiting $\tau_{\geq k}$ as a right adjoint of the functor $\text{D}(\mathcal{A})_{\geq k} \rightarrow \text{D}(\mathcal{A})$ induced by the inclusion. Since the unit of the adjunction is a natural isomorphism, it follows that this inclusion is fully faithful, showing (2).

Parts (3) and (4) are similar, and can be seen as instances of (1) and (2) applied to $\mathcal{A}^{\text{op}}$. The chain complex $\tau_{\leq k}(C)_\bullet$ may be explicitly given as follows:

$$\tau_{\leq k}(C)_n := \begin{cases} 0 & n > k \\ \text{coker}(d_{k+1}: C_{k+1} \rightarrow C_k) & n = k \\ C_n & n < k. \end{cases} \quad \square$$

Definition 6.1.22. Fix an integer $k \in \mathbb{Z}$. A complex A is called *k-connective* if $H_n(A) = 0$ for $n < k$. It is called *k-coconnective* if $H_n(A) = 0$ for $n > k$. When $k = 0$, we simply speak of *connective* and *coconnective* complexes.

Corollary 6.1.23. *The image of the inclusion $D(\mathcal{A})_{\geq k} \hookrightarrow D(\mathcal{A})$ from the previous proposition consists of the k-connective complexes, while the image of $D(\mathcal{A})_{\leq k} \hookrightarrow D(\mathcal{A})$ consists of the k-coconnective complexes.*

Proof. The essential image of the first inclusion consists of those complexes C for which the counit map $\tau_{\geq k}(C) \rightarrow C$ of the adjunction is an isomorphism in $D(\mathcal{A})$, i.e. induces isomorphisms on homology. This happens precisely if the homology of C is trivial below degree k , i.e. if C is k -connective. The argument for the second inclusion is dual. $\square$

Corollary 6.1.24. *Given a k-connective complex $A \in D(\mathcal{A})_{\geq k}$ and a (k-1)-coconnective complex $B \in D(\mathcal{A})_{\leq k-1}$, we have*

$$\text{Hom}_{D(\mathcal{A})}(A, B) = 0.$$

Proof. By adjunction we have $\text{Hom}_{D(\mathcal{A})}(A, B) \simeq \text{Hom}_{D(\mathcal{A})}(A, \tau_{\geq k} B) = 0$, where we use that $\tau_{\geq k} B = 0$ as its homology is trivial. $\square$

We are now finally in a position to deduce the full faithfulness of $\mathcal{A} \hookrightarrow D(\mathcal{A})$.

Proposition 6.1.25. *For every $k \in \mathbb{Z}$, the composite*

$$\mathcal{A} \xrightarrow{A \mapsto A[k]} \text{Ch}(\mathcal{A}) \rightarrow D(\mathcal{A})$$

is fully faithful, with essential image those complexes whose homology is concentrated in degree k.

Proof. Let $\text{Ch}(\mathcal{A})_{=k} \subseteq \text{Ch}(\mathcal{A})$ denote the subcategory of chain complexes $C_\bullet$ satisfying $C_n = 0$ for $n \neq k$. Then the assignment $A \mapsto A[k]$ induces an equivalence $\mathcal{A} \xrightarrow{\sim} \text{Ch}(\mathcal{A})_{=k}$. The adjunction $\text{Ch}(\mathcal{A})_{\geq k} \rightleftarrows \text{Ch}(\mathcal{A})$ restricts to an adjunction $\mathcal{A} \simeq \text{Ch}(\mathcal{A})_{=k} \rightleftarrows \text{Ch}(\mathcal{A})_{\leq k}$, and since both functors preserve quasi-isomorphisms, we get an induced adjunction

$$\mathcal{A} \rightleftarrows D(\mathcal{A})_{\leq k},$$

where the left adjoint is fully faithful as the unit is a natural isomorphism. The functor in question is now given by the composite

$$\mathcal{A} \hookrightarrow D(\mathcal{A})_{\leq k} \hookrightarrow D(\mathcal{A}),$$

hence is fully faithful as well. $\square$

6.1.3 Stability of derived ∞ -categories

We will now show that $D(\mathcal{A})$ is a stable ∞ -category, and that it has small (co)limits whenever $\mathcal{A}$ has small (co)products that interact nicely with the abelian structure. The main input will be to show that short exact sequences of chain complex in $\mathcal{A}$ result in exact sequences in $D(\mathcal{A})$. To get a handle on fibers and cofibers in $D(\mathcal{A})$, we will equip $\text{Ch}(\mathcal{A})$ with the structure of an ‘ ∞ -category with weak equivalences and (co)fibrations’, as introduced in Definition 2.2.4, and then apply the results from Section 2.2.

Proposition 6.1.26. *Let $\mathcal{A}$ be an abelian category. Then the triple $(\text{Ch}(\mathcal{A}), \text{quasi-isos}, \text{monos})$ is an ∞ -category with weak equivalences and cofibrations. It admits functorial factorizations, and is homotopy cocomplete whenever $\mathcal{A}$ satisfies (AB3) and (AB4), i.e. $\mathcal{A}$ has small coproducts and monomorphisms are closed under small coproducts.*

Proof. We verify the (dualized) axioms from Definition 2.2.4:

- (1) The 2-out-of-3 property for quasi-isomorphisms follows immediately from the 2-out-of-3 property for isomorphisms in $\mathcal{A}$.
- (2) All isomorphisms are monomorphisms, and since $\text{Ch}(\mathcal{A})$ is again an abelian category the monomorphisms are closed under compositions and pushouts; see Proposition 6.1.5. Note that every chain complex is cofibrant, since the map $0 \rightarrow C$ is a monomorphism for all $C \in \text{Ch}(\mathcal{A})$.
- (3) Consider a pushout square in $\text{Ch}(\mathcal{A})$ of the form

$$\begin{array}{ccc} C_{\bullet} & \xrightarrow{f} & C'_{\bullet} \\ i \downarrow & \ulcorner & \downarrow i' \\ D_{\bullet} & \xrightarrow{f'} & D'_{\bullet} \end{array}$$

where i and i' are monomorphisms and i is a quasi-isomorphism. We need to show that also i' is a quasi-isomorphism. Let $E_{\bullet}$ be the cokernel of i and let $E'_{\bullet}$ be the cokernel of i' . Since i is a quasi-isomorphism, it follows from Corollary 6.1.17 that the homology of $E_{\bullet}$ is trivial. By the pasting law for pushout squares, we see that the induced map $E_{\bullet} \rightarrow E'_{\bullet}$ is an isomorphism of chain complexes, and so also the homology of $E'_{\bullet}$ is trivial. By applying Corollary 6.1.17 in the other direction, it follows that i' is a quasi-isomorphism.

- (4) We now show that any chain map $f: C_{\bullet} \rightarrow D_{\bullet}$ factors functorially as a monomorphism followed by a quasi-isomorphism. Define the *mapping cylinder* $\text{Cyl}(f)$ of f as the chain complex defined by $(\text{Cyl}(f))_n = C_n \oplus C_{n-1} \oplus D_n$. The differential $d^{\text{Cyl}(f)}: (\text{Cyl}(f))_n \rightarrow (\text{Cyl}(f))_{n-1}$ is given by

$$d^{\text{Cyl}(f)} := \begin{pmatrix} d^C & \text{id}_C & 0 \\ 0 & -d^C & 0 \\ 0 & -f & d^D \end{pmatrix},$$

i.e. in terms of elements we have $d^{\text{Cyl}(f)}(c_n, c_{n-1}, d_n) := (d^C c_n + c_{n-1}, -d^C c_{n-1}, -f(c_{n-1}) + d^D d_n)$. There is a chain map $i_C: C_{\bullet} \rightarrow \text{Cyl}(f)_{\bullet}$ given by $i_C(c_n) = (c_n, 0, 0)$, and a chain map $p: \text{Cyl}(f)_{\bullet} \rightarrow D_{\bullet}$ given by $p(c_n, c_{n-1}, d_n) = f(c_n) + d_n$. Note that the constructions of $\text{Cyl}(f)$, i_C and p are functorial in f and that $f = p \circ i_C$, so this provides a functorial factorization of f . Since i_C is clearly a monomorphism, it remains to show that p is a quasi-isomorphism. By Exercise 6.1.15,

it suffices to show that p is in fact a chain homotopy equivalence. We claim that the chain map $i_D: D_\bullet \rightarrow \text{Cyl}(f)_\bullet$ given by $i_D(d_n) = (0, 0, d_n)$ is a chain homotopy inverse for p . It is clear that $p \circ i_D = \text{id}_{D_\bullet}$, so it remains to construct a chain homotopy H between $i_D \circ p$ and $\text{id}_{\text{Cyl}(f)_\bullet}$. Such a chain homotopy is given by $H_n(c_n, c_{n-1}, d_n) := (0, c_n, 0)$.

We have thus shown that the triple $(\text{Ch}(\mathcal{A}), \text{quasi-isos}, \text{monos})$ is an ∞ -category with weak equivalences and cofibrations, and that it has functorial factorizations. Under the assumption that $\mathcal{A}$ satisfies (AB3) and (AB4), the category $\text{Ch}(\mathcal{A})$ inherits arbitrary coproducts from $\mathcal{A}$, formed degreewise, and monomorphisms in $\text{Ch}(\mathcal{A})$ are again closed under arbitrary coproducts. Moreover, since the coproduct functor $\bigoplus_I: \text{Ch}(\mathcal{A})^I \rightarrow \text{Ch}(\mathcal{A})$ is exact, it commutes with formation of homology, and it follows that an arbitrary coproduct of quasi-isomorphisms is again a quasi-isomorphism. This verifies the (dualized) conditions from Definition 2.2.10, showing that $\text{Ch}(\mathcal{A})$ is homotopy cocomplete. $\square$

Corollary 6.1.27. *Let $\mathcal{A}$ be an abelian category. The triple $(\text{Ch}(\mathcal{A}), \text{quasi-isos}, \text{epis})$ is an ∞ -category with weak equivalences and fibrations. It admits functorial factorizations, and is homotopy complete whenever $\mathcal{A}$ satisfies (AB3*) and (AB4*), i.e. $\mathcal{A}$ has small products and epimorphisms are closed under small products.*

Proof. This is a special case of Proposition 6.1.26 applied to the abelian category $\mathcal{A}^{\text{op}}$, using that $\text{Ch}(\mathcal{A}^{\text{op}}) \simeq \text{Ch}(\mathcal{A})^{\text{op}}$. $\square$

Corollary 6.1.28. *Let $\mathcal{A}$ be an abelian category.*

- (1) *The derived ∞ -category $\text{D}(\mathcal{A})$ admits finite limits and finite colimits.*
- (2) *If $\mathcal{A}$ satisfies (AB3) and (AB4), then $\text{D}(\mathcal{A})$ admits small colimits. If it satisfies (AB3*) and (AB4*) then $\text{D}(\mathcal{A})$ admits small limits.*
- (3) *The localization functor $\gamma: \text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$ preserves pushouts along monomorphisms and preserves pullbacks along epimorphisms.*
- (4) *For a short exact sequence $0 \rightarrow C_\bullet \hookrightarrow D_\bullet \twoheadrightarrow E_\bullet \rightarrow 0$ in $\text{Ch}(\mathcal{A})$, the induced sequence*

$$\gamma(C_\bullet) \rightarrow \gamma(D_\bullet) \rightarrow \gamma(E_\bullet)$$

in $\text{D}(\mathcal{A})$ is both a fiber sequence and a cofiber sequence.

Proof. Parts (1)-(3) follow directly from applying Theorem 2.2.8, Theorem 2.2.11, and Corollary 2.2.9 to the structures established in Proposition 6.1.26 and Proposition 6.1.27. Part (4) is a specific case of (3). $\square$

We are now ready to prove the main result of this section:

Theorem 6.1.29. *For any abelian category $\mathcal{A}$, the derived ∞ -category $\text{D}(\mathcal{A})$ is stable. Moreover, the localization functor $\gamma: \text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$ sends short exact sequences to exact sequences.*

Proof. Assuming stability, the last claim is immediate from part (4) of the previous corollary. For stability, we saw in Corollary 6.1.28 that $\text{D}(\mathcal{A})$ admits finite limits and finite colimits. Moreover, the localization functor $\gamma: \text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$ preserves the zero object, so $\text{D}(\mathcal{A})$ is pointed. By

Theorem 4.2.2, it remains to show that the unit and counit of the adjunction $\Sigma \dashv \Omega$ is an equivalence. By the universal property of $\gamma: \text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$, this amounts to showing that for a chain complex $C_\bullet$, the unit $\gamma(C_\bullet) \rightarrow \Omega\Sigma\gamma(C_\bullet)$ and the counit $\Sigma\Omega\gamma(C_\bullet) \rightarrow \gamma(C_\bullet)$ are isomorphisms in $\text{D}(\mathcal{A})$.

To this end, consider the chain complex $C'_\bullet$ defined by $C'_n := C_n \oplus C_{n-1}$, with differential $d_n^{C'}(c_n, c_{n-1}) = (d^C c_n + c_{n-1}, -d^C c_{n-1})$; note that this is the mapping cylinder of the map $C_\bullet \rightarrow 0$. The maps $f(c_n) := (c_n, 0)$ and $g(c_n, c_{n-1}) := c_{n-1}$ define a short exact sequence

$$0 \rightarrow C_\bullet \xrightarrow{f} C'_\bullet \xrightarrow{g} C[1]_\bullet \rightarrow 0,$$

and by part (4) of Corollary 6.1.28 the resulting commutative square

$$\begin{array}{ccc} \gamma(C_\bullet) & \xrightarrow{\gamma(f)} & \gamma(C'_\bullet) \\ 0 \downarrow & \lrcorner & \downarrow \gamma(g) \\ 0 & \longrightarrow & \gamma(C[1]_\bullet) \end{array}$$

in $\text{D}(\mathcal{A})$ is both a pullback square and a pushout square. Since the map $C'_\bullet \rightarrow 0$ is a quasi-isomorphism, we get $\gamma(C'_\bullet) \simeq 0$, so this square simultaneously exhibits $\gamma(C[1]_\bullet)$ as the suspension of $\gamma(C_\bullet)$ and $\gamma(C_\bullet)$ as the loops of $\gamma(C[1]_\bullet)$. We conclude that the maps $\gamma(C_\bullet) \rightarrow \Omega\Sigma\gamma(C_\bullet)$ and $\Sigma\Omega\gamma(C_\bullet) \rightarrow \gamma(C_\bullet)$ are isomorphisms in $\text{D}(\mathcal{A})$. $\square$

Remark 6.1.30. The proof shows that the localization functor $\text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$ turns the explicit shift functor $(-)[n]$ from Example 6.1.10 into the abstract shift functor $[n]$ in the stable ∞ -category $\text{D}(\mathcal{A})$.

Corollary 6.1.31. *For a complex $A \in \text{D}(\mathcal{A})$ and $n \in \mathbb{Z}$, there exists a functorial exact sequence in $\text{D}(\mathcal{A})$ of the form*

$$\tau_{\geq n} A \rightarrow A \rightarrow \tau_{\leq n-1} A.$$

Proof. For a chain complex $C_\bullet \in \text{Ch}(\mathcal{A})$, put $Q_\bullet := C_\bullet / \tau_{\geq n} C_\bullet$. There is a functorial short exact sequence

$$0 \rightarrow \tau_{\geq n} C_\bullet \hookrightarrow C_\bullet \twoheadrightarrow Q_\bullet \rightarrow 0.$$

The natural map $Q_\bullet \rightarrow \tau_{\leq n-1} C_\bullet$, given in degree $n-1$ by the quotient $C_{n-1} \rightarrow \text{coker}(d_n)$ and by the identity below that degree, is a quasi-isomorphism. The theorem therefore turns the displayed short exact sequence into an exact sequence $\gamma\tau_{\geq n} \rightarrow \gamma \rightarrow \gamma\tau_{\leq n-1}$ in $\text{Fun}(\text{Ch}(\mathcal{A}), \text{D}(\mathcal{A}))$. By the universal property of $\gamma: \text{Ch}(\mathcal{A}) \rightarrow \text{D}(\mathcal{A})$, this descends to the desired exact sequence $\tau_{\geq n} \rightarrow \text{id} \rightarrow \tau_{\leq n-1}$. $\square$

6.2 Ordinary homology and cohomology via $\text{D}(\mathbb{Z})$

We now construct the ordinary homology and cohomology theories that were used axiomatically in Chapter 3. The construction is intrinsic to animae: instead of choosing a topological model and taking singular chains, we use the universal property of An to produce chain and cochain complexes directly in the derived category $\text{D}(\mathbb{Z})$. The bridge from chain complexes to spectra is provided by Eilenberg–MacLane spectra.

6.2.1 Eilenberg-MacLane spaces and spectra

Throughout this section we fix the abelian category $\mathcal{A} = \text{Ab}$ of abelian groups and work in its derived ∞ -category $D(\mathbb{Z})$. By forming mapping spectra out of the complex $\mathbb{Z}[0]$, we may associate a spectrum to every complex:

Definition 6.2.1 (Eilenberg-MacLane spectrum). Given a complex $A \in D(\mathbb{Z})$, we define its *Eilenberg-MacLane spectrum* as the mapping spectrum

$$HA := \text{hom}_{D(\mathbb{Z})}(\mathbb{Z}[0], A) \in \text{Sp}$$

Similarly, for $n \in \mathbb{Z}$ we define its *n-th Eilenberg MacLane anima* as

$$K(A, n) := \text{Hom}_{D(\mathbb{Z})}(\mathbb{Z}[-n], A) \in \text{An}.$$

These define functors $H: D(\mathbb{Z}) \rightarrow \text{Sp}$ and $K(-, n): D(\mathbb{Z}) \rightarrow \text{An}$.

In particular, every abelian group $A \in \text{Ab}$ defines an Eilenberg-MacLane spectrum HA and an n -th Eilenberg MacLane anima $K(A, n)$ by regarding it as a chain complex concentrated in degree 0.

Eilenberg-MacLane spectra play an important role in stable homotopy theory. In the next subsection, we will use them to classify ordinary (co)homology theories with coefficients in an abelian group.

Exercise 6.2.2. Show that for an associative ring R the forgetful functor $R\text{Mod}_R(\text{Ab}) \rightarrow \text{Ab}$ induces a functor $D(R) \rightarrow D(\mathbb{Z})$. Show that the composite $D(R) \rightarrow D(\mathbb{Z}) \xrightarrow{H} \text{Sp}$ may be identified with $\text{hom}_{D(R)}(R[0], -)$.

The main property of the Eilenberg-MacLane anima and spectra is that their homotopy groups are precisely the homology groups of A :

Proposition 6.2.3. *For a complex $A \in D(\mathbb{Z})$ and integers $n \in \mathbb{Z}$, $k \geq 0$, there are natural isomorphisms of abelian groups*

$$\pi_k(K(A, n)) \cong H_{k-n}(A) \quad \text{and} \quad \pi_n(HA) \cong H_n(A).$$

Remark 6.2.4. If A is an abelian group regarded as a complex concentrated in degree 0, then Proposition 6.2.3 shows that HA is connective and that $\Omega^\infty HA$ is the discrete commutative group A . Under the recognition equivalence of Theorem 5.4.5, it therefore corresponds to $A \in \text{CGrp}(\text{Set})$, giving a natural isomorphism

$$HA \xrightarrow{\sim} \mathbf{B}^\infty A.$$

Thus this construction agrees with the Eilenberg-MacLane spectrum of Definition 5.4.8.

This has the following useful consequence:

Lemma 6.2.5. *The functor $H: D(\mathbb{Z}) \rightarrow \text{Sp}$ preserves colimits.*

Proof. Since it is exact, it remains to show it preserves arbitrary coproducts: the map $\bigoplus_{i \in I} H(A_i) \rightarrow H(\bigoplus_{i \in I} A_i)$ is an isomorphism of spectra. This may be tested at the level of homotopy groups. Combining with Proposition 6.2.3, this then follows from the fact that both $\pi_n(-): \text{Sp} \rightarrow \text{Ab}$ and $H_n(-): D(\mathbb{Z}) \rightarrow \text{Ab}$ preserve coproducts. $\square$

To prove the proposition, we introduce an auxiliary ∞ -category in which the complexes $\mathbb{Z}[n]$ corepresent homology directly. We will then transport this corepresentability across the further localization to $D(\mathbb{Z})$.

Definition 6.2.6. Let $\mathcal{A}$ be an abelian category. We define the ∞ -category $\mathcal{K}(\mathcal{A})$ as the localization

$$\mathcal{K}(\mathcal{A}) := \text{Ch}(\mathcal{A})[\{\text{chain homotopy equivalences}^{-1}\}].$$

There are localization functors $\text{Ch}(\mathcal{A}) \rightarrow \mathcal{K}(\mathcal{A}) \rightarrow D(\mathcal{A})$.

A chain map $i: C_\bullet \rightarrow D_\bullet$ is called a *degreewise split monomorphism* if each component $i_n: C_n \rightarrow D_n$ is a split monomorphism in $\mathcal{A}$. The chosen retractions are not required to assemble into a chain map. Degreewise split epimorphisms are defined dually.

Proposition 6.2.7. *For every abelian category $\mathcal{A}$, the ∞ -category $\mathcal{K}(\mathcal{A})$ is stable. Under the localization functor $\text{Ch}(\mathcal{A}) \rightarrow \mathcal{K}(\mathcal{A})$, the suspension of a chain complex $C_\bullet$ is naturally isomorphic to the shifted complex $C[1]_\bullet$.*

Proof. The proof is the split analogue of the proofs of Proposition 6.1.26 and Theorem 6.1.29. Equip $\text{Ch}(\mathcal{A})$ with chain homotopy equivalences as weak equivalences and degreewise split monomorphisms as cofibrations. The mapping-cylinder factorization shows that every map factors as a cofibration followed by a weak equivalence. The pushout axiom follows from the criterion that a degreewise split monomorphism is a chain homotopy equivalence precisely when its cokernel is contractible. The dual argument uses degreewise split epimorphisms, so the localization $\mathcal{K}(\mathcal{A})$ admits finite limits and colimits.

For a chain complex $C_\bullet$, the degreewise split short exact sequence

$$0 \rightarrow C_\bullet \rightarrow \text{Cyl}(C_\bullet \rightarrow 0) \rightarrow C[1]_\bullet \rightarrow 0$$

is sent to a cofiber sequence in $\mathcal{K}(\mathcal{A})$. Its middle term is contractible, so it identifies suspension with the shift $[1]$. Thus suspension is an equivalence, with inverse induced by $[-1]$, and Theorem 4.2.2 implies that $\mathcal{K}(\mathcal{A})$ is stable. $\square$

Lemma 6.2.8. *The homotopy category of $\mathcal{K}(\mathcal{A})$ is the category whose objects are chain complexes and whose morphisms are chain maps modulo chain homotopy.*

Proof. As in the proof of Corollary 2.3.7, this follows from the observation that a functor $\text{Ch}(\mathcal{A}) \rightarrow C$ into a 1-category C inverts chain homotopy equivalences if and only if it identifies chain homotopic chain maps. $\square$

Corollary 6.2.9. *For a chain complex $C_\bullet \in \text{Ch}(\text{Ab})$ and $n \in \mathbb{Z}$ there is a natural isomorphism*

$$\pi_0 \text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[n], C_\bullet) \cong H_n(C).$$

In particular, the functor $\text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[n], -): \mathcal{K}(\text{Ab}) \rightarrow \text{An}$ corepresented by $\mathbb{Z}[n]$ inverts quasi-isomorphisms.

Proof. By Lemma 6.2.8, the left-hand side is the set of chain maps $\mathbb{Z}[n] \rightarrow C_\bullet$ up to chain homotopy. The first claim follows, since a chain map $\mathbb{Z}[n] \rightarrow C_\bullet$ is nothing but an element of $\ker(d_n: C_n \rightarrow C_{n-1})$ and two chain maps are chain homotopic if and only if their difference lies in the image of $d_{n+1}: C_{n+1} \rightarrow C_n$.

The second claim now follows from the stability of $\mathcal{K}(\text{Ab})$ established in Proposition 6.2.7: for $k \geq 0$ we have

$$\pi_k \text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[n], A) \cong \pi_0 \text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[n+k], A) \cong H_{n+k}(A). \quad \square$$

Lemma 6.2.10. *Let $\gamma: C \rightarrow C[W^{-1}]$ be the localization functor inverting a collection of morphisms W in some ∞ -category C . Let $X \in C$ be an object with the property that $\text{Hom}_C(X, -): C \rightarrow \text{An}$ inverts all morphisms in W . Then for every $Y \in C$ the map*

$$\text{Hom}_C(X, Y) \rightarrow \text{Hom}_{C[W^{-1}]}(\gamma X, \gamma Y)$$

is an isomorphism of anima.

Proof. The assumption on X guarantees that $\text{Hom}_C(X, -)$ descends to a functor $C[W^{-1}] \rightarrow \text{An}$. We need to show that this functor is corepresented by $\gamma X \in C[W^{-1}]$. But this follows immediately from the full faithfulness of $\gamma^*: \text{Fun}(C[W^{-1}], \text{An}) \hookrightarrow \text{Fun}(C, \text{An})$: for any other functor $F: C[W^{-1}] \rightarrow \text{An}$ we have

$$\text{Nat}_{C[W^{-1}] \rightarrow \text{An}}(\text{Hom}_C(X, -), F) \xrightarrow{\sim} \text{Nat}_{C \rightarrow \text{An}}(\text{Hom}_C(X, -), F \circ \gamma) \xrightarrow{\sim} F(\gamma(X)). \quad \square$$

We are now ready for the proof of Proposition 6.2.3:

Proof of Proposition 6.2.3. Combining Corollary 6.2.9 and Lemma 6.2.10, we see that for every complex $A \in \mathcal{K}(\text{Ab})$ and $n \in \mathbb{Z}$, the canonical map

$$\text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[-n], A) \rightarrow \text{Hom}_{\mathcal{D}(\mathbb{Z})}(\mathbb{Z}[-n], A) = K(A, n)$$

is an isomorphism of anima. In particular, we have

$$\pi_k(K(A, n)) \cong \pi_k \text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[-n], A) \cong \pi_0 \text{Hom}_{\mathcal{K}(\text{Ab})}(\mathbb{Z}[k-n], A) \cong H_{k-n}(A)$$

and thus

$$\pi_n(HA) \cong \pi_0 H(A[-n]) \cong \pi_0 K(A[-n], 0) \cong H_0(A[-n]) \cong H_n(A). \quad \square$$

6.2.2 Ordinary homology and cohomology

Recall from Section 3.3 the axiomatization of ordinary (co)homology with coefficients in an abelian group A . We can now describe these as the (co)homology theory associated to the Eilenberg-MacLane spectrum HA of A . More generally, we obtain versions of ordinary (co)homology with coefficients in an arbitrary complex of abelian groups.

Definition 6.2.11 (Reduced ordinary (co)homology). Let $A \in \mathcal{D}(\mathbb{Z})$ be a complex of abelian groups and let $X \in \text{An}_*$ be a pointed anima. We define the *reduced ordinary homology of X with coefficients*

in A and the *reduced ordinary cohomology of X with coefficients in A* as the (co)homology theories represented by the Eilenberg-MacLane spectrum HA :

$$\tilde{H}_n(X; A) := HA_n(X) = \pi_n(\Sigma^\infty X \otimes HA)$$

and

$$\tilde{H}^n(X; A) := HA^n(X) = \pi_{-n} \operatorname{hom}(\Sigma^\infty X, HA).$$

We will mostly be interested in the case where $A \in \mathbf{Ab}$ is an abelian group, regarded as a complex concentrated in degree 0.

The groups $\tilde{H}_*(X; A)$ and $\tilde{H}^*(X; A)$ may alternatively be computed as the homology groups of certain chain complexes associated to X and A .

Construction 6.2.12 (Chain and cochain complexes). Let $A \in D(\mathbb{Z})$ be a complex. Since $D(\mathbb{Z})$ admits all small limits and colimits by Corollary 6.1.28, the universal property of $\mathbf{An}$ from Corollary 1.8.10 implies that evaluation at the point induces equivalences

$$\operatorname{ev}_{\text{pt}}: \operatorname{Fun}^{\operatorname{colim}}(\mathbf{An}, D(\mathbb{Z})) \xrightarrow{\sim} D(\mathbb{Z}) \quad \text{and} \quad \operatorname{ev}_{\text{pt}}: \operatorname{Fun}^{\operatorname{lim}}(\mathbf{An}^{\operatorname{op}}, D(\mathbb{Z})) \xrightarrow{\sim} D(\mathbb{Z}).$$

In particular, the object $A \in D(\mathbb{Z})$ determines two functors

$$C_*(-; A): \mathbf{An} \rightarrow D(\mathbb{Z}) \quad \text{and} \quad C^*(-; A): \mathbf{An}^{\operatorname{op}} \rightarrow D(\mathbb{Z}),$$

characterized uniquely by the properties that $C_*(-; A)$ preserves colimits, $C^*(-; A)$ preserves limits, and $C_*(\text{pt}; A) \simeq A \simeq C^*(\text{pt}; A)$.

For a commutative ring R , the same construction in $D(R)$ defines chains and cochains with coefficients in any $A \in D(R)$. Restriction of scalars along $\mathbb{Z} \rightarrow R$ identifies their underlying complexes in $D(\mathbb{Z})$ with the constructions above.

For a pointed anima $(X, x) \in \mathbf{An}_*$, we define the *reduced chain complex* and *reduced cochain complex* as

$$\tilde{C}_*(X; A) := \operatorname{cofib}(A \simeq C_*(\text{pt}; A) \xrightarrow{x_*} C_*(X; A))$$

and

$$\tilde{C}^*(X; A) := \operatorname{fib}(C^*(X; A) \xrightarrow{x^*} C^*(\text{pt}; A) \simeq A).$$

Definition 6.2.13 (Ordinary (co)homology). Given an anima $X \in \mathbf{An}$, a chain complex $A \in D(\mathbb{Z})$, and an integer $n \in \mathbb{Z}$, we define the *ordinary homology* and *ordinary cohomology of X with coefficients in A* as

$$H_n(X; A) := H_n(C_*(X; A)) \quad \text{and} \quad H^n(X; A) := H_{-n}(C^*(X; A)).$$

When $A = \mathbb{Z}$ (regarded as a complex concentrated in degree 0), we simply write $H_n(X) := H_n(X; \mathbb{Z})$ and $H^n(X) := H^n(X; \mathbb{Z})$.

The next result shows that the spectrum-theoretic and chain complex definitions of ordinary (co)homology agree.

Proposition 6.2.14. *Let $A \in D(\mathbb{Z})$ be a chain complex. There are natural equivalences of functors*

$$H(C_*(-; A)) \simeq \Sigma_+^\infty(-) \otimes HA: \mathbf{An} \rightarrow \mathbf{Sp}$$

and

$$H(C^*(-; A)) \simeq \mathrm{hom}(\Sigma_+^\infty(-), HA): \mathbf{An}^{\mathrm{op}} \rightarrow \mathbf{Sp}.$$

In particular, for any anima X there are natural isomorphisms

$$H_n(X; A) \cong \pi_n(\Sigma_+^\infty X \otimes HA) \quad \text{and} \quad H^n(X; A) \cong \pi_{-n} \mathrm{hom}(\Sigma_+^\infty X, HA).$$

Proof. Since the Eilenberg-MacLane functor $H: D(\mathbb{Z}) \rightarrow \mathbf{Sp}$ preserves colimits by Lemma 6.2.5, it follows that $H \circ C_*(-; A): \mathbf{An} \rightarrow \mathbf{Sp}$ preserves colimits. On the other hand, the functor $\Sigma_+^\infty(-) \otimes HA$ also preserves colimits, since both Σ_+^∞ and $- \otimes HA$ do. Since both evaluate to HA on the point, it follows they are naturally equivalent. The equivalence $H(C^*(-; A)) \simeq \mathrm{hom}(\Sigma_+^\infty(-), HA)$ follows similarly, as both sides preserve limits.

The final statement follows by taking homotopy groups and using Proposition 6.2.3. $\square$

Corollary 6.2.15. *For a pointed anima $X \in \mathbf{An}_*$ and a chain complex $A \in D(\mathbb{Z})$, there are natural isomorphisms*

$$\tilde{H}_n(X; A) \cong H_n(\tilde{C}_*(X; A)) \quad \text{and} \quad \tilde{H}^n(X; A) \cong H_{-n}(\tilde{C}^*(X; A)).$$

Proof. We have

$$\tilde{H}_n(X; A) = \pi_n(\Sigma^\infty X \otimes HA) \cong \pi_n(\mathrm{cofib}(\mathbb{S} \otimes HA \rightarrow \Sigma_+^\infty X \otimes HA))$$

since $\Sigma^\infty X = \mathrm{cofib}(\mathbb{S} \rightarrow \Sigma_+^\infty X)$ and $- \otimes HA$ is exact. By Proposition 6.2.14, this is isomorphic to

$$\pi_n(H(\mathrm{cofib}(A \rightarrow C_*(X; A)))) = \pi_n(H(\tilde{C}_*(X; A))) = H_n(\tilde{C}_*(X; A)).$$

The cohomological case is similar. $\square$

Lemma 6.2.16. *For any anima X there is an isomorphism $H_0(X) \cong \bigoplus_{\pi_0(X)} \mathbb{Z}$.*

Proof. Since both $\Sigma_+^\infty X$ and $H\mathbb{Z}$ are connective spectra, so is $\Sigma_+^\infty X \otimes H\mathbb{Z}$. The functor $H_0(-): \mathbf{An} \rightarrow \mathbf{Ab}$ may thus be written as the composite

$$\mathbf{An} \xrightarrow{\Sigma_+^\infty} \mathbf{Sp}_{\geq 0} \xrightarrow{- \otimes H\mathbb{Z}} \mathbf{Sp}_{\geq 0} \xrightarrow{\pi_0} \mathbf{Ab},$$

each of which preserves colimits. Here $\pi_0: \mathbf{Sp}_{\geq 0} \rightarrow \mathbf{Ab}$ preserves colimits because it is left adjoint to the inclusion of abelian groups as Eilenberg-MacLane spectra. On the other hand, the assignment $X \mapsto \bigoplus_{\pi_0(X)} \mathbb{Z}$ may be written as the composite of $\pi_0: \mathbf{An} \rightarrow \mathbf{Set}$ with the free abelian group functor $\mathbf{Set} \rightarrow \mathbf{Ab}$, both of which preserve colimits as well. By the universal property of $\mathbf{An}$, it thus suffices to check the claim for $X = \mathrm{pt}$, where both sides are $\mathbb{Z}$. $\square$

For an abelian group A , one can show that the assignments $X \mapsto C_*^{\mathrm{sing}}(X; A)$ and $X \mapsto C_{\mathrm{sing}}^*(X; A)$ from Remark 3.3.2 are homotopy invariant, send disjoint unions to (co)products, satisfy the Mayer-Vietoris property, and carry skeletal colimits of CW-complexes to colimits and limits, respectively;

see for example Hatcher [Hat02]. By restricting these assignments to CW-complexes, it follows that they descend to functors

$$C_*^{\text{sing}}(-; A): \mathbf{An} \rightarrow \mathbf{D}(\mathbb{Z}) \quad \text{and} \quad C_{\text{sing}}^*(-; A): \mathbf{An}^{\text{op}} \rightarrow \mathbf{D}(\mathbb{Z})$$

that preserve colimits and limits, respectively. Since $C_*^{\text{sing}}(\text{pt}; A) \simeq A[0] \simeq C_{\text{sing}}^*(\text{pt}; A)$, the universal property of $\mathbf{An}$ implies that these functors are naturally equivalent to $C_*(-; A)$ and $C^*(-; A)$ from Construction 6.2.12. Thus the intrinsic ordinary (co)homology constructed here agrees with singular (co)homology for anima coming from topological spaces.

Finally, for an abelian group A and the pointed anima S^0 , we have

$$\widetilde{H}_n(S^0; A) \cong \pi_n(HA) \cong H_n(A) \quad \text{and} \quad \widetilde{H}^n(S^0; A) \cong \pi_{-n}(HA) \cong H_{-n}(A).$$

Both groups are therefore A when $n = 0$ and vanish otherwise. This verifies the dimension axiom and completes the construction of the ordinary theories postulated in Theorem 3.3.1.

6.3 Connective objects and t-structures

In Section 6.1, we saw that the derived ∞ -category $\mathbf{D}(\mathcal{A})$ of an abelian category $\mathcal{A}$ admits a natural decomposition into ‘connective’ and ‘coconnective’ parts. Specifically, a complex $A \in \mathbf{D}(\mathcal{A})$ is called connective if its homology vanishes in negative degrees, and coconnective if its homology vanishes in positive degrees. These two classes of objects interact in a structured way:

- (1) (Closure under shifts) Connective complexes are closed under positive shifts, while coconnective complexes are closed under negative shifts.
- (2) (Orthogonality) There are no nonzero maps from a connective complex to a (-1) -coconnective complex.
- (3) (Decomposition) Every complex A sits in an exact sequence $\tau_{\geq 0}A \rightarrow A \rightarrow \tau_{\leq -1}A$ decomposing it into a connective and a (-1) -coconnective part.

The ∞ -category $\mathbf{Sp}$ of spectra exhibits the same pattern: a spectrum X is called connective if $\pi_k(X) = 0$ for $k < 0$, and coconnective if $\pi_k(X) = 0$ for $k > 0$. These subcategories again satisfy properties (1)–(3).

Structures like these are omnipresent in stable homotopy theory and homological algebra. The axiomatic framework capturing this phenomenon is that of a *t-structure* on a stable ∞ -category, introduced by Beilinson, Bernstein, and Deligne [BBD82]. One of the key features of a t-structure is its *heart*: the full subcategory of objects that are both connective and coconnective. The heart turns out to be an abelian 1-category, providing a bridge between the stable and abelian worlds. This allows many arguments from classical homological algebra to be carried out in any stable ∞ -category equipped with a t-structure, even when that category does not arise as the derived category of an abelian category.

In Section 6.3.1 we give the definition of a t-structure and establish its basic properties, including the existence of truncation functors. In Section 6.3.2 we prove that the heart of a t-structure is an abelian category, and relate exact sequences in $\mathcal{C}$ to short exact sequences in the heart. In Section 6.3.3

we introduce homotopy group objects and establish the long exact sequence of homotopy groups. Finally, in Section 6.3.4 we discuss completeness and boundedness conditions that control how well objects are determined by their truncations.

6.3.1 Definition and basic properties

Definition 6.3.1 ([BBD82]). A *t-structure* on a stable ∞ -category C consists of a pair $(C_{\geq 0}, C_{\leq 0})$ of full subcategories of C satisfying the following conditions:

- (1) (Closure under shifts) We have $C_{\geq 0}[1] \subseteq C_{\geq 0}$ and $C_{\leq 0}[-1] \subseteq C_{\leq 0}$.
- (2) (Orthogonality) If $X \in C_{\geq 0}$ and $Y \in C_{\leq 0}$, then $\mathrm{Hom}_C(X, Y[-1]) = 0$.
- (3) (Decomposition) Every $X \in C$ sits in an exact sequence of the form

$$\tau_{\geq 0}X \rightarrow X \rightarrow \tau_{\leq -1}X,$$

with $\tau_{\geq 0}X \in C_{\geq 0}$ and $\tau_{\leq -1}X \in C_{\leq 0}[-1]$.

Objects in $C_{\geq 0}$ are called *connective* (with respect to the t-structure), while objects in $C_{\leq 0}$ are called *coconnective*. For $n \in \mathbb{Z}$ we set $C_{\geq n} := C_{\geq 0}[n]$ and $C_{\leq n} := C_{\leq 0}[n]$.

If D is another stable ∞ -category equipped with a t-structure, then an exact functor $F: C \rightarrow D$ is called *left t-exact* if it sends $C_{\leq 0}$ to $D_{\leq 0}$, and hence by exactness sends $C_{\leq n}$ to $D_{\leq n}$ for all $n \in \mathbb{Z}$. Similarly, F is called *right t-exact* if it sends $C_{\geq 0}$ to $D_{\geq 0}$, or equivalently sends $C_{\geq n}$ to $D_{\geq n}$ for all n . We say F is *t-exact* if it is both left and right t-exact.

Remark 6.3.2 (Cohomological grading). We index t-structures *homologically*, matching our convention for chain complexes (Remark 6.1.8). A reader preferring *cohomological* grading should negate the index throughout, writing $C^{\geq n} := C_{\leq -n}$ and $C^{\leq n} := C_{\geq -n}$, and likewise $\tau^{\geq n} := \tau_{\leq -n}$ and $\tau^{\leq n} := \tau_{\geq -n}$ for the truncation functors introduced below.

The two main examples of t-structures are the standard t-structure on derived categories and the Postnikov t-structure on spectra.

Proposition 6.3.3 (Standard t-structure on derived categories). *Let $\mathcal{A}$ be an abelian category. The pair $(D(\mathcal{A})_{\geq 0}, D(\mathcal{A})_{\leq 0})$ defines a t-structure on $D(\mathcal{A})$, called the standard t-structure.*

Proof. Property (1) is immediate from the definition of $D(\mathcal{A})_{\geq n}$ and $D(\mathcal{A})_{\leq n}$: a complex with homology concentrated in degrees ≥ 0 has, after a positive shift, homology concentrated in degrees ≥ 1 , so in particular ≥ 0 .

Property (2) was established in Corollary 6.1.24: if A is connective and B is (-1) -coconnective, then $\mathrm{Hom}_{D(\mathcal{A})}(A, B) = 0$.

Property (3) is Corollary 6.1.31: for every complex A , there is an exact sequence $\tau_{\geq 0}A \rightarrow A \rightarrow \tau_{\leq -1}A$ in $D(\mathcal{A})$. $\square$

Proposition 6.3.4 (Postnikov t-structure on spectra). *Define $\mathrm{Sp}_{\geq 0} \subseteq \mathrm{Sp}$ to be the full subcategory of connective spectra, i.e. those X with $\pi_k(X) = 0$ for $k < 0$, and $\mathrm{Sp}_{\leq 0} \subseteq \mathrm{Sp}$ the full subcategory of coconnective spectra, i.e. those with $\pi_k(X) = 0$ for $k > 0$. Then the pair $(\mathrm{Sp}_{\geq 0}, \mathrm{Sp}_{\leq 0})$ defines a t-structure on Sp , called the Postnikov t-structure.*

Proof. Property (1) is immediate from the behavior of homotopy groups under shifts: $\pi_k(X[1]) \cong \pi_{k-1}(X)$.

For property (2), let $X \in \mathrm{Sp}_{\geq 0}$ and $Y \in \mathrm{Sp}_{\leq 0}$. By the recognition principle for connective spectra (Theorem 5.4.5), we may write $X \simeq \mathbf{B}^\infty(A)$ for some $A \in \mathrm{CGrp}(\mathrm{An})$. By adjunction,

$$\mathrm{Hom}_{\mathrm{Sp}}(X, Y[-1]) \cong \mathrm{Hom}_{\mathrm{CGrp}(\mathrm{An})}(A, \Omega^\infty(Y[-1])).$$

But $\Omega^\infty(Y[-1])$ is contractible: its homotopy groups $\pi_k(\Omega^\infty(Y[-1])) \cong \pi_k(Y[-1]) \cong \pi_{k+1}(Y)$ vanish for all $k \geq 0$ since $Y \in \mathrm{Sp}_{\leq 0}$.

For property (3), given a spectrum X , we define $\tau_{\geq 0}X := \mathbf{B}^\infty \Omega^\infty X$. The counit of the adjunction $\mathbf{B}^\infty \dashv \Omega^\infty$ provides a map $\tau_{\geq 0}X \rightarrow X$. Define $\tau_{\leq -1}X$ as its cofiber. From the long exact sequence on homotopy groups (Proposition 4.4.29), one verifies that $\tau_{\geq 0}X$ is connective and $\tau_{\leq -1}X$ has homotopy groups concentrated in degrees ≤ -1 . $\square$

Example 6.3.5. If C admits a t -structure $(C_{\geq 0}, C_{\leq 0})$, then C^{op} can be given the *opposite t -structure*, in which $(C^{\mathrm{op}})_{\geq 0} := (C_{\leq 0})^{\mathrm{op}}$ and $(C^{\mathrm{op}})_{\leq 0} := (C_{\geq 0})^{\mathrm{op}}$.

Example 6.3.6. If C and D both admit t -structures, then so does their product $C \times D$ by setting $(C \times D)_{\geq 0} := C_{\geq 0} \times D_{\geq 0}$ and $(C \times D)_{\leq 0} := C_{\leq 0} \times D_{\leq 0}$.

We now show that the objects $\tau_{\geq 0}X$ and $\tau_{\leq -1}X$ appearing in axiom (3) are in fact functorial in X , arising from adjunctions.

Lemma 6.3.7. *Let $(C_{\geq 0}, C_{\leq 0})$ be a t -structure on C , and let $n \in \mathbb{Z}$.*

- (1) *The inclusion $C_{\geq n} \hookrightarrow C$ admits a right adjoint $\tau_{\geq n}: C \rightarrow C_{\geq n}$.*
- (2) *The inclusion $C_{\leq n} \hookrightarrow C$ admits a left adjoint $\tau_{\leq n}: C \rightarrow C_{\leq n}$.*

Proof. We prove (1); the proof of (2) is dual. By shifting, we may assume that $n = 0$; in general we may then set $\tau_{\geq n}(X) := \tau_{\geq 0}(X[-n])[n]$. Using the pointwise formula for adjunctions, it suffices to show that for every object $X \in C$ there exists some $\tau_{\geq 0}X$ equipped with a morphism $\tau_{\geq 0}X \rightarrow X$ inducing equivalences $\mathrm{Hom}_C(Y, \tau_{\geq 0}X) \xrightarrow{\sim} \mathrm{Hom}_C(Y, X)$ for every $Y \in C_{\geq 0}$. For this, we consider the exact sequence $\tau_{\geq 0}X \rightarrow X \rightarrow \tau_{\leq -1}X$ provided by axiom (3). This gives rise to a long exact sequence

$$\cdots \rightarrow \pi_{n+1} \mathrm{Hom}_C(Y, \tau_{\leq -1}X) \rightarrow \pi_n \mathrm{Hom}_C(Y, \tau_{\geq 0}X) \rightarrow \pi_n \mathrm{Hom}_C(Y, X) \rightarrow \pi_n \mathrm{Hom}_C(Y, \tau_{\leq -1}X) \rightarrow \cdots$$

Since $\mathrm{Hom}_C(Y, \tau_{\leq -1}X) = 0$ by axiom (2), this implies that the map $\mathrm{Hom}_C(Y, \tau_{\geq 0}X) \rightarrow \mathrm{Hom}_C(Y, X)$ induces isomorphisms on all homotopy groups, hence is an equivalence. $\square$

Corollary 6.3.8. *The subcategory $C_{\leq n}$ of C is closed under all limits that exist in C . The subcategory $C_{\geq n}$ of C is closed under all colimits that exist in C .* $\square$

Corollary 6.3.9. *Assume C admits I -indexed colimits. Then so does the subcategory $C_{\leq n}$: they are computed as $\tau_{\leq n}(\mathrm{colim}_{i \in I} X_i)$. Dually, if C admits I -indexed limits, then so does $C_{\geq n}$ via the formula $\tau_{\geq n}(\mathrm{lim}_{i \in I} X_i)$.* $\square$

Lemma 6.3.10. *Let C be a stable ∞ -category equipped with a t -structure, and let $n, m \in \mathbb{Z}$.*

(1) If $X \in C_{\leq m}$, then $\tau_{\geq n}X \in C_{\leq m}$.

(2) If $X \in C_{\geq n}$, then $\tau_{\leq m}X \in C_{\geq n}$.

Proof. We prove (1) first. Let $X \in C_{\leq m}$. If $n > m$, then $\text{Hom}_C(Z, X) = 0$ for every $Z \in C_{\geq n}$ by orthogonality. By adjunction, this gives $\text{Hom}_C(Z, \tau_{\geq n}X) = 0$ for every $Z \in C_{\geq n}$. Taking $Z = \tau_{\geq n}X$, we see that the identity of $\tau_{\geq n}X$ is null, so $\tau_{\geq n}X$ is zero.

Assume instead that $n \leq m$. The truncation sequence

$$\tau_{\geq n}X \rightarrow X \rightarrow \tau_{\leq n-1}X$$

exhibits $\tau_{\geq n}X$ as the fiber of a morphism between objects of $C_{\leq m}$, since $C_{\leq n-1} \subseteq C_{\leq m}$. By Corollary 6.3.8, it follows that $\tau_{\geq n}X \in C_{\leq m}$.

The proof of (2) is dual. □

Warning 6.3.11. The notation $\tau_{\leq n}$ is sometimes also used for the notion of n -truncation in an ∞ -category. This notation is *not* compatible with the left adjoint $\tau_{\leq n}$ from the lemma. In fact, the only n -truncated object in a stable ∞ -category is the zero object. (Compare Lemma 6.3.18.)

Exercise 6.3.12. Let C and D have t-structures and let $F : C \rightleftarrows D : G$ be an adjunction. Show that F is right t-exact if and only if G is left t-exact.

Lemma 6.3.13. Let $F : C \rightarrow D$ be a t-exact functor between two stable ∞ -categories equipped with t-structures. Then for all $X \in C$ and $n \in \mathbb{Z}$, there are natural equivalences

$$\tau_{\leq n}F(X) \xrightarrow{\sim} F(\tau_{\leq n}X) \quad \text{and} \quad F(\tau_{\geq n}X) \xrightarrow{\sim} \tau_{\geq n}F(X).$$

Proof. We prove the first equivalence; the second is dual. Since F is t-exact, applying F to the exact sequence $\tau_{\geq n+1}X \rightarrow X \rightarrow \tau_{\leq n}X$ gives another exact sequence

$$F(\tau_{\geq n+1}X) \rightarrow F(X) \rightarrow F(\tau_{\leq n}X)$$

where the first object lies in $D_{\geq n+1}$ and the last one in $D_{\leq n}$. It follows that the second map induces an isomorphism $\tau_{\leq n}F(X) \xrightarrow{\sim} F(\tau_{\leq n}X)$. □

6.3.2 The heart of a t-structure

Given a t-structure on C , a special role is played by the objects that are both connective and coconnective.

Definition 6.3.14. Let $(C_{\geq 0}, C_{\leq 0})$ be a t-structure on C . We define the *heart* of the t-structure to be the intersection

$$C^\heartsuit := C_{\geq 0} \cap C_{\leq 0}.$$

The key feature of t-structures is that the heart is always an abelian 1-category, and that there is a close relation between exact sequences in $C^\heartsuit$ (in the sense of homological algebra) and exact sequences in C (in the sense of stable homotopy theory).

Proposition 6.3.15. Let C be a stable ∞ -category equipped with a t-structure.

- (1) The heart $C^\heartsuit$ is an abelian 1-category.
- (2) A morphism $g: Y \rightarrow Z$ in $C^\heartsuit$ is an epimorphism in $C^\heartsuit$ if and only if its fiber in C lies in $C^\heartsuit$.
- (3) A morphism $f: X \rightarrow Y$ in $C^\heartsuit$ is a monomorphism in $C^\heartsuit$ if and only if its cofiber in C lies in $C^\heartsuit$.
- (4) Consider a commutative square in $C^\heartsuit$ of the form

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ h \downarrow & & \downarrow g \\ W & \xrightarrow{k} & Z. \end{array}$$

- (a) If g and h are epimorphisms in $C^\heartsuit$, then the square is a pullback square in $C^\heartsuit$ if and only if it is a pullback square in C .
- (b) If f and k are monomorphisms in $C^\heartsuit$, then the square is a pushout square in $C^\heartsuit$ if and only if it is a pushout square in C .

Proof. For (1), first observe that $C^\heartsuit$ is a 1-category, in the sense that the hom animae $\mathrm{Hom}_{C^\heartsuit}(X, Y)$ are sets. Indeed, for $k \geq 1$, we have

$$\pi_k \mathrm{Hom}_{C^\heartsuit}(X, Y) \cong \pi_0 \mathrm{Hom}_C(X, Y[-k]) = 0$$

since $X \in C_{\geq 0}$ and $Y[-k] \in C_{\leq -k} \subseteq C_{\leq -1}$. Further, since both $C_{\leq 0}$ and $C_{\geq 0}$ are closed under direct sums in C , it is clear $C^\heartsuit$ is additive. We also observe that $C^\heartsuit$ admits kernels and cokernels, which are computed as

$$\ker(Y \rightarrow Z) = \tau_{\geq 0}(\mathrm{fib}(Y \rightarrow Z)) \quad \text{and} \quad \mathrm{coker}(X \rightarrow Y) = \tau_{\leq 0}(\mathrm{cofib}(X \rightarrow Y)).$$

Before finishing the proof of (1), let us first address the ‘only if’-directions in (2) and (3). Since replacing C by C^{op} translates (2) into (3) and vice versa, it suffices to prove the ‘only if’-direction in (2). To this end, let $g: Y \rightarrow Z$ be an epimorphism in $C^\heartsuit$. We need to show that $\mathrm{fib}(g) \in C_{\geq 0}$. (Since $C_{\leq 0}$ is closed under limits, the condition $\mathrm{fib}(g) \in C_{\leq 0}$ is automatic.) The assumption on g implies that the commutative square

$$\begin{array}{ccc} Y & \xrightarrow{g} & Z \\ g \downarrow & & \downarrow = \\ Z & \xrightarrow{=} & Z \end{array}$$

is a pushout square in $C^\heartsuit$. Since pushouts in $C^\heartsuit$ are computed by first forming the pushout in $C_{\geq 0}$ and then applying $\tau_{\leq 0}: C_{\geq 0} \rightarrow C^\heartsuit$, it follows that the map $Z \rightarrow Z \sqcup_Y Z$ obtained by forming the pushout in C induces an isomorphism $Z = \tau_{\leq 0}Z \xrightarrow{\sim} \tau_{\leq 0}(Z \sqcup_Y Z)$. Passing to cofibers then gives $\tau_{\leq 0}(\mathrm{cofib}(g)) = 0$, or equivalently $\mathrm{cofib}(g) \in C_{\geq 1}$. But then the relation $\mathrm{fib}(g) \simeq \mathrm{cofib}(g)[-1]$ implies $\mathrm{fib}(g) \in C_{\geq 0}$, which is what we needed to show.

We now prove (4). Again, (a) translates to (b) under replacing C by C^{op} , so it suffices to prove (a). By full faithfulness of the inclusion $C^\heartsuit \hookrightarrow C$, it is clear that if the square is a pullback in C then it is also a pullback in $C^\heartsuit$. Conversely, if the square is a pullback in $C^\heartsuit$, then it in particular induces an isomorphism $\ker(h) \xrightarrow{\sim} \ker(g)$ on kernels in $C^\heartsuit$. But by the direction of (2) that we just proved,

these kernels agree with the fibers in C . We deduce that the induced map $\text{fib}(h) \rightarrow \text{fib}(g)$ is an isomorphism in C , which implies that the square is also a pullback square in C . This finishes the proof of (4).

We now return to (1), finishing the proof that $C^\heartsuit$ is abelian. Consider a commutative square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow g \\ 0 & \longrightarrow & Z \end{array}$$

in $C^\heartsuit$, and assume that f is a monomorphism in $C^\heartsuit$ and that g is an epimorphism in $C^\heartsuit$. We need to show that this square is a pullback square in $C^\heartsuit$ if and only if it is a pushout square in $C^\heartsuit$. But by (4) this is immediate from stability of C .

Finally, we prove the ‘if’-directions in (2) and (3). As before, it suffices to do this for (2). Given a morphism $g: Y \rightarrow Z$ in $C^\heartsuit$ whose fiber lies in $C^\heartsuit$, we must show g is an epimorphism. Since $C^\heartsuit$ is abelian, it suffices to show that its cokernel $\text{coker}(g) = \tau_{\leq 0}(\text{cofib}(g))$ is zero, i.e., that $\text{cofib}(g) \in C_{\geq 1}$. But this follows from the relation $\text{cofib}(g) \simeq \text{fib}(g)[1]$ and the assumption $\text{fib}(g) \in C_{\geq 0}$. $\square$

Corollary 6.3.16. *Let C be a stable ∞ -category equipped with a t -structure. For every short exact sequence*

$$0 \rightarrow X \hookrightarrow Y \twoheadrightarrow Z \rightarrow 0$$

in $C^\heartsuit$, the sequence $X \rightarrow Y \rightarrow Z$ is an exact sequence in C .

Proof. This is an immediate consequence of part (4) of Proposition 6.3.15. $\square$

Corollary 6.3.17. *Let $F: C \rightarrow D$ be an exact functor between stable ∞ -categories with t -structures. Assume F restricts to a functor $F: C^\heartsuit \rightarrow D^\heartsuit$. Then this restriction is an exact functor of abelian categories.*

Proof. Let $0 \rightarrow X \hookrightarrow Y \twoheadrightarrow Z \rightarrow 0$ be a short exact sequence in $C^\heartsuit$. By the previous corollary, the sequence $X \rightarrow Y \rightarrow Z$ is exact in C , hence induces an exact sequence $F(X) \rightarrow F(Y) \rightarrow F(Z)$ in D . Using the various parts of Proposition 6.3.15, we see:

- (i) By (2), the morphism $F(Y) \rightarrow F(Z)$ is an epimorphism in $D^\heartsuit$, since its fiber $F(X)$ lies in $D^\heartsuit$;
- (ii) By (3), the morphism $F(X) \rightarrow F(Y)$ is a monomorphism in $D^\heartsuit$, since its cofiber $F(Z)$ lies in $D^\heartsuit$;
- (iii) By (4), the exact sequence $F(X) \rightarrow F(Y) \rightarrow F(Z)$ in D is also exact in $D^\heartsuit$.

We conclude that the sequence $0 \rightarrow F(X) \hookrightarrow F(Y) \twoheadrightarrow F(Z) \rightarrow 0$ is exact in $D^\heartsuit$, as desired. $\square$

To identify the heart of a t -structure, the following observation is sometimes useful:

Lemma 6.3.18. *Let C be a stable ∞ -category equipped with a t -structure, and let $X \in C_{\geq 0}$ be a connective object. Then X lies in the heart of C if and only if it is a 0-truncated object of $C_{\geq 0}$, in the sense that $\pi_k \text{Hom}_{C_{\geq 0}}(Y, X) = 0$ for all $Y \in C_{\geq 0}$ and $k > 0$.*

Proof. First assume that $X \in C^\heartsuit$. Then for $Y \in C_{\geq 0}$ and $k > 0$ we have $\pi_k \operatorname{Hom}_C(Y, X) \simeq \pi_0 \operatorname{Hom}_C(Y, X[-k]) = 0$ as $X[-k] \in C_{\leq -k} \subseteq C_{\leq -1}$. Conversely, if X is 0-truncated in $C_{\geq 0}$, then we have $\pi_k \operatorname{Hom}_C(Y, X[-1]) \cong \pi_{k+1} \operatorname{Hom}_C(Y, X) \cong 0$ for all $Y \in C_{\geq 0}$ and $k \geq 0$. It follows that $X[-1] \in C_{\leq -1}$, and thus $X \in C^\heartsuit$. $\square$

Corollary 6.3.19. *For a stable ∞ -category equipped with a t-structure, the heart $C^\heartsuit$ is the full subcategory of 0-truncated objects in $C_{\geq 0}$.* $\square$

We now identify the hearts of our two main examples of t-structures.

Example 6.3.20 (Heart of the standard t-structure). Let $\mathcal{A}$ be an abelian category. The heart of the standard t-structure on $D(\mathcal{A})$ is equivalent to $\mathcal{A}$. Indeed, the heart consists of those complexes whose homology is concentrated in degree zero, which by Proposition 6.1.25 is precisely the essential image of the fully faithful functor $\mathcal{A} \hookrightarrow D(\mathcal{A})$.

Proposition 6.3.21 (Heart of the Postnikov t-structure). *The functor $\mathbf{B}^\infty: \operatorname{Ab} \hookrightarrow \operatorname{Sp}$ induces an equivalence*

$$\mathbf{B}^\infty: \operatorname{Ab} \xrightarrow{\sim} \operatorname{Sp}^\heartsuit.$$

Proof. By the recognition principle for connective spectra (Theorem 5.4.5), the functor $\mathbf{B}^\infty$ induces an equivalence $\mathbf{B}^\infty: \operatorname{CGrp}(\operatorname{An}) \xrightarrow{\sim} \operatorname{Sp}_{\geq 0}$. By Corollary 6.3.19, we may identify $\operatorname{Sp}^\heartsuit$ with the subcategory of $\operatorname{CGrp}(\operatorname{An})$ spanned by the 0-truncated objects. Note that object in $\operatorname{CGrp}(\operatorname{An})$ is 0-truncated if and only if its underlying anima is 0-truncated, which in turn happens if and only if it is contained in the full subcategory $\operatorname{Set} \subseteq \operatorname{An}$. All in all, we get $\operatorname{Ab} = \operatorname{CGrp}(\operatorname{Set}) \xrightarrow{\sim} \operatorname{CGrp}(\operatorname{An})_{\leq 0} \xrightarrow{\sim} \operatorname{Sp}^\heartsuit$. $\square$

Remark 6.3.22. Combining Example 6.3.20 and Proposition 6.3.21, we see that the Eilenberg–MacLane functor $H: D(\mathbb{Z}) \rightarrow \operatorname{Sp}$ from Definition 6.2.1 restricts to an equivalence $\operatorname{Ab} \xrightarrow{\sim} \operatorname{Sp}^\heartsuit$ on hearts. In particular, H is t-exact.

More generally, for a connective ring spectrum R , the heart of the module category LMod_R can be identified with the category of discrete modules over $\pi_0(R)$. We will establish this in Section 8.2, after developing the necessary theory of ring spectra and their modules.

6.3.3 Homotopy group objects

We will now define the *homotopy group objects* of an object $X \in C$: the objects of the heart obtained by truncating X both from above and below. First observe that this operation is independent of the order of truncation:

Lemma 6.3.23. *Let C be a stable ∞ -category equipped with a t-structure. For $n, m \in \mathbb{Z}$, the canonical natural transformation*

$$\tau_{\leq m} \tau_{\geq n} \rightarrow \tau_{\geq n} \tau_{\leq m}$$

is a natural isomorphism of functors $C \rightarrow C_{\leq m} \cap C_{\geq n}$.

Proof. By Lemma 6.3.10, both composites take values in $C_{\leq m} \cap C_{\geq n}$. If $n > m$, this intersection is zero by orthogonality, so both composites are zero. We may therefore assume that $n \leq m$.

Fixing $Y \in C$, the map $\tau_{\geq n}Y \rightarrow Y$ induces $\tau_{\leq m}\tau_{\geq n}Y \rightarrow \tau_{\leq m}Y$. Since the source is n -connective, this factors uniquely through a map

$$\tau_{\leq m}\tau_{\geq n}Y \rightarrow \tau_{\geq n}\tau_{\leq m}Y,$$

which is the ‘canonical map’ from the statement. To show it is an isomorphism, we may equivalently show that the cofiber of $\tau_{\leq m}\tau_{\geq n}Y \rightarrow \tau_{\leq m}Y$ is $(n-1)$ -coconnective. To this end, we may compare the truncation sequences for $\tau_{\geq n}Y$ and Y to obtain a commutative diagram

$$\begin{array}{ccccc} \tau_{\geq m+1}\tau_{\geq n}Y & \longrightarrow & \tau_{\geq n}Y & \longrightarrow & \tau_{\leq m}\tau_{\geq n}Y \\ \cong \downarrow & & \downarrow & & \downarrow \\ \tau_{\geq m+1}Y & \longrightarrow & Y & \longrightarrow & \tau_{\leq m}Y. \end{array}$$

Since $m \geq n$, the left vertical map is an isomorphism. By the pasting law for exact squares, it follows that the cofiber of the right vertical map is the cofiber of the middle vertical map. This cofiber is $\tau_{\leq n-1}Y$, which is $(n-1)$ -coconnective, finishing the proof. $\square$

Definition 6.3.24. Let C be a stable ∞ -category equipped with a t -structure and let X be an object of C . For every $n \in \mathbb{Z}$, we define the n -th homotopy group object $\pi_n(X) \in C^\heartsuit$ by

$$\pi_n(X) := \tau_{\geq 0}\tau_{\leq 0}(X[-n]) \in C^\heartsuit.$$

Remark 6.3.25. In the context of homological algebra, for example when C is the derived ∞ -category of an abelian category, the objects $\pi_n(X)$ are often denoted by $H_n(X)$ instead, emphasizing the analogy with the homology groups of a chain complex rather than the analogy with homotopy groups of a spectrum. Under the equivalence $D(\mathcal{A})^\heartsuit \simeq \mathcal{A}$ from Example 6.3.20, the object $\pi_n(A) \in \mathcal{A}$ agrees with the n -th homology $H_n(A)$ of A viewed as a chain complex.

Lemma 6.3.26. Let $F: C \rightarrow D$ be a t -exact functor between two stable ∞ -categories equipped with t -structures. Then for all $X \in C$ and $n \in \mathbb{Z}$, there is a natural isomorphism $F(\pi_n X) \xrightarrow{\sim} \pi_n F(X)$.

Proof. By Lemma 6.3.13 we have natural isomorphisms

$$F(\pi_n X) = F(\tau_{\geq 0}\tau_{\leq 0}(X[-n])) \xrightarrow{\sim} \tau_{\geq 0}\tau_{\leq 0}(F(X)[-n]) = \pi_n(F(X)). \quad \square$$

Just like for spectra, exact sequences in C induce long exact sequences on homotopy groups:

Proposition 6.3.27 (Long exact sequence of homotopy groups). *Let C be a stable ∞ -category equipped with a t -structure. For every exact sequence $X \xrightarrow{f} Y \xrightarrow{g} Z$ in C , we obtain a long exact sequence of homotopy groups in $C^\heartsuit$:*

$$\cdots \rightarrow \pi_{n+1}(Z) \xrightarrow{\partial} \pi_n(X) \xrightarrow{\pi_n(f)} \pi_n(Y) \xrightarrow{\pi_n(g)} \pi_n(Z) \xrightarrow{\partial} \pi_{n-1}(X) \rightarrow \cdots$$

Proof. Since every such exact sequence in C induces exact sequences $Z[-1] \rightarrow X \rightarrow Y$ and $Y \rightarrow Z \rightarrow X[1]$ and we have $\pi_n(Z[-1]) = \pi_{n+1}(Z)$ and $\pi_n(X[1]) = \pi_{n-1}(X)$, it suffices by induction to show that the sequence

$$\pi_0(X) \xrightarrow{\pi_0(f)} \pi_0(Y) \xrightarrow{\pi_0(g)} \pi_0(Z)$$

is exact in $C^\heartsuit$. We will prove this in progressively more general cases.

Case 1: Assume that $X, Y, Z \in C_{\geq 0}$. We claim that in this case, the sequence

$$\pi_0(X) \rightarrow \pi_0(Y) \rightarrow \pi_0(Z) \rightarrow 0$$

is exact. To see this, it suffices to show that for every $W \in C^\heartsuit$, the induced sequence of abelian groups

$$0 \rightarrow \text{Hom}_{C^\heartsuit}(\pi_0(Z), W) \rightarrow \text{Hom}_{C^\heartsuit}(\pi_0(Y), W) \rightarrow \text{Hom}_{C^\heartsuit}(\pi_0(X), W)$$

is exact. Since $Z \in C_{\geq 0}$, we have $\pi_0(Z) = \tau_{\leq 0}Z$, so by adjunction we obtain isomorphisms

$$\text{Hom}_{C^\heartsuit}(\pi_0(Z), W) \cong \pi_0 \text{Hom}_C(Z, W) \cong \pi_0 \text{hom}_C(Z, W),$$

and similarly for X and Y . Furthermore, we have $\pi_1 \text{hom}_C(X, W) \cong \pi_0 \text{hom}_C(X[1], W) \cong 0$ as $\tau_{\leq 0}(X[1]) \cong 0$. The claim now follows from the long exact sequence on homotopy groups associated to the exact sequence of mapping spectra

$$\text{hom}_C(Z, W) \rightarrow \text{hom}_C(Y, W) \rightarrow \text{hom}_C(X, W).$$

Case 2: Assume that $X \in C_{\geq 0}$, but Y and Z are arbitrary. For every $W \in C_{\leq -1}$, orthogonality and the exact sequence $X \rightarrow Y \rightarrow Z$ identify $\text{Hom}_C(Z, W)$ with $\text{Hom}_C(Y, W)$. By adjunction and the Yoneda lemma in $C_{\leq -1}$, the induced map $\tau_{\leq -1}Y \rightarrow \tau_{\leq -1}Z$ is therefore an isomorphism, so we get a cofiber sequence

$$0 \cong \tau_{\leq -1}X \rightarrow \tau_{\leq -1}Y \rightarrow \tau_{\leq -1}Z,$$

showing that $\tau_{\leq -1}Y \cong \tau_{\leq -1}Z$. Consider now the following diagram:

$$\begin{array}{ccccc} \tau_{\geq 0}X & \longrightarrow & \tau_{\geq 0}Y & \longrightarrow & \tau_{\geq 0}Z \\ \cong \downarrow & & \downarrow & & \downarrow \\ X & \longrightarrow & Y & \longrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \tau_{\leq -1}Y & \xrightarrow{\cong} & \tau_{\leq -1}Z. \end{array}$$

Since all three columns are exact and the bottom two rows are exact, it follows that the top row is also exact. Since $\pi_0(X) \cong \pi_0(\tau_{\geq 0}X)$, and similarly for Y and Z , the exactness of $\pi_0(X) \rightarrow \pi_0(Y) \rightarrow \pi_0(Z) \rightarrow 0$ has thus been reduced to Case 1.

Case 3: Assume that $Z \in C_{\leq 0}$, but X and Y are arbitrary. Using an argument dual to that of Cases 1 and 2, we obtain an exact sequence

$$0 \rightarrow \pi_0(X) \rightarrow \pi_0(Y) \rightarrow \pi_0(Z).$$

Case 4: Finally, we consider the case for general X, Y, Z . Let Q denote the cofiber of the composite $\tau_{\geq 0}X \rightarrow X \rightarrow Y$. We then have the following commutative diagram:

$$\begin{array}{ccccc} \tau_{\geq 0}X & \longrightarrow & Y & \longrightarrow & Q \\ \downarrow & & \parallel & & \downarrow \\ X & \longrightarrow & Y & \longrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ \tau_{\leq -1}X & \longrightarrow & 0 & \longrightarrow & (\tau_{\leq -1}X)[1]. \end{array}$$

The three rows are exact, as are the left two columns, and hence so is the third column. Applying Cases 2 and 3, we get exact sequences

$$\pi_0(X) \rightarrow \pi_0(Y) \twoheadrightarrow \pi_0(Q) \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \pi_0(Q) \hookrightarrow \pi_0(Z) \rightarrow \pi_{-1}(X).$$

Gluing them together then gives the desired exact sequence $\pi_0(X) \rightarrow \pi_0(Y) \rightarrow \pi_0(Z)$. $\square$

6.3.4 Complete and bounded t-structures

One of the primary advantages of having a t-structure on a given stable ∞ -category C is that it often allows us to reduce questions about arbitrary objects of C to questions about objects of the heart $C^\heartsuit$, which may be easier to address. In order for this strategy to work, we need to ensure that the truncation functors $\tau_{\leq n}$ and $\tau_{\geq n}$ retain enough information about the objects of C . For this purpose, we now introduce completeness and boundedness conditions on C .

Definition 6.3.28. A t-structure on C is *left complete* if for every $X \in C$ the canonical map

$$X \longrightarrow \lim_n \tau_{\leq n} X$$

is an isomorphism. It is *right complete* if the canonical map

$$\operatorname{colim}_n \tau_{\geq -n} X \longrightarrow X$$

is an isomorphism for every $X \in C$.

We say that the t-structure is *left separated* if every infinitely connective object $X \in C_{\geq \infty} := \bigcap_{n \in \mathbb{Z}} C_{\geq n}$ is zero. Dually, it is *right separated* if every infinitely coconnective object $X \in C_{\leq -\infty} := \bigcap_{n \in \mathbb{Z}} C_{\leq n}$ is zero.

Separatedness says that no nonzero object is invisible to all truncations, while completeness says that every object can be reconstructed from its truncations. Under the hypotheses of the criterion below, these two conditions agree.

Example 6.3.29. Let $\mathcal{A}$ be an abelian category. The standard t-structure on $D(\mathcal{A})$ is left and right separated: an infinitely connective or infinitely coconnective complex has trivial homology and is therefore zero. If $\mathcal{A}$ has exact countable products, then the t-structure is left complete; if it has exact countable coproducts, then it is right complete. Indeed, the countable version of the construction underlying Corollary 6.1.28 supplies the required (co)products in $D(\mathcal{A})$, and the criterion below applies.

Example 6.3.30. The Postnikov t-structure on Sp is left and right separated, since a spectrum with vanishing homotopy groups is zero. Homotopy groups commute with products and coproducts, so the criterion below shows that this t-structure is also left and right complete.

Lemma 6.3.31. *Let C be a stable ∞ -category with a t-structure. Assume that C admits countable products and that $C_{\geq 0}$ is closed under countable products. Then the t-structure is left complete if and only if it is left separated. Dually, if C admits countable coproducts and $C_{\leq 0}$ is closed under countable coproducts, then the t-structure is right complete if and only if it is right separated.*

Proof. This is [Lur17, Proposition 1.2.1.19]. $\square$

Definition 6.3.32. A t-structure on C is *right bounded* if every object $X \in C$ is contained in $C_{\geq n}$ for some $n \in \mathbb{Z}$. It is *left bounded* if every $X \in C$ is contained in $C_{\leq n}$ for some $n \in \mathbb{Z}$. It is *bounded* if it is both left and right bounded.

Definition 6.3.33. Let C be a stable ∞ -category with a t-structure. We define the subcategory of *bounded objects* by

$$C^b := \bigcup_{n \in \mathbb{N}} (C_{\geq -n} \cap C_{\leq n}) \subseteq C.$$

Observe that C^b inherits a t-structure from C which by construction is bounded. Moreover, C is bounded if and only if $C = C^b$. One may similarly define subcategories

$$C^- := \bigcup_{n \in \mathbb{N}} C_{\geq -n} \subseteq C \quad \text{and} \quad C^+ := \bigcup_{n \in \mathbb{N}} C_{\leq n} \subseteq C$$

of *bounded below* and *bounded above* objects, respectively.

Example 6.3.34. For the derived category $D(\mathcal{A})$ of an abelian category, the subcategory $D(\mathcal{A})^b$ is often denoted $D^b(\mathcal{A})$ and called the *bounded derived category*. Similarly, $D(\mathcal{A})^-$ is denoted $D^-(\mathcal{A})$ and $D(\mathcal{A})^+$ is denoted $D^+(\mathcal{A})$.

Exercise 6.3.35. Let C and D be stable ∞ -categories with t-structure, and assume that C is left bounded. Show that the restriction functor

$$\mathrm{Fun}^{t\text{-}\mathrm{rex}}(C, D) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{rex}}(C_{\geq 0}, D_{\geq 0})$$

provides an equivalence between right t-exact functors $F: C \rightarrow D$ and right exact functors $C_{\geq 0} \rightarrow D_{\geq 0}$.

6.4 Projective objects and Ext-groups

In classical algebra, projective modules are the ‘well-behaved’ building blocks essential for homological algebra. Their definition makes sense in an arbitrary abelian category $\mathcal{A}$: an object P is called *projective* if the Hom functor $\mathrm{Hom}_{\mathcal{A}}(P, -): \mathcal{A} \rightarrow \mathrm{Ab}$ is an exact functor of abelian categories. This definition has an immediate analogue in an arbitrary t-structure:

Definition 6.4.1. Let C be a stable ∞ -category equipped with a t-structure. An object $P \in C$ is called *t-projective* if the mapping spectrum functor $\mathrm{hom}_C(P, -): C \rightarrow \mathrm{Sp}$ is t-exact.

Observation 6.4.2. Every t-projective object P is connective. To see this, consider the truncation $\tau_{\leq -1}P$. By t-exactness, the mapping spectrum $\mathrm{hom}_C(P, \tau_{\leq -1}P)$ lies in $\mathrm{Sp}_{\leq -1}$, and it follows that

$$0 = \pi_0 \mathrm{hom}_C(P, \tau_{\leq -1}P) \cong \pi_0 \mathrm{Hom}_C(\tau_{\leq -1}P, \tau_{\leq -1}P),$$

where the last identification holds by adjunction. In particular, the identity on $\tau_{\leq -1}P$ is homotopic to the zero map, forcing $\tau_{\leq -1}P = 0$, hence $P \in C_{\geq 0}$.

Observation 6.4.3. Conversely, if $P \in C_{\geq 0}$ is a connective object, then $\mathrm{hom}_C(P, -): C \rightarrow \mathrm{Sp}$ is always left t-exact: if $X \in C_{\leq 0}$, then for $n > 0$ we have

$$\pi_n(\mathrm{hom}_C(P, X)) \cong \pi_0 \mathrm{Hom}_C(P, X[-n]) = 0,$$

since $P \in C_{\geq 0}$ while $X[-n] \in C_{\leq -1}$. So P is t-projective if and only if $\mathrm{hom}_C(P, -)$ is also *right* t-exact.

In the case $C = D(\mathcal{A})$, this definition does indeed recover the usual notion of projectivity in $\mathcal{A}$:

Lemma 6.4.4. *Let $\mathcal{A}$ be an abelian category and let $M \in \mathcal{A}$. Then M is projective in $\mathcal{A}$ if and only if the complex $M[0] \in D(\mathcal{A})$ is t-projective.*

Proof. Suppose first that $M[0]$ is t-projective, so that $\text{hom}_{D(\mathcal{A})}(M[0], -): D(\mathcal{A}) \rightarrow \text{Sp}$ is t-exact. By Corollary 6.3.17, this functor restricts to an exact functor on hearts $D(\mathcal{A})^\heartsuit \rightarrow \text{Sp}^\heartsuit$, i.e., $\mathcal{A} \rightarrow \text{Ab}$. This restricted functor is $\text{Hom}_{\mathcal{A}}(M, -)$, showing that M is projective in $\mathcal{A}$.

Conversely, assume that $\text{Hom}_{\mathcal{A}}(M, -): \mathcal{A} \rightarrow \text{Ab}$ is exact. Just as in Corollary 6.2.9, this implies that for any chain complex $C_\bullet$ in $\mathcal{A}$ we have an isomorphism

$$\pi_n \text{hom}_{\mathcal{K}(\mathcal{A})}(M[0], C_\bullet) \cong \text{Hom}_{\mathcal{A}}(M, H_n(C)).$$

In particular, the functor $\text{hom}_{\mathcal{K}(\mathcal{A})}(M[0], -): \mathcal{K}(\mathcal{A}) \rightarrow \text{Sp}$ inverts quasi-isomorphisms. The same argument as in Lemma 6.2.10 shows that it is corepresented by $M[0] \in D(\mathcal{A})$, giving a natural isomorphism $\text{hom}_{D(\mathcal{A})}(M[0], -) \cong \text{hom}_{\mathcal{K}(\mathcal{A})}(M[0], -)$. It follows that $\pi_n \text{hom}_{D(\mathcal{A})}(M[0], X) \cong \text{Hom}_{\mathcal{A}}(M, H_n(X))$ for all complexes $X \in D(\mathcal{A})$, showing that $\text{hom}_{D(\mathcal{A})}(M[0], X)$ is connective whenever X is. $\square$

Corollary 6.4.5. *Let $\mathcal{A}$ be an abelian category and let $M \in \mathcal{A}$ be projective. Then for every $X \in D(\mathcal{A})$ and every $n \in \mathbb{Z}$, there is a natural isomorphism*

$$\pi_n \text{hom}_{D(\mathcal{A})}(M[0], X) \cong \text{Hom}_{\mathcal{A}}(M, H_n(X)).$$

Proof. This is the computation in the second half of the proof of Lemma 6.4.4. $\square$

6.4.1 Ext-groups

We will next give a formulation of t-projectivity in terms of the Ext-groups in C .

Definition 6.4.6 (Ext-groups). Let C be a stable ∞ -category. For objects $X, Y \in C$ and $n \in \mathbb{Z}$, we define the n -th Ext-group to be

$$\text{Ext}_C^n(X, Y) := \pi_{-n} \text{hom}_C(X, Y) \simeq \pi_0 \text{Hom}_C(X, Y[n]).$$

For objects $M, N \in \mathcal{A}$, regarded as complexes concentrated in degree zero, we abbreviate $\text{Ext}_{D(\mathcal{A})}^n(M, N)$ to $\text{Ext}_{\mathcal{A}}^n(M, N)$.

Assume now that C comes equipped with a t-structure. For objects $M, N \in C^\heartsuit$, the Ext-groups $\text{Ext}_C^n(M, N)$ admit a direct interpretation in small degrees:

Observation 6.4.7. For $n < 0$, we have $\text{Ext}_C^n(M, N) = 0$, since $M \in C_{\geq 0}$ while $N[n] \in C_{\leq n} \subseteq C_{\leq -1}$.

Observation 6.4.8. For $n = 0$, we have $\text{Ext}_C^0(M, N) = \pi_0 \text{Hom}_C(M, N) = \text{Hom}_{C^\heartsuit}(M, N)$, the set of morphisms in the heart.

The interpretation of $\text{Ext}_C^1(M, N)$ in terms of extensions is developed in Exercise 6.7.

Exercise 6.4.9. Assume that $C^\heartsuit$ has enough projectives, meaning that for every object M there exists an epimorphism $P \twoheadrightarrow M$ from a projective object P . Show that $\text{Ext}_C^n(M, N)$ for $n \geq 2$ can be described in terms of *degree n Yoneda extensions*: exact sequences in $C^\heartsuit$ of the form

$$0 \rightarrow N \rightarrow E_n \rightarrow E_{n-1} \rightarrow \cdots \rightarrow E_2 \rightarrow E_1 \rightarrow M \rightarrow 0.$$

Hint: use Proposition 6.4.14 below.

We now record the fundamental long exact sequences for Ext-groups.

Observation 6.4.10 (Ext long exact sequence). Every exact sequence $Y' \rightarrow Y \rightarrow Y''$ in C induces an exact sequence of mapping spectra $\text{hom}_C(X, Y') \rightarrow \text{hom}_C(X, Y) \rightarrow \text{hom}_C(X, Y'')$, thus giving rise to a long exact sequence of homotopy groups of the form

$$\begin{array}{ccccccc} & & & & \cdots & \longrightarrow & \text{Ext}_C^{n-2}(X, Y'') \\ & \swarrow & & & & & \searrow \\ \text{Ext}_C^{n-1}(X, Y') & \longrightarrow & \text{Ext}_C^{n-1}(X, Y) & \longrightarrow & \text{Ext}_C^{n-1}(X, Y'') & & \\ & \swarrow & & & & & \searrow \\ & \text{Ext}_C^n(X, Y') & \longrightarrow & \text{Ext}_C^n(X, Y) & \longrightarrow & \text{Ext}_C^n(X, Y'') & \\ & \swarrow & & & & & \searrow \\ & \text{Ext}_C^{n+1}(X, Y') & \longrightarrow & \cdots & & & \end{array}$$

Similarly, every exact sequence $X' \rightarrow X \rightarrow X''$ in C induces a long exact sequence of the form

$$\cdots \rightarrow \text{Ext}_C^{n-1}(X', Y) \rightarrow \text{Ext}_C^n(X'', Y) \rightarrow \text{Ext}_C^n(X, Y) \rightarrow \text{Ext}_C^n(X', Y) \rightarrow \text{Ext}_C^{n+1}(X'', Y) \rightarrow \cdots$$

As a special case, given an object $N \in C^\heartsuit$ and a short exact sequence $M' \hookrightarrow P \twoheadrightarrow M$ in $C^\heartsuit$, the negative Ext-groups vanish, giving a one-sided long exact sequence of the form

$$\begin{aligned} 0 \rightarrow \text{Hom}_{C^\heartsuit}(M, N) &\rightarrow \text{Hom}_{C^\heartsuit}(P, N) \rightarrow \text{Hom}_{C^\heartsuit}(M', N) \\ &\rightarrow \text{Ext}_C^1(M, N) \rightarrow \text{Ext}_C^1(P, N) \rightarrow \text{Ext}_C^1(M', N) \\ &\rightarrow \text{Ext}_C^2(M, N) \rightarrow \cdots \end{aligned}$$

Using Ext-groups, we can give several equivalent characterizations of t-projectivity.

Proposition 6.4.11 ([Lur17, Proposition 7.2.2.6]). *Let C be a stable ∞ -category equipped with a t -structure, and let $P \in C_{\geq 0}$ be a connective object. Then the following conditions are equivalent:*

- (1) *The object P is t -projective;*
- (2) *For every $Q \in C_{\geq 0}$ and every $i > 0$, the abelian group $\text{Ext}_C^i(P, Q)$ is zero;*
- (3) *For every $Q \in C_{\geq 0}$, the abelian group $\text{Ext}_C^1(P, Q)$ is zero;*
- (4) *Given an exact sequence*

$$N' \rightarrow N \rightarrow N''$$

in C with $N', N, N'' \in C_{\geq 0}$, the map $\text{Ext}_C^0(P, N) \rightarrow \text{Ext}_C^0(P, N'')$ is surjective.

If C is left complete, this is further equivalent to:

(5) For every $Q \in C^\heartsuit$ and every $i > 0$, the abelian group $\mathrm{Ext}_C^i(P, Q)$ is zero.

Proof. Recall from Observation 6.4.3 that P is t-projective if and only if the mapping spectrum functor $\mathrm{hom}_C(P, -) : C \rightarrow \mathrm{Sp}$ is right t-exact. By definition, this means that for any object $Q \in C_{\geq 0}$, the mapping spectrum $\mathrm{hom}_C(P, Q)$ has vanishing negative homotopy groups $\pi_{-i} \mathrm{hom}_C(P, Q) = \mathrm{Ext}_C^i(P, Q)$ for $i > 0$. This shows the equivalence (1) $\Leftrightarrow$ (2).

It is clear that (2) implies (3), and the implication (3) $\Rightarrow$ (2) follows by replacing Q by $Q[i - 1]$ for all $i \geq 1$.

We next show that (3) $\Leftrightarrow$ (4). It is clear that (3) implies (4): the exact sequence $N' \rightarrow N \rightarrow N''$ induces a long exact sequence of the form

$$\cdots \rightarrow \mathrm{Ext}_C^0(P, N) \rightarrow \mathrm{Ext}_C^0(P, N'') \rightarrow \mathrm{Ext}_C^1(P, N') \rightarrow \cdots,$$

so the vanishing of $\mathrm{Ext}_C^1(P, N')$ implies that the first map must be surjective. Assume now that (4) holds. Given a class $\eta \in \mathrm{Ext}_C^1(P, Q) = \pi_0 \mathrm{Hom}_C(P, Q[1])$, we obtain an exact sequence

$$P' \rightarrow P \xrightarrow{\eta} Q[1]$$

and thus a long exact sequence

$$\cdots \rightarrow \mathrm{Ext}_C^0(P, P') \rightarrow \mathrm{Ext}_C^0(P, P) \xrightarrow{\eta \circ -} \mathrm{Ext}_C^1(P, Q) \rightarrow \cdots$$

As $P, P', Q \in C_{\geq 0}$, the first map is surjective, hence the second map is the zero map. Since it sends id_P to η , we conclude that $\eta = 0$. This shows that (3) $\Leftrightarrow$ (4).

Finally, suppose that C is left complete. It is clear that (2) implies (5). Conversely, assume (5). The exact sequences

$$(\pi_{n+1}Q)[n+1] \rightarrow \tau_{\leq n+1}Q \rightarrow \tau_{\leq n}Q$$

show that the towers of Ext-groups in each positive degree are eventually constant. Since left completeness gives $Q \simeq \lim_n \tau_{\leq n}Q$, the corresponding tower of mapping spectra computes $\mathrm{hom}_C(P, Q)$. Eventual constancy in adjacent degrees eliminates the derived-limit term and gives

$$\mathrm{Ext}_C^i(P, Q) \cong \mathrm{Ext}_C^i(P, \tau_{\leq 0}Q) = 0$$

for $i > 0$. This is the Postnikov-tower argument of [Lur17, Proposition 7.2.2.6]. $\square$

6.4.2 Ext-groups via projective resolutions

As a consequence of Proposition 6.4.11, we may compute Ext-groups in terms of projective resolutions.

Definition 6.4.12. Let C be a stable ∞ -category equipped with a t-structure, and let $M \in C^\heartsuit$ be an object in the heart of C . A *projective resolution* of M is a long exact sequence

$$\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \twoheadrightarrow M \rightarrow 0$$

in $C^\heartsuit$ such that each object P_n is t-projective.

Remark 6.4.13. When $C = D(\mathcal{A})$ is the derived ∞ -category of some abelian category $\mathcal{A}$, this recovers the usual notion of projective resolutions in $\mathcal{A}$ from homological algebra by Lemma 6.4.4.

Proposition 6.4.14 (Ext-groups via projective resolutions). *Let C be a stable ∞ -category equipped with a t -structure, and let $P_\bullet$ be a projective resolution of $M \in C^\heartsuit$. Then for every other object $N \in C^\heartsuit$ and every $n \in \mathbb{Z}$ there is an isomorphism of abelian groups*

$$\mathrm{Ext}_C^n(M, N) \cong H^n(\mathrm{Hom}_{C^\heartsuit}(P_\bullet, N))$$

between the n -th Ext-group and the n -th cohomology group of the cochain complex $\mathrm{Hom}_{C^\heartsuit}(P_\bullet, N)$.

Proof. The claim is clear for $n < 0$ as both sides are zero: $\mathrm{Ext}_C^n(M, N) = \pi_0 \mathrm{Hom}_C(M, N[n]) = 0$ since $M \in C_{\geq 0}$ and $N[n] \in C_{\leq n} \subseteq C_{\leq -1}$.

For the remaining cases, we break the resolution into short exact sequences and use the vanishing of the higher Ext-groups of the P_k to splice the resulting long exact sequences together. Define auxiliary objects $M_{-1} := M$ and $M_k := \ker(P_k \twoheadrightarrow M_{k-1})$ for $k \geq 0$. The projective resolution then admits the following factorization in $C^\heartsuit$, where each of the diagonal sequences are exact:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \searrow & & \searrow & & \searrow \\
 & & & M_2 & & & M_0 \\
 & & \nearrow & \searrow & \nearrow & \searrow & \nearrow \\
 \cdots & \cdots & \cdots & P_3 & \cdots & P_2 & \cdots & P_1 & \cdots & P_0 & \cdots \\
 & & \nearrow & & \nearrow & & \nearrow & & \nearrow & & \nearrow \\
 & & & & & M_1 & & & & M & \\
 & & & & & \searrow & & & & \searrow & \\
 & & & & 0 & & 0 & & & 0 &
 \end{array}$$

For each of these short exact sequences, we obtain a long exact sequence on Ext-groups from Observation 6.4.10. Since the higher Ext-groups of the t -projective objects P_k vanish by Proposition 6.4.11, its initial part is

$$\begin{aligned}
 0 \rightarrow \mathrm{Hom}_{C^\heartsuit}(M_{k-1}, N) &\rightarrow \mathrm{Hom}_{C^\heartsuit}(P_k, N) \rightarrow \mathrm{Hom}_{C^\heartsuit}(M_k, N) \\
 &\rightarrow \mathrm{Ext}_C^1(M_{k-1}, N) \rightarrow 0,
 \end{aligned}$$

and its remaining terms give isomorphisms

$$\mathrm{Ext}_C^{i+1}(M_{k-1}, N) \cong \mathrm{Ext}_C^i(M_k, N) \quad (i \geq 1).$$

Since the map $P_{k+1} \twoheadrightarrow M_k$ is an epimorphism, the induced map $\mathrm{Hom}_{C^\heartsuit}(M_k, N) \hookrightarrow \mathrm{Hom}_{C^\heartsuit}(P_{k+1}, N)$ is injective, and so the initial part of this sequence provides for each $k \geq 0$ an isomorphism

$$\mathrm{Hom}_{C^\heartsuit}(M_{k-1}, N) \cong \ker(\mathrm{Hom}_{C^\heartsuit}(P_k, N) \rightarrow \mathrm{Hom}_{C^\heartsuit}(P_{k+1}, N)).$$

The next part of the sequence then shows that $\mathrm{Ext}_C^1(M_{k-1}, N)$ is the cokernel of the map $\mathrm{Hom}_{C^\heartsuit}(P_k, N) \rightarrow \mathrm{Hom}_{C^\heartsuit}(M_k, N)$. Combining this with the previous identification of $\mathrm{Hom}_{C^\heartsuit}(M_k, N)$ as a kernel, this shows that

$$\mathrm{Ext}_C^1(M_{k-1}, N) \cong \frac{\ker(\mathrm{Hom}_{C^\heartsuit}(P_{k+1}, N) \rightarrow \mathrm{Hom}_{C^\heartsuit}(P_{k+2}, N))}{\mathrm{im}(\mathrm{Hom}_{C^\heartsuit}(P_k, N) \rightarrow \mathrm{Hom}_{C^\heartsuit}(P_{k+1}, N))} = H^{k+1}(\mathrm{Hom}_{C^\heartsuit}(P_\bullet, N)).$$

Observe that setting $k = 0$ in these two isomorphisms proves the statement of the proposition for $n = 0$ and $n = 1$. Iterating the displayed dimension-shifting isomorphisms gives $\mathrm{Ext}_C^n(M, N) \cong \mathrm{Ext}_C^1(M_{n-2}, N)$ for all $n \geq 1$. Setting $k = n - 1$ in the previous isomorphism thus gives

$$\mathrm{Ext}_C^n(M, N) \cong H^n(\mathrm{Hom}_{C^\heartsuit}(P_\bullet, N))$$

for all $n \geq 1$, finishing the proof. $\square$

6.4.3 Projective objects and geometric realizations

There is a second natural way to generalize the notion of projectivity beyond the abelian setting. An object P in an abelian category $\mathcal{A}$ is projective if and only if $\mathrm{Hom}_{\mathcal{A}}(P, -)$ preserves epimorphisms. Since epimorphisms in $\mathcal{A}$ are precisely the reflexive coequalizers, this is equivalent to asking that $\mathrm{Hom}_{\mathcal{A}}(P, -)$ preserves reflexive coequalizers. Since the ∞ -categorical analogues of reflexive coequalizers are geometric realizations, this leads to the following definition:

Definition 6.4.15. Let D be an ∞ -category with geometric realizations. An object $P \in D$ is called *projective* if the functor $\mathrm{Hom}_D(P, -): D \rightarrow \mathbf{An}$ preserves geometric realizations.

If C is a stable ∞ -category with a left complete t-structure, it turns out that an object of C is t-projective if and only if it is a projective object in the connective part $C_{\geq 0}$. We start with the following auxiliary result:

Lemma 6.4.16. *Assume that the t-structure on C is left complete. Then $C_{\geq 0}$ admits geometric realizations. If C' is another stable ∞ -category with a left complete t-structure, then any exact right t-exact functor $F: C \rightarrow C'$ restricts to a functor $F: C_{\geq 0} \rightarrow C'_{\geq 0}$ that preserves geometric realizations.*

Proof. By [Lur17, Lemma 1.3.3.11(2)], the connective part of a left-complete t-structure admits geometric realizations, and a right exact functor between connective parts preserves them. The restriction of an exact right t-exact functor is right exact, which gives the second claim. $\square$

As a second auxiliary result, we record the following general description of pushouts. Besides its application below, it illustrates why geometric realizations play the role of reflexive coequalizers in the preceding discussion.

Lemma 6.4.17 (Pushouts as geometric realizations). *Let D be an ∞ -category with finite coproducts and geometric realizations. Given a span $X \xleftarrow{f} A \xrightarrow{g} Y$, there exists a simplicial object $B_\bullet(X, A, Y)$ with*

$$B_n(X, A, Y) := X \sqcup A^{\sqcup n} \sqcup Y.$$

Its geometric realization is the pushout:

$$|B_\bullet(X, A, Y)| \cong X \sqcup_A Y.$$

Proof. **TO DO.** This is an instance of the Bousfield–Kan formula for colimits. $\square$

Proposition 6.4.18. *Let C be a stable ∞ -category equipped with a left-complete t -structure, and let $P \in C_{\geq 0}$ be a connective object of C . Then P is t -projective if and only if it is a projective object in $C_{\geq 0}$ in the sense of Definition 6.4.15.*

Proof. First assume that P is projective in $C_{\geq 0}$. We will verify condition (3) from Proposition 6.4.11. Given an object $Q \in C_{\geq 0}$, apply Lemma 6.4.17 to the span $0 \leftarrow Q \rightarrow 0$. The resulting simplicial object has $Q^{\oplus n}$ in degree n : its face maps add adjacent summands or omit the first or last summand, and its degeneracy maps insert a zero summand. Thus it begins

$$\dots \rightrightarrows Q \oplus Q \rightrightarrows Q \rightrightarrows 0.$$

Its geometric realization is the pushout $0 \sqcup_Q 0$. By Lemma 6.3.7, the inclusion $C_{\geq 0} \hookrightarrow C$ has a right adjoint and therefore preserves this pushout, which in the stable category C is the suspension $Q[1]$. Since P is projective, $\mathrm{Hom}_C(P, Q[1])$ is therefore the geometric realization of the simplicial anima obtained by mapping out of P . The set of components of a geometric realization is a quotient of the set of components in simplicial degree zero, which here is the singleton $\pi_0 \mathrm{Hom}_C(P, 0) = *$. Thus $\mathrm{Hom}_C(P, Q[1])$ is connected, so $\mathrm{Ext}_C^1(P, Q) = 0$. This shows that P is t -projective.

For the converse, assume now that P is t -projective, giving a t -exact functor $\mathrm{hom}_C(P, -): C \rightarrow \mathrm{Sp}$. We get from Lemma 6.4.16 that the restricted functor $\mathrm{hom}_C(P, -): C_{\geq 0} \rightarrow \mathrm{Sp}_{\geq 0}$ preserves geometric realizations. Since $\Omega^\infty: \mathrm{Sp}_{\geq 0} \xrightarrow{\sim} \mathrm{CGrp}(\mathrm{An}) \rightarrow \mathrm{An}$ preserves geometric realizations as well, it follows that so does $\mathrm{Hom}_C(P, -): C_{\geq 0} \rightarrow \mathrm{An}$, showing that P is projective in $C_{\geq 0}$. $\square$

Warning 6.4.19. It is not reasonable to ask for P to be projective in the stable ∞ -category C itself, since the only projective objects in C are the zero objects. To see this, assume that P is a projective object of C . By Lemma 6.4.17, the shift $N[1] \cong 0 \sqcup_N 0$ of any object $N \in C$ is the geometric realization of the simplicial object with $N^{\oplus n}$ in degree n . We then obtain equivalences

$$\mathrm{Hom}_C(P[-1], N) \simeq \mathrm{Hom}_C(P, N[1]) \simeq |[n] \mapsto \mathrm{Hom}_C(P, N^{\oplus n})|.$$

Since the right-hand side is connected, we see that $\pi_0 \mathrm{Hom}_C(P[-1], N) = 0$. Replacing N by $N[-k]$ for all $k \in \mathbb{N}$ then gives that $\pi_k \mathrm{Hom}_C(P[-1], N) = 0$ for all k , hence $\mathrm{Hom}_C(P[-1], N) = 0$, showing that P is the zero object.

6.5 Flat objects and Tor-groups

In classical algebra, flat modules form another important class of ‘well-behaved’ modules. An R -module N is called *flat* if the tensor product functor $- \otimes_R N$ is exact: short exact sequences of R -modules remain exact after tensoring with N . While projective modules are always flat, the converse does not hold in general. For example, $\mathbb{Q}$ is a flat $\mathbb{Z}$ -module but not projective.

Just as the notion of projective objects generalizes to t -projective objects in a stable ∞ -category with t -structure, the notion of flat objects generalizes to the following setting.

Definition 6.5.1. Let C , D , and E be stable ∞ -categories equipped with t -structures. Let $- \otimes -: C \times D \rightarrow E$ be a functor that is exact in each variable separately, and assume it restricts to $C_{\geq 0} \times D_{\geq 0} \rightarrow E_{\geq 0}$, i.e. the tensor product of connective objects is connective.

A connective object $N \in D_{\geq 0}$ is called *t -flat* (with respect to $\otimes$) if the functor $(-) \otimes N: C \rightarrow E$ is t -exact. Similarly, $M \in C_{\geq 0}$ is t -flat if the functor $M \otimes (-): D \rightarrow E$ is t -exact.

Remark 6.5.2. By assumption, $- \otimes N$ is right t-exact, so N is t-flat if and only if it is additionally left t-exact.

Remark 6.5.3 (Motivation). The three main examples we have in mind for this abstract setup are:

- (1) The tensor product $\text{Mod}_R \times \text{Mod}_R \xrightarrow{\otimes_R} \text{Mod}_R$ for a commutative ring spectrum R .
- (2) More generally the relative tensor product $\text{RMod}_R \times \text{LMod}_R \xrightarrow{\otimes_R} \text{Sp}$ for an associative ring spectrum R .
- (3) The derived tensor product $D^-(\mathcal{A}) \times D^-(\mathcal{A}) \xrightarrow{\otimes^L} D^-(\mathcal{A})$ for a symmetric monoidal abelian category $\mathcal{A}$ with enough projectives (constructed in Section 6.6).

We work in this generality for two reasons. First, it makes clear that the notion of flatness does not depend on the intricate coherences involved in the definition of Mod_R and its tensor product, discussed in Part II of this book. Second, it provides a single formulation that covers all relevant examples.

Exercise 6.5.4. Show that the t-flat objects in C and D are closed under finite direct sums and retracts.

The following exercise highlights the structural analogy between t-projectivity and t-flatness.

Exercise 6.5.5. Let C be a stable ∞ -category with a t-structure. Setting $D := C^{\text{op}}$ and $E := \text{Sp}^{\text{op}}$, the hom spectrum functor in C takes the form $\text{hom}_C : C \times C^{\text{op}} \rightarrow \text{Sp}^{\text{op}}$.

- (1) Show that this functor satisfies the conditions of Definition 6.5.1: For $X \in C_{\geq 0}$ and $Y \in C_{\leq 0}$ we have $\text{hom}_C(X, Y) \in \text{Sp}_{\leq 0}$.
- (2) Show that an object $P \in C$ is t-flat with respect to hom_C if and only if it is t-projective.

Just as the homotopy groups of mapping spectra give rise to Ext-groups, the homotopy groups of tensor products give rise to Tor-groups.

Definition 6.5.6 (Tor-groups). In the situation of Definition 6.5.1, for objects $M \in C$ and $N \in D$, we define the n -th Tor-group to be

$$\text{Tor}_n^{\otimes}(M, N) := \pi_n(M \otimes N) \in E^{\vee}.$$

When $M \in C^{\heartsuit}$ and $N \in D^{\heartsuit}$ both lie in the heart, we also define $M \otimes^{\heartsuit} N := \text{Tor}_0^{\otimes}(M, N) = \pi_0(M \otimes N)$, resulting in a functor

$$- \otimes^{\heartsuit} -: C^{\heartsuit} \times D^{\heartsuit} \rightarrow E^{\heartsuit}.$$

Since $M \otimes N \in E_{\geq 0}$, this agrees with $\tau_{\leq 0}(M \otimes N)$ and it follows that this functor is right exact in both variables.

Observation 6.5.7 (Tor long exact sequence). Every exact sequence $N' \rightarrow N \rightarrow N''$ in D induces an exact sequence $M \otimes N' \rightarrow M \otimes N \rightarrow M \otimes N''$ in E (since $M \otimes (-)$ is exact), and thus a long exact sequence of homotopy groups of the form

$$\begin{aligned} \cdots \rightarrow \text{Tor}_{n+1}^{\otimes}(M, N'') \rightarrow \text{Tor}_n^{\otimes}(M, N'') \rightarrow \text{Tor}_n^{\otimes}(M, N') \rightarrow \text{Tor}_n^{\otimes}(M, N) \\ \rightarrow \text{Tor}_n^{\otimes}(M, N'') \rightarrow \text{Tor}_{n-1}^{\otimes}(M, N'') \rightarrow \cdots \end{aligned}$$

As a special case, given an object $N \in D^\heartsuit$ and a short exact sequence $M' \hookrightarrow M \twoheadrightarrow M''$ in $C^\heartsuit$, the negative Tor-groups vanish, giving a one-sided long exact sequence of the form

$$\begin{aligned} \cdots \rightarrow \mathrm{Tor}_1^\otimes(M, N) \rightarrow \mathrm{Tor}_1^\otimes(M'', N) \rightarrow \mathrm{Tor}_0^\otimes(M', N) \\ \rightarrow \mathrm{Tor}_0^\otimes(M, N) \rightarrow \mathrm{Tor}_0^\otimes(M'', N) \rightarrow 0. \end{aligned}$$

The same remarks apply to the other variable.

Exercise 6.5.8. The derived ∞ -category $D(\mathbb{Z})$ carries a symmetric monoidal structure $\otimes^L$ with unit $\mathbb{Z}$; see Section 6.6. For an abelian group M , use the short exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z} \rightarrow 0$$

to show that $\mathrm{Tor}_1^{\otimes^L}(M, \mathbb{Z}/n\mathbb{Z})$ is the subgroup of M consisting of the n -torsion elements, i.e., those $x \in M$ satisfying $nx = 0$.

We now give several equivalent characterizations of t-flatness, analogous to Proposition 6.4.11 for t-projective objects.

Proposition 6.5.9. *For a connective object $N \in D_{\geq 0}$, the following conditions are equivalent:*

- (1) *The object N is t-flat;*
- (2) *For every $M \in C_{\leq 0}$ and every $i > 0$, the object $\mathrm{Tor}_i^\otimes(M, N)$ is zero;*
- (3) *For every $M \in C_{\leq 0}$, the object $\mathrm{Tor}_1^\otimes(M, N)$ is zero;*

Assume moreover that C is right complete, that the functor $(-) \otimes N$ preserves sequential colimits, and that $E_{\leq 0}$ is closed under sequential colimits. Then these conditions are further equivalent to:

- (4) *For every $M \in C^\heartsuit$ and every $i > 0$, the object $\mathrm{Tor}_i^\otimes(M, N)$ is zero.*

Proof. The functor $(-) \otimes N$ is right t-exact by assumption, so N is t-flat if and only if it sends $C_{\leq 0}$ into $E_{\leq 0}$. This is precisely condition (2). Conditions (2) and (3) are equivalent because, for $i > 0$,

$$\mathrm{Tor}_i^\otimes(M, N) \cong \mathrm{Tor}_1^\otimes(M[1-i], N),$$

and $M[1-i]$ is still coconnective.

It remains to compare with condition (4). For $M \in C_{\leq 0}$, each truncation $\tau_{\geq -n}M$ is obtained by finitely many extensions from shifts $(\pi_{-k}M)[-k]$ of objects of the heart. Condition (4) therefore implies that $(\tau_{\geq -n}M) \otimes N$ is coconnective. Right completeness and preservation of sequential colimits identify $M \otimes N$ with the colimit of these objects, which is coconnective because $E_{\leq 0}$ is closed under sequential colimits. Thus (4) implies (2); the converse is immediate. $\square$

In practice, we may often reduce to the case $i = 1$ in condition (4). This is an instance of the following lemma:

Lemma 6.5.10. *Assume that C is right complete, that $E_{\leq 0}$ is closed under sequential colimits, that $(-) \otimes N$ preserves sequential colimits, and that every object $M \in C^\heartsuit$ admits an epimorphism $P \twoheadrightarrow M$ from a t-flat object P . Then an object $N \in D^\heartsuit$ is t-flat if and only if $\mathrm{Tor}_1^\otimes(M, N) = 0$ for every $M \in C^\heartsuit$.*

Proof. The ‘only if’ is clear from Proposition 6.5.9. Conversely, we show by induction on i that $\mathrm{Tor}_i^\otimes(M, N) = 0$ for all $M \in C^\heartsuit$ and all $i > 0$. The case $i = 1$ holds by assumption. Assuming the claim in degree $i - 1$, consider an epimorphism $P \twoheadrightarrow M$ with P t-flat, and let M' be its kernel. Then the long exact sequence of Tor-groups contains

$$\cdots \rightarrow \mathrm{Tor}_i^\otimes(P, N) \rightarrow \mathrm{Tor}_i^\otimes(M, N) \rightarrow \mathrm{Tor}_{i-1}^\otimes(M', N) \rightarrow \mathrm{Tor}_{i-1}^\otimes(P, N) \rightarrow \cdots$$

The first term vanishes because P is t-flat, and the third vanishes by the induction hypothesis applied to M' . Hence $\mathrm{Tor}_i^\otimes(M, N) = 0$, and Proposition 6.5.9 shows that N is t-flat. $\square$

As a consequence of Proposition 6.5.9, we may compute Tor-groups in terms of flat resolutions.

Definition 6.5.11. Let $N \in D^\heartsuit$ be an object in the heart of D . A *flat resolution* of N is a long exact sequence

$$\cdots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \twoheadrightarrow N \rightarrow 0$$

in $D^\heartsuit$ such that each object F_n is t-flat.

Proposition 6.5.12 (Tor-groups via flat resolutions). *Let $F_\bullet$ be a flat resolution of $N \in D^\heartsuit$. Then for every object $M \in C^\heartsuit$ and every $n \in \mathbb{Z}$ there is an isomorphism*

$$\mathrm{Tor}_n^\otimes(M, N) \cong H_n(M \otimes^\heartsuit F_\bullet)$$

between the n -th Tor-group and the n -th homology group of the chain complex $M \otimes^\heartsuit F_\bullet$ in $E^\heartsuit$.

Proof. The proof is entirely dual to Proposition 6.4.14; we will therefore be rather brief. Set $N_{-1} := N$. The flat resolution splits up into short exact sequences of the form

$$0 \rightarrow N_n \hookrightarrow F_n \twoheadrightarrow N_{n-1} \rightarrow 0.$$

This provides a long exact sequence on Tor-groups by Observation 6.5.7, in which the higher Tor-groups of F_n vanish by Proposition 6.5.9. Using that $M \otimes^\heartsuit -: D^\heartsuit \rightarrow E^\heartsuit$ is right exact, one obtains isomorphisms

$$M \otimes^\heartsuit N_{k-1} \cong \mathrm{coker}(M \otimes^\heartsuit F_{k+1} \rightarrow M \otimes^\heartsuit F_k)$$

for all $k \geq 0$, which in turn give isomorphisms

$$\mathrm{Tor}_1^\otimes(M, N_{k-1}) \cong \frac{\ker(M \otimes^\heartsuit F_{k+1} \rightarrow M \otimes^\heartsuit F_k)}{\mathrm{im}(M \otimes^\heartsuit F_{k+2} \rightarrow M \otimes^\heartsuit F_{k+1})} = H_{k+1}(M \otimes^\heartsuit F_\bullet)$$

for all $k \geq 0$. The preceding cokernel formula for $k = 0$ gives the claim in degree zero. Finally, one can inductively show that $\mathrm{Tor}_n^\otimes(M, N) \cong \mathrm{Tor}_1^\otimes(M, N_{n-2})$ for all $n \geq 1$. Setting $k = n - 1$ in the previous isomorphism thus gives

$$\mathrm{Tor}_n^\otimes(M, N) \cong H_n(M \otimes^\heartsuit F_\bullet)$$

for all $n \geq 1$, finishing the proof. $\square$

6.6 Derived functors

A recurring theme of this chapter is the interplay between the stable and the abelian setting. Exactness is better behaved in a stable ∞ -category, while explicit computations are usually easier in an abelian category, and the derived ∞ -category $D(\mathcal{A})$ relates the two: it is stable, and it contains $\mathcal{A}$ as its heart. This relation may also be used to study functors on $\mathcal{A}$ itself. Consider a right t-exact functor $G: D(\mathcal{A}) \rightarrow D(\mathcal{B})$ between derived categories. Given a short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

in $\mathcal{A}$, we showed in Corollary 6.3.16 that this induces an exact sequence $A \rightarrow B \rightarrow C$ in $D(\mathcal{A})$. Applying G , we obtain an exact sequence $G(A) \rightarrow G(B) \rightarrow G(C)$ in $D(\mathcal{B})$, which in turn gives rise to a long exact sequence of homology groups:

$$\cdots \rightarrow G_1(B) \rightarrow G_1(C) \rightarrow G_0(A) \rightarrow G_0(B) \rightarrow G_0(C) \rightarrow 0,$$

where $G_n := H_n \circ G: \mathcal{A} \rightarrow \mathcal{B}$. In particular, the functor $G_0: \mathcal{A} \rightarrow \mathcal{B}$ is right exact, and the long exact sequence provides a powerful computational tool for understanding it.

The goal of *derived functors* is to reverse this process: given a right exact functor $F: \mathcal{A} \rightarrow \mathcal{B}$, can we obtain it as the zeroth homology of some right t-exact functor $D(\mathcal{A}) \rightarrow D(\mathcal{B})$? One might try to obtain such an extension by simply applying F degreewise to chain complexes. However, this naive approach fails: the degreewise extension $\text{Ch}(F): \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{B})$ does not preserve quasi-isomorphisms unless F is exact. Instead, we could ask: does there exist a universal approximation of $\text{Ch}(F)$ from the left by a functor that descends to $D(\mathcal{A}) \rightarrow D(\mathcal{B})$?

If $\mathcal{A}$ has enough projectives, we will see that such an approximation exists, at least at the level of bounded below complexes $\mathbf{L}F: D^-(\mathcal{A}) \rightarrow D^-(\mathcal{B})$. It may be computed on an arbitrary bounded below complex C by first replacing C with a quasi-isomorphic projective complex P and then applying F : we have $\mathbf{L}F(C) \simeq F(P)$. Dually, a left exact functor gives rise to right derived functors.

6.6.1 Abstract derived functors

We begin by developing the abstract theory of derived functors in terms of universal properties. The key observation is that a derived functor should be a ‘best approximation’ to a given functor by one that factors through a localization. Our discussion here is inspired by [DHKS04, Sections 39-41], [Rie14, Chapter 2] and [Cis19, Section 7.5].

Definition 6.6.1. Let C be an ∞ -category with a collection of morphisms W , and let $\gamma: C \rightarrow C[W^{-1}]$ be the localization functor. Let $F: C \rightarrow D$ be a functor. A functor $\mathbf{L}F: C[W^{-1}] \rightarrow D$ is called a *total left derived functor of F* if it comes equipped with a natural transformation $\alpha: \mathbf{L}F \circ \gamma \Rightarrow F$ of functors $C \rightarrow D$ that exhibits $\mathbf{L}F$ as a right Kan extension of F along γ :

$$\begin{array}{ccc} C & \xrightarrow{F} & D \\ \searrow \gamma & \Uparrow \alpha & \nearrow \mathbf{L}F \\ & C[W^{-1}] & \end{array}$$

Note that the functor $\mathbf{L}F$ is unique whenever it exists.

Dually, a *total right derived functor* of F is a total left derived functor of $F^{\text{op}}: C^{\text{op}} \rightarrow D^{\text{op}}$, i.e. a functor $\mathbf{R}F: C[W^{-1}] \rightarrow D$ equipped with $\beta: F \Rightarrow \mathbf{R}F \circ \gamma$ which exhibits $\mathbf{R}F$ as a left Kan extension of F along γ .

Observation 6.6.2. A priori, the condition on $\mathbf{L}F$ to be a right Kan extension demands that for every functor $H: C[W^{-1}] \rightarrow D$, the composite

$$\text{Nat}(H, \mathbf{L}F) \xrightarrow{\gamma^*} \text{Nat}(H \circ \gamma, \mathbf{L}F \circ \gamma) \xrightarrow{\alpha \circ -} \text{Nat}(H \circ \gamma, F)$$

is an equivalence. But since γ^* is always an equivalence by definition of localizations, this is equivalent to asking the second map to be an equivalence. We conclude that a transformation $\alpha: \mathbf{L}F \circ \gamma \Rightarrow F$ is a total left derived functor of F if and only if for every functor $G: C \rightarrow D$ that inverts all maps in W , the map

$$\alpha \circ -: \text{Nat}(G, \mathbf{L}F \circ \gamma) \rightarrow \text{Nat}(G, F)$$

is an equivalence.

In practice, total left derived functors can often be computed using *cofibrant replacements*. The idea is that while F may not preserve weak equivalences in general, it often does so when restricted to some full subcategory of cofibrant objects. The key ingredient then is the fact that cofibrant objects compute the localization: see Corollary 2.2.9. This turns out to be enough to produce the derived functor, which is the content of the following result of Cisinski [Cis19]. We will use it as a black box.

Proposition 6.6.3 (Cisinski [Cis19, p. 7.5.25]). *Let (C, W, I) be an ∞ -category with weak equivalences and cofibrations, let $i: C_c \hookrightarrow C$ denote the full subcategory of cofibrant objects, and let $F: C \rightarrow D$ be a functor such that the restriction $F|_{C_c}: C_c \rightarrow D$ inverts the weak equivalences between cofibrant objects. Then F admits a total left derived functor*

$$\mathbf{L}F: C[W^{-1}] \rightarrow D.$$

It is uniquely characterized by the existence of an isomorphism $\mathbf{L}F \circ \gamma \circ i \simeq F \circ i$. In particular, if $q: Q \rightarrow X$ is any weak equivalence with Q cofibrant, then

$$\mathbf{L}F(\gamma X) \simeq F(Q).$$

Moreover, the structure transformation $\mathbf{L}F \circ \gamma \Rightarrow F$ exhibits $\mathbf{L}F$ as an absolute right Kan extension: for every functor $H: D \rightarrow E$, the composite $H \circ \mathbf{L}F$ is a total left derived functor of $H \circ F$.

Proof. We give the construction of $\mathbf{L}F$, and refer to [Cis19, p. 7.5.25] for the verification that it has the required universal property. (The reference treats the dual situation of an ∞ -category with weak equivalences and fibrations, and constructs the total *right* derived functor of a functor which inverts the weak equivalences between *fibrant* objects.)

Write W_c for the weak equivalences between cofibrant objects. The restriction $F|_{C_c}$ descends to a functor $F_c: C_c[W_c^{-1}] \rightarrow D$. By part (1) of Corollary 2.2.9, the inclusion induces an equivalence $\bar{i}: C_c[W_c^{-1}] \xrightarrow{\simeq} C[W^{-1}]$, and we may thus define

$$\mathbf{L}F := F_c \circ \bar{i}^{-1}: C[W^{-1}] \rightarrow D$$

by choosing an inverse of $\bar{i}$. By construction, the restriction of $\mathbf{L}F$ to the cofibrant objects is isomorphic to $F|_{C_c}$, that is, $\mathbf{L}F \circ \gamma \circ i \simeq F \circ i$. Since $\bar{i}$ is an equivalence, a functor out of $C[W^{-1}]$ is determined by its restriction along $\gamma \circ i$, which gives the asserted uniqueness. Finally, if $q: Q \rightarrow X$ is a weak equivalence with Q cofibrant, then $\gamma(q)$ is an isomorphism, and hence

$$\mathbf{L}F(\gamma X) \simeq \mathbf{L}F(\gamma Q) \simeq F(Q).$$

The construction of $\mathbf{L}F$ is also compatible with postcomposition: for every $H: D \rightarrow E$, the composite $H \circ \mathbf{L}F$ is obtained by applying the same construction to $H \circ F$. Its structure transformation is therefore again a right Kan extension by [Cis19, (7.5.25.3)], proving absoluteness. $\square$

Remark 6.6.4. Everything in the above proof is formal except for the statement being cited: that the functor $\mathbf{L}F$, which by construction only knows about the cofibrant objects, is a right Kan extension of F along γ . The latter is a statement about *all* objects of C , and it is here that the real work happens: one needs to know that Kan extensions along γ may be computed in terms of the cofibrant objects. This is [Cis19, Corollary 7.5.17], whose proof rests on the theory of finite direct diagrams and their Reedy fibrant replacements developed in [Cis19, Section 7.4].

Remark 6.6.5. The situation simplifies considerably when C admits a *functorial cofibrant replacement*: a functor $Q: C \rightarrow C_c$ together with a natural weak equivalence $q: i \circ Q \Rightarrow \text{id}_C$. Such a replacement exists as soon as (C, W, I) admits functorial factorizations, by factoring the map $\emptyset \rightarrow X$ functorially in X . One may then simply take $\mathbf{L}F \circ \gamma \simeq F \circ Q$, with structure map $F(q): F \circ Q \Rightarrow F$, and the universal property can be deduced directly from the naturality of q , without any of the machinery above. This is the route taken in [DHKS04, Sections 39-41] and [Rie14, Chapter 2], where such data is called a *left deformation*.

We have avoided this hypothesis because it is genuinely restrictive: while Top and $\text{Ch}(\mathcal{A})$ do admit functorial factorizations, the projective cofibration structure on bounded below chain complexes from Proposition 6.6.14 does not, at least not for an arbitrary abelian category $\mathcal{A}$ with enough projectives: a projective resolution is built from a choice of epimorphism $P \twoheadrightarrow A$ from a projective object for each $A \in \mathcal{A}$, and such choices cannot in general be made functorially. (For $\mathcal{A} = \text{Mod}_R$ one can of course take the free module on the underlying set of A .)

Exercise 6.6.6. Let $F: C \rightarrow D$ be a functor which inverts all morphisms in W . By the universal property of $C[W^{-1}]$, F factors through a functor $\mathbf{L}F: C[W^{-1}] \rightarrow D$. Show that $\mathbf{L}F$ is a total left derived functor of F .

Exercise 6.6.7 (Products of cofibration structures). Let (C_1, W_1, I_1) and (C_2, W_2, I_2) be ∞ -categories with weak equivalences and cofibrations, and equip the product $C_1 \times C_2$ with the componentwise weak equivalences and cofibrations.

- (1) Show that $(C_1 \times C_2, W_1 \times W_2, I_1 \times I_2)$ is again an ∞ -category with weak equivalences and cofibrations, whose cofibrant objects are precisely the pairs of cofibrant objects.
- (2) Deduce that a functor $F: C_1 \times C_2 \rightarrow D$ which inverts the weak equivalences between pairs of cofibrant objects admits a total left derived functor

$$\mathbf{L}F: C_1[W_1^{-1}] \times C_2[W_2^{-1}] \rightarrow D,$$

using that localizations commute with finite products of ∞ -categories.

This is how derived functors of several variables, such as the derived tensor product of Definition 6.6.24, are obtained.

6.6.2 Derived functors between derived categories

We now apply the abstract framework to construct derived functors between derived ∞ -categories. The key observation is that bounded below projective chain complexes play the role of cofibrant objects: quasi-isomorphisms between them are automatically chain homotopy equivalences, so any additive functor preserves such quasi-isomorphisms.

Definition 6.6.8 (Projective chain complex). Let $\mathcal{A}$ be an abelian category. We write $\mathrm{Ch}^-(\mathcal{A}) \subseteq \mathrm{Ch}(\mathcal{A})$ for the full subcategory of bounded below chain complexes, those $C_\bullet$ for which $C_n = 0$ for all sufficiently small n . A chain complex $P_\bullet \in \mathrm{Ch}(\mathcal{A})$ is called *projective* if each object P_n is projective in $\mathcal{A}$, in the sense of Section 6.4.¹⁴ We write $\mathrm{Ch}_{\mathrm{proj}}^-(\mathcal{A}) \subseteq \mathrm{Ch}^-(\mathcal{A})$ for the full subcategory of bounded below projective chain complexes.

Remark 6.6.9. The localization functor identifies

$$\mathrm{Ch}^-(\mathcal{A})[\{\text{quasi-isomorphisms}\}^{-1}] \xrightarrow{\sim} \mathrm{D}^-(\mathcal{A}),$$

where $\mathrm{D}^-(\mathcal{A})$ is the full subcategory of bounded below objects from Definition 6.3.33. Indeed, Proposition 6.1.21 identifies the localization of $\mathrm{Ch}(\mathcal{A})_{\geq m}$ with the full subcategory $\mathrm{D}(\mathcal{A})_{\geq m}$ of $\mathrm{D}(\mathcal{A})$ for every m . Taking the union over all m gives the displayed identification. We use it without further comment.

Lemma 6.6.10. *Let $P_\bullet$ be a bounded below projective chain complex with trivial homology, i.e., $H_n(P) = 0$ for all n . Then $P_\bullet$ is contractible: the identity $\mathrm{id}_{P_\bullet}$ is chain homotopic to zero.*

Proof. We construct a contracting homotopy $s_n: P_n \rightarrow P_{n+1}$ satisfying $d_{n+1}s_n + s_{n-1}d_n = \mathrm{id}_{P_n}$ by induction. Say $P_\bullet$ is concentrated in degrees $\geq m$; we set $s_n = 0$ for $n < m$.

For the base case $n = m$: Since $P_{m-1} = 0$, the differential $d_m: P_m \rightarrow P_{m-1}$ is zero. The condition $H_m(P) = 0$ then says that $\ker(d_m)/\mathrm{im}(d_{m+1}) = P_m/\mathrm{im}(d_{m+1}) = 0$, meaning $d_{m+1}: P_{m+1} \rightarrow P_m$ is an epimorphism. Since P_m is projective, the identity id_{P_m} lifts along this epimorphism to give $s_m: P_m \rightarrow P_{m+1}$ with $d_{m+1}s_m = \mathrm{id}_{P_m}$. This satisfies our requirement since $s_{m-1} = 0$.

For the inductive step, suppose we have constructed s_k for all $k < n$ satisfying the homotopy relation. Define

$$\alpha_n := \mathrm{id}_{P_n} - s_{n-1}d_n: P_n \rightarrow P_n.$$

We claim that $\mathrm{im}(\alpha_n) \subseteq \mathrm{im}(d_{n+1})$. Indeed, using the induction hypothesis $d_n s_{n-1} + s_{n-2}d_{n-1} = \mathrm{id}_{P_{n-1}}$, we compute

$$d_n \circ \alpha_n = d_n - d_n s_{n-1} d_n = d_n - (\mathrm{id}_{P_{n-1}} - s_{n-2} d_{n-1}) d_n = s_{n-2} d_{n-1} d_n = 0.$$

Thus $\mathrm{im}(\alpha_n) \subseteq \ker(d_n) = \mathrm{im}(d_{n+1})$, where the last equality uses $H_n(P) = 0$. Since P_n is projective and $d_{n+1}: P_{n+1} \rightarrow \mathrm{im}(d_{n+1})$ is an epimorphism, there exists $s_n: P_n \rightarrow P_{n+1}$ with $d_{n+1}s_n = \alpha_n = \mathrm{id}_{P_n} - s_{n-1}d_n$, as required. $\square$

¹⁴This is weaker than $P_\bullet$ being a projective object of the abelian category $\mathrm{Ch}(\mathcal{A})$: the latter are exactly the contractible complexes that are degreewise projective. Throughout, ‘projective chain complex’ always refers to the degreewise condition; some authors instead say *degreewise projective* or *complex of projectives*.

Proposition 6.6.11. *Let $P_\bullet$ and $Q_\bullet$ be bounded below projective chain complexes. Then any quasi-isomorphism $f: P_\bullet \rightarrow Q_\bullet$ is a chain homotopy equivalence.*

Proof. The mapping cone $\text{Cone}(f)$ fits into a short exact sequence

$$0 \rightarrow Q_\bullet \rightarrow \text{Cone}(f) \rightarrow P_\bullet[1] \rightarrow 0.$$

The associated long exact sequence shows that $\text{Cone}(f)$ is acyclic. It is again bounded below and projective, hence contractible by Lemma 6.6.10. The mapping-cone criterion of Exercise 6.2 now shows that f is a chain homotopy equivalence. $\square$

Corollary 6.6.12. *Let $F: \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{B})$ be a functor that preserves chain homotopies (for example, a functor induced by an additive functor $\mathcal{A} \rightarrow \mathcal{B}$). Then F preserves quasi-isomorphisms between bounded below projective chain complexes.* $\square$

We now assume that $\mathcal{A}$ has enough projectives, meaning that every object of $\mathcal{A}$ admits an epimorphism from a projective object. This is enough to construct projective resolutions one complex at a time.

Proposition 6.6.13 (Existence of projective resolutions). *Let $\mathcal{A}$ be an abelian category with enough projectives. Every bounded below chain complex $C_\bullet$ admits a degreewise epimorphic quasi-isomorphism*

$$q: P_\bullet \xrightarrow{\sim} C_\bullet$$

from a bounded below projective chain complex $P_\bullet$. Moreover, if $C_n = 0$ for all $n < m$, then $P_\bullet$ may be chosen with $P_n = 0$ for all $n < m$.

Proof. This is the standard bounded-below projective-replacement theorem. One construction takes the total complex of a Cartan–Eilenberg projective resolution; it produces a degreewise epimorphic quasi-isomorphism with the stated lower bound. See [Wei94, Theorem 10.4.8 and its proof]. $\square$

The projective complexes may now be characterized intrinsically as the cofibrant objects in a cofibration structure on bounded below chain complexes.

Proposition 6.6.14 (The projective cofibration structure). *Let $\mathcal{A}$ be an abelian category with enough projectives. Equip $\text{Ch}^-(\mathcal{A})$ with the quasi-isomorphisms as weak equivalences and the degreewise split monomorphisms with bounded below projective cokernel as cofibrations. This makes $\text{Ch}^-(\mathcal{A})$ into an ∞ -category with weak equivalences and cofibrations. Its cofibrant objects are precisely the bounded below projective chain complexes.*

Proof. The 2-out-of-3 property is clear. The stated cofibrations contain the isomorphisms and are closed under composition: the cokernel of a composite is a degreewise split extension of the two projective cokernels. They are closed under pushout, since a pushout is again degreewise split with the same cokernel. This also proves that pushouts of trivial cofibrations are weak equivalences, using Corollary 6.1.17.

It remains to verify the factorization axiom. Let $f: P_\bullet \rightarrow C_\bullet$ be a chain map with $P_\bullet$ cofibrant, i.e. bounded below projective. Choose a degreewise epimorphic projective resolution $q: Q_\bullet \xrightarrow{\sim} C_\bullet$ using Proposition 6.6.13. Since $P_\bullet$ is bounded below projective and q is a degreewise epimorphism with

acyclic kernel, the map f lifts along q : by Exercise 6.3 there exists a chain map $\tilde{f}: P_\bullet \rightarrow Q_\bullet$ with $q\tilde{f} = f$. Now consider the mapping-cylinder factorization of $\tilde{f}$ from the proof of Proposition 6.1.26:

$$P_\bullet \hookrightarrow \text{Cyl}(\tilde{f})_\bullet \xrightarrow{\sim} Q_\bullet.$$

The first map is degreewise the inclusion $P_n \hookrightarrow P_n \oplus P_{n-1} \oplus Q_n$, hence a degreewise split monomorphism whose cokernel is degreewise $P_{n-1} \oplus Q_n$ and thus bounded below projective; so it is a cofibration. The second map is a chain homotopy equivalence, and composing it with q therefore exhibits f as a cofibration followed by a quasi-isomorphism.

Finally, $0 \rightarrow P_\bullet$ is a cofibration precisely when $P_\bullet$ is bounded below projective. $\square$

Proposition 6.6.15. *Let $\mathcal{A}$ be an abelian category with enough projectives. Then the inclusion $\text{Ch}_{\text{proj}}^-(\mathcal{A}) \hookrightarrow \text{Ch}^-(\mathcal{A})$ induces an equivalence on localizations:*

$$\mathcal{K}_{\text{proj}}^-(\mathcal{A}) := \text{Ch}_{\text{proj}}^-(\mathcal{A})[\text{chain homotopy equivalences}^{-1}] \xrightarrow{\sim} \text{D}^-(\mathcal{A}).$$

Proof. By Proposition 6.6.14 and part (1) of Corollary 2.2.9, the inclusion of the cofibrant objects induces an equivalence after localizing at the quasi-isomorphisms. By Proposition 6.6.11, the quasi-isomorphisms between bounded below projective complexes are precisely the chain homotopy equivalences, which gives the displayed equivalence. $\square$

We can now construct left derived functors between derived categories.

Construction 6.6.16 (Derived functors between derived categories). Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories, assume that $\mathcal{A}$ has enough projectives, and let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor. Extending F degreewise gives a functor $\text{Ch}^-(F): \text{Ch}^-(\mathcal{A}) \rightarrow \text{Ch}^-(\mathcal{B})$ that preserves chain homotopies. By Corollary 6.6.12, the restriction

$$\text{Ch}^-(F)|_{\text{proj}}: \text{Ch}_{\text{proj}}^-(\mathcal{A}) \rightarrow \text{Ch}^-(\mathcal{B})$$

preserves quasi-isomorphisms. Composing with the localization $\gamma_{\mathcal{B}}: \text{Ch}^-(\mathcal{B}) \rightarrow \text{D}^-(\mathcal{B})$, we obtain a functor that inverts quasi-isomorphisms between the cofibrant objects of the projective cofibration structure on $\text{Ch}^-(\mathcal{A})$. By Proposition 6.6.3, it therefore admits a total left derived functor

$$\mathbf{L}F: \text{D}^-(\mathcal{A}) \rightarrow \text{D}^-(\mathcal{B})$$

If $q: P_\bullet \xrightarrow{\sim} C_\bullet$ is any projective resolution, then

$$\mathbf{L}F(C_\bullet) \simeq F(P_\bullet).$$

Exercise 6.6.17. Let $P_\bullet \rightarrow C_\bullet$ be a quasi-isomorphism in $\text{Ch}(\mathcal{A})_{\geq 0}$, with $P_\bullet$ projective. Show that $\mathbf{L}F(C_\bullet) \simeq F(P_\bullet)$ in $\text{D}^-(\mathcal{B})$. If F is right exact, conclude that $\pi_0 \mathbf{L}F(A) \cong F(A)$ for all $A \in \mathcal{A}$.

Lemma 6.6.18. *In the situation of Construction 6.6.16, the functor $\mathbf{L}F: \text{D}^-(\mathcal{A}) \rightarrow \text{D}^-(\mathcal{B})$ is exact and right t-exact.*

Proof. For right t-exactness, let $C_\bullet$ be concentrated in degrees ≥ 0 . The construction in Proposition 6.6.13 gives a projective resolution $P_\bullet \rightarrow C_\bullet$ which is also concentrated in degrees ≥ 0 . Hence $\mathbf{L}F(C_\bullet) \simeq F(P_\bullet)$ is connective.

For exactness, it suffices to show that $\mathbf{L}F$ preserves cofiber sequences. Every morphism in $\mathbf{D}^-(\mathcal{A})$ can be represented by a chain map $f: P_\bullet \rightarrow Q_\bullet$ between bounded below projective complexes (by first replacing both source and target with projective resolutions). The cofiber in $\mathbf{D}^-(\mathcal{A})$ is represented by the mapping cone $\text{Cone}(f)$, which is again a bounded below projective complex. Since F is additive, it preserves mapping cones: $F(\text{Cone}(f)) \cong \text{Cone}(F(f))$. We then compute

$$\mathbf{L}F(\text{cofib}(f)) \simeq F(\text{Cone}(f)) \cong \text{Cone}(F(f)) \simeq \text{cofib}(\mathbf{L}F(f)),$$

showing that $\mathbf{L}F$ preserves cofiber sequences. $\square$

Remark 6.6.19. Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be a right exact functor between abelian categories. For an object $A \in \mathcal{A}$, viewed as a complex concentrated in degree zero, we may define functors $\mathbf{L}^n F: \mathcal{A} \rightarrow \mathcal{B}$ by

$$\mathbf{L}^n F(A) := H_n(\mathbf{L}F(A))$$

for $n \geq 0$. They may be computed as the homology groups of $F(P_\bullet)$, where $P_\bullet \rightarrow A$ is any choice of projective resolution of A . Moreover, applying $\mathbf{L}F$ to a short exact sequence $0 \rightarrow A \hookrightarrow B \twoheadrightarrow C \rightarrow 0$ in $\mathcal{A}$ yields a cofiber sequence $\mathbf{L}F(A) \rightarrow \mathbf{L}F(B) \rightarrow \mathbf{L}F(C)$ in $\mathbf{D}^-(\mathcal{B})$ by Lemma 6.6.18; its long exact sequence on homology is the classical long exact sequence relating the derived functors $\mathbf{L}^n F$, and in particular the connecting homomorphisms of the latter arise precisely from the exactness of $\mathbf{L}F$. In classical treatments of homological algebra, one often merely considers the functors $\mathbf{L}^n F$ rather than the entire derived functor $\mathbf{L}F$.

Lemma 6.6.20. *Assume that $\mathcal{A}$ and $\mathcal{B}$ satisfy (AB3) and (AB4), and that $F: \mathcal{A} \rightarrow \mathcal{B}$ preserves arbitrary coproducts. Then $\mathbf{L}F: \mathbf{D}^-(\mathcal{A}) \rightarrow \mathbf{D}^-(\mathcal{B})$ preserves coproducts of uniformly bounded below families.*

Proof. A uniformly bounded below family of complexes has a coproduct in $\text{Ch}^-(\mathcal{A})$, formed degree-wise. The assumptions (AB3) and (AB4) imply that this coproduct preserves quasi-isomorphisms, and F commutes with it degreewise. The claim therefore follows by localization. $\square$

The construction of derived functors can be generalized by replacing the target $\mathbf{D}^-(\mathcal{B})$ with an arbitrary left complete t-structure on a stable ∞ -category C : any right exact functor $F: \mathcal{A} \rightarrow C^\heartsuit$ extends uniquely to a right t-exact functor $\mathbf{D}^-(\mathcal{A}) \rightarrow C$ which sends the projective objects of $\mathcal{A}$ into the heart of C . In fact, this property completely characterizes the bounded below derived category:

Theorem 6.6.21 (Universal property of the bounded below derived category). *Let $\mathcal{A}$ be an abelian category with enough projectives and let C be a stable ∞ -category with a left complete t-structure. Then restriction along the inclusion $\mathcal{A} \hookrightarrow \mathbf{D}^-(\mathcal{A})$ induces an equivalence of ∞ -categories*

$$\text{Fun}'(\mathbf{D}^-(\mathcal{A}), C) \xrightarrow{\simeq} \text{Fun}^{\text{rex}}(\mathcal{A}, C^\heartsuit),$$

where the left-hand side denotes the full subcategory of exact functors $\mathbf{D}^-(\mathcal{A}) \rightarrow C$ that are right t-exact and send projective objects of $\mathcal{A}$ into the heart $C^\heartsuit$.

Proof. We refer to Lurie [Lur17, Theorem 1.3.3.2]. $\square$

Given a right exact functor $F: \mathcal{A} \rightarrow C^\heartsuit$, its unique extension obtained by the theorem is denoted $\mathbf{L}F: \mathbf{D}^-(\mathcal{A}) \rightarrow C$, and referred to as the *left derived functor* of F . Note that when $C = \mathbf{D}^-(\mathcal{B})$ for some abelian category $\mathcal{B}$, this agrees with the functor $\mathbf{L}F$ from Construction 6.6.16 by uniqueness: the latter is exact and right t-exact by Lemma 6.6.18, and it sends a projective object $P \in \mathcal{A}$ to $F(P) \in \mathcal{B}$, which lies in the heart of $\mathbf{D}^-(\mathcal{B})$.

6.6.3 Derived tensor product and derived hom

We now apply the derived functor formalism to construct the derived tensor product and derived hom. Throughout this subsection, let $\mathcal{A}$ be a closed symmetric monoidal abelian category with enough projectives, with tensor product $\otimes: \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ and internal hom $[-, -]: \mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \mathcal{A}$. These conditions are satisfied, for example, when $\mathcal{A} = \text{RMod}_R$ for a commutative ring R .

Definition 6.6.22 (Tensor product of chain complexes). The *tensor product* of chain complexes $C_\bullet, C'_\bullet \in \text{Ch}(\mathcal{A})$ is the chain complex $(C \otimes C')_\bullet$ defined by

$$(C \otimes C')_n := \bigoplus_{p+q=n} C_p \otimes C'_q$$

whose differential acts on the summand $C_p \otimes C'_q$ as the sum $d^C \otimes \text{id}_{C'} + (-1)^p \text{id}_C \otimes d^{C'}$ of the two canonical composite maps. This makes $\text{Ch}(\mathcal{A})$ into a symmetric monoidal category. Moreover, tensor products preserve chain homotopies: if $f \simeq g: C_\bullet \rightarrow D_\bullet$ via a homotopy h , then $f \otimes \text{id}_{C'} \simeq g \otimes \text{id}_{C'}$ via $h \otimes \text{id}_{C'}$.

The tensor product of chain complexes does not preserve quasi-isomorphisms in general:

Example 6.6.23. Let $\mathcal{A} = \text{Ab}$ and consider the quasi-isomorphism $f: C_\bullet \rightarrow D_\bullet$ where $C_\bullet$ is the complex $\mathbb{Z} \xrightarrow{2} \mathbb{Z}$ concentrated in degrees 1 and 0, and $D_\bullet = \mathbb{Z}/2$ concentrated in degree 0. Then $\mathbb{Z}/2 \otimes f$ is the map

$$(\mathbb{Z}/2 \xrightarrow{0} \mathbb{Z}/2) \longrightarrow (0 \rightarrow \mathbb{Z}/2),$$

which is not a quasi-isomorphism: the left-hand side has $H_1 = \mathbb{Z}/2$, while the right-hand side has $H_1 = 0$.

Definition 6.6.24 (Derived tensor product). We define the *derived tensor product*

$$- \otimes^{\text{L}} -: \text{D}^-(\mathcal{A}) \times \text{D}^-(\mathcal{A}) \rightarrow \text{D}^-(\mathcal{A})$$

as the total left derived functor of the composite

$$\text{Ch}^-(\mathcal{A}) \times \text{Ch}^-(\mathcal{A}) \xrightarrow{- \otimes -} \text{Ch}^-(\mathcal{A}) \longrightarrow \text{D}^-(\mathcal{A}),$$

where the source is equipped with the product of the projective cofibration structures of Proposition 6.6.14; see Exercise 6.6.7. Indeed, on pairs of bounded below projective complexes, tensor product preserves quasi-isomorphisms, since these are chain homotopy equivalences and tensor product preserves chain homotopies in each variable. Consequently, if $P \rightarrow C$ and $Q \rightarrow D$ are any projective resolutions, then

$$C \otimes^{\text{L}} D \simeq P \otimes Q.$$

As in Lemma 6.6.18, this functor is exact in both variables, and restricts to $\text{D}(\mathcal{A})_{\geq 0} \times \text{D}(\mathcal{A})_{\geq 0} \rightarrow \text{D}(\mathcal{A})_{\geq 0}$. If $\mathcal{A}$ satisfies (AB3) and (AB4), then it also preserves coproducts of uniformly bounded below families in both variables.

For objects $M, N \in \mathcal{A}$, regarded as complexes concentrated in degree zero, we abbreviate

$$\text{Tor}_n^{\mathcal{A}}(M, N) := H_n(M \otimes^{\text{L}} N) = \text{Tor}_n^{\otimes^{\text{L}}}(M, N).$$

Remark 6.6.25. Suppose in addition that the monoidal unit of $\mathcal{A}$ is projective and that the tensor product of two projective objects is again projective. Then the derived tensor product equips $D^-(\mathcal{A})$ with a symmetric monoidal structure. The construction, including the coherent symmetric monoidal structure and the identification of its tensor product with $-\otimes^L -$, is given in Example 20.1.7.

In practice, it usually suffices to resolve only one of the two variables by a projective complex. This is a consequence of the following lemma.

An object $F \in \mathcal{A}$ is called *flat* if the functor $F \otimes (-): \mathcal{A} \rightarrow \mathcal{A}$ is exact.

Lemma 6.6.26. *Let $F_\bullet$ be a bounded below chain complex such that each object F_i is flat. Then the functor*

$$F_\bullet \otimes -: \text{Ch}^-(\mathcal{A}) \rightarrow \text{Ch}^-(\mathcal{A})$$

preserves quasi-isomorphisms.

Proof. Given a quasi-isomorphism $f: C \rightarrow D$, we need to show that the map $F_\bullet \otimes f$ is again a quasi-isomorphism. To this end, filter $F_\bullet$ by the subcomplexes $F^{\leq n}$, where $(F^{\leq n})_k = F_k$ for $k \leq n$ and 0 otherwise. We have degree-wise split short exact sequences $0 \rightarrow F^{\leq n-1} \rightarrow F^{\leq n} \rightarrow F_n[n] \rightarrow 0$. The map $F_n[n] \otimes f$ is a quasi-isomorphism since F_n is flat. By induction on n and the five lemma, each $F^{\leq n} \otimes f$ is a quasi-isomorphism. Since $F_\bullet, C_\bullet$ and $D_\bullet$ are all bounded below, the sequence $(F^{\leq n} \otimes C)_k$ is eventually constant with value $(F \otimes C)_k$ in every degree k , and similarly for D . The map $F_\bullet \otimes f$ is therefore a degree-wise colimit of the maps $F^{\leq n} \otimes f$ along which homology is eventually constant, and so it is a quasi-isomorphism. $\square$

Corollary 6.6.27. *Assume that all projective objects in $\mathcal{A}$ are flat.*

- (1) *The derived tensor product may be computed by resolving only one variable: for any projective resolutions $P \rightarrow C$ and $Q \rightarrow D$, there are isomorphisms*

$$C \otimes^L D \simeq P \otimes D \simeq C \otimes Q$$

in $D^-(\mathcal{A})$.

- (2) *Tor groups in $\mathcal{A}$ may be computed by resolving one variable: given objects $M, N \in \mathcal{A}$, we have*

$$\text{Tor}_n^{\mathcal{A}}(M, N) \cong H_n(P_\bullet \otimes N)$$

for any projective resolution $P_\bullet \rightarrow M$.

- (3) *For a flat object $F \in \mathcal{A}$, the complex $F[0]$ is a t-flat object of $D^-(\mathcal{A})$.*

Proof. Part (1) is immediate from Lemma 6.6.26, as P and Q are bounded below flat chain complexes. Part (2) then follows, as $\text{Tor}_n^{\mathcal{A}}(M, N)$ is defined as the n -th homology group of $M \otimes^L N$. Part (3) also follows, as the derived tensor $F \otimes^L -$ is quasi-isomorphic to the underived tensor $F \otimes -$, which preserves both connective and coconnective complexes. $\square$

We now turn to the derived internal hom.

For the explicit construction below, we assume in addition that countable products in $\mathcal{A}$ are exact, that tensor products of projective objects are projective, and that projective objects are flat. These

hypotheses ensure that bounded below projective complexes are K-projective and that tensoring with such a complex preserves quasi-isomorphisms. They hold, in particular, for modules over a commutative ring.

Definition 6.6.28 (Internal hom of chain complexes). The *internal hom* of chain complexes $C_\bullet, D_\bullet \in \text{Ch}(\mathcal{A})$ is the chain complex $\underline{\text{Hom}}(C, D)_\bullet$ defined by

$$\underline{\text{Hom}}(C, D)_n := \prod_{p \in \mathbb{Z}} [C_p, D_{p+n}]$$

with differential $(d\varphi)_p = d \circ \varphi_p - (-1)^n \varphi_{p-1} \circ d$ for $\varphi = (\varphi_p)_p \in \underline{\text{Hom}}(C, D)_n$.

Exercise 6.6.29. Show that there is a natural isomorphism of chain complexes

$$\underline{\text{Hom}}(A \otimes B, C) \cong \underline{\text{Hom}}(A, \underline{\text{Hom}}(B, C)),$$

and deduce the tensor-hom adjunction

$$\text{Hom}_{\text{Ch}(\mathcal{A})}(A \otimes B, C) \cong \text{Hom}_{\text{Ch}(\mathcal{A})}(A, \underline{\text{Hom}}(B, C)).$$

Lemma 6.6.30. If $P_\bullet$ is a bounded below projective chain complex, then the functor

$$\underline{\text{Hom}}(P, -): \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A})$$

preserves quasi-isomorphisms.

Proof. First observe that $[P_n, -]: \mathcal{A} \rightarrow \mathcal{A}$ is exact for every n . Indeed, exactness may be tested by mapping out of projective objects, and for projective objects Q we have

$$\text{Hom}_{\mathcal{A}}(Q, [P_n, -]) \cong \text{Hom}_{\mathcal{A}}(Q \otimes P_n, -),$$

which is exact because $Q \otimes P_n$ is projective. It follows that $\underline{\text{Hom}}(P, A)$ is acyclic whenever A is acyclic: filter P by its bounded truncations and apply the preceding exactness degree-wise. The transition maps between the resulting internal hom complexes are degree-wise split epimorphisms, so exactness of countable products allows us to pass to their inverse limit. Applying this to the cone of a quasi-isomorphism proves the claim. $\square$

Lemma 6.6.31. Let $P_\bullet$ be a bounded below projective chain complex and let $X_\bullet$ be any chain complex. Then the localization functor induces an isomorphism

$$\text{Hom}_{\mathcal{K}(\mathcal{A})}(P_\bullet, X_\bullet) \xrightarrow{\sim} \text{Hom}_{\text{D}(\mathcal{A})}(P_\bullet, X_\bullet).$$

Proof. By Lemma 6.6.30, the functor $\underline{\text{Hom}}(P_\bullet, -)$ preserves quasi-isomorphisms. Since

$$\pi_k \text{Hom}_{\mathcal{K}(\mathcal{A})}(P_\bullet, X_\bullet) \cong H_k(\underline{\text{Hom}}(P_\bullet, X_\bullet))$$

for every $k \geq 0$, the functor $\text{Hom}_{\mathcal{K}(\mathcal{A})}(P_\bullet, -)$ also preserves quasi-isomorphisms. Viewing $\text{D}(\mathcal{A})$ as the localization of $\mathcal{K}(\mathcal{A})$ at the quasi-isomorphisms, the claim follows from Lemma 6.2.10. $\square$

Definition 6.6.32 (Derived hom). Let $\text{Ch}_{\text{proj}}^-(\mathcal{A})$ denote the full subcategory of bounded below projective chain complexes. The internal hom restricts to a functor

$$\underline{\text{Hom}}(-, -): \text{Ch}_{\text{proj}}^-(\mathcal{A})^{\text{op}} \times \text{Ch}(\mathcal{A}) \rightarrow \text{Ch}(\mathcal{A}).$$

It preserves quasi-isomorphisms in both variables: in the first variable this follows from Proposition 6.6.11, while in the second it follows from Lemma 6.6.30. Consequently, localization and Proposition 6.6.15 produce a functor

$$\mathbf{R}\underline{\text{Hom}}(-, -): D^-(\mathcal{A})^{\text{op}} \times D(\mathcal{A}) \rightarrow D(\mathcal{A}).$$

For any projective resolution $P \rightarrow C$, it is computed by

$$\mathbf{R}\underline{\text{Hom}}(C, D) \simeq \underline{\text{Hom}}(P, D).$$

Just as the derived tensor product computes Tor-groups, the derived hom computes Ext-groups, which ties the present discussion back to Section 6.4:

Observation 6.6.33. Let R be a commutative ring and let $M, N \in \text{RMod}_R$, regarded as complexes concentrated in degree 0. Choosing a projective resolution $P_\bullet \rightarrow M$, the complex $\underline{\text{Hom}}(P, N)$ has $\underline{\text{Hom}}(P, N)_{-n} = [P_n, N]$ with the differential induced by that of $P_\bullet$; it is thus the cochain complex $\text{Hom}_R(P_\bullet, N)$, placed in non-positive degrees. Taking homology, we obtain

$$H_{-n}(\mathbf{R}\underline{\text{Hom}}_R(M, N)) \cong H^n(\text{Hom}_R(P_\bullet, N)) \cong \text{Ext}_R^n(M, N),$$

where the second isomorphism is Proposition 6.4.14. In particular $\mathbf{R}\underline{\text{Hom}}_R(M, N)$ is coconnective, with $H_0 = \text{Hom}_R(M, N)$.

The key relationship between the derived tensor product and derived hom is that they remain adjoint:

Proposition 6.6.34 (Derived tensor-hom adjunction). *For $C, D \in D^-(\mathcal{A})$ and $E \in D(\mathcal{A})$, there is a natural equivalence*

$$\text{Hom}_{D(\mathcal{A})}(C \otimes^{\mathbf{L}} D, E) \simeq \text{Hom}_{D(\mathcal{A})}(C, \mathbf{R}\underline{\text{Hom}}(D, E)).$$

Proof. Choose a chain complex representing E . By Proposition 6.6.15, the ∞ -category $D^-(\mathcal{A})$ is equivalent to the localization of $\text{Ch}_{\text{proj}}^-(\mathcal{A})$ at chain homotopy equivalences. It therefore suffices to establish the adjunction for bounded below projective complexes $P_\bullet$ and $Q_\bullet$. In this case, we have

$$P \otimes^{\mathbf{L}} Q \simeq P \otimes Q \quad \text{and} \quad \mathbf{R}\underline{\text{Hom}}(Q, E) \simeq \underline{\text{Hom}}(Q, E).$$

The complex $P \otimes Q$ is again bounded below projective. Using Lemma 6.6.31 and the internal tensor-hom adjunction of Exercise 6.6.29, we obtain

$$\text{Hom}_{D(\mathcal{A})}(P \otimes Q, E) \xrightarrow{\sim} \text{Hom}_{\mathcal{K}(\mathcal{A})}(P \otimes Q, E) \xrightarrow{\sim} \text{Hom}_{\mathcal{K}(\mathcal{A})}(P, \underline{\text{Hom}}(Q, E)) \xrightarrow{\sim} \text{Hom}_{D(\mathcal{A})}(P, \underline{\text{Hom}}(Q, E)),$$

as required. The middle isomorphism is an isomorphism of mapping anima because its k -th homotopy group is the k -th homology of the internal tensor-hom isomorphism from Exercise 6.6.29. $\square$

Corollary 6.6.35. *Assume in addition that $\mathcal{A}$ satisfies (AB3), (AB4), (AB3*), and (AB4*). For every $C \in D^-(\mathcal{A})$, the functor $\mathbf{R}\underline{\mathrm{Hom}}(C, -): D(\mathcal{A}) \rightarrow D(\mathcal{A})$ preserves all small limits. For every $E \in D(\mathcal{A})$, the functor $\mathbf{R}\underline{\mathrm{Hom}}(-, E)$ sends colimits of uniformly bounded below diagrams in $D^-(\mathcal{A})$ to limits.*

Proof. The functor $\mathbf{R}\underline{\mathrm{Hom}}(C, -) \simeq \underline{\mathrm{Hom}}(P, -)$, for P a projective resolution of C , is exact and satisfies $\underline{\mathrm{Hom}}(P, \prod_i E_i) \cong \prod_i \underline{\mathrm{Hom}}(P, E_i)$ degreeewise. It therefore preserves all limits.

In the first variable, the functor is exact and sends coproducts of uniformly bounded below families to products: if P^i is a projective resolution of C^i , chosen with a common lower bound, then $\bigoplus_i P^i$ is a projective resolution of $\bigoplus_i C^i$, and

$$\underline{\mathrm{Hom}}\left(\bigoplus_i P^i, E\right) \cong \prod_i \underline{\mathrm{Hom}}(P^i, E)$$

degreeewise. Together with exactness, this gives the assertion for uniformly bounded below colimits. $\square$

6.7 Künneth and universal coefficient theorems

The derived tensor product and derived internal hom now give concrete computational consequences for ordinary homology and cohomology. We first record the algebraic input over principal ideal domains, then apply it to the chain complexes of anima constructed in Section 6.2.

6.7.1 Tor and Ext over principal ideal domains

For complexes of modules over a principal ideal domain, the homology groups of the derived tensor product and derived hom admit particularly explicit descriptions in terms of Tor and Ext groups of the homology of the individual complexes. The key simplification is that projective resolutions over a PID have length at most one, which allows for elementary proofs of the Künneth formula and universal coefficient theorem.

Proposition 6.7.1. *Let R be a principal ideal domain. Then every R -module M admits a projective resolution of length at most one:*

$$0 \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0.$$

In particular, $\mathrm{Tor}_n^R(M, N) = 0$ and $\mathrm{Ext}_R^n(M, N) = 0$ for all R -modules M, N and all $n \geq 2$.

Proof. Every R -module M admits a surjection from a free module: take $P_0 := R^{(M)}$, the free module on the underlying set of M , with the canonical surjection $\varepsilon: P_0 \twoheadrightarrow M$. The kernel $P_1 := \ker(\varepsilon)$ is a submodule of the free module P_0 . Over a PID, every submodule of a free module is itself free, hence projective. This gives the desired resolution.

The vanishing of higher Tor and Ext groups follows from the formulas $\mathrm{Tor}_n^R(M, N) \cong H_n(P_\bullet \otimes_R N)$ and $\mathrm{Ext}_R^n(M, N) \cong H^n(\mathrm{Hom}_R(P_\bullet, N))$ of Corollary 6.6.27 and Proposition 6.4.14, which are both zero for $n \geq 2$ since $P_k = 0$ for $k \geq 2$. $\square$

Theorem 6.7.2 (Algebraic Künneth formula). *Let R be a principal ideal domain and let $A, B \in D^-(R)$ be bounded below complexes of R -modules. Then for each $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \rightarrow \bigoplus_{i+j=n} H_i(A) \otimes_R H_j(B) \rightarrow H_n(A \otimes_R^L B) \rightarrow \bigoplus_{i+j=n-1} \text{Tor}_1^R(H_i(A), H_j(B)) \rightarrow 0,$$

and this sequence splits (non-naturally).

Proof. We may represent the complexes A and B up to quasi-isomorphism by bounded below chain complexes C and C' , respectively, such that the R -modules C_i and C'_i are free for all $i \in \mathbb{Z}$. Then the derived tensor product $A \otimes_R^L B$ is represented by the algebraic tensor product $C \otimes_R C'$.

We first treat the special case where all the differentials in C are zero, so that $H_i(C) = C_i$ for all i . In this case, the differential on $C \otimes_R C'$ is given by $d(c \otimes c') = (-1)^i c \otimes dc'$ for $c \in C_i$, and the chain complex $C \otimes_R C'$ decomposes as the direct sum of the complexes $C_i[i] \otimes_R C'$. Since each C_i is free, the complex $C_i \otimes_R C'$ is a direct sum of copies of C' , and hence

$$H_n(C_i[i] \otimes_R C') \cong C_i \otimes_R H_{n-i}(C') = H_i(C) \otimes_R H_{n-i}(C').$$

Summing over i yields an isomorphism $H_n(C \otimes_R C') \cong \bigoplus_{i+j=n} H_i(C) \otimes_R H_j(C')$, which is the statement of the theorem since the Tor terms vanish ($H_i(C) = C_i$ being free, hence flat).

For the general case, let $Z_i \subseteq C_i$ and $B_i \subseteq C_i$ denote the cycles and boundaries in degree i , respectively. These give subchain complexes Z and B of C with trivial differentials. We have a short exact sequence of chain complexes

$$0 \rightarrow Z \rightarrow C \rightarrow B[1] \rightarrow 0$$

arising from the short exact sequences $0 \rightarrow Z_i \rightarrow C_i \rightarrow B_{i-1} \rightarrow 0$ in each degree. Each of these splits since B_{i-1} is free, being a submodule of the free module C_{i-1} . Because of this splitting, tensoring with C' preserves exactness, and we obtain a long exact sequence in homology:

$$\cdots \rightarrow H_n(Z \otimes_R C') \rightarrow H_n(C \otimes_R C') \rightarrow H_{n-1}(B \otimes_R C') \xrightarrow{i_{n-1}} H_{n-1}(Z \otimes_R C') \rightarrow \cdots$$

where $i_k: H_k(B \otimes_R C') \rightarrow H_k(Z \otimes_R C')$ is induced by the inclusion $B \hookrightarrow Z$.

Since Z and B are chain complexes with trivial differentials (and have free terms, being submodules of the free modules C_i), the special case treated before converts this to:

$$\cdots \xrightarrow{i_n} \bigoplus_{i+j=n} Z_i \otimes_R H_j(C') \rightarrow H_n(C \otimes_R C') \rightarrow \bigoplus_{i+j=n-1} B_i \otimes_R H_j(C') \xrightarrow{i_{n-1}} \bigoplus_{i+j=n-1} Z_i \otimes_R H_j(C') \rightarrow \cdots$$

The long exact sequence thus provides short exact sequences

$$0 \rightarrow \text{coker}(i_n) \rightarrow H_n(C \otimes_R C') \rightarrow \ker(i_{n-1}) \rightarrow 0.$$

To identify $\text{coker}(i_n)$ and $\ker(i_{n-1})$, observe that the short exact sequence $0 \rightarrow B_i \rightarrow Z_i \rightarrow H_i(C) \rightarrow 0$ functions as a free resolution of $H_i(C)$. The long exact sequence on Tor-groups between $H_i(C)$ and $H_j(C')$ takes the form

$$0 \rightarrow \text{Tor}_1^R(H_i(C), H_j(C')) \rightarrow B_i \otimes_R H_j(C') \rightarrow Z_i \otimes_R H_j(C') \rightarrow H_i(C) \otimes_R H_j(C') \rightarrow 0.$$

By taking the direct sum over all j , this results in isomorphisms

$$\operatorname{coker}(i_n) \cong \bigoplus_{i+j=n} H_i(C) \otimes_R H_j(C') \quad \text{and} \quad \ker(i_{n-1}) \cong \bigoplus_{i+j=n-1} \operatorname{Tor}_1^R(H_i(C), H_j(C')).$$

All maps used to construct this short exact sequence are functorial in chain maps. Through the projective-complex model of $D^-(R)$ from Proposition 6.6.15, the sequence is therefore natural in A and B .

Finally, we show that the short exact sequence splits. Since each C_i is free, the short exact sequence $0 \rightarrow Z_i \rightarrow C_i \rightarrow B_{i-1} \rightarrow 0$ splits, and the quotient maps $Z_i \rightarrow H_i(C)$ extend to homomorphisms $C_i \rightarrow H_i(C)$. We similarly obtain homomorphisms $C'_j \rightarrow H_j(C')$. Viewing the sequences of homology groups $H_i(C)$ and $H_j(C')$ as chain complexes with trivial differentials, we obtain chain maps $C \rightarrow H(C)$ and $C' \rightarrow H(C')$. Their tensor product is a chain map $C \otimes_R C' \rightarrow H(C) \otimes_R H(C')$. Since the differentials on the target are trivial, the induced map on homology

$$H_n(C \otimes_R C') \rightarrow H_n(H(C) \otimes_R H(C')) = \bigoplus_{i+j=n} H_i(C) \otimes_R H_j(C')$$

provides the desired splitting. $\square$

An analogous argument yields the universal coefficient theorem for the derived hom:

Theorem 6.7.3 (Algebraic universal coefficient theorem). *Let R be a principal ideal domain, let $A \in D^-(R)$, and let M be an R -module, regarded as a complex concentrated in degree zero. For each $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \rightarrow \operatorname{Ext}_R^1(H_{n-1}(A), M) \rightarrow H_{-n}(\mathbf{R}\underline{\operatorname{Hom}}_R(A, M[0])) \rightarrow \operatorname{Hom}_R(H_n(A), M) \rightarrow 0,$$

and this sequence splits non-naturally.

Proof. We refer to [Wei94, Theorem 3.6.5], applied to $\underline{\operatorname{Hom}}_R(C, M)$, which represents $\mathbf{R}\underline{\operatorname{Hom}}_R(A, M[0])$. $\square$

Example 6.7.4. For $R = \mathbb{Z}$, the Künneth formula specializes to the classical result for abelian groups. Given chain complexes A and B of abelian groups, the homology of the derived tensor product fits into the short exact sequence

$$0 \rightarrow \bigoplus_{i+j=n} H_i(A) \otimes_{\mathbb{Z}} H_j(B) \rightarrow H_n(A \otimes_{\mathbb{Z}}^{\mathbf{L}} B) \rightarrow \bigoplus_{i+j=n-1} \operatorname{Tor}_1^{\mathbb{Z}}(H_i(A), H_j(B)) \rightarrow 0.$$

Since $\operatorname{Tor}_1^{\mathbb{Z}}(M, N)$ consists of the common torsion of M and N (see Exercise 6.5.8), this shows that the homology of the derived tensor product is determined by the homology groups of A and B , up to an extension involving their torsion.

Example 6.7.5. Let R be a PID and let $A, B \in D^-(R)$. If all homology groups $H_i(A)$ are flat (equivalently, torsion-free), then

$$H_n(A \otimes_R^{\mathbf{L}} B) \cong \bigoplus_{i+j=n} H_i(A) \otimes_R H_j(B).$$

Example 6.7.6. Let R be a PID, let $A \in D^-(R)$, and let M be an R -module. If all homology groups $H_i(A)$ are projective (equivalently, free), then

$$H_{-n}(\mathbf{R}\underline{\operatorname{Hom}}_R(A, M[0])) \cong \operatorname{Hom}_R(H_n(A), M).$$

6.7.2 Künneth theorem

The Künneth theorem describes the homology of a product of spaces in terms of the homology of the factors. The key observation is that the chain complex of a product is the derived tensor product of the chain complexes of the factors.

Proposition 6.7.7. *Let R be a commutative ring. For animae $X, Y \in \mathbf{An}$, there is a natural equivalence*

$$C_*(X \times Y; R) \simeq C_*(X; R) \otimes_R^{\mathbf{L}} C_*(Y; R)$$

in $\mathbf{D}(R)$.

Proof. Chain complexes of animae are connective, so both sides are functors $\mathbf{An} \times \mathbf{An} \rightarrow \mathbf{D}(R)_{\geq 0}$, and the derived tensor product of Definition 6.6.24 is defined on them. It preserves colimits in each variable: it is exact and preserves coproducts of uniformly bounded below families, and every colimit of connective objects is of this form. Similarly, the functor $- \times -: \mathbf{An} \times \mathbf{An} \rightarrow \mathbf{An}$ preserves colimits in both variables, and $C_*(-; R)$ preserves colimits. It follows that both sides define functors $\mathbf{An} \times \mathbf{An} \rightarrow \mathbf{D}(R)_{\geq 0}$ preserving colimits in each variable. Both send (pt, pt) to $R[0]$, hence they are naturally equivalent by the universal property of $\mathbf{An}$. $\square$

Theorem 6.7.8 (Künneth theorem). *Let R be a principal ideal domain and let $X, Y \in \mathbf{An}$ be animae. For each $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \rightarrow \bigoplus_{i+j=n} H_i(X; R) \otimes_R H_j(Y; R) \rightarrow H_n(X \times Y; R) \rightarrow \bigoplus_{i+j=n-1} \text{Tor}_1^R(H_i(X; R), H_j(Y; R)) \rightarrow 0,$$

and this sequence splits, though not naturally.

Proof. By Proposition 6.7.7, we have $H_n(X \times Y; R) \cong H_n(C_*(X; R) \otimes_R^{\mathbf{L}} C_*(Y; R))$. The result now follows immediately from the algebraic Künneth formula (Theorem 6.7.2). $\square$

Corollary 6.7.9. *If R is a principal ideal domain and $H_i(X; R)$ is a free R -module for all i , then*

$$H_n(X \times Y; R) \cong \bigoplus_{i+j=n} H_i(X; R) \otimes_R H_j(Y; R)$$

for all $n \in \mathbb{Z}$ and all animae Y . $\square$

6.7.3 Universal coefficient theorems

The universal coefficient theorems express the (co)homology with coefficients in a module in terms of the integral (co)homology. We begin with the homological version.

Proposition 6.7.10. *For an anima $X \in \mathbf{An}$ and a bounded below chain complex $A \in \mathbf{D}^-(\mathbb{Z})$, there is a natural equivalence*

$$C_*(X; A) \simeq C_*(X; \mathbb{Z}) \otimes_{\mathbb{Z}}^{\mathbf{L}} A$$

in $\mathbf{D}^-(\mathbb{Z})$.

Proof. Say A is concentrated in degrees $\geq m$. Both $C_*(-; A)$ and $C_*(-; \mathbb{Z}) \otimes_{\mathbb{Z}}^L A$ then take values in $D(\mathbb{Z})_{\geq m}$, which is closed under colimits in $D(\mathbb{Z})$. Both preserve colimits: for the first this is the defining property from Construction 6.2.12, and for the second it follows as in Proposition 6.7.7, since $C_*(-; \mathbb{Z})$ is colimit-preserving with connective values and $-\otimes_{\mathbb{Z}}^L A$ is exact and preserves coproducts of uniformly bounded below families. Finally, both send pt to A . By the universal property of An , they are naturally equivalent. $\square$

Theorem 6.7.11 (Universal coefficient theorem for homology). *Let $X \in \text{An}$ be an anima and let A be an abelian group. For each $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \rightarrow H_n(X) \otimes_{\mathbb{Z}} A \rightarrow H_n(X; A) \rightarrow \text{Tor}_1^{\mathbb{Z}}(H_{n-1}(X), A) \rightarrow 0,$$

and this sequence splits, though not naturally.

Proof. By Proposition 6.7.10, we have $H_n(X; A) \cong H_n(C_*(X; \mathbb{Z}) \otimes_{\mathbb{Z}}^L A)$. Viewing A as a chain complex concentrated in degree 0, the statement is thus an instance of the algebraic Künneth formula (Theorem 6.7.2), using that $H_j(A) = 0$ for $j \neq 0$ and $H_0(A) = A$. $\square$

For cohomology, we use the internal hom in $D(\mathbb{Z})$ instead of the derived tensor product.

Proposition 6.7.12. *For an anima $X \in \text{An}$ and a chain complex $A \in D(\mathbb{Z})$, there is a natural equivalence*

$$C^*(X; A) \simeq \mathbf{R}\underline{\text{Hom}}_{\mathbb{Z}}(C_*(X; \mathbb{Z}), A)$$

in $D(\mathbb{Z})$.

Proof. The chain complexes $C_*(X; \mathbb{Z})$ are connective. Thus every diagram in the image of the colimit-preserving functor $C_*(-; \mathbb{Z}): \text{An} \rightarrow D(\mathbb{Z})_{\geq 0}$ is uniformly bounded below, and Corollary 6.6.35 shows that $\mathbf{R}\underline{\text{Hom}}_{\mathbb{Z}}(C_*(-; \mathbb{Z}), A)$ preserves limits as a functor on An^{op} . It sends pt to $\mathbf{R}\underline{\text{Hom}}_{\mathbb{Z}}(\mathbb{Z}, A) \simeq A$, so the universal property of An identifies it with $C^*(-; A)$. $\square$

Theorem 6.7.13 (Universal coefficient theorem for cohomology). *Let $X \in \text{An}$ be an anima and let A be an abelian group. For each $n \in \mathbb{Z}$, there is a natural short exact sequence*

$$0 \rightarrow \text{Ext}_{\mathbb{Z}}^1(H_{n-1}(X), A) \rightarrow H^n(X; A) \rightarrow \text{Hom}_{\mathbb{Z}}(H_n(X), A) \rightarrow 0,$$

and this sequence splits, though not naturally.

Proof. By Proposition 6.7.12, we have $H^n(X; A) \cong H_{-n}(\mathbf{R}\underline{\text{Hom}}_{\mathbb{Z}}(C_*(X; \mathbb{Z}), A))$. The result is therefore the case $R = \mathbb{Z}$, $M = A$ of Theorem 6.7.3. $\square$

Corollary 6.7.14. *If $H_{n-1}(X)$ is a free abelian group, then*

$$H^n(X; A) \cong \text{Hom}_{\mathbb{Z}}(H_n(X), A)$$

for all abelian groups A . $\square$

Exercises

Exercise 6.1 (Two-term complexes). For an integer $n \geq 1$, let $C(n)_\bullet$ be the chain complex of abelian groups with $\mathbb{Z}$ in degrees 1 and 0, differential

$$d_1: C(n)_1 = \mathbb{Z} \xrightarrow{n} C(n)_0 = \mathbb{Z},$$

and zero in all other degrees.

- (1) Compute the homology groups $H_k(C(n))$.
- (2) Show that the evident map $C(n)_\bullet \rightarrow \mathbb{Z}/n\mathbb{Z}[0]$ is a quasi-isomorphism.
- (3) Deduce that $C(n)_\bullet$ is a projective resolution of $\mathbb{Z}/n\mathbb{Z}$.

Exercise 6.2 (Mapping cones and quasi-isomorphisms). Let $f: C_\bullet \rightarrow D_\bullet$ be a chain map of chain complexes in an abelian category.

- (1) Verify that the mapping cone of Definition 6.1.14 fits into the short exact sequence

$$0 \rightarrow D_\bullet \rightarrow \text{Cone}(f) \rightarrow C_\bullet[1] \rightarrow 0.$$

- (2) Use the associated long exact sequence to show that f is a quasi-isomorphism if and only if $\text{Cone}(f)$ is acyclic (i.e. has trivial homology).
- (3) If $C_\bullet$ and $D_\bullet$ are bounded below complexes of projective objects, show that the same holds for $\text{Cone}(f)$.
- (4) Using that every acyclic bounded below chain complex of projective objects is chain homotopy contractible, deduce that a quasi-isomorphism $f: C_\bullet \rightarrow D_\bullet$ is a chain homotopy equivalence.

Exercise 6.3 (Lifting along a projective resolution). Let $\mathcal{A}$ be an abelian category, let $q: Q_\bullet \rightarrow C_\bullet$ be a degreewise epimorphic quasi-isomorphism between bounded below chain complexes, and let $P_\bullet$ be a bounded below projective chain complex. Write $K_\bullet := \ker(q)$, and write $Z_n(-)$ and $B_n(-)$ for the cycles and the boundaries in degree n .

- (1) Show that $K_\bullet$ is acyclic, and deduce that the differential restricts to an epimorphism $d: K_n \twoheadrightarrow Z_{n-1}(K)$.
- (2) Show that q restricts to an epimorphism $B_n(Q) \twoheadrightarrow B_n(C)$ on boundaries. (Hint: $B_n(C)$ is the image of $d \circ q_{n+1}$, since q_{n+1} is an epimorphism.)
- (3) Deduce that q restricts to an epimorphism $Z_n(Q) \twoheadrightarrow Z_n(C)$ on cycles. (Hint: compare the short exact sequences $0 \rightarrow B_n \rightarrow Z_n \rightarrow H_n \rightarrow 0$ for Q and for C , and use that q is a quasi-isomorphism.)
- (4) The differential $d: C_n \rightarrow C_{n-1}$ factors through the cycles, so we may form the pullback $X_n := C_n \times_{Z_{n-1}(C)} Z_{n-1}(Q)$; the maps q_n and d_n assemble into a map $\psi_n: Q_n \rightarrow X_n$. Show that ψ_n is an epimorphism. (Hint: by (3), the projection $X_n \rightarrow C_n$ is an epimorphism with kernel $Z_{n-1}(K)$. View ψ_n as a map of short exact sequences which is the identity on C_n and which on kernels is the map $d: K_n \rightarrow Z_{n-1}(K)$ from (1).)

- (5) Deduce, by induction on the degree, that every chain map $f: P_\bullet \rightarrow C_\bullet$ admits a lift $\tilde{f}: P_\bullet \rightarrow Q_\bullet$ with $q\tilde{f} = f$. Where is the boundedness of $P_\bullet$ used?
- (6) Show that any two such lifts are chain homotopic through chain homotopies h with $qh = 0$.

Exercise 6.4. Let C and D be stable ∞ -categories with t-structures, let $F: C \rightarrow D$ be an exact functor, and let

$$0 \rightarrow A \hookrightarrow B \twoheadrightarrow C' \rightarrow 0$$

be a short exact sequence in the heart $C^\heartsuit$. Construct a long exact sequence in $D^\heartsuit$ of the form

$$\cdots \rightarrow \pi_{n+1}F(C') \rightarrow \pi_n F(A) \rightarrow \pi_n F(B) \rightarrow \pi_n F(C') \rightarrow \pi_{n-1}F(A) \rightarrow \cdots$$

Show that if F is t-exact, this long exact sequence collapses to the original short exact sequence in the heart.

Exercise 6.5 (Eilenberg–MacLane objects). Let A be an abelian group, regarded as an object of $D(\mathbb{Z})^\heartsuit$.

- (1) Use Proposition 6.2.3 to show that HA has homotopy group A in degree 0 and no other homotopy groups.
- (2) Show that, for $n \geq 0$, the anima $K(A, n)$ has homotopy group A in degree n and no other homotopy groups.
- (3) Show that for $n \geq 1$ there is a natural isomorphism $\Omega K(A, n) \cong K(A, n-1)$.

Exercise 6.6 (Chains on spheres). Use the construction of ordinary chains from Construction 6.2.12.

- (1) Show that the reduced chain functor $\tilde{C}_*(-; \mathbb{Z}): \mathbf{An}_* \rightarrow D(\mathbb{Z})$ sends cofiber sequences of pointed anima to exact sequences in $D(\mathbb{Z})$.
- (2) Deduce that $\tilde{C}_*(\Sigma X; \mathbb{Z}) \cong \tilde{C}_*(X; \mathbb{Z})[1]$.
- (3) Compute $\tilde{C}_*(S^n; \mathbb{Z})$ and $C_*(S^n; \mathbb{Z})$ in $D(\mathbb{Z})$.

Exercise 6.7 (Extensions and addition in Ext^1). Let C be a stable ∞ -category with a t-structure, and let $M, N \in C^\heartsuit$.

- (1) Starting from a morphism $\varphi: M \rightarrow N[1]$, form its fiber and show that this gives an extension

$$0 \rightarrow N \rightarrow E_\varphi \rightarrow M \rightarrow 0$$

in $C^\heartsuit$.

- (2) Starting from an extension of M by N , construct the corresponding morphism $M \rightarrow N[1]$.
- (3) Check that these two constructions are inverse to each other on isomorphism classes.
- (4) Describe the addition law on extensions which corresponds to the abelian group structure on $\text{Ext}_C^1(M, N)$.

Exercise 6.8 (Cyclic Tor and Ext). Let A be an abelian group and let $n \geq 1$.

- (1) Use the projective resolution $0 \rightarrow \mathbb{Z} \xrightarrow{\cdot n} \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z} \rightarrow 0$ to compute $\mathrm{Tor}_i^{\mathbb{Z}}(\mathbb{Z}/n\mathbb{Z}, A)$ for all $i \geq 0$.
- (2) Use the same resolution to compute $\mathrm{Ext}_{\mathbb{Z}}^i(\mathbb{Z}/n\mathbb{Z}, A)$ for all $i \geq 0$.
- (3) Specialize to $A = \mathbb{Z}/m\mathbb{Z}$ and express the answer in terms of $\gcd(m, n)$.

Exercise 6.9 (A derived functor from a homotopy invariant functor). Let $F: \mathrm{Top} \rightarrow D$ be a functor to an ∞ -category D which sends homotopy equivalences of topological spaces to isomorphisms.

- (1) Using CW-complexes as cofibrant objects, explain how Proposition 6.6.3 produces a total left derived functor

$$\mathbf{L}F: \mathrm{An} \rightarrow D.$$

- (2) Show that if X is a CW-complex, then $\mathbf{L}F(\Pi_{\infty} X) \cong F(X)$.
- (3) Explain why, if F already sends weak homotopy equivalences to isomorphisms, then $\mathbf{L}F$ is just the functor induced by the universal property of the localization $\mathrm{Top} \rightarrow \mathrm{An}$.

For the next two exercises, you may use that $H_k(\mathbb{CP}^m) \cong \mathbb{Z}$ for $k = 0, 2, \dots, 2m$ and vanishes otherwise, while $H_0(\mathbb{RP}^2) \cong \mathbb{Z}$, $H_1(\mathbb{RP}^2) \cong \mathbb{Z}/2\mathbb{Z}$, and all other homology groups of $\mathbb{RP}^2$ vanish.

Exercise 6.10 (Künneth computations). Use Theorem 6.7.8 to compute the integral homology groups of the following products:

- (1) $S^m \times S^n$.
- (2) $\mathbb{CP}^m \times S^n$.
- (3) $\mathbb{RP}^2 \times S^1$.

In each case, indicate whether the $\mathrm{Tor}_1^{\mathbb{Z}}$ term vanishes and why.

Exercise 6.11 (Universal coefficient computations). Use Theorem 6.7.11 and Theorem 6.7.13.

- (1) Compute $H_*(\mathbb{RP}^2; \mathbb{Z}/2\mathbb{Z})$ from the integral homology of $\mathbb{RP}^2$.
- (2) Compute $H^*(\mathbb{RP}^2; \mathbb{Z})$.
- (3) Compute $H^*(\mathbb{RP}^2; \mathbb{Z}/2\mathbb{Z})$ and identify the contribution of the $\mathrm{Ext}_{\mathbb{Z}}^1$ term.

7 Localization and completion

An abelian group can often be understood by separating its behavior at different primes. Its p -localization inverts all primes other than p , and its p -completion records compatible information modulo the powers p^n . For finitely generated groups, the arithmetic fracture square recovers the original group from its p -completion and its p -inversion. All three operations have analogues for spectra, and the aim of this chapter is to construct them and prove their basic properties.

The right level of generality is that of a *Bousfield localization*: given a spectrum E , we invert the maps that are isomorphisms on E -homology, and thereby retain exactly those features of a spectrum that are ‘seen by’ E . Rationalization, p -localization and p -completion all arise this way, for suitable choices of E . Bousfield localizations are also the basic organizing device of chromatic homotopy theory, which we do not treat here.

Section 7.1 introduces the E -local and E -acyclic spectra and shows that the E -localization of a spectrum always exists. Section 7.2 treats inverting a set of primes; the homotopy groups localize algebraically, and rationally we obtain Serre’s theorem $\mathbb{S}_{\mathbb{Q}} \simeq H\mathbb{Q}$ and the resulting splitting of rational spectra. Section 7.3 treats p -completion, which is less well behaved: it is not given by a naive algebraic operation on homotopy groups, and instead Theorem 7.3.10 expresses $\pi_*(X_p^\wedge)$ through the Ext- and Hom-groups of Chapter 6. We close with the arithmetic fracture square, which reassembles a spectrum from its p -completions and its rationalization.

7.1 Bousfield localizations

Let E be a spectrum, taken to be fixed throughout this section. As discussed in Definition 4.4.21, this leads to a definition of the *E -homology groups* of an anima:

$$X \mapsto E_*(X) := \pi_*(E \otimes \mathbb{S}[X]).$$

For example, taking $E = HA$ the Eilenberg–MacLane spectrum of an abelian group A results in singular homology with coefficients in A , while taking $E = \mathbb{S}$ the sphere spectrum results in the stable homotopy groups.

The homology groups $E_*(X)$ will generally only see *some* aspects of the given anima X , and may completely ignore others. Consider for example the case of the projective space $X = \mathbb{R}P^2$. If we considered the Eilenberg–MacLane spectrum $E = H\mathbb{Q}$ of the rational numbers, its homology would give no information beyond that of a point: the reduced rational homology of $\mathbb{R}P^2$ is trivial. But taking $E = H\mathbb{Z}/2$ *does* capture interesting information about $\mathbb{R}P^2$. The point of Bousfield localizations is to focus precisely on those parts of an anima (or spectrum) which are ‘seen by’ E .

Definition 7.1.1. A morphism $f: X \rightarrow Y$ is called an *E-equivalence* if the induced map $E \otimes f: E \otimes X \rightarrow E \otimes Y$ is an isomorphism. We say that a spectrum X is *E-acyclic* if the map $X \rightarrow 0$ is an *E-equivalence*, i.e. if $E \otimes X$ is the zero spectrum.

Recall that a morphism of spectra is an isomorphism if and only if it induces isomorphisms on all homotopy groups. So if we write $E_*(X) := \pi_*(E \otimes X)$ for the *E-homology* of X , generalizing the definition for anima, we see that the *E-equivalences* are precisely those maps that induce isomorphisms on *E-homology*. Similarly, the *E-acyclic* spectra are those with trivial *E-homology*.

Example 7.1.2. For $E = H\mathbb{Q}$, the *E-equivalences* are known as the *rational equivalences*: those that induce isomorphisms on rational homology.

We will now introduce a class of spectra that ‘cannot distinguish between *E-equivalent* spectra’:

Definition 7.1.3. A spectrum L is called *E-local* if for every *E-equivalence* $f: X \rightarrow Y$, the induced map $f^*: \text{hom}(Y, L) \rightarrow \text{hom}(X, L)$ on mapping spectra is an isomorphism. We denote by

$$\text{Sp}_E \subseteq \text{Sp}$$

the full subcategory spanned by the *E-local* spectra.

Equivalently, it suffices to require that f induce an isomorphism $\text{Hom}_{\text{Sp}}(Y, L) \rightarrow \text{Hom}_{\text{Sp}}(X, L)$ on hom anima for every *E-equivalence* $f: X \rightarrow Y$. Indeed, every shift of an *E-equivalence* is again an *E-equivalence*, and applying this condition to all shifts of f detects all homotopy groups of the map of mapping spectra.

Exercise 7.1.4. Show the following basic properties of *E-local* spectra:

- (1) A spectrum L is *E-local* if and only if $\text{hom}(X, L) = 0$ for every *E-acyclic* spectrum X ;
- (2) Given an exact sequence $X \rightarrow Y \rightarrow Z$, if two of these three spectra are *E-local* then so is the third;
- (3) In particular, if L is *E-local* then so are all its shifts $L[n]$;
- (4) A limit $\lim_i L_i$ of *E-local* spectra is *E-local*;
- (5) A retract of an *E-local* spectrum is *E-local*.

Exercise 7.1.5. Show that for every spectrum Y , the mapping spectrum $\text{hom}(E, Y)$ is *E-local*. Using this, show that for a morphism $f: X \rightarrow X'$ of spectra the following conditions are equivalent:

- (1) The morphism $f: X \rightarrow X'$ is an *E-equivalence*;
- (2) The induced map $f^*: \text{hom}(X', L) \rightarrow \text{hom}(X, L)$ is an isomorphism for all *E-local* spectra L ;
- (3) The induced map $f^*: \text{Hom}_{\text{Sp}}(X', L) \rightarrow \text{Hom}_{\text{Sp}}(X, L)$ is an isomorphism for all *E-local* spectra L .

Hint: use that we have $\pi_n \text{hom}(X, Y) \cong \pi_0 \text{hom}(X, Y[n]) \cong \pi_0 \text{Hom}_{\text{Sp}}(X, Y[n])$, and similarly for X' .

Lemma 7.1.6 (*' E_* -Whitehead theorem'*). *Let $f: L \rightarrow L'$ be a morphism between E -local spectra. Then f is an isomorphism if and only if it is an E -equivalence.*

Proof. By passing to fibers, we may equivalently show that an E -local spectrum L is zero if and only if it is E -acyclic. One direction is clear. For the other direction, assume L is E -acyclic. Then by assumption we have $\text{hom}(L, L) = 0$, hence $\text{Hom}_{\text{Sp}}(L, L) = *$, hence the identity on L is the zero map. This implies that L itself is zero, as desired. $\square$

We will now talk about the universal way of approximating an arbitrary spectrum by an E -local one:

Definition 7.1.7. A morphism $l: X \rightarrow X'$ of spectra is called an *E -localization* if X' is E -local and the morphism l is an E -equivalence.

Corollary 7.1.8. *Let X' be an E -local spectrum. A morphism $l: X \rightarrow X'$ is an E -localization if and only if for every E -local spectrum Y the induced map*

$$l^*: \text{Hom}_{\text{Sp}}(X', Y) \rightarrow \text{Hom}_{\text{Sp}}(X, Y)$$

is an isomorphism.

Proof. This is immediate from Exercise 7.1.5. $\square$

Note that the conclusion of the corollary says that X' is a *left adjoint object* of X under the inclusion $\text{Sp}_E \hookrightarrow \text{Sp}$, in the sense of Definition 21.1.3. In particular, the E -localization of X is unique if it exists, and is then denoted by $L_E(X)$.

Theorem 7.1.9 (Bousfield [Bou79, Theorem 1.1]). *For every object X , there exists an E -localization $l: X \rightarrow L_E X$. In particular, the inclusion $\text{Sp}_E \hookrightarrow \text{Sp}$ admits a left adjoint*

$$L_E: \text{Sp} \rightarrow \text{Sp}_E$$

called the E -localization functor.

Proof. Let $\text{Sp}^{E\text{-acyc}} \subseteq \text{Sp}$ be the full subcategory of E -acyclic spectra. It is the kernel of the colimit-preserving functor

$$E \otimes -: \text{Sp} \rightarrow \text{Sp},$$

so it is presentable and its inclusion $i: \text{Sp}^{E\text{-acyc}} \hookrightarrow \text{Sp}$ preserves small colimits by Corollary 22.2.4. This kernel is stable, since $E \otimes -$ is exact. The adjoint functor theorem Theorem 22.2.5 therefore supplies a right adjoint Γ_E to i .

For a spectrum X , define $L_E X$ by the exact sequence

$$i\Gamma_E X \longrightarrow X \longrightarrow L_E X,$$

where the first map is the counit of the adjunction. Let A be an E -acyclic spectrum. Full faithfulness of i and the adjunction give isomorphisms

$$\text{Hom}_{\text{Sp}}(A, i\Gamma_E X) \xrightarrow{\sim} \text{Hom}_{\text{Sp}^{E\text{-acyc}}}(A, \Gamma_E X) \xrightarrow{\sim} \text{Hom}_{\text{Sp}}(A, X).$$

Their composite is the map induced by the counit. Mapping out of A carries the defining exact sequence for $L_E X$ to a fiber sequence of anima, so it follows that $\mathrm{Hom}_{\mathrm{Sp}}(A, L_E X) = *$. The same argument applies to every shift of A , which is again E -acyclic. Hence $\mathrm{hom}(A, L_E X) = 0$, and $L_E X$ is E -local by Exercise 7.1.4. On the other hand, $E \otimes i\Gamma_E X = 0$, so applying $E \otimes -$ to the exact sequence shows that $X \rightarrow L_E X$ is an E -equivalence. Thus it is an E -localization. $\square$

The proof packages the required size argument into the general theory of presentable categories. Bousfield's original construction instead builds the reflector explicitly by a transfinite cell-attachment process. The general notion of a Bousfield localization is discussed in Section 21.8.

7.2 Localization at a prime

In algebra, one commonly studies rings and modules through their *localizations*, in which one inverts certain non-zero elements in the ring. The terminology is inspired by the geometric perspective of this operation: it corresponds to studying the ‘local’ behavior of a variety/scheme around a point.

In the case of abelian groups, this operation corresponds to inverting collections of prime numbers. Given an abelian group A and a natural number n , there is a multiplication-by- n map $n: A \rightarrow A$, sending a to the n -fold sum na . Abelian groups for which this map is an isomorphism for certain n are usually easier to study than general abelian groups:

- If multiplication by n is an isomorphism for all $n > 0$, then A acquires a unique structure of a rational vector space, hence is simply a direct sum of copies of $\mathbb{Q}$.
- If for some prime p , multiplication by q is an isomorphism for all primes $q \neq p$, then A is called *p-local*. In particular, A does not admit q -torsion for any $q \neq p$.

More generally, for a set P of primes we write $A[P^{-1}]$ for the universal abelian group under A on which multiplication by every $p \in P$ is invertible. When P consists of all primes, this is the rationalization $A_{\mathbb{Q}}$ of A . When P consists of all primes except a single prime p , it is the p -localization $A_{(p)}$ of A . Some questions about A can be reduced to questions about $A_{\mathbb{Q}}$ or $A_{(p)}$.

The goal of this section is to introduce an analogous theory of localizations for spectra, and to exhibit it as a special case of the general formalism of Bousfield localizations introduced in the previous section.

Notation 7.2.1. Let C be a semiadditive ∞ -category. We saw in Proposition 5.3.24 that the hom anima $\mathrm{Hom}_C(X, Y)$ admits a canonical structure of a commutative monoid for all $X, Y \in C$. In particular, there is for any $n \in \mathbb{N}$ a multiplication-by- n map

$$n \cdot - : \mathrm{Hom}_C(X, Y) \rightarrow \mathrm{Hom}_C(X, Y),$$

given explicitly by sending a map $f: X \rightarrow Y$ to the composite

$$X \xrightarrow{\Delta} \bigoplus_n X \xrightarrow{\bigoplus_n f} \bigoplus_n Y \xrightarrow{\nabla} Y.$$

For $f = \mathrm{id}_X: X \rightarrow X$, we refer to this map as *multiplication by n on X* , and denote it by

$$n: X \rightarrow X.$$

Note that when $C = \mathbf{Ab}$ is the category of abelian groups, this precisely recovers the ordinary multiplication-by- n map.

We record the following compatibility because we will apply it both to spectra and to their homotopy groups.

Exercise 7.2.2. Show that any semiadditive functor $F: C \rightarrow D$ between semiadditive ∞ -categories preserves the multiplication-by- n maps.

We now specialize to spectra. Let $P \subseteq \mathbb{N}$ be a set of primes. We say a spectrum X is *P-divisible* if multiplication by p on X is an isomorphism for every $p \in P$. We write

$$\mathrm{Sp}[P^{-1}] \subseteq \mathrm{Sp}$$

for the full subcategory of P -divisible spectra.

Definition 7.2.3. Consider a spectrum X .

- For $n > 0$, we define $X[\frac{1}{n}]$ as the colimit

$$X[\frac{1}{n}] := \mathrm{colim}(X \xrightarrow{n} X \xrightarrow{n} X \xrightarrow{n} \dots) \in \mathrm{Sp}.$$

- If P consists of finitely many primes, we set $N := \prod_{p \in P} p$ and set

$$X[P^{-1}] := X[\frac{1}{N}] \in \mathrm{Sp}.$$

- If P consists of infinitely many primes, say $p_1, p_2, p_3, \dots$, we define $X[P^{-1}]$ as the colimit

$$X[P^{-1}] := \mathrm{colim}(X \xrightarrow{p_1} X \xrightarrow{p_1 p_2} X \xrightarrow{p_1 p_2 p_3} \dots) \in \mathrm{Sp}.$$

When P consists of all primes, we refer to $X[P^{-1}]$ as the *rationalization of X* and denote it by $X_{\mathbb{Q}}$.

When P consists of all primes except a single prime p , we refer to $X[P^{-1}]$ as the *p-localization of X* , and denote it by $X_{(p)}$.

Note that $X[P^{-1}]$ always comes with a canonical map $X \rightarrow X[P^{-1}]$, by including the first entry of the colimit diagram. Moreover, for every spectrum X , the functor $X \otimes -: \mathrm{Sp} \rightarrow \mathrm{Sp}$ preserves colimits, and so there is an isomorphism

$$X[P^{-1}] \cong X \otimes \mathbb{S}[P^{-1}].$$

Lemma 7.2.4. Let X be a spectrum. Then there is an isomorphism

$$\pi_*(X[P^{-1}]) \cong \pi_*(X)[P^{-1}].$$

Proof. In each case of the definition, $X[P^{-1}]$ is a filtered colimit whose transition maps are multiplication by products of primes in P . By Exercise 7.2.2, the functors $\pi_n: \mathrm{Sp} \rightarrow \mathbf{Ab}$ carry these to the corresponding multiplication maps of abelian groups. Since they preserve filtered colimits by Lemma 4.4.27, and since the resulting colimit of abelian groups has the universal property defining localization, we obtain

$$\pi_*(X[P^{-1}]) \cong \pi_*(X)[P^{-1}]. \quad \square$$

Corollary 7.2.5. *For every spectrum X , the spectrum $X[P^{-1}]$ is P -divisible.*

Proof. We need to show that for every prime $p \in P$, the map $\cdot p: X[P^{-1}] \rightarrow X[P^{-1}]$ is an isomorphism of spectra. We may check this on homotopy groups, where it becomes the statement that the map

$$p: \pi_*(X)[P^{-1}] \rightarrow \pi_*(X)[P^{-1}]$$

is an isomorphism. This holds by elementary properties of localization of abelian groups. $\square$

Recall that an abelian group A is called P -power torsion if every element of A is annihilated by a positive integer all of whose prime factors lie in P . It is not difficult to see that A is P -power torsion if and only if $A[P^{-1}] = 0$. Together with Lemma 7.2.4, this implies:

Corollary 7.2.6. *A spectrum X is $\mathbb{S}[P^{-1}]$ -acyclic if and only if the homotopy groups $\pi_*(X)$ are P -power torsion.* $\square$

We next show that the assignment $X \mapsto X[P^{-1}]$ is a reflection onto the subcategory of P -divisible spectra.

Proposition 7.2.7. *Let X be a spectrum and let Y be a P -divisible spectrum. Then precomposition with the canonical map $X \rightarrow X[P^{-1}]$ induces an isomorphism of mapping spectra*

$$\mathrm{hom}(X[P^{-1}], Y) \xrightarrow{\sim} \mathrm{hom}(X, Y).$$

In particular, the inclusion $\mathrm{Sp}[P^{-1}] \hookrightarrow \mathrm{Sp}$ admits a left adjoint given by

$$(-)[P^{-1}]: \mathrm{Sp} \rightarrow \mathrm{Sp}[P^{-1}].$$

Proof. The mapping spectrum $\mathrm{hom}(X[P^{-1}], Y)$ is the limit obtained by applying $\mathrm{hom}(-, Y)$ to the defining diagram for $X[P^{-1}]$. The transition maps in this limit are given by precomposition with the corresponding multiplication maps on X . By bilinearity of composition in spectra, these maps agree with postcomposition by the corresponding multiplication maps on Y .

All transition maps are induced by multiplication by products of primes in P on Y , hence are isomorphisms since Y is P -divisible. The limit is therefore isomorphic to the first term $\mathrm{hom}(X, Y)$.

Passing to underlying mapping anima and using Corollary 7.2.5 gives the claim. $\square$

The previous proposition establishes the functor $(-)[P^{-1}]: \mathrm{Sp} \rightarrow \mathrm{Sp}$ as the reflector onto the P -divisible spectra. We will now identify it with the E -localization functor for $E = \mathbb{S}[P^{-1}]$.

Lemma 7.2.8. *A spectrum X is P -divisible if and only if it is $\mathbb{S}[P^{-1}]$ -local.*

Proof. First assume X is P -divisible. To see that X is $\mathbb{S}[P^{-1}]$ -local, consider an $\mathbb{S}[P^{-1}]$ -acyclic spectrum Y . We must show that $\mathrm{hom}(Y, X) = 0$. This follows from the following sequence of isomorphisms:

$$\mathrm{hom}(Y, X) \cong \mathrm{hom}(Y[P^{-1}], X) \cong \mathrm{hom}(0, X) \cong 0;$$

Here the first isomorphism follows from Proposition 7.2.7, and the second from the observation that $Y[P^{-1}] \cong Y \otimes \mathbb{S}[P^{-1}]$.

Conversely, assume that X is $\mathbb{S}[P^{-1}]$ -local. To show that $p: X \rightarrow X$ is an isomorphism, it suffices by Lemma 7.1.6 to show that the map $p: X[P^{-1}] = X \otimes \mathbb{S}[P^{-1}] \rightarrow X \otimes \mathbb{S}[P^{-1}] = X[P^{-1}]$ is an isomorphism, which is the content of Corollary 7.2.5. $\square$

Corollary 7.2.9. *For every spectrum X , the map $X \rightarrow X[P^{-1}]$ is an $\mathbb{S}[P^{-1}]$ -localization of X .*

Proof. By Corollary 7.1.8, a map $X \rightarrow Y$ is an $\mathbb{S}[P^{-1}]$ -localization if and only if it exhibits Y as a reflection into the subcategory of $\mathbb{S}[P^{-1}]$ -local spectra. By Lemma 7.2.8, this subcategory agrees with the P -divisible spectra, and so the claim is an instance of Proposition 7.2.7. $\square$

For $E = \mathbb{S}[P^{-1}]$, the previous corollary implies that the E -localization of X is given by the map $X \cong \mathbb{S} \otimes X \rightarrow L_E(\mathbb{S}) \otimes X$. Localizations of this form have a special name:

Definition 7.2.10. The localization functor $L_E: \mathrm{Sp} \rightarrow \mathrm{Sp}_E$ is called a *smashing localization* if the spectrum $L_E \mathbb{S} \otimes X$ is E -local for every spectrum X .

Exercise 7.2.11. Assume L_E is a smashing localization. Construct an isomorphism $L_E(X) \xrightarrow{\sim} L_E(\mathbb{S}) \otimes X$ for every spectrum X .

We next record the expected fact that reducing a spectrum modulo a power of p leaves only p -primary information, in the sense that all other primes act invertibly. The analogous claim for abelian groups is clear, since any prime different from p is invertible modulo p^n . The following proposition extends this claim to spectra by expressing each homotopy group of X/p^n as an extension of two groups killed by p^n , allowing us to reduce to the abelian setting.

Proposition 7.2.12. *Let X be a spectrum, let p be a prime number and let $n \geq 1$. Then the spectrum $X/p^n := \mathrm{cofib}(X \xrightarrow{\cdot p^n} X)$ is p -local.*

Proof. Let q be a prime number different from p . We must show that the map $q: X/p^n \rightarrow X/p^n$ is an isomorphism, which we may check on homotopy groups.

The long exact sequence on homotopy groups associated to the exact sequence $X \xrightarrow{\cdot p^n} X \rightarrow X/p^n$ contains, for every $m \in \mathbb{Z}$, a natural short exact sequence

$$0 \rightarrow \pi_m(X)/p^n \rightarrow \pi_m(X/p^n) \rightarrow \pi_{m-1}(X)[p^n] \rightarrow 0.$$

Here $\pi_m(X)/p^n$ denotes the cokernel of multiplication by p^n , and $\pi_{m-1}(X)[p^n]$ denotes the kernel of multiplication by p^n . By Exercise 7.2.2, homotopy groups carry multiplication by q on spectra to multiplication by q on abelian groups, so this natural short exact sequence is compatible with multiplication by q .

Now multiplication by q is an isomorphism on every abelian group killed by p^n : choose integers a, b with $aq + bp^n = 1$, and then multiplication by a is inverse to multiplication by q . It follows that multiplication by q is an isomorphism on the two outer terms of the short exact sequence above. By the short five lemma, it is therefore also an isomorphism on $\pi_m(X/p^n)$.

Since this holds for all m , multiplication by q is an isomorphism of spectra. As this is true for every prime $q \neq p$, the spectrum X/p^n is p -local. $\square$

Remark 7.2.13. The previous argument shows more generally that X/N is P -divisible whenever $\gcd(q, N) = 1$ for every $q \in P$.

Theorem 7.2.14 (Serre). *The rational sphere $\mathbb{S}_{\mathbb{Q}}$ is isomorphic to the Eilenberg–MacLane spectrum $H\mathbb{Q}$ of the rational numbers.*

Proof. Consider the map $\mathbb{S} \rightarrow H\mathbb{Q}$ corresponding to $1 \in \mathbb{Q} \cong \pi_0(H\mathbb{Q})$. All primes act invertibly on the homotopy groups of $H\mathbb{Q}$, so $H\mathbb{Q}$ is $\mathbb{S}_{\mathbb{Q}}$ -local by Lemma 7.2.8, and this map therefore uniquely refines to a map $\mathbb{S}_{\mathbb{Q}} \rightarrow H\mathbb{Q}$. We have to show that this map induces isomorphisms on homotopy groups. By Lemma 7.2.4, the induced map on π_0 is the isomorphism $\mathbb{Q} \cong \pi_0(\mathbb{S}_{\mathbb{Q}}) \rightarrow \pi_0(H\mathbb{Q}) \cong \mathbb{Q}$ which sends 1 to 1. It thus remains to show that $\pi_n(\mathbb{S}_{\mathbb{Q}}) = 0$ for all $n \neq 0$. For $n > 0$, the stable homotopy group $\pi_n(\mathbb{S}) = \pi_n^{\text{st}}$ is finite by a theorem of Serre [Ser53], while for $n < 0$ it vanishes since $\mathbb{S}$ is connective. In either case it is killed by rationalization. $\square$

Definition 7.2.15. We write $\text{Sp}_{\mathbb{Q}} \subseteq \text{Sp}$ for the full subcategory of *rational spectra*, i.e. the $\mathbb{S}_{\mathbb{Q}}$ -local spectra, or equivalently the spectra on which all primes act invertibly. By Serre’s theorem, these are also precisely the $H\mathbb{Q}$ -local spectra.

Proposition 7.2.16. *For every rational spectrum X , there is a noncanonical isomorphism*

$$\bigoplus_{n \in \mathbb{Z}} H(\pi_n X)[n] \xrightarrow{\sim} X.$$

Proof. For every integer $n \in \mathbb{Z}$, we pick a basis of $\pi_n(X)$ as a rational vector space. An element of $\pi_n(X)$ corresponds to a map of spectra $\mathbb{S}[n] \rightarrow X$. Since X is rational, this uniquely extends to a map $H\mathbb{Q}[n] = \mathbb{S}_{\mathbb{Q}}[n] \rightarrow X$. By taking a direct sum over all basis vectors, this induces a map $H(\pi_n X)[n] \cong \bigoplus H\mathbb{Q}[n] \rightarrow X$ which by construction induces the identity on $\pi_n(X)$. We now take the direct sum over all $n \in \mathbb{Z}$. Direct sums of spectra are filtered colimits of finite biproducts, so Lemma 4.4.27 shows that the resulting map $\bigoplus_{n \in \mathbb{Z}} H(\pi_n X)[n] \rightarrow X$ induces an isomorphism on every homotopy group. It is therefore an isomorphism. $\square$

7.3 Completion at a prime

Beyond localizations, there is another important operation in algebra, known as *p-completion* for a prime p . The process of p -completion is in some sense orthogonal to the process of inverting p . For example, if A is a finitely generated abelian group and $A_p^{\wedge} := \lim_n A/p^n A$ is its p -completion, then A can be recovered from its p -completion and its p -inversion via the *arithmetic fracture square*:¹⁵ a pullback square in Ab of the form

$$\begin{array}{ccc} A & \xrightarrow{\quad} & A_p^{\wedge} \\ \downarrow & \lrcorner & \downarrow \\ A[\frac{1}{p}] & \longrightarrow & A_p^{\wedge}[\frac{1}{p}]. \end{array}$$

The goal of this section is to construct an analogous p -completion operation on spectra, show that it is an instance of a Bousfield localization at some spectrum, compute the homotopy groups of a completion, and establish a spectral form of the arithmetic fracture square.

¹⁵The finite-generation hypothesis is essential here: for arbitrary abelian groups, the corresponding statement requires derived p -completion; see Section 7.3.2.

7.3.1 Completion as a Bousfield localization

Definition 7.3.1. Consider the *mod p Moore spectrum* $\mathbb{S}/p$, defined as the cofiber

$$\mathbb{S}/p := \text{cofib}(p: \mathbb{S} \rightarrow \mathbb{S}).$$

We refer to an $\mathbb{S}/p$ -local spectrum as *p-complete*. We write

$$\text{Sp}_p^\wedge \subseteq \text{Sp}$$

for the full subcategory of *p-complete* spectra, and denote the resulting localization functor by

$$(-)_p^\wedge := L_{\mathbb{S}/p}: \text{Sp} \rightarrow \text{Sp}_p^\wedge.$$

Observation 7.3.2. Note that $X \otimes \mathbb{S}/p \simeq X/p$ for every X . In particular, a spectrum X is $\mathbb{S}/p$ -acyclic if and only if the map $p: X \rightarrow X$ is an isomorphism, i.e. X is $\mathbb{S}[\frac{1}{p}]$ -local by Lemma 7.2.8.

Warning 7.3.3. An $\mathbb{S}[\frac{1}{p}]$ -local spectrum should not be confused with a *p-local* spectrum: in the former, multiplication by p is an isomorphism; in the latter, this holds for every prime other than p . Moreover, unlike the prime-inversion localizations of the previous section, *p-completion* is not smashing in general.

We would like to get a more explicit description of *p-completeness* that makes its connection with *p-completion* in classical algebra clear.

Definition 7.3.4. We define the spectrum $\mathbb{S}/p^\infty$ as the cofiber of the map $\mathbb{S} \rightarrow \mathbb{S}[\frac{1}{p}]$:

$$\mathbb{S}/p^\infty := \text{cofib}(\mathbb{S} \rightarrow \mathbb{S}[\frac{1}{p}] = \text{colim}(\mathbb{S} \xrightarrow{\cdot p} \mathbb{S} \xrightarrow{\cdot p} \mathbb{S} \xrightarrow{\cdot p} \dots)).$$

By writing $\mathbb{S}$ as the colimit over the identity maps $\mathbb{S} \xrightarrow{\text{id}} \mathbb{S}$, we may alternatively write $\mathbb{S}/p^\infty$ as the colimit over the bottom row of the following commutative diagram, where all vertical sequences are cofiber sequences:

$$\begin{array}{ccccccc} \mathbb{S} & \xlongequal{\quad} & \mathbb{S} & \xlongequal{\quad} & \mathbb{S} & \xlongequal{\quad} & \dots \\ \cdot p \downarrow & & \downarrow \cdot p^2 & & \downarrow \cdot p^3 & & \\ \mathbb{S} & \xrightarrow{\cdot p} & \mathbb{S} & \xrightarrow{\cdot p} & \mathbb{S} & \xrightarrow{\cdot p} & \dots \\ \downarrow & & \downarrow & & \downarrow & & \\ \mathbb{S}/p & \xrightarrow{\cdot p} & \mathbb{S}/p^2 & \xrightarrow{\cdot p} & \mathbb{S}/p^3 & \xrightarrow{\cdot p} & \dots \end{array}$$

(This also justifies the notation $\mathbb{S}/p^\infty$.)

Theorem 7.3.5. For a spectrum X , the map

$$X = \text{hom}(\mathbb{S}, X) \rightarrow \text{hom}(\mathbb{S}/p^\infty[-1], X)$$

induced by the map $\mathbb{S}/p^\infty[-1] \rightarrow \mathbb{S}$ exhibits its target as a *p-completion* of X , i.e.

$$X_p^\wedge \xrightarrow{\sim} \text{hom}(\mathbb{S}/p^\infty[-1], X) \xrightarrow{\sim} \text{hom}(\mathbb{S}/p^\infty, X)[1].$$

Proof. We have to show that $\text{hom}(\mathbb{S}/p^\infty[-1], X)$ is p -complete and that the map $X \rightarrow \text{hom}(\mathbb{S}/p^\infty[-1], X)$ is an $\mathbb{S}/p$ -equivalence. For the latter, we may equivalently show that its fiber is $\mathbb{S}/p$ -acyclic, or equivalently that its fiber is $\mathbb{S}[\frac{1}{p}]$ -local by Observation 7.3.2. Because of the exact sequence $\mathbb{S}/p^\infty[-1] \rightarrow \mathbb{S} \rightarrow \mathbb{S}[\frac{1}{p}]$, this fiber is isomorphic to $\text{hom}(\mathbb{S}[\frac{1}{p}], X)$. But this is clearly $\mathbb{S}[\frac{1}{p}]$ -local: for any $\mathbb{S}[\frac{1}{p}]$ -acyclic spectrum Y we have $\text{hom}(Y, \text{hom}(\mathbb{S}[\frac{1}{p}], X)) \simeq \text{hom}(Y \otimes \mathbb{S}[\frac{1}{p}], X) \simeq \text{hom}(0, X) = 0$.

We will now show that $\text{hom}(\mathbb{S}/p^\infty[-1], X)$ is p -complete, i.e. $\mathbb{S}/p$ -local. To this end, let A be an $\mathbb{S}/p$ -acyclic spectrum, i.e. the map $p: A \rightarrow A$ is an isomorphism. We need to show that $0 = \text{hom}(A, \text{hom}(\mathbb{S}/p^\infty[-1], X)) \simeq \text{hom}(A \otimes \mathbb{S}/p^\infty[-1], X)$. For this, it will suffice to show that $A \otimes \mathbb{S}/p^\infty = 0$. Since we may write $\mathbb{S}/p^\infty$ as a colimit of $\mathbb{S}/p^n$, we may similarly write

$$A \otimes \mathbb{S}/p^\infty \simeq \text{colim}_n (A \otimes \mathbb{S}/p^n) \simeq \text{colim}_n (A/p^n).$$

But as A is $\mathbb{S}/p$ -acyclic, the map $p: A \rightarrow A$ and hence also $p^n: A \rightarrow A$ is an isomorphism, and so $A/p^n = 0$. This shows that $A \otimes \mathbb{S}/p^\infty \simeq 0$, as desired. $\square$

As a corollary of the theorem, we may now describe the p -completion of X as a familiar limit:

Corollary 7.3.6. *For a spectrum X , its p -completion is given by the following limit:*

$$X_p^\wedge \simeq \lim(\cdots \rightarrow X/p^3 \rightarrow X/p^2 \rightarrow X/p).$$

Proof. The exact sequence $\mathbb{S}/p^n[-1] \rightarrow \mathbb{S} \xrightarrow{\cdot p^n} \mathbb{S}$ induces an exact sequence

$$\text{hom}(\mathbb{S}, X) \xrightarrow{\cdot p^n} \text{hom}(\mathbb{S}, X) \rightarrow \text{hom}(\mathbb{S}/p^n[-1], X).$$

Since $\text{hom}(\mathbb{S}, X) \simeq X$, it follows that $\text{hom}(\mathbb{S}/p^n[-1], X) \simeq X/p^n$ for all n . Furthermore, under these isomorphisms the maps $\mathbb{S}/p^n \xrightarrow{\cdot p} \mathbb{S}/p^{n+1}$ induce the canonical maps $X/p^{n+1} \rightarrow X/p^n$, as these are the ones induced on vertical cofibers in the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{\cdot p} & X \\ \cdot p^{n+1} \downarrow & & \downarrow \cdot p^n \\ X & \xlongequal{\quad} & X. \end{array}$$

Using Theorem 7.3.5, we may then compute that

$$\begin{aligned} X_p^\wedge &\simeq \text{hom}(\mathbb{S}/p^\infty[-1], X) \simeq \text{hom}(\text{colim}_n \mathbb{S}/p^n[-1], X) \\ &\simeq \lim_n \text{hom}(\mathbb{S}/p^n[-1], X) \simeq \lim_n X/p^n, \end{aligned}$$

as desired. $\square$

Corollary 7.3.7. *Let p be a prime number. Then every p -complete spectrum is p -local.*

Proof. Since X is p -complete, the localization map $X \rightarrow X_p^\wedge$ is an $\mathbb{S}/p$ -equivalence between $\mathbb{S}/p$ -local spectra, hence is an isomorphism by Lemma 7.1.6. By Corollary 7.3.6, we may therefore write $X \cong \lim_n (X/p^n)$. Each X/p^n is p -local by Proposition 7.2.12. By Lemma 7.2.8, p -local spectra are $\mathbb{S}[P^{-1}]$ -local for P the set of all primes different from p , so they are closed under limits by Exercise 7.1.4. $\square$

Corollary 7.3.8. *Let p and q be two distinct primes. If X is both p -complete and q -complete, then X is the zero spectrum.*

Proof. The q -completeness of X means that X is $\mathbb{S}/q$ -local. The p -completeness of X implies, via Corollary 7.3.7, that the map $q: X \rightarrow X$ is an isomorphism, hence that X is $\mathbb{S}/q$ -acyclic. It then follows from Lemma 7.1.6 that $X = 0$. $\square$

7.3.2 Homotopy groups of a p -completion

The inverse limit formula $X_p^\wedge \simeq \lim_k X/p^k$ does not immediately compute the homotopy groups of the p -completion, since homotopy groups need not preserve inverse limits. Instead, we use the mapping spectrum formula of Theorem 7.3.5. The telescope construction will give a resolution of $\mathbb{S}/p^\infty$ by direct sums of spheres which realizes a free resolution of the Prüfer group $\mathbb{Z}/p^\infty := \mathbb{Z}[\frac{1}{p}]/\mathbb{Z}$. After applying $\text{hom}(-, X)$, the resulting long exact sequence computes $\pi_*(X_p^\wedge)$ in terms of Hom and Ext .

Accordingly, for an abelian group A we write

$$L_0^p A := \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/p^\infty, A) \quad \text{and} \quad L_1^p A := \text{Hom}_{\mathbb{Z}}(\mathbb{Z}/p^\infty, A)$$

for its two *derived p -completion groups*. Throughout this subsection we write $\mathbb{Z}_p := \lim_k \mathbb{Z}/p^k$, with respect to the quotient maps, for the ring of p -adic integers. We first record a useful calculation.

Lemma 7.3.9. *Let A be an abelian group, and write $A[p^k] \subseteq A$ for the subgroup of elements annihilated by p^k . Then*

$$L_1^p A \cong \lim_k A[p^k],$$

$$0 \longrightarrow \lim_k^1 A[p^k] \longrightarrow L_0^p A \longrightarrow \lim_k A/p^k A \longrightarrow 0,$$

where the transition maps on $A[p^k]$ are multiplication by p . If A is finitely generated, then $L_1^p A = 0$ and $L_0^p A \cong A \otimes_{\mathbb{Z}} \mathbb{Z}_p$.

Proof. Write $\mathbb{Z}/p^\infty \cong \text{colim}_k \mathbb{Z}/p^k$ for the filtered colimit along the multiplication-by- p injections $\mathbb{Z}/p^k \xrightarrow{\cdot p} \mathbb{Z}/p^{k+1}$. Since filtered colimits are exact in Ab , the colimit in $\text{D}(\mathbb{Z})$ of a filtered diagram of abelian groups concentrated in degree zero is again concentrated in degree zero, with degree-zero homology given by the colimit in Ab . Thus the displayed colimit also computes the colimit in $\text{D}(\mathbb{Z})$. Mapping out of it gives

$$\text{hom}_{\text{D}(\mathbb{Z})}(\mathbb{Z}/p^\infty, A) \simeq \lim_k \text{hom}_{\text{D}(\mathbb{Z})}(\mathbb{Z}/p^k, A).$$

For every $k \geq 1$, the standard resolution of $\mathbb{Z}/p^k$ identifies

$$\text{Hom}_{\mathbb{Z}}(\mathbb{Z}/p^k, A) \cong A[p^k] \quad \text{and} \quad \text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/p^k, A) \cong A/p^k A.$$

Under these identifications, the transition maps are multiplication by p on the first tower and the quotient maps on the second. The two claims now follow by applying the Milnor sequence of Corollary 4.4.36 in degrees 0 and -1 .

If A is finitely generated, its p -primary torsion is bounded, so the transition maps into every fixed stage of the first tower are eventually zero. It follows directly from the description in Definition 4.4.35 that both its limit and its first derived limit vanish. Finally, $\lim_k A/p^k A \cong A \otimes_{\mathbb{Z}} \mathbb{Z}_p$ for every finitely generated abelian group A . $\square$

Theorem 7.3.10 (Homotopy groups of a p -completion). *Let p be a prime and let X be a spectrum. Then for every $n \in \mathbb{Z}$ there is a natural short exact sequence*

$$0 \rightarrow \operatorname{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/p^\infty, \pi_n X) \rightarrow \pi_n(X_p^\wedge) \rightarrow \operatorname{Hom}_{\mathbb{Z}}(\mathbb{Z}/p^\infty, \pi_{n-1} X) \rightarrow 0.$$

Proof. Let $F := \bigoplus_{k \geq 0} \mathbb{S}$, with summand inclusions $\iota_k: \mathbb{S} \rightarrow F$. Splicing the telescope sequence for $\mathbb{S}[\frac{1}{p}]$ from Lemma 4.4.33 with the defining exact sequence $\mathbb{S} \rightarrow \mathbb{S}[\frac{1}{p}] \rightarrow \mathbb{S}/p^\infty$, and then reindexing the source, gives an exact sequence

$$F \xrightarrow{d} F \rightarrow \mathbb{S}/p^\infty,$$

where $d \circ \iota_0 = \iota_0$ and $d \circ \iota_{k+1} = \iota_k - p\iota_{k+1}$ for $k \geq 0$. On π_0 this realizes the free resolution

$$0 \rightarrow \bigoplus_{k \geq 0} \mathbb{Z} \xrightarrow{d_*} \bigoplus_{k \geq 0} \mathbb{Z} \rightarrow \mathbb{Z}/p^\infty \rightarrow 0,$$

in which the k -th basis vector of the middle group maps to $p^{-k} + \mathbb{Z}$. Thus the exact sequence of spectra is obtained by realizing this free resolution by direct sums of spheres.

Apply $\operatorname{hom}(-, X)$ to the exact sequence above. For every $m \in \mathbb{Z}$, the universal property of the coproduct identifies

$$\pi_m \operatorname{hom}(F, X) \cong \operatorname{Hom}_{\mathbb{Z}}\left(\bigoplus_{k \geq 0} \mathbb{Z}, \pi_m X\right),$$

and the map induced by d agrees with precomposition by d_* . By Theorem 7.3.5, we have $X_p^\wedge \simeq \operatorname{hom}(\mathbb{S}/p^\infty, X)[1]$, so that $\pi_n(X_p^\wedge) \cong \pi_{n-1} \operatorname{hom}(\mathbb{S}/p^\infty, X)$. Write d_m^* for the endomorphism induced by d^* on $\pi_m \operatorname{hom}(F, X)$. The long exact sequence associated to the exact sequence $\operatorname{hom}(\mathbb{S}/p^\infty, X) \rightarrow \operatorname{hom}(F, X) \xrightarrow{d^*} \operatorname{hom}(F, X)$ then gives a short exact sequence

$$0 \rightarrow \operatorname{coker}(d_n^*) \rightarrow \pi_n(X_p^\wedge) \rightarrow \ker(d_{n-1}^*) \rightarrow 0.$$

By Proposition 6.4.14, its outer terms are

$$\operatorname{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/p^\infty, \pi_n X) = L_0^P(\pi_n X) \quad \text{and} \quad \operatorname{Hom}_{\mathbb{Z}}(\mathbb{Z}/p^\infty, \pi_{n-1} X) = L_1^P(\pi_{n-1} X),$$

respectively. This gives the asserted short exact sequence. $\square$

Corollary 7.3.11. *Let X be a spectrum and let $n \in \mathbb{Z}$ be such that $\pi_n(X)$ and $\pi_{n-1}(X)$ are finitely generated. Then there is an isomorphism*

$$\pi_n(X_p^\wedge) \cong \pi_n(X) \otimes_{\mathbb{Z}} \mathbb{Z}_p.$$

Proof. By Lemma 7.3.9, the right-hand term of the short exact sequence of Theorem 7.3.10 vanishes, while its left-hand term is $\pi_n(X) \otimes_{\mathbb{Z}} \mathbb{Z}_p$. $\square$

Example 7.3.12. All homotopy groups of the sphere spectrum are finitely generated: this is $\pi_0(\mathbb{S}) = \mathbb{Z}$ together with the finiteness theorem of Serre [Ser53] used in the proof of Theorem 7.2.14. Applying Corollary 7.3.11, we obtain $\pi_0(\mathbb{S}_p^\wedge) \cong \mathbb{Z}_p$, while for $n \neq 0$ the group $\pi_n(\mathbb{S}_p^\wedge) \cong \pi_n(\mathbb{S}) \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is the p -primary part of the stable stem $\pi_n(\mathbb{S})$. Completing at p thus discards precisely the prime-to- p information in the stable stems.

Warning 7.3.13. The group $L_1^P A$ detects compatible systems of p -power torsion, and it need not vanish. For example, let $X := H(\mathbb{Z}/p^\infty)$. Since the Prüfer group is injective, $L_0^P(\mathbb{Z}/p^\infty) = 0$, whereas Lemma 7.3.9 gives $L_1^P(\mathbb{Z}/p^\infty) \cong \mathbb{Z}_p$. Hence

$$X_p^\wedge \simeq H(\mathbb{Z}_p)[1].$$

This example in particular shows that completion is not computed degree-wise on homotopy groups: the p -completion of a nonzero spectrum concentrated in degree 0 may be concentrated in degree 1.

7.3.3 The arithmetic fracture square

We now return to the interaction between p -completion and inversion of p , and prove the spectral form of the arithmetic fracture square announced at the start of this section.

Lemma 7.3.14. *Let P be a set of primes and let X be a spectrum satisfying $X[P^{-1}] = 0$ and $X/p = 0$ for every $p \in P$. Then $X = 0$.*

Proof. For every $p \in P$, the condition $X/p = 0$ means that multiplication by p on X is an isomorphism. Thus X is P -divisible, hence $\mathbb{S}[P^{-1}]$ -local by Lemma 7.2.8. On the other hand, the isomorphism $X[P^{-1}] \cong X \otimes \mathbb{S}[P^{-1}]$ shows that X is $\mathbb{S}[P^{-1}]$ -acyclic. It follows from Lemma 7.1.6 that $X = 0$. $\square$

Theorem 7.3.15 (The p -adic arithmetic fracture square). *Let X be a spectrum. Then the following commutative square is a pullback square:*

$$\begin{array}{ccc} X & \xrightarrow{\quad} & X_p^\wedge \\ \downarrow & \lrcorner & \downarrow \\ X[\frac{1}{p}] & \longrightarrow & X_p^\wedge[\frac{1}{p}]. \end{array}$$

Proof. We have to show that the map $X \rightarrow X[\frac{1}{p}] \times_{X_p^\wedge[\frac{1}{p}]} X_p^\wedge$ is an isomorphism, or equivalently that its fiber F is the zero spectrum. Note that $F[\frac{1}{p}] = 0$, since the square

$$\begin{array}{ccc} X[\frac{1}{p}] & \longrightarrow & X_p^\wedge[\frac{1}{p}] \\ \downarrow & & \downarrow \\ X[\frac{1}{p}][\frac{1}{p}] & \longrightarrow & X_p^\wedge[\frac{1}{p}][\frac{1}{p}] \end{array}$$

is a pullback square (as the two vertical maps are isomorphisms). We claim that also $F/p = 0$, so that $F = 0$ by Lemma 7.3.14, applied with $P = \{p\}$. To see this, we need to show that the commutative square

$$\begin{array}{ccc} X/p & \longrightarrow & X_p^\wedge/p \\ \downarrow & & \downarrow \\ X[\frac{1}{p}]/p & \longrightarrow & X_p^\wedge[\frac{1}{p}]/p \end{array}$$

is a pullback square. The two objects at the bottom are zero, since the maps $p: X[\frac{1}{p}] \rightarrow X[\frac{1}{p}]$ and $p: X_p^\wedge[\frac{1}{p}] \rightarrow X_p^\wedge[\frac{1}{p}]$ are isomorphisms by Corollary 7.2.5. In particular the bottom map is an isomorphism. But the top map is also an isomorphism, since the map $X \rightarrow X_p^\wedge$ is by definition an $\mathbb{S}/p$ -equivalence. This finishes the proof. $\square$

Exercise 7.3.16 (Arithmetic fracture square). Show that for every spectrum X , the commutative squares

$$\begin{array}{ccccc} X & \longrightarrow & \prod_p X_p^\wedge & & X & \longrightarrow & \prod_p X_{(p)} & & X_{(p)} & \longrightarrow & X_p^\wedge \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ X_{\mathbb{Q}} & \longrightarrow & \left(\prod_p X_p^\wedge\right)_{\mathbb{Q}} & & X_{\mathbb{Q}} & \longrightarrow & \left(\prod_p X_{(p)}\right)_{\mathbb{Q}} & & X_{\mathbb{Q}} & \longrightarrow & \left(X_p^\wedge\right)_{\mathbb{Q}} \end{array}$$

are pullback squares. In the first two squares, the products are taken over all prime numbers p ; the third square is asserted for every prime p .

Hint: in each case, apply Lemma 7.3.14 to the fiber of the comparison map, taking P to be the set of all primes.

Exercises

Exercise 7.1. Let E and F be spectra. Show that the following two conditions are equivalent:

- (1) Every E -acyclic spectrum X is also F -acyclic;
- (2) Every F -local spectrum Y is also E -local.

Remark. Two spectra E and F are said to *have the same Bousfield class* if a spectrum X is E -acyclic if and only if it is F -acyclic. By the previous exercise, E and F have the same Bousfield class if and only if the subcategories $\mathrm{Sp}_E \subseteq \mathrm{Sp}$ and $\mathrm{Sp}_F \subseteq \mathrm{Sp}$ agree.

Exercise 7.2. Let $X \in \mathrm{Sp}_{\mathbb{Q}}$ be a rational spectrum. Assume that X is p -complete for some prime p . Show that X is the zero spectrum.

Exercise 7.3. Let E, F and X be spectra. Is it always true that there is an isomorphism $L_E L_F X \simeq L_F L_E X$? If yes, provide a proof. If no, provide a counterexample.

8 Ring spectra and modules

Cohomology theories are almost never encountered as bare graded abelian groups: they usually carry multiplicative structure. Ordinary cohomology $H^*(X; \mathbb{Z})$ comes equipped with a *cup product*, turning it into a graded ring, and the K-theory of a space (introduced in the next chapter) carries a product coming from the tensor product of vector bundles. Combining Theorem 3.2.6 with the Yoneda lemma, the cup product on $H^*(X; \mathbb{Z})$ is induced by a multiplication $H\mathbb{Z} \otimes H\mathbb{Z} \rightarrow H\mathbb{Z}$ on the representing spectrum itself, which is unital, associative and commutative up to homotopy. The resulting structure on $H\mathbb{Z}$ is that of a *homotopy ring spectrum*, and the spectrum representing K-theory carries a similar multiplicative structure.

Once spectra rather than cohomology theories are taken as the primary objects of study, it becomes natural to view this multiplication on $H\mathbb{Z}$ as the shadow of a much richer structure, that of a *ring spectrum*: a spectrum R equipped with a multiplication map $R \otimes R \rightarrow R$ which is coherently unital and associative, much like the coherent multiplication of a monoid object in Chapter 5. Alongside it comes the notion of an *R -module*: a spectrum equipped with a coherent action of R .

Where ordinary algebra takes place over the base ring $\mathbb{Z}$, the algebra of ring spectra takes place over the sphere spectrum $\mathbb{S}$, the monoidal unit for the tensor product on Sp . As the homotopy groups of $\mathbb{S}$ are the stable stems, algebra over the sphere is a substantially richer subject than algebra over $\mathbb{Z}$. Ordinary algebra can be recovered as a fragment of this theory: the Eilenberg–MacLane construction $A \mapsto HA$ promotes an ordinary ring to a ring spectrum. This also recovers the homological algebra from Chapter 6: for a commutative ring R we will prove an equivalence

$$D(R) \simeq \mathrm{Mod}_{HR}(\mathrm{Sp}),$$

identifying the derived ∞ -category of R -modules with the ∞ -category of HR -modules. The tools from homological algebra all admit counterparts over an arbitrary ring spectrum.

We begin in Section 8.1 by recording the package of coherent algebra and module theory that we will use from Part II. Section 8.2 studies the interaction of modules with connectivity, and Section 8.3 carries out the comparison with ordinary rings and with differential graded algebras sketched above. We then construct localizations of commutative ring spectra in Section 8.4. The online version concludes with supplementary treatments of Ore localization, projective and flat modules, and Tor-amplitude.

The homotopy coherent algebra underlying all of this is the theory of ∞ -operads, which is developed at length in Part II. In the present chapter we use that theory as a black box. More precisely, we record in Proposition 8.1.5 the constructions and properties that will be used below. A reader following Part I may take this package as given; none of the later arguments requires familiarity with operads.

This chapter also leans on [Lur17] more heavily than the rest of Part I. Results quoted without proof are flagged as black boxes where they occur. Several further results that are not used elsewhere in the book are retained in the online-only supplements.

8.1 Algebras and modules

The notions of algebras and modules make sense in every symmetric monoidal ∞ -category. We first recall some examples of symmetric monoidal structures, and then state the coherent algebra package from Part II that we will use throughout the chapter.

Definition 8.1.1. A *symmetric monoidal ∞ -category* is defined to be a commutative monoid in Cat_∞ . In particular, it is an ∞ -category C which comes equipped with a *tensor product functor*

$$- \otimes -: C \times C \rightarrow C$$

and a *monoidal unit* $\mathbb{1} \in C$, and the tensor product is coherently unital, associative and commutative. We will often denote symmetric monoidal ∞ -categories as $(C, \otimes, \mathbb{1})$, or often also just as C . Given symmetric monoidal ∞ -categories C and D , a *symmetric monoidal functor* $F: C \rightarrow D$ is a morphism in $\text{CMon}(\text{Cat}_\infty)$. We write $\text{Map}^\otimes(C, D)$ for the hom anima in $\text{CMon}(\text{Cat}_\infty)$.

We write $\text{Fun}^\otimes(C, D)$ for the ∞ -category of symmetric monoidal functors and monoidal natural transformations. We will also use the more flexible notion of a *lax symmetric monoidal functor* $F: C \rightarrow D$. Concretely, such a functor is equipped with coherent structure maps

$$\mathbb{1}_D \longrightarrow F(\mathbb{1}_C), \quad F(X) \otimes F(Y) \longrightarrow F(X \otimes Y)$$

compatible with the unit, associativity and symmetry. A lax symmetric monoidal functor is symmetric monoidal precisely when these structure maps are isomorphisms. We denote the ∞ -category of lax symmetric monoidal functors by $\text{Fun}^{\otimes\text{-lax}}(C, D)$. The formal constructions of these functor categories are given in Definitions 14.2.3 and 14.3.1.

Example 8.1.2. The following are examples of symmetric monoidal ∞ -categories:

- (1) If C is an ∞ -category and D is symmetric monoidal, then $\text{Fun}(C, D)$ admits the *pointwise symmetric monoidal structure*, with tensor product and unit computed pointwise.
- (2) An ∞ -category C with finite products admits the *cartesian monoidal structure* $(C, \times, *)$. Dually, finite coproducts give the *cocartesian monoidal structure* $(C, \amalg, \emptyset)$. These structures are constructed in Sections 15.2 and 15.3.
- (3) Let $(C, \otimes, \mathbb{1})$ be a symmetric monoidal ∞ -category and let $D \subseteq C$ be a full subcategory which contains the monoidal unit and is closed under tensor products. Then D inherits a symmetric monoidal structure for which the inclusion $D \hookrightarrow C$ is symmetric monoidal. See Lemma 14.5.2 for the construction.
- (4) If C and D are symmetric monoidal, C is small, and D is cocomplete with tensor product preserving colimits in both variables, then $\text{Fun}(C, D)$ also admits the *Day convolution* symmetric monoidal structure. We refer to Chapter 16, especially Proposition 16.2.7 and Corollary 16.2.8, for its universal property and construction.

- (5) The ∞ -category Sp of spectra admits a symmetric monoidal structure $\otimes$ satisfying the property that for every spectrum X , the functor $X \otimes - : \mathrm{Sp} \rightarrow \mathrm{Sp}$ preserves colimits. Furthermore, Sp is *universal* with this structure: if D is another symmetric monoidal stable ∞ -category which admits colimits and in which the tensor product preserves colimits in both variables, then the ∞ -category of colimit-preserving symmetric monoidal functors $\mathrm{Sp} \rightarrow D$ is contractible. We construct the symmetric monoidal structure in Chapter 16, and discuss its universal property in Section 18.5.
- (6) The ∞ -categories An_* , $\mathrm{CMon}(\mathrm{An})$ and $\mathrm{CGrp}(\mathrm{An})$ similarly admit universal symmetric monoidal structures for pointed, semiadditive, and additive ∞ -categories. In particular, the functors

$$\mathrm{An} \xrightarrow{(-)_+} \mathrm{An}_* \rightarrow \mathrm{CMon}(\mathrm{An}) \xrightarrow{(-)^{\mathrm{grp}}} \mathrm{CGrp}(\mathrm{An}) \xrightarrow{\mathbf{B}^\infty} \mathrm{Sp}$$

are colimit-preserving and symmetric monoidal; see Proposition 16.6.4.

- (7) Let R be a commutative ring. Then the *derived ∞ -category* of R , defined as the localization

$$\mathrm{D}(R) := \mathrm{Ch}(R)[\{\text{quasi-isomorphisms}\}^{-1}]$$

of the category of chain complexes of R -modules at the quasi-isomorphisms, admits a symmetric monoidal structure $(\mathrm{D}(R), \otimes, R[0])$ whose monoidal unit is the chain complex $R[0]$ given by putting R in degree 0. The tensor product corresponds to the *derived tensor product* of chain complexes. A construction of this symmetric monoidal structure is given in Section 20.2.

- (8) Recall the full subcategory $\mathrm{Cat}_1 \subseteq \mathrm{Cat}_\infty$ from Example 1.8.14(2) spanned by the 1-categories. A commutative monoid in Cat_1 is precisely that of a symmetric monoidal category as defined classically. Since the inclusion $\mathrm{Cat}_1 \hookrightarrow \mathrm{Cat}_\infty$ preserves products, this implies that any symmetric monoidal category defines a symmetric monoidal ∞ -category.

Recall from classical algebra that a *commutative algebra* in a symmetric monoidal category C is an object A of C that comes equipped with a unit $1 : \mathbb{1} \rightarrow A$ and a multiplication map $m : A \otimes A \rightarrow A$ such that the following diagrams commute:

$$\begin{array}{ccccc} A \otimes \mathbb{1} & \xrightarrow{\mathrm{id}_A \otimes 1} & A \otimes A & \xleftarrow{1 \otimes \mathrm{id}_A} & A \\ & \searrow \cong & \downarrow m & \swarrow \cong & \\ & & A & & \end{array} \quad \begin{array}{ccc} A \otimes A \otimes A & \xrightarrow{m \otimes \mathrm{id}_A} & A \otimes A \\ \mathrm{id}_A \otimes m \downarrow & & \downarrow m \\ A \otimes A & \xrightarrow{m} & A, \end{array} \quad \begin{array}{ccc} A \otimes A & \xrightarrow[\cong]{\tau} & A \otimes A \\ m \searrow & & \swarrow m \\ & A & \end{array}$$

In an ∞ -category, these three relations are only the beginning: one must also specify coherent higher homotopies between them. The following definition packages all these coherences at once.

Definition 8.1.3 (Commutative algebra). Let $(C, \otimes, \mathbb{1})$ be a symmetric monoidal ∞ -category. We define a *commutative algebra in C* to be a symmetric monoidal functor

$$A : \mathrm{Fin} \rightarrow C,$$

where we equip the category Fin of finite sets with the cocartesian monoidal structure; this functor packages the coherent n -fold multiplications described below. We refer to the object $A(*)$ in C as the *underlying object of A* . We will frequently abuse notation and simply write A for $A(*)$.

We define the ∞ -category of commutative algebras in C as

$$\mathrm{CAlg}(C) := \mathrm{Fun}^\otimes(\mathrm{Fin}, C).$$

Remark 8.1.4. For the cartesian monoidal structure on an ∞ -category C with finite products, commutative algebras are precisely commutative monoids: $\mathrm{CAlg}(C, \times) \simeq \mathrm{CMon}(C)$ by Corollary 15.3.13.

Since every finite set is a disjoint union of one-point sets, a commutative algebra supplies coherent n -fold multiplication maps

$$A^{\otimes n} \longrightarrow A$$

for all $n \geq 0$. The unit, associativity, and commutativity displayed above are part of this structure, together with all higher compatibility data.

There are analogous coherent notions of associative algebras and of modules. We collect the part of their theory needed in this chapter.

Proposition 8.1.5 (Coherent algebra package). *Let C be a cocomplete symmetric monoidal ∞ -category whose tensor product preserves colimits separately in both variables.*

- (1) *There are ∞ -categories $\mathrm{Alg}(C)$ and $\mathrm{CAlg}(C)$ of associative and commutative algebras in C . Their objects have underlying objects of C equipped with coherently unital and associative multiplication maps; for commutative algebras the multiplication is coherently commutative as well.*
- (2) *For every $A \in \mathrm{Alg}(C)$ there are ∞ -categories $\mathrm{LMod}_A(C)$ and $\mathrm{RMod}_A(C)$ of left and right A -modules. The forgetful functors to C create every limit admitted by C and all colimits, and the free left A -module on $X \in C$ has underlying object $A \otimes X$. If C is stable, then $\mathrm{LMod}_A(C)$ and $\mathrm{RMod}_A(C)$ are stable.*
- (3) *There is a relative tensor product*

$$- \otimes_A -: \mathrm{RMod}_A(C) \times \mathrm{LMod}_A(C) \longrightarrow C,$$

which preserves colimits separately in both variables. It is unital: for every left A -module M and right A -module N there are natural isomorphisms

$$A \otimes_A M \xrightarrow{\sim} M, \quad N \otimes_A A \xrightarrow{\sim} N.$$

- (4) *Every morphism $f: A \rightarrow B$ in $\mathrm{Alg}(C)$ induces a restriction functor*

$$f_*: \mathrm{LMod}_B(C) \longrightarrow \mathrm{LMod}_A(C),$$

which admits the extension-of-scalars functor

$$f^* := B \otimes_A -: \mathrm{LMod}_A(C) \longrightarrow \mathrm{LMod}_B(C)$$

as a left adjoint. There is an analogous adjunction for right modules, whose left adjoint is $- \otimes_A B$.

- (5) *If $R \in \mathrm{CAlg}(C)$, then left and right R -modules agree, and the resulting ∞ -category $\mathrm{Mod}_R(C)$ is symmetric monoidal under $- \otimes_R -$. There are natural equivalences*

$$\mathrm{CAlg}(\mathrm{Mod}_R(C)) \simeq \mathrm{CAlg}(C)_{R/}.$$

- (6) *Lax symmetric monoidal functors preserve algebra objects and their modules. In particular, a symmetric monoidal adjunction $F : C \rightleftarrows D : G$ induces adjunctions on algebras and on their module categories. If $A \in \text{Alg}(C)$ and the unit $A \rightarrow GF(A)$ is an isomorphism, this gives an adjunction*

$$\text{LMod}_A(C) \rightleftarrows \text{LMod}_{F(A)}(D)$$

whose underlying functors are induced by F and G . Right adjoints of symmetric monoidal functors are canonically lax symmetric monoidal.

- (7) *If C is presentable, then $\text{LMod}_A(C)$ and $\text{RMod}_A(C)$ are presentable for every $A \in \text{Alg}(C)$.*

Proof. These statements are discussed in more detail in Part II and Chapter 22. See Sections 19.1 and 19.2, Corollary 19.1.17, Propositions 14.3.6, 19.2.12, 19.2.15 and 22.5.5, and Theorem 19.2.14. $\square$

Definition 8.1.6 (Ring spectra and modules). An *associative ring spectrum* is an associative algebra in $(\text{Sp}, \otimes, \mathbb{S})$, and a *commutative ring spectrum* is a commutative algebra in Sp . Every algebra A is canonically a left and right module over itself via its multiplication $A \otimes A \rightarrow A$; this is called the *regular A -module*. We write

$$\text{Alg} := \text{Alg}(\text{Sp}), \quad \text{CAlg} := \text{CAlg}(\text{Sp})$$

for their ∞ -categories. For an associative ring spectrum R , we abbreviate

$$\text{LMod}_R := \text{LMod}_R(\text{Sp}), \quad \text{RMod}_R := \text{RMod}_R(\text{Sp}).$$

If R is commutative, we write $\text{Mod}_R := \text{Mod}_R(\text{Sp})$ for its symmetric monoidal ∞ -category of modules.

Unless commutativity is specified, a ring spectrum will mean an associative ring spectrum.

The regular module generates an important finiteness class.

Definition 8.1.7 (Perfect module). Let R be an associative ring spectrum. We denote by

$$\text{Perf}(R) \subseteq \text{LMod}_R$$

the smallest thick subcategory containing the regular left R -module R , where a subcategory is called *thick* if it is stable and closed under retracts. A left R -module is called *perfect* if it belongs to $\text{Perf}(R)$.

Thus a perfect module is obtained from finitely many shifts of R by iterating finite cofiber sequences and retracts. The compact-generation results from Chapter 22 identify this intrinsic finite construction with compactness.

Lemma 8.1.8 (Perfect modules and compact objects, [Lur17, Proposition 7.2.4.2]). *Let R be an associative ring spectrum.*

- (1) *The regular left R -module R is compact.*
- (2) *The ∞ -category LMod_R is compactly generated by R .*

(3) A left R -module is perfect if and only if it is compact.

Proof. The functor $\mathrm{Hom}_{\mathrm{LMod}_R}(R, -)$ is naturally isomorphic to the composite

$$\mathrm{LMod}_R \xrightarrow{\mathrm{fgt}} \mathrm{Sp} \xrightarrow{\Omega^\infty} \mathrm{An},$$

which preserves filtered colimits. Thus R is compact. Its shifts jointly detect isomorphisms, since

$$\pi_n(M) \cong \pi_0 \mathrm{Hom}_{\mathrm{LMod}_R}(R[n], M).$$

Hence R is a compact generator of LMod_R . The characterization of compact objects in a compactly generated stable ∞ -category from Proposition 22.3.4 now identifies the compact objects with the thick subcategory generated by R , which is $\mathrm{Perf}(R)$. $\square$

8.2 Connective modules

Let R be a connective ring spectrum. The goal of this section is to establish the Postnikov t-structure on the stable ∞ -category LMod_R of left R -modules. This t-structure will be inherited from the Postnikov t-structure on Sp (Proposition 6.3.4), using the following property of the monoidal structure:

Definition 8.2.1 (Connectivity-preserving symmetric monoidal structure). Let C be a stably symmetric monoidal ∞ -category, meaning that C is stable and its tensor product is exact separately in both variables, and equip C with a t-structure. We say that *the symmetric monoidal structure on C is connectivity-preserving* if the following two conditions are satisfied:

- (1) The monoidal unit $\mathbb{1}$ of C is connective;
- (2) For two connective objects $X, Y \in C_{\geq 0}$, the tensor product $X \otimes Y$ is again connective.

This condition is often expressed by saying that the t-structure is *compatible with the symmetric monoidal structure*. If the symmetric monoidal structure is connectivity-preserving, then $C_{\geq 0}$ inherits a symmetric monoidal structure making the inclusion $C_{\geq 0} \hookrightarrow C$ into a symmetric monoidal functor; this is proved in Part II in Lemma 14.5.2. As a consequence, the truncation functor $\tau_{\geq 0}: C \rightarrow C_{\geq 0}$ is canonically lax symmetric monoidal (see Proposition 14.3.6).

Definition 8.2.2. Let C be a stable ∞ -category equipped with a t-structure and a connectivity-preserving symmetric monoidal structure. Let $R \in \mathrm{Alg}(C_{\geq 0})$ be a connective associative algebra in C . We say that a left R -module M is *connective* if its underlying object in C is connective, and similarly for coconnective modules. This defines full subcategories

$$\mathrm{LMod}_R(C)_{\geq 0}, \quad \mathrm{LMod}_R(C)_{\leq 0} \subseteq \mathrm{LMod}_R(C).$$

Lemma 8.2.3. Let C be a stably symmetric monoidal ∞ -category equipped with a t-structure for which the symmetric monoidal structure is connectivity-preserving, and let $R \in \mathrm{Alg}(C_{\geq 0})$. Then $\mathrm{LMod}_R(C)$ admits a t-structure in which a left R -module is connective or coconnective precisely when its underlying object in C is so. The truncation functors are computed on underlying objects.

Proof. The adjunction between the symmetric monoidal inclusion $C_{\geq 0} \hookrightarrow C$ and its lax symmetric monoidal right adjoint $\tau_{\geq 0}$ induces, after passing to algebras and modules and taking the fiber over R , an adjunction

$$\mathrm{LMod}_R(C_{\geq 0}) \rightleftarrows \mathrm{LMod}_R(C).$$

Its right adjoint is computed on underlying objects by $\tau_{\geq 0}$. Finite limits and colimits of modules are computed on underlying objects by Corollary 19.1.17, so it remains to check the orthogonality and decomposition axioms for a t-structure. If X is connective and Y is coconnective, adjunction gives

$$\mathrm{Hom}_{\mathrm{LMod}_R(C)}(X, Y[-1]) \simeq \mathrm{Hom}_{\mathrm{LMod}_R(C_{\geq 0})}(X, \tau_{\geq 0}(Y[-1])) \simeq *.$$

Finally, for an arbitrary R -module X , the counit $\tau_{\geq 0}X \rightarrow X$ in modules has connective source, and its cofiber is coconnective because this is true after forgetting to C . This supplies the required decomposition. $\square$

We now wish to apply this to the Postnikov t-structure on Sp .

Lemma 8.2.4. *The symmetric monoidal structure on Sp is connectivity-preserving.*

Proof. The sphere spectrum is connective. Given two connective spectra X and Y , we need to show that $X \otimes Y$ is again connective. If we write $X_n := \Omega^{\infty-n}X$ and $Y_m := \Omega^{\infty-m}Y$ for all $n, m \geq 0$, then X_n is n -connective and Y_m is m -connective. The standard connectivity estimate for smash products, which follows by applying the Blakers–Massey theorem to $X_n \vee Y_m \rightarrow X_n \times Y_m$, shows that $X_n \wedge Y_m$ is $(n+m)$ -connective. Using the colimit formula from Section 4.4.3, we have

$$X \otimes Y \simeq \mathrm{colim}_n \mathrm{colim}_m (\Sigma^{\infty}(X_n \wedge Y_m))[-(n+m)].$$

The shift-additivity ambiguity mentioned there is irrelevant here, since we use only the connectivity of the terms. Each spectrum in this colimit is connective, and $\mathrm{Sp}_{\geq 0} \subseteq \mathrm{Sp}$ is closed under colimits, so $X \otimes Y \in \mathrm{Sp}_{\geq 0}$, as desired. $\square$

Corollary 8.2.5. *Let R be a connective associative ring spectrum. Then LMod_R admits a t-structure given by the connective and coconnective R -modules. This t-structure is both left and right complete, and every left R -module M satisfies*

$$M \simeq \mathrm{colim}(\tau_{\geq 0}M \rightarrow \tau_{\geq -1}M \rightarrow \dots) \quad \text{and} \quad M \simeq \lim(\dots \rightarrow \tau_{\leq 1}M \rightarrow \tau_{\leq 0}M).$$

The same statements hold for RMod_R .

Proof. The existence of the t-structure is, in light of the previous lemma, an instance of Lemma 8.2.3. By that same lemma, an R -module is connective or coconnective precisely when its underlying spectrum is, so an infinitely connective or infinitely coconnective R -module has vanishing homotopy groups and is zero. Thus the t-structure is left and right separated. The forgetful functor $\mathrm{LMod}_R \rightarrow \mathrm{Sp}$ preserves limits and colimits, and $\mathrm{Sp}_{\geq 0}$ and $\mathrm{Sp}_{\leq 0}$ are closed under countable products and countable coproducts, respectively. Lemma 6.3.31 therefore upgrades separatedness to left and right completeness. The two displayed formulas express these completeness properties. $\square$

Proposition 8.2.6. *The functor $\pi_0: \mathrm{Sp}_{\geq 0} \rightarrow \mathrm{Ab}$ is symmetric monoidal, where Ab is equipped with the usual tensor product $- \otimes^{\vee} -$.*

Proof. Under the symmetric monoidal equivalence $\mathrm{Sp}_{\geq 0} \simeq \mathrm{CGrp}(\mathrm{An})$ established in Part II (Corollary 16.6.5), this functor corresponds to the symmetric monoidal functor $\pi_0: \mathrm{CGrp}(\mathrm{An}) \rightarrow \mathrm{CGrp}(\mathrm{Set}) = \mathrm{Ab}$ of Proposition 16.4.5. $\square$

Observation 8.2.7. Let R be a connective associative ring spectrum. The fully faithful symmetric monoidal inclusion $\mathrm{Sp}_{\geq 0} \hookrightarrow \mathrm{Sp}$ identifies $\mathrm{LMod}_R(\mathrm{Sp}_{\geq 0})$ with the full subcategory of LMod_R spanned by the connective modules. By Proposition 8.2.6, extracting π_0 therefore determines a functor

$$\pi_0: \mathrm{LMod}_{R, \geq 0} = \mathrm{LMod}_R(\mathrm{Sp}_{\geq 0}) \rightarrow \mathrm{LMod}_{\pi_0(R)}(\mathrm{Ab}).$$

Furthermore, this functor is compatible with relative tensor products: given a connective right R -module M and a connective left R -module N , we have an isomorphism of abelian groups

$$\pi_0(M) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \xrightarrow{\sim} \pi_0(M \otimes_R N), \quad [x] \otimes [y] \mapsto [x \otimes y].$$

Both $\pi_0: \mathrm{Sp}_{\geq 0} \rightarrow \mathrm{Ab}$ and the symmetric monoidal inclusion $\mathrm{Sp}_{\geq 0} \hookrightarrow \mathrm{Sp}$ preserve geometric realizations. Applying Proposition 19.2.11, proved in Part II, to these two functors shows, respectively, that π_0 preserves relative tensor products and that the relative tensor product of connective modules computed in $\mathrm{Sp}_{\geq 0}$ agrees with the one computed in Sp . In particular, the relative tensor product of connective modules is connective, and this gives the asserted isomorphism. The analogous statements hold for right modules.

The following cellular description of connective modules will also identify the heart of the module t-structure.

Proposition 8.2.8 ([Lur17, Proposition 7.1.1.13(1)]). *Let R be a connective associative ring spectrum. Then the ∞ -category $\mathrm{LMod}_{R, \geq 0}$ is the smallest subcategory of LMod_R which contains R and which is closed under colimits. The analogous statement holds for connective right R -modules.*

Proof. It is clear that $\mathrm{LMod}_{R, \geq 0}$ contains R and is closed under colimits. Conversely, given a connective left R -module M , we will show that M can be written as a colimit of a diagram of left R -modules

$$M(0) \rightarrow M(1) \rightarrow M(2) \rightarrow \dots$$

satisfying the following properties:

- (i) The R -module $M(0)$ is a coproduct of copies of R .
- (ii) For $i \geq 0$, there is an exact sequence

$$F[i] \rightarrow M(i) \rightarrow M(i+1),$$

where F is a coproduct of copies of R .

We will construct the modules $M(i)$ inductively, together with their maps $f_i: M(i) \rightarrow M$. In fact, we will inductively ensure that these maps satisfy the following third property:

- (iii) Let $i \geq 0$, and let $K(i)$ be the fiber of the map $M(i) \rightarrow M$. Then $\pi_j K(i) \simeq 0$ for $j < i$.

For $i = 0$, we can choose a map of left R -modules $M(0) \rightarrow M$ where $M(0)$ is a coproduct of copies of R , such that the induced map $\pi_0(M(0)) \rightarrow \pi_0(M)$ is surjective: for instance, we could take the coproduct indexed by the set $\pi_0(M)$. For the induction step, we first consider the exact sequence

$$K(i) \rightarrow M(i) \rightarrow M.$$

The group $\pi_i K(i)$ is a $\pi_0(R)$ -module, and we may choose a map of left R -modules $g: F[i] \rightarrow K(i)$ from a coproduct of copies of $R[i]$ such that the induced map $\pi_i(F[i]) \rightarrow \pi_i(K(i))$ is surjective. Let $\tilde{g}: F[i] \rightarrow M(i)$ be the composite of g with the fiber inclusion $K(i) \rightarrow M(i)$. We define $M(i+1)$ to be the pushout of the diagram

$$\begin{array}{ccc} F[i] & \xrightarrow{\tilde{g}} & M(i) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & M(i+1), \end{array}$$

so that $M(i+1)$ is the cofiber of $\tilde{g}$. Since the composite $F[i] \rightarrow K(i) \rightarrow M(i) \rightarrow M$ is canonically nullhomotopic, there is an induced map $f_{i+1}: M(i+1) \rightarrow M$. The induced fiber sequence

$$F[i] \longrightarrow K(i) \longrightarrow K(i+1)$$

shows that $\pi_j K(i+1) = 0$ for $j < i$, while the surjectivity of $\pi_i(F[i]) \rightarrow \pi_i(K(i))$ gives $\pi_i K(i+1) = 0$. This verifies the inductive step.

It remains to prove that the natural map $\operatorname{colim}_i M(i) \rightarrow M$ is an isomorphism. Since filtered colimits are exact in $\mathbf{Sp}$, the fiber K of this map is $\operatorname{colim}_i K(i)$. By property (iii), the group $\pi_j K(i)$ vanishes for $i > j$, and by Lemma 4.4.27 we conclude that $\pi_j(K) = 0$ for every j . Hence $K = 0$.

The same construction with right free modules proves the assertion for connective right R -modules. $\square$

Lemma 8.2.9. *Let R be a connective associative ring spectrum. Then the functor π_0 from Observation 8.2.7 restricts to equivalences*

$$\operatorname{LMod}_R^\heartsuit \xrightarrow{\sim} \operatorname{LMod}_{\pi_0(R)}(\operatorname{Ab}), \quad \operatorname{RMod}_R^\heartsuit \xrightarrow{\sim} \operatorname{RMod}_{\pi_0(R)}(\operatorname{Ab})$$

between the hearts of the module categories and the abelian categories of discrete modules over $\pi_0(R)$.

Proof. We prove the statement for left modules; the proof for right modules is identical. For essential surjectivity, write a discrete $\pi_0(R)$ -module M as a coequalizer of free modules. The two maps between the free modules lift to maps between coproducts of copies of R , since maps out of R are classified by elements of π_0 . Form their coequalizer $\tilde{M}$ in $\operatorname{LMod}_{R, \geq 0}$ and then apply $\tau_{\leq 0}$. Since π_0 preserves colimits of connective modules, we have $\pi_0(\tilde{M}) \cong M$, and hence $\tau_{\leq 0}\tilde{M}$ is a discrete R -module whose image under π_0 is M .

We will now show that the functor is fully faithful. Let $N \in \operatorname{LMod}_R^\heartsuit$, and consider the subcategory $C \subseteq \operatorname{LMod}_{R, \geq 0}$ spanned by those left R -modules M such that the map

$$\pi_0 \operatorname{Hom}_{\operatorname{LMod}_R}(M, N) \rightarrow \operatorname{Hom}_{\operatorname{LMod}_{\pi_0(R)}(\operatorname{Ab})}(\pi_0(M), \pi_0(N))$$

is an isomorphism. The subcategory C contains R . It is also closed under colimits: for connective M and discrete N , the mapping spectrum $\mathrm{hom}_R(M, N)$ is coconnective, and $\pi_0: \mathrm{Sp}_{\leq 0} \rightarrow \mathrm{Ab}$ preserves limits. Thus both sides of the displayed map carry colimits in M to limits. It follows from Proposition 8.2.8 that $C = \mathrm{LMod}_{R, \geq 0}$, as desired. $\square$

Notation 8.2.10. Let R be a discrete associative ring. We will from now on denote the categories $\mathrm{LMod}_R(\mathrm{Ab})$ and $\mathrm{RMod}_R(\mathrm{Ab})$ of discrete left and right R -modules by $\mathrm{LMod}_R^\heartsuit$ and $\mathrm{RMod}_R^\heartsuit$, respectively. This notation is justified by Lemma 8.2.9: these categories agree with the hearts of the t-structures on $\mathrm{LMod}_R = \mathrm{LMod}_R(\mathrm{Sp})$ and $\mathrm{RMod}_R = \mathrm{RMod}_R(\mathrm{Sp})$.

Tensor products in Ab will from now on always be denoted by $\otimes^\heartsuit$; we reserve $\otimes$ for the tensor product in Sp .

Remark 8.2.11. The ∞ -category LMod_R also admits a t-structure when R is not connective. In this case, the subcategory $\mathrm{LMod}_{R, \geq 0}$ may still be described as the smallest subcategory of LMod_R that contains R and is closed under colimits. The subcategory $\mathrm{LMod}_{R, \leq 0}$ is still given by those R -modules whose underlying spectrum is coconnective. We refer to [Lur17, Proposition 1.4.4.11] for a very general result for producing t-structures on presentable ∞ -categories.

8.3 Eilenberg–MacLane ring spectra and DGAs

The symmetric monoidality of π_0 has an important consequence for the Eilenberg–MacLane functor.

Corollary 8.3.1. *The Eilenberg–MacLane functor $H: \mathrm{Ab} \rightarrow \mathrm{Sp}_{\geq 0}$ is canonically lax symmetric monoidal, hence induces functors*

$$H: \mathrm{Alg}(\mathrm{Ab}) \rightarrow \mathrm{Alg}(\mathrm{Sp}) \quad \text{and} \quad H: \mathrm{CAlg}(\mathrm{Ab}) \rightarrow \mathrm{CAlg}(\mathrm{Sp}).$$

Proof. Since H is right adjoint to the symmetric monoidal functor $\pi_0: \mathrm{Sp}_{\geq 0} \rightarrow \mathrm{Ab}$, it follows from Proposition 14.3.6 that H is canonically lax symmetric monoidal. Every lax symmetric monoidal functor preserves associative and commutative algebra objects, as does the symmetric monoidal inclusion $\mathrm{Sp}_{\geq 0} \hookrightarrow \mathrm{Sp}$. $\square$

Thus every ordinary associative ring A determines an associative ring spectrum HA , and every ordinary commutative ring A determines a commutative ring spectrum HA . In either case, the underlying spectrum is the Eilenberg–MacLane spectrum from Definition 6.2.1.

To incorporate differential graded algebras, we pass through the derived ∞ -category. Let R be a commutative ring. The derived tensor product equips $D(R)$ with a symmetric monoidal structure

$$- \otimes_R^L -: D(R) \times D(R) \rightarrow D(R)$$

whose monoidal unit is $R[0]$. The ∞ -category $D(R)$ is stable and cocomplete, and the derived tensor product preserves colimits separately in both variables. In Proposition 20.2.2 we construct this symmetric monoidal structure as a monoidal Dwyer–Kan localization of $\mathrm{Ch}(R)$. Its compact objects are the perfect complexes, and Corollary 20.2.5 gives an equivalence

$$D(R) \simeq \mathrm{Ind}(\mathrm{Perf}(R)).$$

In particular, $D(R)$ is presentable by Proposition 22.3.6.

We can therefore form the ∞ -categories

$$\mathrm{Alg}(D(R)) \quad \text{and} \quad \mathrm{CAlg}(D(R))$$

of associative and commutative algebra objects in $D(R)$. These objects should be thought of as derived associative and commutative R -algebras. One source of such objects is provided by differential graded algebras.

Definition 8.3.2 (Differential graded algebra). Let R be a commutative ring. An *associative differential graded R -algebra*, or *associative DGA over R* , is an associative algebra object in the symmetric monoidal category $\mathrm{Ch}(R)$ of chain complexes of R -modules. Concretely, it consists of a chain complex $A_\bullet$, a unit $R[0] \rightarrow A_\bullet$, and an associative multiplication

$$A_p \otimes_R A_q \longrightarrow A_{p+q}, \quad x \otimes y \longmapsto xy,$$

which satisfies the Leibniz rule

$$d(xy) = d(x)y + (-1)^p x d(y)$$

for homogeneous $x \in A_p$ and $y \in A_q$.

A *commutative DGA over R* is a commutative algebra object in $\mathrm{Ch}(R)$. Equivalently, it is an associative DGA whose multiplication is graded commutative:

$$xy = (-1)^{pq} yx$$

for homogeneous $x \in A_p$ and $y \in A_q$.

Example 8.3.3. Every associative R -algebra A defines an associative DGA $A[0]$ concentrated in degree 0. If A is commutative, then $A[0]$ is a commutative DGA. As an example with non-zero differential, let $a \in R$ and consider the Koszul DGA

$$K_R(a) := (\Lambda_R(e), d), \quad |e| = 1, \quad d(e) = a.$$

Here $\Lambda_R(e)$ denotes the exterior algebra on one generator, so this is a commutative DGA in homological degrees 1 and 0.

In Part II we show that the localization functor $\gamma: \mathrm{Ch}(R) \rightarrow D(R)$ admits a lax symmetric monoidal refinement (Corollary 20.2.4). Since lax symmetric monoidal functors preserve algebra objects, it induces functors

$$\mathrm{Alg}(\mathrm{Ch}(R)) \longrightarrow \mathrm{Alg}(D(R)) \quad \text{and} \quad \mathrm{CAlg}(\mathrm{Ch}(R)) \longrightarrow \mathrm{CAlg}(D(R)).$$

Thus every associative or commutative DGA determines a corresponding derived algebra. It is a somewhat surprising fact that the first functor is essentially surjective [Lur17, Proposition 7.1.4.6], while the second one is so when R contains the field $\mathbb{Q}$ of rational numbers [Lur17, Proposition 7.1.4.11].

The universal property of the symmetric monoidal structure on spectra recalled in Example 8.1.2 now produces a unique colimit-preserving symmetric monoidal functor

$$L_R: \mathrm{Sp} \longrightarrow D(R).$$

It sends the sphere spectrum to the monoidal unit $R[0]$. Since $D(R)$ and Sp are presentable, Theorem 22.2.5 gives a right adjoint to L_R . The universal enrichment of stable ∞ -categories in spectra identifies this right adjoint as

$$H_R: D(R) \longrightarrow \mathrm{Sp}, \quad H_R(A) = \mathrm{hom}_{D(R)}(R[0], A).$$

By Proposition 14.3.6, the right adjoint H_R has a canonical lax symmetric monoidal structure. It consequently induces functors

$$\mathrm{Alg}(D(R)) \longrightarrow \mathrm{Alg}(\mathrm{Sp}) \quad \text{and} \quad \mathrm{CAlg}(D(R)) \longrightarrow \mathrm{CAlg}(\mathrm{Sp}).$$

For $R = \mathbb{Z}$, the underlying functor of H_R is the Eilenberg–MacLane functor from Definition 6.2.1, since both are defined by the formula $A \mapsto \mathrm{hom}_{D(\mathbb{Z})}(\mathbb{Z}[0], A)$.

The following comparison is an application of the monogenic Morita theorem from Part II. This theorem identifies a stable category generated by its monoidal unit with modules over the endomorphism ring spectrum of that unit.

Corollary 8.3.4 (Eilenberg–MacLane modules). *For every commutative ring R , there is a symmetric monoidal equivalence*

$$D(R) \xrightarrow{\sim} \mathrm{Mod}_{H_R}(\mathrm{Sp}).$$

Under this equivalence, the functor $H_R = \mathrm{hom}_{D(R)}(R[0], -)$ agrees with the forgetful functor $\mathrm{Mod}_{H_R} \rightarrow \mathrm{Sp}$, and the lax symmetric monoidal structure on H_R sends the unit $R[0]$ to the Eilenberg–MacLane ring spectrum HR from Corollary 8.3.1.

Proof. Equip $D(R)$ with the symmetric monoidal structure of Proposition 20.2.2. Its unit $R[0]$ is a compact generator by Corollary 20.2.5. Moreover,

$$\pi_n \mathrm{hom}_{D(R)}(R[0], R[0]) \cong H_n(R[0])$$

by Corollary 6.4.5. This is R for $n = 0$ and zero otherwise. Thus the resulting commutative endomorphism ring spectrum is concentrated in degree 0. The adjunction $\pi_0: \mathrm{Sp}_{\geq 0} \rightleftarrows \mathrm{Ab} : H$ is symmetric monoidal by Proposition 8.2.6 and Corollary 8.3.1. The induced localization of commutative algebras from Part II (Example 14.5.5), together with the fully faithful symmetric monoidal inclusion $\mathrm{Sp}_{\geq 0} \hookrightarrow \mathrm{Sp}$, identifies commutative ring spectra concentrated in degree 0 with ordinary commutative rings. The multiplication on π_0 of the endomorphism ring spectrum is composition of endomorphisms of $R[0]$, hence the ordinary multiplication of R . It is therefore the Eilenberg–MacLane ring spectrum HR . The result now follows from Theorem 19.5.6. The description of H_R is built into the comparison functor in that theorem. $\square$

Definition 8.3.5 (Eilenberg–MacLane ring spectrum). Let A be an associative DGA over a commutative ring R . Its *Eilenberg–MacLane ring spectrum* is the associative ring spectrum

$$HA := H_R(\gamma A) \in \mathrm{Alg}(\mathrm{Sp}).$$

If A is a commutative DGA, the lax symmetric monoidal structure on $H_R \circ \gamma$ refines HA to a commutative ring spectrum.

Remark 8.3.6 (Agreement of the Eilenberg–MacLane constructions). For an ordinary associative R -algebra A , the ring spectrum $H_R(A[0])$ is canonically equivalent to the direct Eilenberg–MacLane ring spectrum HA from Corollary 8.3.1. Indeed, both are concentrated in degree 0, and their multiplications induce the ordinary multiplication of A on π_0 . The classification of associative ring spectra concentrated in degree 0 from Corollary 14.5.4 therefore identifies them. If A is commutative, the same argument gives an equivalence of commutative ring spectra.

Corollary 8.3.7. *Let R be a commutative ring and let M and N be R -modules, regarded as complexes concentrated in degree 0. Write $\mathrm{Tor}_n^R(M, N)$ and $\mathrm{Ext}_R^n(M, N)$ for their classical Tor- and Ext-groups. Equip HM and HN with their natural HR -module structures. Under the equivalence of Corollary 8.3.4, there are natural isomorphisms*

$$\mathrm{Tor}_n^R(M, N) \cong \pi_n(HM \otimes_{HR} HN) \quad \text{and} \quad \mathrm{Ext}_R^n(M, N) \cong \pi_{-n} \mathrm{hom}_{HR}(HM, HN),$$

Proof. The symmetric monoidal equivalence $D(R) \simeq \mathrm{Mod}_{HR}$ identifies $M[0] \otimes_R^L N[0]$ with $HM \otimes_{HR} HN$ and the derived hom with $\mathrm{hom}_{HR}(HM, HN)$. Their homotopy groups compute the classical derived functors by the resolution formulas of Corollary 6.6.27 and Proposition 6.4.14. $\square$

Important examples include $H\mathbb{Z}$, $H\mathbb{Q}$, and $H\mathbb{F}_2$. Their multiplicative structures underlie the products in ordinary cohomology and will be used later in the discussion of orientations, Thom isomorphisms, Poincaré duality, and the Chern character.

8.4 Localizations of ring spectra

In classical algebra, given a commutative ring R and a multiplicative subset $S \subseteq R$, one can form the localization $R[S^{-1}]$, a new ring in which every element of S becomes invertible. We now develop the analogous construction for commutative ring spectra. This is the form needed in Chapter 9, where periodic complex K-theory is obtained by inverting the Bott element in connective complex K-theory.

We have already met a special case in Section 7.2: inverting a set of primes P on the sphere spectrum gives a mapping telescope $\mathbb{S}[P^{-1}]$, and tensoring with this telescope is the corresponding reflection. The same two features persist here. Localization is smashing, and localization at a single homogeneous element is computed by a telescope.

We first treat commutative ring spectra, which is the only case used elsewhere in the book. The online version then continues with the localization of associative ring spectra at multiplicative subsets satisfying an Ore condition.

8.4.1 The commutative case

Let R be a commutative ring spectrum. Its homotopy groups form a graded-commutative ring, and the homotopy groups of an R -module form a graded module over it.

Definition 8.4.1. Let $S \subseteq \pi_*(R)$ be a subset of homogeneous elements. We say that S is *multiplicative* if it contains the unit $1 \in \pi_0(R)$ and is closed under multiplication.

For a homogeneous element $s \in \pi_d(R)$, multiplication by s is already defined at the level of R -modules:

Observation 8.4.2. Multiplication by $s \in \pi_d(R)$ is the natural transformation of functors $\text{Mod}_R \rightarrow \text{Mod}_R$ given by

$$s: M[d] \simeq R[d] \otimes_R M \xrightarrow{s \otimes \text{id}} R \otimes_R M \simeq M.$$

On homotopy groups this is the map $x \mapsto s \cdot x$.

Definition 8.4.3. Let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements. An R -module N is S -local if multiplication by every $s \in S$ is an isomorphism

$$s: N[|s|] \xrightarrow{\sim} N.$$

Equivalently, multiplication by every $s \in S$ is an isomorphism on the graded group $\pi_*(N)$. We write $\text{Mod}_R^{\text{Loc}(S)} \subseteq \text{Mod}_R$ for the full subcategory of S -local modules.

Proposition 8.4.4 (Commutative localization is smashing). *Let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements.*

(1) *The inclusion $\text{Mod}_R^{\text{Loc}(S)} \hookrightarrow \text{Mod}_R$ admits a left adjoint*

$$S^{-1}(-): \text{Mod}_R \longrightarrow \text{Mod}_R^{\text{Loc}(S)}.$$

The subcategory $\text{Mod}_R^{\text{Loc}(S)}$ inherits a symmetric monoidal structure for which this localization functor is symmetric monoidal and the inclusion is lax symmetric monoidal.

(2) *The localization is smashing: for every $M \in \text{Mod}_R$, the map*

$$M \simeq M \otimes_R R \xrightarrow{\text{id}_M \otimes \eta_R} M \otimes_R S^{-1}R$$

exhibits $M \otimes_R S^{-1}R$ as the S -localization of M . In particular,

$$S^{-1}M \simeq M \otimes_R S^{-1}R$$

naturally in M .

Proof. For every $s \in S$ of degree d and every $n \in \mathbb{Z}$, consider the shifted multiplication map

$$s[n]: R[d+n] \longrightarrow R[n].$$

An R -module is local with respect to this small collection of morphisms precisely when it is S -local: allowing all shifts means that locality tests multiplication by s on every homotopy group. Since Mod_R is presentable by Proposition 22.5.5, Theorem 22.2.2(4) provides the asserted reflection $S^{-1}(-)$.

The symmetric monoidal ∞ -category Mod_R is closed, as recalled after Corollary 19.2.17. If N is S -local and P is any R -module, then $\underline{\text{Hom}}_R(P, N)$ is again S -local. Indeed, under the natural isomorphism

$$\underline{\text{Hom}}_R(P, N)[d] \xrightarrow{\sim} \underline{\text{Hom}}_R(P, N[d]),$$

multiplication by s is obtained by applying $\underline{\mathrm{Hom}}_R(P, -)$ to the isomorphism $s: N[d] \xrightarrow{\sim} N$. The internal-hom criterion Lemma 14.5.6 therefore gives the symmetric monoidal structure and the symmetric monoidality of the localization.

It remains to prove the smashing formula. The unit $\eta_R: R \rightarrow S^{-1}R$ is an S -local equivalence, and the proof of the internal-hom criterion shows that S -local equivalences remain so after tensoring with any R -module. Consequently,

$$\mathrm{id}_M \otimes \eta_R: M \longrightarrow M \otimes_R S^{-1}R$$

is an S -local equivalence. Its target is S -local: multiplication by s on $M \otimes_R S^{-1}R$ is obtained by tensoring id_M with the isomorphism $s: (S^{-1}R)[[s]] \xrightarrow{\sim} S^{-1}R$. A local equivalence from M to a local object exhibits that object as the reflection of M , which proves the claim. $\square$

We can now localize in commutative ring spectra.

Corollary 8.4.5 (Localization of a commutative ring spectrum). *Let R be a commutative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements. Then the R -module $S^{-1}R$ carries a canonical structure of a commutative ring spectrum $R[S^{-1}]$, and the localization map refines to a morphism*

$$\eta: R \longrightarrow R[S^{-1}]$$

in $\mathrm{CAlg}(\mathrm{Sp})$. It has the following universal property: for every commutative ring spectrum A , precomposition with η induces a monomorphism of animae

$$- \circ \eta: \mathrm{Hom}_{\mathrm{CAlg}(\mathrm{Sp})}(R[S^{-1}], A) \hookrightarrow \mathrm{Hom}_{\mathrm{CAlg}(\mathrm{Sp})}(R, A)$$

whose image consists of those maps $f: R \rightarrow A$ for which $\pi_(f)$ carries every element of S to a unit of $\pi_*(A)$.*

Proof. By Proposition 8.4.4, the reflection

$$S^{-1}(-): \mathrm{Mod}_R \longrightarrow \mathrm{Mod}_R^{\mathrm{Loc}(S)}$$

is a symmetric monoidal Bousfield localization. The Part II result Corollary 19.2.17 therefore equips $S^{-1}R$ with a commutative R -algebra structure and characterizes it as initial among commutative R -algebras whose underlying modules are S -local. Under the equivalence

$$\mathrm{CAlg}(\mathrm{Mod}_R) \simeq \mathrm{CAlg}(\mathrm{Sp})_{R/},$$

this gives the asserted morphism $\eta: R \rightarrow R[S^{-1}]$ of commutative ring spectra.

Let $f: R \rightarrow A$ be a morphism of commutative ring spectra. Its underlying R -module is S -local if and only if $\pi_*(f)$ sends every $s \in S$ to a unit. One direction is immediate. Conversely, suppose that A is S -local and let $s' = \pi_*(f)(s)$ have degree d . Since multiplication by s' is bijective, there is an element $u \in \pi_{-d}(A)$ such that $s'u = 1$. Graded commutativity gives $us' = (-1)^d$, so $(-1)^d u$ is a left inverse of s' , while u is a right inverse. Since left and right inverses coincide, s' is a unit.

The initial property of $S^{-1}R$ among commutative R -algebras with local underlying module now gives the claimed factorization property. Conversely, every morphism that factors through $R[S^{-1}]$ inverts S , since the image of S in $\pi_*(R[S^{-1}])$ consists of units and ring morphisms preserve units. $\square$

Finally, we record the computation used for Bott inversion.

Lemma 8.4.6 (Inverting a single element). *Let R be a commutative ring spectrum, let $s \in \pi_d(R)$ be a homogeneous element, and let $S := \{1, s, s^2, \dots\}$ be the multiplicative subset it generates. Then for every $M \in \text{Mod}_R$ the canonical map exhibits the telescope*

$$M[s^{-1}] := \text{colim}(M \xrightarrow{s} M[-d] \xrightarrow{s} M[-2d] \xrightarrow{s} \dots)$$

as the S -localization of M . On homotopy groups it is given by graded localization:

$$\pi_*(M[s^{-1}]) \cong \pi_*(M)[s^{-1}].$$

Proof. The transition maps are shifts of the module map of Observation 8.4.2, so the telescope is a colimit in Mod_R . Since homotopy groups preserve filtered colimits by Lemma 4.4.27, we have

$$\pi_k(M[s^{-1}]) \cong \text{colim}(\pi_k(M) \xrightarrow{s} \pi_{k+d}(M) \xrightarrow{s} \pi_{k+2d}(M) \longrightarrow \dots),$$

which is the degree- k part of $\pi_*(M)[s^{-1}]$. In particular, multiplication by s on $M[s^{-1}]$ is an isomorphism, so the telescope is S -local.

Let N be any S -local R -module. Mapping out of the telescope gives

$$\text{Hom}_{\text{Mod}_R}(M[s^{-1}], N) \simeq \lim_j \text{Hom}_{\text{Mod}_R}(M[-jd], N).$$

Every transition map in this limit is an equivalence, because under the shift adjunction it is induced by the isomorphism $s : N[d] \xrightarrow{\sim} N$. Evaluation at the zeroth term therefore gives an equivalence

$$\text{Hom}_{\text{Mod}_R}(M[s^{-1}], N) \xrightarrow{\sim} \text{Hom}_{\text{Mod}_R}(M, N),$$

induced by precomposition with the canonical map $M \rightarrow M[s^{-1}]$. This is the universal property of the S -localization. $\square$

Remark 8.4.7. Combining Corollary 8.4.5 with Lemma 8.4.6 for $M = R$, the telescope

$$R[s^{-1}] = \text{colim}(R \xrightarrow{s} R[-d] \xrightarrow{s} R[-2d] \longrightarrow \dots)$$

is a commutative ring spectrum, the map $R \rightarrow R[s^{-1}]$ is a morphism of commutative ring spectra, and

$$\pi_*(R[s^{-1}]) \cong \pi_*(R)[s^{-1}].$$

8.4.2 Ore localization (online only)

The commutative theory above is the form used elsewhere in the book. We conclude by discussing the more general localization problem for associative ring spectra. In this setting the graded ring $\pi_*(R)$ need not be commutative, and a multiplicative subset must satisfy an Ore condition for localization to have the expected properties.

Definition 8.4.8 (Localization of a ring spectrum). Let R be a ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements. The *localization of R at S* , denoted $R[S^{-1}]$, is

a ring spectrum equipped with a map of ring spectra $\eta: R \rightarrow R[S^{-1}]$ that satisfies the following property: for every ring spectrum A , precomposition with η induces a monomorphism of anima

$$- \circ \eta: \operatorname{Hom}_{\operatorname{Alg}}(R[S^{-1}], A) \hookrightarrow \operatorname{Hom}_{\operatorname{Alg}}(R, A)$$

whose image consists of those algebra maps $f: R \rightarrow A$ for which $\pi_*(f)$ carries every element of S to a unit of the graded ring $\pi_*(A)$.

In other words: an algebra map $f: R \rightarrow A$ factors through η if and only if it inverts S , and in that case the factorization is unique. Note that $R[S^{-1}]$ is unique, if it exists. In an analogous way, we may define localization at the level of modules. Let us first introduce some terminology:

Definition 8.4.9. Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements.

- (1) A left R -module M is called *S -nilpotent* if for every element $x \in \pi_n(M)$, there exists an element $s \in S$ such that $s \cdot x = 0$ in $\pi_*(M)$.
- (2) A left R -module N is called *S -local* if, for every element $s \in S$ of degree d and every $k \in \mathbb{Z}$, left multiplication by s induces an isomorphism

$$s \cdot -: \pi_k(N) \xrightarrow{\sim} \pi_{k+d}(N).$$

We denote by $\operatorname{LMod}_R^{S\text{-nil}}$ and $\operatorname{LMod}_R^{\operatorname{Loc}(S)}$ the full subcategories of LMod_R spanned by the S -nilpotent and S -local modules, respectively.

When R is commutative, the second condition agrees with the definition given in the previous subsection. For a general associative ring spectrum, however, left multiplication by s on N need not be a morphism of left R -modules, which explains the formulation in terms of homotopy groups. Equivalently, the underlying map of spectra $s: N[|s|] \rightarrow N$ is an isomorphism for every $s \in S$. There is also an intrinsic reformulation in LMod_R : right multiplication by $s \in \pi_d(R)$ defines a morphism $\cdot s: R[d] \rightarrow R$ of left R -modules, and N is S -local precisely when it is local with respect to all shifts of these morphisms.

Both conditions are formulated purely in terms of the graded $\pi_*(R)$ -module $\pi_*(M)$, and this makes them stable under all the operations we will need:

Observation 8.4.10. The subcategories $\operatorname{LMod}_R^{S\text{-nil}}$ and $\operatorname{LMod}_R^{\operatorname{Loc}(S)}$ are stable subcategories of LMod_R , closed under all small colimits.

Proof. Throughout we use that for a homogeneous element $s \in \pi_d(R)$ the operation $x \mapsto s \cdot x$ is a natural transformation $\pi_k(-) \rightarrow \pi_{k+d}(-)$ of functors $\operatorname{LMod}_R \rightarrow \operatorname{Ab}$, and that π_* carries exact sequences of left R -modules to long exact sequences of graded $\pi_*(R)$ -modules.

Both subcategories evidently contain 0 and are closed under shifts. For closure under cofibers, consider an exact sequence $M' \rightarrow M \rightarrow M''$ in LMod_R .

If M' and M are S -local, then in the long exact sequence

$$\pi_*(M') \rightarrow \pi_*(M) \rightarrow \pi_*(M'') \rightarrow \pi_{*-1}(M') \rightarrow \pi_{*-1}(M)$$

multiplication by s acts as an isomorphism on four of the five terms, so the five lemma shows that it is an isomorphism on $\pi_*(M'')$ as well; hence M'' is S -local. The same argument applied to the rotated sequence shows closure under fibers, so $\mathbf{LMod}_R^{\mathrm{Loc}(S)}$ is stable.

If M' and M are S -nilpotent, let $x \in \pi_n(M'')$. Its image $\partial(x) \in \pi_{n-1}(M')$ is annihilated by some $s \in S$, so $\partial(s \cdot x) = s \cdot \partial(x) = 0$ and therefore $s \cdot x$ lifts to an element $y \in \pi_{n+|s|}(M)$. Choosing $t \in S$ with $t \cdot y = 0$, we obtain $(ts) \cdot x = t \cdot (s \cdot x) = 0$, and $ts \in S$ since S is multiplicative. Hence M'' is S -nilpotent.

Similarly, suppose that M' and M'' are S -nilpotent and let $x \in \pi_n(M)$. Choose $s \in S$ which annihilates the image of x in $\pi_n(M'')$. Then $s \cdot x$ is the image of an element $y \in \pi_{n+|s|}(M')$. Choosing $t \in S$ with $t \cdot y = 0$ gives $(ts) \cdot x = 0$. Thus M is S -nilpotent as well, so $\mathbf{LMod}_R^{S\text{-nil}}$ is stable and closed under extensions.

Being stable, both subcategories are in particular closed under finite coproducts: a biproduct $M_1 \oplus M_2$ sits in an exact sequence $M_1 \rightarrow M_1 \oplus M_2 \rightarrow M_2$, and both subcategories are closed under extensions. Consequently both are closed under all finite colimits.

Next, both are closed under filtered colimits, since π_* preserves these by Lemma 4.4.27. Indeed, a filtered colimit of isomorphisms is an isomorphism, which handles the S -local case; and in the S -nilpotent case every element of $\pi_n(\mathrm{colim}_i M_i) \cong \mathrm{colim}_i \pi_n(M_i)$ is the image of some $x \in \pi_n(M_i)$, and any $s \in S$ annihilating x also annihilates its image. Since an arbitrary coproduct is the filtered colimit of its finite subcoproducts, both subcategories are closed under small coproducts. As an ∞ -category admitting pushouts and small coproducts admits all small colimits, this completes the proof. $\square$

Warning 8.4.11. The finite-coproduct step above is the one place where some care is needed: given $x_1 \in \pi_n(M_1)$ annihilated by s_1 and $x_2 \in \pi_n(M_2)$ annihilated by s_2 , the product $s_2 s_1$ need *not* annihilate (x_1, x_2) , since $s_2 s_1 \cdot x_2 = s_2 \cdot (s_1 \cdot x_2)$ and $s_1 \cdot x_2$ has no reason to vanish. What saves the argument is that we may instead first apply s_2 , killing the second coordinate, and then use the S -nilpotence of M_1 to annihilate the element $s_2 \cdot x_1$ (rather than x_1 itself). No commutativity or Ore condition is needed.

Definition 8.4.12. Let R be a ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements. Given a left R -module M , its S -localization is an S -local left R -module $S^{-1}M$ equipped with a morphism of left R -modules $\eta: M \rightarrow S^{-1}M$ satisfying the property that for every other S -local left R -module N , precomposition with η induces an equivalence of anima

$$- \circ \eta: \mathrm{Hom}_{\mathbf{LMod}_R}(S^{-1}M, N) \xrightarrow{\sim} \mathrm{Hom}_{\mathbf{LMod}_R}(M, N).$$

Again, $S^{-1}M$ is unique whenever it exists. In the non-commutative setting, this abstract operation of formally inverting elements can be poorly behaved. To ensure a good theory, we need an additional condition on the set S , known as the Ore condition.

Definition 8.4.13 ([Lur17, Definition 7.2.3.1, Remark 7.2.3.7]). Let A_* be a graded associative ring and let $S \subseteq A_*$ be a set of homogeneous elements which contains the unit and is closed under multiplication. We say that S satisfies the *left Ore condition* if the following hold:

- (a) For every pair of elements $x \in A_*$ and $s \in S$, there exist elements $y \in A_*$ and $t \in S$ such that $tx = ys$.

(b) If $xs = 0$ for some $x \in A_*$ and $s \in S$, then there exists some $t \in S$ such that $tx = 0$.

It suffices to check these conditions for homogeneous x , in which case y in part (a) may also be chosen homogeneous. An ordinary ring is regarded as concentrated in degree zero.

This seemingly purely algebraic condition can be conveniently rephrased in terms of S -nilpotent modules. For an element $s \in S$ of degree d , we denote by R/Rs the cofiber of the map of left R -modules $\cdot s: R[d] \rightarrow R$ given by right multiplication by s , i.e., the cofiber of the following composite

$$\cdot s: R[d] \simeq R \otimes \mathbb{S}^d \xrightarrow{\text{id} \otimes s} R \otimes R \xrightarrow{m} R.$$

Lemma 8.4.14 ([Lur17, Lemma 7.2.3.11]). *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset of homogeneous elements. The following conditions are equivalent:*

- (1) *The set S satisfies the left Ore condition.*
- (2) *For every element $s \in S$, the left R -module R/Rs is S -nilpotent.*

Proof. Assume first that (1) is satisfied. Let $x \in \pi_n(R/Rs)$; we wish to show that it is annihilated by some element of S . Using the exact sequence

$$\pi_n(R/Rs) \xrightarrow{\varphi} \pi_{n-d-1}(R) \xrightarrow{\cdot s} \pi_{n-1}(R),$$

we deduce that $\varphi(x) \in \pi_{n-d-1}(R)$ is annihilated by right multiplication by s . Using condition (b) of the left Ore condition, we deduce that there exists an element $t \in S$ of degree d' such that $0 = t\varphi(x) = \varphi(tx) \in \pi_{n+d'-d-1}(R)$. Using the exactness of the sequence

$$\pi_{n+d'}(R) \xrightarrow{\psi} \pi_{n+d'}(R/Rs) \xrightarrow{\varphi} \pi_{n+d'-d-1}(R),$$

we conclude that $tx = \psi(y)$ for some $y \in \pi_{n+d'}(R)$. Using condition (a) of the left Ore condition, we can find an element $u \in S$ of degree d'' and an element $z \in \pi_{n+d'+d''-d}(R)$ such that $uy = zs$. It follows that the image of uy in $\pi_{n+d'+d''}(R/Rs)$ vanishes, so that $(ut)x = u(tx) = u\psi(y) = \psi(uy) = 0$. This completes the proof that (1) implies (2).

Now suppose that (2) is satisfied. We will show that S satisfies conditions (a) and (b) of the left Ore condition. For (a), suppose that $x \in \pi_*(R)$ is a homogeneous element of degree n and $s \in S$ has degree d . We wish to show that there exist $y \in \pi_*(R)$ and $t \in S$ such that $tx = ys$. By S -nilpotence, the image of x in $\pi_n(R/Rs)$ is annihilated by multiplication by some homogeneous element $t \in S$ of degree d' . It follows that tx belongs to the kernel of the map $\pi_{n+d'}(R) \rightarrow \pi_{n+d'}(R/Rs)$, and therefore to the image of the map $\cdot s: \pi_{n+d'-d}(R) \rightarrow \pi_{n+d'}(R)$ given by right multiplication by s . Thus $tx = ys$ for some $y \in \pi_{n+d'-d}(R)$, which proves (a).

We now verify (b). It suffices to show that if $x \in \pi_n(R)$ is annihilated by right multiplication by some element $s \in S$ of degree d , then $tx = 0$ for some $t \in S$. The exactness of the sequence

$$\pi_{n+d+1}(R/Rs) \xrightarrow{\varphi} \pi_n(R) \xrightarrow{\cdot s} \pi_{n+d}(R)$$

shows that $x = \varphi(y)$ for some $y \in \pi_{n+d+1}(R/Rs)$. Since R/Rs is S -nilpotent, there exists an element $t \in S$ such that $ty = 0$, from which it follows immediately that $tx = t\varphi(y) = \varphi(ty) = 0$. $\square$

The previous lemma says that under the Ore condition the modules R/Rs are S -nilpotent. In fact they generate all S -nilpotent modules, in a strong and rather explicit sense. This is the one genuinely difficult input into the theory, and it is the engine behind every result in this section. We use this result as a black box.

Lemma 8.4.15 ([Lur17, Lemma 7.2.3.13]). *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset satisfying the left Ore condition. Then every S -nilpotent left R -module M can be written as the colimit of a sequence of left R -modules*

$$0 = M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow \cdots$$

in which the fiber of each map $M_i \rightarrow M_{i+1}$ is a coproduct of modules of the form $(R/Rs_\alpha)[n_\alpha]$ with $s_\alpha \in S$ and $n_\alpha \in \mathbb{Z}$. In particular, $\mathrm{LMod}_R^{S\text{-nil}}$ is generated under colimits by the set of objects

$$\{ (R/Rs)[n] \mid s \in S, n \in \mathbb{Z} \}.$$

Proposition 8.4.16 ([Lur17, Proposition 7.2.3.14]). *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset satisfying the left Ore condition. Let N be a left R -module. Then the following conditions are equivalent:*

- (1) *The module N is S -local.*
- (2) *For every element $s \in S$ and every integer n , the hom anima $\mathrm{Hom}_{\mathrm{LMod}_R}((R/Rs)[n], N)$ is contractible.*
- (3) *For every S -nilpotent module $M \in \mathrm{LMod}_R$, the hom anima $\mathrm{Hom}_{\mathrm{LMod}_R}(M, N)$ is contractible.*

Proof. Let $s \in S$ be an element of degree d . Applying the mapping spectrum functor $\mathrm{hom}_{\mathrm{LMod}_R}(-, N)$ to the exact sequence

$$R[d] \xrightarrow{\cdot s} R \rightarrow R/Rs$$

and using the identifications $\mathrm{hom}_{\mathrm{LMod}_R}(R, N) \simeq N$ and $\mathrm{hom}_{\mathrm{LMod}_R}(R[d], N) \simeq N[-d]$ gives an exact sequence of spectra

$$\mathrm{hom}_{\mathrm{LMod}_R}(R/Rs, N) \longrightarrow N \xrightarrow{s \cdot -} N[-d].$$

The second map is left multiplication by s : precomposition with right multiplication $\cdot s: R[d] \rightarrow R$ sends a map $f: R \rightarrow N$ to the map determined by $f(s) = s \cdot f(1)$.

Condition (1) says precisely that the second map is an isomorphism for every $s \in S$, which by exactness is equivalent to the vanishing of $\mathrm{hom}_{\mathrm{LMod}_R}(R/Rs, N)$. Since a spectrum vanishes if and only if all the anima $\Omega^\infty(-[-n])$ do, this is in turn equivalent to $\mathrm{Hom}_{\mathrm{LMod}_R}((R/Rs)[n], N)$ being contractible for all n . This proves (1) $\Leftrightarrow$ (2).

To prove the implication (3) $\Rightarrow$ (2), we observe that since S satisfies the left Ore condition, the module R/Rs is S -nilpotent for every $s \in S$, as established in Lemma 8.4.14. As $\mathrm{LMod}_R^{S\text{-nil}}$ is closed under shifts (Observation 8.4.10), condition (3) applied to $M = (R/Rs)[n]$ immediately yields condition (2).

We now prove the implication (2) $\Rightarrow$ (3). Let M be an S -nilpotent left R -module, and write it as the colimit of a sequence

$$0 = M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow \cdots$$

whose successive fibers are coproducts of modules $(R/Rs_\alpha)[n_\alpha]$, as provided by Lemma 8.4.15. Since hom animae turn colimits in the first variable into limits, we have

$$\mathrm{Hom}_{\mathrm{LMod}_R}(M, N) \xrightarrow{\sim} \lim_i \mathrm{Hom}_{\mathrm{LMod}_R}(M_i, N).$$

It will therefore suffice to show that each hom anima $\mathrm{Hom}_{\mathrm{LMod}_R}(M_i, N)$ is contractible. We proceed by induction on i . The case $i = 0$ is clear, since $M_0 = 0$. For the inductive step, assume that $\mathrm{Hom}_{\mathrm{LMod}_R}(M_i, N)$ is contractible. Let K_i denote the fiber of the map $M_i \rightarrow M_{i+1}$. By construction, we have an exact sequence of modules $K_i \rightarrow M_i \rightarrow M_{i+1}$. Applying the functor $\mathrm{Hom}_{\mathrm{LMod}_R}(-, N)$ yields a fiber sequence of animae

$$\mathrm{Hom}_{\mathrm{LMod}_R}(M_{i+1}, N) \rightarrow \mathrm{Hom}_{\mathrm{LMod}_R}(M_i, N) \rightarrow \mathrm{Hom}_{\mathrm{LMod}_R}(K_i, N).$$

By the inductive hypothesis, the middle term is contractible. It therefore suffices to show that the right-hand term, $\mathrm{Hom}_{\mathrm{LMod}_R}(K_i, N)$, is also contractible. By construction, K_i is a coproduct of modules of the form $(R/Rs_\alpha)[n_\alpha]$. Since hom animae send coproducts in the first variable to products, we have

$$\mathrm{Hom}_{\mathrm{LMod}_R}(K_i, N) \xrightarrow{\sim} \prod_{\alpha} \mathrm{Hom}_{\mathrm{LMod}_R}((R/Rs_\alpha)[n_\alpha], N).$$

By condition (2), each factor in this product is contractible. It follows that the entire product is contractible, which completes the inductive step and the proof. $\square$

We will now show the existence of the S -localized R -modules $S^{-1}M$ when S satisfies the left Ore condition, by showing that any left R -module M decomposes into an S -nilpotent part and an S -local part:

Proposition 8.4.17 ([Lur17, Proposition 7.2.3.17, Remark 7.2.3.18]). *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset satisfying the left Ore condition. Then for every left R -module M , there exists an exact sequence in LMod_R*

$$M' \rightarrow M \rightarrow M'',$$

where M' is S -nilpotent and M'' is S -local.

Proof sketch. By Lemma 8.4.15, the ∞ -category $\mathrm{LMod}_R^{S\text{-nil}}$ is the localizing subcategory generated by the set of objects $\{(R/Rs)[n] \mid s \in S, n \in \mathbb{Z}\}$. It is therefore presentable by Proposition 22.2.7(2). Its inclusion into LMod_R preserves colimits by Observation 8.4.10, so Theorem 22.2.5 gives a right adjoint G . For any module M , the counit of this adjunction gives a map $G(M) \rightarrow M$. We define $M' := G(M)$ and we let M'' be the cofiber of this map. It remains to show that M'' is S -local, for which we use criterion (3) from Proposition 8.4.16. Let N be an S -nilpotent left R -module. Applying $\mathrm{hom}_{\mathrm{LMod}_R}(N, -)$ provides an exact sequence of spectra

$$\mathrm{hom}_{\mathrm{LMod}_R}(N, M') \rightarrow \mathrm{hom}_{\mathrm{LMod}_R}(N, M) \rightarrow \mathrm{hom}_{\mathrm{LMod}_R}(N, M'').$$

Both functors in the adjunction are exact, so the adjunction is enriched in spectra. The first map is therefore an isomorphism, and it follows that $\mathrm{hom}_{\mathrm{LMod}_R}(N, M'') = 0$, as desired. $\square$

Remark 8.4.18. The previous two results show that the pair $(\mathrm{LMod}_R^{S\text{-nil}}, \mathrm{LMod}_R^{\mathrm{Loc}(S)})$ satisfies the axioms of a t-structure on LMod_R in the sense of Definition 6.3.1: closure under shifts is Observation 8.4.10, orthogonality is part (3) of Proposition 8.4.16, and the decomposition axiom is Proposition 8.4.17.

This t-structure is, however, of a completely different nature from the ones in Chapter 6: both halves are *stable* subcategories, closed under *all* shifts, so that $\mathrm{LMod}_R^{S\text{-nil}}[n] = \mathrm{LMod}_R^{S\text{-nil}}$ for every n . In particular the heart is zero, and there is no connectivity intuition to be had here. What the pair really encodes is a *semiorthogonal decomposition* of LMod_R : every module is functorially glued from an S -nilpotent part and an S -local part, and there are no maps from the nilpotent part to the local part.

Corollary 8.4.19. *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ satisfy the left Ore condition. For every left R -module M , there exists an S -localization $S^{-1}M$. In particular, the inclusion $\mathrm{LMod}_R^{\mathrm{Loc}(S)} \hookrightarrow \mathrm{LMod}_R$ admits a left adjoint*

$$S^{-1}(-): \mathrm{LMod}_R \rightarrow \mathrm{LMod}_R^{\mathrm{Loc}(S)}.$$

Proof. By Proposition 8.4.17 there is an exact sequence $M' \rightarrow M \rightarrow M''$ of left R -modules such that M' is S -nilpotent and M'' is S -local. We claim that M'' is an S -localization. Indeed, for any other S -local left R -module N , mapping into N provides a fiber sequence of anima

$$\mathrm{Hom}_{\mathrm{LMod}_R}(M'', N) \rightarrow \mathrm{Hom}_{\mathrm{LMod}_R}(M, N) \rightarrow \mathrm{Hom}_{\mathrm{LMod}_R}(M', N).$$

But since M' is S -nilpotent, the anima $\mathrm{Hom}_{\mathrm{LMod}_R}(M', N)$ is contractible by part (3) of Proposition 8.4.16. It follows that the first map is an equivalence, as desired. $\square$

The proof exhibits the fiber of the localization map $\eta_M: M \rightarrow S^{-1}M$ as the S -nilpotent part M' of M . This gives the following recognition criterion:

Observation 8.4.20. Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ satisfy the left Ore condition. Then:

- (1) For $M \in \mathrm{LMod}_R$ we have $S^{-1}M = 0$ if and only if M is S -nilpotent.
- (2) A map $f: M \rightarrow N$ of left R -modules becomes an isomorphism after S -localization if and only if $\mathrm{cofib}(f)$ is S -nilpotent.
- (3) If T is an S -local module and $f: M \rightarrow T$ is a map with S -nilpotent cofiber, then f exhibits T as the S -localization of M .

Proof. (1) Write $M' \rightarrow M \rightarrow S^{-1}M$ for the decomposition of Proposition 8.4.17. If M is S -nilpotent, then the counit $M' = G(M) \rightarrow M$ of the coreflection is an isomorphism, so its cofiber $S^{-1}M$ vanishes. Conversely, if $S^{-1}M = 0$ then $M' \rightarrow M$ is an isomorphism, so M is S -nilpotent.

(2) The functor $S^{-1}(-)$ is a left adjoint between stable ∞ -categories, hence preserves cofibers. Thus $S^{-1}(f)$ is an isomorphism if and only if $S^{-1} \mathrm{cofib}(f) \simeq \mathrm{cofib}(S^{-1}f)$ vanishes, which by (1) happens if and only if $\mathrm{cofib}(f)$ is S -nilpotent.

(3) By (2) the map $S^{-1}(f): S^{-1}M \rightarrow S^{-1}T$ is an isomorphism, and the unit $\eta_T: T \rightarrow S^{-1}T$ is an isomorphism because T is already S -local. Naturality of the unit gives $\eta_T \circ f = S^{-1}(f) \circ \eta_M$. It follows that the isomorphism $\eta_T^{-1} \circ S^{-1}(f): S^{-1}M \rightarrow T$ identifies f with the localization map $\eta_M: M \rightarrow S^{-1}M$. $\square$

The existence of S -localizations of R -modules can be used to deduce the existence of localizations of ring spectra. Regarding R as a module over itself, applying the S -localization functor gives an S -local R -module $S^{-1}R \in \mathbf{LMod}_R$. This object can be endowed with the structure of an associative ring spectrum, which we denote by $R[S^{-1}]$.

Proposition 8.4.21 ([Lur17, Remark 7.2.3.26, Proposition 7.2.3.27]). *Let R be an associative ring spectrum and let $S \subseteq \pi_*(R)$ be a multiplicative subset satisfying the left Ore condition. The S -localization $S^{-1}R$ admits the structure of an associative ring spectrum, which we denote by $R[S^{-1}]$. Furthermore, the canonical map $R \rightarrow S^{-1}R$ can be enhanced to a morphism of associative ring spectra*

$$\varphi: R \rightarrow R[S^{-1}].$$

This morphism exhibits $R[S^{-1}]$ as an S -localization of R , in the sense of Definition 8.4.8.

We use this result as a black box, taking it directly from the cited reference. Constructing the coherently associative multiplication on $S^{-1}R$ is the one step of the theory for which the module-theoretic arguments above do not suffice.

Finally, this abstract homotopical construction recovers the classical construction of fractions when applied to an ordinary associative ring and module.

Proposition 8.4.22 ([Lur17, Proposition 7.2.3.20 and the proof of Proposition 7.2.3.5]). *Let R be an associative ring and let $S \subseteq R$ be a subset satisfying the left Ore condition. For every left R -module M , regard HR as a ring spectrum and HM as a left HR -module. Then $S^{-1}(HM)$ is discrete, and its zeroth homotopy group is naturally isomorphic to the classical left module of fractions $S^{-1}M$, constructed as the set of equivalence classes of symbols $s^{-1}x$ for $s \in S$ and $x \in M$.*

We use this result as a black box, taking it directly from the cited reference.

8.5 Projective and flat modules (online only)

In Sections 6.4 and 6.5 we developed the general theory of t -projective and t -flat objects in stable ∞ -categories equipped with t -structures. In this section, we specialize this theory to modules over ring spectra, obtaining analogues of projective and flat modules from ordinary algebra.

8.5.1 Projective modules

Definition 8.5.1 (Ext-groups for modules). Let R be an associative ring spectrum and let $M, N \in \mathbf{LMod}_R$ be left R -modules. We define the n -th Ext-group of M and N over R to be

$$\mathrm{Ext}_R^n(M, N) \quad := \quad \mathrm{Ext}_{\mathbf{LMod}_R}^n(M, N) \quad = \quad \pi_{-n} \mathrm{hom}_R(M, N) \quad \simeq \quad \pi_0 \mathrm{Hom}_{\mathbf{LMod}_R}(M, N[n]).$$

These are the Ext-groups from Definition 6.4.6 specialized to the stable ∞ -category $\mathbf{LMod}_R$.

The general vanishing, degree-zero identification, and long exact sequence for Ext-groups from Observations 6.4.7, 6.4.8 and 6.4.10 apply verbatim to R -modules.

Definition 8.5.2 (Projective module). Let R be a connective associative ring spectrum. A left R -module P is *projective* if it is a t-projective object of LMod_R in the sense of Definition 6.4.1; that is, if the mapping spectrum functor $\mathrm{hom}_R(P, -) : \mathrm{LMod}_R \rightarrow \mathrm{Sp}$ is t-exact.

Observation 8.5.3. The characterizations of t-projective objects from Proposition 6.4.11 specialize to the following equivalent conditions for a connective left R -module P :

- (1) P is projective;
- (2) For every connective left R -module Q and every $i > 0$, we have $\mathrm{Ext}_R^i(P, Q) = 0$;
- (3) For every connective left R -module Q , we have $\mathrm{Ext}_R^1(P, Q) = 0$;
- (4) Given an exact sequence $N' \rightarrow N \rightarrow N''$ of connective left R -modules, the induced map $\mathrm{Ext}_R^0(P, N) \rightarrow \mathrm{Ext}_R^0(P, N'')$ is surjective.

Since the t-structure on LMod_R is left complete (Corollary 8.2.5), these are further equivalent to:

- (5) For every $Q \in \mathrm{LMod}_R^\heartsuit$ and every $i > 0$, we have $\mathrm{Ext}_R^i(P, Q) = 0$.

The projective R -modules admit a simple description as retracts of free R -modules.

Definition 8.5.4 (Free module). Let R be an associative ring spectrum. A left R -module M is *free* if M is a (possibly infinite) coproduct of (unshifted) copies of R , viewed as a left module over itself. A free left R -module is *finitely generated* if it is equivalent to a finite coproduct of copies of R .

Warning 8.5.5. This is a special case of the free-module construction from Section 19.1.2. There one allows modules of the form $R \otimes X$ for an arbitrary spectrum X ; here $X = \bigoplus_{i \in I} \mathbb{S}$ is required to be a coproduct of unshifted copies of the sphere spectrum.

Proposition 8.5.6 ([Lur17, Proposition 7.2.2.7]). *Let R be a connective associative ring spectrum, and let P be a connective left R -module. The following conditions are equivalent:*

- (1) *The left R -module P is projective.*
- (2) *There exists a free R -module M such that P is a retract of M .*

Proof. Suppose first that P is projective. We can choose a map of left R -modules $p : N \rightarrow P$, where N is free, such that the induced map $\pi_0(p) : \pi_0(N) \rightarrow \pi_0(P)$ is surjective. (For instance, we can take N to be the free module generated by the set $\pi_0(P)$.) Letting N' be the fiber of p , it follows from the long exact sequence on homotopy groups that N' is connective as well. The exact sequence $N' \rightarrow N \rightarrow P$ of connective R -modules induces by Observation 6.4.10 a sequence

$$\mathrm{Ext}_R^0(P, N') \rightarrow \mathrm{Ext}_R^0(P, N) \rightarrow \mathrm{Ext}_R^0(P, P) \rightarrow \mathrm{Ext}_R^1(P, N').$$

Since P is projective, the last term vanishes by criterion (3) of Observation 8.5.3. In particular, there exists a map $s : P \rightarrow N$ such that $p \circ s = \mathrm{id}_P$, exhibiting P as a retract of the free R -module N .

For the converse, we observe that the collection of projective left R -modules is stable under retracts. It will therefore suffice to show that every free left R -module is projective. We check criterion (2)

from Observation 8.5.3. Since $\text{Ext}_R^i(\bigoplus_I R, Q) \cong \prod_I \text{Ext}_R^i(R, Q)$, we may assume $P = R$ is free on a single generator. But then

$$\text{Ext}_R^i(R, Q) = \pi_0 \text{Hom}_{\text{LMod}_R}(R, Q[i]) \simeq \pi_0 \text{Hom}_{\text{Sp}}(\mathbb{S}, Q[i]) = \pi_{-i}(Q).$$

Since Q is connective, this is zero for $i > 0$, as desired. $\square$

Remark 8.5.7. If R is an ordinary associative ring, it follows from the proposition that the projective HR -modules are precisely the Eilenberg–MacLane modules HP associated with projective R -modules in the classical sense.

8.5.2 Flat modules

We now turn to flat modules. Recall from Definition 6.5.1 the general notion of t-flatness with respect to a tensor product $C \times D \rightarrow E$ of stable ∞ -categories with t-structures. We will specialize this to the relative tensor product of modules over a ring spectrum.

Definition 8.5.8 (Tor-groups for modules). Let R be an associative ring spectrum. For a right R -module M and a left R -module N , we define the n -th Tor-group of M and N over R to be

$$\text{Tor}_n^R(M, N) := \pi_n(M \otimes_R N) \in \text{Ab}.$$

These are the Tor-groups from Definition 6.5.6 specialized to the relative tensor product $\text{RMod}_R \times \text{LMod}_R \xrightarrow{\otimes_R} \text{Sp}$.

Observation 8.5.9. Let R be a connective associative ring spectrum. By Observation 8.2.7, the relative tensor product of a connective right R -module M and a connective left R -module N is connective, and

$$\text{Tor}_0^R(M, N) = \pi_0(M \otimes_R N) \cong \pi_0(M) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N).$$

In particular, $\text{Tor}_n^R(M, N) = 0$ for $n < 0$ whenever M and N are connective.

Observation 8.5.10 (Tor long exact sequence). Specializing Observation 6.5.7, every exact sequence of right R -modules induces a long exact sequence of Tor-groups after tensoring with a fixed left R -module. The same holds for exact sequences in the left-module variable.

Definition 8.5.11 (Flat module). Let R be a connective associative ring spectrum. A connective left R -module N is said to be *flat* if the functor $(-) \otimes_R N: \text{RMod}_R \rightarrow \text{Sp}$ is t-exact.

Remark 8.5.12. By Observation 8.5.9, the functor $(-) \otimes_R N$ is automatically right t-exact when N is connective. Thus a connective left R -module N is flat if and only if this functor is left t-exact: for every coconnective right R -module M , the spectrum $M \otimes_R N$ is coconnective.

We now give several complementary tests for flatness. Conditions (2) and (3) describe the homotopy groups of a flat module and of its tensor products, while conditions (4) and (5) reduce flatness to discrete modules. The final two conditions recover the classical criterion for an ordinary ring.

Theorem 8.5.13 (Characterization of flat modules, [Lur17, Proposition 7.2.2.13, Theorem 7.2.2.15]). *Let R be a connective associative ring spectrum and let N be a connective left R -module. The following conditions are equivalent:*

(1) The left R -module N is flat.

(2) For every right R -module M and $n \in \mathbb{Z}$, the following map is an isomorphism:

$$\pi_n(M) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \xrightarrow{\sim} \pi_n(M \otimes_R N), \quad [x] \otimes [y] \mapsto [x \otimes y].$$

(3) The discrete $\pi_0(R)$ -module $\pi_0(N)$ is flat, and for each integer n the map

$$\pi_n(R) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow \pi_n(N), \quad [r] \otimes [x] \mapsto [rx]$$

is an isomorphism of abelian groups.

(4) For every discrete right R -module M , the spectrum $M \otimes_R N$ is discrete. Equivalently, $\mathrm{Tor}_i^R(M, N) = 0$ for all discrete M and all $i > 0$.

(5) Regarding $\pi_0(R)$ as a discrete ring spectrum, the left $\pi_0(R)$ -module $\pi_0(R) \otimes_R N$ is discrete and flat.

If R and N are both discrete, these conditions are further equivalent to:

(6) For every injection $M' \hookrightarrow M$ of discrete right R -modules, the induced map $M' \otimes_R^{\heartsuit} N \rightarrow M \otimes_R^{\heartsuit} N$ is injective.

(7) For every discrete right R -module M , we have $\mathrm{Tor}_1^R(M, N) = 0$.

Proof. We first compare the t-structural conditions. Apply Proposition 6.5.9 to the relative tensor product $\mathrm{RMod}_R \times \mathrm{LMod}_R \rightarrow \mathrm{Sp}$. The t-structure on RMod_R is right complete by Corollary 8.2.5, relative tensor products preserve sequential colimits, and $\mathrm{Sp}_{\leq 0}$ is closed under sequential colimits. Moreover, $M \otimes_R N$ is connective when M and N are connective by Observation 8.5.9. Thus the heart criterion of Proposition 6.5.9 says precisely that (1) $\Leftrightarrow$ (4).

(1) $\Rightarrow$ (2): Assume that $(-) \otimes_R N$ is t-exact. By Lemma 6.3.26, it commutes with taking homotopy group objects: for every right R -module M and $n \in \mathbb{Z}$, there is a natural isomorphism $\pi_n(M) \otimes_R N \xrightarrow{\sim} \pi_n(M \otimes_R N)$ in $\mathrm{Sp}^{\heartsuit} \simeq \mathrm{Ab}$. Furthermore, by Corollary 6.3.17 the t-exact functor $(-) \otimes_R N$ restricts to an exact functor $\mathrm{RMod}_R^{\heartsuit} \rightarrow \mathrm{Ab}$ on hearts. Under the equivalence $\mathrm{RMod}_R^{\heartsuit} \simeq \mathrm{RMod}_{\pi_0(R)}^{\heartsuit}$ from Lemma 8.2.9, this functor corresponds to $(-) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N)$ by Observation 8.5.9. Combining these observations gives (2).

(2) $\Rightarrow$ (3): Taking $M = R$ gives the displayed isomorphisms in (3). To prove that $\pi_0(N)$ is flat, consider a short exact sequence $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ of ordinary right $\pi_0(R)$ -modules, regarded as discrete right R -modules. Tensoring the associated exact sequence with N and applying (2) gives a short exact sequence

$$0 \rightarrow M' \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow M \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow M'' \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow 0.$$

Hence $\pi_0(N)$ is flat over $\pi_0(R)$.

(3) $\Rightarrow$ (2): We prove the formula for progressively more general right R -modules M .

Step 1: The isomorphism holds by assumption when $M = R$, and hence also when $M = F[k]$ is the shift of a free right R -module $F = \bigoplus_I R$.

Step 2: Suppose we have an exact sequence $M' \rightarrow M \rightarrow M''$ of right R -modules such that condition (2) holds for two of the three modules; we claim it also holds for the third. Since $\pi_0(N)$ is a flat $\pi_0(R)$ -module by assumption, the functor $(-) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N)$ is exact on the category of $\pi_0(R)$ -modules. Using the long exact sequence on homotopy groups for $M' \rightarrow M \rightarrow M''$, we obtain an exact sequence

$$\cdots \rightarrow \pi_n(M') \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow \pi_n(M) \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow \pi_n(M'') \otimes_{\pi_0(R)}^{\heartsuit} \pi_0(N) \rightarrow \cdots$$

Comparing this with the long exact sequence for $M' \otimes_R N \rightarrow M \otimes_R N \rightarrow M'' \otimes_R N$ via the maps from (2), the five lemma gives the claim.

Step 3: Let M be connective. The right-module version of Proposition 8.2.8 writes M as the colimit of a sequence $M(0) \rightarrow M(1) \rightarrow \cdots$ in which $M(0)$ and all the cofibers $\text{cofib}(M(i) \rightarrow M(i+1))$ are shifts of free modules. Steps 1 and 2 imply the formula for every $M(i)$. It follows for M because homotopy groups and ordinary tensor products preserve filtered colimits.

Step 4: Since the t-structure on RMod_R is right complete (Corollary 8.2.5), an arbitrary right R -module M may be written as a filtered colimit $M \simeq \text{colim}_{n \leq 0} \tau_{\geq n} M$, where each $\tau_{\geq n} M$ is a shift of a connective module. By Step 3 and the preservation of filtered colimits, (2) holds for M .

(4) $\iff$ (5): Let $A := \pi_0(R)$, regarded as a discrete ring spectrum, and let $L := A \otimes_R N$. Every discrete right R -module is equivalently a discrete right A -module, and for such an M , associativity of relative tensor products gives a natural isomorphism

$$M \otimes_A L \xrightarrow{\sim} M \otimes_R N.$$

If (4) holds, then taking $M = A$ shows that L is discrete. Moreover, the displayed isomorphism shows that $M \otimes_A L$ is discrete for every discrete right A -module M . Thus L satisfies (4) as an A -module, and hence is flat by the equivalence between (1) and (4) already proved above.

Conversely, if L is discrete and flat as an A -module, then for every discrete right R -module M , regarded as a discrete right A -module, the spectrum $M \otimes_A L$ is discrete. By the displayed isomorphism, so is $M \otimes_R N$, which is (4).

Now assume that R and N are both discrete. We show that (4) $\iff$ (6) $\iff$ (7).

(4) $\implies$ (7): This is immediate, since (4) asserts that $\text{Tor}_i^R(M, N) = 0$ for all $i > 0$.

(7) $\implies$ (6): Let $M' \hookrightarrow M$ be an injection of discrete right R -modules, with cokernel $M'' := M/M'$. By Corollary 6.3.16, the sequence $M' \rightarrow M \rightarrow M''$ is exact in RMod_R . The Tor long exact sequence (Observation 8.5.10) gives

$$\text{Tor}_1^R(M'', N) \rightarrow M' \otimes_R^{\heartsuit} N \rightarrow M \otimes_R^{\heartsuit} N \rightarrow M'' \otimes_R^{\heartsuit} N \rightarrow 0.$$

By (7), we have $\text{Tor}_1^R(M'', N) = 0$, so the map $M' \otimes_R^{\heartsuit} N \rightarrow M \otimes_R^{\heartsuit} N$ is injective.

(6) $\implies$ (4): We first show that $\text{Tor}_1^R(M, N) = 0$ for any discrete right R -module M . Choose a free R -module $F = \bigoplus_I R$ with a surjection $F \twoheadrightarrow M$ (for example, take I to be the underlying set of M), and let $K := \ker(F \twoheadrightarrow M)$. The Tor long exact sequence gives

$$\text{Tor}_1^R(F, N) \rightarrow \text{Tor}_1^R(M, N) \rightarrow K \otimes_R^{\heartsuit} N \rightarrow F \otimes_R^{\heartsuit} N.$$

Since $F \otimes_R N \cong \bigoplus_I N$ is discrete, we have $\text{Tor}_1^R(F, N) = 0$. By (6), the inclusion $K \hookrightarrow F$ induces an injection $K \otimes_R^{\heartsuit} N \hookrightarrow F \otimes_R^{\heartsuit} N$. Exactness then forces $\text{Tor}_1^R(M, N) = 0$.

For higher Tor-groups, we use dimension shifting. The short exact sequence $0 \rightarrow K \hookrightarrow F \rightarrow M \rightarrow 0$ gives, for $i \geq 1$,

$$\mathrm{Tor}_{i+1}^R(F, N) \rightarrow \mathrm{Tor}_{i+1}^R(M, N) \rightarrow \mathrm{Tor}_i^R(K, N) \rightarrow \mathrm{Tor}_i^R(F, N).$$

Since F is free, $\mathrm{Tor}_j^R(F, N) = 0$ for $j > 0$. Thus $\mathrm{Tor}_{i+1}^R(M, N) \cong \mathrm{Tor}_i^R(K, N)$ for $i \geq 1$. By induction, $\mathrm{Tor}_i^R(M, N) = 0$ for all $i > 0$ and all discrete M . $\square$

Observation 8.5.14. Let R be a connective associative ring spectrum. Condition (3) of Theorem 8.5.13 immediately gives the following facts:

- (a) The subcategory of LMod_R spanned by the flat R -modules is closed under coproducts, retracts, and filtered colimits.
- (b) Every free module is flat. Combining this with (a) and Proposition 8.5.6, it follows that every projective module is flat.
- (c) If A is an ordinary associative ring, then a connective left HA -module N is flat if and only if N is discrete and $\pi_0 N$ is a flat A -module in the classical sense, i.e., satisfies condition (6) of Theorem 8.5.13.

Lemma 8.5.15. Let $R \in \mathrm{CAlg}(\mathrm{Sp}_{\geq 0})$ be a connective commutative ring spectrum, and let M and N be flat R -modules. Then $M \otimes_R N$ is again flat.

Proof. By Corollary 8.2.5, connectivity and coconnectivity of an R -module are detected on its underlying spectrum. A connective R -module P is therefore flat if and only if the functor $(-) \otimes_R P: \mathrm{Mod}_R \rightarrow \mathrm{Mod}_R$ is t-exact, as opposed to merely its composite with the forgetful functor to Sp . Now associativity of the relative tensor product provides a natural isomorphism

$$(-) \otimes_R (M \otimes_R N) \simeq ((-) \otimes_R M) \otimes_R N$$

of functors $\mathrm{Mod}_R \rightarrow \mathrm{Mod}_R$. Since M and N are flat, the right-hand side is a composite of two t-exact functors, hence t-exact. Finally, $M \otimes_R N$ is connective by Observation 8.5.9, so it is flat. $\square$

Observation 8.5.16. Every free R -module is flat (Observation 8.5.14), so in particular the monoidal unit R is flat. Combined with Lemma 8.5.15, this shows that the full subcategory $\mathrm{Mod}_R^b \subseteq \mathrm{Mod}_{R, \geq 0}$ spanned by the flat R -modules contains the unit and is closed under the tensor product. It therefore inherits a symmetric monoidal structure by the Part II result Lemma 14.5.2.

For connective ring spectra, flat modules admit a characterization analogous to Lazard's theorem in ordinary algebra. It expresses them as filtered colimits of the simplest flat modules, the finitely generated free ones.

Theorem 8.5.17 (Lazard's theorem for spectra, [Lur17, Theorem 7.2.2.15]). Let R be a connective associative ring spectrum and let N be a connective left R -module. The following conditions are equivalent:

- (1) The left R -module N can be obtained as a filtered colimit of finitely generated free left R -modules.

- (2) *The left R -module N can be obtained as a filtered colimit of finitely generated projective left R -modules.*
- (3) *The left R -module N is flat.*

Proof sketch. The implication (1) $\Rightarrow$ (2) is immediate because every free module is projective. The implication (2) $\Rightarrow$ (3) follows from Observation 8.5.14: projective modules are flat, and flat modules are closed under filtered colimits. The reverse implication is the substantive content of [Lur17, Theorem 7.2.2.15], which we use as a black box. Its proof shows that the ∞ -category of maps $F \rightarrow N$, with F finitely generated and free, is filtered and that the canonical map from the colimit of this diagram to N is an isomorphism. The filteredness is ultimately reduced to the classical Lazard theorem over $\pi_0(R)$. $\square$

8.5.3 Tor-amplitude

We conclude with two boundedness properties of perfect modules. The first is immediate from their construction out of the unit.

Lemma 8.5.18. *Let R be a connective associative ring spectrum and let M be a perfect R -module in the sense of Definition 8.1.7. Then M is bounded below: there exists an integer n such that $\pi_i(M) = 0$ for all $i < n$.*

Proof. The ring spectrum R is bounded below, and the bounded below R -modules form a thick subcategory of $\mathbf{LMod}_R$: they are closed under shifts, cofibers, and retracts. The claim follows from the definition of a perfect module. $\square$

Tor-amplitude gives a uniform bound after tensoring with discrete modules. By Theorem 8.5.13, flat modules are precisely the connective modules of Tor-amplitude contained in $[0, 0]$.

Definition 8.5.19 (Tor-amplitude). Let R be a connective associative ring spectrum and let M be a left R -module. We say that M has *Tor-amplitude contained in $[a, b]$* for integers $a \leq b$ if for every discrete right R -module N , the homotopy groups of $N \otimes_R M$ are concentrated in degrees $[a, b]$, that is, $\mathrm{Tor}_i^R(N, M) = 0$ whenever $i < a$ or $i > b$. If such a and b exist, M is said to have *finite Tor-amplitude*.

The last three statements below make this bound effective for perfect modules. A perfect module concentrated in a single Tor degree is a shifted finitely generated projective module; in general, one can successively remove the bottom Tor degree.

Proposition 8.5.20 ([AG14, Proposition 2.13]). *Let R be a connective associative ring spectrum and let M and N be left R -module spectra.*

- (1) *If M has Tor-amplitude contained in $[a, b]$, then the shift $M[k]$ has Tor-amplitude contained in $[a + k, b + k]$ for all $k \in \mathbb{Z}$.*
- (2) *If M is perfect, then M has finite Tor-amplitude.*

- (3) *If R happens to be a commutative ring spectrum, $S \in \mathrm{CAlg}(\mathrm{Sp})_{R/}$ is connective, and M has Tor-amplitude contained in $[a, b]$, then the left S -module $S \otimes_R M$ has Tor-amplitude contained in $[a, b]$.*
- (4) *If M and N have Tor-amplitude contained in $[a, b]$, then the fiber and cofiber of a map $M \rightarrow N$ have Tor-amplitude contained in $[a - 1, b]$ and $[a, b + 1]$.*
- (5) *If M is perfect with Tor-amplitude contained in $[0, b]$, then M is connective and $\pi_0 M \cong \pi_0(\pi_0(R) \otimes_R M)$, where $\pi_0(R)$ is regarded as a discrete ring spectrum.*
- (6) *If M is perfect with Tor-amplitude contained in $[a, a]$, then M is equivalent to $P[a]$ for a finitely generated projective left R -module P .*
- (7) *If M is perfect with Tor-amplitude contained in $[a, b]$, then there exists an exact sequence $P[a] \rightarrow M \rightarrow Q$ with P finitely generated projective and Q perfect with Tor-amplitude contained in $[a + 1, b]$.*

Proof sketch. Part (1) follows directly from the definition. For (2), observe that R has Tor-amplitude contained in $[0, 0]$ and that the left R -modules of finite Tor-amplitude form a thick subcategory of LMod_R : they are closed under shifts and retracts, while closure under cofibers follows from the long exact sequence of Tor-groups (Observation 8.5.10). For (3), if T is a discrete right S -module, then

$$T \otimes_S (S \otimes_R M) \simeq T \otimes_R M.$$

Part (4) follows from the same long exact sequence of Tor-groups. Parts (5)–(7) require the Tor spectral sequence together with the lifting theory of projective modules, so we take them directly from Antieau and Gepner [AG14, Proposition 2.13]. $\square$

Exercises

9 Complex K-theory

One of the most classical cohomology theories is *complex K-theory*, established as part of algebraic topology by Atiyah and Hirzebruch [AH59], building on work of Grothendieck. The starting point is the set of isomorphism classes of complex vector bundles over a space X . This set admits an operation, given by direct sum. The complex K-theory group $K^0(X)$ of X is defined as the group completion of this commutative monoid, formally adding inverses to all vector bundles. Its central property is Bott periodicity: the theory repeats with period 2.

Complex K-theory is represented by a commutative ring spectrum KU . The construction of this spectrum mirrors the definition of $K^0(X)$ via group completion. First, one can refine the set of isomorphism classes of vector bundles to an anima $\mathbf{Vect}(X)^\simeq$, which takes into account the topology on the sets of isomorphisms between vector bundles. The direct sum operation turns this into a coherently commutative monoid in $\mathbf{An}$. Applying the homotopy-theoretic group completion operation of Chapter 5, we obtain a commutative group in $\mathbf{An}$, which under the recognition principle corresponds to a connective spectrum ku . The 2-periodic K-theory spectrum KU is then obtained by inverting a certain Bott class $\beta \in \pi_2(ku)$.

We recall complex vector bundles and their homotopy invariance in Section 9.1, and identify the infinite Grassmannian as their classifying anima in Section 9.2. Section 9.3 defines the groups $K^0(X)$, proves that they are represented by $\mathbb{Z} \times BU$, and states Bott periodicity. Section 9.4 then carries out the spectrum-level construction of ku and KU sketched above. The multiplicative structure comes from the tensor product of vector bundles; making it coherent requires the higher algebra of Part II, and is carried out in Section 19.7. We close by rationalizing KU and deducing that the Chern character is a rational isomorphism.

9.1 Vector bundles

In this section, we introduce the classical theory of complex vector bundles and isolate the input needed later: morphisms, pullbacks, direct sums, tensor products, and homotopy invariance.

Definition 9.1.1. Let X be a topological space. A *finite-rank complex vector bundle* over X is a continuous map $p: E \rightarrow X$, together with a complex vector space structure on every fiber $E_x := p^{-1}(x)$, satisfying the following local triviality condition: every point $x \in X$ admits an open neighborhood U , an integer $k \geq 0$, and a homeomorphism

$$\varphi: U \times \mathbb{C}^k \xrightarrow{\cong} p^{-1}(U)$$

over U whose restriction $\{y\} \times \mathbb{C}^k \rightarrow E_y$ is complex-linear for every $y \in U$. We will usually simply call this a *complex vector bundle*.

The rank function

$$\mathrm{rk}_E: X \longrightarrow \mathbb{N}, \quad x \longmapsto \dim_{\mathbb{C}}(E_x),$$

is locally constant. We say that E has *rank* k if this function is constant with value k . More generally, each subspace $X_k := \mathrm{rk}_E^{-1}(k)$ is clopen in X , and the restriction of E to X_k has rank k .

Given another complex vector bundle $q: E' \rightarrow X$, a *morphism of vector bundles* $E \rightarrow E'$ is a continuous map $f: E \rightarrow E'$ over X whose restriction $E_x \rightarrow E'_x$ is complex-linear for every $x \in X$. We write $\pi_0 \mathrm{Vect}(X)$ for the set of isomorphism classes of finite-rank complex vector bundles over X , and $\pi_0 \mathrm{Vect}_k(X)$ for the subset represented by bundles of rank k .

Let X be a topological space, and let $p_1: E_1 \rightarrow X$ and $p_2: E_2 \rightarrow X$ be two complex vector bundles over X . Their *direct sum* is defined as the fiber product

$$E_1 \oplus E_2 := E_1 \times_X E_2 = \{(e_1, e_2) \mid p_1(e_1) = p_2(e_2)\},$$

equipped on each fiber with the usual direct-sum vector space structure. We leave it to the reader to check that this is again a vector bundle; see Exercise 9.1. The trivial rank-zero bundle $\underline{0} := X \times \mathbb{C}^0$ serves as a unit for direct sum up to the evident isomorphisms, and the usual associativity and symmetry isomorphisms show that direct sum induces an abelian monoid structure on $\pi_0 \mathrm{Vect}(X)$.

We will also use the *tensor product* of two vector bundles E_1 and E_2 . This is the vector bundle $E_1 \otimes E_2 \rightarrow X$ whose fiber over $x \in X$ is $E_{1,x} \otimes_{\mathbb{C}} E_{2,x}$; local trivializations of E_1 and E_2 induce local trivializations of $E_1 \otimes E_2$. Pullback along maps of spaces is compatible with direct sums and tensor products, and tensor product distributes over direct sums up to isomorphism. Thus tensor product induces a multiplication on $\pi_0 \mathrm{Vect}(X)$, compatible with its direct-sum monoid structure.

9.1.1 Homotopy invariance of vector bundles

We will use the following classical homotopy invariance theorem for vector bundles over paracompact spaces.

Definition 9.1.2. A topological space X is called *paracompact* if for every open cover $\{U_i\}_{i \in I}$ of X there exists a *partition of unity subordinate to this cover*, i.e. a collection of continuous maps $\rho_\alpha: X \rightarrow [0, 1]$ satisfying the following three conditions:

- (1) The family $(\rho_\alpha)_\alpha$ is locally finite: every point of X has a neighborhood on which all but finitely many ρ_α vanish;
- (2) The resulting sum satisfies $\sum_\alpha \rho_\alpha(x) = 1$ for all x ;
- (3) For each α , the support $\mathrm{supp}(\rho_\alpha) := \overline{\rho_\alpha^{-1}((0, 1])}$ is contained in one of the open sets U_i .

For Hausdorff spaces, this is equivalent to the usual definition in terms of locally finite refinements of open covers.

Example 9.1.3. The following classes of topological spaces are paracompact:

- Every compact Hausdorff space is paracompact [Hat03, Proposition 1.18];

- Every direct limit, with the colimit topology, of an increasing sequence $X_1 \subseteq X_2 \subseteq \dots$ of compact Hausdorff spaces is paracompact [Hat03, Proposition 1.19];
- Every CW-complex is paracompact [Hat03, Proposition 1.20].

Theorem 9.1.4 (Homotopy invariance of vector bundles, Hatcher [Hat03, Theorem 1.6 and Corollary 1.8]). *Let $p: E \rightarrow X$ be a complex vector bundle and let $f_0, f_1: Y \rightarrow X$ be homotopic continuous maps. If Y is paracompact Hausdorff, then the pullback bundles f_0^*E and f_1^*E are isomorphic.*

In particular, every homotopy equivalence $f: X \rightarrow Y$ between paracompact Hausdorff spaces induces a bijection

$$f^*: \pi_0 \text{Vect}(Y) \xrightarrow{\cong} \pi_0 \text{Vect}(X).$$

Remark 9.1.5. The proof uses the usual partition-of-unity argument: a bundle on $Y \times [0, 1]$ is locally constant in the interval direction after subdividing $[0, 1]$, and paracompactness lets one patch the resulting local isomorphisms. We will use only the conclusion above.

9.1.2 Clutching functions

Let $X = U_+ \cup U_-$ be a topological space written as the union of two open subspaces, and let $E \rightarrow X$ be a rank n complex vector bundle which is trivial over both U_+ and U_- . Choosing trivializations

$$E|_{U_+} \cong U_+ \times \mathbb{C}^n \quad \text{and} \quad E|_{U_-} \cong U_- \times \mathbb{C}^n$$

identifies the gluing of these two trivial bundles along $U_+ \cap U_-$ with a continuous map

$$g: U_+ \cap U_- \rightarrow \text{GL}_n(\mathbb{C}),$$

called the *clutching function* of E with respect to the chosen trivializations. Conversely, any such map g defines a vector bundle by the quotient

$$E_g := (U_+ \times \mathbb{C}^n) \sqcup (U_- \times \mathbb{C}^n) / (x, v)_+ \sim (x, g(x)v)_-$$

for $x \in U_+ \cap U_-$. Thus clutching functions give a concrete way of building vector bundles by gluing trivial bundles.

For spheres this construction gives an elementary classification. Let $m \geq 1$, and let U_+ and U_- be slightly enlarged upper and lower hemispheres in S^m . Then U_+ and U_- are contractible paracompact Hausdorff spaces, and their intersection deformation retracts onto the equator. Informally, we are using the decomposition

$$S^m = D_+^m \cup_{S^{m-1}} D_-^m.$$

By homotopy invariance, every vector bundle over either U_+ or U_- is trivial, so a rank n complex vector bundle over S^m is obtained from a clutching function

$$g: S^{m-1} \rightarrow \text{GL}_n(\mathbb{C}).$$

The clutching construction induces a bijection

$$\pi_0 \text{Vect}_n(S^m) \cong [S^{m-1}, \text{GL}_n(\mathbb{C})]$$

with the set of free homotopy classes [Hat03, Proposition 1.11]. Since $\mathrm{GL}_n(\mathbb{C})$ is a connected topological group, based and free homotopy classes from a sphere agree, and hence this set is $\pi_{m-1}(\mathrm{GL}_n(\mathbb{C}))$. The required connectedness is verified in Exercise 9.1.6; for $m = 1$, the last statement reads $\pi_0 \mathrm{Vect}_n(S^1) \cong \pi_0 \mathrm{GL}_n(\mathbb{C})$.

Exercise 9.1.6 (Vector bundles over the circle). Show that the topological group $\mathrm{GL}_n(\mathbb{C})$ is connected. Deduce that every rank n complex vector bundle over S^1 is trivial, so that

$$\pi_0 \mathrm{Vect}(S^1) \cong \mathbb{N}.$$

Exercise 9.1.7 (Vector bundles over the two-sphere). Compute that $\pi_1(\mathrm{GL}_1(\mathbb{C})) \cong \mathbb{Z}$, with $m \in \mathbb{Z}$ represented by the clutching function $g_m: S^1 \rightarrow \mathbb{C}^\times, z \mapsto z^m$, and construct an explicit representative L_m for the corresponding line bundle over S^2 . Next, show that the inclusion $U(n) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ is a homotopy equivalence. Using the fiber sequence

$$U(n-1) \hookrightarrow U(n) \rightarrow S^{2n-1},$$

deduce that $\pi_1(\mathrm{GL}_n(\mathbb{C})) \cong \mathbb{Z}$ for all $n \geq 1$. Conclude that $\pi_0 \mathrm{Vect}_k(S^2) \cong \mathbb{Z}$ for all $k \geq 1$, with every rank k complex vector bundle over S^2 isomorphic to

$$L_m \oplus \underline{\mathbb{C}}^{k-1}$$

for a unique $m \in \mathbb{Z}$.

9.2 Classifying vector bundles

For paracompact Hausdorff spaces, vector bundles are classified by maps into Grassmannians: there exists a topological space $\mathrm{Gr}_k(\mathbb{C}^\infty)$ such that an isomorphism class of rank k complex vector bundle over X determines and is determined by a homotopy class of continuous maps from X into $\mathrm{Gr}_k(\mathbb{C}^\infty)$. The goal of this section is to explain this result and relate the space $\mathrm{Gr}_k(\mathbb{C}^\infty)$ to the classifying anima $\mathbf{BU}(k)$ of the topological group $U(k)$.

9.2.1 The infinite Grassmannian

Definition 9.2.1 (Infinite grassmannian). Let $\mathbb{C}^\infty$ denote the colimit in $\mathbf{Top}$ of the complex vector spaces $\mathbb{C}^n$ along the standard inclusions $\mathbb{C}^n \hookrightarrow \mathbb{C}^{n+1}$. For $n \geq k$, let $\mathrm{Gr}_k(\mathbb{C}^n)$ be the Grassmannian of k -dimensional linear subspaces of $\mathbb{C}^n$, with its usual topology. We define the *infinite Grassmannian of k -planes* as

$$\mathrm{Gr}_k(\mathbb{C}^\infty) := \bigcup_{n \geq k} \mathrm{Gr}_k(\mathbb{C}^n),$$

equipped with the *colimit topology*: a subset $U \subseteq \mathrm{Gr}_k(\mathbb{C}^\infty)$ is open if and only if $U \cap \mathrm{Gr}_k(\mathbb{C}^n)$ is open for every n .

Similarly, let $V_k(\mathbb{C}^n)$ be the Stiefel manifold of orthonormal k -frames in $\mathbb{C}^n$. The *infinite Stiefel manifold*

$$V_k(\mathbb{C}^\infty) := \bigcup_{n \geq k} V_k(\mathbb{C}^n)$$

is likewise given the colimit topology. Its points may equivalently be regarded as linear isometric embeddings $\mathbb{C}^k \hookrightarrow \mathbb{C}^\infty$. Sending a frame to the subspace it spans defines a surjective map

$$\pi: V_k(\mathbb{C}^\infty) \rightarrow \mathrm{Gr}_k(\mathbb{C}^\infty), \quad (\varphi: \mathbb{C}^k \hookrightarrow \mathbb{C}^\infty) \mapsto \varphi(\mathbb{C}^k),$$

whose restriction at every finite stage is the usual quotient map $V_k(\mathbb{C}^n) \rightarrow \mathrm{Gr}_k(\mathbb{C}^n)$.

The standard Schubert cell structures on the finite Grassmannians are compatible with the inclusions above. Consequently $\mathrm{Gr}_k(\mathbb{C}^\infty)$ is a cell complex whose topology is precisely the colimit topology just defined; see Hatcher [Hat03, Proposition 1.17 and p. 34].

Definition 9.2.2. For $n \geq k$, let

$$E_{k,n} := \{(V, x) \in \mathrm{Gr}_k(\mathbb{C}^n) \times \mathbb{C}^n \mid x \in V\}$$

be the *canonical rank k bundle* over $\mathrm{Gr}_k(\mathbb{C}^n)$. We define

$$E_{\mathrm{univ}} := \bigcup_{n \geq k} E_{k,n}$$

with the colimit topology, and let $p_{\mathrm{univ}}: E_{\mathrm{univ}} \rightarrow \mathrm{Gr}_k(\mathbb{C}^\infty)$ be the projection $(V, x) \mapsto V$.

Exercise 9.2.3. Show that the map p_{univ} is a complex vector bundle of rank k . *Hint:* compare with the proof of Lemma 9.2.12 below.

Let X be a paracompact Hausdorff space. Given a continuous map $f: X \rightarrow \mathrm{Gr}_k(\mathbb{C}^\infty)$, we may pull back the bundle E_{univ} along f , providing a vector bundle

$$f^*(E_{\mathrm{univ}}) := \{(x, \varphi) \in X \times E_{\mathrm{univ}} \mid f(x) = p_{\mathrm{univ}}(\varphi)\}.$$

By Theorem 9.1.4, the isomorphism class of this vector bundle only depends on the homotopy class of f , providing a map

$$[X, \mathrm{Gr}_k(\mathbb{C}^\infty)] \rightarrow \pi_0 \mathrm{Vect}_k(X), \quad [f] \mapsto f^*(E_{\mathrm{univ}}). \quad (9.1)$$

The following classical theorem expresses that p_{univ} is the *universal* rank k complex vector bundle:

Theorem 9.2.4 (Hatcher [Hat03, Theorem 1.16 and p. 31]). *For a paracompact Hausdorff space X , the map (9.1) is a bijection.*

Because of this theorem, $\mathrm{Gr}_k(\mathbb{C}^\infty)$ is known as the *classifying space for rank k complex vector bundles*.

Proof sketch of Theorem 9.2.4. We describe the main ingredients of Hatcher's argument. The key observation is that, for a rank k vector bundle $p: E \rightarrow X$, the following two types of data are equivalent:

- (1) A pair (f, φ) consisting of a continuous map $f: X \rightarrow \mathrm{Gr}_k(\mathbb{C}^\infty)$ and an isomorphism $\varphi: E \cong f^*(E_{\mathrm{univ}})$;
- (2) A continuous map $g: E \rightarrow \mathbb{C}^\infty$ whose restriction to each fiber is a $\mathbb{C}$ -linear injection.

Indeed, an isomorphism $E \cong f^*(E_{\text{univ}})$ gives such a map by projecting $f^*(E_{\text{univ}}) \subseteq X \times \mathbb{C}^\infty$ to $\mathbb{C}^\infty$. Conversely, a fiberwise linear injection g determines $f(x) := g(E_x) \subseteq \mathbb{C}^\infty$, and the evident map $E \rightarrow E_{\text{univ}}$ identifies E with $f^*(E_{\text{univ}})$.

Surjectivity is therefore reduced to constructing a map $g: E \rightarrow \mathbb{C}^\infty$ which is linear and injective on each fiber. Starting from a trivializing cover, one uses the countable-cover lemma for paracompact spaces to obtain a countable trivializing cover $(U_i)_{i \geq 1}$ together with a locally finite subordinate partition of unity $(\rho_i)_{i \geq 1}$. Let $g_i: E|_{U_i} \rightarrow \mathbb{C}^k$ be the coordinate map of a trivialization. The maps

$$v \mapsto \rho_i(p(v))g_i(v),$$

extended by zero outside U_i , assemble into a map $E \rightarrow (\mathbb{C}^k)^\infty \cong \mathbb{C}^\infty$. Local finiteness ensures that this map is continuous and takes values in the algebraic direct sum, while the partition-of-unity condition makes it injective on every fiber. This is the only point at which paracompactness is used in an essential way.

For injectivity, suppose two classifying maps $f_0, f_1: X \rightarrow \text{Gr}_k(\mathbb{C}^\infty)$ pull back E_{univ} to isomorphic bundles. The corresponding fiberwise injections $g_0, g_1: E \rightarrow \mathbb{C}^\infty$ may first be homotoped so that their images land in the odd and even coordinates, respectively. The straight-line homotopy $(1-t)g_0 + tg_1$ then remains fiberwise injective for all t , and hence determines a homotopy from f_0 to f_1 .

For the countable-cover lemma, see Hatcher [Hat03, Lemma 1.21]; the remaining continuity details are part of the proof of Hatcher [Hat03, Theorem 1.16]. $\square$

9.2.2 Classifying animae revisited

Recall that in Construction 5.1.14 we defined the *classifying anima* $\mathbf{B}G$ of a group object $G \in \text{Grp}(\text{An})$. To identify the underlying anima of the infinite Grassmannian, we develop a general recognition criterion for classifying animae. The same criterion will be reused for real vector bundles in Chapter 10.

Recall from Corollary 2.3.20 that the functor $\Pi_\infty: \text{Top} \rightarrow \text{An}$ preserves finite products, hence induces a functor

$$\Pi_\infty: \text{Grp}(\text{Top}) \rightarrow \text{Grp}(\text{An})$$

from the category of topological groups to the ∞ -category of group objects in animae.

Definition 9.2.5. For a topological group G , we define its *classifying anima* $\mathbf{B}G$ as

$$\mathbf{B}G := \mathbf{B}\Pi_\infty(G) \in \text{An}.$$

We will now discuss a concrete way of recognizing topological spaces whose underlying anima is isomorphic to $\mathbf{B}G$.

Definition 9.2.6 (Principal bundle). Let G be a topological group and let B be a topological space. The *trivial G -principal bundle* is the projection map $B \times G \rightarrow B$, where we equip $B \times G$ with the canonical right G -action given by $(b, g)g' := (b, gg')$.

A *G -principal bundle* is a continuous map $\pi: P \rightarrow B$ such that P is equipped with a continuous right G -action such that:

- (1) The right G -action on P preserves the fibers of π : we have $\pi(pg) = \pi(p)$ for all $p \in P$ and $g \in G$;
- (2) The map π is a *locally trivial bundle*: around every point $b \in B$ there is an open neighborhood $U \subseteq B$ such that the restriction $\pi^{-1}(U) \rightarrow U$ of π is a trivial G -principal bundle, i.e. there exists a G -equivariant homeomorphism $\varphi: U \times G \xrightarrow{\cong} \pi^{-1}(U)$ over U :

$$\begin{array}{ccccc} U \times G & \xrightarrow[\cong]{\varphi} & \pi^{-1}(U) & \hookrightarrow & P \\ & \searrow \text{pr}_1 & \downarrow \pi|_U & \lrcorner & \downarrow \pi \\ & & U & \hookrightarrow & B. \end{array}$$

We refer to P as the *total space*, to G as the *structure group*, and to B as the *base space*.

Exercise 9.2.7. Show that every principal $\text{GL}_n(\mathbb{C})$ -bundle $P \rightarrow B$ defines a complex vector bundle of rank n over B by taking

$$E := P \times_{\text{GL}_n(\mathbb{C})} \mathbb{C}^n = (P \times \mathbb{C}^n) / (p, Av) \sim (pA, v),$$

where the equivalence relation identifies the pair (p, Av) with (pA, v) for all $p \in P$, $v \in \mathbb{C}^n$ and $A \in \text{GL}_n(\mathbb{C})$. Show that this defines a bijection between the set of isomorphism classes of principal $\text{GL}_n(\mathbb{C})$ -bundles and the set of isomorphism classes of rank n vector bundles.

Remark 9.2.8. Since $\pi: P \rightarrow B$ is locally trivial, hence locally a Serre fibration, it follows from Theorem 2.3.13 that every principal bundle is a Serre fibration. In particular, the functor Π_∞ preserves pullbacks along principal bundles by Proposition 2.3.19.

Lemma 9.2.9. Let $\pi: P \rightarrow B$ be a principal G -bundle. Then for every $n \geq 0$ the continuous map

$$\Phi_n: P \times G^n \rightarrow P \times_B P \times_B \cdots \times_B P, \quad (p, g_1, \dots, g_n) \mapsto (p, pg_1, \dots, pg_1 \cdots g_n)$$

is a homeomorphism.

Proof. By induction we may immediately reduce to the case $n = 1$, i.e. we must show that the map $\Phi_1: P \times G \rightarrow P \times_B P$, $(p, g) \mapsto (p, pg)$ is a homeomorphism. Observe that this map fits in a commutative diagram of the form

$$\begin{array}{ccc} P \times G & \xrightarrow{\Phi_1} & P \times_B P \\ \text{pr}_1 \searrow & & \swarrow \text{pr}_1 \\ & P, & \end{array}$$

and so it suffices to find an open cover of P such that the given map restricts to a homeomorphism on the preimages. Since π is locally trivial, we may thus assume that it is of the form $\text{pr}: B \times G \rightarrow B$. But in this case, the map Φ_1 takes the form $B \times G \times G \rightarrow B \times G \times G$, $(b, g, h) \mapsto (b, g, gh)$, which is indeed a homeomorphism with inverse $(b, g, h') \mapsto (b, g, g^{-1}h')$. $\square$

Observe that both sides of Lemma 9.2.9 define simplicial objects in $\mathbf{Top}$. The simplicial structure on the right-hand side is that of the Čech nerve $\check{C}(\pi)$, see the proof of Proposition 5.1.15. The one on the left is the bar construction associated to the right action of G on P , with face maps

$$\begin{aligned} d_0(p, g_1, \dots, g_n) &= (pg_1, g_2, \dots, g_n), \\ d_i(p, g_1, \dots, g_n) &= (p, g_1, \dots, g_i g_{i+1}, \dots, g_n) \quad (0 < i < n), \\ d_n(p, g_1, \dots, g_n) &= (p, g_1, \dots, g_{n-1}), \end{aligned}$$

and degeneracy maps

$$s_i(p, g_1, \dots, g_n) = (p, g_1, \dots, g_i, e, g_{i+1}, \dots, g_n) \quad (0 \leq i \leq n),$$

where the initial string $g_1, \dots, g_i$ is empty when $i = 0$.

Lemma 9.2.10. *The homeomorphisms Φ_n from Lemma 9.2.9 define an isomorphism of simplicial objects in $\mathbf{Top}$.*

Proof. It suffices to check the face and degeneracy maps. For the first face map, we have

$$\begin{aligned} \Phi_{n-1}(d_0^*(p, g_1, \dots, g_n)) &= \Phi_{n-1}(pg_1, g_2, \dots, g_n) \\ &= (pg_1, pg_1g_2, \dots, pg_1g_2 \dots g_n) = d_0^*\Phi_n(p, g_1, \dots, g_n). \end{aligned}$$

For a degeneracy map, inserting the identity element among the g_i inserts a repeated partial product among $p, pg_1, \dots, pg_1 \dots g_n$, as required. The remaining face maps either multiply two adjacent g_i or omit the last one, and the same direct calculation proves compatibility. $\square$

We now get to the promised result about recognizing topological spaces whose underlying anima is of the form $\mathbf{BG}$:

Proposition 9.2.11. *Let $\pi: P \rightarrow B$ be a principal G -bundle such that P is contractible.*

- (1) *The Čech nerve of $\Pi_\infty(\pi): \Pi_\infty(P) \rightarrow \Pi_\infty(B)$ is isomorphic to $\Pi_\infty(G) \in \mathbf{Grp}(\mathbf{An})$;*
- (2) *In particular, passing to colimits provides an isomorphism of anima*

$$\Pi_\infty(B) \xrightarrow{\sim} \mathbf{BG}.$$

Proof. A principal bundle is a locally trivial fiber bundle and hence a Serre fibration by Theorem 2.3.13. Since $\Pi_\infty: \mathbf{Top} \rightarrow \mathbf{An}$ preserves pullbacks along Serre fibrations by Proposition 2.3.19, there are isomorphisms

$$\Pi_\infty(P \times_B \dots \times_B P) \xrightarrow{\sim} \Pi_\infty(P) \times_{\Pi_\infty(B)} \dots \times_{\Pi_\infty(B)} \Pi_\infty(P)$$

for all n , which assemble into an isomorphism of simplicial anima $\Pi_\infty(\check{C}(\pi)) \xrightarrow{\sim} \check{C}(\Pi_\infty(\pi))$. By Lemma 9.2.10, the simplicial anima $\Pi_\infty(\check{C}(\pi))$ is isomorphic to $[n] \mapsto \Pi_\infty(P \times G^n)$. By contractibility of P , the projection maps $\Pi_\infty(P \times G^n) \cong \Pi_\infty(P) \times \Pi_\infty(G^n) \rightarrow \Pi_\infty(G^n)$ are isomorphisms, which are compatible with the simplicial structure maps (see Exercise 5.1). This proves part (1). For part (2), observe that $\Pi_\infty(\pi)$ is surjective on path components, since π is surjective. It is therefore an effective epimorphism of anima, so $\Pi_\infty(B)$ is the colimit of its Čech nerve. On the other hand, the colimit of the simplicial anima underlying the group object $\Pi_\infty(G)$ is its classifying anima $\mathbf{BG}$. Part (1) therefore gives the claimed isomorphism. $\square$

9.2.3 The infinite Grassmannian as a classifying anima

We now return to the infinite Grassmannian: we wish to apply Proposition 9.2.11 to the map $\pi: V_k(\mathbb{C}^\infty) \rightarrow \text{Gr}_k(\mathbb{C}^\infty)$ to conclude that $\text{Gr}_k(\mathbb{C}^\infty)$ is a geometric model for the classifying anima of the unitary group $U(k)$, the group of complex isometries $\mathbb{C}^k \xrightarrow{\cong} \mathbb{C}^k$.

Lemma 9.2.12. *The projection map $\pi: V_k(\mathbb{C}^\infty) \rightarrow \text{Gr}_k(\mathbb{C}^\infty)$ is a principal $U(k)$ -bundle. Here $U(k)$ acts on $V_k(\mathbb{C}^\infty)$ by precomposition: given $A \in U(k)$ and $\varphi: \mathbb{C}^k \hookrightarrow \mathbb{C}^\infty$, we define φA as the composite $\mathbb{C}^k \xrightarrow{A} \mathbb{C}^k \xrightarrow{\varphi} \mathbb{C}^\infty$.*

Proof. Let $W \in \text{Gr}_k(\mathbb{C}^\infty)$, and define $U \subseteq \text{Gr}_k(\mathbb{C}^\infty)$ to be the subspace of those k -planes V for which the orthogonal projection $W \rightarrow V$ is an isomorphism. This is open: its intersection with every finite-dimensional stage $\text{Gr}_k(\mathbb{C}^n)$ is open, and $\text{Gr}_k(\mathbb{C}^\infty)$ has the colimit topology. We claim that there exists a $U(k)$ -equivariant homeomorphism

$$\varphi: U \times U(k) \xrightarrow{\cong} \pi^{-1}(U)$$

over U , which will suffice to finish the proof. To construct φ , fix a k -frame $(w_1, \dots, w_k)$ for W . Given any k -plane $V \in U$, the projection $W \rightarrow V$ is an isomorphism, hence sends the k -frame $(w_1, \dots, w_k)$ to some basis for V . Using Gram-Schmidt orthonormalization (which is a continuous operation), we may turn this into a k -frame $(v_1, \dots, v_k)$ for V . All in all, we may then define

$$\varphi(V, A) := (v_1, \dots, v_k) \circ A.$$

Bijectivity of φ holds because two orthonormal frames of a k -plane differ by a unique unitary $(k \times k)$ -matrix. We leave it to the reader to check that this is indeed a well-defined homeomorphism over U . $\square$

Remark 9.2.13. Under the bijection of Exercise 9.2.7, the principal bundle $V_k(\mathbb{C}^\infty) \rightarrow \text{Gr}_k(\mathbb{C}^\infty)$ is in fact the principal bundle associated to the universal rank k vector bundle E_{univ} from Definition 9.2.2.

Lemma 9.2.14. *The topological space $V_k(\mathbb{C}^\infty)$ is contractible.*

Proof. Consider the space $\widetilde{V}_k(\mathbb{C}^\infty)$ of all linear injections $\mathbb{C}^k \hookrightarrow \mathbb{C}^\infty$ which are not necessarily isometries. Equivalently, $\widetilde{V}_k(\mathbb{C}^\infty)$ is the topological space of k -tuples $(v_1, \dots, v_k)$ of linearly independent vectors in $\mathbb{C}^\infty$ which are not necessarily orthogonal to each other. Observe that the inclusion $V_k(\mathbb{C}^\infty) \hookrightarrow \widetilde{V}_k(\mathbb{C}^\infty)$ is a homotopy equivalence: the Gram-Schmidt orthogonalization procedure provides a deformation retract $\widetilde{V}_k(\mathbb{C}^\infty) \rightarrow V_k(\mathbb{C}^\infty)$.

It will thus suffice to show that $\widetilde{V}_k(\mathbb{C}^\infty)$ is contractible. To this end, consider the ‘shift map’ $T^k: \mathbb{C}^\infty \rightarrow \mathbb{C}^\infty$ given on basis vectors by $T^k(e_n) := e_{n+k}$. This induces a map $T^k: \widetilde{V}_k(\mathbb{C}^\infty) \rightarrow \widetilde{V}_k(\mathbb{C}^\infty)$. Note that this map is homotopic to the constant map on $(e_1, \dots, e_k)$ via the straight-line homotopy:

$$H(v, t) := (1 - t)(e_1, \dots, e_k) + tT^k(v).$$

This is well-defined since the first k components of the vectors $T^k(v_i)$ are all zero. Similarly, the map T^k is homotopic to the identity on $\widetilde{V}_k(\mathbb{C}^\infty)$, again via the straight-line homotopy:

$$H(v, t) := (1 - t)v + tT^k(v).$$

To see that this is well-defined, assume for contradiction that for some time t the vectors are no longer linearly independent. Then there are scalars $\lambda_1, \dots, \lambda_k$, not all zero, such that $\sum_{i=1}^k \lambda_i((1-t)v_i + tT^k(v_i)) = 0$. But then the nonzero vector $w := \sum_{i=1}^k \lambda_i v_i$ satisfies $(1-t)w + tT^k(w) = 0$. For $t < 1$, this is impossible by considering the first nonzero coordinate of w ; for $t = 1$, it is impossible because T^k is injective. This contradicts linear independence of v .

It follows that the identity on $\tilde{V}_k(\mathbb{C}^\infty)$ is homotopic to a constant map, and hence this space is contractible. $\square$

Writing $U(k)$ also for the underlying anima of the topological group $U(k)$, we get:

Corollary 9.2.15. *The underlying anima of the Grassmannian $\mathrm{Gr}_k(\mathbb{C}^\infty)$ is isomorphic to the classifying anima $\mathbf{BU}(k)$.*

Proof. In light of Lemma 9.2.12 and Lemma 9.2.14, this is an instance of Proposition 9.2.11. $\square$

Corollary 9.2.16. *Let X be a paracompact Hausdorff space. There is a natural bijection*

$$\left[X, \bigsqcup_{k=0}^{\infty} \mathrm{Gr}_k(\mathbb{C}^\infty) \right] \xrightarrow{\cong} \pi_0 \mathrm{Vect}(X).$$

If X is a cell complex, this bijection can equivalently be written as

$$\pi_0 \mathrm{Hom}_{\mathrm{An}} \left(\Pi_\infty X, \bigsqcup_{k=0}^{\infty} \mathbf{BU}(k) \right) \xrightarrow{\cong} \pi_0 \mathrm{Vect}(X).$$

Proof. Both sides of the first bijection decompose according to locally constant functions $r: X \rightarrow \mathbb{N}$. For such a function, let $X_k := r^{-1}(k)$. These subspaces are clopen, and hence paracompact Hausdorff, and they form a topological disjoint-union decomposition

$$X = \bigsqcup_{k \geq 0} X_k.$$

A continuous map $X \rightarrow \bigsqcup_{k \geq 0} \mathrm{Gr}_k(\mathbb{C}^\infty)$ with component function r is precisely a family of continuous maps $X_k \rightarrow \mathrm{Gr}_k(\mathbb{C}^\infty)$. Similarly, a vector bundle with rank function r is precisely a family of rank k vector bundles over the X_k . The first bijection therefore follows on each clopen stratum from Theorem 9.2.4.

The Schubert cell structures above also make $\bigsqcup_{k \geq 0} \mathrm{Gr}_k(\mathbb{C}^\infty)$ a cell complex. If X is a cell complex as well, then Corollaries 2.3.7 and 9.2.15 identifies the source of the first bijection with the source of the second. $\square$

Remark 9.2.17. More generally, for every topological group G there exists a topological space BG together with a principal G -bundle $EG \rightarrow BG$ whose total space EG is contractible. In [Hus93, Section 4.12], it is shown that Milnor's construction of BG classifies principal G -bundles on paracompact Hausdorff spaces. Since EG is contractible, Proposition 9.2.11 implies that the underlying anima of BG is the classifying anima $\mathbf{BG}$. Thus the terminology ‘classifying anima’ is compatible with the classical terminology ‘classifying space’.

9.2.4 The stable Grassmannian

The passage from vector bundles to K-theory will use the stable Grassmannian rather than a strict point-set model for the whole commutative monoid of vector spaces.

Lemma 9.2.18. *Let*

$$Y_0 \hookrightarrow Y_1 \hookrightarrow Y_2 \hookrightarrow \dots$$

be a sequence of closed embeddings of T_1 topological spaces, and let $Y := \operatorname{colim}_n Y_n$ be the resulting colimit in $\mathbf{Top}$. Then every compact subset $K \subseteq Y$ is contained in some Y_n . In particular, every continuous map $f: X \rightarrow Y$ from a compact space factors through some Y_n .

Proof. Assume for contradiction that K is not contained in any Y_n , and choose $x_n \in K \setminus Y_n$. Set $A := \{x_n \mid n \geq 0\}$. This set is infinite: if it were finite, all its points would lie in some common stage Y_N , contradicting $x_n \notin Y_n$ for $n \geq N$. For every m , the intersection $A \cap Y_m$ is contained in the finite set $\{x_0, \dots, x_{m-1}\}$, hence is closed in the T_1 -space Y_m . The same holds for every subset of A . By the definition of the colimit topology, every subset of A is therefore closed in Y . Thus A is an infinite closed discrete subspace of K , contradicting compactness of K .

The last claim follows by applying the first to the compact subset $f(X) \subseteq Y$. $\square$

Proposition 9.2.19. *In the situation of Lemma 9.2.18, the canonical morphism*

$$\operatorname{colim}_n \Pi_\infty(Y_n) \longrightarrow \Pi_\infty(Y)$$

is an isomorphism of anima.

Proof. We use the Whitehead theorem for anima. The compactness lemma, applied to the point and to the interval, shows that the canonical morphism induces a bijection on path components. Fix a basepoint represented by some $y \in Y_N$. For every $q \geq 1$, compactness of S^q and $S^q \times [0, 1]$ shows that maps $S^q \rightarrow Y$ and homotopies between them factor through finite stages. Consequently,

$$\operatorname{colim}_{n \geq N} \pi_q(Y_n, y) \xrightarrow{\cong} \pi_q(Y, y).$$

By Remark 2.4.18, these are the homotopy groups of the corresponding underlying anima. Homotopy groups of pointed anima commute with filtered colimits, so the left-hand side is also the q -th homotopy group of $\operatorname{colim}_n \Pi_\infty(Y_n)$ at y . The claim now follows from Proposition 2.4.21. $\square$

Definition 9.2.20. We define the *stable Grassmannian*

$$BU := \operatorname{colim} \left(\operatorname{Gr}_0(\mathbb{C}^\infty) \xrightarrow{V \mapsto V \oplus \mathbb{C}} \operatorname{Gr}_1(\mathbb{C}^\infty) \xrightarrow{V \mapsto V \oplus \mathbb{C}} \operatorname{Gr}_2(\mathbb{C}^\infty) \rightarrow \dots \right),$$

where the transition maps are induced by the standard isometric identification $\mathbb{C}^\infty \oplus \mathbb{C} \cong \mathbb{C}^\infty$. The Schubert cell structures make these transition maps inclusions of subcomplexes, so their colimit BU is again a cell complex; see Hatcher [Hat03, p. 34]. In particular, the transition maps are closed embeddings, and we equip BU with the colimit topology.

Proposition 9.2.21. *Let $U := \operatorname{colim}_n U(n)$ be the stable unitary group, where the transition maps are block-sum with the identity on $\mathbb{C}$. Then there is an isomorphism of anima*

$$\Pi_\infty(BU) \simeq \mathbf{B}U.$$

Proof. The principal bundles $V_n(\mathbb{C}^\infty) \rightarrow \text{Gr}_n(\mathbb{C}^\infty)$ identify $\Pi_\infty \text{Gr}_n(\mathbb{C}^\infty)$ with $\mathbf{BU}(n)$ by Corollary 9.2.15. These identifications are compatible with the stabilization maps $\text{Gr}_n(\mathbb{C}^\infty) \rightarrow \text{Gr}_{n+1}(\mathbb{C}^\infty)$ and the block-sum inclusions $U(n) \rightarrow U(n+1)$. Since the Grassmannian stabilization maps are closed embeddings between Hausdorff spaces, Proposition 9.2.19 gives

$$\Pi_\infty(\mathbf{BU}) \simeq \text{colim}_n \mathbf{BU}(n).$$

The classifying-anima equivalence $\mathbf{B}: \text{Grp}(\mathbf{An}) \xrightarrow{\sim} \mathbf{An}_{*, \geq 1}$ preserves colimits. Since connected pointed anima are closed under filtered colimits, the filtered colimit of the $\mathbf{BU}(n)$ agrees whether computed in $\mathbf{An}$, $\mathbf{An}_*$, or $\mathbf{An}_{*, \geq 1}$. Moreover, filtered colimits in $\text{Grp}(\mathbf{An})$ are computed on underlying anima, since finite limits commute with filtered colimits in $\mathbf{An}$. Applying Proposition 9.2.19 also to the closed embeddings $U(n) \hookrightarrow U(n+1)$ therefore gives

$$\text{colim}_n \mathbf{BU}(n) \simeq \mathbf{B}(\text{colim}_n \Pi_\infty U(n)) \simeq \mathbf{B}\Pi_\infty(U),$$

which is the asserted classifying anima $\mathbf{BU}$. $\square$

Remark 9.2.22. The Grassmannian and Stiefel-manifold results above have real analogues with $\mathbb{C}$ replaced by $\mathbb{R}$ and $U(k)$ by $O(k)$, with the same proofs. In particular, $\text{Gr}_k(\mathbb{R}^\infty)$ classifies rank k real vector bundles over paracompact Hausdorff spaces, $V_k(\mathbb{R}^\infty)$ is contractible, and $\Pi_\infty \text{Gr}_k(\mathbb{R}^\infty) \simeq \mathbf{BO}(k)$. We will use these facts in Chapter 10.

9.3 Complex K-theory groups

We now pass from vector bundles to the classical complex K-theory groups. For a compact Hausdorff space, these are obtained by formally adjoining additive inverses to vector bundles. We will prove that K^0 is represented by the space $\mathbb{Z} \times \mathbf{BU}$, define the Bott class, and state the classical Bott periodicity theorem.

Definition 9.3.1 (Complex K-theory). Let X be a compact Hausdorff space. We define the *complex K-theory group* $K^0(X)$ as the Grothendieck group of the abelian monoid $\pi_0 \text{Vect}(X)$:

$$K^0(X) := (\pi_0 \text{Vect}(X))^{\text{grp}}.$$

In other words, $K^0(X)$ consists of equivalence classes of formal differences $E - E'$ of complex vector bundles over X , where we say that $E_1 - E'_1 = E_2 - E'_2$ whenever there is an isomorphism $E_1 \oplus E'_2 \oplus E_3 \cong E'_1 \oplus E_2 \oplus E_3$ of vector bundles over X for some $E_3 \in \pi_0 \text{Vect}(X)$.

Example 9.3.2. For $X = \text{pt}$, a vector bundle is simply a complex vector space, which up to isomorphism is determined by its dimension. We thus see that the abelian monoid $\pi_0 \text{Vect}(\text{pt})$ is isomorphic to $\mathbb{N}$, and hence $K^0(\text{pt}) \cong \mathbb{Z}$.

The abelian group $K^0(X)$ is contravariantly functorial in X : given a continuous map $f: X \rightarrow Y$, pullback along f determines a map

$$f^*: \pi_0 \text{Vect}(Y) \rightarrow \pi_0 \text{Vect}(X),$$

which induces a map $f^*: K^0(Y) \rightarrow K^0(X)$ on Grothendieck groups.

Tensor product distributes over direct sum, so it makes $\pi_0 \text{Vect}(X)$ into a commutative semiring. Its multiplication extends uniquely to a commutative ring structure on $K^0(X)$, with unit the class of the trivial line bundle. Pullback maps are morphisms of rings. The details are left to Exercise 9.4.

Corollary 9.3.3. *Every homotopy equivalence $f: X \rightarrow Y$ between compact Hausdorff spaces induces an isomorphism*

$$f^*: K^0(Y) \xrightarrow{\cong} K^0(X)$$

of abelian groups.

Proof. This follows from Theorem 9.1.4 by applying group completion. $\square$

Definition 9.3.4 (Reduced K-theory). Let X be a compact Hausdorff space. We define the abelian group $\tilde{K}^0(X)$ as the cokernel of the map $\mathbb{Z} \cong K^0(\text{pt}) \rightarrow K^0(X)$. This is called the *reduced K-theory group* of X .

When X has a basepoint x , this is also equivalent to the kernel of $K^0(X) \rightarrow K^0(\{x\}) \cong \mathbb{Z}$.

Exercise 9.3.5. Compute that $K^0(S^1) \cong \mathbb{Z}$ and $\tilde{K}^0(S^1) = 0$. *Hint:* See Exercise 9.1.6.

Exercise 9.3.6. Compute that $K^0(S^2) \cong \mathbb{Z} \times \mathbb{Z}$ and $\tilde{K}^0(S^2) \cong \mathbb{Z}$. Identifying S^2 with $\mathbb{CP}^1$, show that a generator of $\tilde{K}^0(S^2)$ is given by $[H] - [\underline{\mathbb{C}}]$, where $H \rightarrow \mathbb{CP}^1$ is the canonical line bundle. *Hint:* See Exercise 9.1.7.

We will use one more standard fact about vector bundles. It is the geometric reason that virtual bundles can be represented by honest bundles after adding enough trivial summands.

Theorem 9.3.7 (Stable complements, Hatcher [Hat03, Proposition 1.4]). *Let X be a compact Hausdorff space. For every complex vector bundle $E \rightarrow X$ there exists another vector bundle $E' \rightarrow X$ such that the direct sum $E \oplus E'$ is isomorphic to a trivial vector bundle $\underline{\mathbb{C}}^N := X \times \mathbb{C}^N$ for some $N \in \mathbb{N}$.*

Proof sketch. Choose finitely many local trivializations and a subordinate partition of unity. Multiplying the local coordinate functions by the partition-of-unity functions embeds E into a trivial bundle $X \times \mathbb{C}^N$. After choosing a Hermitian inner product on this trivial bundle, the orthogonal complements of the fibers of E assemble to a vector bundle E' with $E \oplus E' \cong \underline{\mathbb{C}}^N$. The details are the usual compactness and partition-of-unity argument; see Hatcher for the full proof. $\square$

Corollary 9.3.8. *There is a surjective morphism of monoids $\pi_0 \text{Vect}(X) \rightarrow \tilde{K}^0(X)$.*

Proof. See Exercise 9.2. $\square$

9.3.1 Representability of K^0

We now prove the classical representability theorem for K^0 using the Grassmannian classification theorem and stable complements.

Choose a linear isometric isomorphism $\mathbb{C}^\infty \oplus \mathbb{C}^\infty \cong \mathbb{C}^\infty$. Direct sum induces, up to homotopy, a multiplication on $\mathbb{Z} \times BU$: the $\mathbb{Z}$ -coordinates are added, while on the BU -coordinate we use the block-sum maps

$$\text{Gr}_n(\mathbb{C}^\infty) \times \text{Gr}_m(\mathbb{C}^\infty) \longrightarrow \text{Gr}_{n+m}(\mathbb{C}^\infty \oplus \mathbb{C}^\infty) \cong \text{Gr}_{n+m}(\mathbb{C}^\infty),$$

passing to the colimit. The proposition below identifies the resulting operation on $[X, \mathbb{Z} \times BU]$ with direct sum in $K^0(X)$, and hence shows in particular that it is independent of the choices and defines an abelian group structure.

Proposition 9.3.9. *Let X be a compact Hausdorff space. There is a natural isomorphism of abelian groups*

$$[X, \mathbb{Z} \times BU] \xrightarrow{\cong} K^0(X).$$

Proof. We first construct the map. Let $f: X \rightarrow \mathbb{Z} \times BU$ be continuous. Since X is compact and $\mathbb{Z}$ is discrete, the composite $X \rightarrow \mathbb{Z}$ has finite image. Write $X_m \subseteq X$ for the clopen subset on which this composite has value m . By Lemma 9.2.18, the restriction $f_m: X_m \rightarrow BU$ factors through some stage $\text{Gr}_n(\mathbb{C}^\infty)$. Let $E_m \rightarrow X_m$ be the pullback of the universal rank n bundle along this factorization. We assign to f_m the class

$$[E_m] - n[\underline{\mathbb{C}}] + m[\underline{\mathbb{C}}] \in K^0(X_m).$$

These classes glue over the finite clopen decomposition $X = \bigsqcup_m X_m$ to a class in $K^0(X)$.

This construction is independent of all choices. Passing from $\text{Gr}_n(\mathbb{C}^\infty)$ to $\text{Gr}_{n+1}(\mathbb{C}^\infty)$ replaces E_m by $E_m \oplus \underline{\mathbb{C}}$, and hence leaves $[E_m] - n[\underline{\mathbb{C}}] + m[\underline{\mathbb{C}}]$ unchanged. If two maps are homotopic, the homotopy has constant $\mathbb{Z}$ -coordinate on each $X_m \times [0, 1]$ and, again by compactness, factors through a common Grassmannian stage after stabilization. Pulling back the universal bundle along the two ends gives isomorphic bundles by Theorem 9.1.4. Thus we get a well-defined natural map

$$\Phi_X: [X, \mathbb{Z} \times BU] \rightarrow K^0(X).$$

We prove that Φ_X is surjective. Let $\alpha \in K^0(X)$, and let $X = \bigsqcup_m X_m$ be the finite clopen decomposition according to the virtual rank of α . It suffices to work on one X_m , so assume α has constant rank m . Choose a presentation $\alpha = [E] - [F]$. By Theorem 9.3.7, choose F' and N such that $F \oplus F' \cong \underline{\mathbb{C}}^N$. Then

$$\alpha = [E \oplus F'] - N[\underline{\mathbb{C}}].$$

The bundle $E \oplus F'$ has rank $N + m$, so Theorem 9.2.4 classifies it by a map $g: X_m \rightarrow \text{Gr}_{N+m}(\mathbb{C}^\infty) \subseteq BU$. Viewed as a map $X_m \rightarrow \{m\} \times BU$, this maps under Φ_{X_m} to

$$[E \oplus F'] - (N + m)[\underline{\mathbb{C}}] + m[\underline{\mathbb{C}}] = [E] - [F] = \alpha.$$

Gluing over the finitely many m proves surjectivity.

For injectivity, suppose $f_0, f_1: X \rightarrow \mathbb{Z} \times BU$ have the same image under Φ_X . Their images in $K^0(X)$ have the same rank function, so the $\mathbb{Z}$ -coordinates of f_0 and f_1 agree. We again restrict to a clopen piece X_m and choose factorizations $f_i: X_m \rightarrow \text{Gr}_{n_i}(\mathbb{C}^\infty)$ classifying bundles E_i . The equality $\Phi_X(f_0) = \Phi_X(f_1)$ gives

$$[E_0] - n_0[\underline{\mathbb{C}}] = [E_1] - n_1[\underline{\mathbb{C}}]$$

in $K^0(X_m)$. By the definition of the Grothendieck group, there exists a vector bundle F and an isomorphism

$$E_0 \oplus \underline{\mathbb{C}}^{n_1} \oplus F \cong E_1 \oplus \underline{\mathbb{C}}^{n_0} \oplus F.$$

Choose F' with $F \oplus F' \cong \underline{\mathbb{C}}^q$ by Theorem 9.3.7. After stabilizing f_0 by $n_1 + q$ steps and f_1 by $n_0 + q$ steps, the two maps land in the same Grassmannian $\text{Gr}_{n_0+n_1+q}(\mathbb{C}^\infty)$ and classify isomorphic vector bundles. By Theorem 9.2.4, they are homotopic. These homotopies glue over the finite clopen decomposition of X , proving injectivity.

The construction is compatible with addition. If maps through $\{m\} \times \text{Gr}_n(\mathbb{C}^\infty)$ and $\{m'\} \times \text{Gr}_{n'}(\mathbb{C}^\infty)$ classify bundles E and E' , then their sum classifies $E \oplus E'$ at stage $n + n'$ and has $\mathbb{Z}$ -coordinate $m + m'$. Therefore

$$[E \oplus E'] - (n + n')[\underline{\mathbb{C}}] + (m + m')[\underline{\mathbb{C}}]$$

equals the sum of the two associated K^0 -classes. Thus Φ_X is an isomorphism of abelian groups. $\square$

9.3.2 The Bott class and Bott periodicity

Definition 9.3.10 (Bott class). Let $H \rightarrow \mathbb{CP}^1$ be the tautological line bundle. The *Bott class* is

$$\beta := [H] - [\underline{\mathbb{C}}] \in \tilde{K}^0(\mathbb{CP}^1).$$

By Exercise 9.3.6, this class generates $\tilde{K}^0(\mathbb{CP}^1) \cong \mathbb{Z}$.

The reduced external tensor product of vector bundles induces a pairing

$$\tilde{K}^0(X) \otimes \tilde{K}^0(Y) \longrightarrow \tilde{K}^0(X \wedge Y)$$

for pointed compact Hausdorff spaces; see Hatcher [Hat03, p. 54]. The following classical theorem is the geometric input behind the periodicity of complex K-theory.

Theorem 9.3.11 (Bott periodicity for $\tilde{K}^0$, Hatcher [Hat03, Theorem 2.11]). *For every pointed compact Hausdorff space X , external multiplication by the Bott class induces an isomorphism*

$$\tilde{K}^0(X) \xrightarrow{\cong} \tilde{K}^0(X \wedge \mathbb{CP}^1) \cong \tilde{K}^0(\Sigma^2 X).$$

9.4 Complex K-theory spectra

The classical construction above uses only isomorphism classes of vector bundles. To construct a representing spectrum, we now retain the automorphisms of vector bundles and all their homotopies, and then apply the homotopical group completion theorem. This produces connective complex K-theory. Inverting the Bott class then produces its periodic counterpart.

9.4.1 The anima of vector bundles

Let X be a topological space. Write $\text{Vect}^{\text{disc}}(X)$ for the 1-category of complex vector bundles over X and bundle maps, with discrete hom sets. Its groupoid core remembers the isomorphisms between bundles, but regards them as discrete and therefore misses their topology. To retain this topology, we instead regard vector bundles and their isomorphisms as forming a topologically enriched groupoid.

For a compact Hausdorff space X , the bundle isomorphisms $E \rightarrow E'$ form a topological space $\text{Iso}_X(E, E')$ with the compact-open topology, and these spaces are the hom objects of the enriched groupoid just described. A continuous n -simplex in $\text{Iso}_X(E, E')$ is equivalently an isomorphism between the pullbacks of E and E' over $X \times |\Delta^n|$. This leads to the formula

$$\text{Vect}(X)^\simeq := \text{colim}_{[n] \in \Delta^{\text{op}}} \text{Vect}^{\text{disc}}(X \times |\Delta^n|)^\simeq.$$

The simplicial formula is a model for the homotopy coherent nerve of the topologically enriched groupoid, while still making sense for arbitrary X . Direct sum and tensor product are available degreewise. In Part II we prove that they assemble, with all the required distributivity and higher coherences, into a functor

$$\mathbf{Vect}(-)^{\simeq} : \mathbf{Top}^{\mathrm{op}} \longrightarrow \mathbf{CRig}(\mathbf{An}),$$

whose addition is direct sum and whose multiplication is tensor product; see Definition 19.7.8 and Proposition 19.7.9. Here $\mathbf{CRig}(\mathbf{An}) := \mathbf{CAlg}(\mathbf{CMon}(\mathbf{An}))$ denotes the ∞ -category of commutative semiring anima; forgetting multiplication recovers the already familiar functor to $\mathbf{CMon}(\mathbf{An})$. For compact Hausdorff X , the resulting anima $\mathbf{Vect}(X)^{\simeq}$ is a model for the homotopy coherent nerve of the topologically enriched groupoid of vector bundles and bundle isomorphisms. If X is paracompact Hausdorff, then there is a bijection

$$\pi_0 \mathbf{Vect}(X)^{\simeq} \cong \pi_0 \mathbf{Vect}(X).$$

For a point, the components and the semiring operations admit the following explicit description.

Proposition 9.4.1. *The underlying anima of $\mathbf{Vect}(\mathrm{pt})^{\simeq}$ is the disjoint union $\bigsqcup_{n \geq 0} \mathbf{BU}(n)$. Under this isomorphism, the addition map*

$$\left(\bigsqcup_{k=0}^{\infty} \mathbf{BU}(k) \right) \times \left(\bigsqcup_{l=0}^{\infty} \mathbf{BU}(l) \right) \rightarrow \left(\bigsqcup_{m=0}^{\infty} \mathbf{BU}(m) \right)$$

is given on the (k, l) -component by the map $\mathbf{BU}(k) \times \mathbf{BU}(l) \rightarrow \mathbf{BU}(k+l)$ induced by the block-sum inclusion

$$U(k) \times U(l) \hookrightarrow U(k+l), \quad (A, B) \mapsto \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}.$$

The multiplication map is given on the (k, l) -component by the map $\mathbf{BU}(k) \times \mathbf{BU}(l) \rightarrow \mathbf{BU}(kl)$ induced by the tensor-product representation

$$U(k) \times U(l) \longrightarrow U(kl), \quad (A, B) \longmapsto A \otimes B.$$

In particular, the induced semiring structure on $\pi_0 \mathbf{Vect}(\mathrm{pt}) \cong \mathbb{N}$ is the usual one.

Proof. By Definition 19.7.8 and Proposition 19.7.9, the rank n component of $\mathbf{Vect}(\mathrm{pt})^{\simeq}$ is the geometric realization of the simplicial groupoid whose value at $[q]$ is the full subgroupoid

$$\left(\mathbf{Vect}^{\mathrm{disc}}(|\Delta^q|)^{\simeq} \right)_{\mathrm{rk}=n}$$

spanned by the rank n bundles. Every rank n vector bundle over $|\Delta^q|$ is trivial. Consequently, the full subgroupoid on the standard trivial bundle $|\Delta^q| \times \mathbb{C}^n$ is an equivalent one-object groupoid. Its automorphism group is

$$\mathrm{Hom}_{\mathrm{Top}}(|\Delta^q|, \mathrm{GL}_n(\mathbb{C})),$$

and these groups assemble into the singular simplicial group of $\mathrm{GL}_n(\mathbb{C})$. Geometric realization preserves finite products by Lemma 21.6.12, so it commutes with the formation of classifying anima of simplicial groups. Since the realization of the singular simplicial group is the underlying

group anima of $\mathrm{GL}_n(\mathbb{C})$, the realization of these one-object groupoids is $\mathbf{B}\mathrm{GL}_n(\mathbb{C})$. Taking the disjoint union over all ranks therefore gives

$$\mathrm{Vect}(\mathrm{pt})^{\simeq} \simeq \bigsqcup_{n \geq 0} \mathbf{B}\mathrm{GL}_n(\mathbb{C}).$$

The inclusions $U(n) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ are homotopy equivalences by polar decomposition and commute with block sums and tensor products. They therefore identify $\mathbf{B}\mathrm{GL}_n(\mathbb{C})$ with $\mathbf{B}U(n)$ compatibly with both operations.

Indeed, direct sum sends $(\mathbb{C}^k, \mathbb{C}^l)$ to $\mathbb{C}^{k+l}$ and induces the block-sum inclusion on automorphism groups. Similarly, tensor product sends $(\mathbb{C}^k, \mathbb{C}^l)$ to $\mathbb{C}^k \otimes \mathbb{C}^l \cong \mathbb{C}^{kl}$ and induces the tensor-product representation on automorphism groups. Passing to classifying anima gives the stated formulas. $\square$

9.4.2 Connective complex K-theory

The definition of the complex K-theory groups via group completion admits the following homotopical refinement.

Definition 9.4.2. We denote by

$$\mathrm{BUP} := (\mathrm{Vect}(\mathrm{pt})^{\simeq})^{\mathrm{grp}} \in \mathrm{CGrp}(\mathrm{An})$$

the group completion of $\mathrm{Vect}(\mathrm{pt})^{\simeq}$ with respect to direct sum, using the left adjoint $(-)^{\mathrm{grp}}$ from Corollary 5.4.4. We define the *connective complex K-theory spectrum* $\mathrm{ku} \in \mathrm{Sp}_{\geq 0}$ to be the connective spectrum corresponding to BUP under the equivalence $\mathrm{Sp}_{\geq 0} \simeq \mathrm{CGrp}(\mathrm{An})$ of Theorem 5.4.5.

While group completion is abstract in general, the telescope form of Theorem 5.5.2 gives an explicit description here.

Corollary 9.4.3. *The underlying anima of BUP is $\mathbb{Z} \times \mathbf{B}U$:*

$$\mathrm{BUP} \simeq (\mathrm{Vect}(\mathrm{pt})^{\simeq})^{\mathrm{grp}} \simeq \mathbb{Z} \times \mathbf{B}U.$$

Proof. By Proposition 9.4.1, the commutative monoid $M = \mathrm{Vect}(\mathrm{pt})^{\simeq}$ has underlying anima $\bigsqcup_{k \geq 0} \mathbf{B}U(k)$, with addition induced by block sum. Let $x \in M$ be the point corresponding to the one-dimensional vector space $\mathbb{C}$. The telescope $T(M, x)$ is obtained by repeatedly adding this line. Its component of virtual rank $m \in \mathbb{Z}$ is the colimit

$$\mathrm{colim}_{n \geq \max(0, -m)} \mathbf{B}U(m + n),$$

which is canonically $\mathbf{B}U$. Thus $T(M, x) \simeq \mathbb{Z} \times \mathbf{B}U$.

The two hypotheses of Theorem 5.5.2 are immediate. The class of x is the generator $1 \in \pi_0(M) \cong \mathbb{N}$. The fundamental group of $\mathbf{B}U$ is $\pi_0(U)$, which is trivial because the stable unitary group is connected. Hence $T(M, x) \rightarrow M^{\mathrm{grp}}$ is an isomorphism. $\square$

The connective spectrum ku can be turned into a commutative ring spectrum; equivalently, the commutative group BUP admits the structure of a commutative ring object in An . This structure ultimately comes from the fact that the anima $\mathrm{Vect}(\mathrm{pt})^{\simeq}$ not only admits an addition via direct sum of vector spaces, but also a multiplication via tensor product of vector spaces. The required distributivity and higher coherences are constructed in Part II, see Section 19.7. The commutative ring spectrum structure on ku is then obtained by passing to group completions; see Proposition 19.7.11.

Corollary 9.4.4. *Let X be a compact Hausdorff space which has the homotopy type of a CW-complex. There is a natural isomorphism of commutative rings*

$$K^0(X) \cong \text{ku}^0(\Pi_\infty(X)).$$

Proof. The space $\mathbb{Z} \times BU$ is a cell complex by the Schubert cell description above. Choose a homotopy equivalence $X' \rightarrow X$ with X' a cell complex. Then

$$[X, \mathbb{Z} \times BU] \cong [X', \mathbb{Z} \times BU] \cong \pi_0 \text{Hom}_{\text{An}}(\Pi_\infty X', \Pi_\infty(\mathbb{Z} \times BU)) \cong \pi_0 \text{Hom}_{\text{An}}(\Pi_\infty X, \Pi_\infty(\mathbb{Z} \times BU)),$$

where the middle isomorphism is Corollary 2.3.7. Combining this with Propositions 9.2.21 and 9.3.9 and Corollary 9.4.3 and the adjunction $\mathbb{S}[-] \dashv \Omega^\infty$ gives

$$\begin{aligned} K^0(X) &\cong [X, \mathbb{Z} \times BU] \\ &\cong \pi_0 \text{Hom}_{\text{An}}(\Pi_\infty(X), \Pi_\infty(\mathbb{Z} \times BU)) \\ &\cong \pi_0 \text{Hom}_{\text{An}}(\Pi_\infty(X), \mathbb{Z} \times \mathbf{B}U) \\ &\cong \pi_0 \text{Hom}_{\text{An}}(\Pi_\infty(X), \mathbf{B}U) \\ &\cong \pi_0 \text{Hom}_{\text{Sp}}(\mathbb{S}[\Pi_\infty(X)], \text{ku}) \\ &= \text{ku}^0(\Pi_\infty(X)). \end{aligned}$$

It remains to check multiplication. Set $M := \text{Vect}(\text{pt})^\simeq$. The unit map $M \rightarrow M^{\text{grp}} = \mathbf{B}U$ is a morphism of commutative semiring objects, since group completion is symmetric monoidal by Proposition 16.6.4. For a vector bundle E over X , its classifying map to M therefore determines a class in $\text{ku}^0(\Pi_\infty X)$, and Proposition 9.4.1 shows that direct sum and tensor product of bundles map to the sum and product of the corresponding classes. This semiring map

$$\pi_0 \text{Vect}(X) \longrightarrow \text{ku}^0(\Pi_\infty X)$$

extends uniquely to a ring map from $K^0(X)$. Under the description $M^{\text{grp}} \simeq \mathbb{Z} \times \mathbf{B}U$ of Corollary 9.4.3, this extension is the isomorphism displayed above: an honest vector bundle is sent to its stable classifying map. Hence that isomorphism is multiplicative. $\square$

For a pointed compact Hausdorff space X of the homotopy type of a CW-complex, naturality with respect to the basepoint shows that the preceding comparison identifies the kernels of the restriction maps to the point. It therefore restricts to a natural isomorphism

$$\widetilde{K}^0(X) \cong \widetilde{\text{ku}}^0(\Pi_\infty X).$$

In particular, for $n \geq 0$ we obtain $\widetilde{K}^0(S^n) \cong \pi_n \text{ku}$.

For pointed spaces X and Y satisfying the same hypotheses, the reduced comparisons are compatible with external products. Indeed, if $E \rightarrow X$ and $F \rightarrow Y$ are vector bundles, then multiplication of their classifying maps classifies

$$E \boxtimes F := \text{pr}_X^* E \otimes \text{pr}_Y^* F \longrightarrow X \times Y.$$

Thus the unreduced comparison preserves external products. Reduced classes correspond to based classifying maps, and their smash product followed by the multiplication on $\mathbf{B}U$ pulls back along

$X \times Y \rightarrow X \wedge Y$ to the ordinary external product. This is also the defining property of the classical reduced external product, so the reduced comparison preserves it.

Under the preceding reduced comparison and the homeomorphism $\mathbb{CP}^1 \cong S^2$, the Bott class of Definition 9.3.10 determines a class

$$\beta \in \pi_2 \text{ku}.$$

Construction 9.4.5 (The Bott self-map). The Bott class is represented by a morphism $\beta: \mathbb{S}[2] \rightarrow \text{ku}$, or equivalently by a morphism $\beta: \mathbb{S} \rightarrow \text{ku}[-2]$. Multiplication by this class defines a morphism of spectra

$$b: \text{ku} \simeq \mathbb{S} \otimes \text{ku} \xrightarrow{\beta \otimes \text{id}} \text{ku}[-2] \otimes \text{ku} \xrightarrow{\mu[-2]} \text{ku}[-2],$$

where μ denotes the multiplication on ku . We call b the *Bott map*.

By the compatibility of the preceding comparison with reduced external products, the map

$$b_*: \pi_n \text{ku} \longrightarrow \pi_{n+2} \text{ku}$$

for $n \geq 0$ is reduced external multiplication with the classical Bott class. More explicitly, if $q: S^n \times \mathbb{CP}^1 \rightarrow S^n \wedge \mathbb{CP}^1$ is the quotient map, then the corresponding reduced class is characterized by the fact that its pullback along q is the ordinary external product.

Corollary 9.4.6 (Bott periodicity for ku). *The Bott map induces an isomorphism*

$$b_*: \pi_n \text{ku} \xrightarrow{\cong} \pi_{n+2} \text{ku}$$

for every $n \geq 0$.

Proof. Apply Theorem 9.3.11 to $X = S^n$. Under the comparison above,

$$\pi_n \text{ku} \cong \widetilde{\text{ku}}^0(S^n) \cong \widetilde{K}^0(S^n),$$

the map b_* is external multiplication by the Bott class. The claim therefore follows from classical Bott periodicity. $\square$

9.4.3 Periodic complex K-theory

The connective spectrum ku admits a 2-periodic variant KU , obtained by inverting the Bott class.

Definition 9.4.7. The *periodic complex K-theory spectrum* is the colimit

$$\text{KU} := \text{ku}[\beta^{-1}] := \text{colim} \left(\text{ku} \xrightarrow{b} \text{ku}[-2] \xrightarrow{b[-2]} \text{ku}[-4] \rightarrow \cdots \right) \in \text{Sp}.$$

We write $\iota: \text{ku} \rightarrow \text{KU}$ for the canonical map into the colimit.

Corollary 9.4.8. *The periodic complex K-theory spectrum KU admits a canonical structure of commutative ring spectrum, and the map $\text{ku} \rightarrow \text{KU}$ is a morphism of commutative ring spectra.*

Proof. By Proposition 19.7.11, the spectrum ku is a commutative ring spectrum, so $\beta \in \pi_2 \mathrm{ku}$ is a homogeneous element of the graded-commutative ring $\pi_* \mathrm{ku}$. Let $S := \{1, \beta, \beta^2, \dots\}$ be the multiplicative subset it generates.

By Corollary 8.4.5, the localization $\mathrm{ku}[\beta^{-1}] := \mathrm{ku}[S^{-1}]$ exists in $\mathrm{CAlg}(\mathrm{Sp})$, and the localization map $\eta: \mathrm{ku} \rightarrow \mathrm{ku}[\beta^{-1}]$ is a morphism of commutative ring spectra. By Lemma 8.4.6, its underlying spectrum is the telescope

$$\mathrm{colim} \left(\mathrm{ku} \xrightarrow{b} \mathrm{ku}[-2] \xrightarrow{b[-2]} \mathrm{ku}[-4] \rightarrow \dots \right),$$

where b is multiplication by β , i.e. the Bott map of Construction 9.4.5. This colimit is precisely the spectrum KU of Definition 9.4.7, and under this identification η is the canonical map $\iota: \mathrm{ku} \rightarrow \mathrm{KU}$. $\square$

Lemma 9.4.9. *For every $k \in \mathbb{Z}$ there is a natural isomorphism*

$$\pi_k \mathrm{KU} \cong \mathrm{colim} \left(\pi_k \mathrm{ku} \xrightarrow{b_*} \pi_{k+2} \mathrm{ku} \xrightarrow{b_*} \pi_{k+4} \mathrm{ku} \rightarrow \dots \right).$$

Proof. By Lemma 4.4.27, the functor π_k preserves the filtered colimit defining KU . Moreover $\pi_k(\mathrm{ku}[-2j]) = \pi_{k+2j} \mathrm{ku}$, and the transition maps are induced by the Bott map. $\square$

Theorem 9.4.10. *The map $\iota: \mathrm{ku} \rightarrow \mathrm{KU}$ exhibits ku as the connective cover of KU : it induces an isomorphism on π_k for $k \geq 0$, and hence an equivalence*

$$\mathrm{ku} \xrightarrow{\cong} \tau_{\geq 0} \mathrm{KU}.$$

Proof. For $k \geq 0$, Lemma 9.4.9 writes $\pi_k \mathrm{KU}$ as a filtered colimit whose transition maps

$$b_*: \pi_{k+2j} \mathrm{ku} \rightarrow \pi_{k+2j+2} \mathrm{ku}$$

are all isomorphisms by Corollary 9.4.6. Hence $\pi_k(\iota)$ is an isomorphism for $k \geq 0$. Since ku is connective, the map ι factors through the connective cover as

$$\mathrm{ku} \xrightarrow{\bar{\iota}} \tau_{\geq 0} \mathrm{KU} \rightarrow \mathrm{KU}$$

by Corollary 5.4.6. The map $\bar{\iota}$ is an isomorphism on all homotopy groups, because both source and target are connective and we have already checked the non-negative homotopy groups. It is therefore an equivalence by Corollary 4.4.26. $\square$

Corollary 9.4.11. *For every anima X and every integer $k \leq 0$, the map $\mathrm{ku} \rightarrow \mathrm{KU}$ induces an isomorphism*

$$\mathrm{ku}^k(X) \xrightarrow{\cong} \mathrm{KU}^k(X).$$

Proof. For every connective spectrum Y , the equivalence $\mathrm{ku} \simeq \tau_{\geq 0} \mathrm{KU}$ induces an equivalence

$$\mathrm{Hom}_{\mathrm{Sp}}(Y, \mathrm{ku}) \xrightarrow{\simeq} \mathrm{Hom}_{\mathrm{Sp}}(Y, \mathrm{KU}).$$

Apply this to $Y = \Sigma^{-k} \mathbb{S}[X]$, which is connective for $k \leq 0$. $\square$

Proposition 9.4.12 (Periodicity). *There is an equivalence $\Sigma^2 \mathrm{KU} \simeq \mathrm{KU}$.*

Proof. The spectrum $KU[-2]$ is the colimit of the diagram defining KU with its first term omitted:

$$ku[-2] \xrightarrow{b[-2]} ku[-4] \rightarrow \cdots$$

This subdiagram is cofinal in the original sequential diagram, so $KU[-2] \simeq KU$. Applying Σ^2 gives the desired equivalence. $\square$

Remark 9.4.13. The periodicity isomorphism of Proposition 9.4.12 is multiplication by the image of β in $\pi_2 KU$. Thus that image is invertible in the graded ring $\pi_*(KU)$.

Exercise 9.4.14. Deduce that there are isomorphisms of graded rings

$$\pi_* ku \cong \mathbb{Z}[\beta] \quad \text{and} \quad \pi_* KU \cong \mathbb{Z}[\beta^{\pm 1}],$$

with $|\beta| = 2$. Equivalently,

$$\pi_k(KU) \cong \begin{cases} \mathbb{Z} & k \text{ even,} \\ 0 & k \text{ odd.} \end{cases}$$

Hint: Use $\pi_0 ku \cong \mathbb{Z}$ and $\pi_1 ku \cong 0$ from Corollary 9.4.3, together with Corollary 9.4.6. Then apply the graded-localization calculation of Lemma 8.4.6 to KU .

The homotopy-group calculation and Proposition 7.2.16 give an abstract noncanonical isomorphism of spectra

$$KU_{\mathbb{Q}} \simeq \bigoplus_{n \in \mathbb{Z}} H\mathbb{Q}[2n].$$

The chosen Bott class allows us to refine this to an isomorphism of commutative ring spectra. Let $\mathbb{Q}[\beta^{\pm 1}]$ denote the commutative DGA with zero differential on an invertible generator β in degree 2, and let $H\mathbb{Q}[\beta^{\pm 1}]$ be its Eilenberg–MacLane ring spectrum in the sense of Section 8.3. Its underlying spectrum is $\bigoplus_{n \in \mathbb{Z}} H\mathbb{Q}[2n]$. The canonical map

$$\bigoplus_{n \in \mathbb{Z}} H\mathbb{Q}[2n] \longrightarrow \prod_{n \in \mathbb{Z}} H\mathbb{Q}[2n]$$

is an isomorphism, since it induces an isomorphism on every homotopy group.

Proposition 9.4.15 (Rational periodic complex K-theory). *The Bott class determines an isomorphism of commutative ring spectra*

$$H\mathbb{Q}[\beta^{\pm 1}] \xrightarrow{\simeq} KU_{\mathbb{Q}}$$

which sends the Laurent polynomial generator β to the rationalized Bott class.

Proof. By the smashing description of rationalization and Theorem 7.2.14, we have $KU_{\mathbb{Q}} \simeq KU \otimes H\mathbb{Q}$, so $KU_{\mathbb{Q}}$ is canonically a commutative $H\mathbb{Q}$ -algebra. Under the symmetric monoidal equivalence $D(\mathbb{Q}) \simeq \text{Mod}_{H\mathbb{Q}}$ of Corollary 8.3.4, the polynomial algebra $\mathbb{Q}[\beta]$ is the free commutative $\mathbb{Q}$ -algebra on a generator in degree 2. The rationalized Bott class therefore determines a morphism of commutative $H\mathbb{Q}$ -algebras

$$H\mathbb{Q}[\beta] \longrightarrow KU_{\mathbb{Q}}.$$

The canonical map $H\mathbb{Q}[\beta] \rightarrow H\mathbb{Q}[\beta^{\pm 1}]$ exhibits its target as $(H\mathbb{Q}[\beta])[\beta^{-1}]$: indeed, the induced map from this localization is an isomorphism on homotopy groups by Lemma 8.4.6. Since the Bott

class is invertible in $\pi_*(\mathrm{KU}_{\mathbb{Q}})$, the preceding morphism therefore extends uniquely to $H\mathbb{Q}[\beta^{\pm 1}]$. It induces the isomorphism

$$\mathbb{Q}[\beta^{\pm 1}] \xrightarrow{\cong} \pi_*(\mathrm{KU}_{\mathbb{Q}})$$

from Exercise 9.4.14 and Lemma 7.2.4, and hence is an isomorphism of commutative ring spectra. $\square$

Definition 9.4.16 (Chern character). The *Chern character* is the stable multiplicative cohomology operation induced by the composite

$$\mathrm{KU} \longrightarrow \mathrm{KU}_{\mathbb{Q}} \xrightarrow{\cong} H\mathbb{Q}[\beta^{\pm 1}],$$

where the second map is the inverse of the isomorphism in Proposition 9.4.15. Thus, for every anima X and every $k \in \mathbb{Z}$, it gives a natural morphism

$$\mathrm{ch}: \mathrm{KU}^k(X) \longrightarrow \prod_{n \in \mathbb{Z}} H^{k+2n}(X; \mathbb{Q}).$$

Remark 9.4.17. Normalize the generator of $H^2(\mathbb{CP}^{\infty}; \mathbb{Q})$ by requiring that its restriction to $\mathbb{CP}^1$ is $c_1(H)$, where H is the tautological line bundle of Definition 9.3.10. With this convention, the operation above agrees with the classical Chern character characterized on line bundles by

$$\mathrm{ch}(L) = e^{c_1(L)}.$$

We do not develop the theory of Chern classes here; see Milnor and Stasheff [MS74] and Hatcher [Hat03, Section 4.1].

Corollary 9.4.18. *Let X be a finite anima and let $k \in \mathbb{Z}$. Then the Chern character induces an isomorphism*

$$\mathrm{KU}^k(X)_{\mathbb{Q}} \xrightarrow{\cong} \bigoplus_{n \in \mathbb{Z}} H^{k+2n}(X; \mathbb{Q}).$$

Proof. Since X is finite, the functor $\mathrm{hom}(\mathbb{S}[X], -): \mathrm{Sp} \rightarrow \mathrm{Sp}$ commutes with rationalization and with direct sums. Applying it to Proposition 9.4.15 gives

$$\mathrm{hom}(\mathbb{S}[X], \mathrm{KU})_{\mathbb{Q}} \simeq \mathrm{hom}(\mathbb{S}[X], \mathrm{KU}_{\mathbb{Q}}) \simeq \bigoplus_{n \in \mathbb{Z}} \mathrm{hom}(\mathbb{S}[X], H\mathbb{Q}[2n]).$$

The resulting map is induced by the Chern character of Definition 9.4.16. Since X is finite, only finitely many of the groups $H^{k+2n}(X; \mathbb{Q})$ are nonzero, so the product in its definition agrees with the direct sum above. Passing to π_{-k} gives the claim. $\square$

Exercises

Exercise 9.1. Let E_1 and E_2 be complex vector bundles over a topological space X . Show that their direct sum

$$E_1 \oplus E_2 := E_1 \times_X E_2$$

is again a complex vector bundle over X .

Exercise 9.2. Let X be a compact Hausdorff space and consider the composite

$$\pi_0 \text{Vect}(X) \rightarrow K^0(X) \rightarrow \tilde{K}^0(X),$$

which is a morphism of abelian monoids. Show that this map is surjective, and that two vector bundles E and E' are sent to the same element of $\tilde{K}^0(X)$ if and only if there exists an isomorphism $E \oplus \underline{\mathbb{C}}^n \cong E' \oplus \underline{\mathbb{C}}^m$ for some n and m .

Hint: Use Theorem 9.3.7.

Exercise 9.3. Let X be a cell complex. Using the commutative monoid structure on

$$\text{Vect}(\text{pt})^{\simeq} \simeq \bigsqcup_{k=0}^{\infty} \mathbf{BU}(k),$$

show that $\text{Hom}_{\text{An}}(\Pi_{\infty} X, \text{Vect}(\text{pt})^{\simeq})$ admits a canonical commutative monoid structure. Under the classification of Corollary 9.2.16, identify the induced abelian monoid structure on its set of components with direct sum of vector bundles.

Exercise 9.4. Construct for two complex vector bundles E_1 and E_2 over X their *tensor product* $E_1 \otimes E_2$ whose fiber over $x \in X$ is given by $(E_1)_x \otimes_{\mathbb{C}} (E_2)_x$, and show that it is again a vector bundle. Show that your construction defines an operation

$$- \otimes -: \pi_0 \text{Vect}(X) \times \pi_0 \text{Vect}(X) \rightarrow \pi_0 \text{Vect}(X)$$

which is unital, commutative and associative. Show that it is distributive with respect to $- \oplus -$, making $\pi_0 \text{Vect}(X)$ into a *commutative semiring*.

Deduce from this that the complex K-theory group $K^0(X)$ of X is in fact a *commutative ring*.

Exercise 9.5. For compact Hausdorff spaces X and Y , show that pullback along the inclusions induces an isomorphism of groups

$$K^0(X \sqcup Y) \xrightarrow{\sim} K^0(X) \times K^0(Y).$$

Exercise 9.6. For a connected compact Hausdorff space X , produce a ‘virtual rank map’, i.e. a group homomorphism

$$\text{rk}: K^0(X) \rightarrow \mathbb{Z}.$$

Show that there is an isomorphism of groups

$$\tilde{K}^0(X) \xrightarrow{\sim} \{\alpha \in K^0(X) \mid \text{rk}(\alpha) = 0\}.$$

10 Thom spectra and bordism theory

In this chapter, we study a family of (co)homology theories called *(co)bordism theory*. Their representing spectra are known as the *Thom spectra*.

The origins behind these theories lie in Thom’s paper [Tho54]. Thom introduced the relation of *cobordism* on the set of closed smooth n -manifolds, with M and N being identified if their disjoint union $M \sqcup N$ is the boundary of some compact $(n + 1)$ -manifold W . He completely determined unoriented bordism and made decisive progress on two problems posed by Steenrod: which homology classes of a space are represented by maps from closed manifolds, and which closed smooth manifolds are boundaries.

Given a space X , we may equip the manifolds with maps to X , producing the unoriented bordism groups $\Omega_n^O(X)$. These groups assemble into a homology theory, a viewpoint developed systematically by Atiyah [Ati61a]. The spectrum MO representing this theory can be constructed using the notion of the *Thom space* of a vector bundle. Thom invented this notion in order to generalize the result of Pontryagin [Pon59] that the groups of *framed* manifolds up to cobordism are isomorphic to the stable stems $\pi_*(\mathbb{S})$. Thom realized that one could dispense with the framings at the cost of replacing the spheres in Pontryagin’s computation by the Thom spaces of certain vector bundles. Forming the Thom spaces of the universal rank k vector bundles for all k produces the Thom spectrum MO .

While Thom’s constructions are geometric in nature, the homotopical perspective on Thom spectra was given its modern ∞ -categorical form by Ando et al. [And+14]: they may be thought of as *twisted suspension spectra*. Given a rank n vector bundle $E \rightarrow X$, forming fiberwise one-point compactifications determines an X -indexed family of pointed spheres S^{E_x} , and (the underlying anima of) its Thom space is the colimit of this family. When E is a trivial bundle, this family is the constant functor $X \rightarrow \mathrm{An}_*$ with constant value S^n , and this colimit is the n -th suspension $\Sigma^n(X_+) = X_+ \wedge S^n$. There is an analogous perspective on Thom spectra as colimits of functors $X \rightarrow \mathrm{Sp}$.

The modern perspective also gives a uniform notion of orientation for vector bundles $E \rightarrow X$ with respect to a commutative ring spectrum R : one considers the associated spherical fibration $\Pi_\infty(X) \rightarrow \mathrm{Sp}, x \mapsto \Sigma^\infty S^{E_x}$, and asks for a trivialization after tensoring with R . Passing to homotopy groups, this recovers the classical Thom isomorphism.

We start the chapter by recalling the classical Thom spaces in Section 10.1. The modern perspective on Thom spaces/spectra is the topic of Section 10.2. In Section 10.3 we introduce the geometric bordism groups and record the Pontryagin–Thom theorem that shows they form the homology theory represented by the Thom spectrum. Finally, Section 10.4 treats orientations and the Thom isomorphism, following Ando et al. [And+14]. The technical proof that one-point compactification

is coherently symmetric monoidal is postponed to Section 10.5, which may be skipped on a first reading.

10.1 Thom spaces and the collapse construction

Let us make Pontryagin's construction from the introduction precise, so as to see what has to be modified in the absence of a framing. Suppose a closed n -manifold M is embedded in $\mathbb{R}^{n+k}$ and that its normal bundle has been trivialized, so that M admits a tubular neighborhood identified with $M \times \mathbb{R}^k$. Collapsing everything outside this tubular neighborhood gives a pointed map

$$S^{n+k} \longrightarrow (M \times \mathbb{R}^k)^+ \cong M_+ \wedge S^k \longrightarrow S^k,$$

and hence, after passing to stabilizations, an element of the stable stem $\pi_n(\mathbb{S}) = \pi_0 \operatorname{Hom}_{\operatorname{Sp}}(\mathbb{S}[n], \mathbb{S})$. Without a trivialization there is no map to S^k : the collapse map remembers the normal directions of M , and these vary from point to point. Thom's remedy is to replace the target S^k by a pointed space built out of the normal bundle itself, obtained by compactifying each fiber separately and then gluing all of the resulting points at infinity into a single basepoint.

Construction 10.1.1 (Fiberwise one-point compactification). Let $p: E \rightarrow X$ be a real vector bundle of rank n . We define its *fiberwise one-point compactification* S^E as the topological space obtained from E by forming the one-point compactification fiberwise, replacing $\mathbb{R}^n$ with $S^n = \mathbb{R}^n \cup \{\infty\}$ in each fiber. More precisely, given a local trivialization $\{\varphi_i: U_i \times \mathbb{R}^n \rightarrow p^{-1}(U_i)\}_{i \in I}$ of E , we define S^E by gluing together the spaces $U_i \times S^n$ along the transition functions: for $x \in U_i \cap U_j$, the map $\varphi_j^{-1} \circ \varphi_i: \{x\} \times \mathbb{R}^n \rightarrow \{x\} \times \mathbb{R}^n$ extends uniquely to a homeomorphism $\{x\} \times S^n \rightarrow \{x\} \times S^n$ fixing the point at infinity. One can show S^E is independent of the choice of local trivializations of $E \rightarrow X$. The space S^E comes equipped with a map $S^E \rightarrow X$. The 'points at infinity' assemble into a continuous section $s_\infty: X \rightarrow S^E$.

Definition 10.1.2 (Thom space). Let $p: E \rightarrow X$ be a real vector bundle. We define the *Thom space* $\operatorname{Th}(E)$ as the quotient of S^E by the points at infinity:

$$\operatorname{Th}(E) := S^E / s_\infty(X).$$

Remark 10.1.3. Given a metric on E , one obtains a homeomorphic description of $\operatorname{Th}(E)$ as the quotient $D(E)/S(E)$, where $D(E)$ and $S(E)$ denote the unit disk and sphere bundles with respect to the metric. The homeomorphism is given by sending a vector v to $v/(1 + \|v\|)$ in each fiber. This is often taken as the definition, but has the disadvantage of requiring an auxiliary choice of metric.

Remark 10.1.4. If X is compact Hausdorff, the inclusion of E induces a homeomorphism $E^+ \cong \operatorname{Th}(E)$. Indeed, a neighborhood of the collapsed section in $\operatorname{Th}(E)$ has compact complement because S^E is compact Hausdorff, so the quotient topology at the basepoint is the one-point compactification topology.

The key observation is that the Thom space of a trivialized bundle is an honest n -fold suspension:

Example 10.1.5 (Suspension as Thom space). Let $E = X \times \mathbb{R}^n$ be the trivial rank n vector bundle over X . Then $\operatorname{Th}(E) \cong X_+ \wedge S^n \cong \Sigma^n(X_+)$.

For a general rank n bundle E , the Thom space $\text{Th}(E)$ differs from $\Sigma^n(X_+)$ only in the way the fiber spheres are glued to one another. We will therefore think of $\text{Th}(E)$ as a *twisted suspension* of X_+ , twisted by E .

Example 10.1.6. Let $E \rightarrow X$ and $E' \rightarrow X'$ be vector bundles. Then their product $E \times E' \rightarrow X \times X'$ is again a vector bundle (called their *external direct sum*) and there is an isomorphism of pointed spaces

$$\text{Th}(E \times E') \cong \text{Th}(E) \wedge \text{Th}(E').$$

To see this, equip $E \times E'$ with the metric $\|(v, w)\| = \max(\|v\|, \|w\|)$. Then the disk bundle $D(E \times E')$ is homeomorphic to the product $D(E) \times D(E')$, and under this homeomorphism the sphere bundle $S(E \times E')$ corresponds to the subspace

$$D(E) \times S(E') \cup S(E) \times D(E')$$

of the product. Passing to quotient spaces then proves the claim.

Construction 10.1.7 (Functoriality of Thom spaces). Consider a fiberwise injective morphism of rank n vector bundles, i.e. a commutative diagram

$$\begin{array}{ccc} E & \xrightarrow{\tilde{f}} & E' \\ p \downarrow & & \downarrow p' \\ X & \xrightarrow{f} & X' \end{array}$$

in which $\tilde{f}$ induces injective $\mathbb{R}$ -linear maps $E_x \hookrightarrow E'_{f(x)}$ on fibers. Then the map $\tilde{f}$ induces a map $S^E \rightarrow S^{E'}$ on fiberwise one-point compactifications, which preserves the sections at infinity and hence induces a continuous map

$$\text{Th}(\tilde{f}): \text{Th}(E) \rightarrow \text{Th}(E').$$

We can now carry out the collapse construction in the generality we will need. We isolate it as a separate construction, since it is used again in the proof of Atiyah duality in Section 11.3.

Construction 10.1.8 (Pontryagin–Thom collapse map). Let $i: N \hookrightarrow N'$ be a smooth embedding of smooth manifolds, with N compact, and let $\nu(i)$ denote its normal bundle. Choose a tubular neighborhood, given by an open neighborhood $U \subseteq N'$ of N and a diffeomorphism from a neighborhood of the zero section in $\nu(i)$ onto U . Collapsing the complement of a smaller tubular neighborhood to the basepoint gives a pointed map

$$\text{PT}(i): N'_+ \longrightarrow \text{Th}(\nu(i)),$$

called the *Pontryagin–Thom collapse map*. Its homotopy class is independent of the auxiliary choices. If N' is equipped with a basepoint away from N , we may choose the tubular neighborhood of N to avoid this basepoint, giving a pointed map

$$\text{PT}(i): N' \longrightarrow \text{Th}(\nu(i)).$$

For a framed embedding, the normal bundle is trivialized and $\text{Th}(\nu(i)) \cong \Sigma^k(N_+)$, so that we recover Pontryagin’s construction. In general the target depends on $\nu(i)$. The next step is to make it independent of the manifold by passing to the universal bundle.




Figure 10.1: The Pontryagin–Thom collapse map of an embedding $N \hookrightarrow N'$, which crushes the complement of a tubular neighborhood of N to the basepoint. In the case drawn, N is a circle in a surface with trivial normal bundle. Cutting the band U along N exhibits $\text{Th}(\nu(i))$ as two cones on N glued along N , that is, as a sphere with equator N whose two cone points have both become the basepoint $*$.

Example 10.1.9. Let us consider the Grassmannian $\text{Gr}_n(\mathbb{R}^\infty)$ of n -dimensional subspaces of $\mathbb{R}^\infty$. By the real analogue of the Grassmannian classification theorem recorded in Remark 9.2.22, this space is a model for the classifying space $BO(n)$ and classifies rank n real vector bundles over paracompact Hausdorff spaces. It is a cell complex by its Schubert cell decomposition. The universal bundle γ_n over $\text{Gr}_n(\mathbb{R}^\infty)$ has as total space

$$E_n = \{(V, v) \in \text{Gr}_n(\mathbb{R}^\infty) \times \mathbb{R}^\infty \mid v \in V\}$$

with the projection map sending (V, v) to V . The fiber over a point $V \in \text{Gr}_n(\mathbb{R}^\infty)$ is precisely the n -dimensional vector space V itself. The Thom space $\text{Th}(\gamma_n)$ is the space of pairs (V, v) where $v \in V$ has length ≤ 1 , modulo those pairs where v has length exactly 1.

The Thom spaces $\text{Th}(\gamma_n)$ are particularly important: every rank n vector bundle over a paracompact Hausdorff space is pulled back from γ_n by Remark 9.2.22, so its Thom space admits a map to $\text{Th}(\gamma_n)$ via Construction 10.1.7. Composing this with the collapse map from Construction 10.1.8 thus produces maps from manifolds into these Thom spaces. They are moreover compatible as n varies, in the following sense.

Construction 10.1.10 (The prespectrum MO^{pre}). Consider the canonical inclusion

$$\text{Gr}_n(\mathbb{R}^\infty) \hookrightarrow \text{Gr}_{n+1}(\mathbb{R} \times \mathbb{R}^\infty) \cong \text{Gr}_{n+1}(\mathbb{R}^\infty), \quad V \mapsto \mathbb{R} \oplus V.$$

The pullback of γ_{n+1} along this inclusion is the product bundle $\gamma_n \times \underline{\mathbb{R}}$, and in particular we obtain a pointed continuous map

$$\Sigma(\text{Th}(\gamma_n)) \cong \text{Th}(\gamma_n \times \underline{\mathbb{R}}) \rightarrow \text{Th}(\gamma_{n+1}).$$

By adjunction, this defines a map $\sigma_n: \text{Th}(\gamma_n) \rightarrow \Omega(\text{Th}(\gamma_{n+1}))$, turning the sequence of pointed animae $(\Pi_\infty \text{Th}(\gamma_n))$ into a prespectrum, which we will denote by MO^{pre} .

Applying spectrification (Proposition 4.3.29) to MO^{pre} produces a spectrum, and this is the classical construction of MO going back to Thom [Tho54]. Instead of taking this as our definition, the next section gives a construction of the Thom spectrum which requires no auxiliary choices and applies to virtual bundles; we shall identify the two in Theorem 10.2.23.

10.2 Thom spectra

The construction of the previous section is tied to point-set models, does not exhibit the coherence of one-point compactification with respect to direct sums, and treats each bordism spectrum separately. More importantly, it applies only to honest vector bundles. The normal bundle ν of an embedding $M \hookrightarrow \mathbb{R}^{n+k}$ of a closed n -manifold is not an invariant of M : enlarging the ambient space changes it by trivial summands. What is intrinsic is the *stable normal bundle*, the virtual vector bundle

$$\nu - \underline{\mathbb{R}}^k = \underline{\mathbb{R}}^n - T_M$$

of virtual rank 0, which is the class classified by a map $M \rightarrow \mathbf{BO}$. Closely related is the virtual bundle $-T_M = \nu - \underline{\mathbb{R}}^{n+k}$ of virtual rank $-n$, which is the one occurring in Atiyah duality in Section 11.3. Neither is an honest vector bundle, and neither has a Thom space in the sense of Definition 10.1.2.

The remedy is to describe the Thom construction in a way which never refers to the total space of the bundle. Recall the picture from the introduction: a rank n vector bundle over X determines a family of n -spheres parametrized by X , and the Thom space should be the colimit of that family. Formulated this way, the input is a functor into pointed animae, and once we stabilize it becomes a functor into invertible spectra, where inverses exist and allow for an extension to virtual bundles.

10.2.1 Thom animae

We start by making precise the claim that Thom spaces are colimits of parametrized families of spheres.

Definition 10.2.1 (Spherical fibration). Let B be an anima and let $n \geq 0$. A *rank n spherical fibration* over B is a functor $\xi: B \rightarrow \mathbf{An}_*$ which lands in the full subcategory spanned by the n -sphere S^n . A *spherical fibration* is a rank n spherical fibration for some n .

Definition 10.2.2 (Thom anima). We define the *Thom anima* of a spherical fibration $\xi: B \rightarrow \mathbf{An}_*$ as its colimit:

$$\mathrm{Th}(\xi) := \mathrm{colim}(\xi: B \rightarrow \mathbf{An}_*).$$

Recall from Corollary 23.2.6 the straightening-unstraightening equivalence $\mathbf{An}/_B \simeq \mathrm{Fun}(B, \mathbf{An})$. Passing to pointed objects, this provides an equivalence

$$\mathrm{Fun}(B, \mathbf{An}_*) \simeq \mathrm{Fun}(B, \mathbf{An})_* \simeq (\mathbf{An}/_B)_*,$$

so that a rank n spherical fibration over B is the same as a pair (p, s) of a morphism $p: A \rightarrow B$ of animae and a section $s: B \rightarrow A$ of p such that each fiber A_b of p is isomorphic to the n -sphere S^n .

Example 10.2.3 (Sphere bundle). Let X be a topological space and consider an n -dimensional sphere bundle $Y \rightarrow X$ over X , i.e. a fiber bundle whose generic fiber is S^n and whose structure group is $\mathrm{Homeo}_*(S^n)$. Then the underlying map of animae $\Pi_\infty(Y) \rightarrow \Pi_\infty(X)$ is a spherical fibration: by Proposition 2.3.19 and Corollary 2.4.15 the fibers of this map are the spheres S^n .

Example 10.2.4 (Spherical fibration of vector bundle). Given a rank n vector bundle $p: E \rightarrow X$, we may apply the previous example to the fiberwise one-point compactification $S^E \rightarrow X$ from Construction 10.1.1. We denote the resulting spherical fibration by

$$\xi_E: \Pi_\infty(X) \rightarrow \mathbf{An}_*.$$

Informally, ξ_E is the functor which sends a point $x \in X$ to $\Pi_\infty(S^{E_x})$, where S^{E_x} is the one-point compactification of the fiber E_x of p over x .

We now compare the geometric Thom space with the homotopical Thom anima:

Proposition 10.2.5. *Let $p: E \rightarrow X$ be a vector bundle over a paracompact Hausdorff space of the homotopy type of a cell complex. Then the underlying pointed anima of its Thom space is isomorphic to the Thom anima of ξ_E :*

$$\Pi_\infty(\mathrm{Th}(E)) \cong \mathrm{Th}(\xi_E).$$

Proof. We first identify the colimit of any pointed family with the cofiber of its section. We then compare this homotopy cofiber with the point-set quotient defining the classical Thom space.

Set $B := \Pi_\infty(X)$. Under the straightening-unstraightening equivalence $\mathrm{Fun}(B, \mathrm{An}_*) \xrightarrow{\sim} (\mathrm{An}/B)_*$, the spherical fibration ξ_E corresponds to the pointed object of An/B given by

$$\Pi_\infty(X) \xrightarrow{\Pi_\infty(s_\infty)} \Pi_\infty(S^E) \longrightarrow \Pi_\infty(X).$$

More generally, suppose that a functor $\xi: B \rightarrow \mathrm{An}_*$ corresponds to a pointed object $B \xrightarrow{s} A \rightarrow B$ of An/B . By Proposition 23.5.2, the colimit of the underlying functor $B \rightarrow \mathrm{An}$ is A , while the colimit of the constant basepoint diagram is B . Since colimits in pointed objects are obtained by forming a pushout over the basepoint, it follows that

$$\mathrm{colim}_B \xi \cong \mathrm{cofib}(s: B \rightarrow A).$$

Applied to ξ_E , this gives an isomorphism

$$\mathrm{Th}(\xi_E) \cong \mathrm{cofib}(\Pi_\infty(X) \xrightarrow{\Pi_\infty(s_\infty)} \Pi_\infty(S^E)).$$

It remains to identify this cofiber with the classical Thom space. Choose a fiberwise metric on E , which exists by paracompactness. It identifies the fiberwise one-point compactification S^E with the unit sphere bundle of $E \oplus \mathbb{R}$. Under this identification, the section s_∞ is the section given by the north pole in each fiber. The pair (S^n, ∞) admits an NDR structure (cf. Corollary 2.3.11), which may be chosen $O(n)$ -equivariantly and therefore induces an NDR structure on $(S^E, s_\infty(X))$. Moreover, S^E has the homotopy type of a cell complex, since it is the total space of a fibration over a space of the homotopy type of a cell complex with fiber S^n ; see Milnor [Mil59, Theorem 3]. Thus X and S^E satisfy the hypotheses of Corollary 2.3.11, giving the isomorphism

$$\mathrm{cofib}(\Pi_\infty(X) \xrightarrow{\Pi_\infty(s_\infty)} \Pi_\infty(S^E)) \simeq \Pi_\infty(S^E/s_\infty(X)) = \Pi_\infty(\mathrm{Th}(E)),$$

as desired. □

10.2.2 Thom spectra

The homotopical perspective of Thom spaces directly generalizes to the stable setting.

Definition 10.2.6 (Stable spherical fibration). Let $\mathrm{Pic}(\mathbb{S}) \subseteq \mathrm{Sp}^\sim$ denote the full subcategory spanned by shifts $\mathbb{S}[n]$ of the sphere spectrum. For an anima A , we define a *stable spherical fibration* on A to be a functor $\xi: A \rightarrow \mathrm{Sp}$ which lands in $\mathrm{Pic}(\mathbb{S})$. For an integer $n \in \mathbb{Z}$ we say that ξ has *rank* n if it in fact lands in the full subanima of $\mathrm{Pic}(\mathbb{S})$ spanned by $\mathbb{S}[n]$.

One can show that objects of $\text{Pic}(\mathbb{S})$ are precisely the *invertible spectra*: those spectra X for which there is a spectrum X^{-1} and an isomorphism $X \otimes X^{-1} \cong \mathbb{S}$; see for example [HPS97]. In particular, $\text{Pic}(\mathbb{S})$ is precisely the Picard anima of Sp in the sense of Definition 11.1.4 below.

If $\xi: A \rightarrow \text{An}_*$ is a rank n spherical fibration, then $\Sigma^\infty \xi$ is a rank n stable spherical fibration. In particular, a rank n vector bundle $E \rightarrow X$ determines the stable spherical fibration $\Sigma^\infty \xi_E$ on $\Pi_\infty(X)$.

Definition 10.2.7 (Thom spectrum). Given a stable spherical fibration $\xi: A \rightarrow \text{Sp}$, we define its *Thom spectrum* as its colimit:

$$\text{th}(\xi) := \text{colim}(\xi: A \rightarrow \text{Sp}).$$

For a vector bundle $E \rightarrow X$, we write

$$\text{th}(E) := \Sigma^\infty \text{Th}(\xi_E)$$

for its Thom spectrum. Since the suspension spectrum functor $\Sigma^\infty(-): \text{An}_* \rightarrow \text{Sp}$ preserves colimits, this agrees with $\text{th}(\Sigma^\infty \xi_E)$, and by Proposition 10.2.5 it is the suspension spectrum of the underlying pointed anima of the classical Thom space. Throughout, Th denotes a Thom space or Thom anima and th a Thom spectrum.

Construction 10.2.8 (Multiplicative structures on Thom spectra). Let A be a small commutative monoid in animae , regarded as a symmetric monoidal ∞ -category, and let

$$\xi: A \longrightarrow \text{Pic}(\mathbb{S}) \subseteq \text{Sp}$$

be a symmetric monoidal functor. We equip the Thom spectrum $\text{th}(\xi)$ with a canonical commutative ring spectrum structure, following Antolín-Camarena and Barthel [AB19].

By Corollary 16.2.8, the functor category $\text{Fun}(A, \text{Sp})$ admits the Day convolution symmetric monoidal structure. The universal property of Day convolution from Lemma 16.1.4 identifies the underlying lax symmetric monoidal functor of ξ with a commutative algebra in $\text{Fun}(A, \text{Sp})$. The colimit functor

$$\text{colim}_A: \text{Fun}(A, \text{Sp}) \longrightarrow \text{Sp}$$

is pointwise left Kan extension along the unique symmetric monoidal functor $A \rightarrow *$, so it is symmetric monoidal by Lemma 16.2.11. Here $\text{Fun}(*, \text{Sp})$ is identified with Sp equipped with its usual tensor product. It therefore carries ξ to a commutative algebra in spectra whose underlying spectrum is

$$\text{colim}_A \xi = \text{th}(\xi).$$

Remark 10.2.9. There is an analogous multiplicative Thom-spectrum construction with *Comm* replaced by another ∞ -operad, once the corresponding operadic Day convolution has been developed. We do not need this generalization here. For the case of the little-cubes operads $\mathcal{E}_n$, see Antolín-Camarena and Barthel [AB19].

The symmetric monoidal stable spherical fibrations used below arise from the stable J-homomorphism. We apply the construction to bordism spectra in Construction 10.2.25.

10.2.3 The J-homomorphism

As previously indicated, passing to spectra allows us to extend the definition of Thom spectra to virtual vector bundles. The idea is simple: the assignment $E \mapsto \Sigma^\infty \xi_E$ is symmetric monoidal and lands in invertible spectra, so it should extend over group completion. The subtlety is coherence: the familiar homeomorphisms $S^{V \oplus W} \cong S^V \wedge S^W$ must assemble into a symmetric monoidal functor from vector spaces to pointed anima.

For a topological space X , let $\text{Vect}_{\mathbb{R}}^{\text{disc}}(X)$ denote the 1-category of finite-rank real vector bundles over X and morphisms of vector bundles. By Corollary 19.7.10, direct sum and pullback give a simplicial diagram

$$[q] \mapsto \text{Vect}_{\mathbb{R}}^{\text{disc}}(X \times |\Delta^q|)^{\simeq} \in \text{CMon}(\text{An}).$$

Definition 10.2.10. We define the *moduli anima of real vector bundles over X* by

$$\text{Vect}_{\mathbb{R}}(X)^{\simeq} := \text{colim}_{[q] \in \Delta^{\text{op}}} \text{Vect}_{\mathbb{R}}^{\text{disc}}(X \times |\Delta^q|)^{\simeq} \in \text{CMon}(\text{An}).$$

This colimit is computed on underlying anima and is functorial in X , giving a functor $\text{Vect}_{\mathbb{R}}(-)^{\simeq} : \text{Top}^{\text{op}} \rightarrow \text{CMon}(\text{An})$. When $X = \text{pt}$, we refer to $\text{Vect}_{\mathbb{R}}(\text{pt})^{\simeq}$ as the *moduli anima of finite-dimensional real vector spaces*.

The following proposition supplies all the coherence needed below. Its proof occupies Section 10.5.

Proposition 10.2.11 (One-point compactification). *For every topological space X , fiberwise one-point compactification defines a symmetric monoidal functor*

$$S_X^{(-)} : \text{Vect}_{\mathbb{R}}(X)^{\simeq} \longrightarrow (\text{An}/\Pi_{\infty}(X))_* \simeq \text{Fun}(\Pi_{\infty}(X), \text{An}_*).$$

These functors are the components of a natural transformation

$$S^{(-)} : \text{Vect}_{\mathbb{R}}(-)^{\simeq} \Longrightarrow (\text{An}/\Pi_{\infty}(-))_*$$

between functors $\text{Top}^{\text{op}} \rightarrow \text{CMon}(\text{Cat}_{\infty})$, where the source is regarded as taking values in symmetric monoidal ∞ -groupoids. For $X = \text{pt}$, there is an isomorphism

$$\text{Vect}_{\mathbb{R}}(\text{pt})^{\simeq} \cong \bigsqcup_{n \geq 0} \mathbf{BO}(n)$$

of commutative monoids, where addition is induced by the block-sum homomorphisms $O(k) \times O(l) \rightarrow O(k+l)$. The restriction of $S_{\text{pt}}^{(-)}$ to the rank n component is classified by the standard pointed action of $O(n)$ on $S^{\mathbb{R}^n}$.

We also write $S^{(-)}$ for the component $S_{\text{pt}}^{(-)}$ when the base is a point.

Definition 10.2.12. Write $O := \text{colim}_n O(n)$ for the stable orthogonal group and $\mathbf{BO}$ for its classifying anima. We define

$$\mathbf{BOP} := (\text{Vect}_{\mathbb{R}}(\text{pt})^{\simeq})^{\text{grp}} \in \text{CGrp}(\text{An})$$

to be the group completion of the commutative monoid of finite-dimensional real vector spaces. The telescope calculation of Corollary 9.4.3, with real vector spaces in place of complex ones, identifies its underlying anima as

$$\mathbf{BOP} \simeq \mathbb{Z} \times \mathbf{BO}.$$

The only additional point in applying Theorem 5.5.2 is that $\pi_1(\mathbf{BO}) \cong \pi_0(O) \cong C_2$ is abelian. The virtual-rank-zero component has the presentation

$$\mathbf{BO} \cong \operatorname{colim}_n \mathbf{BO}(n),$$

where the transition maps add a trivial line. Under this presentation, the map from the n -th stage to the virtual-rank-zero component sends V to $V - \mathbb{R}^n$.

Construction 10.2.13 (Stable J-homomorphism). Applying reduced suspension spectra to Proposition 10.2.11 gives a symmetric monoidal functor

$$\operatorname{Vect}_{\mathbb{R}}(\operatorname{pt}) \xrightarrow{\simeq} \operatorname{An}_* \xrightarrow{\Sigma^\infty} \operatorname{Sp}.$$

Its image lies in $\operatorname{Pic}(\mathbb{S})$, since $\Sigma^\infty S^V \cong \mathbb{S}[\dim(V)]$ is invertible. Since tensor product of spectra turns $\operatorname{Pic}(\mathbb{S})$ into a commutative group in animae , the resulting functor to $\operatorname{Pic}(\mathbb{S})$ extends uniquely over group completion, giving a symmetric monoidal functor

$$J: \mathbf{BOP} \longrightarrow \operatorname{Pic}(\mathbb{S}).$$

We call J the *stable J-homomorphism*. We use the same notation for its restriction to any component of $\mathbf{BOP}$, in particular for

$$J: \mathbf{BO} \simeq \{0\} \times \mathbf{BO} \longrightarrow \operatorname{Pic}(\mathbb{S}).$$

Remark 10.2.14 (Relation to the classical J-homomorphism). The connected component of $\mathbb{S}$ in $\operatorname{Pic}(\mathbb{S})$ is equivalent to $\mathbf{BGL}_1(\mathbb{S})$, where $\operatorname{GL}_1(\mathbb{S}) := \operatorname{Aut}_{\operatorname{Sp}}(\mathbb{S})$ is the automorphism group of the sphere spectrum. The restriction of J to the virtual-rank-zero component lands in this connected component. Passing to loops gives a group homomorphism $O \rightarrow \operatorname{GL}_1(\mathbb{S})$ in An , whose effect on positive homotopy groups is the classical stable J-homomorphism $\pi_n(O) \rightarrow \pi_n(\mathbb{S})$. We refer to Ravenel [Rav86, Chapter 1] for a classical discussion of this map.

The complex analogue is the composite $\mathbf{BUP} \rightarrow \mathbf{BOP} \xrightarrow{J} \operatorname{Pic}(\mathbb{S})$ induced by forgetting the complex structure.

The point of constructing J on the full group completion is that it assigns stable spherical fibrations, and hence Thom spectra, to virtual vector bundles of arbitrary rank.

Definition 10.2.15 (Virtual vector bundle). Let A be an anima. A *virtual vector bundle* on A is a map of animae

$$V: A \longrightarrow \mathbf{BOP}.$$

Its *rank* is the composite $\operatorname{rk}(V): A \rightarrow \mathbb{Z}$ with the projection to $\mathbb{Z}$, and we say that V has rank $d \in \mathbb{Z}$ if this composite is constant with value d . A virtual vector bundle on a topological space X means one on $\Pi_\infty(X)$.

Since $\mathbf{BOP}$ is a commutative group in animae , so is the anima $\operatorname{Hom}_{\operatorname{An}}(A, \mathbf{BOP})$ of virtual vector bundles on A . We write $V + W$ for the sum and $-V$ for the inverse; ranks add, and $\operatorname{rk}(-V) = -\operatorname{rk}(V)$.

Construction 10.2.16 (Thom spectrum of a virtual vector bundle). Let V be a virtual vector bundle on an anima A . Composing with the J-homomorphism gives a stable spherical fibration

$$\xi_V := J \circ V: A \longrightarrow \operatorname{Pic}(\mathbb{S}).$$

If V has rank d , then so does ξ_V , since J carries the class of $\mathbb{R}$ to $\mathbb{S}[1]$. We define the *Thom spectrum* of V as

$$\mathrm{th}(V) := \mathrm{th}(\xi_V) = \mathrm{colim} (J \circ V: A \rightarrow \mathrm{Sp}).$$

Remark 10.2.17. As J is a homomorphism of commutative groups in $\mathbf{animae}$, the assignment $V \mapsto \xi_V$ carries sums to pointwise tensor products and inverses to pointwise inverses:

$$\xi_{V+W} \cong \xi_V \otimes \xi_W \quad \text{and} \quad \xi_{-V} \cong \xi_V^{-1}.$$

The Thom spectrum itself is of course not additive: $\mathrm{th}(V + W)$ is a colimit over A , not a tensor product of colimits.

We next relate this construction to honest vector bundles. A pointed action of a topological group G on a topological space Y , together with a principal G -bundle $P \rightarrow X$, gives an associated pointed bundle $P \times_G Y \rightarrow X$. Passing to fundamental $\mathbf{animae}$ and applying pointed straightening produces a functor $\Pi_\infty(X) \rightarrow \mathbf{An}_*$. For a universal principal bundle, we refer to the resulting functor $\mathbf{BG} \rightarrow \mathbf{An}_*$ as the functor classified by the pointed G -action on Y .

Construction 10.2.18 (One-point compactification of vector spaces). By Proposition 10.2.11, the rank n component of $\mathrm{Vect}_{\mathbb{R}}(\mathrm{pt})^\simeq$ is isomorphic to $\mathbf{BO}(n)$. We denote the restriction of one-point compactification to this component by

$$J_n^{\mathrm{un}}: \mathbf{BO}(n) \longrightarrow \mathbf{An}_*.$$

It is the functor classified by the standard pointed action of $O(n)$ on $S^{\mathbb{R}^n}$. Consequently, the restriction of J along the rank n component $\mathbf{BO}(n) \rightarrow \mathbf{BOP}$ is $\Sigma^\infty J_n^{\mathrm{un}}$. Its restriction along the stable map

$$\mathbf{BO}(n) \longrightarrow \{0\} \times \mathbf{BO} \subseteq \mathbf{BOP}, \quad V \longmapsto V - \mathbb{R}^n,$$

is $\Sigma^\infty J_n^{\mathrm{un}}[-n]$.

Lemma 10.2.19. *Let X be a paracompact Hausdorff space and let $f: X \rightarrow \mathrm{Gr}_n(\mathbb{R}^\infty)$ be a continuous map. Let $p: E \rightarrow X$ be the rank n vector bundle classified by f . Then the map $\xi_E: \Pi_\infty(X) \rightarrow \mathbf{An}_*$ is given by the composite*

$$\Pi_\infty(X) \xrightarrow{\Pi_\infty(f)} \Pi_\infty(\mathrm{Gr}_n(\mathbb{R}^\infty)) \xrightarrow{\sim} \mathbf{BO}(n) \xrightarrow{J_n^{\mathrm{un}}} \mathbf{An}_*,$$

where the middle isomorphism is the one from Proposition 9.2.11.

Proof. By definition, $p: E \rightarrow X$ is the pullback of the universal bundle $p_{\mathrm{univ}}: E_{\mathrm{univ}} \rightarrow \mathrm{Gr}_n(\mathbb{R}^\infty)$. The induced square of sphere bundles is carried to a pullback of $\mathbf{animae}$ by Proposition 2.3.19. Under straightening-unstraightening, pullback corresponds to precomposition, so it suffices to prove the claim for p_{univ} . The Stiefel manifold $V_n(\mathbb{R}^\infty)$ of orthonormal n -frames is contractible, and its projection to $\mathrm{Gr}_n(\mathbb{R}^\infty)$ is a principal $O(n)$ -bundle. It therefore identifies $\Pi_\infty(\mathrm{Gr}_n(\mathbb{R}^\infty))$ with $\mathbf{BO}(n)$ by Proposition 9.2.11. Under this identification, the fiberwise one-point compactification of the universal bundle is the associated pointed sphere bundle

$$V_n(\mathbb{R}^\infty) \times_{O(n)} S^n \longrightarrow \mathrm{Gr}_n(\mathbb{R}^\infty).$$

Its straightening is precisely J_n^{un} . □

Example 10.2.20 (The class of a vector bundle). Let X be a paracompact Hausdorff space and let $E \rightarrow X$ be a rank n vector bundle, classified by a map $f: X \rightarrow \mathrm{Gr}_n(\mathbb{R}^\infty)$. Combining the identification $\Pi_\infty(\mathrm{Gr}_n(\mathbb{R}^\infty)) \xrightarrow{\sim} \mathbf{BO}(n)$ of Proposition 9.2.11 with the preceding construction, we obtain a virtual vector bundle

$$[E]: \Pi_\infty(X) \xrightarrow{\Pi_\infty(f)} \mathbf{BO}(n) \subseteq \mathrm{Vect}_{\mathbb{R}}(\mathrm{pt})^\simeq \longrightarrow \mathrm{BOP}$$

of rank n . For the trivial bundle $\underline{\mathbb{R}}^N$ this is the constant map with value $\mathbb{R}^N$.

This construction is compatible with direct sums. Indeed, under the identifications $\Pi_\infty(\mathrm{Gr}_n(\mathbb{R}^\infty)) \cong \mathbf{BO}(n)$, the Whitney-sum map corresponds to the map induced by the block-sum homomorphism $O(n) \times O(m) \rightarrow O(n+m)$. To see this, observe that the orthonormal frame bundle of $E \oplus E'$ is obtained from the product over X of the orthonormal frame bundles of E and E' by extension of structure group along this homomorphism. Since the addition on $\mathrm{Vect}_{\mathbb{R}}(\mathrm{pt})^\simeq$ is induced by block sum, it follows that

$$[E \oplus E'] \cong [E] + [E'].$$

Finally, we record the compatibility with the construction for honest vector bundles, and the interaction between adding trivial summands and shifts; we will use these properties in our proof of Atiyah duality in Section 11.3.

Lemma 10.2.21. *Let X be a paracompact Hausdorff space of the homotopy type of a cell complex and let $E \rightarrow X$ be a rank n vector bundle. Then there is an isomorphism $\xi_{[E]} \cong \Sigma^\infty(\xi_E)$ of stable spherical fibrations, and hence an isomorphism $\mathrm{th}([E]) \cong \mathrm{th}(E)$. More generally, for every $N \geq 0$ we have $\xi_{[E] - [\underline{\mathbb{R}}^N]} \cong \Sigma^\infty(\xi_E)[-N]$ and therefore*

$$\mathrm{th}([E] - [\underline{\mathbb{R}}^N]) \cong \mathrm{th}(E)[-N] \cong \Sigma^{\infty-N} \mathrm{Th}(E).$$

Proof. By Construction 10.2.18, the restriction of J to the rank n component is $\Sigma^\infty J_n^{\mathrm{un}}$. The first isomorphism is therefore precisely Lemma 10.2.19, and passing to colimits gives $\mathrm{th}([E]) \cong \Sigma^\infty \mathrm{Th}(\xi_E) = \mathrm{th}(E)$, since Σ^∞ preserves colimits.

The virtual vector bundle $[\underline{\mathbb{R}}^N]$ is constant with value $\mathbb{R}^N$, so $\xi_{[\underline{\mathbb{R}}^N]}$ is constant with value $\Sigma^\infty S^{\mathbb{R}^N} \cong \mathbb{S}[N]$. By Remark 10.2.17 we thus obtain

$$\xi_{[E] - [\underline{\mathbb{R}}^N]} \cong \Sigma^\infty(\xi_E) \otimes \mathbb{S}[-N] \cong \Sigma^\infty(\xi_E)[-N],$$

and the remaining claims follow because shifts commute with colimits. $\square$

10.2.4 Bordism spectra

We now apply the Thom-spectrum construction to bordism spectra. We start with MO , which comes from the virtual-rank-zero component of BOP , and relate it to the prespectrum $\mathrm{MO}^{\mathrm{pre}}$ constructed in Section 10.1.

Definition 10.2.22 (Unoriented bordism spectrum). We define the *unoriented bordism spectrum* MO as the Thom spectrum of the restriction of J to the virtual-rank-zero component:

$$\mathrm{MO} := \mathrm{th}(J|_{\mathbf{BO}}) = \mathrm{colim}(J: \mathbf{BO} \rightarrow \mathrm{Sp}).$$

Theorem 10.2.23. *The spectrum associated to the prespectrum $\mathrm{MO}^{\mathrm{pre}}$ of Construction 10.1.10 via Proposition 4.3.29 is MO :*

$$(\mathrm{MO}^{\mathrm{pre}})^{\mathrm{sp}} \cong \mathrm{MO}.$$

Proof. We first identify the levels and structure maps of the classical prespectrum with the unstable Thom construction. We then pass to the colimit presentation of the virtual-rank-zero component of BOP.

Step 1: Consider the functor $\xi_{E_n} : \Pi_\infty(\mathrm{Gr}_n(\mathbb{R}^\infty)) \rightarrow \mathrm{An}_*$ associated to γ_n . Applying Lemma 10.2.19 to the identity of $\mathrm{Gr}_n(\mathbb{R}^\infty)$ identifies ξ_{E_n} with J_n^{un} . In particular, Proposition 10.2.5 provides an isomorphism

$$\mathrm{MO}_n^{\mathrm{pre}} = \Pi_\infty(\mathrm{Th}(\gamma_n)) \cong \mathrm{Th}(J_n^{\mathrm{un}})$$

of animae. The identification $E_{n+1}|_{\mathrm{Gr}_n(\mathbb{R}^\infty)} \cong E_n \times \mathbb{R}$ corresponds to the monoidal identification

$$J_{n+1}^{\mathrm{un}}|_{\mathbf{BO}(n)} \cong \Sigma J_n^{\mathrm{un}}.$$

Thus the structure maps of $\mathrm{MO}^{\mathrm{pre}}$ correspond to the canonical comparison maps

$$\Sigma \mathrm{Th}(J_n^{\mathrm{un}}) \cong \mathrm{Th}(J_{n+1}^{\mathrm{un}}|_{\mathbf{BO}(n)}) \longrightarrow \mathrm{Th}(J_{n+1}^{\mathrm{un}}).$$

Step 2: Applying Theorem 21.2.11 to the presentation $\mathbf{BO} \simeq \mathrm{colim}_{n \in \mathbb{N}} \mathbf{BO}(n)$ of Definition 10.2.12 gives

$$\mathrm{MO} \cong \mathrm{colim}_n \Sigma^\infty \mathrm{Th}(J_n^{\mathrm{un}})[-n].$$

The structure maps in this colimit diagram are precisely those of $\mathrm{MO}^{\mathrm{pre}}$, so the right-hand side is $(\mathrm{MO}^{\mathrm{pre}})^{\mathrm{sp}}$. $\square$

The abstract definition also makes the variants of MO formal.

Definition 10.2.24 (The Thom spectrum MB). Let $\varphi : B \rightarrow \mathrm{BOP}$ be a map of animae. We define its associated Thom spectrum MB by

$$MB := \mathrm{colim} (B \xrightarrow{\varphi} \mathrm{BOP} \xrightarrow{J} \mathrm{Sp}).$$

Equivalently, φ is a virtual vector bundle on B , and $MB = \mathrm{th}(\varphi)$ in the notation of Construction 10.2.16. The notation suppresses φ , which will always be clear from the context. We allow B to be an arbitrary anima. For the Pontryagin–Thom theorem of Section 10.3, we will specialize to maps which land in the virtual-rank-zero component $\mathbf{BO} \subseteq \mathrm{BOP}$.

Construction 10.2.25 (Multiplicative structures on bordism spectra). Let B be a commutative monoid in animae and let

$$\varphi : B \longrightarrow \mathrm{BOP}$$

be a morphism of commutative monoids. Since the J -homomorphism is symmetric monoidal, so is the composite

$$B \xrightarrow{\varphi} \mathrm{BOP} \xrightarrow{J} \mathrm{Pic}(\mathbb{S}) \subseteq \mathrm{Sp}.$$

Consequently, Construction 10.2.8 equips MB with a canonical commutative ring spectrum structure.

Example 10.2.26. A continuous group homomorphism $G \rightarrow O$ induces a map $\mathbf{B}G \rightarrow \mathbf{B}O$, and we write MG for the resulting Thom spectrum. Taking $B = \mathbf{B}O$ with φ the identity recovers $\mathbf{M}O$. Further examples are

$$\mathbf{M}SO, \quad \mathbf{M}U, \quad \mathbf{M}SU, \quad \mathbf{M}Sp, \quad \text{and} \quad \mathbf{M}Spin.$$

Here $\mathbf{M}U$ is the *complex bordism spectrum*. For $\mathbf{M}O$ and $\mathbf{M}U$, the defining maps to $\mathbf{B}OP$ are restrictions of morphisms of commutative groups constructed above. Hence Construction 10.2.25 equips these Thom spectra with commutative ring structures. The other classical Thom spectra in the display also admit commutative ring structures, but their construction requires compatible infinite-loop refinements of the maps $\mathbf{B}G \rightarrow \mathbf{B}O$, which we do not develop here.

The full group completions give the *periodic unoriented and complex bordism spectra*

$$\mathbf{M}OP := \text{th} (J: \mathbf{B}OP \rightarrow \mathbf{Sp}) \quad \text{and} \quad \mathbf{M}UP := \text{th} (\mathbf{B}UP \longrightarrow \mathbf{B}OP \xrightarrow{J} \mathbf{Sp}),$$

respectively. Thus $\mathbf{M}OP$ is obtained from the identity of $\mathbf{B}OP$, while $\mathbf{M}UP$ is obtained from the forgetful map $\mathbf{B}UP \rightarrow \mathbf{B}OP$. Restricting these defining diagrams to their virtual-rank-zero components recovers $\mathbf{M}O$ and $\mathbf{M}U$. Translation by the classes of $\mathbb{R}$ and $\mathbb{C}$ permutes the components of $\mathbf{B}OP$ and $\mathbf{B}UP$ while shifting the associated stable spherical fibrations by 1 and 2, respectively. It follows that there are isomorphisms of spectra

$$\mathbf{M}OP \cong \mathbf{M}OP[1] \quad \text{and} \quad \mathbf{M}UP \cong \mathbf{M}UP[2],$$

which explains the terminology.

Exercise 10.2.27. Construct the stabilization-compatible homomorphisms defining the five nonperiodic Thom spectra in the first display above.

10.3 Bordism and the Pontryagin–Thom theorem

We now return to the geometric problem which motivated the whole construction: the classification of manifolds up to bordism. Given a map $B \rightarrow \mathbf{B}O$ and a topological space X , we may consider bordism classes of manifolds equipped with a normal B -structure and a map to X . The Pontryagin–Thom theorem identifies these groups with the homology theory represented by MB . We state this theorem as a black box and only describe the collapse map which gives the comparison.

We fix a map $\varphi: B \rightarrow \mathbf{B}O$ throughout.

Definition 10.3.1 (Normal B -structure). Let M be a compact smooth n -manifold. Its *stable normal bundle* $\underline{\mathbb{R}}^n - T_M$ is a virtual vector bundle of rank 0, with classifying map $\nu_M: M \rightarrow \mathbf{B}O$. A *normal B -structure* on M is a lift of this map along φ :

$$\begin{array}{ccc} & & B \\ & \nearrow & \downarrow \varphi \\ M & \xrightarrow{\nu_M} & \mathbf{B}O. \end{array}$$

Concretely, this consists of a map $\sigma: M \rightarrow B$ together with a homotopy $\varphi \circ \sigma \simeq \nu_M$. If M has boundary, then the stable normal bundle of ∂M is identified with the restriction of the stable

normal bundle of M using the outward normal direction. Consequently, a normal B -structure on M restricts to one on ∂M . A diffeomorphism of manifolds with normal B -structures is understood to be equipped with an identification in the anima of such lifts.

Example 10.3.2. For $B = \mathbf{BO}$ with φ the identity, a normal B -structure is no data at all. For $B = \mathbf{BSO}$ it is an orientation of the stable normal bundle, and for $B = \mathbf{pt}$ it is a trivialization of it, that is, a stable framing of M .

Definition 10.3.3 (B -bordism). Let X be a topological space and let $n \geq 0$. A B -bordism cycle of degree n over X is a triple (M, f, σ) consisting of a closed smooth n -manifold M , a continuous map $f: M \rightarrow X$, and a normal B -structure σ on M . Two cycles (M_0, f_0, σ_0) and (M_1, f_1, σ_1) are *bordant* if there exist compact smooth $(n+1)$ -manifolds W_0 and W_1 with normal B -structures, together with maps $F_i: W_i \rightarrow X$, and a diffeomorphism

$$M_0 \amalg \partial W_0 \cong M_1 \amalg \partial W_1$$

of manifolds with normal B -structures over X .

We write $\Omega_n^B(X)$ for the set of bordism classes. Disjoint union defines an addition, with the empty manifold as zero. For $n < 0$, we set $\Omega_n^B(X) = 0$.

Remark 10.3.4. We used the formulation of Switzer [Swi75, Chapter 12]. Note it is an equivalence relation: reflexivity takes $W_0 = W_1 = \emptyset$, symmetry is built into the statement, and transitivity follows by taking disjoint unions. It is also visibly additive, so $\Omega_n^B(X)$ is a commutative monoid in which every class of the form $[\partial W]$ vanishes.

Inverses are obtained via the cylinder construction. Pulling back σ along the projection equips $M \times [0, 1]$ with a normal B -structure whose restriction to one end is σ ; write σ' for the structure induced on the second end. Then

$$(M, \sigma) \amalg (M, \sigma') \cong \partial(M \times [0, 1])$$

as manifolds with normal B -structures, and hence

$$[(M, f, \sigma)] + [(M, f, \sigma')] = 0$$

in $\Omega_n^B(X)$. Thus $\Omega_n^B(X)$ is an abelian group. For $B = \mathbf{BO}$ there is no additional structure, so $\sigma' = \sigma$ and every class is its own inverse; for $B = \mathbf{BSO}$ the structure σ' is the reversed orientation, and the groups are not 2-torsion in general.

A map $g: X \rightarrow Y$ induces a homomorphism $g_*: \Omega_n^B(X) \rightarrow \Omega_n^B(Y)$ by postcomposition. The comparison with MB is given by the collapse construction.

Construction 10.3.5 (Pontryagin–Thom transformation). Let (M, f, σ) be a B -bordism cycle of degree n over X . The closed smooth manifold M has the homotopy type of a finite cell complex by [Mil63, Theorem 3.5 and Corollary 6.7], so the Thom-spectrum constructions above apply to its vector bundles. Choose a smooth embedding $i: M \hookrightarrow \mathbb{R}^{n+k}$ for some k , compactify the ambient space to S^{n+k} with basepoint at infinity, and let ν be the rank k normal bundle. Applying Σ^∞ to the pointed collapse map of Construction 10.1.8 and shifting gives a map of spectra

$$\mathbb{S}[n] \longrightarrow \mathrm{th}(\nu)[-k].$$

By Lemma 10.2.21, the target is the Thom spectrum of the stable normal bundle. The map $(\sigma, f): M \rightarrow B \times X$ therefore induces a map

$$\mathrm{th}(\nu)[-k] \longrightarrow MB \otimes \mathbb{S}[X].$$

The composite determines an element of $\pi_n(MB \otimes \mathbb{S}[X])$. The standard arguments proving independence of the embedding and bordism invariance are included in the Pontryagin–Thom theorem quoted below. They show that the construction defines a natural homomorphism

$$\mathrm{PT}: \Omega_n^B(X) \longrightarrow \pi_n(MB \otimes \mathbb{S}[X]).$$

Theorem 10.3.6 (Pontryagin–Thom theorem). *For every cell complex X and every $n \in \mathbb{Z}$, the Pontryagin–Thom transformation is a natural isomorphism*

$$\Omega_n^B(X) \xrightarrow{\cong} \pi_n(MB \otimes \mathbb{S}[X]).$$

In particular, the groups $\Omega_n^B(X)$ are the absolute groups of the generalized homology theory represented by MB .

For $n < 0$, both sides vanish: the left-hand side does so by definition, while $MB \otimes \mathbb{S}[X]$ is connective because $J \circ \varphi$ takes values in connective spectra and connective spectra are closed under colimits and tensor products. This is the generalized Pontryagin–Thom theorem for normal B -structures. We use it as a black box; see Stong [Sto68, Chapter II] and Switzer [Swi75, Theorem 12.30 and Proposition 12.35]. These sources formulate the theorem using a stabilization-compatible sequence $B_n \rightarrow BO(n)$. A map $B \rightarrow \mathbf{BO}$ determines such a sequence by pullback along the maps $\mathbf{BO}(n) \rightarrow \mathbf{BO}$.

Remark 10.3.7. The spectrum MB also represents a cohomology theory $MB^*(X)$, traditionally called *cobordism*. We do not develop a geometric cocycle model for it; see Atiyah [Ati61a] for the relation with bordism.

Remark 10.3.8 (Coefficient rings). The Pontryagin–Thom theorem turns computations of bordism groups into computations of homotopy groups of spectra. For MO and MU , Construction 10.2.25 constructs the multiplication on the Thom spectrum from Whitney sum. Under the Pontryagin–Thom isomorphism, this multiplication corresponds to cartesian product of bordism classes. We will not prove this geometric compatibility here.

Thom computed the unoriented coefficient ring as

$$\pi_*(\mathrm{MO}) \cong \mathbb{F}_2[a_i \mid i \geq 1, i \neq 2^j - 1], \quad |a_i| = i;$$

see Thom [Tho54]. The complex bordism spectrum has coefficient ring

$$\pi_*(\mathrm{MU}) \cong \mathbb{Z}[x_1, x_2, \dots], \quad |x_i| = 2i,$$

a theorem of Milnor and Novikov; see Ravenel [Rav86]. By Theorem 10.3.6 this is the ring of complex bordism classes. The spectrum MU is central to chromatic homotopy theory, where its additional structure and the algebra of its cooperations organize stable homotopy according to height. At the other extreme, take $B = \mathrm{pt}$ with the constant map to $\mathbf{BO}$, so that $MB = \mathbb{S}$. By Example 10.3.2 the cycles are then stably framed manifolds, and the Pontryagin–Thom theorem becomes

$$\Omega_n^{\mathrm{fr}} := \Omega_n^{\mathrm{pt}}(\mathrm{pt}) \cong \pi_n(\mathbb{S}),$$

recovering Pontryagin’s construction from the beginning of Section 10.1.

10.4 Orientations and the Thom isomorphism

Throughout this section, we fix a commutative ring spectrum $R \in \mathbf{CAlg}(\mathbf{Sp})$.

A virtual vector bundle V of rank d on an anima A determines a stable spherical fibration $\xi_V: A \rightarrow \mathbf{Pic}(\mathbb{S})$. After shifting by $-d$ and tensoring with R , its fibers are R -modules isomorphic to R . An orientation is a coherent choice of such isomorphisms. This formulation applies directly to virtual bundles and specializes to the classical notion of a Thom class for an honest bundle. It follows the orientation theory of Ando et al. [And+14]; for the classical account, see May [May99a, Chapter 23, Section 5].

Definition 10.4.1 (R -lines). We denote by

$$\mathbf{Line}_R \subseteq \mathbf{Mod}_R^{\cong}$$

the full subanima spanned by those R -modules that are isomorphic to R . Its objects are called *R -lines*.

Remark 10.4.2. The pointed connected anima $\mathbf{Line}_R$ is the classifying anima of the group of R -linear automorphisms of R :

$$\mathbf{Line}_R \simeq \mathbf{B} \operatorname{Aut}_R(R).$$

Example 10.4.3. For an ordinary commutative ring k , the group $\operatorname{Aut}_{Hk}(Hk)$ is the discrete group $k^\times$. In particular,

$$\mathbf{Line}_{H\mathbb{F}_2} \simeq * \quad \text{and} \quad \mathbf{Line}_{H\mathbb{Z}} \simeq \mathbf{BC}_2.$$

Definition 10.4.4 (Local systems and trivializations). A *local system of R -lines* on an anima A is a functor $\xi: A \rightarrow \mathbf{Line}_R$. A *trivialization* of ξ is an isomorphism $\xi \cong R_A$, where R_A is the constant functor with value R .

Definition 10.4.5 (Orientation). Let V be a virtual vector bundle of rank d on an anima A . Its associated local system of R -lines is

$$\xi_{V,R}: A \xrightarrow{\xi_V[-d]} \mathbf{Line}_{\mathbb{S}} \xrightarrow{-\otimes R} \mathbf{Line}_R.$$

Here $\xi_V[-d]$ lands in the virtual-rank-zero component $\mathbf{Line}_{\mathbb{S}} \subseteq \mathbf{Pic}(\mathbb{S})$. An *R -orientation* of V is a trivialization $\xi_{V,R} \cong R_A$.

An R -orientation of V induces a morphism of R -modules

$$\operatorname{th}(V)[-d] \otimes R \cong \operatorname{colim}_A \xi_{V,R} \cong R[A] \longrightarrow R,$$

where the last map is the augmentation. By adjunction, this corresponds to a class

$$u_V: \operatorname{th}(V) \longrightarrow R[d],$$

which we call the *Thom class* of the oriented virtual bundle V .

Lemma 10.4.6 (Orientations of inverse virtual bundles). *Let V be a virtual vector bundle on A . An R -orientation of V canonically induces an R -orientation of $-V$.*

Proof. By Remark 10.2.17, including the normalization by virtual rank, there is an isomorphism $\xi_{-V,R} \cong \xi_{V,R}^{-1}$. Inverting a trivialization of $\xi_{V,R}$ gives the required trivialization. $\square$

For an honest vector bundle, this recovers the familiar definition in terms of a Thom class.

Proposition 10.4.7. *Let $E \rightarrow X$ be a rank n vector bundle over a paracompact Hausdorff space of the homotopy type of a cell complex. Under the identification $\mathrm{th}([E]) \cong \Sigma^\infty \mathrm{Th}(E)$ of Lemma 10.2.21, an R -orientation of $[E]$ is the same as a class $\theta \in \widetilde{R}^n(\mathrm{Th}(E))$ whose restriction to every fiber is a unit in $\pi_0(R)$.*

Proof. A class $\theta \in \widetilde{R}^n(\mathrm{Th}(E))$ is a morphism $\mathrm{th}([E])[-n] \otimes R \rightarrow R$ of R -modules. Since the source is the colimit of $\xi_{[E],R}$, adjunction identifies this with a morphism $\xi_{[E],R} \rightarrow R_{\Pi_\infty(X)}$ of local systems. It is an isomorphism precisely when each restriction $\theta_x \in \pi_0(R)$ is a unit. $\square$

Example 10.4.8. Since $\mathrm{Line}_{H\mathbb{F}_2}$ is contractible, every virtual vector bundle of fixed rank has a unique $H\mathbb{F}_2$ -orientation. For $R = H\mathbb{Z}$ and an honest real vector bundle $E \rightarrow X$, the local system $\xi_{[E],H\mathbb{Z}}$ is classified by

$$X \longrightarrow \mathbf{BO}(n) \xrightarrow{\det} \mathbf{BC}_2.$$

Thus an $H\mathbb{Z}$ -orientation is a classical orientation, equivalently a nullhomotopy of $w_1(E)$.

Example 10.4.9 (The universal MO-orientation). The universal rank-zero virtual bundle on $\mathbf{BO}$ is canonically MO-oriented. Indeed, for every $W \in \mathbf{BO}$, symmetric monoidality of J and the definition $\mathrm{MO} = \mathrm{colim}_{V \in \mathbf{BO}} J(V)$ give

$$J(W) \otimes \mathrm{MO} \cong \mathrm{colim}_{V \in \mathbf{BO}} J(W + V) \cong \mathrm{MO}.$$

The second isomorphism is induced by the equivalence $W + (-) : \mathbf{BO} \rightarrow \mathbf{BO}$, and these isomorphisms vary coherently with W . They therefore trivialize the associated local system of MO-lines. Pulling this orientation back along the classifying map of an honest real vector bundle shows that every real vector bundle is canonically MO-orientable.

Theorem 10.4.10 (Thom isomorphism). *Let V be an R -oriented virtual vector bundle of rank d on an anima A . Then there is an isomorphism of R -modules*

$$\mathrm{th}(V) \otimes R \cong (\mathbb{S}[A] \otimes R)[d],$$

and an isomorphism of spectra

$$\mathrm{hom}(\mathrm{th}(V), R) \cong \mathrm{hom}(\mathbb{S}[A][d], R).$$

Proof. Passing to colimits in the chosen trivialization $\xi_{V,R} \cong R_A$ gives

$$\mathrm{th}(V)[-d] \otimes R \cong \mathrm{colim}_A \xi_{V,R} \cong \mathrm{colim}_A R_A \cong R[A],$$

which proves the first isomorphism. The second follows by applying $\mathrm{hom}_R(-, R)$ and using the free-forgetful adjunction between spectra and R -modules. $\square$

Corollary 10.4.11 (Classical Thom isomorphism). *Let $E \rightarrow X$ be a rank n vector bundle over a paracompact Hausdorff space of the homotopy type of a cell complex, equipped with an R -Thom class θ . Then*

$$\widetilde{R}_m(\mathrm{Th}(E)) \cong R_{m-n}(X) \quad \text{and} \quad \widetilde{R}^m(\mathrm{Th}(E)) \cong R^{m-n}(X).$$

The inverse of the cohomological isomorphism is the Thom homomorphism $x \mapsto x \smile \theta$.

Proof. Apply Theorem 10.4.10 to $V = [E]$ and use Proposition 10.4.7 and Lemma 10.2.21. The description of the cohomological map follows from the Thom diagonal

$$\Delta_E: \mathrm{Th}(E) \longrightarrow X_+ \wedge \mathrm{Th}(E), \quad [v] \longmapsto (p(v), [v]).$$

□

Remark 10.4.12 (The mod 2 homology of $\mathbf{MO}$). The universal rank-zero virtual bundle on $\mathbf{BO}$ has a unique $H\mathbb{F}_2$ -orientation. The Thom isomorphism gives

$$H\mathbb{F}_2 \otimes \mathbf{MO} \cong H\mathbb{F}_2[\mathbf{BO}]$$

of $H\mathbb{F}_2$ -modules. This computes the mod 2 homology of $\mathbf{MO}$ in terms of that of $\mathbf{BO}$, whose mod 2 cohomology is the polynomial algebra $\mathbb{F}_2[w_1, w_2, \dots]$ on the Stiefel–Whitney classes. It is the first step of Thom’s calculation of $\pi_*(\mathbf{MO})$ recorded in Remark 10.3.8, but not the whole of it: splitting $\mathbf{MO}$ into a wedge of shifts of $H\mathbb{F}_2$, and identifying the multiplicative structure of the coefficient ring, requires further input.

10.5 Coherence for one-point compactification (online only)

We now prove Proposition 10.2.11. This section supplies the coherence behind the construction and may be skipped without interrupting the subsequent applications.

Let us first describe the strategy. For every topological space X , fiberwise one-point compactification gives a symmetric monoidal functor from vector bundles over X to pointed bundles over X . Passing to fundamental animae gives a lax symmetric monoidal functor

$$\xi_X: \mathrm{Vect}_{\mathbb{R}}^{\mathrm{disc}}(X)^{\simeq} \longrightarrow \mathrm{Fun}(\Pi_{\infty}(X), \mathbf{An}_*), \quad E \longmapsto (x \mapsto \Pi_{\infty}(S^{E_x})),$$

and the structure maps of this functor are pointwise isomorphisms. This only gives a construction on the discrete groupoid of vector bundles, so it does not yet encode the topology on the spaces of bundle isomorphisms. To retain that topology, we apply the construction to $X \times |\Delta^q|$ in every simplicial degree. The projection $X \times |\Delta^q| \rightarrow X$ induces an isomorphism on fundamental animae, so left Kan extension transports the resulting family back to $\Pi_{\infty}(X)$. These functors form a cocone on the simplicial diagram defining $\mathrm{Vect}_{\mathbb{R}}(X)^{\simeq}$, and hence descend to the desired functor. Carrying out the construction naturally in X supplies all the required coherence.

For this coherent construction we need two consequences of the pointed-object formalism developed in Part II. We record them here because their only use is the present application. Their proofs assume reasonable familiarity with the operadic material of Part II and may be treated as black boxes on a first reading.

Lemma 10.5.1 (Pointed realization). *Every finite-product-preserving functor $F: C \rightarrow D$ between ∞ -categories with finite products induces a canonical morphism of pointed ∞ -operads*

$$F_*: (\mathcal{M}_{(C, \times)})_* \longrightarrow (\mathcal{M}_{(D, \times)})_*,$$

whose color functor sends $(\rightarrow X)$ to $(* \rightarrow F(X))$. If D admits finite colimits and its cartesian product preserves them separately in both variables, then its target is naturally equivalent to $\mathcal{M}_{(D_*, \wedge)}$.*

Proof. By Proposition 15.3.6, the functor F has a unique symmetric monoidal refinement for the cartesian monoidal structures. Applying the pointed-object construction $(-)_*: \text{Op}_\infty^* \rightarrow \text{Op}_\infty^{\text{pt}}$ of Theorem 18.4.12 gives the asserted morphism. The identification of the target with the smash-product operad is Lemma 18.4.10. $\square$

Lemma 10.5.2 (Pointwise smash products). *For every anima B , pointed straightening is a symmetric monoidal equivalence*

$$\text{Fun}(B, \text{An}_*) \simeq (\text{An}_B)_*,$$

where the source carries the pointwise smash product and the target carries the smash product induced from the cartesian monoidal structure on An_B .

Proof. The equivalence

$$\text{Ar}(\text{Fun}(B, \text{An})) \simeq \text{Fun}(B, \text{Ar}(\text{An}))$$

identifies pushout products and cofibers on the left with their pointwise counterparts on the right. The symmetric monoidal localization of Lemma 16.3.4 therefore upgrades the canonical equivalence $\text{Fun}(B, \text{An}_*) \simeq \text{Fun}(B, \text{An}_*)$ to a symmetric monoidal equivalence, where the right-hand side carries the pointwise smash product. Unpointed straightening $\text{Fun}(B, \text{An}) \simeq \text{An}_B$ preserves finite products, since pointwise products correspond to fiber products over B . Applying Lemma 10.5.1 finishes the proof. $\square$

These lemmas keep the point-set topology and the higher coherence separate: point-set topology enters through fiberwise one-point compactification, while pointed realization and straightening supply the coherent lax symmetric monoidal structure after passage to anima.

Fix a topological space X . Let $\text{Bun}(X)$ be the 1-category of locally trivial fiber bundles over X and continuous maps over X . Its terminal object is the identity bundle $X \rightarrow X$, and its finite products are fiber products over X . Let $\text{Bun}_{*,c}(X)$ be the 1-category of fiber bundles over X with compact Hausdorff pointed fibers and pointed bundle maps. Its symmetric monoidal structure is the fiberwise smash product, and we refer to the distinguished section of such a bundle as its section at infinity.

Lemma 10.5.3 (Fiberwise one-point compactification). *Fiberwise one-point compactification defines a symmetric monoidal functor*

$$\text{Vect}_\mathbb{R}^{\text{disc}}(X)^\simeq \longrightarrow \text{Bun}_{*,c}(X)^\simeq, \quad E \longmapsto (X \xrightarrow{s_\infty} S^E \rightarrow X).$$

Proof. The total-space functor from vector bundles over X to fiber bundles over X preserves finite products: the total space of $E \oplus F$ is $E \times_X F$. Fiberwise one-point compactification carries this product to the fiberwise smash product. More precisely, removal of the section at infinity gives a natural isomorphism

$$(K \wedge_X L) \setminus \{\infty\} \cong (K \setminus \{\infty\}) \times_X (L \setminus \{\infty\}).$$

Fiberwise one-point compactification and removal of the section at infinity are inverse equivalences between the groupoid of bundles with locally compact Hausdorff fibers and the groupoid $\text{Bun}_{*,c}(X)^\simeq$. Since removal of the section is fully faithful, the displayed identification is compatible with the associativity, unit, and symmetry constraints. This proves the claim. $\square$

We next pass from pointed bundles to families of pointed animae. Taking fundamental animae of total spaces defines a functor

$$\Pi_{\infty,X}: \text{Bun}(X) \longrightarrow \text{An}_{/\Pi_{\infty}(X)}.$$

Lemma 10.5.4 (Parametrized realization). *The functor $\Pi_{\infty,X}$ preserves finite products. On pointed compact bundles it induces a canonical lax symmetric monoidal functor*

$$\Pi_{\infty,X,*}: (\text{Bun}_{*,c}(X), \wedge_X) \longrightarrow ((\text{An}_{/\Pi_{\infty}(X)})_*, \wedge_{\Pi_{\infty}(X)}).$$

Proof. The terminal bundle is sent to the terminal object $\Pi_{\infty}(X) \rightarrow \Pi_{\infty}(X)$. If $E \rightarrow X$ and $F \rightarrow X$ are fiber bundles, their product is $E \times_X F \rightarrow X$. Since every fiber bundle is a Serre fibration by Theorem 2.3.13, we obtain from Proposition 2.3.19 a natural isomorphism

$$\Pi_{\infty}(E \times_X F) \cong \Pi_{\infty}(E) \times_{\Pi_{\infty}(X)} \Pi_{\infty}(F).$$

Thus $\Pi_{\infty,X}$ preserves finite products. Straightening identifies $\text{An}_{/\Pi_{\infty}(X)}$ with $\text{Fun}(\Pi_{\infty}(X), \text{An})$, so it has finite colimits and its cartesian product preserves them separately in both variables. The lax symmetric monoidal refinement on pointed objects is therefore supplied by Lemma 10.5.1. On compact pointed bundles, the tensor operations in this pointed construction are represented by fiberwise smash products: a bundle map $\prod_{i=1}^r K_i \rightarrow L$ which sends the union on which some coordinate is the section at infinity to the section at infinity factors uniquely through a pointed bundle map

$$K_1 \wedge_X \cdots \wedge_X K_r \longrightarrow L.$$

Restricting pointed realization to these bundles therefore gives the asserted lax symmetric monoidal functor. $\square$

Pointed straightening is a symmetric monoidal equivalence

$$((\text{An}_{/\Pi_{\infty}(X)})_*, \wedge_{\Pi_{\infty}(X)}) \simeq \text{Fun}(\Pi_{\infty}(X), \text{An}_*)$$

by Lemma 10.5.2. Composing this equivalence with the functors of Lemmas 10.5.3 and 10.5.4 gives the promised lax symmetric monoidal functor

$$\xi_X: \text{Vect}_{\mathbb{R}}^{\text{disc}}(X)^\simeq \longrightarrow \text{Fun}(\Pi_{\infty}(X), \text{An}_*), \quad E \longmapsto \xi_E.$$

At this stage no preservation of pushouts by $\Pi_{\infty,X}$ has been asserted or used. The lax structure is supplied by pointed realization; we now check pointwise that its structure maps are isomorphisms on sphere bundles.

Lemma 10.5.5. *The lax symmetric monoidal functor ξ_X is symmetric monoidal.*

Proof. It is enough to show that its unit and binary lax structure maps are isomorphisms. Isomorphisms in $\text{Fun}(\Pi_\infty(X), \text{An}_*)$ are detected pointwise. For a point $x \in X$, the bundle projections are Serre fibrations, so taking the fiber of the pointed object associated with $S^E \rightarrow X$ gives the pointed anima $\Pi_\infty(S^{E_x})$. The fiber at x of the binary lax structure map is therefore the canonical map

$$\Pi_\infty(S^{E_x}) \wedge \Pi_\infty(S^{F_x}) \longrightarrow \Pi_\infty(S^{E_x} \wedge S^{F_x}) \cong \Pi_\infty(S^{E_x \oplus F_x}).$$

The pointed spheres are cell complexes and their basepoints are NDR inclusions. Hence Corollaries 2.3.11 and 2.3.20 and Proposition 2.3.10 identify the first map as an isomorphism. The unit comparison is the corresponding isomorphism

$$S^0 \cong \Pi_\infty(S^0).$$

Thus the unit and binary structure maps of the already constructed lax symmetric monoidal functor are isomorphisms, so the functor is symmetric monoidal. $\square$

Lemma 10.5.6 (Naturality in the base). *The symmetric monoidal functors ξ_X assemble into a natural transformation*

$$\xi: \text{Vect}_{\mathbb{R}}^{\text{disc}}(-)^{\simeq} \Longrightarrow \text{Fun}(\Pi_\infty(-), \text{An}_*)$$

between functors $\text{Top}^{\text{op}} \rightarrow \text{CMon}(\text{Cat}_\infty)$, where the functor categories on the right carry their pointwise symmetric monoidal structures.

Proof. The assignments $X \mapsto \text{Vect}_{\mathbb{R}}^{\text{disc}}(X)^{\simeq}$ and $X \mapsto \text{Bun}(X)$ are classified by cartesian fibrations over Top , whose cartesian morphisms are the pullback squares of vector bundles and fiber bundles, respectively. The total-space functor and fiberwise one-point compactification preserve cartesian morphisms.

Taking fundamental anima of the horizontal and vertical maps gives a functor from the cartesian fibration of fiber bundles to the codomain cartesian fibration

$$\text{Ar}(\text{An}) \longrightarrow \text{An}.$$

It preserves cartesian morphisms: for a continuous map $f: Y \rightarrow X$ and a fiber bundle $E \rightarrow X$, the defining pullback square for f^*E is sent by Π_∞ to a pullback square, since $E \rightarrow X$ is a Serre fibration. After cartesian straightening, pointed realization, and pointed straightening, these cartesian functors therefore give the asserted natural transformation. The constructions preserve the fiberwise symmetric monoidal structures, so this is a natural transformation of functors valued in $\text{CMon}(\text{Cat}_\infty)$. $\square$

Lemma 10.5.7 (Transport along simplicial thickenings). *Let X be a topological space and let*

$$p_{X,q}: \Pi_\infty(X \times |\Delta^q|) \longrightarrow \Pi_\infty(X)$$

be induced by the projection. Left Kan extension along $p_{X,q}$ gives symmetric monoidal functors

$$\Phi_{X,q} := (p_{X,q})! \circ \xi_{X \times |\Delta^q|}: \text{Vect}_{\mathbb{R}}^{\text{disc}}(X \times |\Delta^q|)^{\simeq} \longrightarrow \text{Fun}(\Pi_\infty(X), \text{An}_*).$$

As $[q]$ varies, these functors form a cocone on the simplicial diagram defining $\text{Vect}_{\mathbb{R}}(X)^{\simeq}$. This cocone is natural in X .

Proof. The projection $X \times |\Delta^q| \rightarrow X$ is a homotopy equivalence, so $p_{X,q}$ is an isomorphism of anima. Consequently, restriction along $p_{X,q}$ is a symmetric monoidal equivalence

$$p_{X,q}^*: \text{Fun}(\Pi_\infty(X), \text{An}_*) \longrightarrow \text{Fun}(\Pi_\infty(X \times |\Delta^q|), \text{An}_*).$$

Its inverse $(p_{X,q})_!$ inherits the unique symmetric monoidal structure of an inverse equivalence. It follows that $\Phi_{X,q}$ is symmetric monoidal.

A morphism $\alpha: [m] \rightarrow [q]$ in Δ induces a commutative square

$$\begin{array}{ccc} X \times |\Delta^m| & \xrightarrow{\text{id}_X \times \bar{\alpha}} & X \times |\Delta^q| \\ \downarrow & & \downarrow \\ X & \xlongequal{\quad} & X, \end{array}$$

where $\bar{\alpha}$ is the corresponding affine map. All three induced maps of fundamental anima in this square are isomorphisms. The natural transformation of Lemma 10.5.6, together with functoriality of the inverse equivalences given by left Kan extension, therefore gives a symmetric monoidal natural isomorphism

$$\Phi_{X,m} \circ (\text{id}_X \times \bar{\alpha})^* \cong \Phi_{X,q}.$$

These isomorphisms are compatible with composition because both the naturality of ξ and left Kan extension are functorial. They therefore define the asserted cocone. For a continuous map $f: Y \rightarrow X$, the analogous commutative square has the projections to Y and X as its vertical maps. Since these induce isomorphisms of fundamental anima, the same argument gives a symmetric monoidal natural isomorphism

$$\Phi_{Y,q} \circ (f \times \text{id}_{|\Delta^q|})^* \cong (\Pi_\infty f)^* \circ \Phi_{X,q}.$$

The coherence of these isomorphisms under composition follows from the same functoriality, proving that the cocone is natural in X . $\square$

Proof of Proposition 10.2.11. Fix a topological space X . Since every morphism in the source of $\Phi_{X,q}$ is a bundle isomorphism, the cocone of Lemma 10.5.7 factors through the groupoid core of $\text{Fun}(\Pi_\infty(X), \text{An}_*)$. By the defining colimit of Definition 10.2.10, it therefore induces a symmetric monoidal functor

$$S_X^{(-)}: \text{Vect}_{\mathbb{R}}(X)^\simeq \longrightarrow \text{Fun}(\Pi_\infty(X), \text{An}_*) \simeq (\text{An}/\Pi_\infty(X))_*.$$

In simplicial degree zero, this functor sends a vector bundle $E \rightarrow X$ to the spherical fibration ξ_E . The naturality in X asserted in the proposition follows from the naturality of the cocones in Lemma 10.5.7.

It remains to identify the rank components and the restriction of $S_{\text{pt}}^{(-)}$. Real vector bundles over a simplex are trivial, and the automorphisms of the trivial rank n bundle form the singular simplicial group of $\text{GL}_n(\mathbb{R})$. Its realization is $\mathbf{BGL}_n(\mathbb{R})$. Polar decomposition gives a homotopy equivalence $O(n) \hookrightarrow \text{GL}_n(\mathbb{R})$, and hence an isomorphism

$$\mathbf{BO}(n) \cong \mathbf{BGL}_n(\mathbb{R}).$$

Thus the rank n component of $\text{Vect}_{\mathbb{R}}(\text{pt})^\simeq$ is $\mathbf{BO}(n)$. The construction sends a linear isomorphism to its extension over one-point compactifications, so its restriction to this component classifies the standard pointed action of $O(n)$ on $S^{\mathbb{R}^n}$. The block-sum maps give compatibility with the commutative-monoid structures. $\square$

Exercises

11 Duality phenomena

Duality is a recurring phenomenon in algebraic topology. We start in Section 11.1 with its categorical formulation in a symmetric monoidal ∞ -category. We then compare dualizability and compactness in Section 11.2, showing in particular that the dualizable modules over a commutative ring spectrum are precisely the perfect modules. In Section 11.3, we specialize to spectra: Spanier–Whitehead duality relates the cohomology of a dualizable spectrum to the homology of its dual, while Atiyah duality identifies the dual of the suspension spectrum of a closed smooth manifold M with the Thom spectrum $\mathrm{th}(-T_M)$ of its negative virtual tangent bundle. The latter result was first proved by Atiyah [Ati61b], with a precursor in [MS60]. Combining it with the Thom isomorphism, we recover generalized Poincaré duality in Section 11.4.

We refer to Becker and Gottlieb [BG99] for a historical overview of duality phenomena in algebraic topology.

11.1 Symmetric monoidal duality

The modern way to interpret duality phenomena is in terms of the notion of dualizable objects in a symmetric monoidal ∞ -category. Throughout this section, we fix a symmetric monoidal ∞ -category $(C, \otimes, \mathbb{1})$.

Definition 11.1.1 (Dualizable object). Consider objects X and Y of C . We say that a morphism $\mathrm{ev}: Y \otimes X \rightarrow \mathbb{1}$ exhibits Y as a dual to X if there exists another morphism $\mathrm{coev}: \mathbb{1} \rightarrow X \otimes Y$ such that the triangle identities are satisfied: there exist commutative diagrams

$$\begin{array}{ccc} & X \otimes Y \otimes X & \\ \mathrm{coev} \otimes \mathrm{id} \nearrow & & \searrow \mathrm{id} \otimes \mathrm{ev} \\ X & \xlongequal{\quad} & X \end{array} \quad \text{and} \quad \begin{array}{ccc} & Y \otimes X \otimes Y & \\ \mathrm{id} \otimes \mathrm{coev} \nearrow & & \searrow \mathrm{ev} \otimes \mathrm{id} \\ Y & \xlongequal{\quad} & Y. \end{array}$$

In this case we refer to ev as the *evaluation map* and to coev as the *coevaluation map*, and call (Y, ev) a *duality datum* for X . We say that an object X is *dualizable* if it admits such a duality datum.

The following result guarantees that the duality datum (Y, ev) is unique if it exists:

Proposition 11.1.2. *Consider objects X and Y of C and consider a morphism $\mathrm{ev}: Y \otimes X \rightarrow \mathbb{1}$ in C . The following conditions are equivalent:*

- (1) *The morphism $\mathrm{ev}: Y \otimes X \rightarrow \mathbb{1}$ exhibits Y as a dual to X ;*

(2) For all objects Z and W of C , the composite

$$\mathrm{Hom}_C(W, Z \otimes Y) \xrightarrow{- \otimes X} \mathrm{Hom}_C(W \otimes X, Z \otimes Y \otimes X) \xrightarrow{\mathrm{ev} \circ -} \mathrm{Hom}_C(W \otimes X, Z)$$

is an equivalence.

(3) The map in (2) is an equivalence for $(W, Z) = (\mathbb{1}, X)$ and for $(W, Z) = (Y, \mathbb{1})$.

Proof. We first show that (1) implies (2). Let $\mathrm{coev}: \mathbb{1} \rightarrow X \otimes Y$ be the corresponding coevaluation map. It then follows directly from the triangle identities that an inverse to the map in (2) is given by the composite

$$\mathrm{Hom}_C(W \otimes X, Z) \xrightarrow{- \otimes Y} \mathrm{Hom}_C(W \otimes X \otimes Y, Z \otimes Y) \xrightarrow{- \circ \mathrm{coev}} \mathrm{Hom}_C(W, Z \otimes Y).$$

It is clear that (2) implies (3). Finally, assume that (3) is satisfied. Taking $W = \mathbb{1}$ and $Z = X$, we get that the composite

$$\mathrm{Hom}_C(\mathbb{1}, X \otimes Y) \xrightarrow{- \otimes X} \mathrm{Hom}_C(X, X \otimes Y \otimes X) \xrightarrow{\mathrm{ev} \circ -} \mathrm{Hom}_C(X, X)$$

is an equivalence. In particular, there exists a morphism $\mathrm{coev}: \mathbb{1} \rightarrow X \otimes Y$ that is mapped to id_X under this equivalence. In particular, the composite

$$X \xrightarrow{\mathrm{coev} \otimes \mathrm{id}} X \otimes Y \otimes X \xrightarrow{\mathrm{id} \otimes \mathrm{ev}} X$$

is homotopic to id_X , showing one of the two triangle identities. To show that the other triangle identity is also satisfied, consider $W = Y$ and $Z = \mathbb{1}$, so that the composite

$$\mathrm{Hom}_C(Y, Y) \xrightarrow{- \otimes X} \mathrm{Hom}_C(Y \otimes X, Y \otimes X) \xrightarrow{\mathrm{ev} \circ -} \mathrm{Hom}_C(Y \otimes X, \mathbb{1})$$

is an equivalence. We claim that both the map $\mathrm{id}_Y: Y \rightarrow Y$ as well as the composite $Y \xrightarrow{\mathrm{id} \otimes \mathrm{coev}} Y \otimes X \otimes Y \xrightarrow{\mathrm{ev} \otimes \mathrm{id}} Y$ are sent to $\mathrm{ev}: Y \otimes X \rightarrow \mathbb{1}$ under this equivalence. This is clear for id_Y . For $(\mathrm{ev} \otimes \mathrm{id}) \circ (\mathrm{id} \otimes \mathrm{coev})$ this follows from the following commutative diagram:

$$\begin{array}{ccccc} Y \otimes X & \xrightarrow{\mathrm{id} \otimes \mathrm{coev} \otimes \mathrm{id}} & Y \otimes X \otimes Y \otimes X & \xrightarrow{\mathrm{ev} \otimes \mathrm{id} \otimes \mathrm{id}} & Y \otimes X \\ & \searrow & \downarrow \mathrm{id} \otimes \mathrm{id} \otimes \mathrm{ev} & & \downarrow \mathrm{ev} \\ & & Y \otimes X & \xrightarrow{\mathrm{ev}} & \mathbb{1} \end{array}$$

This shows that also the second triangle identity is satisfied, finishing the proof. $\square$

Thus being a duality datum is a property of the evaluation map ev : once condition (2) holds, the required coevaluation and triangle identities follow. We will use this criterion below to establish duality by constructing and analyzing an evaluation map.

Convention 11.1.3. Let X be a dualizable object of C . By condition (2) of Proposition 11.1.2, its dual represents the functor $\mathrm{Hom}_C(- \otimes X, \mathbb{1}): C^{\mathrm{op}} \rightarrow \mathbf{An}$. In particular, the dual is unique, and we denote it by $X^\vee$. An object representing this functor need not be a dual when X is not known to be dualizable; see the discussion of weak duals in Section 11.1.1.

Definition 11.1.4 (Invertible object). An object X of C is called *invertible* if there exists another object X^{-1} and an isomorphism $X \otimes X^{-1} \cong \mathbb{1}$. Equivalently, X is invertible if the functor $X \otimes -: C \rightarrow C$ is an equivalence.

The full subcategory $\text{Pic}(C) \subseteq C^\simeq$ spanned by the invertible objects is called the *Picard anima* of C .

Lemma 11.1.5. *Every invertible object is dualizable.*

Proof. Let X be an invertible object and let X^{-1} be an inverse with isomorphism $\text{ev}: X^{-1} \otimes X \xrightarrow{\sim} \mathbb{1}$. We claim that ev exhibits X^{-1} as a dual to X . Using Proposition 11.1.2 we must show that for all objects Z and W of C the composite

$$\text{Hom}_C(W, Z \otimes X^{-1}) \xrightarrow{- \otimes X} \text{Hom}_C(W \otimes X, Z \otimes X^{-1} \otimes X) \xrightarrow{\text{ev} \circ -} \text{Hom}_C(W \otimes X, Z)$$

is an equivalence. But both maps are equivalences: the first because the functor $- \otimes X: C \rightarrow C$ is an equivalence, and the second because ev is an isomorphism. $\square$

Corollary 11.1.6. *The unit object $\mathbb{1} \in C$ is dualizable.*

Proof. It is invertible, using $\mathbb{1} \otimes \mathbb{1} \cong \mathbb{1}$. $\square$

Lemma 11.1.7. *Symmetric monoidal functors preserve dualizable objects and their duals. Moreover, if $X, Y \in C$ are dualizable, then $X \otimes Y$ is dualizable with dual $X^\vee \otimes Y^\vee$.*

Proof. A symmetric monoidal functor preserves evaluation and coevaluation maps and the triangle identities. For the second claim, apply this observation to the tensor-product functor $C \times C \rightarrow C$, which is symmetric monoidal for the componentwise tensor product on $C \times C$, and to the dualizable object (X, Y) . $\square$

11.1.1 Internal homs

The notion of duality is closely related to that of *internal homs*. Internal homs provide a candidate $D(X)$ for the dual of every object; dualizability is the additional condition that tensoring with this weak dual actually represents mapping out of X . This perspective will allow us to prove that dualizable objects form a thick subcategory.

Definition 11.1.8. Given two objects $X, Z \in C$, an object $\underline{\text{Hom}}(X, Z) \in C$ equipped with a map $\text{ev}: \underline{\text{Hom}}(X, Z) \otimes X \rightarrow Z$ is said to *exhibit $\underline{\text{Hom}}(X, Z)$ as an internal hom from X to Z* if for every third object W the composite

$$\text{Hom}_C(W, \underline{\text{Hom}}(X, Z)) \xrightarrow{- \otimes X} \text{Hom}_C(W \otimes X, \underline{\text{Hom}}(X, Z) \otimes X) \xrightarrow{\text{ev} \circ -} \text{Hom}_C(W \otimes X, Z)$$

is an equivalence.

Remark 11.1.9. Internal homs are unique if they exist. When they exist for every Z , the assignment $Z \mapsto \underline{\text{Hom}}(X, Z)$ defines a right adjoint to $- \otimes X: C \rightarrow C$. A symmetric monoidal ∞ -category admitting all internal homs is also called *closed*.

Lemma 11.1.10. Assume that C admits internal homs. Then for every object $X \in C$, the adjunction between $- \otimes X$ and $\underline{\text{Hom}}(X, -)$ gives a currying isomorphism: for $Y, Z \in C$ there is a natural equivalence

$$\underline{\text{Hom}}(X \otimes Y, Z) \simeq \underline{\text{Hom}}(X, \underline{\text{Hom}}(Y, Z)).$$

Proof. This follows from the Yoneda lemma, and the observation that for every fourth object $W \in C$ there are natural equivalences

$$\begin{aligned} \text{Hom}_C(W, \underline{\text{Hom}}(X \otimes Y, Z)) &\simeq \text{Hom}_C(W \otimes X \otimes Y, Z) \\ &\simeq \text{Hom}_C(W \otimes X, \underline{\text{Hom}}(Y, Z)) \\ &\simeq \text{Hom}_C(W, \underline{\text{Hom}}(X, \underline{\text{Hom}}(Y, Z))). \end{aligned} \quad \square$$

We now provide a formulation of duality in terms of internal homs.

Lemma 11.1.11. Let X and Y be objects and let $\text{ev}: Y \otimes X \rightarrow \mathbb{1}$ be a morphism in C . Then the following conditions are equivalent:

- (1) The morphism ev exhibits Y as a dual to X .
- (2) For every object Z of C , the morphism $\text{id}_Z \otimes \text{ev}: Z \otimes Y \otimes X \rightarrow Z$ exhibits $Z \otimes Y$ as an internal hom $\underline{\text{Hom}}(X, Z)$.
- (3) The functor $X \otimes -: C \rightarrow C$ admits a right adjoint $\underline{\text{Hom}}(X, -): C \rightarrow C$, and for every object Z of C the map

$$Z \otimes Y \rightarrow \underline{\text{Hom}}(X, Z)$$

induced by the morphism $\text{id} \otimes \text{ev}: Z \otimes Y \otimes X \rightarrow Z$ is an isomorphism in C .

Proof. The equivalence between (1) and (2) is a reformulation of Proposition 11.1.2. The equivalence between (2) and (3) is clear. $\square$

There is often an a priori candidate for the dual of X , sometimes called the *weak dual*:

Definition 11.1.12. Assume that C admits internal homs. We define the *internal duality functor* $D: C^{\text{op}} \rightarrow C$ as

$$D(-) := \underline{\text{Hom}}(-, \mathbb{1}): C^{\text{op}} \rightarrow C.$$

If X is dualizable, the evaluation map $\text{ev}: X^\vee \otimes X \rightarrow \mathbb{1}$ induces an isomorphism $X^\vee \xrightarrow{\sim} \underline{\text{Hom}}(X, \mathbb{1}) = D(X)$.

Corollary 11.1.13. Let X be an object of C for which the internal hom functor $\underline{\text{Hom}}(X, -): C \rightarrow C$ exists. Then X is dualizable if and only if for every $Z \in C$ the natural map $Z \otimes D(X) \rightarrow \underline{\text{Hom}}(X, Z)$ adjoint to the composite $Z \otimes D(X) \otimes X \xrightarrow{Z \otimes \text{ev}} Z \otimes \mathbb{1} \simeq Z$ is an equivalence.

Remark 11.1.14. The objects called dualizable here are called *strongly dualizable* in Lewis, May, and Steinberger [LMS86, Chapter III, Definition 1.1]. That reference also gives the following useful criterion: if C is closed, then X is dualizable precisely when the morphism $\mathbb{1} \rightarrow \underline{\text{Hom}}(X, X)$ adjoint to id_X factors through the natural map $X \otimes D(X) \rightarrow \underline{\text{Hom}}(X, X)$.

Definition 11.1.15. A *stably symmetric monoidal ∞ -category* is a symmetric monoidal ∞ -category C which is stable and whose tensor product $- \otimes -: C \times C \rightarrow C$ is exact in both variables.

Lemma 11.1.16. *Let C be a stably symmetric monoidal ∞ -category which admits internal homs. Then the collection of dualizable objects forms a thick subcategory of C , i.e. it is a stable subcategory closed under retracts.*

Proof. By Corollary 11.1.13, an object X is dualizable if and only if the natural map

$$Z \otimes D(X) \rightarrow \underline{\mathrm{Hom}}(X, Z)$$

is an equivalence for every $Z \in C$. Since both sides are exact functors in X , it follows that the collection of objects X for which it is an equivalence forms a stable subcategory of C . It is also closed under retracts since equivalences in C are closed under retracts. $\square$

11.2 Dualizability and compactness

Both compactness and dualizability express a finiteness property of an object, but they are defined differently: compactness concerns mapping out of the object, whereas dualizability concerns tensoring with it. We will now give conditions under which these notions agree, and then apply them to modules over ring spectra.

Lemma 11.2.1. *Let C be a symmetric monoidal ∞ -category with filtered colimits, and assume that the tensor product in C preserves filtered colimits in both variables. If the monoidal unit $\mathbb{1} \in C$ is compact, then every dualizable object is compact.*

Proof. Let X be a dualizable object in C , and let $X^\vee$ denote its dual. Then the functor $X \otimes -: C \rightarrow C$ admits a right adjoint of the form $X^\vee \otimes -: C \rightarrow C$. In particular, we have

$$\mathrm{Hom}_C(X, -) \simeq \mathrm{Hom}_C(X \otimes \mathbb{1}, -) \simeq \mathrm{Hom}_C(\mathbb{1}, X^\vee \otimes -),$$

and the right-hand side preserves filtered colimits by the assumptions on C . $\square$

In practice, the converse frequently holds as well: all compact objects are dualizable.

Lemma 11.2.2 (cf. Hovey, Palmieri, and Strickland [HPS97]). *Let C be a presentably symmetric monoidal stable ∞ -category in the sense of Definition 22.5.1, and let $\mathcal{G}$ be a small collection of compact generators of C . If every object of $\mathcal{G}$ is dualizable, then every compact object of C is dualizable.*

Proof. The tensor product admits internal homs by Theorem 22.2.5. The result on compact generation, Proposition 22.3.4 from Chapter 22, which we use as a black box, identifies the compact objects of C with the thick subcategory generated by $\mathcal{G}$. The claim now follows from Lemma 11.1.16. $\square$

Corollary 11.2.3. *Let C be a presentably symmetric monoidal stable ∞ -category in the sense of Definition 22.5.1, and let $\mathcal{G}$ be a small collection of compact generators. Then the compact objects of C are precisely its dualizable objects if and only if the unit $\mathbb{1}$ is compact and every object of $\mathcal{G}$ is dualizable.*

Proof. If the unit is compact and every object of $\mathcal{G}$ is dualizable, the claim follows from Lemmas 11.2.1 and 11.2.2. Conversely, if the compact and dualizable objects coincide, then the unit is compact because it is always dualizable, while every object of $\mathcal{G}$ is dualizable because it is compact. $\square$

We now apply this discussion to the perfect modules introduced in Definition 8.1.7 as the thick subcategory of left R -modules generated by R . When R is commutative, Proposition 8.1.5 identifies left modules with the symmetric monoidal category Mod_R , and perfectness admits a third characterization in addition to its descriptions by finite construction and compactness from Lemma 8.1.8.

Proposition 11.2.4. *Let R be a commutative ring spectrum. An R -module is perfect if and only if it is dualizable in the symmetric monoidal ∞ -category $(\text{Mod}_R, \otimes_R)$.*

Proof. The category Mod_R is presentable and its tensor product preserves colimits, so it admits internal homs by Proposition 8.1.5 and Theorem 22.2.5. By Lemma 11.1.16, the dualizable R -modules therefore form a thick subcategory of Mod_R . Since this subcategory contains the monoidal unit R , every perfect R -module is dualizable. Conversely, the unit R is compact and the tensor product of R -modules preserves colimits in both variables. Hence every dualizable R -module is compact by Lemma 11.2.1, and therefore perfect by Lemma 8.1.8. $\square$

11.3 Spanier–Whitehead and Atiyah duality

The abstract notion of duality is especially useful in stable homotopy theory because it interchanges homology and cohomology. Indeed, if X is a dualizable spectrum, then mapping out of X is equivalent to tensoring with its dual $X^\vee$. Consequently, the cohomology of X can be computed as the homology of $X^\vee$, and conversely the cohomology of $X^\vee$ can be computed as the homology of X . For spectra X and E , we use the notation

$$E_k(X) := \pi_k(E \otimes X) \quad \text{and} \quad E^k(X) := \pi_{-k} \text{hom}(X, E).$$

For $X = \mathbb{S}[A]$, this recovers the conventions for the E -(co)homology of an anima A .

Lemma 11.3.1. *Let X and E be spectra and assume that X is dualizable. Then there are natural isomorphisms*

$$E_*(X^\vee) \cong E^{-*}(X) \quad \text{and} \quad E^*(X^\vee) \cong E_{-*}(X).$$

Proof. The first isomorphism follows from

$$E_*(X^\vee) \cong \pi_*(E \otimes X^\vee) \cong \pi_*(\text{hom}(X, E)) \cong E^{-*}(X).$$

The second follows by applying the first to $X^\vee$, whose dual is X . $\square$

The dual of a dualizable spectrum is therefore also called its *Spanier–Whitehead dual*. The thick-subcategory property from Lemma 11.1.16 supplies the basic class of examples.

Corollary 11.3.2. *Every finite spectrum is dualizable and hence admits a Spanier–Whitehead dual.*

Proof. The category $\mathbf{Sp}$ admits internal homs, the sphere spectrum is dualizable, and the dualizable spectra form a thick subcategory by Lemma 11.1.16. $\square$

Corollary 11.3.3. *Let X be a finite anima. Then the spectrum $\mathbb{S}[X]$ is dualizable.*

Proof. The collection of anima X for which $\mathbb{S}[X]$ is dualizable is closed under finite colimits and contains $*$ $\in \mathbf{An}$; consequently it contains all finite anima. $\square$

This gives an abstract existence result, but does not by itself identify the dual geometrically. For a closed smooth manifold, Atiyah duality provides such an identification in terms of the tangent bundle.

Let M be a closed smooth manifold of dimension n , i.e. a compact smooth manifold without boundary. By Morse theory, M has the homotopy type of a finite CW-complex [Mil63, Theorem 3.5 and Corollary 6.7]. It thus follows from Corollary 11.3.3 that its suspension spectrum $\mathbb{S}[M]$ is dualizable. We will give an explicit description of its dual, due to Atiyah [Ati61b]: there is an isomorphism

$$D(\mathbb{S}[M]) \cong \mathrm{th}(-T_M)$$

between the dual of $\mathbb{S}[M]$ and the Thom spectrum of the *negative stable tangent bundle* $-T_M$. Here $-T_M$ denotes the inverse of the virtual vector bundle $[T_M]$ of Example 10.2.20, which has rank $-n$, and $\mathrm{th}(-T_M)$ is its Thom spectrum in the sense of Construction 10.2.16. No new construction is required: the group completion entering the stable J-homomorphism already assigns a Thom spectrum to every virtual vector bundle, of any rank.

To get a better understanding of the Thom spectrum $\mathrm{th}(-T_M)$, we will now show that we can write it as $\Sigma^{\infty-N}(\mathrm{Th}(v_\varphi))$ for a suitable honest vector bundle v_φ over M . The main ingredient is the following simple observation:

Lemma 11.3.4. *Let M be a compact smooth n -manifold. Then there exists a natural number N and a smooth embedding $\varphi: M \hookrightarrow \mathbb{R}^N$.*

Proof. By compactness, we may choose finitely many coordinate charts $\varphi_i: U_i \rightarrow \mathbb{R}^n$, smaller open subsets V_i which still cover M and satisfy $\overline{V_i} \subseteq U_i$, and smooth functions $\rho_i: M \rightarrow [0, 1]$ supported in U_i and strictly positive on V_i . Extend $\rho_i \varphi_i$ by zero outside U_i and consider

$$\varphi := (\rho_i \varphi_i, \rho_i)_{i=1}^k: M \longrightarrow (\mathbb{R}^n \times \mathbb{R})^k.$$

If $x \neq y$, choose i with $x \in V_i$. Equality of the i th pair of coordinates would force $\rho_i(y) = \rho_i(x) > 0$, hence $y \in U_i$, and then $\varphi_i(y) = \varphi_i(x)$, a contradiction. Thus φ is injective. The same coordinate pair shows that $d\varphi$ is injective at x : if it killed a tangent vector, then both $d\rho_i$ and $d(\rho_i \varphi_i) = \rho_i d\varphi_i + \varphi_i d\rho_i$ would vanish, and hence so would $d\varphi_i$. Thus φ is an injective immersion. Since M is compact, it is a smooth embedding. $\square$

Remark 11.3.5. If $n > 0$, the *Whitney embedding theorem* shows that N can be chosen to be $2n$ in the previous lemma.

For the remainder of this section, we will fix once and for all a smooth embedding $\varphi: M \hookrightarrow \mathbb{R}^N$. We will denote by $\nu := \nu_\varphi \rightarrow M$ the normal bundle of this embedding, i.e., the cokernel of the map $T_M \rightarrow \varphi^*(T_{\mathbb{R}^N}) \cong \underline{\mathbb{R}}^N$ of vector bundles over M :

$$T_M \rightarrow \underline{\mathbb{R}}^N \rightarrow \nu.$$

By choosing a metric on the vector bundle $\underline{\mathbb{R}}^N$ one obtains an orthogonal decomposition

$$\underline{\mathbb{R}}^N \cong T_M \oplus \nu.$$

Passing to classes of virtual vector bundles and using Example 10.2.20, this gives $[T_M] + [\nu] \cong [\underline{\mathbb{R}}^N]$, so that $-T_M$ is the virtual vector bundle $[\nu] - [\underline{\mathbb{R}}^N]$ of rank $-n$. Hence Lemma 10.2.21 identifies

$$\text{th}(-T_M) \cong \text{th}(\nu)[-N] \cong \Sigma^{\infty-N}(\text{Th}(\nu)). \quad (11.1)$$

We will now construct the duality data exhibiting $\text{th}(-T_M)$ as dual to $\mathbb{S}[M]$. In the order fixed by Definition 11.1.1, these are maps

$$\text{ev}: \text{th}(-T_M) \otimes \mathbb{S}[M] \rightarrow \mathbb{S} \quad \text{and} \quad \text{coev}: \mathbb{S} \rightarrow \mathbb{S}[M] \otimes \text{th}(-T_M)$$

satisfying the triangle identities. Using Equation (11.1), it is enough to construct maps of pointed spaces

$$\text{Th}(\nu) \wedge M_+ \rightarrow S^N \quad \text{and} \quad S^N \rightarrow M_+ \wedge \text{Th}(\nu).$$

The constructions below naturally produce these maps with the two smash factors reversed. We use the symmetry of the smash product to put them in the displayed order, but suppress these symmetry maps in the point-set formulas. We will use Thom-space functoriality and the Pontryagin–Thom collapse map.

Construction 11.3.6. Let $f: X \rightarrow Y$ be a continuous map of topological spaces. Let $p: E \rightarrow Y$ be a vector bundle over Y , and let $f^*(E) \rightarrow X$ be its pullback bundle. By applying functoriality of the Thom space construction (Construction 10.1.7) to the commutative diagram

$$\begin{array}{ccc} f^*(E) & \longrightarrow & E \\ f^*(p) \downarrow & & \downarrow p \\ X & \xrightarrow{f} & Y, \end{array}$$

we obtain a continuous map

$$f_*: \text{Th}(f^*E) \rightarrow \text{Th}(E).$$

Example 11.3.7. Consider the terminal map $p: M \rightarrow \text{pt}$, and consider the trivial vector bundle $E = \mathbb{R}^N \rightarrow \text{pt}$. Then the pullback bundle $p^*(\mathbb{R}^N)$ is the trivial vector bundle $\underline{\mathbb{R}}^N$ over M , and we obtain a map

$$p_*: \Sigma^N(M_+) \xrightarrow{\text{Example 10.1.5}} \text{Th}(\underline{\mathbb{R}}^N) \xrightarrow{p_*} \text{Th}(\mathbb{R}^N) \xrightarrow{\text{Example 10.1.5}} S^N.$$

Example 11.3.8. Consider the diagonal map $\Delta: M \rightarrow M \times M$, and let $E = \nu \times \underline{0}$ be the product bundle over $M \times M$, where ν is the normal bundle of our fixed embedding $\varphi: M \hookrightarrow \mathbb{R}^N$ and $\underline{0}$ is the zero bundle. Observe that the pullback bundle $\Delta^*(E) = \Delta^*(\nu \times \underline{0})$ is isomorphic to the direct sum $\nu \oplus \underline{0}$ of these two bundles, and hence is isomorphic to ν itself. The above construction thus produces a map

$$\Delta_*: \text{Th}(\nu) \rightarrow \text{Th}(\nu \times \underline{0}) \xrightarrow{\text{Example 10.1.6}} \text{Th}(\nu) \wedge \text{Th}(\underline{0}) \cong \text{Th}(\nu) \wedge M_+.$$

Recall the Pontryagin–Thom collapse map $\text{PT}(i): N'_+ \rightarrow \text{Th}(\nu(i))$ of Construction 10.1.8. We will also need the following twisted version.

Construction 11.3.9 (Twisted Pontryagin–Thom collapse map). The Pontryagin–Thom collapse map also has a ‘twisted’ version, where in addition to a smooth embedding $i: N \hookrightarrow N'$ of compact manifolds without boundary we are given a vector bundle E over N' . It takes the form

$$\text{PT}(i, E): \text{Th}(E) \rightarrow \text{Th}(\nu(i) \oplus i^*E).$$

To define it, choose a metric on E and consider the composite embedding $N \xrightarrow{i} N' \xrightarrow{s_0} E$, where the second map is the zero-section. Its normal bundle is the direct sum $\nu(i) \oplus i^*E$, so the ordinary Pontryagin–Thom construction gives a collapse map

$$E_+ \longrightarrow \text{Th}(\nu(i) \oplus i^*E).$$

We may choose its tubular neighborhood inside the open unit disk bundle of E . The collapse map then sends the complement of that disk bundle, and in particular the unit sphere bundle, to the basepoint. It therefore factors through the quotient $E_+ \rightarrow E^+ \cong \text{Th}(E)$ of Remark 10.1.4, giving the desired twisted collapse map. Its based homotopy class is independent of the metric and tubular-neighborhood choices.

Example 11.3.10. Consider our fixed embedding $\varphi: M \hookrightarrow \mathbb{R}^N$. Since M is compact, its image is closed in S^N and misses the point at infinity. Thus $M \hookrightarrow S^N$ is a pointed smooth embedding, and the Pontryagin–Thom construction provides a map

$$\text{PT}(\varphi): S^N \rightarrow \text{Th}(\nu).$$

Example 11.3.11. Consider the diagonal embedding $\Delta: M \hookrightarrow M \times M$. Its normal bundle is isomorphic to the tangent bundle T_M of M . As a twist, we use the external direct sum $E = \underline{0} \times \nu$ over $M \times M$, where $\nu = \nu(\varphi)$ as before. The pullback of E along Δ is isomorphic to ν , as in Example 11.3.8, and we see that there is a trivialization $\nu(\Delta) \oplus \Delta^*(E) \cong T_M \oplus \nu \cong \underline{\mathbb{R}}^N$. The twisted Pontryagin–Thom map in this case takes the form

$$\text{PT}(\Delta, \underline{0} \times \nu): M_+ \wedge \text{Th}(\nu) \cong \text{Th}(\underline{0}) \wedge \text{Th}(\nu) \cong \text{Th}(\underline{0} \oplus \nu) \rightarrow \text{Th}(\underline{\mathbb{R}}^N) \cong \Sigma^N(M_+).$$

Combining the four examples, we obtain our evaluation and coevaluation maps:

$$\text{ev}: M_+ \wedge \text{Th}(\nu) \xrightarrow{\text{PT}(\Delta, \underline{0} \times \nu)} \Sigma^N(M_+) \xrightarrow{P_*} \Sigma^N(S^0) = S^N.$$

$$\text{coev}: S^N \xrightarrow{\text{PT}(\varphi)} \text{Th}(\nu) \xrightarrow{\Delta_*} \text{Th}(\nu) \wedge M_+.$$

To finish the proof of Atiyah duality, it remains to show that these maps satisfy the space-level analogues of the triangle identities. We will use the following standard compatibilities of the Pontryagin–Thom construction. We formulate them only for compact manifolds without boundary, which is the case needed below.

Proposition 11.3.12 (Pontryagin–Thom calculus). *Pontryagin–Thom collapse maps have the following properties, up to based homotopy.*

- (1) Isotopy invariance. *Isotopic embeddings induce homotopic collapse maps. The same holds for twisted collapse maps, provided the twisting bundle is fixed throughout the isotopy.*
- (2) Compatibility with composition. *Let $N \xhookrightarrow{i} P \xhookrightarrow{j} Q$ be smooth embeddings of compact manifolds, and let E be a vector bundle over Q . After choosing a splitting of the normal-bundle sequence*

$$0 \rightarrow \nu(i) \rightarrow \nu(ji) \rightarrow i^* \nu(j) \rightarrow 0,$$

the composite

$$\mathrm{Th}(E) \xrightarrow{\mathrm{PT}(j,E)} \mathrm{Th}(\nu(j) \oplus j^* E) \xrightarrow{\mathrm{PT}(i,\nu(j) \oplus j^* E)} \mathrm{Th}(\nu(i) \oplus i^* \nu(j) \oplus (ji)^* E)$$

agrees up to homotopy with $\mathrm{PT}(ji, E)$ under the resulting identification of target Thom spaces.

- (3) Transverse base change. *Let $f: M_1 \rightarrow M_2$ be a smooth map between compact smooth manifolds without boundary and let $i_2: N_2 \hookrightarrow M_2$ be a closed smooth submanifold. Assume that f is transverse to N_2 , and set $N_1 := f^{-1}(N_2)$, with inclusion $i_1: N_1 \hookrightarrow M_1$ and induced map $\bar{f}: N_1 \rightarrow N_2$. Then N_1 is a smooth submanifold and there is a natural isomorphism $\nu(i_1) \cong \bar{f}^* \nu(i_2)$. Under this identification, the square*

$$\begin{array}{ccc} (M_1)_+ & \xrightarrow{f_+} & (M_2)_+ \\ \mathrm{PT}(i_1) \downarrow & & \downarrow \mathrm{PT}(i_2) \\ \mathrm{Th}(\nu(i_1)) & \xrightarrow{\bar{f}_*} & \mathrm{Th}(\nu(i_2)) \end{array}$$

commutes up to based homotopy. More generally, for a vector bundle E over M_2 , the square

$$\begin{array}{ccc} \mathrm{Th}(f^* E) & \xrightarrow{f_*} & \mathrm{Th}(E) \\ \mathrm{PT}(i_1, f^* E) \downarrow & & \downarrow \mathrm{PT}(i_2, E) \\ \mathrm{Th}(\nu(i_1) \oplus i_1^* f^* E) & \xrightarrow{\bar{f}_*} & \mathrm{Th}(\nu(i_2) \oplus i_2^* E) \end{array}$$

commutes up to based homotopy.

Proof sketch. Choose Riemannian metrics and tubular neighborhoods. An isotopy of embeddings has a tubular neighborhood over the parameter interval, which gives the homotopy in (1). For (2), choose nested tubular neighborhoods for $N \subseteq P \subseteq Q$; collapsing successively then agrees with collapsing the resulting tubular neighborhood of N in Q . For (3), transversality identifies the normal bundle of N_1 with $\bar{f}^* \nu(i_2)$. Compactness allows the tubular neighborhoods to be chosen compatibly over the whole square. Comparing the corresponding collapse quotients gives the displayed homotopies. The same choices made in the total space of E prove the twisted statements. $\square$

We are finally ready to prove Atiyah duality. Consider the composite

$$\begin{aligned} \mathrm{th}(-T_M) \otimes \mathbb{S}[M] &\cong \Sigma^{\infty-N} \mathrm{Th}(\nu) \otimes \mathbb{S}[M] \cong \Sigma^{\infty-N} (\mathrm{Th}(\nu) \wedge M_+) \\ &\cong \Sigma^{\infty-N} (M_+ \wedge \mathrm{Th}(\nu)) \\ &\xrightarrow{\Sigma^{\infty-N} \mathrm{PT}(\Delta, \underline{0} \times \nu)} \Sigma^{\infty-N} \Sigma^N(M_+) \\ &\xrightarrow{P_*} \Sigma^{\infty-N} \Sigma^N(S^0) \cong \mathbb{S}. \end{aligned}$$

Its homotopy class is independent of the metrics and tubular neighborhoods used in its construction.

Theorem 11.3.13 (Atiyah duality). *The displayed map exhibits $\text{th}(-T_M)$ as a dual of $\mathbb{S}[M]$, and hence determines an isomorphism:*

$$D(\mathbb{S}[M]) \cong \text{th}(-T_M).$$

Proof. We verify the triangle identities before applying suspension spectra. Each triangle composite is built from twisted collapse maps and a Thom-space pushforward. Transverse base change rewrites it as the composite of a collapse map for a graph embedding and the pushforward along an identity map. Compatibility with composition and isotopy invariance then identify both factors with identity maps.

Observe that the map we wrote is obtained by applying $\Sigma^{\infty-N}(-): \text{Top}_* \rightarrow \text{An}_* \rightarrow \text{Sp}$ to the following composite:

$$\text{ev}: M_+ \wedge \text{Th}(\nu) \xrightarrow{\text{PT}(\Delta, \underline{0} \times \nu)} \Sigma^N(M_+) \xrightarrow{P_*} \Sigma^N(S^0) = S^N.$$

For the candidate coevaluation map, we similarly apply $\Sigma^{\infty-N}(-)$ to the following composite:

$$\text{coev}: S^N \xrightarrow{\text{PT}(\varphi)} \text{Th}(\nu) \xrightarrow{\Delta_*} \text{Th}(\nu) \wedge M_+.$$

To see that the triangle identities are satisfied, it is enough to prove corresponding homotopies at the level of topological spaces. More precisely, the triangle identity on $\text{th}(-T_M)$ is obtained by applying $\Sigma^{\infty-2N}$ to the composite

$$S^N \wedge \text{Th}(\nu) \xrightarrow{\text{coev} \wedge \text{id}} \text{Th}(\nu) \wedge M_+ \wedge \text{Th}(\nu) \xrightarrow{\text{id} \wedge \text{ev}} \text{Th}(\nu) \wedge S^N$$

while the triangle identity on $\mathbb{S}[M]$ is obtained by applying $\Sigma^{\infty-N}$ to

$$M_+ \wedge S^N \xrightarrow{\text{id} \wedge \text{coev}} M_+ \wedge \text{Th}(\nu) \wedge M_+ \xrightarrow{\text{ev} \wedge \text{id}} S^N \wedge M_+$$

Thus it suffices to show that these two composites are homotopic to the respective identities, up to the displayed symmetries of the smash product. We will start with the former. Consider the following pullback square:

$$\begin{array}{ccc} M & \xrightarrow{\Delta} & M \times M \\ \Delta \int \downarrow & \lrcorner & \downarrow \int \text{id} \times \Delta \\ M \times M & \xrightarrow{\Delta \times \text{id}} & M \times M \times M. \end{array}$$

Let us check the transversality hypothesis in this instance. At a point (x, y) of the inverse image of $\text{im}(\text{id} \times \Delta)$, the image of $d(\Delta \times \text{id})$ consists of triples (v, v, w) , while the tangent space to $\text{im}(\text{id} \times \Delta)$ consists of triples (a, b, b) . Every triple (r, s, t) decomposes as

$$(r, s, t) = (s, s, t) + (r - s, 0, 0),$$

with the first summand of the former type and the second of the latter type. Thus $\Delta \times \text{id}$ is transversal to $\text{im}(\text{id} \times \Delta)$. Applying Proposition 11.3.12 to this square (i.e. we take $M_2 := M \times M \times M$, $N_2 := M \times M$, $i_2 = \text{id} \times \Delta$, $M_1 := M \times M$, $N_1 := M$, $f = \Delta \times \text{id}$ and $E = \nu \times \underline{0} \times \nu$) gives the following homotopy-commutative diagram:

$$\begin{array}{ccc} \text{Th}(\nu) \wedge \text{Th}(\nu) & \xrightarrow{(\Delta \times \text{id})_*} & \text{Th}(\nu) \wedge M_+ \wedge \text{Th}(\nu) \\ \text{PT}(\Delta, \nu \times \nu) \downarrow & & \downarrow \text{PT}(\text{id} \times \Delta, \nu \times \underline{0} \times \nu) \\ \Sigma^N \text{Th}(\nu) & \xrightarrow{\Delta_*} & \text{Th}(\nu) \wedge \Sigma^N(M_+). \end{array}$$

All collapse maps below whose source contains S_+^N factor through the reduced quotient, since the relevant embeddings lie in $\mathbb{R}^N \times M$. We use these reduced maps without further notation. Unwinding the definitions of $\text{coev} \wedge \text{id}$ and $\text{id} \wedge \text{ev}$, we then get the following homotopy-commutative diagram. Its top row is the triangle composite, its southeast-pointing arrows are Pontryagin–Thom collapse maps, and its northeast-pointing arrows are Thom-space pushforwards:

$$\begin{array}{ccccc}
S^N \wedge \text{Th}(\nu) & \xrightarrow{\text{coev} \wedge \text{id}} & \text{Th}(\nu) \wedge M_+ \wedge \text{Th}(\nu) & \xrightarrow{\text{id} \wedge \text{ev}} & \text{Th}(\nu) \wedge S^N \\
\searrow \text{PT}(\varphi \times \text{id}, \underline{0} \times \nu) & & \nearrow \Delta_* \wedge \text{id} = (\Delta \times \text{id})_* & & \nearrow \text{id} \wedge \text{PT}(\Delta, \underline{0} \times \nu) = \text{PT}(\text{id} \times \Delta, \nu \times \underline{0} \times \nu) \\
& & \text{Th}(\nu) \wedge \text{Th}(\nu) & & \text{Th}(\nu) \wedge \Sigma^N(M_+) \\
& \searrow \text{PT}(\Delta, \nu \times \nu) & & \nearrow \Delta_* & \\
& & \Sigma^N \text{Th}(\nu) & &
\end{array}$$

$\text{id} \wedge p_* = (\text{id} \times p)_*$

The upper two triangles commute by the definitions of coev and ev , while the central diamond is the transverse base-change square displayed above. By compatibility with composition in Proposition 11.3.12, the left diagonal composite is induced on the reduced quotient by the twisted Pontryagin–Thom collapse map for the composite

$$j: M \xrightarrow{\Delta} M \times M \xrightarrow{\varphi \times \text{id}} S^N \times M,$$

where we twist by the bundle $\underline{0} \times \nu$ on $S^N \times M$. The corresponding unreduced twisted collapse map is

$$\text{PT}(j, \underline{0} \times \nu): S_+^N \wedge \text{Th}(\nu) \rightarrow \Sigma^N \text{Th}(\nu),$$

Since the image of j is contained in $\mathbb{R}^N \times M \subseteq S^N \times M$, this map collapses the entire fiber over the point at infinity and therefore factors through the quotient $S_+^N \wedge \text{Th}(\nu) \twoheadrightarrow S^N \wedge \text{Th}(\nu)$. The embedding $j = (\varphi, \text{id})$ is isotopic, via the straight-line isotopy in $\mathbb{R}^N \subseteq S^N$, to the embedding $(0, \text{id}): M \hookrightarrow \mathbb{R}^N \times M \subseteq S^N \times M$. By isotopy invariance, the induced map on the reduced quotient is therefore homotopic to the one obtained from $(0, \text{id})$. Using $\mathbb{R}^N \times M$ as a tubular neighborhood for this inclusion, we see that the latter is the identity on $S^N \wedge \text{Th}(\nu)$.

The right diagonal is the pushforward functoriality of Thom spaces applied to the composite

$$M \xrightarrow{\Delta} M \times M \xrightarrow{\text{id} \times p} M \times \text{pt} = M.$$

But this composite is simply the identity on M , hence it induces the identity on $S^N \wedge \text{Th}(\nu)$. We conclude that the composite $(\text{id} \wedge \text{ev}) \circ (\text{coev} \wedge \text{id})$ is homotopic to the identity on $S^N \wedge \text{Th}(\nu)$, proving the first triangle identity.

The second triangle identity follows by the same argument, applied to the transposed pullback square. More precisely, we now take $i_2 = \Delta \times \text{id}$, $f = \text{id} \times \Delta$ and use the twist $E = \underline{0} \times \nu \times \underline{0}$ over $M \times M \times M$. Transverse base change and compatibility with composition identify the composite $(\text{ev} \wedge \text{id}) \circ (\text{id} \wedge \text{coev})$ with the composite of the Pontryagin–Thom collapse map for the graph embedding

$$(\text{id}, \varphi): M \hookrightarrow M \times S^N$$

and the Thom-space pushforward induced by $(p \times \text{id}) \circ \Delta = \text{id}_M$. The graph embedding is isotopic to $(\text{id}, 0): M \hookrightarrow M \times \mathbb{R}^N$, whose collapse map is the identity on $M_+ \wedge S^N$ under the standard Thom identifications. Hence the second triangle composite is also homotopic to the identity. $\square$

11.4 Poincaré duality

Poincaré duality for closed smooth manifolds is an immediate consequence of Atiyah duality. Recall that Poincaré duality says that when M is a closed orientable smooth n -manifold, then for each $k \in \mathbb{Z}$ there is an isomorphism

$$- \cap [M]: H^{n-k}(M) \xrightarrow{\sim} H_k(M).$$

Many readers will have encountered this statement in their algebraic topology courses, where it is usually proved in quite a down-to-earth way, based on manipulations of simplices in the singular simplicial complex of M and using a couple of reduction arguments. This proof makes it look like Poincaré duality is intrinsically a statement about singular (co)homology. However, this is not the case: there is a more general version of Poincaré duality for cohomology theories represented by commutative ring spectra.

The orientation theory of Section 10.4 applies to the virtual bundles occurring in Atiyah duality, and this is all that is needed.

Theorem 11.4.1 (Poincaré duality). *Let E be a commutative ring spectrum, and let M be a closed smooth n -manifold equipped with an E -orientation of the virtual tangent bundle $[T_M]$, or equivalently of $-T_M$ by Lemma 10.4.6. Then there is an isomorphism of spectra*

$$\mathbb{S}[M] \otimes E \cong \text{hom}(\mathbb{S}[M], E[n]).$$

In particular, for every $k \in \mathbb{Z}$ there is an isomorphism

$$E_k(M) \cong E^{n-k}(M).$$

Proof. By Theorem 11.3.13, the dual of the spectrum $\mathbb{S}[M]$ in Sp is $\text{th}(-T_M)$, and in particular we obtain an isomorphism of spectra

$$\mathbb{S}[M] \otimes E \cong \text{hom}(\text{th}(-T_M), E).$$

By Lemma 10.4.6, the given orientation of $[T_M]$ induces an orientation of $-T_M$. Since this virtual bundle has rank $-n$, Theorem 10.4.10 gives an isomorphism $\text{th}(-T_M) \otimes E \cong (\mathbb{S}[M] \otimes E)[-n]$ in Mod_E , and it follows that

$$\begin{aligned} \text{hom}(\text{th}(-T_M), E) &\cong \text{hom}_E(\text{th}(-T_M) \otimes E, E) \\ &\cong \text{hom}_E((\mathbb{S}[M] \otimes E)[-n], E) \\ &\cong \text{hom}(\mathbb{S}[M][-n], E) \\ &\cong \text{hom}(\mathbb{S}[M], E[n]). \end{aligned}$$

Combining these two isomorphisms thus gives the first claim. The second claim immediately follows by applying $\pi_k(-)$ to both sides. $\square$

The preceding proof is the abstract form of Poincaré duality. We next relate it to the classical fundamental-class formulation. The orientation of $-T_M$ gives a Thom class $u_{-T_M}: \text{th}(-T_M) \rightarrow E[-n]$. Define the E -fundamental class

$$[M]_E \in E_n(M) = \pi_n(E \otimes \mathbb{S}[M])$$

to be represented, after shifting, by the composite

$$\mathbb{S} \xrightarrow{\text{coev}} \mathbb{S}[M] \otimes \text{th}(-T_M) \xrightarrow{\text{id} \otimes u_{-T_M}} \mathbb{S}[M] \otimes E[-n].$$

Using the embedding $M \hookrightarrow \mathbb{R}^N$ chosen in the proof of Theorem 11.3.13, this is concretely the class determined by the Pontryagin–Thom collapse $S^N \rightarrow \text{Th}(\nu)$, followed by the diagonal $\Delta_*: \text{Th}(\nu) \rightarrow \text{Th}(\nu) \wedge M_+$ and then the Thom class of the normal bundle on the first factor.

Proposition 11.4.2. *Under the above identification, the isomorphism $E^{n-k}(M) \rightarrow E_k(M)$ inverse to the one in Theorem 11.4.1 is cap product with the fundamental class:*

$$- \cap [M]_E: E^{n-k}(M) \xrightarrow{\sim} E_k(M).$$

Proof sketch. Represent a cohomology class by a map $a: \mathbb{S}[M] \rightarrow E[n-k]$. Cap product with $[M]_E$ is represented, after shifting, by the composite

$$\mathbb{S} \xrightarrow{[M]_E} \mathbb{S}[M] \otimes E[-n] \xrightarrow{\mathbb{S}[\Delta] \otimes \text{id}} \mathbb{S}[M] \otimes \mathbb{S}[M] \otimes E[-n] \xrightarrow{\text{id} \otimes a \otimes \text{id}} \mathbb{S}[M] \otimes E[n-k] \otimes E[-n] \xrightarrow{\text{id} \otimes \mu} \mathbb{S}[M] \otimes E[-k].$$

Substituting the definition of $[M]_E$ expresses this as the composite obtained from the Atiyah coevaluation, the class a , and the Thom class. The triangle identity for the Atiyah duality datum then identifies it with the inverse of the isomorphism in Theorem 11.4.1. $\square$

Example 11.4.3. For $E = H\mathbb{F}_2$, every real vector bundle is E -orientable because $\text{Line}_{H\mathbb{F}_2} \simeq *$. Thus every closed smooth manifold satisfies mod-2 Poincaré duality. For $E = H\mathbb{Z}$, Example 10.4.8 identifies an $H\mathbb{Z}$ -orientation with a classical orientation, or equivalently with a nullhomotopy of $w_1(T_M)$. This recovers integral Poincaré duality for orientable manifolds.

Remark 11.4.4. A Spin^c -structure on M determines a KU-orientation of $[T_M]$ through the construction of Atiyah, Bott, and Shapiro [ABS64]. At the other extreme, Example 10.4.9 shows that every real vector bundle is canonically MO-orientable. This ties generalized Poincaré duality directly to the bordism theory of Section 10.3.

Example 11.4.5. The manifold $\mathbb{R}P^2$ is not integrally orientable, and integral Poincaré duality indeed fails: $H_0(\mathbb{R}P^2; \mathbb{Z}) \cong \mathbb{Z}$, whereas $H^2(\mathbb{R}P^2; \mathbb{Z}) \cong \mathbb{Z}/2$. With $\mathbb{F}_2$ -coefficients, it is orientable and Poincaré duality holds.

Remark 11.4.6. For a compact manifold with boundary, the corresponding statement is Poincaré–Lefschetz duality:

$$E_k(M) \cong E^{n-k}(M, \partial M).$$

under the appropriate E -orientation hypothesis. At the spectrum level, Atiyah duality identifies the Spanier–Whitehead dual of $M/\partial M$ with the Thom spectrum $\text{th}(-T_M)$. Together with the Thom isomorphism, this gives the displayed identification of the homology of M with its relative cohomology.

Exercises

Exercise 11.1. Let C be the 1-category Vect_k of vector spaces over some field k . Show that every finite-dimensional vector space V is dualizable in Vect_k , with dual given by $V^\vee = V^* := \text{Hom}_k(V, k)$.

Exercise 11.2. Give a direct proof of Lemma 11.1.5 by constructing the coevaluation map from an inverse X^{-1} and checking the two triangle identities.

Exercise 11.3. Let C be a symmetric monoidal ∞ -category, and assume that C is stable and that the tensor product $X \otimes - : C \rightarrow C$ is exact for every object X of C .

- (1) Show that the zero object is dualizable;
- (2) Deduce that the subcategory of C consisting of the dualizable objects is closed under pushouts (*Hint*: show it is closed under direct sums and cofibers);
- (3) Give a direct proof of Corollary 11.3.3 from the preceding two parts.

Part II

Operads and higher algebra

Introduction to Part II

In Definition 8.1.1, we introduced the notion of a *symmetric monoidal ∞ -category*, defined as a commutative monoid in Cat_∞ . This definition, while conceptually clean, is rather implicit: it tells us that a symmetric monoidal structure *exists* as an object of a certain ∞ -category, but does not directly provide us with tools for constructing or manipulating such structures. For example, even for something as natural as the cartesian monoidal structure on the ∞ -category $\mathbf{An}$ it is not a priori clear how to write down the corresponding commutative monoid in Cat_∞ . As a result, we are in need of a general framework that lets us construct and manipulate coherent algebraic structures more easily. This is precisely the purpose of *∞ -operads*, the central notion of this part.

Our approach differs from the one in Lurie’s foundational text [Lur17] in that we define ∞ -operads as functors to the *span category* $\text{Span}(\mathbf{Fin})$ of finite sets, rather than to the category $\mathbf{Fin}_*$ of finite pointed sets. While the two approaches are equivalent (see Section 17.4), the span-based perspective has the advantage that all the required conditions can be formulated in terms of finite products, which is conceptually cleaner.

The part is organized as follows. We start in Chapter 12 with the classical theory of colored operads, gradually motivating the ∞ -categorical definition; a reader already comfortable with colored operads may skim this chapter. The span category $\text{Span}(\mathbf{Fin})$ plays a central role in our approach, and its construction as an ∞ -category requires some care; this is carried out in Chapter 13. With these preparations in place, Chapter 14 introduces ∞ -operads, symmetric monoidal ∞ -categories, algebras, lax monoidal functors, and monoidal Bousfield localizations; it is the most important chapter in this part. We then turn to applications: Chapter 15 equips any ∞ -category with finite products or coproducts with its natural symmetric monoidal structure, and Chapter 16 develops Day convolution and uses it to construct the *tensor product* on the ∞ -category of spectra, one of the main goals of this book. In Chapter 17, we construct the *envelope* $\text{Env}(O)$ of an ∞ -operad, which freely promotes it to a symmetric monoidal ∞ -category; this chapter also establishes the comparison with Lurie’s definition. The general operadic Day convolution and stabilization constructions are then developed in Chapter 18. Next, Chapter 19 develops the abstract theory of associative algebras and their modules, on which the theory of ring spectra in Part I (Chapter 8) is based, and concludes by constructing connective complex K-theory as a commutative ring spectrum. Finally, Chapter 20 develops monoidal Dwyer–Kan localizations and applies them to derived categories and Eilenberg–MacLane modules.

Prerequisites. This part makes heavy use of the theory of cocartesian fibrations and the straightening/unstraightening equivalence from Chapter 23. The construction of span categories in Chapter 13 also relies on the presentation of ∞ -categories by complete Segal anima from Chapter 24. Readers unfamiliar with either topic are encouraged to consult the corresponding chapter in Part III as needed.

12 Operads: the heuristics

Before diving into the abstract theory, the goal of the present chapter is to provide the intuitions behind the definition of ∞ -operads. Roughly speaking, we may think of an ∞ -operad as encoding a type of ‘algebraic structure’; for example, there will be an ∞ -operad for commutative algebras, one for associative algebras, one for left modules over an associative algebra, and so on and so forth. We will start by explaining the 1-categorical situation: In Section 12.1 we will introduce non-colored operads as structures encoding collections of ‘ n -ary operations’ $A^{\otimes n} \rightarrow A$, and in Section 12.2 we discuss the colored variant that allows for more general operations of the form $A_1 \otimes \cdots \otimes A_n \rightarrow B$. In order to motivate the ∞ -categorical analogue of operads, we will explain in Section 12.3 how to rephrase the classical notion of colored operads in a way that robustly encodes all the coherences without needing to specify them by hand. This will then pave the way for our definition of ∞ -operads, which we give in Section 12.4. We conclude with brief pointers to other models for ∞ -operads.

12.1 Non-colored operads

Let us start by thinking about how to describe some algebraic structure on an object A of a symmetric monoidal category $(C, \otimes, \mathbb{1})$. Often, such algebraic structures are given in terms of a collection of *n -ary operations* for natural numbers $n \geq 0$, i.e., maps of the form $A^{\otimes n} \rightarrow A$ from the n -fold tensor product of A to itself. For example, if A is an associative algebra object in C , we have the multiplication map $\cdot : A^{\otimes 2} \rightarrow A$, $(x, y) \mapsto xy$. Also the unit is of this form: when $n = 0$, an n -fold tensor product of A is by convention the monoidal unit $\mathbb{1} \in C$, and we often want to treat the unit element $1 \in A$ as a 0-ary operation $1 : \mathbb{1} \rightarrow A$.

Observe that operations may be combined to obtain more of them. For example, assume that for natural numbers $k, n_1, \dots, n_k \geq 0$ we are given operations on A of the form

$$P : A^{\otimes k} \rightarrow A, \quad Q_i : A^{\otimes n_i} \rightarrow A, \quad i = 1, \dots, k.$$

Then we may combine them into an n -ary operation $P(Q_1, \dots, Q_k)$ for $n := \sum_{i=1}^k n_i$ by forming the following composite:

$$A^{\otimes n} \cong \bigotimes_{i=1}^k A^{\otimes n_i} \xrightarrow{\bigotimes_{i=1}^k Q_i} \bigotimes_{i=1}^k A = A^{\otimes k} \xrightarrow{P} A.$$

We will refer to the operation $P(Q_1, \dots, Q_k)$ as the *composition* of P with $(Q_1, \dots, Q_k)$. In similar fashion we may also always consider the *identity operation* $\text{id} : A \rightarrow A, a \mapsto a$. Finally, we may permute the inputs of an n -ary operation by any permutation $\sigma \in \Sigma_n$:

$$(P\sigma)(x_1, \dots, x_n) := P(x_{\sigma^{-1}(1)}, \dots, x_{\sigma^{-1}(n)}).$$

If we apply this process to our associative algebra A , we find that there are precisely $n!$ formal ways of constructing an n -ary operation: for every permutation $\sigma \in \Sigma_n := \text{Aut}_{\text{Set}}(\{1, \dots, n\})$, we obtain a multiplication map $m_\sigma: A^{\otimes n} \rightarrow A$ given by $m_\sigma(x_1, \dots, x_n) := x_{\sigma(1)} \cdots x_{\sigma(n)}$. For a particular algebra, some of these maps may of course agree.

The notion of a (non-colored, symmetric, non-enriched) *operad* is trying to axiomatize the collections of possible operations one may have, together with their various compositions and permutations:

Definition 12.1.1 ([May72; BV73]). A *non-colored operad* $\mathcal{O}$ consists of the following data:

- A set $\mathcal{O}(n)$ of n -ary operations, for every $n \geq 0$;
- A *composition map*

$$-\circ -: \mathcal{O}(k) \times \prod_{i=1}^k \mathcal{O}(n_i) \rightarrow \mathcal{O}(n_1 + \cdots + n_k), \quad (P, Q_1, \dots, Q_k) \mapsto P(Q_1, \dots, Q_k)$$

for all natural numbers $k, n_1, \dots, n_k \geq 0$;

- An *identity operation* $\text{id} \in \mathcal{O}(1)$;
- A right action of the symmetric group Σ_n on $\mathcal{O}(n)$ for every $n \geq 0$.

These data are subject to the following conditions:

- Composition is *unital* and *associative*:

$$P(\text{id}, \dots, \text{id}) = P = \text{id}(P),$$

$$R(P_1(Q_1^1, \dots, Q_{k_1}^1), \dots, P_l(Q_1^l, \dots, Q_{k_l}^l)) = (R(P_1, \dots, P_l))(Q_1^1, \dots, Q_{k_1}^1, Q_1^2, \dots, Q_{k_l}^l).$$

- Composition is compatible with the permutation actions; see Exercise 12.1 for details.

The following are some examples of non-colored operads:

Example 12.1.2. The *associative operad* $\mathcal{A}ssoc$ has $\mathcal{A}ssoc(n) := \Sigma_n$, i.e., there is one n -ary operation for every permutation $\sigma \in \Sigma_n$.

Example 12.1.3. The *commutative operad* $Comm$ has $Comm(n) := *$, i.e., there is a *single* n -ary operation for every $n \geq 0$.

Example 12.1.4. There is the *trivial operad* $\mathcal{T}riv$, whose only operation is the identity operation:

$$\mathcal{T}riv(1) = \{\text{id}\}, \quad \mathcal{T}riv(n) = \emptyset \quad \text{for } n \neq 1.$$

Example 12.1.5. The *pointed operad* $\mathcal{E}_0$ has only one non-identity operation:

$$\mathcal{E}_0(1) = \{\text{id}\}, \quad \mathcal{E}_0(0) = \{\eta\}, \quad \mathcal{E}_0(n) = \emptyset \quad \text{for } n \geq 2.$$

Example 12.1.6. For every field K there is an operad enriched in K -vector spaces called the *Lie operad*, denoted $\mathcal{L}ie$, which captures the operations present in a Lie algebra. The vector space $\mathcal{L}ie(n)$ is the subspace of the free Lie algebra $L(x_1, \dots, x_n)$ spanned by the ‘multilinear’ elements: formal sums of iterated Lie brackets that use each variable x_i exactly once. Lie algebras are algebras over $\mathcal{L}ie$ in the enriched sense; forgetting the enrichment gives an underlying set-valued operad.

Example 12.1.7. For every $k \geq 0$, there is a topological operad¹⁶ called the *little k -cubes operad*, often denoted $\mathcal{E}_k$. Its n -ary operations are the so-called ‘rectilinear’ embeddings $\bigsqcup_{i=1}^n [0, 1]^k \rightarrow [0, 1]^k$ of a disjoint union of n copies of the k -dimensional cube into the k -dimensional cube. Here ‘rectilinear’ means that on each of the n components it is given by an affine function, i.e., maps of the form

$$[0, 1]^k \rightarrow [0, 1]^k, \quad (x_i)_{i=1}^k \mapsto (a_i x_i + b_i)_{i=1}^k, \quad a_i > 0,$$

which allow us to scale and shift the cube but not rotate or distort it in any other way. The composition in $\mathcal{E}_k$ is given by composition of rectilinear embeddings. For $k = 0$, this recovers the pointed operad $\mathcal{E}_0$ described above.

Remark 12.1.8. For $k = 1$, there is an operad morphism $\mathcal{E}_1 \rightarrow \mathcal{A}ssoc$ such that for every $n \geq 0$ the map $\mathcal{E}_1(n) \rightarrow \mathcal{A}ssoc(n)$ is a homotopy equivalence, see Exercise 12.6. In particular, after passing from sets and topological spaces to animae (as we will do when working with ∞ -operads) there is no longer a distinction between $\mathcal{E}_1$ and $\mathcal{A}ssoc$. In more classical setups, for example when working with model categories, the homotopically correct operad to work with is often $\mathcal{E}_1$. This is the reason why in older literature you will often find the phrase ‘ $\mathcal{E}_1$ -algebras’ in places where we would say ‘associative algebra’.

Example 12.1.9 (Endomorphism operad). For an object A of a symmetric monoidal category C , we may form the *endomorphism operad* $End_C(A)$, given by $End_C(A)(n) := Hom_C(A^{\otimes n}, A)$. Composition is induced by composition in C and the associativity isomorphisms for tensor products. The right action of Σ_n is given by precomposition with the symmetry isomorphisms of $A^{\otimes n}$, using the convention from the display above.

Given the structure an operad has, it is not difficult to guess the correct notion of *morphisms* between them: if $\mathcal{O}$ and $\mathcal{P}$ are operads, then an operad morphism $f: \mathcal{O} \rightarrow \mathcal{P}$ is a collection of maps $f(n): \mathcal{O}(n) \rightarrow \mathcal{P}(n)$ that preserves the identity operation, the composition of operations and the permutation of operations.

Definition 12.1.10. Let $\mathcal{O}$ be a non-colored operad and let C be a symmetric monoidal category. We define an *$\mathcal{O}$ -algebra in C* as a pair (A, f_A) consisting of an object A of C together with an operad morphism $f_A: \mathcal{O} \rightarrow End_C(A)$. In particular, f_A consists of an n -ary operation $P_A: A^{\otimes n} \rightarrow A$ for every $P \in \mathcal{O}(n)$ satisfying the condition that composition of operations in $\mathcal{O}$ corresponds to the composition of operations on A .

The basic examples of algebras over these operads are worked out in Exercise 12.4 at the end of the chapter.

Remark 12.1.11. Note that $Comm$ is the *terminal* operad: there is precisely one operad map $\mathcal{O} \rightarrow Comm$ given by the unique maps $\mathcal{O}(n) \rightarrow Comm(n) = *$. Also note that $Triv$ is the *initial* operad: there is precisely one operad map $Triv \rightarrow \mathcal{O}$ sending the unique operation id of $Triv$ to the identity operation $id \in \mathcal{O}(1)$.

¹⁶This means that we remember the topology on each set of operations $\mathcal{O}(n)$.

12.2 Colored operads

While the notion of non-colored operads introduced previously is elegant, it is insufficient for our purposes. To illustrate this, let us discuss the algebraic structure of an associative algebra A acting on some module M . To encode such an action, we need a map in C of the form

$$\text{act}: A \otimes M \rightarrow M, \quad (a, m) \mapsto am,$$

and its source is not a tensor-power of a single object. More generally, given a collection of objects (A_x) in C we would like to have some notion of operad which can deal with operations of the form

$$P: A_{y_1} \otimes \cdots \otimes A_{y_n} \rightarrow A_x$$

for certain choices of indices $y_1, \dots, y_n, x$. This leads to the following ‘multicolored’ version of Definition 12.1.1:

Definition 12.2.1. A *colored operad* (or *multicategory*) O consists of the following structure:

- There is a set $O^\simeq$ of *colors* (sometimes also called the *objects* of O);
- For colors $x, y_1, \dots, y_n \in O^\simeq$, there is a set $O((y_1, \dots, y_n); x)$ of *multimorphisms* from $(y_1, \dots, y_n)$ to x . We refer to x as the *output color* and to the y_i as the *input colors*;
- If we are additionally given colors $z_j^i \in O^\simeq$ for $i = 1, \dots, n$ and $j = 1, \dots, k_i$, there is a *composition map*


$$\circ: O((y_1, \dots, y_n); x) \times \prod_{i=1}^n O((z_1^i, \dots, z_{k_i}^i); y_i) \rightarrow O((z_1^1, \dots, z_{k_1}^1, \dots, z_1^n, \dots, z_{k_n}^n); x).$$

- For every color x there is an identity operation $\text{id}_x \in O((x); x)$;
- Composition should be unital and associative like before;
- For every permutation $\sigma \in \Sigma_n$ there is a ‘permutation isomorphism’


$$O((y_1, \dots, y_n); x) \xrightarrow{\sim} O((y_{\sigma(1)}, \dots, y_{\sigma(n)}); x), \quad P \mapsto P\sigma,$$

which is compatible with the multiplication in the group Σ_n and is furthermore compatible with composition in O . We leave a precise formulation of this condition to the reader; see Exercise 12.1.


We may pictorially think of a multimorphism P from $(y_1, \dots, y_n)$ to x as follows:




The composition of multimorphisms may then be illustrated as follows (with $n = k_1 = 3$, $k_2 = 0$ and $k_3 = 2$):



The identity multimorphism will be pictured as follows:



Finally, we have the permutation operation on multimorphisms, which we may picture as follows (with $z_i := y_{\sigma(i)}$):



Definition 12.2.2. If $\mathcal{P}$ is another colored operad, then a *morphism* $f: \mathcal{O} \rightarrow \mathcal{P}$ of colored operads consists of a map of sets $f: \mathcal{O}^\approx \rightarrow \mathcal{P}^\approx$ on colors and for all colors $x, y_1, \dots, y_n \in \mathcal{O}^\approx$ a map

$$f: \mathcal{O}((y_1, \dots, y_n); x) \rightarrow \mathcal{P}((fy_1, \dots, fy_n); fx)$$

on multimorphisms, in such a way that f preserves identity operations, composition of operations and permutations.

Let us go through some examples of colored operads:

Example 12.2.3 (From non-colored to colored). Every non-colored operad $\mathcal{O}$ may be regarded as a colored operad with a single object: we take $\mathcal{O}^\approx := \{*\}$ and we set

$$\mathcal{O}(\underbrace{(*, \dots, *)}_{n \text{ times}}; *) := \mathcal{O}(n).$$

From now on, we will abuse notation and regard non-colored operads as colored operads in this way. In particular, we obtain colored operads $\mathcal{A}ssoc$, $\mathcal{C}omm$, $\mathcal{L}ie$ and $\mathcal{E}_k$.

Warning 12.2.4. While $Comm$ is still terminal as a colored operad, $\mathcal{T}riv$ is no longer initial as a colored operad: the initial colored operad has an empty set of colors.

Example 12.2.5 (From colored to non-colored). Conversely, for every colored operad O and every color $x \in O^\approx$ we obtain a non-colored operad O_x by setting

$$O_x(n) := O(\underbrace{(x, \dots, x)}_{n \text{ times}}; x).$$

The operad O_x comes with an obvious inclusion map $O_x \hookrightarrow O$ of colored operads.

Example 12.2.6 (Trivial operad on a 1-category). Let C be a 1-category. We define a colored operad $\mathcal{T}riv_C$, called the *trivial operad generated by C* , as follows: its colors are the objects of C , and its multimorphisms are given by

$$\mathcal{T}riv_C((y_1, \dots, y_n); x) := \begin{cases} \text{Hom}_C(y_1, x) & \text{if } n = 1, \\ \emptyset & \text{otherwise.} \end{cases}$$

The identity operations are the identity morphisms of C , the composition is the composition in C , and the permutation operations are trivial since Σ_1 is the trivial group. In other words, $\mathcal{T}riv_C$ has no operations besides the unary ones, and these are precisely the morphisms of C .

The trivial operad $\mathcal{T}riv$ is the special case $C = *$, while $C = \emptyset$ gives the initial colored operad from the warning above.

Example 12.2.7 (Multimorphism operad). Let C be a symmetric monoidal category. We define a colored operad $\mathcal{M}_C$, called the *multimorphism operad*¹⁷ of C , as follows:

- The colors of $\mathcal{M}_C$ are the objects of C .
- For objects $x, y_1, \dots, y_n$ of C , the multimorphisms in $\mathcal{M}_C$ are given by maps from the tensor product:

$$\mathcal{M}_C((y_1, \dots, y_n); x) := \text{Hom}_C(y_1 \otimes \dots \otimes y_n; x);$$

- The identity operations are given by the identity maps $\text{id}_x \in \text{Hom}_C(x, x)$;
- The composition of operations is given by the composition in C ;
- The permutation operations are induced by the isomorphisms $y_1 \otimes \dots \otimes y_n \simeq y_{\sigma(1)} \otimes \dots \otimes y_{\sigma(n)}$ coming from the symmetry isomorphisms in C .

Note that for an object A of C , the non-colored operad $(\mathcal{M}_C)_A$ obtained by applying Example 12.2.5 is precisely the endomorphism operad $\text{End}_C(A)$ introduced in Example 12.1.9.

Example 12.2.8 (Module operad). There is the *module operad* Mod which has two colors a and m , so that $Mod^\approx = \{a, m\}$. The operations are given as follows:

¹⁷This operad is usually not given a name in the literature. Some sources refer to it as the ‘colored endomorphism operad of C ’, generalizing the terminology from Example 12.1.9. The terminology ‘multimorphism operad’ was suggested to the author by Jan Steinebrunner.

- If the output color is a , the operations should encode a commutative algebra structure, so we only want to allow operations all of whose input colors are a as well:

$$\text{Mod}((x_1, \dots, x_n); a) := \begin{cases} * & \text{if } x_j = a \text{ for all } j = 1, \dots, n, \\ \emptyset & \text{otherwise,} \end{cases}$$

- If the output color is m , the operations should encode the action of the commutative algebra on a module, so we only want operations that have precisely one input color that is m , while all others are a :

$$\text{Mod}((x_1, \dots, x_n); m) := \begin{cases} * & \text{if for some } i \text{ we have } x_i = m \text{ and } x_j = a \text{ for } j \neq i, \\ \emptyset & \text{otherwise.} \end{cases}$$

The composition is uniquely determined.

Note that restricting the operations to the color a gives the commutative operad, while restricting the operations to the color m gives the trivial operad:

$$\text{Mod}_a \cong \text{Comm} \quad \text{and} \quad \text{Mod}_m \cong \text{Triv}.$$

Example 12.2.9 (Left module operad). There is also a non-commutative version of the module operad, capturing the structure of a left module over an associative algebra. We will denote it by $\mathcal{LMod}$ and call it the *left module operad*. It again has two colors, $\mathcal{LMod}^\approx = \{a, m\}$. The operations are defined as follows:

- If the output color is a , we again only want operations all of whose input colors are a , but we should allow them to be ordered arbitrarily:

$$\mathcal{LMod}((x_1, \dots, x_n); a) := \begin{cases} \{\text{linear orders on } \{1, \dots, n\}\} & \text{if } x_j = a \text{ for all } j = 1, \dots, n, \\ \emptyset & \text{otherwise.} \end{cases}$$

- If the output color is m , we should again only allow operations such that precisely one of the inputs is m . To allow for permutations of the other inputs, let us write $I_m := \{i \in \{1, \dots, n\} \mid x_i = m\}$ for the set of indices corresponding to module inputs, and let $I_a := \{j \in \{1, \dots, n\} \mid x_j = a\}$ be the set of indices corresponding to algebra inputs. We then set

$$\mathcal{LMod}((x_1, \dots, x_n); m) := \begin{cases} \{\text{linear orders on } I_a\} & \text{if } |I_m| = 1 \text{ (which implies } |I_a| = n - 1), \\ \emptyset & \text{if } |I_m| \neq 1. \end{cases}$$

Note that this set has cardinality $(n - 1)!$.

Composition substitutes an ordered list into each algebra input and concatenates the resulting lists in the order specified by the outer operation. For an operation with output color m , the ordered lists substituted into the algebra inputs are followed by the ordered list of algebra inputs belonging to the operation substituted into the unique module input; the unique module input itself is carried along. The symmetric group action transports these linear orders along the induced bijections between the sets of algebra input positions. These rules are unital, associative and compatible with the symmetric group actions.

As in the commutative case, restricting operations to the color a recovers the underlying algebra operad, while restricting to the color m yields the trivial operad:

$$\mathcal{L}Mod_a \cong \mathcal{A}ssoc \quad \text{and} \quad \mathcal{L}Mod_m \cong \mathcal{T}riv.$$

Just as in the non-colored case, we may define algebras over operads in terms of operad morphisms:

Definition 12.2.10. Let $\mathcal{O}$ be a colored operad and let $\mathcal{C}$ be a symmetric monoidal category. An $\mathcal{O}$ -algebra in $\mathcal{C}$ is a morphism $\mathcal{O} \rightarrow \mathcal{M}_{\mathcal{C}}$ of colored operads.

For a non-colored operad, this agrees with the preceding definition because $(\mathcal{M}_{\mathcal{C}})_A = \mathcal{E}nd_{\mathcal{C}}(A)$ for every object $A \in \mathcal{C}$.

Example 12.2.11. For specific instances of $\mathcal{O}$, the $\mathcal{O}$ -algebras often have specialized names:

- For $\mathcal{O} = \mathcal{A}ssoc$ these are called *associative algebras*;
- For $\mathcal{O} = \mathcal{C}omm$ they are *commutative algebras*;
- For $\mathcal{O} = \mathcal{E}_k$ they are called $\mathcal{E}_k$ -algebras;
- For $\mathcal{O} = \mathcal{M}od$, an $\mathcal{O}$ -algebra is called a *module*. Restricting along the inclusion $\mathcal{C}omm \hookrightarrow \mathcal{M}od$ gives the *underlying commutative algebra* of the module, generically denoted A . Restricting along the inclusion $\mathcal{T}riv \hookrightarrow \mathcal{M}od$ gives the *underlying object*, generically denoted M .
- Similarly, $\mathcal{O} = \mathcal{L}Mod$ encodes *left modules*: pairs (A, M) where A is an *associative algebra* rather than a commutative one.

12.3 Encoding the coherences: the envelope of an operad

The notion of ∞ -operad we are after is a homotopical version of the notion of colored operad introduced in the previous section: the set $\mathcal{O}^\approx$ of colors should be replaced by an anima, and similarly the sets $\mathcal{O}((y_1, \dots, y_n); x)$ of multimorphisms should be replaced by anima. This leads to some subtleties regarding higher coherences: the unitality and associativity conditions required in Definition 12.2.1 should now become additional *data*, and this data should again satisfy further higher compatibilities. Writing down all the required structure explicitly would be impractical.

Fortunately, there is a way to circumvent this difficulty: the unitality, associativity and permutation laws required in an operad $\mathcal{O}$ can be reformulated in terms of a certain symmetric monoidal category $\mathcal{E}nv(\mathcal{O})$ called the *envelope* of the operad. This alternative description of operads can immediately be adapted to the ∞ -categorical setting, thus providing a definition of ∞ -operads. Let us therefore begin by describing this envelope construction in the classical setting.

We start with an observation regarding the formulation of colored operads: while we have formulated the notion of multimorphisms only for ordered tuples $(y_1, \dots, y_n) \in (\mathcal{O}^\approx)^n$, it is actually more natural to formulate them for *unordered* tuples:

Definition 12.3.1. An *unordered tuple* of the set $\mathcal{O}^\approx$ is a pair (I, x) where I is a finite set and where $x: I \rightarrow \mathcal{O}^\approx$ is a family of colors x_i indexed by I . We will often denote such unordered tuples by $\{x_i\}_{i \in I}$.

Given two unordered tuples $\{x_i\}_{i \in I}$ and $\{y_j\}_{j \in J}$, we can form their *unordered concatenation*

$$\{x_i\}_{i \in I} \sqcup \{y_j\}_{j \in J} := \{z_k\}_{k \in K}, \quad K := I \sqcup J, \quad z_i := x_i, \quad z_j := y_j.$$

More generally, if we are given a finite family $\{x_i^j\}_{i \in I_j}$ of unordered tuples for $j \in J$, then their unordered concatenation is $\{z_k\}_{k \in K}$ with $K := \bigsqcup_{j \in J} I_j$ and $z_{(j,i)} := x_i^j$.

Convention 12.3.2. Given a color $x \in \mathcal{O}^\approx$ and an unordered tuple $\{y_i\}_{i \in I}$, we define the set

$$\mathcal{O}(\{y_i\}_{i \in I}; x)$$

of multimorphisms from $\{y_i\}_{i \in I}$ to x by choosing a bijection $I \cong \{1, \dots, n\}$ and using it to order the inputs. The permutation isomorphisms canonically identify the sets obtained from different choices, so the notation is independent of the chosen ordering. Under this convention, the composition map in the operad $\mathcal{O}$ has the form

$$\circ: \mathcal{O}(\{y_i\}_{i \in I}; x) \times \prod_{i \in I} \mathcal{O}(\{z_j^i\}_{j \in J_i}; y_i) \rightarrow \mathcal{O}(\{z_j^i\}_{(i,j) \in \bigsqcup_{i \in I} J_i}; x).$$


The permutation isomorphism then says that for every bijection $\sigma: J \xrightarrow{\cong} I$, there is an isomorphism

$$\mathcal{O}(\{y_i\}_{i \in I}; x) \xrightarrow{\cong} \mathcal{O}(\{y_{\sigma(j)}\}_{j \in J}; x).$$

Henceforth, we shall use the unordered notation for colored operads.


Definition 12.3.3. Let $\mathcal{O}$ be a colored operad. We define a symmetric monoidal category $\text{Env}(\mathcal{O})$, called the *envelope* of $\mathcal{O}$.

- The objects of $\text{Env}(\mathcal{O})$ are unordered tuples $\{x_i\}_{i \in I}$ of colors of $\mathcal{O}$;
- A morphism $\Phi: \{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ in $\text{Env}(\mathcal{O})$ consists of a morphism of finite sets $f: I \rightarrow J$ together with a multimorphism $\varphi_j \in \mathcal{O}(\{x_i\}_{i \in I_j}; y_j)$ for every $j \in J$. Here we write $I_j := f^{-1}(j)$ for the preimage of j under f . We will visualize this data as follows:



Here the function f is indicated by the lines from left to right, so that $f(1) = f(4) = 3$, $f(2) = f(3) = 1$ and $f(5) = 2$.

- The identity of $\{x_i\}_{i \in I}$ is given by taking $f = \text{id}_I: I \rightarrow I$, and taking $\varphi_j = \text{id}_{x_j} \in \mathcal{O}(\{x_i\}_{i \in \{j\}}; x_j)$:



- The composition of $\Phi = (f, \varphi_j): \{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ and $\Psi = (g, \psi_k): \{y_j\}_{j \in J} \rightarrow \{z_k\}_{k \in K}$ is given by $\Psi \circ \Phi = (h, \xi_k)$ where $h := g \circ f: I \rightarrow K$ is the composition of g and f , and for every $k \in K$ the multimorphism $\xi_k \in \mathcal{O}(\{x_i\}_{i \in I_k}; z_k)$ is the image of $(\psi_k, (\varphi_j)_{j \in J_k})$ under the composition map


$$\circ: \mathcal{O}(\{y_j\}_{j \in J_k}; z_k) \times \prod_{j \in J_k} \mathcal{O}(\{x_i\}_{i \in I_j}; y_j) \rightarrow \mathcal{O}(\{x_i\}_{i \in I_k}; z_k);$$

here we are implicitly using that $I_k = (g \circ f)^{-1}(k) = f^{-1}(g^{-1}(k))$ may be written as the disjoint union of the sets $I_j = f^{-1}(j)$ for $j \in J_k = g^{-1}(k)$.

- The fact that composition in $\text{Env}(\mathcal{O})$ is unital and associative precisely amounts to the unitality and associativity conditions for colored operads.


The category $\text{Env}(\mathcal{O})$ admits a symmetric monoidal structure given by unordered concatenation of tuples.

The composition in $\text{Env}(\mathcal{O})$ may look somewhat difficult when written in formulas, but it becomes clear what one needs to do if one visually represents the morphisms using diagrams. For illustration, let us examine the composition of the following two morphisms in $\text{Env}(\mathcal{O})$:



First, we determine the composite map $h := g \circ f: \{1, \dots, 5\} \rightarrow \{1, \dots, 4\}$; in this case, this is given by $h(1) = h(2) = h(3) = h(4) = 2$ and $h(5) = 4$. Note that the preimages $h^{-1}(1)$ and $h^{-1}(3)$ are empty, so that ξ_1 and ξ_3 will be nullary operations. The multimorphism ξ_2 will have x_1, x_2, x_3 and x_4 as input colors, and is obtained by composing ψ_2 with (φ_1, φ_3) , where we need to remember which of the four inputs should go into φ_1 and which ones go into φ_3 . Finally, the multimorphism ξ_4 has only x_5 as input color, and is simply the composite $\psi_4 \circ \varphi_2$. The resulting composite morphism

can be illustrated as follows:



Note: we draw the permutation σ here to match the ordered tuple convention used in Section 12.2, since the nodes must be drawn in some fixed order on the page. When working with unordered tuples, the σ would not appear: the composition of multimorphisms simply remembers that its inputs are indexed by the disjoint union of $\{1, 3\}$ and $\{2, 4\}$ without fixing any order between them.

Example 12.3.4. The envelope $\text{Env}(\text{Comm})$ of the commutative operad is the category Fin of finite sets. The symmetric monoidal structure is given by disjoint union.

The envelope of $\mathcal{A}\text{ssoc}$, the functoriality of $\text{Env}(-)$, and its adjunction with the multimorphism-operad construction are treated in Exercises 12.9 to 12.11 at the end of the chapter. Since Comm is terminal and $\text{Env}(\text{Comm}) = \text{Fin}$, functoriality supplies every envelope with a symmetric monoidal functor

$$p: \text{Env}(\mathcal{O}) \longrightarrow \text{Env}(\text{Comm}) = \text{Fin}.$$

The nice feature about the envelope construction is that it allows us to formulate the unitality and associativity conditions for composition of multimorphisms in $\mathcal{O}$ in terms of unitality and associativity of morphisms in an actual category. In particular, this could potentially provide a convenient way to generalize the definition of operads to the ∞ -categorical setting. The main question we would then need to address is: what structure do we need on a symmetric monoidal category $\mathcal{C}$ to guarantee that it is of the form $\text{Env}(\mathcal{O})$ for some operad $\mathcal{O}$? We identify the three crucial properties that $\text{Env}(\mathcal{O})$ has:

- (1) The colors of $\mathcal{O}$ are precisely the objects that are sent to a one-point set $\{i\}$ under the symmetric monoidal functor $p: \text{Env}(\mathcal{O}) \rightarrow \text{Fin}$.
- (2) Given an object $X \in \text{Env}(\mathcal{O})$ with $p(X) = I$, we may uniquely write X as an I -tuple $\{x_i\}_{i \in I}$ for some colors $x_i \in p^{-1}(\{i\})$. Similarly, the morphisms in the fiber $p^{-1}(I)$ are I -tuples of maps $\{f_i: x_i \rightarrow y_i\}_{i \in I}$. In other words, the fiber over I is the I -fold product of the fiber over the one-point set.
- (3) The morphisms into an arbitrary tuple $\{y_j\}_{j \in J}$ are completely determined by morphisms into 1-tuples: If we consider two unordered tuples $\{x_i\}_{i \in I}$ and $\{y_j\}_{j \in J}$, then the fiber of the induced map

$$\text{Hom}_{\text{Env}(\mathcal{O})}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J}) \xrightarrow{p} \text{Hom}_{\text{Fin}}(I, J)$$

over some map $f: I \rightarrow J$ is given by the product

$$\prod_{j \in J} \text{Hom}_{\text{Env}(O)}(\{x_i\}_{i \in I_j}, y_j).$$

The first property explains how to recover the colors from p , while the remaining two characterize the morphisms and tensor products in the envelope. One can verify that a symmetric monoidal category C equipped with a symmetric monoidal functor $p: C \rightarrow \text{Fin}$ is of the form $\text{Env}(O)$ for some O if and only if it satisfies the corresponding analogues of conditions (2) and (3):

(2') For every finite set I , the tensor product in C defines an equivalence

$$\prod_{i \in I} C_{\{i\}} \xrightarrow{\sim} C_I,$$

where C_I is the fiber of p over $I \in \text{Fin}$;

(3') For every finite collection $(I_j)_{j \in J}$ of finite sets and every collection of objects $X_j \in p^{-1}(I_j)$ and $y_j \in p^{-1}(\{j\})$, the commutative square

$$\begin{array}{ccc} \prod_{j \in J} \text{Hom}_C(X_j, y_j) & \xrightarrow{\otimes_{j \in J}} & \text{Hom}_C(\bigotimes_{j \in J} X_j, \bigotimes_{j \in J} y_j) \\ \downarrow & & \downarrow p \\ * & \xrightarrow{\quad \quad \quad} & \text{Hom}_{\text{Fin}}(\bigsqcup_{j \in J} I_j, J) \end{array}$$

is a pullback square, where the bottom map hits the map $\bigsqcup_{j \in J} I_j \rightarrow \bigsqcup_{j \in J} * = J$ induced by the maps $I_j \rightarrow *$.

Exercise 12.3.5. Given a symmetric monoidal category C with $p: C \rightarrow \text{Fin}$ satisfying (2') and (3'), construct a colored operad O and a symmetric monoidal equivalence $C \xrightarrow{\sim} \text{Env}(O)$ over Fin .

12.4 Writing concatenations as products

The characterization of envelopes of operads obtained in the previous section is neat, and immediately generalizes to the ∞ -categorical setting. Nevertheless, this is not the definition of ∞ -operads that usually appears in the literature. One inconvenience about this characterization is that it requires us to equip an ∞ -category with a symmetric monoidal structure, which is a task substantially more involved than for classical categories since we can no longer ‘just write it down by hand’. It would be convenient if we could avoid reliance on symmetric monoidal structures.

Fortunately, this is possible: there is a variation of the envelope construction in which the concatenation of tuples becomes the *categorical product*, and hence does not need to be recorded as additional data. We have already seen this idea in action in our definition of commutative monoids in Chapter 5. Recall from Definition 8.1.3 that a commutative algebra in a symmetric monoidal ∞ -category C is encoded by a symmetric monoidal functor

$$\text{Fin} \rightarrow C,$$

so its definition uses the given tensor product on C . By contrast, if C admits finite products, a commutative monoid in C is encoded by a product-preserving functor

$$\text{Span}(\text{Fin}) \rightarrow C.$$

The additional left-pointing morphisms ensure that disjoint union in Fin becomes the categorical product in $\text{Span}(\text{Fin})$, so no separately specified symmetric monoidal structure on C is needed.

We will apply the same mechanism to the envelope $\text{Env}(\mathcal{O})$: by adjoining left-pointing morphisms, we obtain a variant of $\text{Env}(\mathcal{O})$ in which concatenation of tuples is the categorical product, and therefore need not be encoded separately. Let us emphasize that this maneuver is merely about encoding the operad structure; it does not impose a cartesian tensor product on a category in which we later consider $\mathcal{O}$ -algebras. Just as in the case of the span category $\text{Span}(\text{Fin})$, the construction naturally produces a $(2, 1)$ -category rather than a $(1, 1)$ -category, since composition is defined using pullbacks and is therefore associative only up to canonical isomorphism.

Construction 12.4.1. Let $\mathcal{O}$ be a colored operad. We describe a $(2, 1)$ -category $\mathcal{O}^\otimes$ as follows:

- The objects of $\mathcal{O}^\otimes$ are unordered tuples $\{x_i\}_{i \in I}$;
- A morphism $\{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ in $\mathcal{O}^\otimes$ consists of a span

$$I \xleftarrow{f} K \xrightarrow{g} J$$

of finite sets together with a multimorphism $\varphi_j \in \mathcal{O}(\{x_{f(k)}\}_{k \in K_j}; y_j)$ for every $j \in J$; here we write $K_j := g^{-1}(j)$ for the preimage of j under g . Together with the canonical notion of isomorphisms between such labeled spans, these form a groupoid $\text{Hom}_{\mathcal{O}^\otimes}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J})$.

- The identity of $\{x_i\}_{i \in I}$ is given by taking $f = g = \text{id}_I: I \rightarrow I$, and taking $\varphi_j = \text{id}_{x_j} \in \mathcal{O}(\{x_i\}_{i \in \{j\}}; x_j)$.
- The composition of $(f, g, \varphi_j): \{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ and $(h, k, \psi_l): \{y_j\}_{j \in J} \rightarrow \{z_l\}_{l \in L}$ is given by $(f \circ \text{pr}_K, k \circ \text{pr}_M, \xi_l)$ where pr_K and pr_M are the projection maps in the following pullback diagram

$$\begin{array}{ccccc} & & K \times_J M & & \\ & \swarrow & \downarrow \vee & \searrow & \\ f \circ \text{pr}_K & & & & k \circ \text{pr}_M \\ & \swarrow & & \searrow & \\ & K & & M & \\ & \swarrow & \downarrow g & \downarrow h & \searrow k \\ I & & J & & L \end{array}$$

and for every $l \in L$ the multimorphism $\xi_l \in \mathcal{O}(\{x_{f(u)}\}_{(u,m) \in (K \times_J M)_l}; z_l)$ is the image of $(\psi_l, (\varphi_{h(m)})_{m \in M_l})$ under the composition map


$$\circ: \mathcal{O}(\{y_{h(m)}\}_{m \in M_l}; z_l) \times \prod_{m \in M_l} \mathcal{O}(\{x_{f(u)}\}_{u \in K_{h(m)}}; y_{h(m)}) \rightarrow \mathcal{O}(\{x_{f(u)}\}_{(u,m) \in (K \times_J M)_l}; z_l).$$

- The operad axioms make composition unital and associative up to the canonical isomorphisms induced by pullbacks of spans.

Unordered concatenation gives finite products in $O^\otimes$; the details are included in the following exercise.

Exercise 12.4.2. Convince yourself that $O^\otimes$ is a well-defined (2,1)-category and that it is functorial in O . Argue that it admits finite products given by the unordered concatenation of tuples.

This time, morphisms in $O^\otimes$ may be visualized as follows:



The first part of the data consists of two maps $f: \{1, \dots, 5\} \rightarrow \{1, \dots, 4\}$ and $g: \{1, \dots, 5\} \rightarrow \{1, 2, 3\}$; in this case f is given by $f(1) = f(5) = 2$, $f(2) = 1$, $f(3) = 3$ and $f(4) = 4$, while g is given by $g(2) = g(3) = 1$, $g(5) = 2$ and $g(1) = g(4) = 3$. In particular, the preimages of 1, 2 and 3 under g are $\{2, 3\}$, $\{5\}$ and $\{1, 4\}$, respectively. The second part of the data consists of multimorphisms $\varphi_1: (x_{f(2)}, x_{f(3)}) \rightarrow y_1$, $\varphi_2: (x_{f(5)}) \rightarrow y_2$ and $\varphi_3: (x_{f(1)}, x_{f(4)}) \rightarrow y_3$.

Exercise 12.4.3. Make a visualization of how composition in $O^\otimes$ works, generalizing the picture for $\text{Env}(O)$ by adding in the left-pointing arrows. Explain how the process of forming the pullback $K \times_J M$ is reflected in the visualization.


For $O = \text{Comm}$ we get $\text{Comm}^\otimes = \text{Span}(\text{Fin})$. Since the construction of $O^\otimes$ is functorial in O , we obtain a functor

$$p: O^\otimes \rightarrow \text{Comm}^\otimes = \text{Span}(\text{Fin})$$

which preserves finite products. As in the previous section, we may ask: can we characterize those (2, 1)-categories over $\text{Span}(\text{Fin})$ that are of the form $O^\otimes$ for some operad? The fiber decomposition from (2') remains valid, now in terms of categorical products. The product universal property decomposes morphisms according to the factors of their target, but it does not by itself isolate the corresponding factors in the source. Moreover, the morphisms over right-pointing spans do not determine the morphisms over a general span $I \leftarrow K \rightarrow J$. To reduce the general case to the right-pointing case, we need distinguished lifts of left-pointing spans.

To reduce the general case to the special case, consider an object $\{x_i\}_{i \in I}$ of $O^\otimes$ and consider a morphism $f: K \rightarrow I$ of finite sets. We may then form the morphism $\tilde{f}: \{x_i\}_{i \in I} \rightarrow \{x_{f(k)}\}_{k \in K}$ in $O^\otimes$ whose underlying span of finite sets is $I \xleftarrow{f} K \xrightarrow{\text{id}} K$ and whose multimorphisms $\varphi_k \in O(\{x_{f(k)}\}, x_{f(k)})$ are the identity morphisms $\text{id}_{x_{f(k)}}$. This morphism $\tilde{f}$ may be visualized as

follows:



This morphism has the following special feature, which is the shadow of the general notion of a p -cocartesian morphism that we will meet in Definition 23.1.1:

(3'') Consider objects $\{x_i\}_{i \in I}$ and $\{y_j\}_{j \in J}$ in $\mathcal{O}^\otimes$, and consider a span $I \xleftarrow{f} K \xrightarrow{g} J$. Given a morphism $\tilde{h}: \{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ in $\mathcal{O}^\otimes$ lifting¹⁸ the given span, there exists a unique morphism $\tilde{g}: \{x_{f(k)}\}_{k \in K} \rightarrow \{y_j\}_{j \in J}$ which lifts the span $K \xleftarrow{=} K \xrightarrow{g} J$ and satisfies $\tilde{h} = \tilde{g} \circ \tilde{f}$:

$$\begin{array}{ccc}
 & \{x_{f(k)}\}_{k \in K} & \\
 \tilde{f} \nearrow & & \searrow \exists! \tilde{g} \\
 \{x_i\}_{i \in I} & \xrightarrow{\tilde{h}} & \{y_j\}_{j \in J}
 \end{array}$$

Indeed, the lift $\tilde{h}$ is encoded by certain multimorphisms $\varphi_j \in \mathcal{O}(\{x_{f(k)}\}_{k \in K_j}; y_j)$, and this is precisely the data we need for defining the lift $\tilde{g}$. The full cocartesian condition is the pullback-square condition appearing as condition (3) in Definition 12.4.5.

In a sense, we see that these morphisms $\tilde{f}$ in $\mathcal{O}^\otimes$ do not really contribute very much apart from encoding the categorical product structure. These morphisms, and more generally the cocartesian morphisms lying over left-pointing spans, are called *inert morphisms*. In contrast, the morphisms in $\mathcal{O}^\otimes$ living over *right*-pointing spans are the ones that encode the actual operad structure, and are called the *active morphisms*.

Exercise 12.4.4. Regarding $\{x_i\}_{i \in I}$ as the I -indexed product of the objects x_i and similarly for $\{x_{f(k)}\}_{k \in K}$ (as in Exercise 12.4.2), show that the morphism $\tilde{f}: \{x_i\}_{i \in I} \rightarrow \{x_{f(k)}\}_{k \in K}$ is the map whose k -th component is the projection map onto $x_{f(k)}$.

We have now finally achieved what we were after: the entire data of the operad $\mathcal{O}$ is encoded by the $(2, 1)$ -category $\mathcal{O}^\otimes$ together with its functor to $\text{Span}(\text{Fin})$, and the properties we singled out tell us which such functors are obtained via this procedure. These properties still make sense in the ∞ -categorical setting, and we will use them in our definition of ∞ -operads:

Definition 12.4.5. An ∞ -operad is a pair $\mathcal{O} = (\mathcal{O}^\otimes, p_\mathcal{O})$ consisting of an ∞ -category $\mathcal{O}^\otimes$ equipped with a functor $p_\mathcal{O}: \mathcal{O}^\otimes \rightarrow \text{Span}(\text{Fin})$ satisfying the following conditions:

- (1) The ∞ -category $\mathcal{O}^\otimes$ admits finite products, and the functor $p_\mathcal{O}$ preserves finite products;

¹⁸This means that the image of $\tilde{h}$ under p is the given span $I \xleftarrow{f} K \xrightarrow{g} J$.

(2) For every finite set I , the product in $O^\otimes$ defines an equivalence

$$\prod_{i \in I} O_{\{i\}}^\otimes \xrightarrow{\sim} O_I^\otimes.$$

Here we write $O_I^\otimes$ for the fiber of p_O over the set I .

(3) For a morphism $f: J \rightarrow I$ in $\mathbf{Fin}$ and objects $X_i \in O^\otimes$, the map $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ whose j -th component is the projection to $X_{f(j)}$ is p_O -cocartesian, in the sense that for every $Y \in O^\otimes$ the following square is a pullback square:

$$\begin{array}{ccc} \mathrm{Hom}_{O^\otimes}(\prod_{j \in J} X_{f(j)}, Y) & \xrightarrow{- \circ \tilde{f}} & \mathrm{Hom}_{O^\otimes}(\prod_{i \in I} X_i, Y) \\ p_O \downarrow & & \downarrow p_O \\ \mathrm{Hom}_{\mathrm{Span}(\mathbf{Fin})}(p_O(\prod_{j \in J} X_{f(j)}), p_O(Y)) & \xrightarrow{- \circ p_O(\tilde{f})} & \mathrm{Hom}_{\mathrm{Span}(\mathbf{Fin})}(p_O(\prod_{i \in I} X_i), p_O(Y)). \end{array}$$

Remark 12.4.6. Our definition of ∞ -operad is a priori different from that of Lurie [Lur17], who instead works with functors p_O whose target is the category $\mathbf{Fin}_*$ of finite pointed sets. To relate the two approaches, recall from Lemma 5.3.8 that $\mathbf{Fin}_*$ is equivalent to the subcategory $\mathrm{Span}(\mathbf{Fin}, \mathrm{inj}, \mathrm{all})$ of $\mathrm{Span}(\mathbf{Fin})$ in which the left-pointing morphisms are required to be injective. We will show in Section 17.4 below that pullback along the resulting inclusion $\mathbf{Fin}_* \hookrightarrow \mathrm{Span}(\mathbf{Fin})$ induces an equivalence between our ∞ -category of ∞ -operads and Lurie's one.

One advantage of working over $\mathbf{Fin}_*$ is that it is a strict 1-category. In Lurie's model, the category $O^\otimes$ from Construction 12.4.1 also becomes a strict 1-category, and can therefore be directly imported into the world of ∞ -categories, showing that every colored operad gives rise to an ∞ -operad. While the same can in principle be done at the level of $(2, 1)$ -categories, we will not develop the theory of $(2, 1)$ -categories in this book, and instead import classical operads via Lurie's model; see Proposition 17.4.9.

On the other hand, there are a variety of reasons to work over $\mathrm{Span}(\mathbf{Fin})$ rather than $\mathbf{Fin}_*$. First, the span approach makes clear the orthogonal roles played by the backwards and forwards maps; this separation is less visible in $\mathbf{Fin}_*$. Second, allowing only injective backwards maps breaks the property that concatenation of unordered tuples is a categorical product, so conditions on $O^\otimes$ need to be formulated in a slightly more complicated way. And third, the span perspective generalizes more easily to other settings, making it easier to connect the theory developed here to concepts like Mackey functors, normed ∞ -categories and six-functor formalisms. For example, it allows us to treat the theory of (co)cartesian monoidal structures as an instance of Barwick's more general unfurling construction in Chapter 15.

Remark 12.4.7 (Other models for ∞ -operads). The span-based model used in this book and Lurie's model over $\mathbf{Fin}_*$ from Section 17.4 are not the only available descriptions of ∞ -operads. In the *dendroidal* approach, introduced by Moerdijk and Weiss [MW07], the simplex category is replaced by a category of trees. The presentation of ∞ -operads by complete dendroidal Segal anima is due to Cisinski and Moerdijk [CM13]. A direct comparison with Lurie's model is given by Hinich and Moerdijk [HM24b]; see [HM22] for a comprehensive account of dendroidal homotopy theory.

The classical description of one-colored operads as associative algebras in symmetric sequences also admits a homotopy-coherent generalization due to Haugseng [Hau22]. More generally, this construction treats varying anima of colors by means of a double ∞ -category of symmetric collections; see

[Hau23a] for a general introduction to ∞ -operads. A related viewpoint describes pinned ∞ -operads by analytic monads on slices $\mathbf{An}/_S$, with the anima S of colors allowed to vary [GHK22; Hau23b]. This description is compared directly with both Lurie’s model and the span-based model through their Lawvere theories by Haugseng [Hau23b]. More general Segal-type formalisms, notably the theory of algebraic patterns developed by Chu and Haugseng [CH21], provide a common framework for these and many similar homotopy-coherent algebraic structures. The general theory of envelopes for algebraic patterns developed by Barkan, Haugseng, and Steinebrunner [BHS22] contains our construction of operadic envelopes in Section 17.3 as a special case. We will not otherwise use the alternative presentations discussed in this remark.

Before studying ∞ -operads in more detail, we will give a proper introduction to span ∞ -categories and their basic properties in the next chapter. We return to ∞ -operads and their algebras in Chapter 14.

Exercises

Exercise 12.1 (Equivariance conditions). In the definition of an operad, we have not specified how the permutation actions on $O(n)$ interact with the composition structure. In this exercise, you will formulate the missing conditions.

Let O be a non-colored operad, and consider operations $P \in O(k)$ and $Q_i \in O(n_i)$ for $i = 1, \dots, k$.

- (1) Consider a permutation $\sigma \in \Sigma_k$. Using the motivating example of the endomorphism operad $O = \mathcal{E}nd_C(A)$, write down a rule for expressing the operation

$$(P\sigma)(Q_1, \dots, Q_k)$$

as an operation of the form $P(R_1, \dots, R_k)\tau$ for certain operations R_i and a certain permutation $\tau \in \Sigma_n$, where $n := \sum_{i=1}^k n_i$.

- (2) Consider permutations $\sigma_i \in \Sigma_{n_i}$ for all $i = 1, \dots, k$. Again using $O = \mathcal{E}nd_C(A)$ as motivating example, write down a rule for expressing the operation

$$P(Q_1\sigma_1, \dots, Q_k\sigma_k)$$

as an operation of the form $P(Q_1, \dots, Q_k)\tau$ for a certain permutation $\tau \in \Sigma_n$, where $n := \sum_{i=1}^k n_i$.

- (3) Formulate the corresponding equivariance conditions for a colored operad. In doing so, keep track of the reindexing of the input colors under each permutation.

Exercise 12.2 (The associative operad). Write out the composition maps and the Σ_n -actions in the case of the associative operad $\mathcal{A}ssoc$.

Exercise 12.3 (Lie operad). Look up the definition of the Lie operad $\mathcal{L}ie$ over a given field K . Give a precise description of the vector spaces $\mathcal{L}ie(n)$ for all n and of the composition structure of the operad.

Bonus question: Look up the definition of an *enriched* operad, and explain in what sense the Lie operad over the field K is enriched in K -vector spaces. Also explain why its algebras in the symmetric monoidal category $\mathbf{Vect}_K$ are precisely the Lie algebras.

Exercise 12.4. Let C be a symmetric monoidal category. Show that:

- $\mathcal{Triv}$ -algebras are simply objects of C ;
- $\mathcal{E}_0$ -algebras are ‘pointed objects’ of C , i.e., objects A equipped with a ‘unit map’ $\eta: \mathbb{1} \rightarrow A$.
- $\mathcal{Comm}$ -algebras are objects A equipped with maps $\eta: \mathbb{1} \rightarrow A$ and $m: A \otimes A \rightarrow A$ which are unital, associative and commutative.
- $\mathcal{Assoc}$ -algebras are similar but without commutativity.

Exercise 12.5 (Universal property of trivial operads). Let $\mathcal{O}$ be a colored operad. Show that the colors of $\mathcal{O}$ together with the sets $\mathcal{O}((y);x)$ of unary operations form a 1-category $\mathcal{O}_{\langle 1 \rangle}$, called the *underlying category* of $\mathcal{O}$, where $\langle 1 \rangle := \{1\}$. Show furthermore that for every 1-category C , restriction to the unary operations defines a bijection between the morphisms of colored operads $\mathcal{Triv}_C \rightarrow \mathcal{O}$ and the functors $C \rightarrow \mathcal{O}_{\langle 1 \rangle}$, where $\mathcal{Triv}_C$ is the trivial operad of Example 12.2.6.

Exercise 12.6 (Little cubes operad). Spend some time understanding the 1-dimensional little cubes operad $\mathcal{E}_1$. Explain how we may think of $\mathcal{E}_1$ as a *topological* operad, i.e., each $\mathcal{E}_1(n)$ is a topological space and all the structure maps are continuous. Show that there is an operad morphism $\mathcal{E}_1 \rightarrow \mathcal{Assoc}$ which induces homotopy equivalences $\mathcal{E}_1(n) \rightarrow \mathcal{Assoc}(n)$ for all $n \geq 0$.

Exercise 12.7 (Left module operad). Verify that the composition and permutation rules described in Example 12.2.9 satisfy the colored-operad axioms.

Exercise 12.8. Let $\mathcal{O}$ be a colored operad, and let C be a symmetric monoidal category. Recall that we defined an $\mathcal{O}$ -algebra as a morphism of colored operads $A: \mathcal{O} \rightarrow \mathcal{M}_C$. Spell out in explicit terms all the data contained in such an $\mathcal{O}$ -algebra. In the special case of $\mathcal{O} = \mathcal{LMod}$, verify that this structure precisely encodes a pair (A, M) of an associative algebra A in C together with a left module M over it.

Remark: Saying ‘left’ versus ‘right’ is just a convention; the same operad also encodes right actions.

Exercise 12.9. Compute the envelope $\text{Env}(\mathcal{Assoc})$ of the associative operad, and show that it may be described as follows:

- The objects are finite sets I ;
- The morphisms $I \rightarrow J$ are tuples $(f, \preceq_j)$ consisting of a morphism $f: I \rightarrow J$ of finite sets and a collection of linear orders $\preceq_j$ on each of the preimages $f^{-1}(j)$ of f ;
- The symmetric monoidal structure is disjoint union.

Exercise 12.10. Show that the assignment $\mathcal{O} \mapsto \text{Env}(\mathcal{O})$ is functorial: every morphism of operads $\mathcal{O} \rightarrow \mathcal{P}$ induces a symmetric monoidal functor $\text{Env}(\mathcal{O}) \rightarrow \text{Env}(\mathcal{P})$. Deduce from the terminal morphism $\mathcal{O} \rightarrow \mathcal{Comm}$ the symmetric monoidal functor $p: \text{Env}(\mathcal{O}) \rightarrow \text{Fin}$ used above.

Exercise 12.11. Show that the assignment $\mathcal{O} \mapsto \text{Env}(\mathcal{O})$ is left adjoint to the assignment $C \mapsto \mathcal{M}_C$: for an operad $\mathcal{O}$ and a symmetric monoidal category C , there is a bijection between operad morphisms $\mathcal{O} \rightarrow \mathcal{M}_C$ and symmetric monoidal functors $\text{Env}(\mathcal{O}) \rightarrow C$.

13 Span categories

In the approach to higher algebra taken in this book, a prominent role is played by the notion of a *span category*. In recent years, this notion has gained a lot of prominence among researchers, as it provides a convenient way to encode algebraic structures in a variety of settings, like those of *spectral Mackey functors* [Bar17], *normed ∞ -categories* [BH21] or *six-functor formalisms* [Sch25; HM24a]. So far, the use of span categories in higher algebra has not yet become mainstream, which is probably in large part due to the fact that the main textbook on higher algebra, [Lur17], does not use them. For this reason, we will dedicate some time in this chapter to a proper introduction of span categories.

Let C be an ∞ -category with pullbacks, and let C_L and C_R be collections of morphisms of C closed under composition and pullback. Then the *span category* $\text{Span}_{L,R}(C)$ associated with this triple is an ∞ -category whose objects are the objects of C , while the morphisms from X to Y are given by *spans*

$$\begin{array}{ccc} & U & \\ f \swarrow & & \searrow g \\ X & & Y, \end{array}$$

where $f \in C_L$ and $g \in C_R$. The identity morphism is the *identity span* $X \xleftarrow{\text{id}_X} X \xrightarrow{\text{id}_X} X$. The composition of a span $X \leftarrow U \rightarrow Y$ with a span $Y \leftarrow V \rightarrow Z$ is given by forming pullbacks:

$$\begin{array}{ccccc} & & U \times_Y V & & \\ & \swarrow & \downarrow \smile & \searrow & \\ & U & & V & \\ \swarrow & & \downarrow & & \searrow \\ X & & Y & & Z. \end{array}$$

In this chapter, we will give a formal definition of $\text{Span}_{L,R}(C)$, following the approach of Barwick [Bar17] and Haugseng, Hebestreit, Linskens, and Nuiten [HHLN23b]. We will further discuss the existence of products and coproducts in $\text{Span}_{L,R}(C)$.

13.1 Constructing span categories

So far, we could reduce all of the constructions of ∞ -categories we encountered to a handful of elementary constructions, like pullbacks, subcategories and functor categories. The situation for span categories is different: while the heuristic description given before tells us what the objects and morphisms should be, it is not immediately clear how to assemble these into an ∞ -category.

The technique we will use is that of *complete Segal anima*, discussed in detail in Chapter 24. Let us briefly recall the main idea. Given an ∞ -category C , we may consider for every $n \geq 0$ the anima $\text{Map}([n], C)$ of ‘strings of n composable morphisms in C ’. Sending $[n]$ to $\text{Map}([n], C)$ for all n provides a simplicial anima $N(C): \Delta^{\text{op}} \rightarrow \text{An}$, called the *nerve of C* . A major insight due to Rezk is that we may use this assignment to identify ∞ -categories with suitable simplicial anima. More precisely, the resulting nerve functor

$$N: \text{Cat}_{\infty} \hookrightarrow \text{sAn}$$

from ∞ -categories to simplicial anima is fully faithful. The essential image of this inclusion is given by the *complete Segal anima*, a class of simplicial anima satisfying analogues of the Segal Axiom (Proposition 1.4.5) and the Rezk Axiom (Axiom F).

For the purpose of constructing span categories, we will use this process in reverse: if we manage to write down a simplicial anima $\text{NSpan}_{L,R}(C): \Delta^{\text{op}} \rightarrow \text{An}$ encoding strings of n composable morphisms in the span category for all n , and if we manage to verify that it is a complete Segal anima, then we have managed to construct our desired ∞ -category $\text{Span}_{L,R}(C)$. This leads to the next question: what data should be contained in a functor $[n] \rightarrow \text{Span}_{L,R}(C)$? Such a functor should first of all encode n composable spans. Furthermore, it should provide a compatible family of choices for their compositions. Given that we want composition in the span category to be given by pullback, we conclude that $\text{NSpan}_{L,R}(C)_n$ should be the anima of diagrams in C of the following form:

$$\begin{array}{ccccccc}
 & & & X_{0n} & & & \\
 & & \swarrow & \downarrow & \searrow & & \\
 & & \dots & & X_{1n} & & \\
 & \swarrow & \downarrow & \searrow & \swarrow & \downarrow & \searrow \\
 & X_{02} & & \dots & & \dots & \\
 \swarrow & \downarrow & \searrow & \swarrow & \downarrow & \searrow & \swarrow \\
 X_{01} & & X_{12} & & \dots & & \dots \\
 \swarrow & \downarrow & \searrow & \swarrow & \downarrow & \searrow & \swarrow \\
 X_{00} & & X_{11} & & X_{22} & & \dots & & X_{nn}.
 \end{array}$$

Here all the left-pointing morphisms should be in the given class C_L , all the right-pointing morphisms should be in C_R , and all squares should be pullback squares.

In the remainder of this section, we will make the construction of the simplicial anima $\text{NSpan}_{L,R}(C)$ more precise, and verify that it is indeed a complete Segal anima. We will also prove various basic properties about the resulting ∞ -categories of spans. Our exposition closely follows that of Haugseng, Hebestreit, Linskens, and Nuiten [HHLN23b].

Definition 13.1.1. A subcategory $C' \subseteq C$ is called *wide* if it contains all objects of C .

We will usually think of a wide subcategory C' as specifying a collection of morphisms in C closed under composition. Since a collection of morphisms is a union of components of $\text{Map}([1], C)$, every wide subcategory contains all isomorphisms of C .

Definition 13.1.2 ([Bar17]). An *adequate triple* (C, C_L, C_R) consists of an ∞ -category C equipped with two wide subcategories $C_L, C_R \subseteq C$ satisfying the condition that for every morphism $l: X \rightarrow Y$

in C_L and $r: Y' \rightarrow Y$ in C_R , there exists a pullback square of the form

$$\begin{array}{ccc} X' & \xrightarrow{r'} & X \\ l' \downarrow & \lrcorner & \downarrow l \\ Y' & \xrightarrow{r} & Y, \end{array} \quad (13.1)$$

and we have $l' \in C_L$ and $r' \in C_R$. We refer to C_L as the *class of backwards morphisms* and to C_R as the *class of forward morphisms*.

Given another adequate triple (D, D_L, D_R) , a *morphism of adequate triples* is a functor $F: C \rightarrow D$ that sends C_L to D_L and C_R to D_R , and sends every pullback square of the form (13.1), with $l \in C_L$ and $r \in C_R$, to a pullback square in D . We denote by

$$\mathbf{AdTrip} \quad \subseteq \quad \mathbf{Fun}(\perp, \mathbf{Cat}_\infty)$$

the non-full subcategory spanned by those diagrams $C_L \hookrightarrow C \leftarrow C_R$ in Cat_∞ that correspond to adequate triples (C, C_L, C_R) and those morphisms that correspond to morphisms of adequate triples.

Example 13.1.3. For any wide subcategory C' of an ∞ -category C , we have the adequate triples $(C, C', C^\simeq)$ and $(C, C^\simeq, C')$. In particular, for any ∞ -category C we always have the adequate triples $(C, C, C^\simeq)$ and $(C, C^\simeq, C)$.

Example 13.1.4. If C admits all pullbacks, we have the adequate triple (C, C, C) where we take all maps to be both forward and backward maps. This gives rise to a fully faithful embedding

$$\mathrm{Cat}_\infty^{\mathrm{pb}} \hookrightarrow \mathrm{AdTrip},$$

where $\text{Cat}_\infty^{\text{pb}} \subseteq \text{Cat}_\infty$ is the subcategory spanned by those ∞ -categories which admit pullbacks, and those functors which preserve pullbacks.

Example 13.1.5. If (C, C_L, C_R) is an adequate triple, then so is (C, C_R, C_L) . The resulting endomorphism of AdTrip will be denoted

$$(-)_{\text{rev}}: \text{AdTrip} \rightarrow \text{AdTrip}.$$

We will now introduce the indexing diagrams that classify the functors $[n] \rightarrow \text{Span}_{L,R}(C)$.

Definition 13.1.6 (Twisted arrow category). For $n \geq 0$, let $\text{Tw}^r([n])$ be the partially ordered set of tuples (i, j) with $0 \leq i \leq j \leq n$, where the partial order is given by

$$(i, j) \leq (k, l) \quad \text{if and only if} \quad i \leq k \quad \text{and} \quad l \leq j.$$

We will generally denote the pair (i, j) by $(i \leq j)$. The poset looks as follows:

Given a morphism of posets $\varphi: [n] \rightarrow [m]$, we obtain a map $\text{Tw}^r(\varphi): \text{Tw}^r([n]) \rightarrow \text{Tw}^r([m])$ by sending $(i \leq j)$ to $(\varphi(i) \leq \varphi(j))$. This assignment is clearly functorial, resulting in a functor

$$\text{Tw}^r: \mathbf{\Delta} \rightarrow \text{Poset} \subseteq \text{Cat} \subseteq \text{Cat}_\infty.$$

Exercise 13.1.7. Show that the functors $s: \text{Tw}^r([n]) \rightarrow [n], (i \leq j) \mapsto i$ and $t: \text{Tw}^r([n]) \rightarrow [n]^{\text{op}}, (i \leq j) \mapsto j$ are cartesian fibrations.

Lemma 13.1.8. For integers $0 \leq i \leq k \leq l \leq j \leq n$, the commutative square

$$\begin{array}{ccc} & (i \leq j) & \\ \swarrow & & \searrow \\ (i \leq l) & & (k \leq j) \\ \searrow & & \swarrow \\ & (k \leq l) & \end{array}$$

is a pullback square in $\text{Tw}^r([n])$.

Proof. Given $0 \leq i' \leq j' \leq n$ we need to show that we have $(i', j') \leq (i, j)$ if and only if $(i', j') \leq (i, l)$ and $(i', j') \leq (k, j)$. This is clear: in both cases this amounts to asking that for both of the relations $i' \leq i$ and $j \leq j'$. $\square$

Corollary 13.1.9. The poset $\text{Tw}^r([n])$ is an adequate triple, with $\text{Tw}^r([n])_L$ consisting of morphisms of the form $(i \leq j) \leq (i \leq l)$ and $\text{Tw}^r([n])_R$ consisting of morphisms of the form $(i \leq j) \leq (k \leq j)$.

Proof. The two classes are closed under composition, and every cospan consisting of a morphism in $\text{Tw}^r([n])_L$ and a morphism in $\text{Tw}^r([n])_R$ has the form treated in Lemma 13.1.8. $\square$

From now on, we will equip $\text{Tw}^r([n])$ with the adequate triple structure from Corollary 13.1.9. Note that a morphism of posets $\varphi: [n] \rightarrow [m]$ induces a morphism of adequate triples $\text{Tw}^r(\varphi): \text{Tw}^r([n]) \rightarrow \text{Tw}^r([m])$, resulting in a functor

$$\text{Tw}^r: \mathbf{\Delta} \rightarrow \text{AdTrip}.$$

Observe that the diagrams in C we drew in our informal description of the functors $[n] \rightarrow \text{Span}_{L,R}(C)$ given in the introduction to this chapter are precisely the morphisms of adequate triples $\text{Tw}^r([n]) \rightarrow (C, C_L, C_R)$. This leads to the following definition:

Construction 13.1.10. Let (C, C_L, C_R) be an adequate triple. We define the simplicial anima $\text{NSpan}_{L,R}(C) \in \text{sAn}$ as the following composite:

$$\mathbf{\Delta}^{\text{op}} \xrightarrow{\text{Tw}^r} \text{AdTrip}^{\text{op}} \xrightarrow{\text{Hom}_{\text{AdTrip}}(-, (C, C_L, C_R))} \text{An}.$$

In other words, for $n \geq 0$ we consider the subanima


$$\text{NSpan}_{L,R}(C)_n \subseteq \text{Map}(\text{Tw}^r([n]), C)$$

consisting of the morphisms of adequate triples. This construction is natural in the adequate triple, hence defines a functor

$$\text{NSpan}: \text{AdTrip} \rightarrow \text{sAn}.$$

Proposition 13.1.11 (Barwick [Bar17, Proposition 5.6], Haugseng, Hebestreit, Linskens, and Nuiten [HHLN23b, Theorem 2.1.3]). *For every adequate triple (C, C_L, C_R) the simplicial anima $\mathbf{NSpan}_{L,R}(C)$ is a complete Segal anima.*

Proof. We start by proving the Segal condition. Let us write $X := \mathbf{NSpan}_{L,R}(C)$ for ease of notation. For every $n \geq 0$, consider the subposet $J_n \subseteq \mathbf{Tw}^r([n])$ spanned by the objects $(i \leq j)$ with $j \leq i + 1$:



Note that a functor $\mathbf{Tw}^r([n]) \rightarrow C$ preserves the pullback squares from Lemma 13.1.8 if and only if it is right Kan extended along the inclusion $J_n \hookrightarrow \mathbf{Tw}^r([n])$: if we compute this Kan extension row by row, this follows from the pointwise formula for Kan extensions (Theorem 21.4.3). Concretely, right Kan extension from J_n fills the diagram by iterated pullbacks, and hence encodes the successive composites of the original string of spans.

We now use the adequacy of (C, C_L, C_R) . Starting with a functor $J_n \rightarrow C$ that sends the left-pointing morphisms to C_L and the right-pointing morphisms to C_R , construct its right Kan extension by induction on $j - i$. At each step, the value at $(i \leq j)$ is obtained by pulling back the diagram formed by the previously constructed values at $(i \leq j - 1)$, $(i + 1 \leq j)$ and $(i + 1 \leq j - 1)$. The adequate-triple axiom guarantees that this pullback exists and that its new left- and right-pointing legs again lie in C_L and C_R , respectively. Thus restriction to J_n induces an inclusion of anima

$$X_n = \mathbf{NSpan}_{L,R}(C)_n \hookrightarrow \mathbf{Map}(J_n, C)$$

whose image consists precisely of those functors $J_n \rightarrow C$ sending the left-pointing morphisms to C_L and the right-pointing morphisms to C_R . We now observe that the Segal maps $e_i: [1] \hookrightarrow [n]$ induce inclusions $\mathbf{Tw}^r([1]) \hookrightarrow J_n$ which assemble into an equivalence

$$\mathbf{Tw}^r([1]) \sqcup_{\mathbf{Tw}^r([0])} \mathbf{Tw}^r([1]) \sqcup_{\mathbf{Tw}^r([0])} \cdots \sqcup_{\mathbf{Tw}^r([0])} \mathbf{Tw}^r([1]) \xrightarrow{\sim} J_n,$$

as may be checked directly in posets: the objects $(i \leq i)$ along which we glue are maximal in the adjacent copies of $\mathbf{Tw}^r([1])$, so the pushout creates no additional relations. In particular, we get

$$\mathbf{Map}(J_n, C) \xrightarrow{\sim} \mathbf{Map}(\mathbf{Tw}^r([1]), C) \times_{C^\simeq} \mathbf{Map}(\mathbf{Tw}^r([1]), C) \times_{C^\simeq} \cdots \times_{C^\simeq} \mathbf{Map}(\mathbf{Tw}^r([1]), C).$$

Under this equivalence, the image of the restriction map $X_n \hookrightarrow \mathbf{Map}(J_n, C)$ corresponds precisely to the target of the Segal map

$$(e_i^*)_{i=1}^n: X_n \rightarrow X_1 \times_{X_0} \cdots \times_{X_0} X_1,$$


showing that X satisfies the Segal condition.

For completeness, first note that the functor


$$C^\simeq = X_0 \rightarrow X_1 \subseteq \mathbf{Map}(\mathbf{Tw}^r([1]), C)$$

is an inclusion of animae. Indeed, since $\text{Tw}^r([1])$ has an initial object we have $\|\text{Tw}^r([1])\| \simeq *$, and thus the constant diagram functor $C \rightarrow \text{Fun}(\text{Tw}^r([1]), C)$ may be identified with the fully faithful inclusion $\text{Fun}(\|\text{Tw}^r([1])\|, C) \hookrightarrow \text{Fun}(\text{Tw}^r([1]), C)$. It thus remains to show that a span $X \xleftarrow{l} U \xrightarrow{r} Y$ is an isomorphism in the Segal anima $\text{NSpan}_{L,R}(C)$ if and only if both l and r are isomorphisms in C .

We first show that an invertible span has invertible legs. Let $Y \xleftarrow{l'} V \xrightarrow{r'} X$ be a left inverse, and let $Y \xleftarrow{l''} W \xrightarrow{r''} X$ be a right inverse. The two diagrams below exhibit these inverses; pulling them back against each other will produce the isomorphisms needed for a 2-out-of-6 argument. The identities $(l, r) \circ (l', r') = \text{id}$ and $(l'', r'') \circ (l, r) = \text{id}$ are exhibited by commutative diagrams in C of the following form:



Putting them into one single diagram and forming another pullback, we then obtain:



Since isomorphisms in C are closed under pullbacks, the maps $P \rightarrow V$ and $P \rightarrow W$ are isomorphisms, so by the 2-out-of-6 property all morphisms along the outer two edges of the diagram are isomorphisms. By the 2-out-of-3 property, then so are the maps $r': V \rightarrow X$ and $l'': W \rightarrow Y$, hence by pullback also $Y \rightarrow U$ and $X \rightarrow U$. Finally, another instance of the 2-out-of-3 property shows that $l: U \rightarrow X$ and $r: U \rightarrow Y$ are isomorphisms. Conversely, if l and r are isomorphisms, then the span is isomorphic in X_1 to the identity span on U , and hence lies in the image of $X_0 \rightarrow X_1$. This proves completeness. $\square$

Notation 13.1.12. By Proposition 13.1.11 and Theorem 24.1.2, the functor $\text{NSpan}: \text{AdTrip} \rightarrow \text{sAn}$ factors uniquely as

$$\begin{array}{ccc} \text{AdTrip} & \xrightarrow{\text{NSpan}} & \text{sAn} \\ & \searrow \text{Span} & \nearrow N \\ & \text{Cat}_\infty & \end{array}$$

We refer to the resulting ∞ -category $\text{Span}_{L,R}(C)$ as the *span category* of the adequate triple.

If C admits pullbacks, we write

$$\text{Span}(C) := \text{Span}_{\text{all}, \text{all}}(C)$$

for the span category associated with the adequate triple (C, C, C) . In particular, $\text{Span}(\text{Fin})$ is the rigorous construction of the span category of finite sets promised in Construction 5.3.1.

Lemma 13.1.13. *Let (C, C_L, C_R) be an adequate triple and consider objects $X, Y \in C$. Then there is an equivalence*

$$\mathrm{Hom}_{\mathrm{Span}_{L,R}(C)}(X, Y) \simeq ((C_L)_X)^{\simeq} \times_{C^{\simeq}} ((C_R)_Y)^{\simeq}.$$

Proof. Observe that the twisted arrow category $\mathrm{Tw}^r([1])$ is the walking span $\lrcorner$ from Definition 1.4.1. We may then compute:

$$\mathrm{Map}([1], \mathrm{Span}_{L,R}(C)) \simeq \mathrm{NSpan}_{L,R}(C)_1 \simeq \mathrm{Hom}_{\mathrm{AdTrip}}(\lrcorner, (C, C_L, C_R)) \simeq \mathrm{Map}([1], C_L) \times_{\mathrm{ev}_0, C^{\simeq}, \mathrm{ev}_0} \mathrm{Map}([1], C_R).$$

The hom anima $\mathrm{Hom}_{\mathrm{Span}_{L,R}(C)}(X, Y)$ is then given as the fiber over (X, Y) of the map

$$\mathrm{Map}([1], C_L) \times_{\mathrm{ev}_0, C^{\simeq}, \mathrm{ev}_0} \mathrm{Map}([1], C_R) \xrightarrow{(\mathrm{ev}_1, \mathrm{ev}_1)} C^{\simeq} \times C^{\simeq},$$

which is indeed equivalent to $((C_L)_X)^{\simeq} \times_{C^{\simeq}} ((C_R)_Y)^{\simeq}$. $\square$

Thus the case $n = 1$ of Construction 13.1.10 recovers the informal description from the beginning of the chapter: the objects are those of C , morphisms are spans with left leg in C_L and right leg in C_R , and identities are the identity spans. Under the Segal equivalence, the face map $d_1: \mathrm{NSpan}_{L,R}(C)_2 \rightarrow \mathrm{NSpan}_{L,R}(C)_1$ sends a pullback grid to its outer span. Consequently, composition in $\mathrm{Span}_{L,R}(C)$ is given by pullback.

Corollary 13.1.14. *Suppose that $C'_L \subseteq C_L$ and $C'_R \subseteq C_R$ are wide subcategories such that both (C, C'_L, C'_R) and (C, C_L, C_R) are adequate triples. Then the induced functor*

$$\mathrm{Span}_{L',R'}(C) \hookrightarrow \mathrm{Span}_{L,R}(C)$$

is a wide subcategory inclusion.

Proof. Both span categories have the same objects, and Lemma 13.1.13 identifies the induced maps on hom animae with inclusions of components. $\square$

Let us now prove various basic properties of span categories.

Lemma 13.1.15. *There is a natural equivalence $\mathrm{Span}_{L,R}(C)^{\mathrm{op}} \simeq \mathrm{Span}_{R,L}(C)$, in the sense that the following diagram commutes:*

$$\begin{array}{ccc} \mathrm{AdTrip} & \xrightarrow{(-)^{\mathrm{rev}}} & \mathrm{AdTrip} \\ \mathrm{Span} \downarrow & & \downarrow \mathrm{Span} \\ \mathrm{Cat}_{\infty} & \xrightarrow{(-)^{\mathrm{op}}} & \mathrm{Cat}_{\infty}. \end{array}$$

Proof. By definition of $(-)^{\mathrm{op}}: \mathrm{Cat}_{\infty} \rightarrow \mathrm{Cat}_{\infty}$, there is a commutative square

$$\begin{array}{ccc} \mathrm{Cat}_{\infty} & \xrightarrow{(-)^{\mathrm{op}}} & \mathrm{Cat}_{\infty} \\ N \downarrow & & \downarrow N \\ \mathrm{sAn} & \xrightarrow{(-)^{\mathrm{op}}} & \mathrm{sAn}, \end{array}$$

where the bottom functor is given by precomposition with $(-)^{\text{op}} : \Delta \xrightarrow{\sim} \Delta$. We thus need to produce a natural equivalence $\text{NSpan}_{L,R}(C)^{\text{op}} \simeq \text{NSpan}_{R,L}(C)$ of simplicial animae. Unwinding definitions, this amounts to producing a natural equivalence

$$\text{Hom}_{\text{AdTrip}}(\text{Tw}^r([n]^{\text{op}}), (C, C_L, C_R)) \simeq \text{Hom}_{\text{AdTrip}}(\text{Tw}^r([n]), (C, C_R, C_L)),$$

natural in $[n] \in \Delta^{\text{op}}$ and (C, C_L, C_R) . Equivalently, we must produce a natural equivalence $\text{Tw}^r([n]^{\text{op}}) \simeq \text{Tw}^r([n])_{\text{rev}}$. Such an equivalence is given by sending an object $(i \geq j)$ in $\text{Tw}^r([n]^{\text{op}})$ to the object $(j \leq i)$ in $\text{Tw}^r([n])$. $\square$

The span category $\text{Span}_{L,R}(C)$ contains C_R as a wide subcategory on the right-pointing spans $X \xleftarrow{=} X \xrightarrow{r} Y$. Similarly it contains C_L^{op} as a wide subcategory on the left-pointing spans $X \xleftarrow{l} Y \xrightarrow{=} Y$. This is made precise by the following result:

Lemma 13.1.16. *Let $C_L, C_R \subseteq C$ be wide subcategories. Then the span categories of the adequate triples $(C, C^{\simeq}, C_R)$ and $(C, C_L, C^{\simeq})$ from Example 13.1.3 are given by:*

$$\text{Span}_{\simeq,R}(C) \simeq C_R \quad \text{and} \quad \text{Span}_{L,\simeq}(C) \simeq C_L^{\text{op}}.$$

In particular, there are canonical inclusions

$$C_R \simeq \text{Span}_{\simeq,R}(C) \hookrightarrow \text{Span}_{L,R}(C) \quad \text{and} \quad C_L^{\text{op}} \simeq \text{Span}_{L,\simeq}(C) \hookrightarrow \text{Span}_{L,R}(C).$$

Proof. It suffices to prove the first equivalence; the second one follows by combining the first one with Lemma 13.1.15. For $[n] \in \Delta^{\text{op}}$, consider the source functor $s : \text{Tw}^r([n]) \rightarrow [n]$, $(i \leq j) \mapsto i$. Note that this functor admits a fully faithful left adjoint $[n] \rightarrow \text{Tw}^r([n])$ sending $i \in [n]$ to $(i \leq n) \in \text{Tw}^r([n])$. Indeed, there is a morphism $(i \leq n) \rightarrow (k \leq l)$ in $\text{Tw}^r([n])$ if and only if $i \leq k$. It thus follows from Lemma 21.8.8 that restriction along s induces an inclusion of animae

$$\text{Hom}_{\text{Cat}_{\infty}}([n], C_R) \subseteq \text{Hom}_{\text{Cat}_{\infty}}([n], C) \hookrightarrow \text{Hom}_{\text{Cat}_{\infty}}(\text{Tw}^r([n]), C)$$

whose essential image consists of those functors $\text{Tw}^r[n] \rightarrow C$ that send backwards morphisms to isomorphisms and forward morphisms into C_R . Since these are precisely the morphisms of adequate triples into $(C, C^{\simeq}, C_R)$, we obtain a natural equivalence

$$N(C_R)_n = \text{Hom}_{\text{Cat}_{\infty}}([n], C_R) \xrightarrow{\sim} \text{Hom}_{\text{AdTrip}}(\text{Tw}^r([n]), (C, C^{\simeq}, C_R)) = \text{NSpan}_{\simeq,R}(C)_n.$$

This shows that $N(C_R) \simeq \text{NSpan}_{\simeq,R}(C)$, and hence that $C_R \simeq \text{Span}_{\simeq,R}(C)$ by full faithfulness of the nerve functor. $\square$

The construction of span categories is summarized conceptually by the following adjunction. We will not need the adjunction itself below, but it explains the role of the right twisted-arrow construction in the definition of Span .

Proposition 13.1.17 ([HHLN23b, Theorem 2.18]). *The right twisted-arrow construction $\text{Tw}^r : \Delta \rightarrow \text{AdTrip}$ extends to a functor*

$$\text{Tw}^r : \text{Cat}_{\infty} \rightarrow \text{AdTrip},$$

and this extension is left adjoint to the span construction:

$$\text{Tw}^r : \text{Cat}_{\infty} \rightleftarrows \text{AdTrip} : \text{Span}.$$

Lemma 13.1.18. *The ∞ -category $\mathbf{AdTrip}$ admits limits and filtered colimits, and the inclusion $\mathbf{AdTrip} \hookrightarrow \mathbf{Fun}(\sqcup, \mathbf{Cat}_\infty)$ preserves limits and filtered colimits.*

Proof. We start with the case for limits. Recall that limits in $\mathbf{Fun}(\sqcup, \mathbf{Cat}_\infty)$ are computed pointwise. Given a functor $C: I \rightarrow \mathbf{AdTrip}$, we thus need to show that the triple $(C, C_L, C_R) := (\lim_i C(i), \lim_i C(i)_L, \lim_i C(i)_R)$ is again adequate. Note that subcategory inclusions are closed under limits in $\mathbf{Cat}_\infty$: a functor $D \hookrightarrow D'$ is a subcategory inclusion if and only if the commutative square

$$\begin{array}{ccc} D & \xlongequal{\quad} & D \\ \parallel & & \downarrow \\ D & \longrightarrow & D' \end{array}$$

is a pullback square, and limits commute with pullback squares. Furthermore, *wide* subcategory inclusions are closed under limits: an inclusion $D \hookrightarrow D'$ is a wide subcategory if and only if the induced map $D^\simeq \rightarrow D'^\simeq$ on groupoid cores is an equivalence of animae, and $(-)^{\simeq}$ preserves limits.

Consider a cospan $X \xrightarrow{l} X'' \xleftarrow{r} X'$ in $C = \lim_i C(i)$ with $l \in C_L$ and $r \in C_R$. Since the transition functors preserve the pullbacks specified by the adequate-triple structures, the pointwise pullbacks in the $C(i)$ assemble to an object of C . The identity

$$\mathrm{Hom}_C(W, -) \simeq \lim_i \mathrm{Hom}_{C(i)}(W_i, -)$$

shows that this object is the pullback in C , and the pointwise criterion for membership in C_L and C_R shows that its legs lie in the required classes. Thus (C, C_L, C_R) is an adequate triple. Furthermore, a functor into C is a morphism of adequate triples if and only if its composite with each projection $C \rightarrow C(i)$ is, so (C, C_L, C_R) is a limit in $\mathbf{AdTrip}$.

For filtered colimits, note that subcategory inclusions are closed under filtered colimits in $\mathbf{Cat}_\infty$ since filtered colimits commute with pullbacks by Lemma 24.2.2. Since every object in a filtered colimit of ∞ -categories $D(i)$ comes from one of the $D(i)$, it follows that a filtered colimit of wide subcategory inclusions is again wide. Any cospan in $C = \mathrm{colim}_i C(i)$ lifts to a cospan in some stage $C(j)$. Its pullback exists there by adequacy, and the transition functors carry this pullback to the corresponding pullbacks at every later stage. Mapping animae in a filtered colimit of ∞ -categories are filtered colimits of the mapping animae, and finite limits of animae commute with filtered colimits. It follows that the resulting square in C remains a pullback. Thus (C, C_L, C_R) is an adequate triple. A functor out of C is a morphism of adequate triples if and only if its composite with each $C(j) \rightarrow C$ is, so (C, C_L, C_R) is a colimit in $\mathbf{AdTrip}$. $\square$

Proposition 13.1.19. *The functor $\mathrm{Span}: \mathbf{AdTrip} \rightarrow \mathbf{Cat}_\infty$ preserves limits and filtered colimits.*

Proof. The inclusion $N: \mathbf{Cat}_\infty \hookrightarrow \mathbf{sAn}$ preserves limits and filtered colimits (see Lemma 24.2.2 for the latter claim). It will thus suffice to show that $N\mathrm{Span}: \mathbf{AdTrip} \rightarrow \mathbf{sAn}$ preserves these (co)limits. Since these are computed pointwise, this boils down to showing that the functor

$$\mathrm{Hom}_{\mathbf{AdTrip}}(\mathrm{Tw}^r([n]), -): \mathbf{AdTrip} \rightarrow \mathbf{An}$$

preserves them. The case for limits is clear, since the functor is corepresented. For filtered colimits, we use again that filtered colimits commute with pullbacks, so that we may reduce to the cases $n = 0$ and $n = 1$ via the Segal condition. Note that the ∞ -categories $\mathrm{Tw}^r([0]) = *$ and $\mathrm{Tw}^r([1]) = \sqcap$ are

finite, so $\text{Hom}_{\text{Cat}_\infty}(\text{Tw}^r([n]), -) : \text{Cat}_\infty \rightarrow \text{An}$ preserves filtered colimits by Corollary 24.2.3. This proves the claim for $n = 0$. For $n = 1$, one observes that the subanimaes of morphisms of adequate triples are closed under filtered colimits. $\square$

13.2 Adjunctions of span categories

Before studying products and coproducts in span categories, we develop a general criterion for an adjunction between adequate triples to induce an adjunction between their span categories.

Construction 13.2.1. Consider adequate triples (C, C_L, C_R) and (D, D_L, D_R) and let $F, G : C \rightarrow D$ be morphisms of adequate triples. Let $\alpha : F \Rightarrow G$ be a natural transformation, and assume that the following two conditions are satisfied:

- (1) The morphism $\alpha(X) : F(X) \rightarrow G(X)$ lies in D_R for every object X of C ;
- (2) For every morphism $l : Z \rightarrow X$ in C_L the commutative square

$$\begin{array}{ccc} F(Z) & \xrightarrow{\alpha(Z)} & G(Z) \\ F(l) \downarrow & & \downarrow G(l) \\ F(X) & \xrightarrow{\alpha(X)} & G(X) \end{array}$$

is a pullback square in D .

We will construct a natural transformation

$$\text{Span}^R(\alpha) : \text{Span}(F) \Rightarrow \text{Span}(G)$$

of functors $\text{Span}_{L,R}(C) \rightarrow \text{Span}_{L,R}(D)$ which is objectwise given by the forward spans

$$F(X) \xleftarrow{=} F(X) \xrightarrow{\alpha(X)} G(X)$$

for $X \in C$. For this, equip $[1]$ with the adequate triple structure $([1], [1]^\approx, [1])$. Then the product $C \times [1]$ is an adequate triple with backward and forward classes $C_L \times [1]^\approx$ and $C_R \times [1]$, respectively. Regarding α as a functor $C \times [1] \rightarrow D$, conditions (1) and (2) say precisely that it is a morphism of adequate triples. Indeed, the only additional pullback squares to check are sent to pastings of images under F or G of adequate pullback squares with the naturality squares from (2). We therefore obtain a functor

$$\text{Span}_{L,R}(C \times [1]) \rightarrow \text{Span}_{L,R}(D).$$

Using that $\text{Span}(-)$ preserves products by Proposition 13.1.19 and that $\text{Span}_{\text{iso,all}}([1]) \simeq [1]$ by Lemma 13.1.16, this functor takes the form

$$\text{Span}_{L,R}(C) \times [1] \rightarrow \text{Span}_{L,R}(D).$$

The restrictions to 0 and 1 are precisely $\text{Span}(F)$ and $\text{Span}(G)$, so this produces the desired natural transformation $\text{Span}^R(\alpha)$.

Remark 13.2.2. The previous construction can be dualized, if we assume instead that the morphism $\alpha(X): F(X) \rightarrow G(X)$ is in D_L for every $X \in C$ and that for every morphism $r: Z \rightarrow Y$ in C_R the commutative square

$$\begin{array}{ccc} F(Z) & \xrightarrow{\alpha(Z)} & G(Z) \\ F(r) \downarrow & & \downarrow G(r) \\ F(Y) & \xrightarrow{\alpha(Y)} & G(Y) \end{array}$$

is a pullback square in D . By applying Lemma 13.1.15 to the previous construction, we deduce that α induces a natural transformation

$$\text{Span}^L(\alpha): \text{Span}(G) \Longrightarrow \text{Span}(F)$$

of functors $\text{Span}_{L,R}(C) \rightarrow \text{Span}_{L,R}(D)$ given on an object $X \in C$ by the backwards span

$$G(X) \xleftarrow{\alpha(X)} F(X) \xrightarrow{=} F(X)$$

Lemma 13.2.3. *Given a third morphism of adequate triples $H: (C, C_L, C_R) \rightarrow (D, D_L, D_R)$, let $\alpha: F \Rightarrow G$ and $\beta: G \Rightarrow H$ be natural transformations satisfying the conditions (1) and (2) from Construction 13.2.1. Then we have natural equivalences*

$$\text{Span}^R(\text{id}_F) \simeq \text{id}_{\text{Span}(F)} \quad \text{and} \quad \text{Span}^R(\beta \circ \alpha) \simeq \text{Span}^R(\beta) \circ \text{Span}^R(\alpha).$$

The construction is also compatible with whiskering: if $K: (B, B_L, B_R) \rightarrow (C, C_L, C_R)$ and $J: (D, D_L, D_R) \rightarrow (E, E_L, E_R)$ are morphisms of adequate triples, then

$$\text{Span}^R(\alpha K) \simeq \text{Span}^R(\alpha) \circ \text{Span}^R(K) \quad \text{and} \quad \text{Span}^R(J\alpha) \simeq \text{Span}^R(J) \circ \text{Span}^R(\alpha).$$

The dual claims for $\text{Span}^L(-)$ also hold.

Proof. Composition and identities of natural transformations are formed using the posets $[n]$ for $n = 0, 2$. Whiskering is given by pre- and postcomposition with morphisms of adequate triples. The claims therefore follow from the naturality in both $[n] \in \Delta$ and $(C, C_L, C_R) \in \text{AdTrip}$ of the equivalence $\text{Span}_{L,R}(C \times [n]) \simeq \text{Span}_{L,R}(C) \times [n]$. $\square$

Corollary 13.2.4. *Let (C, C_L, C_R) and (D, D_L, D_R) be adequate triples. Let $F: C \rightleftarrows D : G$ be an adjunction of ∞ -categories and assume that F and G are morphisms of adequate triples.*

(1) *Assume that the unit map $\eta_X: X \rightarrow G(F(X))$ is in C_R for all $X \in C$ and the counit map $\varepsilon_Y: F(G(Y)) \rightarrow Y$ is in D_R for all $Y \in D$. Furthermore, assume that for every morphism $l: Z \rightarrow X$ in C_L and every morphism $l': Z' \rightarrow X'$ in D_L , the commutative squares*

$$\begin{array}{ccc} Z & \xrightarrow{\eta_Z} & G(F(Z)) \\ l \downarrow & & \downarrow G(F(l)) \\ X & \xrightarrow{\eta_X} & G(F(X)) \end{array} \quad \text{and} \quad \begin{array}{ccc} F(G(Z')) & \xrightarrow{\varepsilon_{Z'}} & Z' \\ F(G(l')) \downarrow & & \downarrow l' \\ F(G(X')) & \xrightarrow{\varepsilon_{X'}} & X' \end{array}$$

are pullback squares. Then F and G induce an adjunction

$$\text{Span}(F): \text{Span}_{L,R}(C) \rightleftarrows \text{Span}_{L,R}(D) : \text{Span}(G).$$

- (2) Dually, assume that the unit map $\eta_X: X \rightarrow G(F(X))$ is in C_L for all $X \in C$ and the counit map $\varepsilon_Y: F(G(Y)) \rightarrow Y$ is in D_L for all $Y \in D$. Furthermore, assume that for every morphism $r: Z \rightarrow Y$ in C_R and every morphism $r': Z' \rightarrow Y'$ in D_R , the commutative squares

$$\begin{array}{ccc} Z & \xrightarrow{\eta_Z} & G(F(Z)) \\ r \downarrow & & \downarrow G(F(r)) \\ Y & \xrightarrow{\eta_Y} & G(F(Y)) \end{array} \quad \text{and} \quad \begin{array}{ccc} F(G(Z')) & \xrightarrow{\varepsilon_{Z'}} & Z' \\ F(G(r')) \downarrow & & \downarrow r' \\ F(G(Y')) & \xrightarrow{\varepsilon_{Y'}} & Y' \end{array}$$

are pullback squares. Then F and G induce an adjunction

$$\text{Span}(G): \text{Span}_{L,R}(D) \rightleftarrows \text{Span}_{L,R}(C) : \text{Span}(F).$$

Proof. We prove (1), as (2) is dual. By Construction 13.2.1 and Lemma 13.2.3, the unit $\eta: \text{id} \rightarrow GF$ and counit $\varepsilon: FG \rightarrow \text{id}$ induce transformations

$$\text{Span}^R(\eta): \text{Span}(\text{id}) \rightarrow \text{Span}(G) \circ \text{Span}(F) \quad \text{and} \quad \text{Span}^R(\varepsilon): \text{Span}(F) \circ \text{Span}(G) \rightarrow \text{id}.$$

Compatibility with vertical composition and whiskering from Lemma 13.2.3 shows that these transformations satisfy the triangle identities. They therefore exhibit the asserted adjunction by Proposition 21.1.2. $\square$

13.3 Products and coproducts in span categories

We now apply the previous results on adjunctions of span categories to study products and coproducts in span categories.

Definition 13.3.1. An adequate triple (C, C_L, C_R) is called *weakly extensive* if the following conditions are satisfied:

- (1) The ∞ -category C admits finite coproducts;
- (2) The coproduct functor $C \times C \rightarrow C$ is a morphism of adequate triples (i.e., morphisms in C_L and C_R are closed under coproducts, and coproducts of pullback squares of morphisms in C_L along morphisms in C_R are again pullback squares);
- (3) For every object $X \in C$, the morphisms $\emptyset \rightarrow X$ and $\nabla: X \sqcup X \rightarrow X$ are in C_L ;
- (4) For every morphism $r: X \rightarrow Y$ in C_R , the commutative squares

$$\begin{array}{ccc} X \sqcup X & \xrightarrow{r \sqcup r} & Y \sqcup Y \\ \nabla \downarrow & & \downarrow \nabla \\ X & \xrightarrow{r} & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} \emptyset & \xrightarrow{\quad} & \emptyset \\ \downarrow & & \downarrow \\ X & \xrightarrow{r} & Y \end{array}$$

are pullback squares.

We say that the triple is *weakly coextensive* if (C, C_R, C_L) is weakly extensive, and we say it is *extensive* if it is both weakly extensive and weakly coextensive. An ∞ -category C is called *extensive* if it admits a wide subcategory C_R such that (C, C, C_R) is extensive. This agrees with the usual definition in terms of slice categories, as shown in Exercise 13.5; in particular, it is independent of the choice of C_R witnessing extensivity.

Example 13.3.2. The triple $(\text{Fin}, \text{Fin}, \text{Fin})$ is extensive. More generally, for a finite group G , the triple $(\text{Fin}_G, \text{Fin}_G, \text{Fin}_G)$ is extensive.

Example 13.3.3. The triple $(\text{Top}, \text{Top}, \text{Top})$ is extensive.

Example 13.3.4. If a full subcategory $C \subseteq \text{An}$ is closed under finite coproducts and pullbacks, then the triple (C, C, C) is extensive.

Lemma 13.3.5. *Let (C, C_L, C_R) be an adequate triple.*

- (1) *If the triple is weakly extensive, then $\text{Span}_{L,R}(C)$ admits finite products, and the inclusion $C_L^{\text{op}} \hookrightarrow \text{Span}_{L,R}(C)$ preserves finite products;*
- (2) *If the triple is weakly coextensive, then $\text{Span}_{L,R}(C)$ admits finite coproducts, and the inclusion $C_R \hookrightarrow \text{Span}_{L,R}(C)$ preserves finite coproducts;*
- (3) *If the triple is extensive, then $\text{Span}_{L,R}(C)$ is semiadditive.*

Proof. For part (1), note that C has binary coproducts if and only if the diagonal functor $\Delta: C \rightarrow C \times C$ admits a left adjoint $\sqcup: C \times C \rightarrow C$. The counit of the adjunction is the fold map $\nabla: X \sqcup X \rightarrow X$ and the unit is the pair of canonical inclusions $(X \hookrightarrow X \sqcup Y, Y \hookrightarrow X \sqcup Y)$. Conditions (1) to (4) supply the hypotheses of part (2) of Corollary 13.2.4, so this adjunction induces an adjunction

$$\text{Span}_{L,R}(C) \rightleftarrows \text{Span}_{L,R}(C \times C) \simeq \text{Span}_{L,R}(C) \times \text{Span}_{L,R}(C),$$

where the adequate triple on $C \times C$ has classes $C_L \times C_L$ and $C_R \times C_R$. Let us spell out the hypotheses of Corollary 13.2.4. The coproduct functor is a morphism of adequate triples by condition (2), while the diagonal functor preserves the relevant classes and pullbacks componentwise. The fold map lies in C_L by condition (3). Each coprojection, for example

$$X = X \sqcup \emptyset \xrightarrow{\text{id}_X \sqcup (\emptyset \rightarrow Y)} X \sqcup Y,$$

lies in C_L by conditions (2) and (3). The naturality square for the fold map is the first pullback square in condition (4). The naturality squares for the coprojections decompose as coproducts of an identity square and the second pullback square in condition (4), so they are pullbacks by condition (2). This proves that the displayed adjunction exists, and hence that $\text{Span}_{L,R}(C)$ admits binary products. For the terminal object, we similarly regard the initial object $\emptyset$ of C as a left adjoint to the functor $C \rightarrow *$. Conditions (3) and (4) allow us to apply Corollary 13.2.4 and obtain an adjunction $\text{Span}_{L,R}(C) \rightleftarrows \text{Span}(*) \simeq *$ exhibiting $\emptyset$ as a terminal object of $\text{Span}_{L,R}(C)$.

It remains to prove the claim about the inclusion $C_L^{\text{op}} \hookrightarrow \text{Span}_{L,R}(C)$. The subcategory C_L admits finite coproducts computed as in C . Indeed, the coprojections lie in C_L by the preceding argument. Given morphisms $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ in C_L , their copairing factors as

$$X \sqcup Y \xrightarrow{f \sqcup g} Z \sqcup Z \xrightarrow{\nabla} Z$$

and therefore lies in C_L . Conversely, a morphism $X \sqcup Y \rightarrow Z$ in C_L restricts to morphisms in C_L because the coprojections lie in C_L . The initial object is initial in C_L by condition (3). Under the inclusion $C_L^{\text{op}} \hookrightarrow \text{Span}_{L,R}(C)$, the resulting product projections are the backwards spans associated

with the coprojections, which are precisely the product projections constructed above. Thus this inclusion preserves finite products.

Part (2) is an immediate consequence of (1) using $\text{Span}_{L,R}(C)^{\text{op}} \simeq \text{Span}_{R,L}(C)$. For part (3), note first that the initial object $\emptyset$ in C becomes both initial and terminal in $\text{Span}_{L,R}(C)$, and that the zero morphism between objects X and Y is given by the span

$$X \leftarrow \emptyset \rightarrow Y.$$

Also notice that the coproduct $X \sqcup Y$ in C becomes both the product and the coproduct in $\text{Span}_{L,R}(C)$, via the spans

$$X \sqcup Y \leftarrow X \xrightarrow{=} X, \quad X \sqcup Y \leftarrow Y \xrightarrow{=} Y, \quad X \xleftarrow{=} X \hookrightarrow X \sqcup Y, \quad Y \xleftarrow{=} Y \hookrightarrow X \sqcup Y.$$

It thus remains to compute the four composites and show that they are id_X , 0 , 0 and id_Y . The required pullback squares follow from the extensivity assumptions. The first square below is the coproduct of the identity square on X and the empty pullback square from condition (4) for the morphism $\emptyset \rightarrow Y$, which lies in C_R by weak coextensivity; the second is obtained symmetrically. The third is the coproduct of the empty pullback square from condition (4) for id_X and the identity square on the morphism $\emptyset \rightarrow Y$. Condition (2) guarantees that these coproducts are again pullback squares. Thus we have:

$$\begin{array}{ccc} \begin{array}{ccc} X & \xrightarrow{\quad} & X \\ \downarrow \lrcorner & & \downarrow \\ X & \xrightarrow{\quad} & X \sqcup Y \end{array} & \begin{array}{ccc} Y & \xrightarrow{\quad} & Y \\ \downarrow \lrcorner & & \downarrow \\ Y & \xrightarrow{\quad} & X \sqcup Y \end{array} & \text{and} \quad \begin{array}{ccc} \emptyset & \xrightarrow{\quad} & Y \\ \downarrow \lrcorner & & \downarrow \\ X & \xrightarrow{\quad} & X \sqcup Y \end{array} \end{array} \quad \square$$

Corollary 13.3.6. *The ∞ -category $\text{Span}(\text{Fin})$ is semiadditive, with biproducts given by disjoint unions of finite sets.*

Proof. Apply part (3) of Lemma 13.3.5 to the extensive triple $(\text{Fin}, \text{Fin}, \text{Fin})$. This recovers the direct argument of Lemma 5.3.3. $\square$

Exercises

Exercise 13.1. Show that the category Fin_G of finite G -sets is extensive for every finite group G .

Exercise 13.2. Show that the categories Top and Sch of topological spaces and schemes, respectively, are extensive.

Exercise 13.3. Show that An is extensive.

Exercise 13.4. Let C be an ∞ -category with finite coproducts. Show that C is extensive if and only if it satisfies the following two conditions:

- *Coproducts in C are disjoint:* for objects X and Y , the commutative square

$$\begin{array}{ccc} \emptyset & \xrightarrow{\quad} & X \\ \downarrow & & \downarrow \\ Y & \xrightarrow{\quad} & X \sqcup Y \end{array}$$

is a pullback square.

- *Coproduct decompositions are stable under pullback:* given a morphism $f: Z \rightarrow X \sqcup Y$, the map $Z_X \sqcup Z_Y \rightarrow Z$ is an isomorphism, where Z_X and Z_Y are defined by the following two pullback squares:

$$\begin{array}{ccc} Z_X & \longrightarrow & Z \\ \downarrow & \lrcorner & \downarrow f \\ X & \longrightarrow & X \sqcup Y \end{array} \quad \text{and} \quad \begin{array}{ccc} Z_Y & \longrightarrow & Z \\ \downarrow & \lrcorner & \downarrow f \\ Y & \longrightarrow & X \sqcup Y. \end{array}$$

Exercise 13.5. Let C be an ∞ -category. Show that C is extensive if and only if it has finite coproducts and for all $n \geq 0$ and objects $X_1, \dots, X_n$ the coproduct functor

$$\prod_{i=1}^n C_{/X_i} \rightarrow C_{/\sqcup_i X_i}$$

is an equivalence.

Exercise 13.6 (The span nerve in low degrees). Let (C, C_L, C_R) be an adequate triple.

- (1) Describe explicitly the anima $\mathbf{NSpan}_{L,R}(C)_n$ for $0 \leq n \leq 3$.
- (2) Show that the face map

$$d_1: \mathbf{NSpan}_{L,R}(C)_2 \rightarrow \mathbf{NSpan}_{L,R}(C)_1$$

sends a pair of composable spans to their composite formed by pullback.

- (3) For three composable spans with apexes U_{01} , U_{12} , and U_{23} , explain how an element of $\mathbf{NSpan}_{L,R}(C)_3$ determines the canonical isomorphism

$$(U_{01} \times_{X_1} U_{12}) \times_{X_2} U_{23} \simeq U_{01} \times_{X_1} (U_{12} \times_{X_2} U_{23})$$

that witnesses the associativity of composition.

Exercise 13.7 (Spans of finite sets as matrices). Let I and J be finite sets. Given a span $I \leftarrow K \rightarrow J$, let $K_{ji} \subseteq K$ be the fiber over $(i, j) \in I \times J$.

- (1) Show that there is an equivalence

$$\mathrm{Hom}_{\mathrm{Span}(\mathrm{Fin})}(I, J) \simeq \prod_{(i,j) \in I \times J} \mathrm{Fin}^{\sim}.$$

- (2) Deduce that the isomorphism classes of morphisms $I \rightarrow J$ in $\mathrm{Span}(\mathrm{Fin})$ are naturally identified with matrices $(a_{ji})_{(j,i) \in J \times I}$ over $\mathbb{N}$, where $a_{ji} := |K_{ji}|$.
- (3) Show that composition of spans corresponds to matrix multiplication and that identity spans correspond to identity matrices. Explain what information in the hom anima is forgotten by passing to matrices over $\mathbb{N}$.

Exercise 13.8. Use Proposition 5.3.16 to prove that $\mathrm{Span}(\mathrm{Fin})$ is the free semiadditive ∞ -category on a single generator: for every semiadditive ∞ -category C , evaluation at the one-point set $\langle 1 \rangle \in \mathrm{Fin}$ induces an equivalence

$$\mathrm{Fun}^{\times}(\mathrm{Span}(\mathrm{Fin}), C) \xrightarrow{\sim} C$$

of ∞ -categories.

14 Basics on ∞ -operads

Now that we have a solid grasp on span categories, we return to the topic of ∞ -operads. The definition itself was already given in Definition 12.4.5; this chapter supplies the vocabulary that makes it usable, and everything later in the part depends on it.

The central construction is the *multimorphism operad* $\mathcal{M}_C$ of a symmetric monoidal ∞ -category C , whose multimorphism anima are the $\mathrm{Hom}_C(x_1 \otimes \cdots \otimes x_n, y)$. It turns symmetric monoidal ∞ -categories into ∞ -operads, and thereby supplies both of the notions we will use constantly: an *$\mathcal{O}$ -algebra* in C is an operad map $\mathcal{O} \rightarrow \mathcal{M}_C$, and a *lax symmetric monoidal functor* $C \rightarrow D$ is an operad map $\mathcal{M}_C \rightarrow \mathcal{M}_D$. Neither has to be defined by hand as a list of coherence data; both fall out of the formalism. From this chapter onwards we make constant use of cocartesian fibrations and the straightening/unstraightening equivalence of Chapter 23.

In Section 14.1 we give the formal definition, establish a useful characterization in terms of cocartesian morphisms, and discuss the first examples. Section 14.2 constructs $\mathcal{M}_C$ and introduces $\mathcal{O}$ -algebras, and Section 14.3 treats lax symmetric monoidal functors, including the (lax) symmetric monoidality of the left and right adjoints of a (lax) symmetric monoidal functor. In Section 14.4 we generalize this discussion to *$\mathcal{O}$ -monoids* and *$\mathcal{O}$ -monoidal ∞ -categories* for an arbitrary ∞ -operad $\mathcal{O}$. Finally, Section 14.5 applies the operadic description to construct symmetric monoidal structures on suitable subcategories and Bousfield localizations, as needed in Part I.

14.1 Definitions and examples

Let us recall the definition of an ∞ -operad from Definition 12.4.5. Since cocartesian morphisms are now available to us (Definition 23.1.1), we may state the third condition directly, rather than unwinding it into a pullback square as we did in Chapter 12. We take the opportunity to also introduce the morphisms between ∞ -operads.

Definition 14.1.1. An *∞ -operad* (or: *$\mathrm{Span}(\mathrm{Fin})$ - ∞ -operad*) is a pair $\mathcal{O} = (\mathcal{O}^\otimes, p_{\mathcal{O}})$ consisting of an ∞ -category $\mathcal{O}^\otimes$ equipped with a functor $p_{\mathcal{O}}: \mathcal{O}^\otimes \rightarrow \mathrm{Span}(\mathrm{Fin})$ satisfying the following conditions:

- (1) The ∞ -category $\mathcal{O}^\otimes$ admits finite products, and the functor $p_{\mathcal{O}}$ preserves finite products;
- (2) For every finite set I , the product in $\mathcal{O}^\otimes$ defines an equivalence

$$\prod_{i \in I} \mathcal{O}_{\{i\}}^\otimes \xrightarrow{\sim} \mathcal{O}_I^\otimes,$$

where $\mathcal{O}_I^\otimes$ denotes the fiber of $p_{\mathcal{O}}$ over the set I ;

- (3) For a morphism $f: J \rightarrow I$ in $\mathbf{Fin}$ and objects $X_i \in \mathcal{O}^\otimes$, the map $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ whose j -th component is the projection to $X_{f(j)}$ is $p_{\mathcal{O}}$ -cocartesian.

The objects X_i in condition (3) are not required to be colors; this more general form will be used in Proposition 14.1.7.

If $\mathcal{P} = (\mathcal{P}^\otimes, p_{\mathcal{P}})$ is another ∞ -operad, then a *morphism of ∞ -operads* $f: \mathcal{O} \rightarrow \mathcal{P}$ is a commutative diagram

$$\begin{array}{ccc} \mathcal{O}^\otimes & \xrightarrow{f^\otimes} & \mathcal{P}^\otimes \\ & \searrow p_{\mathcal{O}} \quad \swarrow p_{\mathcal{P}} & \\ & \text{Span}(\mathbf{Fin}) & \end{array}$$

such that the functor $f^\otimes$ preserves finite products.

If we write $\mathbf{Cat}_\infty^{\text{prod}}$ for the (very large) ∞ -category of ∞ -categories that admit finite products and functors that preserve finite products, we obtain a full subcategory

$$\mathbf{Op}_\infty \subseteq (\mathbf{Cat}_\infty^{\text{prod}})_{/\text{Span}(\mathbf{Fin})}$$

spanned by the ∞ -operads. Given two ∞ -operads $\mathcal{O}$ and $\mathcal{P}$, we will denote by $\mathbf{Fun}_{\mathbf{Op}_\infty}(\mathcal{O}, \mathcal{P})$ the ∞ -category of morphisms of ∞ -operads from $\mathcal{O}$ to $\mathcal{P}$, defined as the following fiber:

$$\begin{array}{ccc} \mathbf{Fun}_{\mathbf{Op}_\infty}(\mathcal{O}, \mathcal{P}) & \xrightarrow{\quad} & \mathbf{Fun}^\times(\mathcal{O}^\otimes, \mathcal{P}^\otimes) \\ \downarrow & \lrcorner & \downarrow p_{\mathcal{P}} \circ - \\ \{p_{\mathcal{O}}\} & \hookrightarrow & \mathbf{Fun}^\times(\mathcal{O}^\otimes, \text{Span}(\mathbf{Fin})). \end{array}$$

In certain contexts, morphisms of ∞ -operads $\mathcal{O} \rightarrow \mathcal{P}$ are also known as *$\mathcal{O}$ -algebras in $\mathcal{P}$* , and a common alternative notation for the ∞ -category $\mathbf{Fun}_{\mathbf{Op}_\infty}(\mathcal{O}, \mathcal{P})$ is $\mathbf{Alg}_{\mathcal{O}}(\mathcal{P})$.

Observation 14.1.2. The ∞ -category $\mathbf{Op}_\infty$ admits finite products. The terminal object is *Comm*, and the product of ∞ -operads $\mathcal{O}$ and $\mathcal{P}$ has total category

$$(\mathcal{O} \times \mathcal{P})^\otimes = \mathcal{O}^\otimes \times_{\text{Span}(\mathbf{Fin})} \mathcal{P}^\otimes.$$

Indeed, finite products, the decomposition of the fibers, and the required cocartesian lifts are all computed componentwise in this fiber product.

Warning 14.1.3. As mentioned in Remark 12.4.6, our definition of ∞ -operads does not agree with the standard one in the literature, due to Lurie, which we will discuss in Section 17.4 below. However, the difference is mostly inconsequential for practical purposes, as most constructions with Lurie's definition go through with our definition. The main difference is that with our definition we can state various conditions on the ∞ -category $\mathcal{O}^\otimes$ in terms of finite products rather than via a Segal condition.

Terminology 14.1.4. We refer to the fiber $\mathcal{O}_{\langle 1 \rangle} := p_{\mathcal{O}}^{-1}(\langle 1 \rangle)$ over the 1-element set $\langle 1 \rangle := \{1\}$ as the *underlying ∞ -category* of the ∞ -operad $\mathcal{O}$. We denote its groupoid core by

$$\mathcal{O}^\simeq := (\mathcal{O}_{\langle 1 \rangle})^\simeq$$

and refer to it as the *anima of colors of $\mathcal{O}$* . In the classical notation of Definition 12.2.1, $\mathcal{O}^\approx$ denoted only the set of colors; here it also remembers the invertible unary morphisms between them. For a finite set I , condition (2) guarantees that every object in $\mathcal{O}_I^\approx$ may be uniquely written as a product $\prod_{i \in I} x_i$ for some colors $x_i \in \mathcal{O}^\approx$. In analogy with the classical case, we may also denote such a product as an unordered tuple $\{x_i\}_{i \in I}$. Given another color $y \in \mathcal{O}^\approx$, we define the *anima of multimorphisms in $\mathcal{O}$* from $\{x_i\}_{i \in I}$ to y as the anima of morphisms in $\mathcal{O}^\approx$ that map to the span $I \xleftarrow{=} I \rightarrow \langle 1 \rangle$:

$$\begin{array}{ccc} \mathcal{O}(\{x_i\}_{i \in I}; y) & \longrightarrow & \mathrm{Hom}_{\mathcal{O}^\approx}(\{x_i\}_{i \in I}, y) \\ \downarrow & \lrcorner & \downarrow p_{\mathcal{O}} \\ * & \longrightarrow & \mathrm{Hom}_{\mathrm{Span}(\mathrm{Fin})}(I, \langle 1 \rangle). \end{array}$$

A morphism $\varphi: X \rightarrow Y$ in $\mathcal{O}^\approx$ whose image in $\mathrm{Span}(\mathrm{Fin})$ is a forward map $I \xleftarrow{=} I \xrightarrow{g} J$ is called an *active morphism*. We say φ is an *inert morphism* if it is $p_{\mathcal{O}}$ -cocartesian and its image in $\mathrm{Span}(\mathrm{Fin})$ is a backwards map $I \xleftarrow{f} J \xrightarrow{=} J$. We want to think of the active morphisms as the ones that encode the operad structure, and of the inert morphisms as merely capturing the product structure in $\mathcal{O}^\approx$.

Definition 14.1.5. A morphism of ∞ -operads $f: \mathcal{O} \rightarrow \mathcal{P}$ is called *fully faithful* if for every finite set I and every collection of colors $\{x_i\}_{i \in I}$ and y of $\mathcal{O}$, the induced map

$$\mathcal{O}(\{x_i\}_{i \in I}; y) \rightarrow \mathcal{P}(\{f(x_i)\}_{i \in I}; f(y))$$

is an equivalence of anima.

Lemma 14.1.6. *Let $f: \mathcal{O} \rightarrow \mathcal{P}$ be a morphism of ∞ -operads. If f is fully faithful and the induced functor $\mathcal{O}_{\langle 1 \rangle} \rightarrow \mathcal{P}_{\langle 1 \rangle}$ is essentially surjective, then f is an equivalence of ∞ -operads.*

Proof. For every finite set I , the fiber $\mathcal{O}_I^\approx$ is equivalent to $\mathcal{O}_{\langle 1 \rangle}^I$, and similarly for $\mathcal{P}$. It follows that $f^\approx: \mathcal{O}^\approx \rightarrow \mathcal{P}^\approx$ is essentially surjective on objects. It remains to see that it is fully faithful.

Let $\{x_i\}_{i \in I}$ and $\{y_j\}_{j \in J}$ be objects of $\mathcal{O}^\approx$, and let α be a span $I \xleftarrow{u} K \xrightarrow{v} J$. By factoring α as a backwards span followed by a forward span and using inert cocartesian transport, the anima of morphisms over α is naturally equivalent to

$$\prod_{j \in J} \mathcal{O}(\{x_{u(k)}\}_{k \in v^{-1}(j)}; y_j).$$

The same description holds in $\mathcal{P}$. Since f is fully faithful on multimorphism anima, it induces an equivalence on the fibers of hom anima over every span α . It follows that $f^\approx$ is fully faithful. Together with essential surjectivity, this shows that $f^\approx$ is an equivalence over $\mathrm{Span}(\mathrm{Fin})$, as claimed. $\square$

14.1.1 An alternative characterization of ∞ -operads

Our definition of ∞ -operads was formulated in terms of finite products in the total category $\mathcal{O}^\approx$. We now discuss an alternative characterization which does not make any a priori assumptions on products in $\mathcal{O}^\approx$ and instead uses a formulation in terms of cocartesian morphisms. This will be our main practical criterion for recognizing ∞ -operads. It is used in the next section to produce the ∞ -operad associated to a symmetric monoidal ∞ -category, and in Section 17.4 to compare our definition of ∞ -operads to Lurie's definition.

Proposition 14.1.7. *Consider a functor $p_O: O^\otimes \rightarrow \text{Span}(\text{Fin})$. Then $(O^\otimes, p_O)$ is an ∞ -operad if and only if the following conditions are satisfied:*

- (i) $O^\otimes$ admits all p_O -cocartesian lifts of backwards morphisms $I \xleftarrow{f} J \xrightarrow{\cong} J$ in $\text{Span}(\text{Fin})$;
- (ii) For every finite set I , the functor

$$O_I^\otimes \rightarrow \prod_{i \in I} O_{\{i\}}^\otimes$$

given by cocartesian transport along the maps $\rho_i: I \hookrightarrow \{i\} \xrightarrow{\cong} \{i\}$ in $\text{Span}(\text{Fin})$ is an equivalence, where the transport functors are supplied by Proposition 23.1.4;

- (iii) Consider an object $Y \in O^\otimes$ with image $J := p_O(Y) \in \text{Span}(\text{Fin})$ and cocartesian lifts $Y \rightarrow Y_j$ over the maps ρ_j in $\text{Span}(\text{Fin})$. Then the maps $Y \rightarrow Y_j$ exhibit Y as a product $\prod_{j \in J} Y_j$ in $O^\otimes$.

Proof. Throughout the proof we write $p = p_O$ for simplicity. First assume that $O^\otimes$ is an ∞ -operad. We show that (i)–(iii) are satisfied:

- (i) Consider a morphism $f: J \rightarrow I$ in Fin and consider $X \in O_I^\otimes$. By condition (2), we may uniquely write X as a product $\prod_{i \in I} X_i$ of objects $X_i \in O_{\{i\}}^\otimes$. We then define $Y := \prod_{j \in J} X_{f(j)}$ and define $\tilde{f}: X \rightarrow Y$ to be the map whose j -th component $X = \prod_{i \in I} X_i \rightarrow X_{f(j)}$ is the projection map onto $X_{f(j)}$. Then assumption (3) tells us that $\tilde{f}$ is a p -cocartesian lift of the span $I \xleftarrow{f} J \xrightarrow{\cong} J$.
- (iii) Using (2), we may write the object $Y \in O_J^\otimes$ uniquely as a product $\prod_{j \in J} Y_j$. By assumption (3), the projection maps $Y = \prod_{j \in J} Y_j \rightarrow Y_j$ are cocartesian lifts of the maps $\rho_j: J \hookrightarrow \{j\} \xrightarrow{\cong} \{j\}$ in $\text{Span}(\text{Fin})$, verifying (iii).
- (ii) To show that the map $O_I^\otimes \rightarrow \prod_{i \in I} O_{\{i\}}^\otimes$ is an equivalence, it suffices to show it is a left-inverse to the product functor $\prod_{i \in I} O_{\{i\}}^\otimes \xrightarrow{\cong} O_I^\otimes$, as this was assumed to be an equivalence. This is again a consequence of the fact that the projection maps $\prod_{i \in I} X_i \rightarrow X_i$ are cocartesian lifts of ρ_i .

We now show that conditions (i)–(iii) imply that $O^\otimes$ is an ∞ -operad. We check conditions (1)–(3):

- (1) For finite products in $O^\otimes$, consider objects $Y^1, \dots, Y^n$ in $O^\otimes$ and set $T_i := p(Y^i)$ and $T := \bigsqcup_i T_i$. Picking cocartesian lifts $Y^i \rightarrow Y_t^i$ over each map ρ_t , condition (iii) tells us that these maps exhibit each Y^i as a product $\prod_{t \in T_i} Y_t^i$. Using (ii), we may pick some object $Y \in O_T^\otimes$ whose cocartesian transport along each ρ_t is Y_t^i , and we similarly see that the maps $Y \rightarrow Y_t^i$ exhibit Y as a product $\prod_{i=1}^n \prod_{t \in T_i} Y_t^i$. Consequently

$$\text{Hom}_{O^\otimes}(Z, Y) \simeq \prod_{i=1}^n \prod_{t \in T_i} \text{Hom}_{O^\otimes}(Z, Y_t^i) \simeq \prod_{i=1}^n \text{Hom}_{O^\otimes}(Z, Y^i)$$

for every $Z \in O^\otimes$, so $Y \simeq \prod_{i=1}^n Y^i$. Thus $O^\otimes$ admits finite products. It is also clear from this calculation that $p: O^\otimes \rightarrow \text{Span}(\text{Fin})$ preserves products.

- (2) It will suffice to argue that the product functor $\prod_{s \in S} \mathcal{O}_s^\otimes \rightarrow \mathcal{O}_S^\otimes$ is a right-inverse to the assumed equivalence $\mathcal{O}_S^\otimes \xrightarrow{\sim} \prod_{s \in S} \mathcal{O}_{\{s\}}^\otimes$ from (ii). But (iii) implies that for $X_s \in \mathcal{O}_{\{s\}}^\otimes$ the projection maps $\prod_{s \in S} X_s \rightarrow X_s$ are the cocartesian lifts of the ρ_s , which gives the claim.
- (3) Consider a morphism $f: J \rightarrow I$ and let $X_i \in \mathcal{O}^\otimes$. We have to show that the map $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ is p -cocartesian.

Let us first prove the special case where $p(X_i) = *$ for all $i \in I$. Let $\bar{f}: \prod_{i \in I} X_i \rightarrow Y$ be a cocartesian lift of f . By the universal property of cocartesian lifts, there is a morphism $Y \rightarrow \prod_{j \in J} X_{f(j)}$ which we wish to show is an equivalence. Note that this map lives in the fiber over J , so by condition (2) it suffices to check this after applying cocartesian transport along each of the maps ρ_j . Note that we have $(\rho_j)!Y \simeq X_{f(j)}$ since the composites of the spans

$$I \xleftarrow{f} J \xrightarrow{\sim} J \quad \text{and} \quad \rho_j: J \hookrightarrow \{j\} \rightarrow *$$

is the span $\rho_{f(j)}: I \hookrightarrow \{f(j)\} \xrightarrow{\sim} \{f(j)\}$. The projections from $\prod_{j \in J} X_{f(j)}$ are cocartesian lifts of the maps ρ_j by the construction of products in part (1), so also $(\rho_j)!(\prod_{j \in J} X_{f(j)}) \simeq X_{f(j)}$. This proves the claim in the special case.

The general case is an immediate consequence by writing each X_i as a product $\prod_{t \in T_i} X_i^t$ and applying the previous argument to $\prod_{i \in I} \prod_{t \in T_i} X_i^t$. $\square$

It is sometimes convenient to use the following alternative formulation of part (iii).

Notation 14.1.8. Given a functor $p_O: \mathcal{O}^\otimes \rightarrow \text{Span}(\text{Fin})$ and a morphism $\alpha: I \rightarrow J$ in $\text{Span}(\text{Fin})$, we write

$$\text{Hom}_{\mathcal{O}^\otimes}^\alpha(X, Y) \quad := \quad \text{fib}_\alpha(p_O: \text{Hom}_{\mathcal{O}^\otimes}(X, Y) \rightarrow \text{Hom}_{\text{Span}(\text{Fin})}(I, J))$$

for all $X, Y \in \mathcal{O}^\otimes$ with $p_O(X) \cong I$ and $p_O(Y) \cong J$.

Lemma 14.1.9. *Given a functor $p_O: \mathcal{O}^\otimes \rightarrow \text{Span}(\text{Fin})$ and cocartesian morphisms $Y \rightarrow Y_j$ over the maps ρ_j , these maps exhibit Y as a product of the Y_j in $\mathcal{O}^\otimes$ if and only if for every $X \in \mathcal{O}_I^\otimes$ and every $\alpha \in \text{Hom}_{\text{Span}(\text{Fin})}(I, J)$ they induce an equivalence*

$$\text{Hom}_{\mathcal{O}^\otimes}^\alpha(X, Y) \rightarrow \prod_{j \in J} \text{Hom}_{\mathcal{O}^\otimes}^{\rho_j \circ \alpha}(X, Y_j).$$

Proof. The maps $Y \rightarrow Y_j$ exhibit Y as a product if and only if for every $X \in \mathcal{O}_I^\otimes$ the top map in the following square is an equivalence:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{O}^\otimes}(X, Y) & \longrightarrow & \prod_{j \in J} \text{Hom}_{\mathcal{O}^\otimes}(X, Y_j) \\ p_O \downarrow & & \downarrow p_O \\ \text{Hom}_{\text{Span}(\text{Fin})}(I, J) & \longrightarrow & \prod_{j \in J} \text{Hom}_{\text{Span}(\text{Fin})}(I, \{j\}). \end{array}$$

Since the bottom map is an equivalence, this is equivalent to the square being a pullback square, which is in turn equivalent to the condition that the top map induces isomorphisms on all fibers over $\alpha \in \text{Hom}_{\text{Span}(\text{Fin})}(I, J)$. $\square$

Corollary 14.1.10. *If O is an ∞ -operad, then the inert morphisms in $O^\otimes$ are precisely the morphisms of the form $\tilde{f}: \prod_{i \in I} x_i \rightarrow \prod_{j \in J} x_{f(j)}$ for maps $f: J \rightarrow I$ in $\mathbf{Fin}$ and colors $x_i \in O^\otimes$.*

Similarly, if $\mathcal{P}$ is another ∞ -operad, and $\varphi^\otimes: O^\otimes \rightarrow \mathcal{P}^\otimes$ is a functor over $\mathbf{Span}(\mathbf{Fin})$, then $\varphi^\otimes$ preserves finite products (i.e., corresponds to a morphism $\varphi: O \rightarrow \mathcal{P}$ of ∞ -operads) if and only if it preserves the inert morphisms.

Proof. By assumption $\tilde{f}$ is inert for every f . Conversely, we saw in the proof of (i) of the previous proposition that for every backwards span $I \xleftarrow{f} J \xrightarrow{g} J$ and every $X \in O_I^\otimes$ the map $\tilde{f}: X \simeq \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ is a cocartesian lift of f starting in X , so by uniqueness it follows that every cocartesian lift is of this form.

If $\varphi^\otimes$ preserves finite products, it in particular preserves the maps $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$, hence all inert morphisms. Conversely, if $\varphi^\otimes$ preserves inert morphisms, it preserves the projection maps $Y \rightarrow Y_j$ for $Y \in O_J^\otimes$ exhibiting Y as a product of the Y_j , hence it preserves every nonempty finite product. It also preserves the terminal object: since $\varphi^\otimes$ lies over $\mathbf{Span}(\mathbf{Fin})$, it carries the unique object of $O_\emptyset^\otimes \simeq *$ to the unique object of $\mathcal{P}_\emptyset^\otimes \simeq *$. $\square$

Remark 14.1.11. Let $O = (O^\otimes, p_O)$ be an ∞ -operad. Then every morphism $\varphi: X \rightarrow Z$ in $O^\otimes$ factors uniquely as

$$X \xrightarrow{\tilde{f}} Y \xrightarrow{\psi} Z,$$

where $\tilde{f}$ is an inert morphism and ψ is an active morphism. To see this, let us denote by $I \xleftarrow{f} K \xrightarrow{g} J$ the image of φ under p . This span may be factored uniquely as a composite of a backwards span $I \xleftarrow{f} K \xrightarrow{g} K$ followed by a forward span $K \xleftarrow{g} K \xrightarrow{g} J$. We then let $\tilde{f}: X \rightarrow Y$ be the unique cocartesian lift of f with domain X . It follows that there is a unique morphism $\psi: Y \rightarrow Z$ over g such that $\psi \circ \tilde{f} = \varphi$.

The argument above produces a factorization and shows that any two of them are isomorphic. With more care, it in fact shows that the factorization is unique, in the sense that the *anima of factorizations* of φ is contractible. Rather than spelling this out by hand, we will obtain it in Section 17.1 from the general theory of *factorization systems* on an ∞ -category, where we show that the inert and active morphisms define such a system on $O^\otimes$.

14.1.2 Examples of ∞ -operads

The examples of ∞ -operads that we can write down directly at this point are of a rather simple nature; the more interesting ones require the constructions developed in the remainder of this part.

Example 14.1.12. The *commutative ∞ -operad* $Comm$ is given by $Comm^\otimes = \mathbf{Span}(\mathbf{Fin})$ and $p_{Comm} = \text{id}: \mathbf{Span}(\mathbf{Fin}) \rightarrow \mathbf{Span}(\mathbf{Fin})$. This is the terminal ∞ -operad.

Example 14.1.13. The ∞ -operad $\mathcal{E}_0$ of *pointed objects* is given by

$$\mathcal{E}_0^\otimes := \mathbf{Span}_{\text{all, inj}}(\mathbf{Fin}) \quad \text{and} \quad p_{\mathcal{E}_0} = \text{incl}: \mathbf{Span}_{\text{all, inj}}(\mathbf{Fin}) \hookrightarrow \mathbf{Span}(\mathbf{Fin}).$$

The active maps in $\mathcal{E}_0$ are the injections $I \hookrightarrow J$ of finite sets. The finite-product conditions in the definition of an ∞ -operad follow from part (1) of Lemma 13.3.5; the fibers of $p_{\mathcal{E}_0}$ are contractible, and the backwards spans remain cocartesian under the inclusion into $\mathbf{Span}(\mathbf{Fin})$.

Example 14.1.14. By Corollary 17.4.10, the classical associative operad determines the associative ∞ -operad $\mathcal{A}ssoc$. The topological little n -cubes operad of Example 12.1.7 similarly determines an ∞ -operad $\mathcal{E}_n$; see [Lur17, Section 5.1.0] for the construction. For $n = 1$, the operad morphism $\mathcal{E}_1 \rightarrow \mathcal{A}ssoc$ from Exercise 12.6 is an equivalence of ∞ -operads by [Lur17, Example 5.1.0.7].

14.1.3 Trivial operads

The simplest ∞ -operads are those without any operations beyond the unary ones, generalizing the trivial colored operads of Example 12.2.6. Given an ∞ -category C , we will construct an ∞ -operad $\mathcal{T}riv_C$ with underlying ∞ -category C and multimorphism animae

$$\mathcal{T}riv_C(\{x_i\}_{i \in I}; y) \simeq \begin{cases} \mathrm{Hom}_C(x_{i_0}, y) & \text{if } I = \{i_0\} \text{ is a singleton;} \\ \emptyset & \text{otherwise,} \end{cases}$$

with composition given by composition in C . There is no operadic structure left to specify: operations of arity $\neq 1$ do not exist, and the unary operations are simply the morphisms of C . To construct $\mathcal{T}riv_C$, note that multimorphisms from $\{x_i\}_{i \in I}$ to y live over the active span $I \xleftarrow{=} I \rightarrow \langle 1 \rangle$, which lies in the subcategory $\mathrm{Fin}^{\mathrm{op}} \subseteq \mathrm{Span}(\mathrm{Fin})$ precisely when I is a singleton. We should therefore look for an ∞ -operad whose structure map factors through the inclusion $\mathrm{Fin}^{\mathrm{op}} \hookrightarrow \mathrm{Span}(\mathrm{Fin})$.

The construction is based on the following ∞ -category of finite families, which will reappear in Section 15.2 as the basis for cocartesian monoidal structures.

Construction 14.1.15 (Finite families). Let C be an ∞ -category. There is a unique finite-product-preserving functor

$$C^{(-)}: \mathrm{Fin}^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}$$

which sends the one-point set to C ; explicitly, it sends a finite set I to the ∞ -category C^I of I -indexed families and a map $f: I \rightarrow J$ to the restriction functor $f^*: C^J \rightarrow C^I$. Equivalently, this functor is the right Kan extension of C from the one-point set. We write

$$q: \mathrm{Fin}(C) \rightarrow \mathrm{Fin}$$

for its cartesian unstraightening. Thus objects of $\mathrm{Fin}(C)$ are finite unordered tuples $\{x_i\}_{i \in I}$, while morphisms $\{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ are pairs $(f, (\varphi_i)_{i \in I})$ consisting of a map $f: I \rightarrow J$ and morphisms $\varphi_i: x_i \rightarrow y_{f(i)}$ in C for all $i \in I$.

We record the following basic observations about the ∞ -category of finite families:

- The fiber of q over $\langle 1 \rangle$ is equivalent to C , and the resulting inclusion $i: C \hookrightarrow \mathrm{Fin}(C)$ is fully faithful.
- A morphism $(f, (\varphi_i)_{i \in I}): \{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ is q -cartesian if and only if each φ_i is an isomorphism in C . We denote by

$$\mathrm{Fin}(C)_{\mathrm{ct}} \subseteq \mathrm{Fin}(C)$$

the wide subcategory spanned by the q -cartesian morphisms.

- The ∞ -category $\text{Fin}(C)$ admits finite coproducts, given by concatenation of unordered tuples, and the functor $q: \text{Fin}(C) \rightarrow \text{Fin}$ preserves finite coproducts. More precisely, given a finite collection of objects $X_a \in \text{Fin}(C)$ for $a \in A$, we may uniquely write each X_a as $\{x_j^a\}_{j \in J_a}$ for some finite set J_a , and then the coproduct is given by $X = \{x_j^a\}_{(a,j) \in \bigsqcup_{a \in A} J_a}$, with the maps $X_a \rightarrow X$ given by the q -cartesian morphisms over the inclusions $J_a \hookrightarrow \bigsqcup_{a \in A} J_a$. This follows from Corollary 23.2.9.

Together with the fully faithful inclusion i , these coproducts exhibit $\text{Fin}(C)$ as the free finite-coproduct completion of C :

Lemma 14.1.16. *Let C be an ∞ -category. For every ∞ -category D with finite coproducts, restriction along the fully faithful inclusion $i: C \hookrightarrow \text{Fin}(C)$ induces an equivalence*

$$i^*: \text{Fun}^{\sqcup}(\text{Fin}(C), D) \xrightarrow{\sim} \text{Fun}(C, D),$$

where $\text{Fun}^{\sqcup}(-, -)$ denotes the full subcategory of functors preserving finite coproducts.

Proof. We claim that an inverse is given by left Kan extension.

Step 1: Given a functor $F: C \rightarrow D$, we show that its left Kan extension $i_! F$ along i exists. By the pointwise formula for left Kan extensions, it suffices to show that each of the colimits

$$(i_! F)(x_1, \dots, x_n) = \text{colim}_{y \in C_{/(x_1, \dots, x_n)}} F(y)$$

exists. Observe that the canonical map

$$\bigsqcup_{k=1}^n C_{/x_k} \rightarrow C_{/(x_1, \dots, x_n)} := C \times_{\text{Fin}(C)} \text{Fin}(C)_{/(x_1, \dots, x_n)}$$

sending $(\varphi: y \rightarrow x_k)$ to the map $(\{k\} \hookrightarrow \langle n \rangle, \varphi)$ is an equivalence. Since D has finite coproducts, we see that colimits indexed by the slices $C_{/(x_1, \dots, x_n)}$ exist in D if and only if colimits indexed by each $C_{/x_k}$ exist. But the latter is clear: as $C_{/x_k}$ admits a terminal object, colimits always exist and are given by evaluation at that terminal object. This shows that $i_! F$ exists and that it is given on objects by

$$(i_! F)(x_1, \dots, x_n) = F(x_1) \sqcup \dots \sqcup F(x_n).$$

Step 2: It follows from the pointwise description of $i_! F$ that it preserves finite coproducts. In particular, the restriction functor thus admits a left adjoint

$$i_!: \text{Fun}(C, D) \rightarrow \text{Fun}^{\sqcup}(\text{Fin}(C), D).$$

Step 3: We now show that the unit $\text{id} \rightarrow i^* i_!$ and counit $i_! i^* \rightarrow \text{id}$ are natural isomorphisms. The unit is a natural isomorphism because i is fully faithful. If $G: \text{Fin}(C) \rightarrow D$ preserves finite coproducts, then the counit evaluated at $(x_1, \dots, x_n)$ is the canonical map

$$G(x_1) \sqcup \dots \sqcup G(x_n) \longrightarrow G(x_1, \dots, x_n),$$

which is an isomorphism because $(x_1, \dots, x_n)$ is the coproduct of the singleton families (x_i) in $\text{Fin}(C)$. $\square$

Construction 14.1.17 (Trivial operads). Let C be an ∞ -category. Applying Construction 14.1.15 to C^{op} gives a cartesian fibration $q: \text{Fin}(C^{\text{op}}) \rightarrow \text{Fin}$; passing to opposite ∞ -categories, we obtain a functor $q^{\text{op}}: \text{Fin}(C^{\text{op}})^{\text{op}} \rightarrow \text{Fin}^{\text{op}}$. We define

$$\mathcal{Triv}_C^{\otimes} := \text{Fin}(C^{\text{op}})^{\text{op}}, \quad p_{\mathcal{Triv}_C}: \text{Fin}(C^{\text{op}})^{\text{op}} \xrightarrow{q^{\text{op}}} \text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin}),$$

and refer to $\mathcal{Triv}_C$ as the *trivial ∞ -operad generated by C* . Unwinding Construction 14.1.15, the objects of $\mathcal{Triv}_C^{\otimes}$ are finite unordered tuples $\{x_i\}_{i \in I}$ of objects of C , while a morphism $\{x_i\}_{i \in I} \rightarrow \{y_j\}_{j \in J}$ lying over the backwards span $I \xleftarrow{g} J \xrightarrow{=} J$ is a collection of morphisms $x_{g(j)} \rightarrow y_j$ in C . The opposite in the definition ensures that the underlying ∞ -category is C itself, and not C^{op} .

Remark 14.1.18. Equivalently, $q^{\text{op}}: \mathcal{Triv}_C^{\otimes} \rightarrow \text{Fin}^{\text{op}}$ is the *cocartesian* unstraightening of the functor $C^{(-)}$ from Construction 14.1.15: it is a cocartesian fibration and its cocartesian straightening $\text{Fin}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ preserves finite products and sends the point to C .

Lemma 14.1.19. *Let C be an ∞ -category. Then $\mathcal{Triv}_C$ is an ∞ -operad with underlying ∞ -category $(\mathcal{Triv}_C)_{\langle 1 \rangle} \simeq C$, and its multimorphism animae are the ones described at the beginning of this subsection. Moreover, for every ∞ -operad O , restriction to the underlying ∞ -categories induces an equivalence*

$$\text{Fun}_{\text{Op}_{\infty}}(\mathcal{Triv}_C, O) \xrightarrow{\sim} \text{Fun}(C, O_{\langle 1 \rangle}).$$

Proof. Let us first check that $\mathcal{Triv}_C$ is an ∞ -operad. By Construction 14.1.15, the ∞ -category $\text{Fin}(C^{\text{op}})$ admits finite coproducts and q preserves them, so $\mathcal{Triv}_C^{\otimes} = \text{Fin}(C^{\text{op}})^{\text{op}}$ admits finite products and q^{op} preserves them; as the inclusion $\text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin})$ preserves finite products by part (1) of Lemma 13.3.5, the same is true for $p_{\mathcal{Triv}_C}$. Since the products of $\mathcal{Triv}_C^{\otimes}$ are the concatenations of tuples, the fiber over a finite set I is C^I , whose decomposition $C^I \simeq \prod_{i \in I} C$ is the one required in the definition of an ∞ -operad.

For the final condition, consider objects $X_i = \{x_a^i\}_{a \in A_i}$ of $\mathcal{Triv}_C^{\otimes}$ for $i \in I$ and a map $f: J \rightarrow I$ of finite sets. Read in $\text{Fin}(C^{\text{op}})$, the morphism $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ is the map $\coprod_{j \in J} X_{f(j)} \rightarrow \coprod_{i \in I} X_i$ which restricts on the j -th summand to the inclusion of $X_{f(j)}$; it lies over the map $\coprod_{j \in J} A_{f(j)} \rightarrow \coprod_{i \in I} A_i$, $(j, a) \mapsto (f(j), a)$, and all of its components in C^{op} are identities. It is therefore q -cartesian, and hence $\tilde{f}$ is q^{op} -cocartesian. It remains $p_{\mathcal{Triv}_C}$ -cocartesian after composing with the inclusion $\text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin})$: every morphism of $\mathcal{Triv}_C^{\otimes}$ lies over a backwards span, and composing a non-backwards span with a backwards span is again non-backwards, so the mapping animae over the remaining spans are empty on both sides.

The description of the multimorphism animae follows: the active span $I \xleftarrow{=} I \rightarrow \langle 1 \rangle$ lies in Fin^{op} only when I is a singleton, in which case it is the identity of $\langle 1 \rangle$. Thus $\mathcal{Triv}_C(\{x_i\}_{i \in I}; y)$ is empty unless $I = \{i_0\}$, in which case it is the anima of morphisms $x_{i_0} \rightarrow y$ in the fiber $(\mathcal{Triv}_C)_{\langle 1 \rangle} \simeq C$.

We turn to the universal property, which will follow from the universal property of $\text{Fin}(C^{\text{op}})$. Indeed, applying Lemma 14.1.16 to C^{op} and passing to opposite categories, we see that restriction along the inclusion $j: C \hookrightarrow \mathcal{Triv}_C^{\otimes}$ of the fiber over $\langle 1 \rangle$ induces an equivalence

$$j^*: \text{Fun}^{\times}(\mathcal{Triv}_C^{\otimes}, E) \xrightarrow{\sim} \text{Fun}(C, E)$$

for every ∞ -category E that admits finite products. This equivalence is natural in E , so applying it to $E = \mathcal{O}^\otimes$ and to $E = \text{Span}(\text{Fin})$ yields a commutative square

$$\begin{array}{ccc} \text{Fun}^\times(\mathcal{T}\text{riv}_C^\otimes, \mathcal{O}^\otimes) & \xrightarrow[\simeq]{J^*} & \text{Fun}(C, \mathcal{O}^\otimes) \\ p_{\mathcal{O}} \circ - \downarrow & & \downarrow p_{\mathcal{O}} \circ - \\ \text{Fun}^\times(\mathcal{T}\text{riv}_C^\otimes, \text{Span}(\text{Fin})) & \xrightarrow[\simeq]{J^*} & \text{Fun}(C, \text{Span}(\text{Fin})). \end{array}$$

The composite $p_{\mathcal{T}\text{riv}_C} \circ j$ is the constant functor at $\langle 1 \rangle$, so passing to the fibers over $p_{\mathcal{T}\text{riv}_C}$ and $\text{const}_{\langle 1 \rangle}$ gives equivalences

$$\text{Fun}_{\text{Op}_\infty}(\mathcal{T}\text{riv}_C, \mathcal{O}) \xrightarrow{\sim} \text{Fun}(C, \mathcal{O}^\otimes) \times_{\text{Fun}(C, \text{Span}(\text{Fin}))} \{\text{const}_{\langle 1 \rangle}\} \simeq \text{Fun}(C, \mathcal{O}_{\langle 1 \rangle}),$$

where the second equivalence holds because $\text{Fun}(C, -)$ preserves pullbacks. By construction, this equivalence is given by restriction to the underlying ∞ -categories. $\square$

Note that the construction $C \mapsto \mathcal{T}\text{riv}_C$ is functorial in C and thus defines a functor $\mathcal{T}\text{riv}_{(-)} : \text{Cat}_\infty \rightarrow \text{Op}_\infty$. It may be characterized as follows:

Corollary 14.1.20. *The functor $\mathcal{T}\text{riv}_{(-)} : \text{Cat}_\infty \rightarrow \text{Op}_\infty$ is left adjoint to the functor*

$$(-)_{\langle 1 \rangle} : \text{Op}_\infty \rightarrow \text{Cat}_\infty, \quad \mathcal{O} \mapsto \mathcal{O}_{\langle 1 \rangle},$$

and it is fully faithful.

Proof. The equivalence of Lemma 14.1.19 is given by passing to underlying ∞ -categories and precomposing with the isomorphism $C \xrightarrow{\sim} (\mathcal{T}\text{riv}_C)_{\langle 1 \rangle}$, and is thus natural in both variables. Passing to groupoid cores yields the corresponding equivalence of mapping animae, so it exhibits $\mathcal{T}\text{riv}_{(-)}$ as a left adjoint of $(-)_{\langle 1 \rangle}$. Taking $\mathcal{O} = \mathcal{T}\text{riv}_D$ for an ∞ -category D , the equivalence

$$\text{Fun}_{\text{Op}_\infty}(\mathcal{T}\text{riv}_C, \mathcal{T}\text{riv}_D) \xrightarrow{\sim} \text{Fun}(C, (\mathcal{T}\text{riv}_D)_{\langle 1 \rangle}) \simeq \text{Fun}(C, D)$$

is inverse to the map induced by $\mathcal{T}\text{riv}_{(-)}$, which is therefore fully faithful. $\square$

Example 14.1.21. Taking $C = *$ recovers the *trivial ∞ -operad*

$$\mathcal{T}\text{riv} := \mathcal{T}\text{riv}_*, \quad \mathcal{T}\text{riv}^\otimes = \text{Fin}^{\text{op}}, \quad p_{\mathcal{T}\text{riv}} = \text{incl} : \text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin}).$$

Taking $C = \emptyset$ gives the *empty ∞ -operad* $\mathcal{T}\text{riv}_\emptyset$, with $\mathcal{T}\text{riv}_\emptyset^\otimes = *$ and $p_{\mathcal{T}\text{riv}_\emptyset} : * \xrightarrow{\langle 0 \rangle} \text{Span}(\text{Fin})$ the inclusion of the empty set $\langle 0 \rangle$. Since $\mathcal{T}\text{riv}_{(-)}$ is a left adjoint by Corollary 14.1.20, it preserves initial objects; as $\emptyset$ is initial in Cat_∞ , we conclude that $\mathcal{T}\text{riv}_\emptyset$ is the initial ∞ -operad.

Lemma 14.1.22. *For every ∞ -operad $\mathcal{O}$, there is a canonical equivalence*

$$\mathcal{T}\text{riv} \times \mathcal{O} \simeq \mathcal{T}\text{riv}_{\mathcal{O}_{\langle 1 \rangle}}$$

of ∞ -operads.

Proof. Recall that $\mathcal{T}riv^\otimes = \mathbf{Fin}^{\mathrm{op}}$ and that $p_{\mathcal{T}riv}: \mathbf{Fin}^{\mathrm{op}} \hookrightarrow \mathbf{Span}(\mathbf{Fin})$ is the inclusion. The product operad $\mathcal{T}riv \times \mathcal{O}$ is therefore represented by the pullback

$$\begin{array}{ccc} \mathbf{Fin}^{\mathrm{op}} \times_{\mathbf{Span}(\mathbf{Fin})} \mathcal{O}^\otimes & \longrightarrow & \mathcal{O}^\otimes \\ \downarrow & \lrcorner & \downarrow p_{\mathcal{O}} \\ \mathbf{Fin}^{\mathrm{op}} & \hookrightarrow & \mathbf{Span}(\mathbf{Fin}). \end{array}$$

By the defining properties of an ∞ -operad, the left vertical map is a cocartesian fibration whose cocartesian straightening preserves finite products and sends $\langle 1 \rangle$ to $\mathcal{O}_{\langle 1 \rangle}$. It is therefore equivalent to

$$\mathbf{Fin}^{\mathrm{op}} \rightarrow \mathbf{Cat}_\infty, \quad I \mapsto \mathcal{O}_{\langle 1 \rangle}^I.$$

By Remark 14.1.18, its cocartesian unstraightening is precisely

$$q_{\mathcal{O}_{\langle 1 \rangle}}^{\mathrm{op}}: \mathbf{Fin}(\mathcal{O}_{\langle 1 \rangle}^{\mathrm{op}})^{\mathrm{op}} \rightarrow \mathbf{Fin}^{\mathrm{op}},$$

which is the total category of $\mathcal{T}riv_{\mathcal{O}_{\langle 1 \rangle}}$. □

Remark 14.1.23. More generally, restricting an arbitrary ∞ -operad $\mathcal{O}$ to the subcategory $\mathbf{Fin}^{\mathrm{op}} \subseteq \mathbf{Span}(\mathbf{Fin})$ discards precisely the operations of arity $\neq 1$: by Lemma 14.1.22, the pullback $\mathcal{O}^\otimes \times_{\mathbf{Span}(\mathbf{Fin})} \mathbf{Fin}^{\mathrm{op}}$ is the total category of $\mathcal{T}riv_{\mathcal{O}_{\langle 1 \rangle}}$. The resulting projection $\mathcal{T}riv_{\mathcal{O}_{\langle 1 \rangle}} \rightarrow \mathcal{O}$ induces the identity on underlying ∞ -categories, and is thus the counit of the adjunction of Corollary 14.1.20.

14.2 Symmetric monoidal ∞ -categories and algebras over operads

Let $\mathcal{O}$ be a colored operad and let C be a symmetric monoidal 1-category. Recall that an $\mathcal{O}$ -algebra in C is defined as an operad map $\mathcal{O} \rightarrow \mathcal{M}_C$ into the *multimorphism operad* of C , whose multimorphisms are given by

$$\mathcal{M}_C(\{x_i\}_{i \in I}; y) := \mathrm{Hom}_C\left(\bigotimes_{i \in I} x_i, y\right);$$

see Definition 12.2.10. The goal of this section is to define the analogous notion of algebras over operads in the ∞ -categorical setting.

14.2.1 Symmetric monoidal ∞ -categories as ∞ -operads

We start by constructing the multimorphism operad $\mathcal{M}_C$ of a symmetric monoidal ∞ -category C . Recall from Definition 8.1.1 that a symmetric monoidal ∞ -category is a commutative monoid in $\mathbf{Cat}_\infty$, i.e. a finite-product-preserving functor $C: \mathbf{Span}(\mathbf{Fin}) \rightarrow \mathbf{Cat}_\infty$. These naturally form an ∞ -category that we denote by

$$\mathbf{Cat}_\infty^\otimes := \mathbf{CMon}(\mathbf{Cat}_\infty).$$

Since the opposite-category involution $(-)^{\mathrm{op}}: \mathbf{Cat}_\infty \rightarrow \mathbf{Cat}_\infty$ preserves finite products, postcomposition with it defines an involution on $\mathbf{Cat}_\infty^\otimes$. For a symmetric monoidal ∞ -category $(C, \otimes)$, its image under this involution is called the *opposite symmetric monoidal ∞ -category* and is denoted $(C, \otimes)^{\mathrm{op}}$.

Given symmetric monoidal ∞ -categories C and D , write $\mathrm{Map}^\otimes(C, D)$ for their hom anima in $\mathbf{Cat}_\infty^\otimes$.

Example 14.2.1 (Terminal category). The terminal ∞ -category $*$ admits a unique symmetric monoidal structure, given by the constant functor $\text{Span}(\text{Fin}) \rightarrow \text{Cat}_\infty, S \mapsto *$.

Example 14.2.2 (Pointwise monoidal structure). If C is a symmetric monoidal ∞ -category and I is an ∞ -category, then $\text{Fun}(I, C)$ admits a symmetric monoidal structure, called the *pointwise monoidal structure*. It is given by the composite

$$\text{Span}(\text{Fin}) \xrightarrow{(C, \otimes)} \text{Cat}_\infty \xrightarrow{\text{Fun}(I, -)} \text{Cat}_\infty.$$

This is functorial in both I and C , resulting in a functor

$$\text{Fun}(-, -): \text{Cat}_\infty^{\text{op}} \times \text{Cat}_\infty^\otimes \rightarrow \text{Cat}_\infty^\otimes.$$

The symmetric monoidal functors and their monoidal natural transformations themselves form an ∞ -category:

Definition 14.2.3. Let C and D be symmetric monoidal ∞ -categories. We define $\text{Fun}^\otimes(C, D)$ to be the ∞ -category whose associated complete Segal anima is

$$N(\text{Fun}^\otimes(C, D))_n := \text{Map}^\otimes(C, \text{Fun}([n], D)),$$

where $\text{Fun}([n], D)$ carries the pointwise symmetric monoidal structure. This is a complete Segal anima because $\text{Map}^\otimes(C, -)$ preserves limits; see Proposition 1.8.6.

Given a symmetric monoidal ∞ -category $C: \text{Span}(\text{Fin}) \rightarrow \text{Cat}_\infty$, we may unstraighten this to a cocartesian fibration $p_C: C^\otimes \rightarrow \text{Span}(\text{Fin})$. The next result implies that this is always an ∞ -operad.

Lemma 14.2.4. *Let $p_O: O^\otimes \rightarrow \text{Span}(\text{Fin})$ be a cocartesian fibration. Then $(O^\otimes, p_O)$ is an ∞ -operad if and only if the cocartesian straightening of p_O*

$$\text{Str}^{\text{cc}}(p_O): \text{Span}(\text{Fin}) \rightarrow \text{Cat}_\infty$$

preserves finite products.

Proof. We check conditions (i)–(iii) of Proposition 14.1.7. Condition (i) is part of the assumption on p_O . Since the spans $\rho_i: I \leftarrow \{i\} \xrightarrow{=} \{i\}$ exhibit I as the product of the one-element sets in $\text{Span}(\text{Fin})$, condition (ii) is equivalent to preservation of finite products by $\text{Str}^{\text{cc}}(p_O)$. Under this assumption, Lemma 23.2.8 shows that the cocartesian lifts of the maps ρ_i exhibit every object over I as the product of its components, which is condition (iii). $\square$

Definition 14.2.5. Given a symmetric monoidal ∞ -category C , we denote its cocartesian unstraightening by

$$C^\otimes := \text{Un}^{\text{cc}}(C: \text{Span}(\text{Fin}) \rightarrow \text{Cat}_\infty).$$

By the previous lemma, the resulting cocartesian fibration $p_C: C^\otimes \rightarrow \text{Span}(\text{Fin})$ defines an ∞ -operad, which we will denote by $\mathcal{M}_C := (C^\otimes, p_C)$ and refer to as the *multimorphism operad* associated to C . By functoriality of unstraightening, this defines a (non-full) inclusion

$$\mathcal{M}: \text{Cat}_\infty^\otimes \hookrightarrow \text{Op}_\infty.$$

Morphisms in its essential image are studied in the next section.

Lemma 14.2.6. *Given $C \in \text{Cat}_\infty^\otimes$ and objects $x_1, \dots, x_n, y \in C^\simeq$, there is a natural equivalence*

$$\mathcal{M}_C((x_1, \dots, x_n); y) \simeq \text{Hom}_C(x_1 \otimes \cdots \otimes x_n, y).$$

Proof. Let α denote the active span $\langle n \rangle \xleftarrow{=} \langle n \rangle \rightarrow \langle 1 \rangle$. Cocartesian transport along α sends $(x_1, \dots, x_n)$ to $x_1 \otimes \cdots \otimes x_n$. The universal property of this cocartesian lift gives

$$\text{Hom}_{C^\otimes}^\alpha((x_1, \dots, x_n), y) \simeq \text{Hom}_C(x_1 \otimes \cdots \otimes x_n, y),$$

and the left-hand side is the indicated multimorphism anima. $\square$

Observation 14.2.7. Cocartesian transport in $C^\otimes$ is compatible with finite products. More precisely, suppose that $\alpha_j: I_j \rightarrow J_j$ are morphisms in $\text{Span}(\text{Fin})$ and that $\varphi_j: X_j \rightarrow Y_j$ are p_C -cocartesian lifts of α_j . Since the straightening of p_C preserves finite products, the product

$$\prod_j \varphi_j: \prod_j X_j \longrightarrow \prod_j Y_j$$

is a p_C -cocartesian lift of the product $\prod_j \alpha_j$. In particular, cocartesian transport along a forward span is computed separately over the elements of its target.

We have the following useful criterion for checking that an ∞ -operad corresponds to a symmetric monoidal ∞ -category:

Lemma 14.2.8. *Let O be an ∞ -operad. Assume that the following two conditions are satisfied:*

- (1) *For every object $\{x_i\}_{i \in I}$ of $O^\otimes$ the functor*

$$O(\{x_i\}_{i \in I}; -): O_{\langle 1 \rangle} \rightarrow \text{An}$$

is corepresentable, in the sense that there exists a multimorphism $\varphi: \{x_i\}_{i \in I} \rightarrow y$ such that for every other color $z \in O^\simeq$ precomposition with φ induces an equivalence

$$\text{Hom}_{O_{\langle 1 \rangle}}(y, z) \xrightarrow{\sim} O(\{x_i\}_{i \in I}; z).$$

This condition uniquely determines y , and we will denote it by $\bigotimes_{i \in I} x_i$.

- (2) *For every morphism $f: I \rightarrow J$ of finite sets, the canonical comparison map*

$$\bigotimes_{i \in I} x_i \rightarrow \bigotimes_{j \in J} \bigotimes_{i \in f^{-1}(j)} x_i$$

is an equivalence.

Then $p_O: O^\otimes \rightarrow \text{Span}(\text{Fin})$ is a cocartesian fibration, corresponding to a symmetric monoidal structure on the underlying ∞ -category $O_{\langle 1 \rangle}^\otimes$.

Proof. We need to produce cocartesian lifts of morphisms in $\text{Span}(\text{Fin})$. Cocartesian lifts of backwards morphisms exist by the defining property of ∞ -operads, so it suffices to produce cocartesian lifts for active spans $\alpha = (I \xleftarrow{=} I \xrightarrow{f} J)$. Let $X = \{x_i\}_{i \in I}$ be an object in $O_I^\otimes$. Applying the assumption on O to each $X_j := \{x_i\}_{i \in f^{-1}(j)}$, we may find multimorphisms $\varphi_j: \{x_i\}_{i \in f^{-1}(j)} \rightarrow y_j$

in $\mathcal{O}$ corepresenting the functor $\mathcal{O}(\{x_i\}_{i \in f^{-1}(j)}; -) : \mathcal{O}_{\langle 1 \rangle} \rightarrow \mathbf{An}$. The multimorphisms φ_j define a morphism $\varphi : X \rightarrow Y := \{y_j\}_{j \in J}$ in $\mathcal{O}^\otimes$. We wish to show that φ is a cocartesian lift of α .

To this end, consider any other span $\beta = (J \xleftarrow{g} K \xrightarrow{h} L)$ of finite sets, and consider an object $Z = \{z_l\}_{l \in L}$ of $\mathcal{O}_L^\otimes$. We need to show that precomposition with φ induces an equivalence

$$- \circ \varphi : \mathrm{Hom}_{\mathcal{O}^\otimes}^\beta(Y, Z) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{O}^\otimes}^{\beta \circ \alpha}(X, Z).$$

Since Z is a product in $\mathcal{O}^\otimes$ of the one-element tuples (z_l) , we may assume that $L = \langle 1 \rangle$, so that there is only a single color $z := z_1$. Using the inert morphisms $\{y_j\} \rightarrow \{y_{g(k)}\}_{k \in K}$ and $\{x_i\}_{(i,k) \in I \times_J K}$, we must equivalently show that composition with the map φ induces an equivalence

$$- \circ \varphi : \mathcal{O}(\{y_{g(k)}\}_{k \in K}; z) \xrightarrow{\sim} \mathcal{O}(\{x_i\}_{(i,k) \in I \times_J K}; z).$$

Now, applying condition (1) another time to the families $\{y_{g(k)}\}_{k \in K}$ and $\{x_i\}_{(i,k) \in I \times_J K}$, both sides are corepresented by objects $\bigotimes_{k \in K} y_{g(k)}$ and $\bigotimes_{(i,k) \in I \times_J K} x_i$, respectively. But since $y_{g(k)} = \bigotimes_{i \in f^{-1}(g(k))} x_i$, the claim now follows from the equivalence

$$\bigotimes_{(i,k) \in I \times_J K} x_i \xrightarrow{\sim} \bigotimes_{k \in K} \bigotimes_{i \in f^{-1}(g(k))} x_i$$

obtained by applying condition (2) to the projection map $I \times_J K \rightarrow K$. Under the two corepresenting equivalences, the map induced by composition with φ is precisely precomposition with this comparison map, by the definition of the latter through operadic composition with the multimorphisms φ_j . Thus φ is cocartesian.

We have now produced cocartesian lifts of all active spans. Together with the inert lifts and the factorization of Remark 14.1.11, these make $p_{\mathcal{O}}$ a cocartesian fibration. Since $\mathcal{O}$ is already an ∞ -operad, Lemma 14.2.4 shows that its straightening preserves finite products, and hence defines the asserted symmetric monoidal structure. $\square$

14.2.2 Algebras over ∞ -operads

Definition 14.2.9. Let $\mathcal{O}$ be an ∞ -operad and let C be a symmetric monoidal ∞ -category. An $\mathcal{O}$ -algebra in C is an operad morphism $\mathcal{O} \rightarrow \mathcal{M}_C$. We write

$$\mathrm{Alg}_{\mathcal{O}}(C) := \mathrm{Fun}_{\mathrm{Op}_{\infty}}(\mathcal{O}, \mathcal{M}_C)$$

for the ∞ -category of $\mathcal{O}$ -algebras in C .

Example 14.2.10 (Commutative/associative algebras). Algebras over the operads Comm and Assoc are referred to as *commutative algebras* and *associative algebras*, respectively, and their ∞ -categories are denoted by

$$\mathrm{CAlg}(C) := \mathrm{Alg}_{\mathit{Comm}}(C) \quad \text{and} \quad \mathrm{Alg}(C) := \mathrm{Alg}_{\mathit{Assoc}}(C).$$

Example 14.2.11 (Trivial algebras). Let D be an ∞ -category. Applying Lemma 14.1.19 to the multimorphism operad $\mathcal{M}_C$, we see that algebras over the trivial operad Triv_D correspond to functors $D \rightarrow C$:

$$\mathrm{Alg}_{\mathit{Triv}_D}(C) \xrightarrow{\sim} \mathrm{Fun}(D, C).$$

In particular, taking $D = *$ shows that a $\mathcal{T}riv$ -algebra is simply an object of C :

$$\mathrm{Alg}_{\mathcal{T}riv}(C) \xrightarrow{\sim} C.$$

Taking $D = \emptyset$ shows that the ∞ -category of $\mathcal{T}riv_{\emptyset}$ -algebras is terminal.

Lemma 14.2.12 (Algebras in pointwise monoidal structure). *Let C be a symmetric monoidal ∞ -category, let I be an ∞ -category, and let O be an ∞ -operad. Then an O -algebra in $\mathrm{Fun}(I, C)$ with the pointwise monoidal structure is the same data as a functor $I \rightarrow \mathrm{Alg}_O(C)$: there is an equivalence of ∞ -categories*

$$\mathrm{Alg}_O(\mathrm{Fun}(I, C)) \xrightarrow{\sim} \mathrm{Fun}(I, \mathrm{Alg}_O(C)).$$

Proof. The pointwise symmetric monoidal structure of Example 14.2.2 is classified by the composite

$$\mathrm{Span}(\mathrm{Fin}) \xrightarrow{\mathrm{Str}^{\mathrm{cc}}(p_C)} \mathrm{Cat}_{\infty} \xrightarrow{\mathrm{Fun}(I, -)} \mathrm{Cat}_{\infty}.$$

By Proposition 23.2.2, its total category is $\mathrm{Span}(\mathrm{Fin}) \times_{\mathrm{Fun}(I, \mathrm{Span}(\mathrm{Fin}))} \mathrm{Fun}(I, C^{\otimes})$. Consequently, before imposing the finite-product condition, a functor over $\mathrm{Span}(\mathrm{Fin})$ from $O^{\otimes}$ to this total category amounts to a functor

$$O^{\otimes} \longrightarrow \mathrm{Fun}(I, C^{\otimes})$$

whose composite with $\mathrm{Fun}(I, p_C)$ is the constant I -diagram on p_O . Currying identifies these with functors

$$I \longrightarrow \mathrm{Fun}_{/\mathrm{Span}(\mathrm{Fin})}(O^{\otimes}, C^{\otimes}).$$

Under this identification, preservation of finite products is pointwise in I : the functor into the pointwise multimorphism operad is an operad map precisely when the corresponding functor $O^{\otimes} \rightarrow C^{\otimes}$ is an operad map at every object of I . Thus the equivalence restricts to the asserted equivalence of algebra categories. $\square$

14.3 Lax symmetric monoidal functors

The definition of symmetric monoidal ∞ -categories as commutative monoids in Cat_{∞} makes it easy to talk about symmetric monoidal functors. It is a priori less clear how to define *lax* symmetric monoidal functors: functors $F: C \rightarrow D$ that come equipped with *lax structure maps*

$$F(X_1) \otimes \cdots \otimes F(X_n) \rightarrow F(X_1 \otimes \cdots \otimes X_n),$$

for $X_1, \dots, X_n \in C$, that are coherently compatible with the symmetric monoidal structures of C and D . The operadic perspective on symmetric monoidal ∞ -categories provides a quick definition:

Definition 14.3.1. Let C and D be symmetric monoidal ∞ -categories. A *lax symmetric monoidal functor* $C \rightarrow D$ is a morphism of ∞ -operads $\mathcal{M}_C \rightarrow \mathcal{M}_D$. An *oplax symmetric monoidal functor* is a morphism of ∞ -operads $\mathcal{M}_{C^{\mathrm{op}}} \rightarrow \mathcal{M}_D$. We denote by

$$\mathrm{Cat}_{\infty}^{\otimes\text{-lax}} \subseteq \mathrm{Op}_{\infty}$$

the full subcategory spanned by the (multimorphism operads of) symmetric monoidal ∞ -categories.

We write

$$\mathrm{Fun}^{\otimes\text{-lax}}(C, D) := \mathrm{Fun}_{\mathrm{Op}_{\infty}}(\mathcal{M}_C, \mathcal{M}_D)$$

for the ∞ -category of lax symmetric monoidal functors from C to D .

The ∞ -category $\text{Fun}^\otimes(C, D)$ identifies with the full subcategory of $\text{Fun}^{\otimes\text{-lax}}(C, D)$ spanned by the lax symmetric monoidal functors whose structure maps are isomorphisms.

Let us unwind how to recover the lax structure maps. An operad map $\mathcal{M}_C \rightarrow \mathcal{M}_D$ is encoded by a commutative triangle of the form

$$\begin{array}{ccc} C^\otimes & \xrightarrow{F^\otimes} & D^\otimes \\ p_C \searrow & & \swarrow p_D \\ & \text{Span}(\text{Fin}) & \end{array}$$

Passing to fibers over $\langle 1 \rangle$ gives an underlying functor $F: C \rightarrow D$. The fact that $F^\otimes$ preserves finite products means that on objects it is given as $F^\otimes(X_1, \dots, X_n) = (F(X_1), \dots, F(X_n))$, and this also fully specifies $F^\otimes$ on inert maps. To understand what $F^\otimes$ does on active maps, it suffices to consider the case $(X_1, \dots, X_n) \rightarrow X_1 \otimes \dots \otimes X_n$, for which the image in $D^\otimes$ takes the form of a map

$$(F(X_1), \dots, F(X_n)) \rightarrow F(X_1 \otimes \dots \otimes X_n)$$

lifting the span $\langle n \rangle \xleftarrow{=} \langle n \rangle \rightarrow \langle 1 \rangle$. This map uniquely factors through the cocartesian morphism $(F(X_1), \dots, F(X_n)) \rightarrow F(X_1) \otimes \dots \otimes F(X_n)$, resulting in the expected lax structure map.

For oplax symmetric monoidal functors the structure maps point the other way.

Construction 14.3.2. For an ∞ -operad $\mathcal{O}$, the construction $C \mapsto \text{Alg}_{\mathcal{O}}(C)$ defines a functor

$$\text{Alg}_{\mathcal{O}}(-) := \text{Fun}_{\text{Op}_{\infty}}(\mathcal{O}, -): \text{Cat}_{\infty}^{\otimes\text{-lax}} \rightarrow \text{Cat}_{\infty}.$$

Given a lax symmetric monoidal functor $F: C \rightarrow D$, the induced functor $\text{Alg}_{\mathcal{O}}(C) \rightarrow \text{Alg}_{\mathcal{O}}(D)$ is given by postcomposition with $\mathcal{M}_C \rightarrow \mathcal{M}_D$. Its functoriality follows directly from functoriality of postcomposition in the defining fibers of functor categories.

14.3.1 Symmetric monoidal adjunctions

The main goal of this subsection is to show that the right adjoint of a symmetric monoidal functor is canonically lax symmetric monoidal. We will then use the resulting symmetric monoidal adjunction to obtain adjunctions on categories of algebras.

To formulate the compatibility of the lifted adjunction with the maps to $\text{Span}(\text{Fin})$, we first record the small amount of relative adjunction theory that we will need. Given two functors $p: C \rightarrow B$ and $q: D \rightarrow B$, we write

$$\text{Fun}_{/B}(C, D) := \text{Fun}(C, D) \times_{\text{Fun}(C, B)} \{p\},$$

where the map to $\text{Fun}(C, B)$ is given by postcomposition with q . Thus an object of $\text{Fun}_{/B}(C, D)$ consists of a functor $H: C \rightarrow D$ together with a specified identification $qH \simeq p$.

Definition 14.3.3 (Relative adjunction). Let L be an object of $\text{Fun}_{/B}(C, D)$. A *right adjoint to L relative to B* consists of an object $R \in \text{Fun}_{/B}(D, C)$ together with a morphism

$$\varepsilon: LR \rightarrow \text{id}_D$$

in $\text{Fun}_{/B}(D, D)$ whose image in $\text{Fun}(D, D)$ exhibits the underlying functor of R as a right adjoint to the underlying functor of L . We then call $L \dashv R$ an *adjunction relative to B* .

The definition is counit-oriented. Equivalently, we may specify the unit as a morphism over the base:

Lemma 14.3.4 (Unit formulation of relative adjunctions). *Let $L \in \text{Fun}_{/B}(C, D)$ and $R \in \text{Fun}_{/B}(D, C)$. The following data are equivalent:*

- (1) *A morphism $\varepsilon: LR \rightarrow \text{id}_D$ in $\text{Fun}_{/B}(D, D)$ which exhibits R as a right adjoint to L after forgetting to $\text{Fun}(D, D)$.*
- (2) *A morphism $\eta: \text{id}_C \rightarrow RL$ in $\text{Fun}_{/B}(C, C)$ which exhibits L as a left adjoint to R after forgetting to $\text{Fun}(C, C)$.*

Proof. Suppose first that the data in (1) are given. By Proposition 21.1.2, composition with the counit gives an equivalence

$$\text{Nat}(\text{id}_C, RL) \xrightarrow{\simeq} \text{Nat}(L, L), \quad \alpha \mapsto (\varepsilon L)(L\alpha).$$

Under the specified identifications over B , the fact that ε is a morphism in $\text{Fun}_{/B}(D, D)$ gives a commutative square

$$\begin{array}{ccc} \text{Nat}(\text{id}_C, RL) & \xrightarrow{\simeq} & \text{Nat}(L, L) \\ \downarrow & & \downarrow \\ \text{Nat}(p, p) & \xlongequal{\quad} & \text{Nat}(p, p). \end{array}$$

It therefore restricts to an equivalence on the fibers over id_p , which are the hom animae

$$\text{Hom}_{\text{Fun}_{/B}(C, C)}(\text{id}_C, RL) \xrightarrow{\simeq} \text{Hom}_{\text{Fun}_{/B}(C, D)}(L, L).$$

The preimage of id_L gives a morphism $\eta: \text{id}_C \rightarrow RL$ over B , together with an identification $(\varepsilon L)(L\eta) \simeq \text{id}_L$ in $\text{Fun}_{/B}(C, D)$. After forgetting the maps to B , this is the unit corresponding to ε under Proposition 21.1.2, so it gives the data in (2).

Conversely, starting from (2), composition with the unit gives an equivalence

$$\text{Nat}(LR, \text{id}_D) \xrightarrow{\simeq} \text{Nat}(R, R), \quad \beta \mapsto (R\beta)(\eta R).$$

It likewise restricts to the fibers over the identity transformation of q , and the preimage of id_R supplies the counit in (1), together with the other triangle identification over B . These constructions are inverse because they are the usual equivalence between unit and counit data from Proposition 21.1.2, restricted to the indicated fibers. $\square$

Lemma 14.3.5 (Relative adjunctions on functor categories). *Let $L: C \rightleftarrows D: R$ be an adjunction relative to B . For every functor $E \rightarrow B$, postcomposition induces an adjunction*

$$L_*: \text{Fun}_{/B}(E, C) \rightleftarrows \text{Fun}_{/B}(E, D): R_*.$$

Proof. The counit in Definition 14.3.3 and the corresponding unit of Lemma 14.3.4 may both be whiskered with a functor $E \rightarrow C$ or $E \rightarrow D$ over B . They therefore give a unit and counit between the displayed functors, with triangle identifications inherited from those of $L \dashv R$. The result follows from Proposition 21.1.2. $\square$

We now produce the promised lax symmetric monoidal structure on the right adjoint of a symmetric monoidal functor.

Proposition 14.3.6. *Let $L: C \rightarrow D$ be a symmetric monoidal functor between symmetric monoidal ∞ -categories. Assume that L (as a plain functor from C to D) admits a right adjoint $R: D \rightarrow C$. Then the following statements hold:*

- (1) *The functor $L^\otimes: C^\otimes \rightarrow D^\otimes$ admits a right adjoint $R^\otimes: D^\otimes \rightarrow C^\otimes$.*
- (2) *The functor $R^\otimes$ has a canonical structure as a functor over $\text{Span}(\text{Fin})$ for which $L^\otimes \dashv R^\otimes$ is a relative adjunction over $\text{Span}(\text{Fin})$.*
- (3) *With this structure, $R^\otimes$ defines a morphism of ∞ -operads from $\mathcal{M}_D$ to $\mathcal{M}_C$.*
- (4) *The underlying functor of this operad map is $R: D \rightarrow C$.*

In particular, the right adjoint of a symmetric monoidal functor is canonically lax symmetric monoidal.

Proof. (1) By the pointwise criterion for right adjoints from Lemma 21.1.4, it will suffice to show that for every object $Y = \{y_j\}_{j \in J} \in D^\otimes$ there exists an object $R^\otimes Y$ together with a counit map $\varepsilon_Y: L^\otimes R^\otimes Y \rightarrow Y$ such that for every other $X \in C^\otimes$ the composite

$$\text{Hom}_{C^\otimes}(X, R^\otimes Y) \xrightarrow{L^\otimes} \text{Hom}_{D^\otimes}(L^\otimes X, L^\otimes R^\otimes Y) \xrightarrow{\varepsilon_Y \circ -} \text{Hom}_{D^\otimes}(L^\otimes X, Y) \quad (14.1)$$

is an equivalence. We define $R^\otimes Y := \{Ry_j\}_{j \in J}$. We then have $L^\otimes R^\otimes Y \simeq \{LRy_j\}_{j \in J}$ and so the counit map ε_Y may be taken to be the collection of morphisms $\varepsilon_{y_j}: LRy_j \rightarrow y_j$, resulting in a morphism in $D_J^\otimes \simeq \prod_{j \in J} D$.

To show that the composite (14.1) is an equivalence, it suffices to show that it induces an equivalence on the fibers over any span $\alpha: I \xleftarrow{f} K \xrightarrow{g} J$ in $\text{Span}(\text{Fin})$. Let $\tilde{\alpha}: X \rightarrow \alpha_! X$ denote a cocartesian lift of the span α . Since $L^\otimes$ preserves cocartesian morphisms, also the map $L^\otimes(\tilde{\alpha}): L^\otimes X \rightarrow L^\otimes \alpha_! X$ is cocartesian. Precomposition with these maps then induces a commutative diagram

$$\begin{array}{ccccc} \text{Hom}_{C^\otimes}(\alpha_! X, R^\otimes Y) & \xrightarrow{L^\otimes} & \text{Hom}_{D^\otimes}(L^\otimes \alpha_! X, L^\otimes R^\otimes Y) & \xrightarrow{\varepsilon_Y \circ -} & \text{Hom}_{D^\otimes}(L^\otimes \alpha_! X, Y) \\ \downarrow - \circ \tilde{\alpha} & & \downarrow - \circ L^\otimes \tilde{\alpha} & & \downarrow - \circ L^\otimes \tilde{\alpha} \\ \text{Hom}_{C^\otimes}(X, R^\otimes Y) & \xrightarrow{L^\otimes} & \text{Hom}_{D^\otimes}(L^\otimes X, L^\otimes R^\otimes Y) & \xrightarrow{\varepsilon_Y \circ -} & \text{Hom}_{D^\otimes}(L^\otimes X, Y). \end{array}$$

Since the vertical maps induce equivalences on fibers over $\text{Hom}_{\text{Span}(\text{Fin})}(J, J) \xrightarrow{- \circ \alpha} \text{Hom}_{\text{Span}(\text{Fin})}(I, J)$, we have thus reduced to the case of the identity span $\alpha = \text{id}_J$. In this case, the fibers are the hom animaes in the fibers $C_J^\otimes \simeq \prod_{j \in J} C$ and $D_J^\otimes \simeq \prod_{j \in J} D$ respectively. The claim thus follows from the fact that for every $j \in J$ the composite

$$\text{Hom}_C((\alpha_! X)_j, Ry_j) \xrightarrow{L} \text{Hom}_D(L(\alpha_! X)_j, LRy_j) \xrightarrow{\varepsilon_{y_j} \circ -} \text{Hom}_D(L(\alpha_! X)_j, y_j)$$

is an equivalence due to the adjunction $L \dashv R$.

(2) We will now equip $R^\otimes$ with the structure of a functor over $\text{Span}(\text{Fin})$. We claim that it makes the following diagram commute:

$$\begin{array}{ccc} D^\otimes & \xrightarrow{R^\otimes} & C^\otimes \\ & \searrow p_D \quad \swarrow p_C & \\ & \text{Span}(\text{Fin}). & \end{array}$$

To this end, we claim that the adjunction counit $\varepsilon: L^\otimes R^\otimes \rightarrow \text{id}_{D^\otimes}$ induces an equivalence

$$p_C R^\otimes \simeq p_D L^\otimes R^\otimes \xrightarrow{p_D \varepsilon} p_D \text{id}_{D^\otimes} \simeq p_D.$$

This may be checked objectwise for $X \in D^\otimes$, where it is clear from the construction of ε_X . The displayed identification equips $R^\otimes$ with the structure of a functor over $\text{Span}(\text{Fin})$ and lifts ε to a morphism in $\text{Fun}_{/\text{Span}(\text{Fin})}(D^\otimes, D^\otimes)$. Thus $L^\otimes \dashv R^\otimes$ is a relative adjunction over $\text{Span}(\text{Fin})$ by Definition 14.3.3.

(3) The functor $R^\otimes$ preserves finite products because it is a right adjoint. Thus $R^\otimes$ is an operad map.

(4) By uniqueness of adjoints, the restriction of $R^\otimes$ to $D = D_{\langle 1 \rangle}^\otimes$ is R , and we deduce that $R^\otimes$ is a lax symmetric monoidal refinement of R . $\square$

Remark 14.3.7. In the previous proposition, it does not suffice for L to be a lax symmetric monoidal functor itself. It would suffice for L to be an *oplax* symmetric monoidal functor, but producing the lax symmetric monoidal right adjoint is more subtle in this case. Dually, the left adjoint to a lax symmetric monoidal functor is oplax symmetric monoidal. We refer to [HHLN23a, Proposition A] for a detailed discussion.

Definition 14.3.8. A *symmetric monoidal adjunction* is an adjunction

$$L: C \rightleftarrows D : R$$

between symmetric monoidal ∞ -categories in which L is symmetric monoidal and R is equipped with the canonical lax symmetric monoidal structure of Proposition 14.3.6.

By part (2) of Proposition 14.3.6, the induced adjunction $L^\otimes \dashv R^\otimes$ is relative over $\text{Span}(\text{Fin})$. We may deduce from this that symmetric monoidal adjunctions induce adjunctions on algebras over every ∞ -operad.

Lemma 14.3.9 (Adjunctions on $\mathcal{O}$ -algebras). *Every symmetric monoidal adjunction $L \dashv R$ induces, for each ∞ -operad $\mathcal{O}$, an adjunction*

$$\text{Alg}_{\mathcal{O}}(L): \text{Alg}_{\mathcal{O}}(C) \rightleftarrows \text{Alg}_{\mathcal{O}}(D) : \text{Alg}_{\mathcal{O}}(R).$$

The underlying functors are obtained by applying L and R , respectively, at every color of $\mathcal{O}$.

Proof. By part (2) of Proposition 14.3.6, the adjunction $L^\otimes \dashv R^\otimes$ is relative to $\text{Span}(\text{Fin})$. Applying Lemma 14.3.5 to $p_{\mathcal{O}}: \mathcal{O}^\otimes \rightarrow \text{Span}(\text{Fin})$ gives an adjunction

$$\text{Fun}_{/\text{Span}(\text{Fin})}(\mathcal{O}^\otimes, C^\otimes) \rightleftarrows \text{Fun}_{/\text{Span}(\text{Fin})}(\mathcal{O}^\otimes, D^\otimes).$$

Postcomposition with both functors preserves finite-product-preserving functors because $L^\otimes$ and $R^\otimes$ are morphisms of ∞ -operads. Restricting the adjunction to these subcategories gives the asserted adjunction on $\text{Alg}_{\mathcal{O}}(C)$ and $\text{Alg}_{\mathcal{O}}(D)$. $\square$

As an application, it follows that limits of $\mathcal{O}$ -algebras may be computed pointwise.

Proposition 14.3.10 (Limits of $\mathcal{O}$ -algebras). *Let C be a symmetric monoidal ∞ -category, let $\mathcal{O}$ be an ∞ -operad, and let I be a small ∞ -category such that C admits I -indexed limits. Then $\text{Alg}_{\mathcal{O}}(C)$ admits I -indexed limits, and the evaluation functors*

$$\text{ev}_x: \text{Alg}_{\mathcal{O}}(C) \rightarrow C, \quad A \mapsto A_x \quad (x \in \mathcal{O}^{\otimes}),$$

jointly create them.

Proof. By Corollary 21.2.2, the constant-diagram functor fits into an adjunction

$$\text{const}: C \rightleftarrows \text{Fun}(I, C) : \lim_I.$$

Being given by restriction along $I \rightarrow *$, the functor const is symmetric monoidal for the pointwise symmetric monoidal structure, so Proposition 14.3.6 makes this a symmetric monoidal adjunction. By Lemma 14.3.9, it induces an adjunction

$$\text{Alg}_{\mathcal{O}}(C) \rightleftarrows \text{Alg}_{\mathcal{O}}(\text{Fun}(I, C)) \simeq \text{Fun}(I, \text{Alg}_{\mathcal{O}}(C)).$$

Under the equivalence of Lemma 14.2.12, the left adjoint is the constant-diagram functor. The right adjoint consequently computes limits in $\text{Alg}_{\mathcal{O}}(C)$, and its value at every color x is $\lim_I A_x$. Thus all the evaluation functors preserve these limits. They jointly detect isomorphisms: every object of $\mathcal{O}^{\otimes}$ is a finite product of colors, and algebra maps preserve these products, so a natural transformation that is invertible at every color is invertible at every object. Consequently the evaluation functors jointly reflect limit cones and hence jointly create the limits. $\square$

14.3.2 Symmetric monoidal left adjoints

As discussed in Remark 14.3.7, it is possible to show that the left adjoint of a lax symmetric monoidal functor admits a canonical oplax symmetric monoidal structure. While this is subtle to construct in general, the construction substantially simplifies when this left adjoint happens to be actually symmetric monoidal.

Proposition 14.3.11. *Let $R: D \rightarrow C$ be a lax symmetric monoidal functor between symmetric monoidal ∞ -categories. Assume that R (as a plain functor from D to C) admits a left adjoint $L: C \rightarrow D$ such that for all $n \geq 0$ and objects $x_1, \dots, x_n \in C$ the composite*

$$L\left(\bigotimes_{i=1}^n x_i\right) \xrightarrow{L(\bigotimes_{i=1}^n \eta_{x_i})} L\left(\bigotimes_{i=1}^n RLx_i\right) \xrightarrow{L(\text{lax}_R)} LR\left(\bigotimes_{i=1}^n Lx_i\right) \xrightarrow{\varepsilon} \bigotimes_{i=1}^n L(x_i)$$

is an isomorphism in D . Then the following statements hold:

- (1) *The functor $R^{\otimes}: D^{\otimes} \rightarrow C^{\otimes}$ admits a left adjoint $L^{\otimes}: C^{\otimes} \rightarrow D^{\otimes}$.*
- (2) *The functor $L^{\otimes}$ has a canonical structure as a functor over $\text{Span}(\text{Fin})$ for which $L^{\otimes} \dashv R^{\otimes}$ is a relative adjunction over $\text{Span}(\text{Fin})$.*
- (3) *With this structure, $L^{\otimes}$ defines a morphism of ∞ -operads from $\mathcal{M}_C$ to $\mathcal{M}_D$.*

- (4) This morphism of ∞ -operads comes from a symmetric monoidal functor from C to D refining L .

In particular, R admits a strong symmetric monoidal left adjoint $L: (C, \otimes_C) \rightarrow (D, \otimes_D)$.

Note that by induction it suffices to check the condition on L for $n = 0$ and $n = 2$, i.e. it suffices to show that the two canonical maps

$$L(\mathbb{1}_C) \rightarrow \mathbb{1}_D \quad \text{and} \quad L(x \otimes y) \rightarrow L(x) \otimes L(y)$$

are isomorphisms.

Proof. (1) As in the proof of Proposition 14.3.6, we use the pointwise criterion for left adjoints from Lemma 21.1.4. We define $L^\otimes$ objectwise as $L^\otimes(\{x_i\}_{i \in I}) := \{L(x_i)\}_{i \in I}$; the unit maps $\eta_{x_i}: x_i \rightarrow RLx_i$ induce a candidate unit map $\eta_X: X \rightarrow R^\otimes L^\otimes X$. We must show that for every object $Y \in D_J^\otimes$ the composite

$$\mathrm{Hom}_{D^\otimes}(L^\otimes X, Y) \xrightarrow{R^\otimes} \mathrm{Hom}_{C^\otimes}(R^\otimes L^\otimes X, R^\otimes Y) \xrightarrow{- \circ \eta_X} \mathrm{Hom}_{C^\otimes}(X, R^\otimes Y)$$

is an equivalence. This may be tested fiberwise over any span $\alpha: I \xleftarrow{f} K \xrightarrow{g} J$. Let

$$X \rightarrow \alpha_! X, \quad L^\otimes X \rightarrow \alpha_! L^\otimes X, \quad \text{and} \quad R^\otimes L^\otimes X \rightarrow \alpha_! R^\otimes L^\otimes X$$

denote cocartesian lifts of α in $C^\otimes$ and $D^\otimes$, and let $\mathrm{lax}_R: \alpha_! R^\otimes L^\otimes X \rightarrow R^\otimes \alpha_! L^\otimes X$ denote the lax structure map of R associated to α . We then get a commutative diagram as follows:

$$\begin{array}{ccc} \mathrm{Hom}_{D^\otimes}^{\mathrm{id}_J}(\alpha_! L^\otimes X, Y) & \xrightarrow{\simeq} & \mathrm{Hom}_{D^\otimes}^\alpha(L^\otimes X, Y) \\ \downarrow R^\otimes & & \downarrow R^\otimes \\ \mathrm{Hom}_{C^\otimes}^{\mathrm{id}_J}(R^\otimes \alpha_! L^\otimes X, R^\otimes Y) & & \\ \downarrow - \circ \mathrm{lax}_R & & \downarrow \\ \mathrm{Hom}_{C^\otimes}^{\mathrm{id}_J}(\alpha_! R^\otimes L^\otimes X, R^\otimes Y) & \xrightarrow{\simeq} & \mathrm{Hom}_{C^\otimes}^\alpha(R^\otimes L^\otimes X, R^\otimes Y) \\ \downarrow - \circ \alpha_! \eta_X & & \downarrow - \circ \eta_X \\ \mathrm{Hom}_{C^\otimes}^{\mathrm{id}_J}(\alpha_! X, R^\otimes Y) & \xrightarrow{\simeq} & \mathrm{Hom}_{C^\otimes}^\alpha(X, R^\otimes Y). \end{array}$$

It will thus suffice to show that the left vertical composite is an equivalence. Under the identifications $C_J^\otimes \simeq \prod_{j \in J} C$ and $D_J^\otimes \simeq \prod_{j \in J} D$, this composite is the product over $j \in J$ of the analogous composites associated to the restricted spans

$$\alpha_j: I \xleftarrow{f|_{g^{-1}(j)}} g^{-1}(j) \longrightarrow \langle 1 \rangle.$$

Indeed, $R^\otimes$ is a morphism of ∞ -operads and therefore respects these product decompositions; in particular, the j -th component of lax_R is the lax structure map associated to α_j . The map $\alpha_! \eta_X$ decomposes in the same way. It therefore suffices to treat the case $J = \langle 1 \rangle$. Composing the vertical composite with the adjunction equivalence $\mathrm{Hom}_C(\alpha_! X, RY) \simeq \mathrm{Hom}_D(L\alpha_! X, Y)$, we obtain a map

$$\mathrm{Hom}_D(\alpha_! L^\otimes X, Y) \rightarrow \mathrm{Hom}_D(L\alpha_! X, Y)$$

which we must show is an equivalence. But unwinding definitions shows that this map is given by precomposition with the composite

$$L\alpha_!X \xrightarrow{L\alpha_!\eta_X} L\alpha_!R^\otimes L^\otimes X \xrightarrow{L(\text{lax}_R)} LR\alpha_!L^\otimes X \xrightarrow{\varepsilon_{\alpha_!L^\otimes X}} \alpha_!L^\otimes X.$$

This is precisely the comparison map from the hypothesis for the family $\{x_{f(k)}\}_{k \in K}$, and is therefore an isomorphism. This proves the existence of the left adjoint $L^\otimes$.

(2) The unit maps $\eta_X: X \rightarrow R^\otimes L^\otimes X$ used in the construction lie in the same fibers of $C^\otimes \rightarrow \text{Span}(\text{Fin})$ as their sources. Their images in $\text{Span}(\text{Fin})$ therefore give a natural identification $p_D L^\otimes \simeq p_C$ which equips $L^\otimes$ with the structure of a functor over $\text{Span}(\text{Fin})$ and lifts the unit to a morphism in $\text{Fun}/\text{Span}(\text{Fin})(C^\otimes, C^\otimes)$. By Lemma 14.3.4, this makes $L^\otimes \dashv R^\otimes$ a relative adjunction over $\text{Span}(\text{Fin})$.

(3) The objectwise formula on tuples shows that $L^\otimes$ preserves finite products. Thus it is a morphism of ∞ -operads. Finally, the comparison isomorphism above identifies $L^\otimes \alpha_!X$ with $\alpha_!L^\otimes X$ for every span α . Hence $L^\otimes$ preserves cocartesian morphisms and is strong symmetric monoidal.

(4) By its objectwise definition, the restriction of $L^\otimes$ to the fiber over $\langle 1 \rangle$ is L . $\square$

14.4 $\mathcal{O}$ -monoidal ∞ -categories

In the previous sections, we have discussed various results for symmetric monoidal ∞ -categories, i.e., commutative monoids in Cat_∞ . All of this can directly be generalized to monoids over an arbitrary ∞ -operad $\mathcal{O}$:

Definition 14.4.1. Let D be an ∞ -category with finite products. An $\mathcal{O}$ -monoid in D is a finite-product-preserving functor $\mathcal{O}^\otimes \rightarrow D$. We write

$$\text{Mon}_{\mathcal{O}}(D) := \text{Fun}^\times(\mathcal{O}^\otimes, D) \subseteq \text{Fun}(\mathcal{O}^\otimes, D)$$

for the full subcategory of $\mathcal{O}$ -monoids in D .

When D is equipped with its cartesian monoidal structure, $\mathcal{O}$ -monoids in D will be identified with $\mathcal{O}$ -algebras in Theorem 15.3.11.

Definition 14.4.2. An $\mathcal{O}$ -monoidal ∞ -category is an $\mathcal{O}$ -monoid in Cat_∞ . When $\mathcal{O} = \mathcal{A}ssoc$, we refer to $\mathcal{O}$ -monoidal ∞ -categories simply as *monoidal ∞ -categories*.

Example 14.4.3. When $\mathcal{O} = \mathcal{T}riv$, an $\mathcal{O}$ -monoidal ∞ -category is just an ∞ -category.

The multimorphism operad functor $\mathcal{M}: \text{Cat}_\infty^\otimes \rightarrow \text{Op}_\infty$ admits an analogue $\mathcal{M}_{-/ \mathcal{O}}: \text{Mon}_{\mathcal{O}}(\text{Cat}_\infty) \rightarrow (\text{Op}_\infty)_{/\mathcal{O}}$ for $\mathcal{O}$ -monoidal ∞ -categories. This relies on the following relative analogue of Lemma 14.2.4.

Lemma 14.4.4. Let $\mathcal{O}$ be an ∞ -operad and let $q: C^\otimes \rightarrow \mathcal{O}^\otimes$ be a cocartesian fibration. Then the composite

$$C^\otimes \xrightarrow{q} \mathcal{O}^\otimes \xrightarrow{p_{\mathcal{O}}} \text{Span}(\text{Fin})$$

exhibits $C^\otimes$ as an ∞ -operad if and only if the functor $\text{Str}^{\text{cc}}(q): \mathcal{O}^\otimes \rightarrow \text{Cat}_\infty$ preserves finite products. In this case, the morphism $q: C^\otimes \rightarrow \mathcal{O}^\otimes$ is automatically a morphism of ∞ -operads.

Proof. Write $F := \text{Str}^{\text{cc}}(q)$ and $p := p_O$. Condition (i) of Proposition 14.1.7 always holds for pq : given a backwards span in $\text{Span}(\text{Fin})$ and an object $X \in C^\otimes$, first choose a p -cocartesian lift starting at $q(X)$ and then a q -cocartesian lift starting at X . The resulting morphism in $C^\otimes$ is (pq) -cocartesian, as follows by pasting the two pullback squares that express the respective cocartesian universal properties.

For every finite set I , cocartesian transport along the maps ρ_i gives a commutative square

$$\begin{array}{ccc} C_I^\otimes & \longrightarrow & \prod_{i \in I} C_{\{i\}}^\otimes \\ q \downarrow & & \downarrow \prod_{i \in I} q \\ O_I^\otimes & \longrightarrow & \prod_{i \in I} O_{\{i\}}^\otimes. \end{array}$$

The top horizontal functor is the one appearing in condition (ii), since a q -cocartesian lift lying over a p -cocartesian lift is (pq) -cocartesian. The bottom horizontal functor is an equivalence because O is an ∞ -operad. Consequently the top horizontal functor is an equivalence if and only if the square is a pullback. This is a square of cocartesian fibrations, so the pullback condition may be checked on fibers. Indeed, over an object $o \in O_I^\otimes$ with components o_i , the i -th component sends $X \in F(o)$ to the target of a q -cocartesian lift over $o \rightarrow o_i$, which under straightening is precisely $F(o \rightarrow o_i)(X)$. Thus the induced functor on fibers is the canonical comparison

$$F(o) \longrightarrow \prod_{i \in I} F(o_i).$$

The maps $o \rightarrow o_i$ are the product projections in $O^\otimes$. It follows that these fiberwise comparisons are equivalences whenever F preserves finite products. Conversely, if condition (ii) holds, the square is a pullback, so all the component comparisons are equivalences. Every object of $O^\otimes$ is the product of its components, and the components of a product $\prod_a o_a$ are obtained by concatenating those of the objects o_a . The product comparison for the family $(o_a)_a$ becomes an equivalence after composing with the product of their component comparisons, so it is itself an equivalence. Thus condition (ii) for pq holds if and only if F preserves finite products.

Under these equivalent conditions, condition (iii) is automatic. Indeed, for $X \in C_I^\otimes$ over o , let $X \rightarrow X_i$ be q -cocartesian lifts of the product projections $o \rightarrow o_i$. By Lemma 23.2.8, these maps exhibit X as the product of the X_i in $C^\otimes$. They are also (pq) -cocartesian over the maps ρ_i , as observed above. The characterization of ∞ -operads now shows that pq exhibits $C^\otimes$ as an ∞ -operad if and only if F preserves finite products. Finally, Lemma 23.2.8 also shows that q preserves finite products, so it is a morphism of ∞ -operads. $\square$

Definition 14.4.5. Given an O -monoidal ∞ -category C , we let $p_C: C^\otimes \rightarrow O^\otimes$ denote its cocartesian unstraightening, which by the lemma defines an ∞ -operad $\mathcal{M}_{C/O} = (C^\otimes, p_O \circ p_C)$. Since p_C is a morphism of ∞ -operads, it turns $\mathcal{M}_{C/O}$ into an object of the slice category $(\text{Op}_\infty)_{/O}$, producing a functor

$$\mathcal{M}_{-/O}: \text{Mon}_O(\text{Cat}_\infty) \rightarrow (\text{Op}_\infty)_{/O}.$$

If $\mathcal{P} \rightarrow \mathcal{O}$ and $\mathcal{Q} \rightarrow \mathcal{O}$ are two ∞ -operads over $\mathcal{O}$, we denote by $\mathrm{Fun}_{(\mathrm{Op}_\infty)/\mathcal{O}}(\mathcal{P}, \mathcal{Q})$ the ∞ -category of operad maps $\mathcal{P} \rightarrow \mathcal{Q}$ over $\mathcal{O}$, defined as the following fiber:

$$\begin{array}{ccc} \mathrm{Fun}_{(\mathrm{Op}_\infty)/\mathcal{O}}(\mathcal{P}, \mathcal{Q}) & \longrightarrow & \mathrm{Fun}_{\mathrm{Op}_\infty}(\mathcal{P}, \mathcal{Q}) \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \mathrm{Fun}_{\mathrm{Op}_\infty}(\mathcal{P}, \mathcal{O}). \end{array}$$

As a special case, we define the ∞ -category $\mathcal{P}/\mathcal{O}$ -algebras in $\mathcal{C}$ as

$$\mathrm{Alg}_{\mathcal{P}/\mathcal{O}}(\mathcal{C}) := \mathrm{Fun}_{(\mathrm{Op}_\infty)/\mathcal{O}}(\mathcal{P}, \mathcal{M}_{\mathcal{C}/\mathcal{O}}).$$

When $\mathcal{O} = \mathcal{A}ssoc$, we usually drop $\mathcal{O}$ from the notation and abusively denote the $\mathcal{P}/\mathcal{A}ssoc$ -algebras in $\mathcal{C}$ by $\mathrm{Alg}_{\mathcal{P}}(\mathcal{C})$. When $\mathcal{P} = \mathcal{A}ssoc$, we also write this as $\mathrm{Alg}(\mathcal{C})$.

If $\mathcal{C}$ is symmetric monoidal and $\mathcal{P} \rightarrow \mathcal{C}omm$ is an ∞ -operad over $\mathcal{C}omm$, then this relative definition agrees with the earlier one:

$$\mathrm{Alg}_{\mathcal{P}/\mathcal{C}omm}(\mathcal{C}) \simeq \mathrm{Alg}_{\mathcal{P}}(\mathcal{C}).$$

Indeed, $\mathcal{C}omm$ is terminal in Op_∞ , so passing to the slice over $\mathcal{C}omm$ imposes no additional condition on operad maps.

Definition 14.4.6. Given $\mathcal{O}$ -monoidal ∞ -categories $\mathcal{C}$ and $\mathcal{D}$, a lax $\mathcal{O}$ -monoidal functor $\mathcal{C} \rightarrow \mathcal{D}$ is an operad morphism $\mathcal{M}_{\mathcal{C}/\mathcal{O}} \rightarrow \mathcal{M}_{\mathcal{D}/\mathcal{O}}$ over $\mathcal{O}^\otimes$.

Proposition 14.4.7 (cf. [Lur17, Corollary 7.3.2.7]). *Let $L: \mathcal{C} \rightarrow \mathcal{D}$ be an $\mathcal{O}$ -monoidal functor between $\mathcal{O}$ -monoidal ∞ -categories. Assume that for every color $o \in \mathcal{O}^\otimes$, the functor $L_o: \mathcal{C}_o \rightarrow \mathcal{D}_o$ admits a right adjoint $R_o: \mathcal{D}_o \rightarrow \mathcal{C}_o$. Then the functor $L^\otimes: \mathcal{C}^\otimes \rightarrow \mathcal{D}^\otimes$ admits a right adjoint $R^\otimes: \mathcal{D}^\otimes \rightarrow \mathcal{C}^\otimes$. The functor $R^\otimes$ has a canonical structure over $\mathcal{O}^\otimes$ for which $L^\otimes \dashv R^\otimes$ is a relative adjunction, and with this structure it is a morphism of operads over $\mathcal{O}$. In particular, L admits a lax $\mathcal{O}$ -monoidal right adjoint R .*

Proof. The proof is entirely analogous to that of Proposition 14.3.6 and is left to the reader. $\square$

Proposition 14.4.8 (cf. [Lur17, Corollary 7.3.2.12]). *Let $R: \mathcal{D} \rightarrow \mathcal{C}$ be an $\mathcal{O}$ -monoidal functor between $\mathcal{O}$ -monoidal ∞ -categories. Assume that the following conditions hold:*

- (1) *For every color $x \in \mathcal{O}^\otimes$, the functor $R_x: \mathcal{D}_x \rightarrow \mathcal{C}_x$ admits a left adjoint $L_x: \mathcal{C}_x \rightarrow \mathcal{D}_x$.*
- (2) *For every multimorphism $\alpha: X = (x_1, \dots, x_n) \rightarrow y$ in $\mathcal{O}^\otimes$, the composite transformation*

$$L_y \alpha!(-) \xrightarrow{\eta} L_y \alpha!(R_X L_X(-)) \xrightarrow{L_y(\mathrm{lax}_R^\alpha)} L_y R_y \alpha! L_X(-) \xrightarrow{\varepsilon} \alpha! L_X(-)$$

of functors $\mathcal{C}_X^\otimes \rightarrow \mathcal{D}_y^\otimes$ is an isomorphism.

Then the functor $R^\otimes: \mathcal{D}^\otimes \rightarrow \mathcal{C}^\otimes$ admits a left adjoint $L^\otimes: \mathcal{C}^\otimes \rightarrow \mathcal{D}^\otimes$. The functor $L^\otimes$ has a canonical structure over $\mathcal{O}^\otimes$ for which $L^\otimes \dashv R^\otimes$ is a relative adjunction, and with this structure it is a morphism of cocartesian fibrations over $\mathcal{O}^\otimes$. In particular, R admits a strong $\mathcal{O}$ -monoidal left adjoint L .

Proof. The proof is entirely analogous to that of Proposition 14.3.11 and is left to the reader. $\square$

The limit result for algebras also admits the following relative form.

Proposition 14.4.9 (cf. [Lur17, Corollary 3.2.2.5]). *Let $q: \mathcal{P} \rightarrow \mathcal{O}$ be a morphism of ∞ -operads, let C be an $\mathcal{O}$ -monoidal ∞ -category, and let I be a small ∞ -category. Assume that each fiber C_x , for $x \in \mathcal{O}^\simeq$, admits I -indexed limits. Then $\mathrm{Alg}_{\mathcal{P}/\mathcal{O}}(C)$ admits I -indexed limits, and the evaluation functors*

$$\mathrm{ev}_y: \mathrm{Alg}_{\mathcal{P}/\mathcal{O}}(C) \rightarrow C_{q(y)}, \quad A \mapsto A_y \quad (y \in \mathcal{P}^\simeq),$$

jointly create them.

Proof. The proof is analogous to that of Proposition 14.3.10. Applying $\mathrm{Fun}(I, -)$ fiberwise gives an $\mathcal{O}$ -monoidal category whose fiber at x is $\mathrm{Fun}(I, C_x)$, and algebras in it identify with I -diagrams in $\mathrm{Alg}_{\mathcal{P}/\mathcal{O}}(C)$. The fiberwise constant-diagram functors form an $\mathcal{O}$ -monoidal functor, with lax $\mathcal{O}$ -monoidal right adjoint given in each fiber by $\lim_I$. By Proposition 14.4.7, these functors form a relative adjunction over $\mathcal{O}^\otimes$, so Lemma 14.3.5 gives the required right adjoint to the constant-diagram functor after applying $\mathrm{Fun}_{/\mathcal{O}^\otimes}(\mathcal{P}^\otimes, -)$ and restricting to operad maps. Its evaluation at y is $\lim_I$ in $C_{q(y)}$. The evaluation functors jointly detect isomorphisms, so they jointly create the limits. $\square$

Corollary 14.4.10. *Let C be a monoidal ∞ -category and let I be a small ∞ -category such that C admits I -indexed limits. Then $\mathrm{Alg}(C)$ admits I -indexed limits, and the forgetful functor $\mathrm{Alg}(C) \rightarrow C$ creates and preserves them. The same holds for the forgetful functor $\mathrm{CAlg}(C) \rightarrow C$ when C is symmetric monoidal.*

Proof. This is the one-color case of Proposition 14.4.9: a monoidal ∞ -category is an $\mathcal{A}\mathrm{ssoc}$ -monoidal ∞ -category, with $\mathrm{Alg}(C) = \mathrm{Alg}_{\mathcal{A}\mathrm{ssoc}/\mathcal{A}\mathrm{ssoc}}(C)$, and a symmetric monoidal ∞ -category is a Comm -monoidal ∞ -category, with $\mathrm{CAlg}(C) = \mathrm{Alg}_{\mathrm{Comm}/\mathrm{Comm}}(C)$. $\square$

Corollary 14.4.11. *Let C be a monoidal ∞ -category with a terminal object T . Then T carries a unique associative algebra structure, and it is terminal in $\mathrm{Alg}(C)$. When C is symmetric monoidal, the same holds for the commutative algebra structure and $\mathrm{CAlg}(C)$.*

Proof. Apply Corollary 14.4.10 to the empty diagram. It produces a terminal algebra A whose underlying object is T . If B is any other algebra with underlying object T , the unique map $B \rightarrow A$ has an isomorphism as its underlying morphism, and is therefore itself an isomorphism. Moreover, both $\mathrm{Hom}_{\mathrm{Alg}(C)}(B, A)$ and $\mathrm{Hom}_C(T, T)$ are contractible, so the anima of such maps lying over id_T is contractible. It follows that the fiber of algebra structures on T is contractible. The commutative case is identical. $\square$

14.5 Monoidal subcategories and monoidal Bousfield localizations

In this section we will introduce conditions for subcategories and Bousfield localizations of symmetric monoidal ∞ -categories to again be symmetric monoidal.

Construction 14.5.1. Let $\mathcal{O}$ be an ∞ -operad. Given a subanima $\mathcal{P}^\simeq \subseteq \mathcal{O}^\simeq$, let $\mathcal{P}^\otimes \subseteq \mathcal{O}^\otimes$ be the full subcategory spanned by objects $\{x_i\}_{i \in I}$ with $x_i \in \mathcal{P}^\simeq$ for all i . Observe that the composite

$p_{\mathcal{P}}: \mathcal{P}^{\otimes} \hookrightarrow \mathcal{O}^{\otimes} \xrightarrow{p_{\mathcal{O}}} \text{Span}(\text{Fin})$ exhibits $\mathcal{P}^{\otimes}$ as an ∞ -operad and the inclusion $\mathcal{P}^{\otimes} \hookrightarrow \mathcal{O}^{\otimes}$ is a morphism of ∞ -operads:

- (1) $\mathcal{P}^{\otimes}$ admits products inherited from $\mathcal{O}^{\otimes}$;
- (2) By definition the equivalence $\prod_{i \in I} \mathcal{O}_{\{i\}}^{\otimes} \xrightarrow{\sim} \mathcal{O}_I^{\otimes}$ restricts to $\prod_{i \in I} \mathcal{P}_{\{i\}}^{\otimes} \xrightarrow{\sim} \mathcal{P}_I^{\otimes}$;
- (3) By full faithfulness, $p_{\mathcal{O}}$ -cocartesian morphisms $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ in $\mathcal{O}^{\otimes}$ are still $p_{\mathcal{P}}$ -cocartesian as morphisms in $\mathcal{P}^{\otimes}$.

We refer to the pair $\mathcal{P} = (\mathcal{P}^{\otimes}, p_{\mathcal{P}})$ as the *full suboperad of $\mathcal{O}$ spanned by the colors $\mathcal{P}^{\otimes}$* .

If $\mathcal{O} = \mathcal{M}_C$ comes from a symmetric monoidal category C , then the choice of a subanima of colors corresponds to a choice of a full subcategory $D \subseteq C$. We will now discuss various conditions that imply that D inherits a symmetric monoidal structure from C , either by restricting the monoidal structure or by localizing it.

Lemma 14.5.2. *Let $(C, \otimes_C, \mathbb{1}_C)$ be a symmetric monoidal ∞ -category. If $D \subseteq C$ is a full subcategory whose collection of objects $D^{\simeq} \subseteq C^{\simeq}$ contains $\mathbb{1}_C$ and is closed under $\otimes_C$, then D inherits a symmetric monoidal structure $(D, \otimes_D, \mathbb{1}_D)$ from C such that the inclusion $i: D \hookrightarrow C$ is a (strong) symmetric monoidal functor.*

Proof. The previous construction provides a suboperad $D^{\otimes} \subseteq C^{\otimes}$ and it remains to show that the functor $D^{\otimes} \hookrightarrow C^{\otimes} \xrightarrow{p} \text{Span}(\text{Fin})$ is still a cocartesian fibration. By full faithfulness of the inclusion, it will suffice to show that the p -cocartesian morphisms in $C^{\otimes}$ starting in an object of $D^{\otimes}$ have their targets in $D^{\otimes}$. Using the inert-active factorization and Observation 14.2.7, we may reduce this to cocartesian morphisms of the form

$$(x_1, \dots, x_n) \longrightarrow x_1 \otimes \cdots \otimes x_n.$$

The claim is clear for $n = 1$. For $n = 0$ it is the condition that $\mathbb{1}_C \in D$, and for $n = 2$ it is the condition that $x \otimes y \in D$ whenever $x, y \in D$. The case $n \geq 3$ reduces to $n = 2$ by induction, using that composites of cocartesian morphisms are again cocartesian. $\square$

Recall that a *Bousfield localization* is a functor $L: C \rightarrow D$ that admits a fully faithful right adjoint $i: D \hookrightarrow C$. In this case a morphism $x \rightarrow y$ is called an *L-equivalence* if the map $L(x) \rightarrow L(y)$ in D is an isomorphism.

Proposition 14.5.3. *Let $(C, \otimes_C, \mathbb{1}_C)$ be a symmetric monoidal ∞ -category, and let $i: D \hookrightarrow C$ be the inclusion of a full subcategory such that i admits a left adjoint $L: C \rightarrow D$. Assume that for every L -equivalence $f: x \rightarrow y$ in C and for every $z \in C$, the morphism $f \otimes_C \text{id}_z: x \otimes z \rightarrow y \otimes z$ is also an L -equivalence. Then the following hold:*

- (1) *The suboperad $D^{\otimes}$ of $C^{\otimes}$ is a symmetric monoidal structure on D ;*
- (2) *The inclusion $i^{\otimes}: D^{\otimes} \hookrightarrow C^{\otimes}$ admits a left adjoint $L^{\otimes}: C^{\otimes} \rightarrow D^{\otimes}$ which is a strong symmetric monoidal refinement of the functor L ;*

(3) If C admits small colimits and its tensor product preserves them separately in both variables, then the same holds for D .

Proof. Step 1: Let us start by observing that the assumption on L implies that for every span $\alpha: I \xleftarrow{f} K \xrightarrow{g} J$ of finite sets the induced functor

$$\alpha_!: C^I \xrightarrow{f^*} C^K \xrightarrow{g_\otimes} C^J, \quad (x_i)_{i \in I} \mapsto \left(\bigotimes_{k \in g^{-1}(j)} x_{f(k)} \right)_{j \in J}$$

preserves componentwise L -equivalences. This is clear for the restriction functor f^* . For the functor $g_\otimes$, it remains to show that if $x_k \rightarrow y_k$ are L -equivalences in C for $k = 1, \dots, n$, then their tensor product $x_1 \otimes_C \dots \otimes_C x_n \rightarrow y_1 \otimes_C \dots \otimes_C y_n$ is also an L -equivalence. For $n = 2$ this follows by writing this map as a composite

$$x_1 \otimes x_2 \rightarrow y_1 \otimes x_2 \simeq x_2 \otimes y_1 \rightarrow y_2 \otimes y_1 \simeq y_1 \otimes y_2,$$

where the two non-invertible arrows are L -equivalences by assumption, and the remaining arrows are the braiding equivalences. The general case follows by induction.

Step 2: We will now show that $i^\otimes$ has a left adjoint $L^\otimes$. To simplify notation, we will treat objects of D as objects in C without writing the inclusion functor i . By the pointwise criterion for adjunctions from Lemma 21.1.4, we may construct $L^\otimes$ objectwise, and we set $L^\otimes(\{x_k\}_{k \in K}) := \{L(x_k)\}_{k \in K}$. The unit map $\eta_X: X \rightarrow L^\otimes X$ is given by the collection of maps $\{\eta_{x_k}: x_k \rightarrow L(x_k)\}_{k \in K}$. We must show that for any other $Y \in D^\otimes$ the precomposition map

$$\mathrm{Hom}_{D^\otimes}(L^\otimes X, Y) \xrightarrow{- \circ \eta_X} \mathrm{Hom}_{C^\otimes}(X, Y)$$

is an equivalence. It suffices to check this on fibers $\mathrm{Hom}_{C^\otimes}^\alpha(-, -)$ over any span $\alpha: K \xleftarrow{f} S \xrightarrow{g} J$. As in the proof of Proposition 14.3.11, we will reduce this to the case of the identity span where it will follow from the adjunction $L \dashv i$. Let $X \rightarrow \alpha_! X$ and $L^\otimes X \rightarrow \alpha_! L^\otimes X$ denote p_C -cocartesian lifts of α starting in X and $L^\otimes X$, respectively, so that precomposition with these maps induces two vertical equivalences as follows:

$$\begin{array}{ccc} \mathrm{Hom}_{D^\otimes}^{\mathrm{id}_J}(\alpha_! L^\otimes X, Y) & \xrightarrow{- \circ \alpha_!(\eta_X)} & \mathrm{Hom}_{C^\otimes}^{\mathrm{id}_J}(\alpha_! X, Y) \\ \simeq \downarrow & & \downarrow \simeq \\ \mathrm{Hom}_{D^\otimes}^\alpha(L^\otimes X, Y) & \xrightarrow{- \circ \eta_X} & \mathrm{Hom}_{C^\otimes}^\alpha(X, Y). \end{array}$$

It thus suffices to show that the top map is an equivalence. Using the identification $C_J^\otimes \simeq C^J$, we may identify this map with the product over $j \in J$ of the maps

$$\mathrm{Hom}_C((\alpha_! L^\otimes X)_j, Y_j) \xrightarrow{- \circ \alpha_!(\eta_X)_j} \mathrm{Hom}_C((\alpha_! X)_j, Y_j),$$

so we may show that each of these is an equivalence. Applying the adjunction $L \dashv i$, this map is in turn equivalent to the map

$$\mathrm{Hom}_D(L(\alpha_! L^\otimes X)_j, Y_j) \xrightarrow{- \circ L\alpha_!(\eta_X)_j} \mathrm{Hom}_D(L(\alpha_! X)_j, Y_j).$$

But now note that the map $\eta_{x_k} : x_k \rightarrow L(x_k)$ is an L -equivalence for each k due to full faithfulness of $i : D \hookrightarrow C$, and so by step 1 the maps $\alpha_!(\eta_X)_j : \alpha_!(X)_j \rightarrow \alpha_!(L^\otimes X)_j$ are also L -equivalences for all j , i.e., each $L\alpha_!(\eta_X)_j$ is an isomorphism. This shows the existence of the left adjoint $L^\otimes$. The same argument as in Proposition 14.3.11 shows that $L^\otimes$ is a morphism of ∞ -operads.

Step 3: We will now show that the suboperad $D^\otimes \subseteq C^\otimes$ is a symmetric monoidal ∞ -category. Given an object $X \in D^\otimes$ and a span α , we need to show that there exists a p_D -cocartesian lift of α in $D^\otimes$ starting in X . We construct this as the composite

$$X \rightarrow \alpha_! X \xrightarrow{\eta} L^\otimes \alpha_! X,$$

where the first map is a p_C -cocartesian lift of α in $C^\otimes$, and the second map is the unit for the adjunction $L^\otimes \dashv \alpha_!$. To show that this map is p_D -cocartesian, we have to show that for every other object $Y \in D^\otimes$ the following commutative square is a pullback square:

$$\begin{array}{ccc} \mathrm{Hom}_{D^\otimes}(L^\otimes \alpha_! X, Y) & \longrightarrow & \mathrm{Hom}_{D^\otimes}(X, Y) \\ \downarrow & & \downarrow p \\ \mathrm{Hom}_{\mathrm{Span}(\mathrm{Fin})}(J, K) & \xrightarrow{-\circ \alpha} & \mathrm{Hom}_{\mathrm{Span}(\mathrm{Fin})}(I, K). \end{array}$$

But by the adjunction equivalence $\mathrm{Hom}_{D^\otimes}(L^\otimes \alpha_! X, Y) \simeq \mathrm{Hom}_{C^\otimes}(\alpha_! X, Y)$ this immediately follows from p_C -cocartesianness of the map $X \rightarrow \alpha_! X$. Thus p_D is a cocartesian fibration. Since $D^\otimes$ is already an ∞ -operad, Lemma 14.2.4 shows that it encodes a symmetric monoidal ∞ -category. This proves part (1).

Step 4: For (2), it remains to show that $L^\otimes$ is strong symmetric monoidal, i.e., that it sends p_C -cocartesian morphisms to p_D -cocartesian morphisms. Given a p_C -cocartesian morphism $X \rightarrow \alpha_! X$, applying $L^\otimes$ gives $L^\otimes X \rightarrow L^\otimes \alpha_! X$, and we need to show that this agrees with the composite $L^\otimes X \rightarrow \alpha_! L^\otimes X \xrightarrow{\eta} L^\otimes \alpha_! L^\otimes X$. But this again follows from the assumption on L : since $\eta_X : X \rightarrow L^\otimes X$ is an L -equivalence, so is the map $\alpha_! X \rightarrow \alpha_! L^\otimes X$, hence applying $L^\otimes$ gives an isomorphism.

Step 5: Assume that C admits small colimits and that its tensor product preserves them separately. Then D admits small colimits, with

$$\mathrm{colim}_{j \in J} X_j \simeq L(\mathrm{colim}_{j \in J} iX_j)$$

for every small diagram $(X_j)_{j \in J}$ in D . Together with the formula

$$X \otimes_D Y \simeq L(iX \otimes_C iY),$$

the assumptions on the tensor product and on L now imply part (3) by an easy computation. $\square$

A symmetric monoidal localization descends to algebras.

Corollary 14.5.4 (cf. [Lur17, Proposition 2.2.1.9]). *Let $L : C \rightleftarrows D : i$ be as in Proposition 14.5.3, and let $\mathcal{O}$ be an ∞ -operad. Then postcomposition with the symmetric monoidal functors $i^\otimes$ and $L^\otimes$ induces an adjunction*

$$\mathrm{Alg}_{\mathcal{O}}(L) : \mathrm{Alg}_{\mathcal{O}}(C) \rightleftarrows \mathrm{Alg}_{\mathcal{O}}(D) : \mathrm{Alg}_{\mathcal{O}}(i)$$

in which the right adjoint is fully faithful. Its essential image consists of those $\mathcal{O}$ -algebras A in C whose underlying objects $A(o) \in C$ lie in D , for every color $o \in \mathcal{O}^\simeq$. The left adjoint is computed by L on underlying objects, so that $\mathrm{Alg}_{\mathcal{O}}(D)$ is the Bousfield localization of $\mathrm{Alg}_{\mathcal{O}}(C)$ at those maps of $\mathcal{O}$ -algebras whose underlying morphisms are L -equivalences.

Proof. By part (1) of Proposition 14.5.3, the symmetric monoidal ∞ -category $D^\otimes$ is a *full* suboperad of $C^\otimes$. Hence postcomposition with $i^\otimes$ is fully faithful, and an ∞ -operad map $O^\otimes \rightarrow C^\otimes$ factors through $D^\otimes$ if and only if it does so on the colors: an object of $C^\otimes$ lying over $\langle n \rangle$ decomposes, via the inert morphisms, into an n -tuple of objects of C , and an ∞ -operad map preserves inert morphisms. This gives full faithfulness of $\text{Alg}_O(i)$ together with the description of the essential image.

By part (2) of Proposition 14.5.3, the adjunction $L^\otimes \dashv i^\otimes$ is a symmetric monoidal adjunction. Applying Lemma 14.3.9 gives the asserted adjunction on O -algebras. The unit is computed colorwise by the unit of $L \dashv i$, so the left adjoint is computed by L on underlying objects and inverts precisely the maps whose underlying morphisms are L -equivalences. $\square$

Example 14.5.5. Taking $O = \text{Comm}$ in Corollary 14.5.4, we find that $\text{CAlg}(D) \subseteq \text{CAlg}(C)$ is a Bousfield localization: a commutative algebra of C lies in $\text{CAlg}(D)$ precisely when its underlying object is local, and the reflection is computed by applying L to the underlying object.

We record a special case for which the conditions of Proposition 14.5.3 are satisfied:

Lemma 14.5.6. *Let C be a symmetric monoidal ∞ -category which admits internal homs in the sense that the functor $- \otimes x: C \rightarrow C$ admits a right adjoint $\underline{\text{Hom}}_C(x, -): C \rightarrow C$ for every object x of C . Let D be a full subcategory of C such that the inclusion $D \hookrightarrow C$ admits a left adjoint, and assume that for objects $x \in D$ and $z \in C$ the internal hom $\underline{\text{Hom}}_C(z, x)$ is again in D . Then D obtains a symmetric monoidal structure such that the localization functor $L: C \rightarrow D$ is symmetric monoidal.*

Proof. We must prove the condition from Proposition 14.5.3. Let $f: x \rightarrow y$ be an L -equivalence and let $z \in C$ be an arbitrary object. We must show that the morphism $L(f \otimes \text{id}_z): L(x \otimes z) \rightarrow L(y \otimes z)$ is invertible in D . By the Yoneda lemma, this amounts to showing that for every object $w \in D$ the map

$$(f \otimes \text{id}_z)^*: \text{Hom}_C(y \otimes z, w) \rightarrow \text{Hom}_C(x \otimes z, w)$$

is an equivalence of anima. Applying the adjunction $- \otimes z \dashv \underline{\text{Hom}}_C(z, -)$, this map can be identified with the map

$$f^*: \text{Hom}_C(y, \underline{\text{Hom}}_C(z, w)) \rightarrow \text{Hom}_C(x, \underline{\text{Hom}}_C(z, w)).$$

Since $\underline{\text{Hom}}_C(z, w) \in D$ by assumption, this map is an equivalence due to the fact that f is an L -equivalence. $\square$

The preceding results concern reflective localizations inside an ∞ -category. The complementary problem of constructing a monoidal structure on a Dwyer–Kan localization at a class of weak equivalences is treated later in Chapter 20.

Exercises

Exercise 14.1. Show that the following functors are examples of ∞ -operads:

- $\text{Comm} = (\text{Span}(\text{Fin}), \text{id})$;
- $\mathcal{T}\text{riv} = (\text{Fin}^{\text{op}}, \text{inclusion})$;
- $\mathcal{T}\text{riv}_\emptyset = (*, \text{inclusion})$;

- $\mathcal{E}_0 = (\text{Span}_{\text{all}, \text{inj}}(\text{Fin}), \text{inclusion})$.

In each case, compute the multimorphism sets and the corresponding composition of multimorphisms, and show that these ∞ -operads agree with their classical counterparts from Section 12.1.

Exercise 14.2. Show that Comm is a terminal object in Op_∞ .

Exercise 14.3. (1) Prove the converse of Lemma 14.2.8: if an ∞ -operad $\mathcal{O}$ is the multimorphism operad of a symmetric monoidal ∞ -category, then conditions (1) and (2) of that lemma are satisfied. Conclude that these two conditions characterize the essential image of $\mathcal{M}: \text{Cat}_\infty^\otimes \rightarrow \text{Op}_\infty$ on objects.

(2) Given symmetric monoidal ∞ -categories C and D , show that an operad map $f: \mathcal{M}_C \rightarrow \mathcal{M}_D$ lies in the image of the functor $\mathcal{M}$ if and only if the map $f^\otimes: C^\otimes \rightarrow D^\otimes$ preserves the cocartesian lifts $(x_1, \dots, x_n) \rightarrow x_1 \otimes \dots \otimes x_n$ of the span $\langle n \rangle \xleftarrow{=} \langle n \rangle \rightarrow \langle 1 \rangle$ for every $n \geq 0$. Use Observation 14.2.7 to pass from these one-output lifts to arbitrary active spans.

Exercise 14.4. Prove the following two statements:

- (1) The right adjoint R of a symmetric monoidal functor $L: C \rightarrow D$ admits a canonical lax symmetric monoidal structure.
- (2) The left adjoint L of a lax symmetric monoidal functor $R: D \rightarrow C$ admits a canonical symmetric monoidal structure whenever certain comparison maps

$$L\left(\bigotimes_{i=1}^n X_i\right) \rightarrow \bigotimes_{i=1}^n L(X_i)$$

are isomorphisms in D .

Formulate and prove versions of these statements for $\mathcal{O}$ -monoidal ∞ -categories for an arbitrary ∞ -operad $\mathcal{O}$.

15 Cartesian and cocartesian monoidal structures

Every ∞ -category C with finite products admits a *cartesian monoidal structure* in which the tensor product of two objects is their cartesian product; the higher coherences required to make this an honest symmetric monoidal structure are uniquely determined by the universal property of the product. Dually, if C admits finite coproducts, it admits a *cocartesian monoidal structure*. The goal of this chapter is to construct these two symmetric monoidal structures.

As a first step, we reformulate the problem in terms of the construction of coherent adjoints. Recall that C admits finite products if and only if for every finite set I the diagonal functor $C \rightarrow C^I$ admits a right adjoint; for $I = \emptyset$ this encodes the terminal object, while for $I = \{1, 2\}$ this returns the cartesian product. As a consequence, the restriction functors $C^I \rightarrow C^J$ have right adjoints for all morphisms of finite sets $J \rightarrow I$. Our task in constructing the cartesian monoidal structure on C is then to assemble both the contravariant restrictions as well as their covariant right adjoints into a single functor

$$\mathrm{Span}(\mathrm{Fin}) \rightarrow \mathrm{Cat}_\infty, \quad I \mapsto C^I.$$

The cocartesian monoidal structure should be constructed similarly by extending by left adjoints instead.

The general construction that allows us to carry this out is due to Barwick [Bar17], and is known as *unfurling*. Its input is a contravariant functor whose restriction functors admit left or right adjoints satisfying a certain compatibility known as the *Beck–Chevalley condition*. Its output is a functor on a span category, which on backwards legs restricts to the original functor and on forward legs implements the adjoints. Applying the unfurling construction to $I \mapsto C^I$ then produces the two monoidal structures we are after.

If C is not assumed to admit coproducts, the first steps in the unfurling procedure still make sense and produce an ∞ -operad O_C^{II} , which we refer to as the *cocartesian operad of C* . The reason for this is that the universal property of a coproduct $x_1 \sqcup \cdots \sqcup x_n$ tells us how to map out of it: we need to specify maps out of each object x_i . Even if the coproduct does not actually exist, the formula

$$O_C^{\mathrm{II}}(x_1, \dots, x_n; y) \simeq \prod_{i=1}^n \mathrm{Hom}_C(x_i, y)$$

can be used to define the multimorphism anima of an ∞ -operad.

The algebras over ∞ -operads of the form O_C^{II} are surprisingly simple: they correspond to C -indexed families of commutative algebras. To see why this is the case at an intuitive level, note that the

cocartesian operad for the terminal ∞ -category $*$ is precisely the commutative operad $Comm$. The assignment $C \mapsto O_C^{\amalg}$ is functorial, and so every object of C determines an operad map $Comm \rightarrow O_C^{\amalg}$, and hence evaluating an $O_C^{\amalg}$ -algebra at this object produces a commutative algebra. We will show in Section 15.2 that an $O_C^{\amalg}$ -algebra is completely captured by the resulting C -indexed family of commutative algebras.

The classification of algebras over cocartesian operads directly leads to a classification of cocartesian operads themselves: an ∞ -operad O is of the form $O_C^{\amalg}$ for some C if and only if the multimorphism anima in O from a disjoint union of unordered tuples is the product of the individual multimorphism animae out of each of the constituents. This then gives a direct classification of the cocartesian monoidal structures, resulting in an equivalence of ∞ -categories

$$\mathrm{Cat}_{\infty}^{\mathrm{coprod}} \xrightarrow{\sim} \mathrm{Cat}_{\infty}^{\otimes, \mathrm{cocart}}$$

between ∞ -categories with finite coproducts and cocartesian symmetric monoidal ∞ -categories. The analogous result for cartesian monoidal structures follows by duality.

Here and below, $\mathrm{Cat}_{\infty}^{\mathrm{coprod}}$ denotes the ∞ -category of ∞ -categories with finite coproducts and finite-coproduct-preserving functors.

We end the chapter by proving a characterization of algebras in cartesian monoidal structures: for every ∞ -category with finite products C and every ∞ -operad O , we construct an equivalence of ∞ -categories

$$\mathrm{Alg}_O(C, \times) \xrightarrow{\sim} \mathrm{Mon}_O(C)$$

between O -algebras and O -monoids, i.e. finite-product-preserving functors $O^{\otimes} \rightarrow C$. In particular, commutative algebras in $(C, \times)$ are nothing but the commutative monoids studied in Chapter 5. The proof of this equivalence proceeds by constructing an alternative model for the cartesian monoidal structure on C .

The unfurling theorem is the technical input for the chapter. On a first reading, the proofs in Section 15.1.1 may be skipped. Several statements established there will be used later, notably the criterion for cocartesian morphisms in a span category and the adequate triple associated with a cartesian fibration.

15.1 Unfurling

In this section, we explain Barwick’s *unfurling construction* [Bar17], which extends a contravariant functor $F: C^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}$ to the span category by coherently assembling adjoints. We begin by isolating the required adjointability condition.

Definition 15.1.1. Let (C, C, C_R) be an adequate triple and let $F: C^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}$ be a functor.

- (1) The functor F is *left C_R -adjointable* if, for every $r: X \rightarrow Y$ in C_R , the functor $r^* := F(r): F(Y) \rightarrow F(X)$ admits a left adjoint $r_!$, and for every pullback square

$$\begin{array}{ccc} X' & \xrightarrow{r'} & X \\ \downarrow \iota' & \lrcorner & \downarrow \iota \\ Y' & \xrightarrow{r} & Y \end{array}$$

with $r, r' \in C_R$, the Beck–Chevalley transformation

$$r'_! l'^* \xrightarrow{r'_! l'^* \eta_r} r'_! l'^* r^* r_! \simeq r'_! r'^* l^* r_! \xrightarrow{\varepsilon_{r'} l^* r_!} l^* r_!$$

is a natural isomorphism.

- (2) The functor F is *right C_R -adjointable* if every r^* admits a right adjoint r_* and, for every pullback square as above, the dually defined Beck–Chevalley transformation $l^* r_* \rightarrow r'_! l'^*$ is a natural isomorphism.

When $C_R = C$, we simply say that F is *left adjointable* or *right adjointable*.

The trick for extending F to the span category is by reformulating the problem in terms of (co)cartesian fibrations: starting with the unstraightening of F , we wish to construct the unstraightening of its extension. We now provide a formulation of the adjointability conditions on F in terms of its associated fibration over C .

Definition 15.1.2. Let (C, C_L, C_R) be an adequate triple. A functor $p: E \rightarrow C$ is called a *Beck–Chevalley fibration* with respect to (C_L, C_R) if the following three conditions are satisfied:

- (1) Every morphism in C_L admits p -cartesian lifts with given target.
- (2) Every morphism in C_R admits p -cocartesian lifts with given source.
- (3) In a commutative square

$$\begin{array}{ccc} \tilde{X}' & \xrightarrow{\tilde{r}'} & \tilde{X} \\ \tilde{l}' \downarrow & & \downarrow \tilde{l} \\ \tilde{Y}' & \xrightarrow{\tilde{r}} & \tilde{Y} \end{array}$$

whose image in C is a pullback square with $p(\tilde{l}) \in C_L$ and $p(\tilde{r}) \in C_R$, assume that $\tilde{r}$ is p -cocartesian and $\tilde{l}'$ is p -cartesian. Then $\tilde{r}'$ is p -cocartesian if and only if $\tilde{l}$ is p -cartesian.

Conditions (1) and (2) say that the restrictions $E \times_C C_L \rightarrow C_L$ and $E \times_C C_R \rightarrow C_R$ are a cartesian and a cocartesian fibration, respectively. They provide contravariant transport l^* along C_L , covariant transport $r_!$ along C_R , and Beck–Chevalley transformations $r'_! l'^* \rightarrow l^* r_!$ for pullback squares as above. By choosing cartesian and cocartesian lifts around such a square, one sees that condition (3) is precisely the invertibility of these transformations.

The following result is the key technical input to the unfurling construction.

Theorem 15.1.3. Let (C, C_L, C_R) be an adequate triple and let $p: E \rightarrow C$ be a Beck–Chevalley fibration with respect to (C_L, C_R) . Set $E_L^{p\text{-cart}} := p^{-1}(C_L) \cap E^{p\text{-cart}}$ and $E_R := p^{-1}(C_R)$. Then $(E, E_L^{p\text{-cart}}, E_R)$ is an adequate triple, and the induced functor

$$\text{Span}(p): \text{Span}_{\text{ct}, R}(E) \longrightarrow \text{Span}_{L, R}(C)$$

is a cocartesian fibration with fiber E_X over $X \in C$. Its transport along a span $X \xleftarrow{l} U \xrightarrow{r} Y$ is $r_! l^*$.

Here $\text{Span}_{\text{ct}, R}(E)$ denotes the span category of the adequate triple just defined. For expository purposes, we postpone the proof to Section 15.1.1.

Corollary 15.1.4 (Contravariant left-adjoint unfurling). *Let (C, C, C_R) be an adequate triple. Every left C_R -adjointable functor $F: C^{\text{op}} \rightarrow \text{Cat}_\infty$ extends canonically to a functor*

$$\text{Unf}^L(F): \text{Span}_{\text{all}, R}(C) \longrightarrow \text{Cat}_\infty$$

which sends a span $X \xleftarrow{l} U \xrightarrow{r} Y$ to $r_! l^$. Its restriction along $C^{\text{op}} \hookrightarrow \text{Span}_{\text{all}, R}(C)$ is F itself.*

Proof. Let $p: \text{Un}^{\text{cl}}(F) \rightarrow C$ be the cartesian unstraightening. The dual of Lemma 23.1.20, applied over each $r \in C_R$, identifies the existence of $r_!$ with that of p -cocartesian lifts over r . The Beck–Chevalley condition for F is condition (3) of Definition 15.1.2, so Theorem 15.1.3 gives the extension and its transport formula. The descriptions of the fibers and backwards transport identify its restriction with F . $\square$

Corollary 15.1.5 (Contravariant right-adjoint unfurling). *Let (C, C, C_R) be an adequate triple. Every right C_R -adjointable functor $F: C^{\text{op}} \rightarrow \text{Cat}_\infty$ extends canonically to a functor*

$$\text{Unf}^R(F): \text{Span}_{\text{all}, R}(C) \longrightarrow \text{Cat}_\infty$$

which sends a span $X \xleftarrow{l} U \xrightarrow{r} Y$ to $r_ l^*$. Its restriction along $C^{\text{op}} \hookrightarrow \text{Span}_{\text{all}, R}(C)$ is F itself.*

Proof. The pointwise opposite $F^{\text{op}}: C^{\text{op}} \rightarrow \text{Cat}_\infty$ is left C_R -adjointable: the left adjoint to $(r^*)^{\text{op}}$ is $(r_*)^{\text{op}}$, and its Beck–Chevalley transformations are the opposites of those for F . Apply Corollary 15.1.4 to F^{op} and take opposites pointwise once more; the transport functor becomes $r_* l^*$. $\square$

For completeness, we record the covariant form which appears in Barwick’s original construction. Let $F: C \rightarrow \text{Cat}_\infty$ be a functor and let $C_L \subseteq C$ be a wide subcategory such that (C, C_L, C) is an adequate triple. Write $l_! := F(l)$. We call F *covariantly right C_L -adjointable* if every $l_!$ admits a right adjoint l^* and, for every pullback square as in Definition 15.1.1 with $l, l' \in C_L$, the Beck–Chevalley transformation $r'_! l'^* \rightarrow l^* r_!$ is a natural isomorphism.

Proposition 15.1.6 (Covariant right-adjoint unfurling). *The cocartesian unstraightening $p: \text{Un}^{\text{cc}}(F) \rightarrow C$ of a functor $F: C \rightarrow \text{Cat}_\infty$ is a Beck–Chevalley fibration with respect to (C_L, C) if and only if F is covariantly right C_L -adjointable. Consequently, such an F extends to $\text{Span}_{L, \text{all}}(C)$ by sending $X \xleftarrow{l} U \xrightarrow{r} Y$ to $r_! l^*$.*

Proof. Condition (2) of Definition 15.1.2 holds automatically, since p is a cocartesian fibration. For every $l \in C_L$, we may apply Lemma 23.1.20 after pulling p back along $l: [1] \rightarrow C$. It follows that cartesian lifts over l exist precisely when $l_!$ admits a right adjoint l^* .

Now consider a pullback square as in Definition 15.1.1, with $l, l' \in C_L$. The canonical commutativity isomorphism $r_! l'_! \simeq l_! r'_!$, together with the units and counits of the adjunctions, determines the Beck–Chevalley transformation

$$r'_! l'^* \longrightarrow l^* r_!.$$

Under the correspondence between right adjoints and cartesian lifts, the invertibility of this transformation is exactly condition (3) of Definition 15.1.2. This proves the first claim. The final assertion now follows from Theorem 15.1.3. $\square$

Exercise 15.1.7. Formulate and prove a covariant left-adjoint unfurling result.

15.1.1 Cocartesian morphisms in span categories

We now provide a proof of Theorem 15.1.3. The key input is a general criterion for morphisms in span categories to be cocartesian. This criterion is useful beyond the unfurling application, notably when constructing cocartesian operads that need not arise from symmetric monoidal structures. The results here are of a technical nature and may be skipped on a first reading.

Theorem 15.1.8 (Barwick [Bar17], [HHLN23b, Theorem 3.1]). *Let $p : (E, E_L, E_R) \rightarrow (C, C_L, C_R)$ be a morphism of adequate triples, and consider a span*

$$X \xleftarrow{\varphi} U \xrightarrow{\psi} Y,$$

with $\varphi \in E_L$ and $\psi \in E_R$. Then this morphism is cocartesian with respect to $\text{Span}(p) : \text{Span}_{L,R}(E) \rightarrow \text{Span}_{L,R}(C)$ whenever the following conditions are satisfied:

- (1) *The morphism φ is p -cartesian and the morphism ψ is p -cocartesian;*
- (2) *(Left cancellation for E_L) Given a commutative triangle*

$$\begin{array}{ccc} & U & \\ \xi \nearrow & & \searrow \varphi \\ T & \xrightarrow{\varphi\xi} & X \end{array}$$

such that $p(\xi) \in C_L$ and $\varphi\xi \in E_L$, we have $\xi \in E_L$;

- (3) *(Right cancellation for E_R) Given a commutative triangle*

$$\begin{array}{ccc} & Y & \\ \psi \nearrow & & \searrow \xi \\ U & \xrightarrow{\xi\psi} & T \end{array}$$

such that $p(\xi) \in C_R$ and $\xi\psi \in E_R$, we have $\xi \in E_R$;

- (4) *(Enough p -cocartesian lifts) For any pullback square in C of the form*

$$\begin{array}{ccc} K & \xrightarrow{g} & J \\ l' \downarrow & \lrcorner & \downarrow l \\ pU & \xrightarrow{p\psi} & pY, \end{array}$$


with $l \in C_L$, and for any $W \in E_K$, the morphism g admits a p -cocartesian lift $\psi' : W \rightarrow V$ which lies in E_R and also satisfies the condition from (3);

- (5) *(Beck–Chevalley) Consider a commutative square in E of the form*

$$\begin{array}{ccc} W & \xrightarrow{\psi'} & V \\ \xi' \downarrow & & \downarrow \xi \\ U & \xrightarrow{\psi} & Y \end{array}$$

such that its image in C is a pullback square, $\xi' \in E_L$, $\psi' \in E_R$, and $p(\xi) \in C_L$. Then ψ' is p -cocartesian if and only if $\xi \in E_L$ and the square is a pullback square.

Proof. We consider any other span $X \leftarrow W \rightarrow Z$ in E . We have to show that we may uniquely fill the following dashed diagram



whenever we are given a similar filling for the resulting diagram in C ; here all left-pointing maps are in E_L and all right-pointing maps are in E_R . By the description of the hom anima of a span category from Lemma 13.1.13, the anima of such fillings is precisely the fiber of the comparison map

$$\mathrm{Hom}_{\mathrm{Span}_{L,R}(E)}(Y, Z) \longrightarrow \mathrm{Hom}_{\mathrm{Span}_{L,R}(E)}(X, Z) \times_{\mathrm{Hom}_{\mathrm{Span}_{L,R}(C)}(pX, pZ)} \mathrm{Hom}_{\mathrm{Span}_{L,R}(C)}(pY, pZ)$$

over the given point, so that showing all these anima to be contractible is exactly showing that our span is $\mathrm{Span}(p)$ -cocartesian. We produce the filling in three steps, each of which is unique up to a contractible choice.

- (i) Since φ is p -cartesian, there exists a unique lift $W \rightarrow U$ making the left triangle commute, which lies in E_L by condition (2).
- (ii) We claim that this lift $W \rightarrow U$ may uniquely be written as the pullback along ψ of some morphism $V \rightarrow Y$ in E_L compatible with the given pullback square in C ; more precisely, we claim that the following commutative square is a pullback square:

$$\begin{array}{ccc} (E/Y)^L & \xrightarrow{\psi^*} & (E/U)^L \\ p \downarrow & & \downarrow p \\ (C/pY)^L & \xrightarrow{(p\psi)^*} & (C/pU)^L. \end{array}$$

Here the superscript $(-)^L$ means that we take the full subcategories spanned by the morphisms in E_L and C_L , respectively.

- (a) For essential surjectivity of $(E/Y)^L \rightarrow (E/U)^L \times_{(C/pU)^L} (C/pY)^L$, we consider a morphism $\xi': W \rightarrow U$ in E_L and assume we are given a pullback square in C of the form

$$\begin{array}{ccc} pW & \xrightarrow{g} & J \\ p\xi' \downarrow & \lrcorner & \downarrow l \\ pU & \xrightarrow{p\psi} & pY \end{array}$$

with $l \in C_L$. By assumption (4) we can find a p -cocartesian lift $\psi': W \rightarrow V$ of g which lies in E_R and satisfies condition (3). The composite $\psi \circ \xi': W \rightarrow Y$ then factors through a map $\xi: V \rightarrow Y$, giving a commutative square in E of the form considered in (5). By assumption (5), the map ξ then lies in E_L and the resulting square is a pullback square. As ξ lifts l , this shows that the pair (ξ', l) lies in the essential image, as desired.

- (b) For full faithfulness, consider two objects $\xi_1: V_1 \rightarrow Y$ and $\xi_2: V_2 \rightarrow Y$ of $(E/Y)^L$. We must show that the commutative square

$$\begin{array}{ccc} \mathrm{Hom}_{E/Y}(V_1, V_2) & \xrightarrow{\psi^*} & \mathrm{Hom}_{E/U}(U \times_Y V_1, U \times_Y V_2) \\ \downarrow p & & \downarrow p \\ \mathrm{Hom}_{C/pY}(pV_1, pV_2) & \xrightarrow{(p\psi)^*} & \mathrm{Hom}_{C/pU}(pU \times_{pY} pV_1, pU \times_{pY} pV_2) \end{array}$$

is a pullback square. By the universal property of pullbacks, the right vertical map is isomorphic to the map

$$p: \mathrm{Hom}_{E/Y}(U \times_Y V_1, V_2) \rightarrow \mathrm{Hom}_{C/pY}(pU \times_{pY} pV_1, pV_2),$$

hence the claim follows from the fact that the projection map $U \times_Y V_1 \rightarrow V_1$ is p -cocartesian: this is the implication from right to left in assumption (5), applied to the pullback square of ξ_1 along ψ , whose remaining two sides lie in E_L and E_R by adequacy. This morphism remains cocartesian after passing to the slice over Y , since its cocartesian universal property is unchanged when the target is fixed.

- (iii) Finally, given a p -cocartesian morphism $W \rightarrow V$, the given map $W \rightarrow Z$ uniquely factors through a map $V \rightarrow Z$, which lies in E_R by condition (3). $\square$

In practice, we will often be interested in the situation where the subcategories E_L and E_R of E are determined by C_L , C_R and the functor $p: E \rightarrow C$. To set this up, we start with the following auxiliary lemma:

Lemma 15.1.9. *Let $p: E \rightarrow C$ be a functor, and consider a commutative square*

$$\begin{array}{ccc} X' & \xrightarrow{r'} & X \\ l' \downarrow & & \downarrow l \\ Y' & \xrightarrow{r} & Y \end{array}$$

in E where l is a p -cartesian morphism. Assume that the image of this square in C is a pullback square. Then this square is a pullback square if and only if the morphism l' is p -cartesian.

Proof. Consider for any object $Z \in E$ the following two diagrams:

$$\begin{array}{ccccc} \mathrm{Hom}_E(Z, X') & \xrightarrow{r' \circ -} & \mathrm{Hom}_E(Z, X) & \xrightarrow{p} & \mathrm{Hom}_C(pZ, pX) \\ l' \circ - \downarrow & & \downarrow l \circ - & \lrcorner & \downarrow pl \circ - \\ \mathrm{Hom}_E(Z, Y') & \xrightarrow{r \circ -} & \mathrm{Hom}_E(Z, Y) & \xrightarrow{p} & \mathrm{Hom}_C(pZ, pY), \end{array}$$

$$\begin{array}{ccccc} \mathrm{Hom}_E(Z, X') & \xrightarrow{p} & \mathrm{Hom}_C(pZ, pX') & \xrightarrow{pr' \circ -} & \mathrm{Hom}_C(pZ, pX) \\ l' \circ - \downarrow & & \downarrow pl' \circ - & \lrcorner & \downarrow pl \circ - \\ \mathrm{Hom}_E(Z, Y') & \xrightarrow{p} & \mathrm{Hom}_C(pZ, pY') & \xrightarrow{pr \circ -} & \mathrm{Hom}_C(pZ, pY). \end{array}$$

The right-hand squares of both diagrams are pullback squares: for the first one since l is p -cartesian, for the second one since the image of the given square in C is a pullback square. As the functor

p preserves composition, the outer rectangles of both diagrams agree, so by the pasting law the left-hand square in the first diagram is a pullback square if and only if the left-hand square in the second diagram is a pullback square. Since the former holds for all Z if and only if the given diagram is a pullback square and the latter holds for all Z if and only if l' is p -cartesian, this finishes the proof. $\square$

Proposition 15.1.10. *Let (C, C_L, C_R) be an adequate triple and let $p: E \rightarrow C$ be a functor such that every morphism in C_L admits p -cartesian lifts with given target (for example: p is a cartesian fibration). Then:*

- (1) *The triple $(E, E_L^{p\text{-cart}}, E_R)$ is adequate, where $E_R := p^{-1}(C_R)$ consists of morphisms over C_R and $E_L^{p\text{-cart}} := p^{-1}(C_L) \cap E^{p\text{-cart}}$ consists of p -cartesian morphisms over C_L .*
- (2) *The functor $p: E \rightarrow C$ is a morphism of adequate triples.*

Proof. Both classes are wide subcategories of E : this is clear for $E_R = p^{-1}(C_R)$, while $E_L^{p\text{-cart}}$ contains all identities and is closed under composition since C_L is and since composites of p -cartesian morphisms are again p -cartesian.

We next construct the pullback in E of a morphism $l \in E_L^{p\text{-cart}}$ along some $r \in E_R$. Since $pl \in C_L$ and $pr \in C_R$, we may form the pullback

$$\begin{array}{ccc} pX \times_{pY} pY' & \longrightarrow & pX \\ \downarrow & \lrcorner & \downarrow pl \\ pY' & \xrightarrow{pr} & pY \end{array}$$

in C . By assumption on p , there exists a p -cartesian lift $l': X' \rightarrow Y'$ of the morphism $pX \times_{pY} pY' \rightarrow pY'$. The composite $rl': X' \rightarrow Y$ then uniquely factorizes through some $r': X' \rightarrow X$ lifting the projection $pX \times_{pY} pY' \rightarrow pX$, using that l is p -cartesian. All in all, we have lifted the pullback square in C to a commutative square in E of the form

$$\begin{array}{ccc} X' & \xrightarrow{r'} & X \\ l' \downarrow & & \downarrow l \\ Y' & \xrightarrow{r} & Y. \end{array} \tag{15.1}$$

Since l and l' are p -cartesian, it follows from Lemma 15.1.9 that this square is a pullback square in E , showing that the required pullbacks exist. Moreover, in this square we have $l' \in E_L^{p\text{-cart}}$ and $r' \in E_R$, so both classes are closed under the relevant base changes. This shows part (1).

For part (2), consider a pullback square in E of the form (15.1). We need to show that applying p to this square gives a pullback square in C . Proceeding as before, we may always form the pullback of pl and pr in C and lift this to a pullback square in E . But by uniqueness of pullback squares of l and r , this new square must then agree with the given one. In particular their images in C agree, hence this image is a pullback square. $\square$

Remark 15.1.11. For the morphism of adequate triples in the previous result, parts (2) and (3) of Theorem 15.1.8 follow from cancellation of p -cartesian morphisms and the definition of E_R ,

respectively. For a p -cocartesian morphism $\psi: U \rightarrow Y$ satisfying part (4), part (5) is precisely the invertibility of the Beck–Chevalley transformations

$$g_! l'^* \longrightarrow l^*(p\psi)_!: E_{pU} \longrightarrow E_J$$

for the pullback squares in part (4). Here $(p\psi)_!$ exists by taking $J = pY$ and $l = \text{id}$ in that condition.

We are now ready for the main result of this section.

Proof of Theorem 15.1.3. Condition (1) of Definition 15.1.2 and Proposition 15.1.10 show that $(E, E_L^{p\text{-cart}}, E_R)$ is adequate and that

$$\text{Span}(p): \text{Span}_{\text{ct}, R}(E) \longrightarrow \text{Span}_{L, R}(C)$$

is defined. Given a span $X \xleftarrow{l} U \xrightarrow{r} Y$ and an object over X , choose a p -cartesian lift of l followed by a p -cocartesian lift of r . In the criterion of Theorem 15.1.8, parts (2) and (3) hold by the preceding remark, part (4) follows from condition (2) of Definition 15.1.2 because g is a base change of r , and part (5) follows from condition (3) and Lemma 15.1.9. Hence the chosen span is cocartesian, with transport $r_! l^*$.

Finally, by Lemma 13.1.13, a morphism in the fiber over X is represented by a span whose backwards leg is p -cartesian over id_X , hence an isomorphism, and whose forward leg is an arbitrary morphism of E_X . Thus this fiber is E_X . $\square$

15.2 Cocartesian monoidal structures

We now specialize the unfurling construction to construct cocartesian monoidal structures on ∞ -categories with finite coproducts. Throughout, we will make use of the ∞ -category of finite families from Construction 14.1.15: for an ∞ -category C , we write $q: \text{Fin}(C) \rightarrow \text{Fin}$ for the cartesian unstraightening of the finite-product-preserving functor $C^{(-)}: \text{Fin}^{\text{op}} \rightarrow \text{Cat}_{\infty}$, $I \mapsto C^I$, so that the objects of $\text{Fin}(C)$ are the finite unordered tuples $\{x_i\}_{i \in I}$ of objects of C .

Assume now that C admits finite coproducts. For every map $f: I \rightarrow J$, the restriction functor $f^*: C^J \rightarrow C^I$ admits a left adjoint $f_!: C^J \rightarrow C^I$, which forms coproducts over the fibers of f . These adjunctions satisfy the Beck–Chevalley condition. Indeed, consider a pullback square as in Definition 15.1.1 and a family $A = (A_{y'})_{y' \in Y'}$. For every $x \in X$, the Beck–Chevalley transformation is the natural isomorphism

$$(r'_! l'^* A)_x \simeq \coprod_{x' \in (r')^{-1}(x)} A_{l'(x')} \simeq \coprod_{y' \in r^{-1}(l(x))} A_{y'} \simeq (l^* r_! A)_x,$$

where the middle isomorphism follows from the pullback property. Thus $C^{(-)}$ is left adjointable, and we may apply contravariant unfurling.

Proposition 15.2.1. *Let C be an ∞ -category with finite coproducts. The unfurling of $C^{(-)}$ is a finite-product-preserving functor*

$$(C, \amalg) := \text{Unf}^L(C^{(-)}): \text{Span}(\text{Fin}) \longrightarrow \text{Cat}_{\infty}$$

which defines a symmetric monoidal structure on C whose unit is the initial object and whose tensor product is the coproduct.

Proof. By Corollary 15.1.4, the functor $C^{(-)}$ extends to $\text{Span}(\text{Fin})$. To check whether it preserves finite products, recall from part (1) of Lemma 13.3.5 that the inclusion $\text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin})$ preserves finite products. Since this inclusion is the identity on objects, we may test whether the unfurling preserves finite products after restricting along it. By the last statement of Corollary 15.1.4, the resulting functor is the original functor $C^{(-)}$, which preserves finite products by construction. This finishes the construction of the monoidal structure.

Given the description of coproducts in terms of left adjoints, we see that transport along $\emptyset \rightarrow \langle 1 \rangle$ selects the initial object, while transport along the active span $\langle 2 \rangle \xleftarrow{=} \langle 2 \rangle \xrightarrow{\nabla} \langle 1 \rangle$ is the binary coproduct. $\square$

Definition 15.2.2 (Cocartesian monoidal structure). The symmetric monoidal ∞ -category (C, Π) from Proposition 15.2.1 is called the *cocartesian monoidal structure on C* .

15.2.1 Cocartesian operads

The construction of the cartesian fibration $q: \text{Fin}(C) \rightarrow \text{Fin}$ did not use the existence of finite coproducts in C . Those coproducts were needed only to supply cocartesian transport along all morphisms of Fin and thereby make q a Beck–Chevalley fibration. For an arbitrary ∞ -category C , we may still apply the span-category construction to q . The resulting map need not be a cocartesian fibration over all of $\text{Span}(\text{Fin})$, but it retains the cocartesian lifts over inert morphisms required of an ∞ -operad. We now make this precise.

Recall from Construction 14.1.15 that $\text{Fin}(C)$ admits finite coproducts, given by unordered concatenation of tuples, that q preserves them, and that a morphism $(f, (\varphi_i)_{i \in I})$ of $\text{Fin}(C)$ is q -cartesian if and only if every φ_i is an isomorphism in C ; we write $\text{Fin}(C)_{\text{ct}} \subseteq \text{Fin}(C)$ for the wide subcategory spanned by the q -cartesian morphisms.

Lemma 15.2.3. *Let C be an ∞ -category. Then the triple $(\text{Fin}(C), \text{Fin}(C)_{\text{ct}}, \text{Fin}(C))$ is an adequate triple. Furthermore, it is extensive, in the sense of Definition 13.3.1.*

Proof. The adequate triple is obtained by applying Proposition 15.1.10 to the cartesian fibration $q: \text{Fin}(C) \rightarrow \text{Fin}$. For extensiveness, we check the conditions from Definition 13.3.1:

- (1) We have just recalled that $\text{Fin}(C)$ admits finite coproducts, given by unordered concatenation.
- (2) The q -cartesian morphisms are closed under coproducts. It readily follows from Lemma 15.1.9 that coproducts of pullback squares along q -cartesian morphisms are again pullback squares.
- (3) The morphisms $\emptyset \rightarrow X$ and $\nabla: X \sqcup X \rightarrow X$ in $\text{Fin}(C)$ are q -cartesian.
- (4) The fact that the commutative squares

$$\begin{array}{ccc} X \sqcup X & \xrightarrow{f \sqcup f} & Y \sqcup Y \\ \nabla \downarrow & & \downarrow \nabla \\ X & \xrightarrow{f} & Y \end{array} \quad \text{and} \quad \begin{array}{ccc} \emptyset & \xrightarrow{=} & \emptyset \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

are pullback squares again follows from Lemma 15.1.9, since their images in Fin are pullback squares and the vertical maps are q -cartesian.

This proves weak extensivity. The same arguments also prove weak coextensivity: conditions (1) and (3) are automatic when the left class is all morphisms, while conditions (2) and (4) are precisely the corresponding assertions verified above. Hence the triple is extensive. $\square$

Construction 15.2.4 (Cocartesian operad). Let C be an ∞ -category. We define the ∞ -category $C^{\amalg}$ as

$$C^{\amalg} := \text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C)).$$

Combining Lemma 15.2.3 with Lemma 13.3.5, we see that this is a semiadditive ∞ -category, with finite products and finite coproducts both induced by the unordered concatenation of tuples in C . We denote by

$$p_C^{\amalg} := \text{Span}(q): C^{\amalg} \rightarrow \text{Span}(\text{Fin})$$

the induced functor to $\text{Span}(\text{Fin})$, which preserves finite products.

Proposition 15.2.5. *Let C be an ∞ -category. Then the pair $(C^{\amalg}, p_C^{\amalg})$ is an ∞ -operad.*

Proof. We have already argued that $C^{\amalg} = \text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C))$ has finite products and that $p_C^{\amalg} = \text{Span}(q)$ preserves them, giving condition (1) of the definition. The objects in the fiber over a finite set I form the fiber $\text{Fin}(C)_I = C^I$. Moreover, by Lemma 13.1.13, a morphism over the identity span of I is represented by a span whose backwards leg is q -cartesian over id_I , hence an isomorphism, and therefore reduces to a morphism in C^I . Thus the fiber of $C^{\amalg}$ over I is C^I , giving condition (2). For condition (3), consider objects $X_i \in \text{Fin}(C)$ and a morphism $f: J \rightarrow I$ of finite sets. Then the lift $\tilde{f}: \prod_{i \in I} X_i \rightarrow \prod_{j \in J} X_{f(j)}$ in $\text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C))$ is a backwards span of the form

$$\bigsqcup_{i \in I} X_i \xleftarrow{(f, (\text{id})_f)} \bigsqcup_{j \in J} X_{f(j)} \xrightarrow{(\text{id}, (\text{id}))} \bigsqcup_{j \in J} X_{f(j)}.$$

The left-pointing morphism is q -cartesian and the right-pointing morphism is an identity. In particular, condition (4) of Theorem 15.1.8 only asks for cocartesian lifts of base changes of an identity, which are again isomorphisms; the remaining conditions follow directly from cartesianness and Lemma 15.1.9. Hence this span is $\text{Span}(q)$ -cocartesian, as desired. $\square$

The proof only uses cocartesian lifts over inert morphisms, for which the forward leg of the displayed span is an identity. Thus it does not require the covariant transport $f_!$ used in the full unfurling.

Definition 15.2.6 (Cocartesian operad). For an ∞ -category C , we refer to the ∞ -operad $O_C^{\amalg} := (C^{\amalg}, p_C^{\amalg})$ from Proposition 15.2.5 as the *cocartesian operad of C* .

Warning 15.2.7. In various sources, notably in Lurie [Lur17], no distinction is made between the ∞ -operad $O_C^{\amalg}$ and its total ∞ -category $C^{\amalg}$: both are denoted by $C^{\amalg}$. Although the notation $O_C^{\amalg}$ is slightly more verbose, we find it useful to distinguish between the operad and its total category.

The operad $O_C^{\amalg}$ does precisely what we want it to do: its multimorphisms $(x_1, \dots, x_n) \rightarrow y$ are simply the morphisms $(x_1, \dots, x_n) \rightarrow y$ in $\text{Fin}(C)$, which correspond to tuples $(\varphi_i: x_i \rightarrow y)$ of morphisms in C . If C admits finite coproducts, cocartesian transport along the active fold map $\langle n \rangle \rightarrow \langle 1 \rangle$ sends $(x_1, \dots, x_n)$ to $x_1 \amalg \dots \amalg x_n$. Factoring through this cocartesian lift gives the familiar isomorphism

$$\text{Hom}_C(x_1 \amalg \dots \amalg x_n, y) \simeq \prod_{i=1}^n \text{Hom}_C(x_i, y).$$

The construction of $O_C^{\mathbb{I}}$ retains the expression on the right even when the coproduct representing it does not exist.

Proposition 15.2.8. *Let C be an ∞ -category. Then the following conditions are equivalent:*

- (1) *The ∞ -category C admits finite coproducts;*
- (2) *The cartesian fibration $q: \text{Fin}(C) \rightarrow \text{Fin}$ is a cocartesian fibration;*
- (3) *The functor $p_C^{\mathbb{I}} = \text{Span}(q): C^{\mathbb{I}} \rightarrow \text{Span}(\text{Fin})$ is a cocartesian fibration.*

If these conditions hold, $p_C^{\mathbb{I}}$ is the cocartesian unstraightening of the unfurling $(C, \mathbb{I})$ from Proposition 15.2.1.

Proof. If C admits finite coproducts, we showed above that $q: \text{Fin}(C) \rightarrow \text{Fin}$ is a Beck–Chevalley fibration. The construction in Theorem 15.1.3 identifies the cocartesian unstraightening of the unfurling from Proposition 15.2.1 with $\text{Span}(q)$. This proves that (1) implies (3), as well as the final assertion.

We next show that (3) implies (2). Recall from the dual of Exercise 23.1.15 that every q -cartesian morphism in $\text{Fin}(C)$ whose image in Fin is an isomorphism is itself an isomorphism. Consequently, the square

$$\begin{array}{ccccc} \text{Fin}(C) & \xrightarrow{\simeq} & \text{Span}_{\simeq, \text{all}}(\text{Fin}(C)) & \hookrightarrow & \text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C)) \\ q \downarrow & & \downarrow & \lrcorner & \downarrow \text{Span}(q) \\ \text{Fin} & \xrightarrow{\simeq} & \text{Span}_{\simeq, \text{all}}(\text{Fin}) & \hookrightarrow & \text{Span}(\text{Fin}) \end{array}$$

is a pullback square. Since cocartesian fibrations are closed under pullback, (3) implies (2).

Finally, suppose that q is a cocartesian fibration. By Lemma 23.1.20, its cocartesian transport $f_!: C^I \rightarrow C^J$ along a map $f: I \rightarrow J$ is left adjoint to the cartesian transport $f^*: C^J \rightarrow C^I$. For the map $\emptyset \rightarrow \langle 1 \rangle$ this produces an initial object of C , while for the fold map $\langle 2 \rangle \rightarrow \langle 1 \rangle$ it produces binary coproducts. Thus C admits finite coproducts, proving that (2) implies (1). $\square$

Thus, when C admits finite coproducts, the symmetric monoidal structure associated to $O_C^{\mathbb{I}}$ is precisely the cocartesian monoidal structure constructed by unfurling at the beginning of the section.

Remark 15.2.9. Note that all the steps in the construction of $O_C^{\mathbb{I}} = (C^{\mathbb{I}}, p_C^{\mathbb{I}})$ are functorial in C . Using Proposition 15.1.10, we may write it as the composite

$$\text{Cat}_{\infty} \xrightarrow{\sim} \text{Fun}^{\times}(\text{Fin}^{\text{op}}, \text{Cat}_{\infty}) \xrightarrow{\text{Un}^{\text{ct}}} \text{Cart}(\text{Fin}) \xrightarrow{E \mapsto (E, E^{p\text{-cart}}, E)} \text{AdTrip}_{/\text{Fin}} \xrightarrow{\text{Span}} (\text{Cat}_{\infty})_{/\text{Span}(\text{Fin})}.$$

In particular, we obtain a functor

$$O^{\mathbb{I}}: \text{Cat}_{\infty} \rightarrow \text{Op}_{\infty}.$$

15.2.2 Algebras over cocartesian operads

Algebras over a cocartesian operad $O_C^{\mathbb{I}}$ are simple: as we will show next, they correspond to C -indexed families of commutative algebras.

Recall from Lemma 14.1.16 that $\text{Fin}(C)$ is freely generated by C under finite coproducts; this will be the key input below.

Construction 15.2.10. Let C be an ∞ -category. We construct a functor

$$\iota: \text{Span}(\text{Fin}) \times C \rightarrow C^{\mathbb{I}} = \text{Span}_{\text{ct,all}}(\text{Fin}(C)),$$

informally given by sending a pair (I, x) to the ‘constant’ I -tuple $(x)_{i \in I}$, and by sending a pair $(I \xleftarrow{f} J \xrightarrow{g} K, \varphi: x \rightarrow y)$ of a span in Fin and a morphism in C to the span

$$(x)_{i \in I} \xleftarrow{(f, (\text{id}_x)_{j \in J})} (x)_{j \in J} \xrightarrow{(g, (\varphi)_{j \in J})} (y)_{k \in K}.$$

(Note that the left-pointing morphism is indeed q -cartesian.) Formally, this map is constructed as follows:

- Consider the natural transformation $\Delta: \text{const}_C \rightarrow C^{(-)}$ of functors $\text{Fin}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ which on the one-point set is given by the identity $\text{id}_C: C \rightarrow C$; this property determines the entire transformation since $C^{(-)}$ is by definition right Kan extended from the point. For a finite set I , it is given by the diagonal functor

$$\Delta: C \rightarrow C^I, \quad x \mapsto (x)_{i \in I}.$$

- By cartesian unstraightening, we obtain a morphism of cartesian fibrations over Fin :

$$\begin{array}{ccc} \text{Fin} \times C & \xrightarrow{\text{Un}^{\text{ct}}(\Delta)} & \text{Fin}(C) \\ \text{pr}_1 \searrow & & \swarrow q \\ & \text{Fin}. & \end{array}$$

- By using Proposition 15.1.10, both sides become adequate triples with left class given by the cartesian morphisms. The functor $\text{Un}^{\text{ct}}(\Delta)$ preserves cartesian morphisms and becomes a morphism of adequate triples. Since a morphism on the left-hand side is cartesian if and only if the C -component is an isomorphism, we may use the equivalence $\text{Span}_{\simeq, \text{all}}(C) \simeq C$ to get a map on spans:

$$\begin{array}{ccc} \text{Span}(\text{Fin}) \times C & \xrightarrow{\iota} & \text{Span}_{\text{ct,all}}(\text{Fin}(C)) = C^{\mathbb{I}} \\ \text{pr}_1 \searrow & & \swarrow \text{Span}(q) \\ & \text{Span}(\text{Fin}). & \end{array}$$

Notation 15.2.11. Let C, D and E be ∞ -categories and assume D and E admit finite coproducts. We denote by

$$\text{Fun}^{\mathbb{I}, -}(E \times C, D) \subseteq \text{Fun}(E \times C, D)$$

the full subcategory consisting of functors $F: E \times C \rightarrow D$ that preserve finite coproducts in the first variable, in the sense that $F(-, x): E \rightarrow D$ preserves finite coproducts for all $x \in C$.

Lemma 15.2.12 ([BH21, Lemma C.4], cf. [CLL25, Proof of Proposition 3.30]). *Let C and D be ∞ -categories and assume D admits finite coproducts. Then restriction along ι induces an equivalence*

$$\iota^*: \text{Fun}^{\mathbb{I}}(\text{Span}_{\text{ct,all}}(\text{Fin}(C)), D) \xrightarrow{\sim} \text{Fun}^{\mathbb{I}, -}(\text{Span}(\text{Fin}) \times C, D),$$

with inverse given by left Kan extension along ι .

Proof. Step 1: We start with some auxiliary diagrams. Consider the following commutative diagram:

$$\begin{array}{ccccc}
 C & \xrightarrow{m} & \text{Fin} \times C & \xrightarrow{l} & \text{Span}(\text{Fin}) \times C \\
 & \searrow i & \downarrow \iota' & & \downarrow \iota \\
 & & \text{Fin}(C) & \xrightarrow{k} & \text{Span}_{\text{ct,all}}(\text{Fin}(C)).
 \end{array}$$

In particular, passing to restriction functors induces a commutative diagram as follows:

$$\begin{array}{ccccc}
 \text{Fun}(C, D) & \xleftarrow[m^*]{\simeq} & \text{Fun}^{\text{II},-}(\text{Fin} \times C, D) & \xleftarrow[l^*]{} & \text{Fun}^{\text{II},-}(\text{Span}(\text{Fin}) \times C, D) \\
 & \nwarrow i^* & \uparrow \iota'^* & & \uparrow \iota^* \\
 & & \text{Fun}^{\text{II}}(\text{Fin}(C), D) & \xleftarrow[k^*]{} & \text{Fun}^{\text{II}}(\text{Span}_{\text{ct,all}}(\text{Fin}(C)), D).
 \end{array}$$

The functor m^* is an equivalence, since Fin is the free ∞ -category with finite coproducts. The functor i^* was shown to be an equivalence in Lemma 14.1.16. It follows by 2-out-of-3 that ι'^* is an equivalence as well. Since the inverses to m^* and i^* are given by left Kan extension along m and i , the inverse to ι'^* is given by left Kan extension along ι' .

Step 2: We now prove the following auxiliary statement: for every object $X = \{x_i\}_{i \in I}$ of $\text{Fin}(C)$, the relative slice category of ι' over X embeds fully faithfully into the relative slice category of ι over X , and this inclusion admits a left adjoint. To see this, let us denote these two relative slice categories as follows:

$$\begin{aligned}
 P_X &:= \text{Span}_{\text{ct,all}}(\text{Fin}(C))_{/X} \times_{\text{Span}_{\text{ct,all}}(\text{Fin}(C))} (\text{Span}(\text{Fin}) \times C); \\
 Q_X &:= \text{Fin}(C)_{/X} \times_{\text{Fin}(C)} (\text{Fin} \times C).
 \end{aligned}$$

The inclusions l and k induce a fully faithful inclusion $Q_X \hookrightarrow P_X$. We claim that this inclusion admits a left adjoint $P_X \rightarrow Q_X$. To this end, note that an object of P_X consists of a pair (J, y) and a span

$$\{y\}_{j \in J} \xleftarrow{(f, (\text{id}_y))} \{y\}_{k \in K} \xrightarrow{(g, (\varphi_k: y \rightarrow x_i))} \{x_i\}_{i \in I} = X$$

in $\text{Fin}(C)$, where the left-pointing map is q -cartesian. We will send this to the object of Q_X given by the morphism

$$\{y\}_{k \in K} \xrightarrow{(g, (\varphi_k: y \rightarrow x_i))} \{x_i\}_{i \in I} = X.$$

If we include this back into P_X by taking the left-pointing map to be the identity, the span $J \xleftarrow{f} K \xrightarrow{g} K$ and the identity of y define a morphism from the original object to the resulting object. Let $L: P_X \rightarrow Q_X$ denote the construction just described and let $j: Q_X \hookrightarrow P_X$ denote the inclusion. Composition with this morphism induces natural equivalences

$$\text{Hom}_{Q_X}(L(p), q) \simeq \text{Hom}_{P_X}(p, j(q)).$$

Indeed, a morphism on the right is uniquely determined by its map from the middle object of the left leg, because that leg is q -cartesian. Thus L is left adjoint to j .

Step 3: We now return to the question of interest by showing that ι^* admits a left adjoint given by left Kan extension. Consider a functor $F: \text{Span}(\text{Fin}) \times C \rightarrow D$ which preserves finite coproducts in the first variable. We need to show that the left Kan extension $\iota_! F: \text{Span}_{\text{ct,all}}(\text{Fin}(C)) \rightarrow D$ exists

and preserves finite coproducts. For the former, we may use the pointwise criterion for left Kan extensions and show that for each object $\{x_i\}_{i \in I}$ of $\text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C))$ the colimit

$$(\iota_! F)(\{x_i\}_{i \in I}) = \text{colim}_{(J, y) \in P_X} F(J, y)$$

exists in D , where P_X denotes the relative slice category of ι considered in Step 2. Since the inclusion $Q_X \hookrightarrow P_X$ is a right adjoint, it is in particular final, hence the above colimit exists in D if and only if so does $\text{colim}_{(J, y) \in Q_X} F(J, y)$. But this is indeed the case, since Q_X is the relative slice category of ι' and the pointwise left Kan extension of $l^* F: \text{Fin} \times C \rightarrow D$ along ι' exists by Step 1.

We conclude that $\iota_! F$ exists, and that the canonical map $\iota'_! l^* F \rightarrow k^* \iota_! F$ is a natural isomorphism. It remains to show that $\iota_! F$ preserves finite coproducts, but this can be checked after restricting along the inclusion $k: \text{Fin}(C) \hookrightarrow \text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C))$, and thus follows from the fact that $\iota'_! l^* F$ preserves finite coproducts.

Step 4: Finally, we show that the resulting adjunction

$$\iota_!: \text{Fun}^{\text{II}, -}(\text{Span}(\text{Fin}) \times C, D) \rightleftarrows \text{Fun}^{\text{II}}(\text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C)), D) : \iota^*$$

is an adjoint equivalence. Since $\text{Fin}(C)$ is generated under coproducts by the image of $\text{Fin} \times C$, the functor ι^* is conservative, hence it suffices to show that for every $F \in \text{Fun}^{\text{II}, -}(\text{Span}(\text{Fin}) \times C, D)$ the unit $F \rightarrow \iota^* \iota_! F$ is an isomorphism. By essential surjectivity of l , this in turn reduces to $l^* F \xrightarrow{\sim} l^* \iota^* \iota_! F$. Under the isomorphisms $l^* \iota^* \iota_! F = \iota'^* k^* \iota_! F \simeq \iota'^* \iota'_! l^* F$, this corresponds to the unit of the adjunction $\iota'_! \dashv \iota'^*$, hence is an isomorphism by Step 1. $\square$

We may now deduce the characterization of algebras over cocartesian operads we are after:

Proposition 15.2.13. *Let C be an ∞ -category.*

(1) *If E is an ∞ -category with finite products, restriction along ι induces an equivalence*

$$\text{Fun}^\times(C^{\text{II}}, E) \xrightarrow{\sim} \text{Fun}(C, \text{CMon}(E)).$$

(2) *If O is an ∞ -operad, we obtain an equivalence*

$$\text{Fun}_{\text{Op}_\infty}(O_C^{\text{II}}, O) \simeq \text{Fun}(C, \text{CAlg}(O)).$$

(3) *If D is a symmetric monoidal ∞ -category, we obtain an equivalence*

$$\text{Alg}_{O_C^{\text{II}}}(D) \simeq \text{Fun}(C, \text{CAlg}(D)).$$

Proof. Part (3) is an instance of (2) by taking $O = \mathcal{M}_D$ to be the multimorphism operad of D . Part (2) follows by applying Part (1) to $E = O^\otimes$ and $E = \text{Span}(\text{Fin})$, and passing to horizontal fibers:

$$\begin{array}{ccccc} \text{Fun}_{\text{Op}_\infty}(O_C^{\text{II}}, O) & \longrightarrow & \text{Fun}^\times(C^{\text{II}}, O^\otimes) & \xrightarrow{p_O \circ -} & \text{Fun}^\times(C^{\text{II}}, \text{Span}(\text{Fin})) \\ \downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\ \text{Fun}(C, \text{CAlg}(O)) & \longrightarrow & \text{Fun}(C, \text{CMon}(O^\otimes)) & \xrightarrow{p_O \circ -} & \text{Fun}(C, \text{CMon}(\text{Span}(\text{Fin}))). \end{array}$$

It thus remains to prove part (1). For this, recall from Construction 15.2.4 that C^Π is semiadditive. The left vertical map in the following commutative diagram is an equivalence by Corollary 5.3.23. The right vertical map is an equivalence by Proposition 5.3.16, since $\mathbf{CMon}(E)$ is semiadditive:

$$\begin{array}{ccc} \mathrm{Fun}^\times(C^\Pi, \mathbf{CMon}(E)) & \longrightarrow & \mathrm{Fun}(C, \mathbf{CMon}(\mathbf{CMon}(E))) \\ \cong \downarrow & & \downarrow \cong \\ \mathrm{Fun}^\times(C^\Pi, E) & \longrightarrow & \mathrm{Fun}(C, \mathbf{CMon}(E)). \end{array}$$

Showing that the bottom functor in this square is an equivalence thus reduces to showing that the top functor is an equivalence. Since a functor between semiadditive ∞ -categories preserves finite products if and only if it preserves finite coproducts, this top functor may be rewritten as the restriction functor

$$\iota^*: \mathrm{Fun}^\Pi(C^\Pi, \mathbf{CMon}(E)) \rightarrow \mathrm{Fun}^{\Pi, -}(\mathrm{Span}(\mathrm{Fin}) \times C, \mathbf{CMon}(E)) \simeq \mathrm{Fun}(C, \mathrm{Fun}^\Pi(\mathrm{Span}(\mathrm{Fin}), \mathbf{CMon}(E))).$$

This is an equivalence by Lemma 15.2.12. $\square$

15.2.3 Classification of cocartesian operads

As a consequence of the results from the previous subsection, we obtain an explicit classification of ∞ -operads equivalent to O_C^Π for some ∞ -category C .

Definition 15.2.14. An ∞ -operad O is called *cocartesian* if there exists an equivalence $O \simeq O_C^\Pi$ for some ∞ -category C . We denote by $\mathrm{Op}_\infty^{\mathrm{cocart}} \subseteq \mathrm{Op}_\infty$ the full subcategory spanned by the cocartesian ∞ -operads.

Example 15.2.15. The commutative operad *Comm* is cocartesian: it is equivalent to O_{pt}^Π .

Proposition 15.2.16. Let O be a cocartesian ∞ -operad and let $\mathcal{P}$ be an ∞ -operad such that the ∞ -category $\mathcal{P}^\otimes$ is semiadditive. Then the underlying category functor induces an equivalence

$$\mathrm{Fun}_{\mathrm{Op}_\infty}(O, \mathcal{P}) \xrightarrow{\sim} \mathrm{Fun}(O_{\langle 1 \rangle}, \mathcal{P}_{\langle 1 \rangle})$$

between operad maps $O \rightarrow \mathcal{P}$ and functors $O_{\langle 1 \rangle} \rightarrow \mathcal{P}_{\langle 1 \rangle}$.

Proof. Picking an equivalence $O \simeq O_C^\Pi$ for some C , we may identify $O_{\langle 1 \rangle}$ with C . The inclusion $O_{\langle 1 \rangle} \hookrightarrow O^\otimes$ then corresponds to the composite inclusion $C \hookrightarrow \mathrm{Span}(\mathrm{Fin}) \times C \xrightarrow{\iota} C^\Pi$, where ι is the inclusion from Construction 15.2.10. The functor in question then factors as

$$\mathrm{Fun}_{\mathrm{Op}_\infty}(O_C^\Pi, \mathcal{P}) \xrightarrow{\sim} \mathrm{Fun}(C, \mathrm{CAlg}(\mathcal{P})) \rightarrow \mathrm{Fun}(C, \mathcal{P}_{\langle 1 \rangle}),$$

where the first functor is the equivalence from Proposition 15.2.13(2) and the second functor is induced by $\mathrm{CAlg}(\mathcal{P}) \rightarrow \mathcal{P}_{\langle 1 \rangle}$ given by evaluation at $\langle 1 \rangle \in \mathrm{Span}(\mathrm{Fin})$. It thus remains to show that the latter is an equivalence. Unwinding definitions, we see that this map sits as the left vertical map in a diagram of fiber sequences in Cat_∞ as follows:

$$\begin{array}{ccccc} \mathrm{CAlg}(\mathcal{P}) & \hookrightarrow & \mathbf{CMon}(\mathcal{P}^\otimes) & \xrightarrow{(p\mathcal{P})^\sharp} & \mathbf{CMon}(\mathrm{Span}(\mathrm{Fin})) \\ \downarrow \mathrm{ev}_{\langle 1 \rangle} & & \simeq \downarrow \mathrm{ev}_{\langle 1 \rangle} & & \simeq \downarrow \mathrm{ev}_{\langle 1 \rangle} \\ \mathcal{P}_{\langle 1 \rangle} & \hookrightarrow & \mathcal{P}^\otimes & \xrightarrow{p\mathcal{P}} & \mathrm{Span}(\mathrm{Fin}). \end{array}$$

Since $\mathcal{P}^\otimes$ and $\text{Span}(\text{Fin})$ are semiadditive, Proposition 5.3.16 shows that the middle and right vertical maps are equivalences, and the claim follows. $\square$

Corollary 15.2.17. *The functor $O^\Pi: \text{Cat}_\infty \rightarrow \text{Op}_\infty$ is fully faithful and induces an equivalence*

$$\text{Cat}_\infty \xrightarrow{\sim} \text{Op}_\infty^{\text{cocart}}.$$

Proof. Observe that $O^\Pi: \text{Cat}_\infty \rightarrow \text{Op}_\infty$ admits a retraction, given by $(-)_\langle 1 \rangle: \text{Op}_\infty \rightarrow \text{Cat}_\infty$. It follows that for two ∞ -categories C and D , the composite

$$\text{Hom}_{\text{Cat}_\infty}(C, D) \rightarrow \text{Hom}_{\text{Op}_\infty}(O_C^\Pi, O_D^\Pi) \rightarrow \text{Hom}_{\text{Cat}_\infty}(C, D)$$

is the identity. Since the second map is an equivalence by Proposition 15.2.16, the first map is an equivalence as well, showing full faithfulness. Since $\text{Op}_\infty^{\text{cocart}}$ is by definition the image of O^Π , the second claim follows immediately. $\square$

Corollary 15.2.18. *The following conditions are equivalent for an ∞ -operad O :*

- (1) *The ∞ -operad O is cocartesian;*
- (2) *The ∞ -category $O^\otimes$ is semiadditive;*
- (3) *The following two conditions are satisfied:*
 - (a) *For every color $x \in O^\simeq$, the anima $O(\emptyset; x)$ of nullary operations is contractible;*
 - (b) *For objects $\{x_i\}_{i \in I}, \{y_j\}_{j \in J}$ of $O^\otimes$ and a color $z \in O^\simeq$, composition with the morphisms $\{x_i\}_{i \in I} \rightarrow \{x_i\}_{i \in I} \sqcup \{y_j\}_{j \in J}$ and $\{y_j\}_{j \in J} \rightarrow \{x_i\}_{i \in I} \sqcup \{y_j\}_{j \in J}$ obtained from condition (a) provides an equivalence*

$$O(\{x_i\}_{i \in I} \sqcup \{y_j\}_{j \in J}; z) \simeq O(\{x_i\}_{i \in I}; z) \times O(\{y_j\}_{j \in J}; z).$$

Proof. The equivalence between (2) and (3) is a simple unwinding of definitions: condition (a) says that the terminal object of $O^\otimes$ is also initial, and condition (b) says that the maps $X \simeq X \times * \rightarrow X \times Y$ and $Y \simeq * \times Y \rightarrow X \times Y$ in $O^\otimes$ exhibit the product also as a coproduct.

Since $C^\Pi = \text{Span}_{\text{ct}, \text{all}}(\text{Fin}(C))$ is semiadditive, it is clear that (1) implies (2). To see that (2) implies (1), let O be such an ∞ -operad, and write $C := O_{\langle 1 \rangle}^\otimes$ for its underlying ∞ -category. By Proposition 15.2.16, the identity $\text{id}_C: C \rightarrow O_{\langle 1 \rangle}$ extends to a morphism of ∞ -operads $O_C^\Pi \rightarrow O$. It is clear that it induces an equivalence on colors, so it remains to show that it induces equivalences on all multimorphism animae:

$$O_C^\Pi(\{x_i\}_{i \in I}; y) \simeq O(\{x_i\}_{i \in I}; y).$$

But using condition (3), we may inductively reduce this to the case of the one-point set $I = \langle 1 \rangle$, where it is simply the equality $C = O_{\langle 1 \rangle}^\otimes$. The resulting fully faithful operad map is an equivalence by Lemma 14.1.6. $\square$

15.2.4 Classification of cocartesian monoidal structures

We now determine the morphisms between cocartesian monoidal structures and deduce their classification from that of cocartesian operads.

Lemma 15.2.19. *Let C and D be ∞ -categories with finite coproducts and let $F: C \rightarrow D$ be a functor. Then F preserves finite coproducts if and only if the induced map $\text{Span}(F): \text{Span}_{\text{ct,all}}(\text{Fin}(C)) \rightarrow \text{Span}_{\text{ct,all}}(\text{Fin}(D))$ preserves cocartesian morphisms over $\text{Span}(\text{Fin})$.*

Proof. The map $\text{Span}(F)$ is always a morphism of ∞ -operads, so it remains to check cocartesian morphisms over forward spans. Under the unfurling description from Propositions 15.2.1 and 15.2.8, their restrictions along $\text{Fin} \hookrightarrow \text{Span}(\text{Fin})$ are the cocartesian morphisms in $\text{Fin}(C)$ and $\text{Fin}(D)$ which form coproducts over the fibers of a map of finite sets. The functor $\text{Fin}(F)$ preserves these morphisms if and only if F preserves finite coproducts. $\square$

Definition 15.2.20. Let $\text{Cat}_{\infty}^{\otimes, \text{cocart}} \subseteq \text{Cat}_{\infty}^{\otimes}$ denote the full subcategory spanned by the symmetric monoidal ∞ -categories of the form $(C, \amalg)$ for some ∞ -category C . We say that a symmetric monoidal ∞ -category $(D, \otimes)$ is *cocartesian monoidal* if it is equivalent to $(C, \amalg)$ for some ∞ -category C with finite coproducts.

Corollary 15.2.21. *The equivalence $O^{\amalg}: \text{Cat}_{\infty} \xrightarrow{\sim} \text{Op}_{\infty}^{\text{cocart}}$ from Corollary 15.2.17 restricts to an equivalence*

$$\text{Cat}_{\infty}^{\text{coprod}} \xrightarrow{\sim} \text{Cat}_{\infty}^{\otimes, \text{cocart}}.$$

Proof. We need to check that the equivalence from Corollary 15.2.17 restricts to objects and morphisms. On objects this is the content of Proposition 15.2.8. On morphisms this is the content of Lemma 15.2.19. $\square$

Lemma 15.2.22. *A symmetric monoidal ∞ -category $(D, \otimes)$ is cocartesian monoidal if and only if the following two conditions are satisfied:*

- (1) *The monoidal unit $\mathbb{1} \in D$ is an initial object;*
- (2) *For any two objects X and Y of D , the two maps $X \simeq X \otimes \mathbb{1} \rightarrow X \otimes Y$ and $Y \simeq \mathbb{1} \otimes Y \rightarrow X \otimes Y$ induced by the maps $\mathbb{1} \rightarrow X$ and $\mathbb{1} \rightarrow Y$ from (1) exhibit the tensor product $X \otimes Y$ as a coproduct of X and Y .*

Proof. In light of the equivalence

$$\mathcal{M}_D(\{x_i\}_{i \in I}; y) \simeq \text{Hom}_D\left(\bigotimes_{i \in I} x_i, y\right),$$

this is an immediate consequence of Corollary 15.2.18, together with Proposition 15.2.8. $\square$

15.3 Cartesian monoidal structures

The previous section constructed cocartesian monoidal structures, and we now do the same for cartesian ones. Existence comes for free by duality: an ∞ -category with finite products carries a

symmetric monoidal structure whose tensor product is the product and whose unit is the terminal object, obtained from the same unfurling construction by extending along right adjoints rather than left ones. The characterization of cartesian monoidal structures and their uniqueness dualize as well. The description of the algebras does not, and proving it is what most of this section is for.

Let C be an ∞ -category with finite products. The functor $C^{(-)}: \text{Fin}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ is now right adjointable: restriction along $f: I \rightarrow J$ admits a right adjoint f_* , computed by taking products over the fibers of f . The Beck–Chevalley isomorphisms again follow by reindexing fiberwise products along pullback squares.

Proposition 15.3.1. *Let C be an ∞ -category with finite products. Right-adjoint unfurling extends $C^{(-)}$ to a finite-product-preserving functor*

$$(C, \times) := \text{Unf}^R(C^{(-)}): \text{Span}(\text{Fin}) \longrightarrow \text{Cat}_{\infty}$$

*which sends a span $I \xleftarrow{f} K \xrightarrow{g} J$ to g_*f^* . It defines a symmetric monoidal structure on C whose unit is the terminal object and whose tensor product is the cartesian product.*

Proof. Apply Corollary 15.1.5 to $C^{(-)}$. The resulting transport sends an I -indexed family to

$$(x_i)_{i \in I} \mapsto \left(\prod_{k \in g^{-1}(j)} x_{f(k)} \right)_{j \in J}.$$

The argument of Proposition 15.2.1 applies verbatim to show that the construction preserves finite products: the inclusion $\text{Fin}^{\text{op}} \hookrightarrow \text{Span}(\text{Fin})$ preserves finite products and is the identity on objects, so product preservation may be tested on the original functor $C^{(-)}$. The empty product gives the terminal object as unit, and transport along $\langle 2 \rangle \xleftarrow{=} \langle 2 \rangle \xrightarrow{\nabla} \langle 1 \rangle$ gives the binary product. The inert projections in $C^{\times}$ induce in the fiber C the two projection maps from this product. $\square$

Definition 15.3.2 (Cartesian monoidal structure). The symmetric monoidal ∞ -category $(C, \times)$ from Proposition 15.3.1 is called the *cartesian monoidal structure on C* . We write $\mathcal{O}_C^{\times} := \mathcal{M}_{(C, \times)}$ for its underlying ∞ -operad and $C^{\times} := (\mathcal{O}_C^{\times})^{\otimes}$ for its total category.

We first use duality to characterize cartesian monoidal structures and prove their uniqueness. Afterwards we construct a concrete model for $C^{\times}$. That model is not needed for existence, but it gives the explicit control required to prove the universal property of the cartesian operad.

15.3.1 Cartesian monoidal categories

The preceding construction gives a distinguished symmetric monoidal structure on every ∞ -category with finite products. We next compare this construction with the intrinsic notion of a cartesian monoidal ∞ -category.

Definition 15.3.3. A symmetric monoidal ∞ -category $(D, \otimes)$ is called *cartesian monoidal* if the opposite symmetric monoidal ∞ -category $(D, \otimes)^{\text{op}}$ is cocartesian monoidal. We denote by

$$\text{Cat}_{\infty}^{\otimes, \text{cart}} \subseteq \text{Cat}_{\infty}^{\otimes}$$

the full subcategory spanned by the cartesian monoidal ∞ -categories.

Lemma 15.3.4. *A symmetric monoidal ∞ -category $(D, \otimes)$ is cartesian monoidal if and only if the following two conditions are satisfied:*

- (1) *The monoidal unit $\mathbb{1} \in D$ is a terminal object;*
- (2) *For any two objects X and Y of D , the two maps $X \otimes Y \rightarrow X \otimes \mathbb{1} \simeq X$ and $X \otimes Y \rightarrow \mathbb{1} \otimes Y \simeq Y$ induced by the maps $X \rightarrow \mathbb{1}$ and $Y \rightarrow \mathbb{1}$ from the first condition exhibit the tensor product $X \otimes Y$ as a product of X and Y .*

Proof. This is an instance of Lemma 15.2.22 applied to the opposite symmetric monoidal ∞ -category. $\square$

The characterization shows that the underlying ∞ -category of a cartesian monoidal ∞ -category admits finite products. Moreover, every symmetric monoidal functor between cartesian monoidal ∞ -categories preserves these finite products. Hence there is a forgetful functor

$$U: \text{Cat}_{\infty}^{\otimes, \text{cart}} \rightarrow \text{Cat}_{\infty}^{\text{prod}}.$$

Proposition 15.3.5. *Let C be an ∞ -category with finite products. Then the cartesian monoidal structure $(C, \times)$ is cartesian monoidal.*

Proof. The transport formula in Proposition 15.3.1 shows that the monoidal unit is the terminal object of C and the binary tensor product is the product in C . Under the identification of the fiber over $\langle 1 \rangle$ with C , the two inert projections are the two maps from this product induced by the terminal maps of its factors, hence are precisely the product projections. Thus $(C, \times)$ satisfies the two conditions of Lemma 15.3.4. $\square$

Proposition 15.3.6. *The forgetful functor*

$$U: \text{Cat}_{\infty}^{\otimes, \text{cart}} \rightarrow \text{Cat}_{\infty}^{\text{prod}}$$

is an equivalence, with inverse given by the assignment $C \mapsto (C, \times)$.

Proof. Passing to opposites identifies $\text{Cat}_{\infty}^{\otimes, \text{cart}}$ with $\text{Cat}_{\infty}^{\otimes, \text{cocart}}$, and identifies the forgetful functor U with the corresponding forgetful functor

$$\text{Cat}_{\infty}^{\otimes, \text{cocart}} \rightarrow \text{Cat}_{\infty}^{\text{coprod}}.$$

The latter is an equivalence by Corollary 15.2.21: the functor $C \mapsto (C, \amalg)$ is an equivalence whose composite with the forgetful functor is the identity on $\text{Cat}_{\infty}^{\text{coprod}}$. Hence U is an equivalence as well.

By Proposition 15.3.5, the assignment $C \mapsto (C, \times)$ lands in $\text{Cat}_{\infty}^{\otimes, \text{cart}}$. Full faithfulness of U uniquely lifts every finite-product-preserving functor $C \rightarrow D$ to a symmetric monoidal functor $(C, \times) \rightarrow (D, \times)$. These lifts define a functor whose composite with U is the identity, and hence an inverse to U . $\square$

15.3.2 A concrete model for the cartesian operad

The unfurling construction gives the cartesian monoidal structure directly from the functor $I \mapsto C^I$. To prove its universal property, it is useful to replace an I -indexed family by a functor that also records all products of its subfamilies coherently. This leads to the following model based on the span-functoriality of $I \mapsto \text{Fin}_I$.

Construction 15.3.7. The target projection $\text{ev}_1 : \text{Ar}(\text{Fin}) \rightarrow \text{Fin}$ is a cartesian fibration whose cartesian straightening sends I to Fin_I . Thus pullback defines a functor $\text{Fin}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ with value Fin_I at I . Each pullback functor $f^* : \text{Fin}_J \rightarrow \text{Fin}_I$ has a left adjoint $f_!$, given by postcomposition with f . The Beck–Chevalley isomorphisms follow from the pasting law for pullback squares. Its unfurling therefore sends a span $I \xleftarrow{f} K \xrightarrow{g} J$ to $g_! f^* : \text{Fin}_I \rightarrow \text{Fin}_J$.

Precomposing this unfurling with the leg-swap equivalence $\text{Span}(\text{Fin}) \simeq \text{Span}(\text{Fin})^{\text{op}}$ from Lemma 13.1.15 and then passing to opposites gives a functor $\text{Span}(\text{Fin})^{\text{op}} \rightarrow \text{Cat}_{\infty}$. Let

$$r : A \rightarrow \text{Span}(\text{Fin})$$

be its cartesian unstraightening. Its fiber over I is $(\text{Fin}_I)^{\text{op}}$, and its cartesian transport along a span $I \xleftarrow{f} K \xrightarrow{g} J$ is $(f_! g^*)^{\text{op}}$.

Applying $B \mapsto \text{Fun}(B, C)$ to the straightening of r gives a functor $\text{Span}(\text{Fin}) \rightarrow \text{Cat}_{\infty}$. It sends I to $\text{Fun}((\text{Fin}_I)^{\text{op}}, C)$ and a span $I \xleftarrow{f} K \xrightarrow{g} J$ to restriction along

$$(f_! g^*)^{\text{op}} : (\text{Fin}_J)^{\text{op}} \longrightarrow (\text{Fin}_I)^{\text{op}}.$$

Let $\widetilde{C}^{\times} \rightarrow \text{Span}(\text{Fin})$ be its cocartesian unstraightening, and let $C_{\text{con}}^{\times} \subseteq \widetilde{C}^{\times}$ be the fiberwise full subcategory whose fiber over I consists of the finite-product-preserving functors $(\text{Fin}_I)^{\text{op}} \rightarrow C$. The subscript con indicates that this is the concrete model.

Lemma 15.3.8. *The subcategory $C_{\text{con}}^{\times} \subseteq \widetilde{C}^{\times}$ is closed under cocartesian transport. In particular, $C_{\text{con}}^{\times} \rightarrow \text{Span}(\text{Fin})$ is a cocartesian fibration.*

Proof. For a span $I \xleftarrow{f} K \xrightarrow{g} J$, the functor $f_! g^* : \text{Fin}_J \rightarrow \text{Fin}_I$ preserves finite coproducts: pullback along g does so because Fin is extensive, while postcomposition along f plainly does so. Hence $(f_! g^*)^{\text{op}}$ preserves finite products, and restriction along it preserves finite-product-preserving functors. $\square$

Lemma 15.3.9. *For every finite set I , restriction along the functor $I \hookrightarrow (\text{Fin}_I)^{\text{op}}$ given by $i \mapsto (\{i\} \hookrightarrow I)$ induces an equivalence*

$$\text{Fun}^{\times}((\text{Fin}_I)^{\text{op}}, C) \xrightarrow{\sim} C^I.$$

Proof. Regarding I as a discrete ∞ -category, the description of finite families in Construction 14.1.15 gives a finite-coproduct-preserving equivalence

$$\text{Fin}(I) \xrightarrow{\sim} \text{Fin}_I$$

which sends a finite family of elements of I to its indexing map to I . The dual of Lemma 14.1.16 therefore identifies restriction to the singleton families with an equivalence

$$\text{Fun}^{\times}((\text{Fin}_I)^{\text{op}}, C) \simeq \text{Fun}(I, C) = C^I.$$

Under this equivalence, a functor F corresponds to its values on the singleton inclusions, and one has

$$F(U \rightarrow I) \simeq \prod_{u \in U} F(\{u\} \rightarrow I). \quad \square$$

Proposition 15.3.10. *There is a canonical equivalence over $\text{Span}(\text{Fin})$ between the concrete model $C_{\text{con}}^\times$ and the total category $C^\times$ of the cartesian monoidal structure.*

Proof. By the uniqueness of cartesian monoidal structures from Proposition 15.3.6, it suffices to show that $C_{\text{con}}^\times$ defines a cartesian monoidal structure on C .

By Lemma 15.3.8, the projection $C_{\text{con}}^\times \rightarrow \text{Span}(\text{Fin})$ is a cocartesian fibration. Under the fiber equivalences of Lemma 15.3.9, the product-comparison maps of its straightening are the canonical equivalences

$$C^{I \sqcup J} \simeq C^I \times C^J,$$

and its value at $\emptyset$ is the terminal category. Thus the straightening preserves finite products, so Lemma 14.2.4 makes $C_{\text{con}}^\times$ into a symmetric monoidal ∞ -category with underlying ∞ -category C .

Transport along a span $I \xleftarrow{f} K \xrightarrow{g} J$ sends an I -indexed family to

$$(x_i)_{i \in I} \mapsto \left(\prod_{k \in g^{-1}(j)} x_{f(k)} \right)_{j \in J}.$$

Hence its unit is terminal, its binary tensor product is the product in C , and its inert structure maps are the product projections. By Lemma 15.3.4, it is therefore cartesian monoidal. $\square$

15.3.3 Algebras in cartesian monoidal structures

The classification of cartesian monoidal structures followed from the cocartesian classification by passing to opposite symmetric monoidal ∞ -categories. The corresponding statement about algebras does not follow by the same argument: passing to the opposite of the total category does not define an involution of Op_∞ that turns maps out of a cocartesian operad into maps into a cartesian operad. We therefore prove the cartesian universal property directly.

Let C be an ∞ -category with finite products, and let $\mathcal{O}$ be an ∞ -operad. We have previously introduced two a priori different algebraic structures associated to $\mathcal{O}$:

- We may consider $\mathcal{O}$ -monoids in C : finite-product-preserving functors $M: \mathcal{O}^\otimes \rightarrow C$;
- We may also consider $\mathcal{O}$ -algebras in C with respect to the cartesian monoidal structure $(C, \times)$: operad maps $M: \mathcal{O} \rightarrow \mathcal{M}_{(C, \times)}$.

At an informal level, these two notions encode precisely the same structure: an object M_x for every color $x \in \mathcal{O}^\simeq$ and a multimorphism

$$P_M: M_{x_1} \times M_{x_2} \times \cdots \times M_{x_n} \rightarrow M_y$$

for every multimorphism $P \in \mathcal{O}((x_1, \dots, x_n); y)$. This makes it plausible that $\mathcal{O}$ -monoids and $\mathcal{O}$ -algebras define equivalent ∞ -categories. We will first prove this equivalence directly; the product functor $C^\times \rightarrow C$ will then be recovered from it by naturality.

Theorem 15.3.11 (Universal property of cartesian monoidal structures). *Let C be an ∞ -category with finite products. For every ∞ -operad O , there is a natural equivalence of ∞ -categories*

$$\mathrm{Alg}_O(C, \times) = \mathrm{Fun}_{\mathrm{Op}_\infty}(O, O_C^\times) \xrightarrow{\sim} \mathrm{Fun}^\times(O^\otimes, C) = \mathrm{Mon}_O(C).$$

In particular, the functor $\mathrm{Op}_\infty \rightarrow \mathrm{Cat}_\infty^{\mathrm{prod}}, O \mapsto O^\otimes$ is left adjoint to the cartesian operad functor $O^\times: \mathrm{Cat}_\infty^{\mathrm{prod}} \rightarrow \mathrm{Op}_\infty$.

Proof. We use the concrete model of the cartesian operad. Evaluation at the identity maps id_I will send an operad map $O \rightarrow O_C^\times$ to an O -monoid, and restriction along inert morphisms will provide the inverse construction.

Let $r: A \rightarrow \mathrm{Span}(\mathrm{Fin})$ be the cartesian fibration from Construction 15.3.7. For an ∞ -operad O , put

$$D_O := O^\otimes \times_{\mathrm{Span}(\mathrm{Fin})} A.$$

The fiber of the projection $D_O \rightarrow O^\otimes$ over $X \in O_I^\otimes$ is $(\mathrm{Fin}/I)^{\mathrm{op}}$, whose initial object is id_I . By Proposition 23.3.4, these initial objects determine a unique left-adjoint section

$$s: O^\otimes \rightarrow D_O, \quad X \mapsto (X, \mathrm{id}_I).$$

The cocartesian fibration $\tilde{C}^\times \rightarrow \mathrm{Span}(\mathrm{Fin})$ of Construction 15.3.7 is the parametrized functor category associated to r . Hence Theorem 23.8.3 gives an equivalence

$$\mathrm{Fun}_{/\mathrm{Span}(\mathrm{Fin})}(O^\otimes, \tilde{C}^\times) \xrightarrow{\sim} \mathrm{Fun}(D_O, C). \quad (15.2)$$

It sends Φ to the evaluation functor

$$G_\Phi(X, u: U \rightarrow I) := \Phi(X)(u).$$

By construction of r , an object $u: U \rightarrow I$ of its fiber over I indexes the backwards span

$$I \xleftarrow{u} U \xrightarrow{=} U.$$

The functoriality of partial cocartesian lifts from Proposition 23.1.4 therefore gives, naturally in O , a functor

$$\theta_O: D_O \rightarrow O^\otimes, \quad (X, u: U \rightarrow I) \mapsto X_U,$$

and a natural transformation

$$\delta: \mathrm{id}_{D_O} \longrightarrow s\theta_O$$

whose component at (X, u) has $O^\otimes$ -component the inert lift $X \rightarrow X_U$ of the backwards span above and A -component the cartesian lift from u to id_U over the same span. Restriction along identity maps gives a natural isomorphism $\theta_O s \simeq \mathrm{id}_{O^\otimes}$.

Let $\Phi: O \rightarrow O_C^\times$ be a morphism of ∞ -operads. Regarding its total functor as a functor to the concrete model and applying Equation (15.2), define

$$R(\Phi) := G_\Phi s, \quad R(\Phi)(X) = \Phi(X)(\mathrm{id}_I)$$

for $X \in \mathcal{O}_I^\otimes$. This functor preserves finite products. Indeed, for a finite collection of objects $X_a \in \mathcal{O}_{I_a}^\otimes$, product preservation of Φ and Lemma 15.3.9 give

$$R(\Phi) \left(\prod_a X_a \right) \simeq \left(\prod_a \Phi(X_a) \right) (\text{id}_{\sqcup_a I_a}) \simeq \prod_a \Phi(X_a) (\text{id}_{I_a}) = \prod_a R(\Phi)(X_a).$$

The same formula includes the empty product. We have therefore obtained a functor

$$R: \text{Fun}_{\text{Op}_\infty}(\mathcal{O}, \mathcal{O}_C^\times) \longrightarrow \text{Fun}^\times(\mathcal{O}^\otimes, C).$$

Conversely, let $M: \mathcal{O}^\otimes \rightarrow C$ preserve finite products. The functor

$$M\theta_{\mathcal{O}}: D_{\mathcal{O}} \rightarrow C$$

corresponds under Equation (15.2) to a functor

$$\widehat{M}: \mathcal{O}^\otimes \rightarrow \widetilde{C}^\times.$$

This functor factors through $C_{\text{con}}^\times$. Indeed, fix $X \in \mathcal{O}_I^\otimes$. A finite product in $(\text{Fin}/I)^{\text{op}}$ is represented by a disjoint union $\sqcup_a U_a \rightarrow I$ in Fin/I , and operadic inert restriction gives a natural isomorphism

$$X_{\sqcup_a U_a} \simeq \prod_a X_{U_a}.$$

Consequently,

$$\widehat{M}(X) \left(\sqcup_a U_a \rightarrow I \right) = M(X_{\sqcup_a U_a}) \simeq \prod_a M(X_{U_a}),$$

including the empty product. Thus $\widehat{M}(X): (\text{Fin}/I)^{\text{op}} \rightarrow C$ preserves finite products.

We identify $C_{\text{con}}^\times$ with $C^\times$ using Proposition 15.3.10. The resulting total functor $\widehat{M}: \mathcal{O}^\otimes \rightarrow C^\times$ preserves finite products. Indeed, let $\{X_a\}_{a \in S}$ lie over finite sets $\{I_a\}_{a \in S}$, and let $u: U \rightarrow \sqcup_a I_a$. Writing $U_a := U \times_{\sqcup_a I_a} I_a$, compatibility of inert restriction with products gives

$$\left(\prod_a X_a \right)_U \simeq \prod_a (X_a)_{U_a}.$$

After applying M , these isomorphisms identify, at every u , the product-comparison map for $\widehat{M}$ with an isomorphism. They are natural in u by the coherence of $\theta_{\mathcal{O}}$, and isomorphisms in a functor category are detected pointwise. Thus the comparison is an isomorphism in the relevant functor category. Hence $\widehat{M}$ is a morphism of ∞ -operads. We have constructed a functor

$$\widehat{(-)}: \text{Fun}^\times(\mathcal{O}^\otimes, C) \longrightarrow \text{Fun}_{\text{Op}_\infty}(\mathcal{O}, \mathcal{O}_C^\times).$$

The natural isomorphism $\theta_{\mathcal{O}} s \simeq \text{id}_{\mathcal{O}^\otimes}$ gives, naturally in M ,

$$R(\widehat{M}) = M\theta_{\mathcal{O}} s \simeq M.$$

In the other direction, apply G_Φ to δ . This gives a natural transformation

$$G_\Phi \longrightarrow G_\Phi s \theta_{\mathcal{O}} = R(\Phi) \theta_{\mathcal{O}}.$$

Its component at $(X, u: U \rightarrow I)$ is the canonical map

$$\Phi(X)(u) \longrightarrow \Phi(X_U)(\text{id}_U).$$

It is an isomorphism: the $O^\otimes$ -component of δ is inert, so Φ carries it to an inert, hence cocartesian, morphism in $C^\times$ by Corollary 14.1.10; the concrete transport formula then identifies its target, evaluated at id_U , with its source, evaluated at u . Currying therefore gives a natural isomorphism

$$\Phi \simeq \widehat{R(\Phi)}.$$

Thus R and $\widehat{(-)}$ are inverse equivalences.

The construction is natural in O by pullback and the naturality of θ and δ . It is natural in C because postcomposition with a finite-product-preserving functor commutes with currying and preserves the concrete subcategories $C_{\text{con}}^\times$. Under Proposition 15.3.10, the resulting map between the concrete models is the morphism of cartesian operads supplied by Proposition 15.3.6. Hence we obtain a natural equivalence of functors

$$\text{Op}_\infty^{\text{op}} \times \text{Cat}_\infty^{\text{prod}} \longrightarrow \text{Cat}_\infty.$$

Passing to groupoid cores gives the natural equivalence of hom animae which exhibits $O \mapsto O^\otimes$ as left adjoint to $O^\times$. $\square$

Corollary 15.3.12. *There is a finite-product-preserving functor*

$$\Pi_C: C^\times \rightarrow C$$

such that the equivalence of Theorem 15.3.11 is given by postcomposition with Π_C .

Proof. Apply Theorem 15.3.11 to $O = O_C^\times$. The identity of $O_C^\times$ corresponds to a finite-product-preserving functor $\Pi_C: C^\times \rightarrow C$. By the naturality in O , the equivalence for a general O is given by postcomposition with this functor.

Unwinding the construction of the equivalence, if $F: (\text{Fin}_I)^{\text{op}} \rightarrow C$ is an object of $C^\times$ over I , then

$$\Pi_C(F) \simeq F(\text{id}_I).$$

Under the equivalence of Lemma 15.3.9, this sends the family $\{x_i\}_{i \in I}$ to $\prod_{i \in I} x_i$. $\square$

Corollary 15.3.13. *Let C be an ∞ -category with finite products. Then associative algebras in the cartesian monoidal structure agree with associative operadic monoids in C :*

$$\text{Alg}(C, \times) \xrightarrow{\sim} \text{Mon}_{\mathcal{A}ssoc}(C).$$

Similarly, commutative algebras agree with commutative monoids:

$$\text{CAlg}(C, \times) \xrightarrow{\sim} \text{Mon}_{\text{Comm}}(C) = \text{CMon}(C).$$

Proof. This is the special case of Theorem 15.3.11 where $O = \mathcal{A}ssoc$ and $O = \text{Comm}$, respectively. $\square$

The first equivalence uses the operadic notation $\text{Mon}_{\mathcal{A}ssoc}(C) = \text{Fun}^\times(\mathcal{A}ssoc^\otimes, C)$ from Definition 14.4.1. Its comparison with the simplicial ∞ -category $\text{Mon}(C)$ of Chapter 5 follows from the simplicial model for associative algebras discussed in Section 19.3.

Exercises

16 Day convolution and the tensor product of spectra

Let C and D be symmetric monoidal ∞ -categories. It is natural to wonder whether the ∞ -category $\mathrm{Fun}(C, D)$ of functors from C to D inherits a symmetric monoidal structure. The *Day convolution*, first introduced by Day [Day71] for classical symmetric monoidal categories, provides an affirmative answer to this question under suitable assumptions on C and D .

Let us first recall the classical construction. Let $(C, \otimes_C, \mathbb{1}_C)$ and $(D, \otimes_D, \mathbb{1}_D)$ be symmetric monoidal 1-categories, and assume that C is small and that D admits all small colimits. For any two functors $F, G: C \rightarrow D$, their *Day convolution* $F \otimes_{\mathrm{Day}} G$ is a functor $C \rightarrow D$ defined as the left Kan extension of the composite functor

$$C \times C \xrightarrow{F \times G} D \times D \xrightarrow{\otimes_D} D$$

along the tensor product functor $\otimes_C: C \times C \rightarrow C$. More explicitly, for an object $X \in C$, the value of $F \otimes_{\mathrm{Day}} G$ is given by the colimit

$$(F \otimes_{\mathrm{Day}} G)(X) = \mathrm{colim}_{X_0 \otimes_C X_1 \rightarrow X} F(X_0) \otimes_D G(X_1),$$

where the colimit is taken over the relative slice category $(\otimes_C)_{/X}$, whose objects are morphisms $X_0 \otimes_C X_1 \rightarrow X$ and whose morphisms are compatible maps of pairs (X_0, X_1) . Two properties of this construction are relevant for us:

- (a) If the tensor product $\otimes_D: D \times D \rightarrow D$ preserves small colimits in each variable separately, then the Day convolution equips the functor category $\mathrm{Fun}(C, D)$ with a symmetric monoidal structure.
- (b) Under this monoidal structure, the category of commutative algebra objects $\mathrm{CAlg}(\mathrm{Fun}(C, D))$ is equivalent to the category of lax symmetric monoidal functors from C to D .

The goal of this chapter is to discuss an ∞ -categorical analogue of Day convolution. A first ∞ -categorical version was constructed by Glasman [Gla16] and subsequently generalized by Lurie [Lur17, Section 2.2.6]. We will prove the form needed here using the Hinich–Winges model for Day convolution between symmetric monoidal ∞ -categories. The description of the Hinich–Winges model in Theorem 16.2.2 is the principal input about Day convolution that we take as a black box from the literature. We use the presentability results of Chapter 22 when constructing reflective subcategories.

Our main application is the symmetric monoidal tensor product on the ∞ -category Sp of spectra, upgrading the bifunctor $- \otimes -: \mathrm{Sp} \times \mathrm{Sp} \rightarrow \mathrm{Sp}$ from Proposition 4.4.16. To this end, we use an equivalent description of Sp as the ∞ -category of certain *reduced excisive functors* from $\mathrm{An}_*^{\mathrm{fin}}$ to An . We can then exhibit Sp as a symmetric monoidal Bousfield localization of a Day convolution category.

Section 16.1 characterizes Day convolution operads by a universal property, and Section 16.2 constructs them for symmetric monoidal ∞ -categories; basic examples follow in Section 16.3. In Section 16.4 we use Day convolution to construct tensor products of commutative monoids and groups, leading to general notions of rings and semirings. We then prove the reduced excisive model of spectra in Section 16.5, and use it in Section 16.6 to equip Sp with its symmetric monoidal structure; that section closes with the multiplicative infinite loop space machine and the lax symmetric monoidal graded homotopy functor.

16.1 Day convolution operads

Lurie’s construction of the Day convolution monoidal structure on $\mathrm{Fun}(C, D)$ has a nice feature, characterizing it as an internal hom in the ∞ -category of ∞ -operads. We will take this universal property as the defining property of Day convolution operads:

Definition 16.1.1. Let O and $\mathcal{P}$ be ∞ -operads. An ∞ -operad $\mathrm{Day}(O, \mathcal{P})$ is called a *Day convolution operad for O and $\mathcal{P}$* if it comes equipped with a morphism of ∞ -operads

$$\mathrm{ev}: \mathrm{Day}(O, \mathcal{P}) \times O \rightarrow \mathcal{P}$$

satisfying the following universal property: if Q is another ∞ -operad, the composite

$$\mathrm{Fun}_{\mathrm{Op}_{\infty}}(Q, \mathrm{Day}(O, \mathcal{P})) \xrightarrow{- \times O} \mathrm{Fun}_{\mathrm{Op}_{\infty}}(Q \times O, \mathrm{Day}(O, \mathcal{P}) \times O) \xrightarrow{\mathrm{ev} \circ -} \mathrm{Fun}_{\mathrm{Op}_{\infty}}(Q \times O, \mathcal{P})$$

is an equivalence. Informally, this means we have the following classification of operad morphisms into $\mathrm{Day}(O, \mathcal{P})$:

$$\{\text{Operad morphisms } Q \rightarrow \mathrm{Day}(O, \mathcal{P})\} \quad \leftrightarrow \quad \{\text{Operad morphisms } Q \times O \rightarrow \mathcal{P}\}.$$

Note that this property uniquely determines $\mathrm{Day}(O, \mathcal{P})$ if it exists.

Remark 16.1.2. Let O be an ∞ -operad for which $\mathrm{Day}(O, \mathcal{P})$ exists for all $\mathcal{P}$. Then the functor $- \times O: \mathrm{Op}_{\infty} \rightarrow \mathrm{Op}_{\infty}$ admits a right adjoint

$$\mathrm{Day}(O, -): \mathrm{Op}_{\infty} \rightarrow \mathrm{Op}_{\infty}.$$

Whenever the relevant Day convolution operads exist, this internal-hom description is functorial in both variables. An operad map $u: O' \rightarrow O$ induces by precomposition an operad map

$$u^*: \mathrm{Day}(O, \mathcal{P}) \longrightarrow \mathrm{Day}(O', \mathcal{P}),$$

and an operad map $v: \mathcal{P} \rightarrow \mathcal{P}'$ induces by postcomposition an operad map

$$v_*: \mathrm{Day}(O, \mathcal{P}) \longrightarrow \mathrm{Day}(O, \mathcal{P}').$$

The first map is adjoint to evaluation after precomposition with $\text{id} \times u$, while the second is adjoint to postcomposition of evaluation with v . Their functoriality follows from the universal property. The same universal property gives a natural exponential law

$$\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{D}\text{ay}(\mathcal{P}, \mathcal{Q})) \simeq \mathcal{D}\text{ay}(\mathcal{O} \times \mathcal{P}, \mathcal{Q}).$$

Indeed, maps from an operad $\mathcal{R}$ into either side are naturally equivalent to maps $\mathcal{R} \times \mathcal{O} \times \mathcal{P} \rightarrow \mathcal{Q}$.

The universal property of $\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})$ allows us to deduce most of its basic structure. Let us first describe its underlying ∞ -category.

Lemma 16.1.3. *Assume $\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})$ exists. Then the underlying ∞ -category of $\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})$ is the ∞ -category of functors $\mathcal{O}_{\langle 1 \rangle} \rightarrow \mathcal{P}_{\langle 1 \rangle}$ between the underlying ∞ -categories of $\mathcal{O}$ and $\mathcal{P}$:*

$$\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})_{\langle 1 \rangle} \simeq \text{Fun}(\mathcal{O}_{\langle 1 \rangle}, \mathcal{P}_{\langle 1 \rangle}).$$

Proof. By Lemma 14.1.19, the universal property of Day convolution, and Lemma 14.1.22, we have natural equivalences

$$\begin{aligned} \mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})_{\langle 1 \rangle} &\simeq \text{Fun}_{\text{Op}_{\infty}}(\text{Triv}, \mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})) \\ &\simeq \text{Fun}_{\text{Op}_{\infty}}(\text{Triv} \times \mathcal{O}, \mathcal{P}) \\ &\simeq \text{Fun}_{\text{Op}_{\infty}}(\text{Triv}_{\mathcal{O}_{\langle 1 \rangle}}, \mathcal{P}) \\ &\simeq \text{Fun}(\mathcal{O}_{\langle 1 \rangle}, \mathcal{P}_{\langle 1 \rangle}), \end{aligned}$$

where the final equivalence is another instance of Lemma 14.1.19. □

As another consequence of the universal property, we may describe the commutative algebras in Day convolution operads:

Lemma 16.1.4. *Assume $\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})$ exists. Then commutative algebras in $\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})$ correspond to operad morphisms $\mathcal{O} \rightarrow \mathcal{P}$:*

$$\text{CAlg}(\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})) \simeq \text{Fun}_{\text{Op}_{\infty}}(\mathcal{O}, \mathcal{P}).$$

Proof. Since Comm is the terminal ∞ -operad, we have $\mathcal{O} \simeq \text{Comm} \times \mathcal{O}$, and so

$$\begin{aligned} \text{CAlg}(\mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})) &= \text{Fun}_{\text{Op}_{\infty}}(\text{Comm}, \mathcal{D}\text{ay}(\mathcal{O}, \mathcal{P})) \\ &\simeq \text{Fun}_{\text{Op}_{\infty}}(\text{Comm} \times \mathcal{O}, \mathcal{P}) \\ &\simeq \text{Fun}_{\text{Op}_{\infty}}(\mathcal{O}, \mathcal{P}). \end{aligned} \quad \square$$

16.2 Day convolution between symmetric monoidal categories

The goal of this section is to construct, under suitable conditions on symmetric monoidal ∞ -categories $\mathcal{C}$ and $\mathcal{D}$, the Day convolution symmetric monoidal structure on $\text{Fun}(\mathcal{C}, \mathcal{D})$. More precisely, we will construct the Day convolution operad $\mathcal{D}\text{ay}(\mathcal{M}_{\mathcal{C}}, \mathcal{M}_{\mathcal{D}})$ and determine when it is the multimorphism operad of a symmetric monoidal ∞ -category. If this operad exists, its universal property already determines two of its fundamental features:

- By Lemma 16.1.3, the underlying ∞ -category of $\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)$ is the ∞ -category of functors from C to D :

$$\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)_{\langle 1 \rangle} \simeq \text{Fun}(C, D).$$

In particular, the colors of $\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)$ may be identified with functors $C \rightarrow D$.

- By Lemma 16.1.4, commutative algebras in $\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)$ are lax symmetric monoidal functors from C to D :

$$\text{CAlg}(\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)) \simeq \text{Fun}^{\otimes\text{-lax}}(C, D).$$

To establish existence and determine when this operad defines a symmetric monoidal structure, we will use the concrete model introduced in the next subsection.

All constructions below are understood in an ambient universe containing the symmetric monoidal ∞ -categories under discussion. If C or D is large relative to the basic universe, we enlarge once as in Remark 1.8.1 and carry out the construction there. Further enlargement does not change the resulting Day convolution operad, since its universal property is expressed entirely in terms of the same operads and their mapping animae.

16.2.1 The Hinich–Winges model

While the universal property of $\mathcal{D}ay(\mathcal{M}_C, \mathcal{M}_D)$ gives us access to some of its abstract properties, it does not provide an explicit description of it. We will make this operad concrete by discussing a model for it given by Hinich [Hin20]. We will follow the ‘synthetic’ exposition of it by Winges [Win25].

The main ingredient for this is an explicit description of the unstraightening of the functor

$$(\text{Cat}_\infty^\otimes)^{\text{op}} \times \text{Cat}_\infty^\otimes \rightarrow \text{Cat}_\infty, \quad (C, D) \mapsto \text{Fun}^{\otimes\text{-lax}}(C, D).$$

Definition 16.2.1. We denote by

$$\text{Ar}^{\text{oplax}} \subseteq (\text{Cat}_\infty)_{/[1]}$$

the full subcategory spanned by the cartesian fibrations $E \rightarrow [1]$.

Thus Ar^{oplax} has the same objects as the ordinary arrow category $\text{Ar}(\text{Cat}_\infty)$, but more morphisms. Under cartesian straightening, an object of Ar^{oplax} corresponds to a functor $[1]^{\text{op}} \rightarrow \text{Cat}_\infty$. We write such a functor as $F: C \rightarrow D$, with C as the fiber over 1 and D as the fiber over 0. Since the morphisms in Ar^{oplax} do not need to preserve the cartesian edges, they correspond to *laxly* commutative squares of the form

$$\begin{array}{ccc} C & \longrightarrow & C' \\ F \downarrow & \swarrow & \downarrow F' \\ D & \longrightarrow & D'. \end{array}$$

Observe that Ar^{oplax} is closed under finite products in $(\text{Cat}_\infty)_{/[1]}$, and that pulling back along the inclusions $\{1\} \hookrightarrow [1]$ and $\{0\} \hookrightarrow [1]$ provides a finite-product-preserving functor

$$(s, t): \text{Ar}^{\text{oplax}} \rightarrow \text{Cat}_\infty \times \text{Cat}_\infty.$$

Theorem 16.2.2 (Winges [Win25, Theorem 2.7]). *Let $\mathcal{O}$ be an ∞ -operad, and consider the induced functor*

$$(s, t): \text{Mon}_{\mathcal{O}}(\text{Ar}^{\text{oplax}}) \rightarrow \text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty}) \times \text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty}).$$

Then:

- (1) *The fiber over a pair (C, D) of $\mathcal{O}$ -monoidal ∞ -categories is the ∞ -category $\text{Fun}^{\mathcal{O}\text{-lax}}(C, D)$ of lax $\mathcal{O}$ -monoidal functors $C \rightarrow D$.*
- (2) *For fixed C , the restriction of (s, t) to $\{C\} \times \text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty})$ is a cocartesian fibration classifying*

$$\text{Fun}^{\mathcal{O}\text{-lax}}(C, -): \text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty}) \rightarrow \text{Cat}_{\infty}.$$

- (3) *For fixed D , the restriction of (s, t) to $\text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty}) \times \{D\}$ is a cartesian fibration classifying*

$$\text{Fun}^{\mathcal{O}\text{-lax}}(-, D): \text{Mon}_{\mathcal{O}}(\text{Cat}_{\infty})^{\text{op}} \rightarrow \text{Cat}_{\infty}.$$

Remark 16.2.3. Winges in fact shows the stronger claim that this identification of the fibers is natural in both C and D simultaneously. Formulating this precisely requires the notion of *orthofibrations*, which are functors of the form $E \rightarrow C \times D$ that correspond under a generalized form of straightening to functors $C^{\text{op}} \times D \rightarrow \text{Cat}_{\infty}$. What Winges shows is that the functor (s, t) from Theorem 16.2.2 is the orthofibration classifying the assignment $(C, D) \mapsto \text{Fun}^{\mathcal{O}\text{-lax}}(C, D)$.

With this result at hand, we may now prove the explicit description of $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_D)$:

Theorem 16.2.4 (Hinich [Hin20, Proposition 2.8.9], Winges [Win25, Theorem 1.1]). *Let C and D be symmetric monoidal ∞ -categories, which we identify with commutative algebras in Cat_{∞} with the cartesian monoidal structure. Then the Day convolution operad $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_D)$ exists, and is given by the following pullback square:*

$$\begin{array}{ccc} \mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_D) & \longrightarrow & \mathcal{O}_{\text{Ar}^{\text{oplax}}}^{\times} \\ \downarrow & \lrcorner & \downarrow (s, t) \\ \text{Comm} & \xrightarrow{(C, D)} & \mathcal{O}_{\text{Cat}_{\infty} \times \text{Cat}_{\infty}}^{\times} \end{array}$$

Proof. Let us write $\mathcal{D}$ for this pullback. We will show directly that $\mathcal{D}$ satisfies the defining universal property of $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_D)$. It will suffice to produce for any ∞ -operad $\mathcal{P}$ a natural equivalence

$$\text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, \mathcal{D}) \simeq \text{Fun}_{\text{Op}_{\infty}}(\mathcal{P} \times \mathcal{M}_C, \mathcal{M}_D).$$

For the left-hand side, note that $\text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, -)$ commutes with pullbacks. Since Comm is the terminal ∞ -operad, we have $\text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, \text{Comm}) \simeq *$. Since operad morphisms from $\mathcal{P}$ into a cartesian operad correspond to $\mathcal{P}$ -monoids by Theorem 15.3.11, we obtain equivalences

$$\text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, \mathcal{O}_{\text{Ar}^{\text{oplax}}}^{\times}) \simeq \text{Mon}_{\mathcal{P}}(\text{Ar}^{\text{oplax}}), \quad \text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, \mathcal{O}_{\text{Cat}_{\infty} \times \text{Cat}_{\infty}}^{\times}) \simeq \text{Mon}_{\mathcal{P}}(\text{Cat}_{\infty} \times \text{Cat}_{\infty}).$$

It follows that the ∞ -category $\text{Fun}_{\text{Op}_{\infty}}(\mathcal{P}, \mathcal{D})$ is equivalent to the fiber of the map

$$\text{Mon}_{\mathcal{P}}(\text{Ar}^{\text{oplax}}) \rightarrow \text{Mon}_{\mathcal{P}}(\text{Cat}_{\infty}) \times \text{Mon}_{\mathcal{P}}(\text{Cat}_{\infty})$$

over the pair $(C_{\mathcal{P}}, D_{\mathcal{P}})$, where $C_{\mathcal{P}}$ and $D_{\mathcal{P}}$ are obtained from C and D by precomposition with the operad map $p_{\mathcal{P}} : \mathcal{P} \rightarrow \text{Comm}$. By Theorem 16.2.2, this fiber is the ∞ -category $\text{Fun}^{\mathcal{P}\text{-lax}}(C_{\mathcal{P}}, D_{\mathcal{P}}) = \text{Fun}_{(\text{Op}_{\infty})/\mathcal{P}}(\mathcal{M}_{C_{\mathcal{P}}/\mathcal{P}}, \mathcal{M}_{D_{\mathcal{P}}/\mathcal{P}})$ of lax $\mathcal{P}$ -monoidal functors $C_{\mathcal{P}} \rightarrow D_{\mathcal{P}}$. By the definition of $C_{\mathcal{P}}$ and $D_{\mathcal{P}}$ we have

$$\mathcal{M}_{C_{\mathcal{P}}/\mathcal{P}} \simeq \mathcal{P} \times_{\text{Comm}} \mathcal{M}_C \quad \text{and} \quad \mathcal{M}_{D_{\mathcal{P}}/\mathcal{P}} \simeq \mathcal{P} \times_{\text{Comm}} \mathcal{M}_D,$$

hence

$$\text{Fun}_{(\text{Op}_{\infty})/\mathcal{P}}(\mathcal{M}_{C_{\mathcal{P}}/\mathcal{P}}, \mathcal{M}_{D_{\mathcal{P}}/\mathcal{P}}) \simeq \text{Fun}_{(\text{Op}_{\infty})/\mathcal{P}}(\mathcal{P} \times_{\text{Comm}} \mathcal{M}_C, \mathcal{P} \times_{\text{Comm}} \mathcal{M}_D) \simeq \text{Fun}_{\text{Op}_{\infty}}(\mathcal{P} \times_{\text{Comm}} \mathcal{M}_C, \mathcal{M}_D),$$

where the second equivalence uses the universal property of pullbacks. Since Comm is terminal, the fiber product $\mathcal{P} \times_{\text{Comm}} \mathcal{M}_C$ is the product $\mathcal{P} \times \mathcal{M}_C$ in Op_{∞} . This was precisely what we wanted to show. $\square$

Remark 16.2.5. The above description of $\text{Day}(\mathcal{M}_C, \mathcal{M}_D)$ immediately generalizes to a description of the $\mathcal{O}$ -monoidal Day convolution operad $\text{Day}_{/\mathcal{O}}(\mathcal{M}_{C/\mathcal{O}}, \mathcal{M}_{D/\mathcal{O}}) \in (\text{Op}_{\infty})_{/\mathcal{O}}$ for $\mathcal{O}$ -monoidal ∞ -categories C and D . The proof is essentially identical, up to replacing Op_{∞} with $(\text{Op}_{\infty})_{/\mathcal{O}}$ everywhere. It is this generality in which Hinich and Winges state and prove the result. Note that Theorem 16.2.2, the principal external input about Day convolution, is already stated for an arbitrary ∞ -operad $\mathcal{O}$.

Concretely, the universal property reads

$$\text{Fun}_{(\text{Op}_{\infty})_{/\mathcal{O}}}(\mathcal{Q}, \text{Day}_{/\mathcal{O}}(\mathcal{M}_{C/\mathcal{O}}, \mathcal{M}_{D/\mathcal{O}})) \simeq \text{Fun}_{(\text{Op}_{\infty})_{/\mathcal{O}}}(\mathcal{Q} \times_{\mathcal{O}} \mathcal{M}_{C/\mathcal{O}}, \mathcal{M}_{D/\mathcal{O}})$$

for every ∞ -operad $\mathcal{Q}$ over $\mathcal{O}$. Moreover, since both sides are described by the same pullback square, base change along an operad map $\mathcal{O} \rightarrow \text{Comm}$ identifies

$$\mathcal{O} \times_{\text{Comm}} \text{Day}(\mathcal{M}_C, \mathcal{M}_D) \simeq \text{Day}_{/\mathcal{O}}(\mathcal{M}_{C/\mathcal{O}}, \mathcal{M}_{D/\mathcal{O}}).$$

In other words, the underlying $\mathcal{O}$ -monoidal ∞ -category of the Day convolution is the $\mathcal{O}$ -monoidal Day convolution. For $\mathcal{O} = \mathcal{A}ssoc$, this is the relative version underlying the final assertion of Observation 16.6.7, where the relevant functor is lax monoidal but *not* lax symmetric monoidal.

As a consequence of this result, we may explicitly compute the multimorphism anima of $\text{Day}(\mathcal{M}_C, \mathcal{M}_D)$:

Proposition 16.2.6. *Let C and D be symmetric monoidal ∞ -categories, let I be a finite set, and let $\{F_i : C \rightarrow D\}_{i \in I}$ and $G : C \rightarrow D$ be functors. Then there is an equivalence*

$$\text{Day}(\mathcal{M}_C, \mathcal{M}_D)(\{F_i\}_{i \in I}; G) \simeq \text{Nat} \left(\bigotimes_D^I \circ \prod_{i \in I} F_i, G \circ \bigotimes_C^I \right)$$

between the anima of multimorphisms $\{F_i\}_{i \in I} \rightarrow G$ in the Day convolution operad and the anima of natural transformations α fitting in the following diagram:

$$\begin{array}{ccc} C^I & \xrightarrow{\prod_{i \in I} F_i} & D^I \\ \otimes_C^I \downarrow & \swarrow & \downarrow \otimes_D^I \\ C & \xrightarrow{G} & D. \end{array}$$

Proof. Let us write $\mathcal{O} := \mathcal{Day}(\mathcal{M}_C, \mathcal{M}_D)$ for simplicity. By definition, the anima of multimorphisms $\mathcal{O}(\{F_i\}_{i \in I}; G)$ is the fiber of the projection map

$$p_{\mathcal{O}}: \text{Hom}_{\mathcal{O}^{\otimes}}(\{F_i\}_{i \in I}, G) \rightarrow \text{Hom}_{\text{Span}(\text{Fin})}(I, \langle 1 \rangle)$$

over the active span $I \xleftarrow{\text{id}} I \rightarrow \langle 1 \rangle$. By Theorem 16.2.4, the operad $\mathcal{O}$ sits in the following pullback square of ∞ -operads:

$$\begin{array}{ccc} \mathcal{O} & \xrightarrow{\quad} & \mathcal{O}_{\text{Ar}^{\text{oplax}}}^{\times} \\ \downarrow & \lrcorner & \downarrow (s, t) \\ \text{Comm} & \xrightarrow{(C, D)} & \mathcal{O}_{\text{Cat}_{\infty} \times \text{Cat}_{\infty}}^{\times} \end{array}$$

Since hom animae in a pullback of ∞ -categories form a pullback of animae, $\text{Hom}_{\mathcal{O}^{\otimes}}(\{F_i\}_{i \in I}, G)$ is itself the fiber in a pullback square of hom animae. Taking the fibers of these hom animae over the active span $I \rightarrow \langle 1 \rangle$, we find that $\mathcal{O}(\{F_i\}_{i \in I}; G)$ is the fiber in the following pullback square of animae:

$$\begin{array}{ccc} \mathcal{O}(\{F_i\}_{i \in I}; G) & \xrightarrow{\quad} & \mathcal{O}_{\text{Ar}^{\text{oplax}}}^{\times}(\{\text{Un}(F_i)\}_{i \in I}; \text{Un}(G)) \\ \downarrow & \lrcorner & \downarrow (s_*, t_*) \\ * & \xrightarrow{(\otimes_C^I, \otimes_D^I)} & \mathcal{O}_{\text{Cat}_{\infty}}^{\times}(\{C\}_{i \in I}; C) \times \mathcal{O}_{\text{Cat}_{\infty}}^{\times}(\{D\}_{i \in I}; D). \end{array}$$

Here the object $\text{Un}(F_i) \in \text{Ar}^{\text{oplax}}$ is the cartesian unstraightening of the functor $[1]^{\text{op}} \rightarrow \text{Cat}_{\infty}$ which takes the arrow $1 \rightarrow 0$ to $F_i: C \rightarrow D$, and similarly for $\text{Un}(G)$. Since Cat_{∞} and Ar^{oplax} have finite products, we may write the three multimorphism animae in the source and target of the right vertical map as hom animae in Ar^{oplax} and Cat_{∞} out of the corresponding products, which results in a pullback square as follows:

$$\begin{array}{ccc} \mathcal{O}(\{F_i\}_{i \in I}; G) & \xrightarrow{\quad} & \text{Hom}_{\text{Ar}^{\text{oplax}}}(\text{Un}(\prod_{i \in I} F_i), \text{Un}(G)) \\ \downarrow & \lrcorner & \downarrow (s_*, t_*) \\ * & \xrightarrow{(\otimes_C^I, \otimes_D^I)} & \text{Hom}_{\text{Cat}_{\infty}}(C^I, C) \times \text{Hom}_{\text{Cat}_{\infty}}(D^I, D). \end{array} \quad (16.1)$$

By applying Theorem 16.2.2 to the trivial ∞ -operad, we get that the functor $(s, t): \text{Ar}^{\text{oplax}} \rightarrow \text{Cat}_{\infty} \times \text{Cat}_{\infty}$ is a cartesian fibration over $\text{Cat}_{\infty} \times \{D\}$ and a cocartesian fibration over $\{C\} \times \text{Cat}_{\infty}$. The fibers of (s, t) are given by the functor categories $\text{Fun}(C, D)$, with functoriality in C given by precomposition and functoriality in D given by postcomposition. The fiber of the upper-right hom anima over a pair (f, g) is therefore the anima of natural transformations $g \circ \prod_i F_i \rightarrow G \circ f$. In particular, Equation (16.1) identifies the required fiber as

$$\mathcal{O}(\{F_i\}_{i \in I}; G) \simeq \text{Hom}_{\text{Fun}(C^I, D)}\left(\bigotimes_D^I \circ \prod_{i \in I} F_i, G \circ \bigotimes_C^I\right),$$

as claimed. $\square$

16.2.2 The Day convolution monoidal structure

With the explicit description of the multimorphism animae of $\mathcal{Day}(\mathcal{M}_C, \mathcal{M}_D)$ at hand, we may now return to the question of when this ∞ -operad corresponds to a symmetric monoidal structure on the functor category $\text{Fun}(C, D)$:

Proposition 16.2.7 (Lurie [Lur17, Proposition 2.2.6.16], Winges [Win25, Proposition 4.1]). *Let $(C, \otimes_C, \mathbb{1}_C)$ and $(D, \otimes_D, \mathbb{1}_D)$ be symmetric monoidal ∞ -categories. Let $\mathcal{K}$ be the collection of all relative slice categories $(\bigotimes_C^n)_{/c}$ defined by the pullback squares*

$$\begin{array}{ccc} (\bigotimes_C^n)_{/c} & \longrightarrow & \mathrm{Ar}(C) \\ \downarrow & & \downarrow (s,t) \\ C^n & \xrightarrow{(\bigotimes_{C,c}^n)} & C \times C \end{array}$$

for every natural number $n \geq 0$ and every object $c \in C$. Assume that the following two conditions are satisfied:

- (1) *The ∞ -category D admits all $\mathcal{K}$ -indexed colimits.*
- (2) *The tensor product functor $\otimes_D : D \times D \rightarrow D$ preserves $\mathcal{K}$ -indexed colimits in each variable separately.*

Then the Day convolution operad $\mathcal{D}\mathrm{ay}(\mathcal{M}_C, \mathcal{M}_D)$ defines a symmetric monoidal structure $\otimes_{\mathrm{Day}}$ on $\mathrm{Fun}(C, D)$.

In practice, we will usually apply Proposition 16.2.7 through the following corollary, so we record it before proving the proposition.

Corollary 16.2.8. *Let C and D be symmetric monoidal ∞ -categories. Assume that C is small, that D is cocomplete (i.e., admits all small colimits), and that the tensor product of D preserves small colimits in each variable. Then the Day convolution operad $\mathcal{D}\mathrm{ay}(\mathcal{M}_C, \mathcal{M}_D)$ defines a symmetric monoidal structure $\otimes_{\mathrm{Day}}$ on $\mathrm{Fun}(C, D)$.*

Proof. Every relative slice category $(\bigotimes_C^n)_{/c}$ is a pullback of the diagram $C^n \rightarrow C \times C \leftarrow \mathrm{Ar}(C)$. Since C is small, so are C^n and $\mathrm{Ar}(C)$, and hence so is $(\bigotimes_C^n)_{/c}$. The collection $\mathcal{K}$ of Proposition 16.2.7 therefore consists of small ∞ -categories, so both of its conditions are implied by the assumptions on D . $\square$

Proof of Proposition 16.2.7. We wish to show that the Day convolution operad $\mathcal{O} := \mathcal{D}\mathrm{ay}(\mathcal{M}_C, \mathcal{M}_D)$ defines a symmetric monoidal ∞ -category, i.e., that the functor $p_{\mathcal{O}} : \mathcal{O}^{\otimes} \rightarrow \mathrm{Span}(\mathrm{Fin})$ is a cocartesian fibration. Recall from Lemma 14.2.8 that it is enough to show that the following two conditions are satisfied:

- (a) For every finite collection of functors $\{F_i\}_{i \in I}$, there exists a functor $G = \bigotimes_{i \in I}^{\mathrm{Day}} F_i$ and a multimorphism $\{F_i\}_{i \in I} \rightarrow G$ in $\mathcal{O}$ inducing for every other functor $H : C \rightarrow D$ an equivalence

$$\mathrm{Hom}_{\mathcal{O}_{(\mathbb{1})}}(G, H) \xrightarrow{\sim} \mathcal{O}(\{F_i\}; H).$$

- (b) For a map $f : I \rightarrow J$ of finite sets, if we let $G_j := \bigotimes_{i \in f^{-1}(j)}^{\mathrm{Day}} F_i$ for all $j \in J$, then the canonical map

$$\bigotimes_{i \in I}^{\mathrm{Day}} F_i \rightarrow \bigotimes_{j \in J}^{\mathrm{Day}} G_j$$

is a natural isomorphism of functors $C \rightarrow D$.

For part (a), we define G as the left Kan extension of the composite functor

$$\prod_{i \in I} C \xrightarrow{\prod_{i \in I} F_i} \prod_{i \in I} D \xrightarrow{\otimes_D} D$$

along the tensor product functor $\otimes_C: C^I \rightarrow C$. Explicitly, for an object $c \in C$, its value is given by the colimit:

$$G(c) := \operatorname{colim}_{\{c'_i\}_{i \in I} \in (\otimes_C^I)_{/c}} \left(\bigotimes_{i \in I} F_i(c'_i) \right).$$

The existence of this left Kan extension is guaranteed by condition (1) of our hypothesis, as the indexing category $(\otimes_C^I)_{/c}$ is in our collection $\mathcal{K}$. The universal property of this left Kan extension provides a natural transformation

$$\eta: \bigotimes_D^I \circ \prod_{i \in I} F_i \Rightarrow G \circ \bigotimes_C^I,$$

which by Proposition 16.2.6 corresponds to a multimorphism $\eta: \{F_i\}_{i \in I} \rightarrow G$ in $\mathcal{O}$. The universal property of left Kan extension precisely says that for any other functor $H: C \rightarrow D$, composition with η induces an equivalence

$$\operatorname{Hom}_{\mathcal{O}_{(1)}}(G, H) \simeq \operatorname{Nat}(G, H) \xrightarrow{\sim} \operatorname{Nat}\left(\bigotimes_D^I \circ \prod_{i \in I} F_i, H \circ \bigotimes_C^I\right) \simeq \mathcal{O}(\{F_i\}_{i \in I}; H),$$

as desired.

We now show that also (b) holds. Let $f: I \rightarrow J$ be a map in $\mathbf{Fin}$, and write $G_j := \bigotimes_{i \in f^{-1}(j)}^{\text{Day}} F_i$ for all $j \in J$. We need to show that the canonical transformation

$$\bigotimes_{i \in I}^{\text{Day}} F_i \rightarrow \bigotimes_{j \in J}^{\text{Day}} G_j$$

is a natural isomorphism of functors $C \rightarrow D$. To help visualize this, consider the following diagram:

$$\begin{array}{ccc} C^I & \xrightarrow{\prod_{i \in I} F_i} & D^I \\ \left(\begin{array}{ccc} \downarrow (\otimes_C^f) & & \downarrow (\otimes_D^f) \\ C^J & \xrightarrow{\prod_{j \in J} G_j} & D^J \\ \downarrow \otimes_C^J & & \downarrow \otimes_D^J \end{array} \right) & & \\ \otimes_C^I & & \otimes_D^I \end{array} \quad \begin{array}{ccc} C & \dashrightarrow & D. \end{array}$$

Since left Kan extensions compose, it will suffice to show that the composite $\bigotimes_D^J \circ \prod_{j \in J} G_j: C^J \rightarrow D$ is the left Kan extension of the composite $\bigotimes_D^I \circ \prod_{i \in I} F_i: C^I \rightarrow D$ along $(\otimes_C^f): C^I \rightarrow C^J$. By the pointwise formula for Kan extensions, this amounts to showing that for every tuple $(c_j)_{j \in J} \in C^J$, the map

$$\operatorname{colim}_{(c'_i) \in (\otimes_C^f)_{/c}} \bigotimes_{i \in I}^D F_i(c'_i) \rightarrow \bigotimes_{j \in J} G_j(c_j)$$

is a natural isomorphism. Since the functor $\bigotimes_C^f: C^I \rightarrow C^J$ is a product of the individual functors $\bigotimes_C^{I_j}: C^{I_j} \rightarrow C$, the indexing category $(\bigotimes_C^f)_{/c}$ on the left is a product over $j \in J$ of the relative slice categories $(\bigotimes_C^{I_j})_{/c_j}$. But we also have

$$G_j(c_j) = \operatorname{colim}_{(c_i'') \in (\bigotimes_C^{I_j})_{/c_j}} \bigotimes_{i \in I_j}^D F_i(c_i''),$$

and since the functor $\bigotimes_{j \in J}^D: D^J \rightarrow D$ preserves the relevant colimits in each variable separately it follows that the map is a natural isomorphism. This finishes the proof. $\square$

Remark 16.2.9. The proof of the proposition provides an explicit description of the tensor product functor

$$\bigotimes_{\text{Day}}^I: \operatorname{Fun}(C, D)^I \rightarrow \operatorname{Fun}(C, D)$$

for every finite set I . For $I = \langle 0 \rangle$ and $I = \langle 2 \rangle$, this specializes as follows:

- The monoidal unit $\mathbb{1}_{\text{Day}}: C \rightarrow D$ of the Day convolution monoidal structure is the left Kan extension of $\mathbb{1}_D: * \rightarrow D$ along $\mathbb{1}_C: * \rightarrow C$.
- The Day convolution $F \otimes_{\text{Day}} G$ of two functors $F, G: C \rightarrow D$ is the left Kan extension of $C \times C \xrightarrow{F \times G} D \times D \xrightarrow{\otimes_D} D$ along $\otimes_C: C \times C \rightarrow C$.

Proposition 16.2.10 (Functoriality of monoidal Day convolution). *Assume the relevant Day convolution monoidal structures exist.*

- (1) *A lax symmetric monoidal functor $u: C' \rightarrow C$ induces a lax symmetric monoidal precomposition functor*

$$u^*: \operatorname{Fun}(C, D) \longrightarrow \operatorname{Fun}(C', D).$$

- (2) *Let $\mathcal{K}$ be the collection of relative slice categories from Proposition 16.2.7. If $v: D \rightarrow D'$ is symmetric monoidal and preserves $\mathcal{K}$ -indexed colimits, then postcomposition induces a symmetric monoidal functor*

$$v_*: \operatorname{Fun}(C, D) \longrightarrow \operatorname{Fun}(C, D').$$

Proof. Part (1) is induced by precomposition on Day convolution operads. For part (2), postcomposition first gives a lax symmetric monoidal refinement. Since v is symmetric monoidal and preserves the colimits in the pointwise formulas of Remark 16.2.9, it carries each Day convolution product and the Day convolution unit to the corresponding product and unit in $\operatorname{Fun}(C, D')$. Its structure maps are therefore isomorphisms. $\square$

Lemma 16.2.11 (Left Kan extensions and Day convolution). *Let $u: A \rightarrow B$ be a symmetric monoidal functor, and let D be a symmetric monoidal ∞ -category. Assume that the Day convolution monoidal structures on $\operatorname{Fun}(A, D)$ and $\operatorname{Fun}(B, D)$ exist. Suppose that for every object $b \in B$, the ∞ -category D admits colimits indexed by the relative slice category $u_{/b}$, and that its tensor product preserves*

these colimits separately in each variable. Then pointwise left Kan extension along u defines a symmetric monoidal functor

$$u_! : \text{Fun}(A, D) \longrightarrow \text{Fun}(B, D).$$

In particular, this applies when A and B are small, while D is cocomplete and its tensor product preserves small colimits separately in each variable.

Proof. The hypotheses on the relative slice categories ensure that pointwise left Kan extension along u exists. Its right adjoint u^* is lax symmetric monoidal by Proposition 16.2.10. By Proposition 14.3.11, it therefore suffices to show that for every finite collection of functors $F_i : A \rightarrow D$, indexed by a finite set I , the canonical comparison

$$u_! \left(\bigotimes_{i \in I}^{\text{Day}} F_i \right) \longrightarrow \bigotimes_{i \in I}^{\text{Day}} u_!(F_i)$$

is an isomorphism.

Set

$$H := \bigotimes_D^I \circ \prod_{i \in I} F_i : A^I \rightarrow D.$$

By the description of Day convolution in Remark 16.2.9 and the composition law for left Kan extensions, the source of the comparison is the left Kan extension of H along

$$A^I \xrightarrow{\otimes_A^I} A \xrightarrow{u} B.$$

On the other hand, the pointwise formula for left Kan extensions and the assumptions on the tensor product of D show that

$$\bigotimes_D^I \circ \prod_{i \in I} u_!(F_i) : B^I \rightarrow D$$

is the left Kan extension of H along $u^I : A^I \rightarrow B^I$: the relative slice of u^I over a tuple $(b_i)_{i \in I}$ is the product of the relative slices u/b_i , and its colimit may be computed iteratively. Applying Day convolution and composing left Kan extensions therefore identifies the target of the comparison with the left Kan extension of H along

$$A^I \xrightarrow{u^I} B^I \xrightarrow{\otimes_B^I} B.$$

Since u is symmetric monoidal, these two composite functors are naturally equivalent. The canonical comparison is therefore an isomorphism, as desired. $\square$

As a basic instance of Corollary 16.2.8, we obtain:

Corollary 16.2.12. *Let C be a small symmetric monoidal ∞ -category. Then the functor category $\text{Fun}(C, \text{An})$ admits a symmetric monoidal structure via Day convolution.*

Proof. The cited corollary applies since An admits small colimits and the functor $-\times- : \text{An} \times \text{An} \rightarrow \text{An}$ preserves colimits in both variables. $\square$

For later use, we record the following description of the internal hom for Day convolution.

Proposition 16.2.13. *Let C be a small symmetric monoidal ∞ -category and let D be a complete closed symmetric monoidal ∞ -category for which the Day convolution symmetric monoidal structure on $\text{Fun}(C, D)$ exists. Then this symmetric monoidal structure is closed. Its internal hom is pointwise given by the end formula*

$$\underline{\text{Hom}}_{\text{Fun}(C, D)}(F, G)(X) \simeq \int_{Y \in C} \underline{\text{Hom}}_D(F(Y), G(X \otimes_C Y)).$$

Here the integral denotes the end of Definition 23.6.2.

Proof. It suffices to show that for every third functor $H: C \rightarrow D$ there is a natural equivalence

$$\text{Nat}(H \otimes_{\text{Day}} F, G) \simeq \text{Nat}\left(H, \int_{Y \in C} \underline{\text{Hom}}_D(F(Y), G(- \otimes_C Y))\right).$$

Since the Day convolution is a left Kan extension, the left-hand side is equivalent to the anima

$$\text{Nat}(H(-) \otimes_D F(-), G(- \otimes_C -))$$

of natural transformations of functors $C \times C \rightarrow D$. By Proposition 23.6.5, this anima is the end

$$\int_{(X, Y) \in C \times C} \text{Hom}_D(H(X) \otimes_D F(Y), G(X \otimes_C Y)).$$

Using the Fubini rule for ends, Lemma 23.6.4, the adjunction $- \otimes_D F(Y) \dashv \underline{\text{Hom}}_D(F(Y), -)$, and the fact that $\text{Hom}_D(H(X), -)$ preserves limits, we may rewrite this as

$$\begin{aligned} & \int_{X \in C} \int_{Y \in C} \text{Hom}_D(H(X), \underline{\text{Hom}}_D(F(Y), G(X \otimes_C Y))) \\ & \simeq \int_{X \in C} \text{Hom}_D\left(H(X), \int_{Y \in C} \underline{\text{Hom}}_D(F(Y), G(X \otimes_C Y))\right). \end{aligned}$$

Using the end description of anima of natural transformations once more gives the desired result. $\square$

16.3 Examples of Day convolution

We now discuss several Day convolution structures that occur elsewhere in the book.

16.3.1 Graded and filtered objects

Example 16.3.1 (Graded objects). Regard $\mathbb{Z}$ as a discrete symmetric monoidal category under addition, and let C be a cocomplete symmetric monoidal ∞ -category whose tensor product preserves colimits separately in each variable. Day convolution equips the ∞ -category $\text{Fun}(\mathbb{Z}, C)$ of $\mathbb{Z}$ -graded objects with a symmetric monoidal structure. Explicitly,

$$(F \otimes_{\text{Day}} G)_n \simeq \coprod_{i+j=n} F_i \otimes G_j.$$

For $C = \text{Ab}$ this is the usual tensor product of graded abelian groups, and for $C = \text{Sp}$ it is the corresponding tensor product of graded spectra.

Example 16.3.2 (Filtered objects). Under the same hypotheses on C , regard $(\mathbb{Z}, \leq)$ as a poset with its usual ordering and as a symmetric monoidal category under addition. The Day convolution on $\text{Fun}((\mathbb{Z}, \leq), C)$ is the standard tensor product of increasing filtered objects:

$$(F \otimes_{\text{Day}} G)_n \simeq \text{colim}_{i+j \leq n} F_i \otimes G_j.$$

The transition maps are induced by the order on $\mathbb{Z}$. Replacing $(\mathbb{Z}, \leq)$ by $(\mathbb{N}, \leq)$ gives the analogous tensor product of nonnegatively filtered objects.

16.3.2 Pushout products and smash products

Lemma 16.3.3 (Pushout product). *Let C be a symmetric monoidal ∞ -category that admits finite colimits and such that $X \otimes - : C \rightarrow C$ preserves finite colimits for every $X \in C$. Then $\text{Ar}(C)$ admits a symmetric monoidal structure via Day convolution. Its unit is $\emptyset \rightarrow \mathbb{1}$, and its tensor product is the pushout product*

$$(f : X \rightarrow Y) \square (f' : X' \rightarrow Y') := (X \otimes Y') \sqcup_{X \otimes X'} (Y \otimes X') \longrightarrow Y \otimes Y'.$$

Proof. Equip $[1] = \{0 \leq 1\}$ with the symmetric monoidal structure given by the minimum. Since the relative tensor-product slices of $([1], \min)$ are finite, Proposition 16.2.7 applies to $\text{Fun}([1], C) = \text{Ar}(C)$. By Remark 16.2.9, the unit is the left Kan extension of $\mathbb{1} : * \rightarrow C$ along $\{1\} \hookrightarrow [1]$, hence is $\emptyset \rightarrow \mathbb{1}$. The binary Day convolution is the left Kan extension along $\min : [1] \times [1] \rightarrow [1]$; the pointwise formula identifies its value at 0 with the displayed pushout and its value at 1 with $Y \otimes Y'$. $\square$

Lemma 16.3.4 (Smash product). *Let C be a symmetric monoidal ∞ -category that admits finite colimits and a terminal object, and assume that $X \otimes - : C \rightarrow C$ preserves finite colimits for every $X \in C$. Equip $\text{Ar}(C)$ with the pushout-product monoidal structure of Lemma 16.3.3. Then $C_* \subseteq \text{Ar}(C)$ is symmetric monoidal, and the localization functor*

$$\text{cofib} : \text{Ar}(C) \rightarrow C_*, \quad (f : X \rightarrow Y) \longmapsto (* \rightarrow \text{cofib}(f))$$

is symmetric monoidal. The tensor product of pointed objects is therefore

$$(x : * \rightarrow X) \otimes (y : * \rightarrow Y) = \text{cofib}(X \otimes * \sqcup_{* \otimes *} * \otimes Y \rightarrow X \otimes Y).$$

Proof. The inclusion $C_* \hookrightarrow \text{Ar}(C)$ admits a left adjoint given by taking cofibers. By Proposition 14.5.3, it thus remains to show that tensoring in $\text{Ar}(C)$ with an object preserves those morphisms (= commutative squares in C) which induce isomorphisms on cofibers. To see this, consider such a morphism

$$\begin{array}{ccc} X & \longrightarrow & X' \\ f \downarrow & & \downarrow f' \\ Y & \longrightarrow & Y'. \end{array}$$

Let us write $Z := \text{cofib}(f : X \rightarrow Y)$ and $Z' := \text{cofib}(f' : X' \rightarrow Y')$, so that the condition on the square is that the induced map $Z \rightarrow Z'$ is an isomorphism. Our task is to show that for every other

morphism $(g: A \rightarrow B)$ in C the induced commutative square

$$\begin{array}{ccc} (X \otimes B) \sqcup_{X \otimes A} (Y \otimes A) & \longrightarrow & (X' \otimes B) \sqcup_{X' \otimes A} (Y' \otimes A) \\ f \square g \downarrow & & \downarrow f' \square g \\ Y \otimes B & \longrightarrow & Y' \otimes B \end{array}$$

again induces an isomorphism on (vertical) cofibers. Since $Z \xrightarrow{\sim} Z'$, it will suffice to express the two vertical cofibers purely in terms of g , Z and Z' . We will do so by showing that the bottom front square in the following commutative diagram is a pushout square:

$$\begin{array}{ccccc} X \otimes A & \xrightarrow{\quad} & * \otimes A & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & X \otimes B & & * \otimes B & \\ & \downarrow & \downarrow & \downarrow & \\ Y \otimes A & \xrightarrow{\quad} & Z \otimes A & & \\ & \searrow & \searrow & \searrow & \\ & (X \otimes B) \sqcup_{X \otimes A} (Y \otimes A) & \longrightarrow & (* \otimes B) \sqcup_{* \otimes A} (Z \otimes A) & \\ & \downarrow f \square g & & \downarrow & \\ & Y \otimes B & \longrightarrow & Z \otimes B & \end{array}$$

But this follows from the pasting law for pushout squares: the left and right faces of the diagram are pushouts by definition, while the back square and front rectangle are pushout squares since $- \otimes A$ and $- \otimes B$ preserve the pushout square defining Z as a cofiber of $X \rightarrow Y$. $\square$

Definition 16.3.5. Let C be an ∞ -category with finite products and finite colimits such that the functor $X \times -: C \rightarrow C$ preserves finite colimits for all $X \in C$. We may then apply Lemma 16.3.4 to the cartesian monoidal structure on C . We refer to the resulting symmetric monoidal structure $(C_*, \wedge, S^0)$ on the pointed objects in C as the *smash product*. The monoidal unit is $S^0 := * \sqcup *$, while the smash product $X \wedge Y$ is given by

$$X \wedge Y := \text{cofib}(X \vee Y \rightarrow X \times Y),$$

where $X \vee Y := X \sqcup_* Y$.

Example 16.3.6. Since the cartesian product in $\mathbf{An}$ preserves colimits separately in both variables, this equips $\mathbf{An}_*$ with the smash product monoidal structure. Its underlying tensor product is the smash product introduced in Definition 2.4.10.

16.4 Rings and semirings

Recall that an ordinary commutative ring may be equivalently described as a commutative algebra object in the symmetric monoidal category $(\mathbf{Ab}, \otimes)$ of abelian groups. Given two abelian groups A and B , their tensor product $A \otimes B$ is characterized by the property that group homomorphisms

$A \otimes B \rightarrow C$ correspond to bilinear maps $A \times B \rightarrow C$. We now use Day convolution to construct analogous tensor products on $\mathbf{CGrp}(C)$ and $\mathbf{CMon}(C)$ for any presentably symmetric monoidal ∞ -category C . This gives general notions of commutative ring and semiring objects.

Proposition 16.4.1. *Let C be a presentably symmetric monoidal ∞ -category. The inclusions*

$$\mathbf{CGrp}(C) \hookrightarrow \mathbf{CMon}(C) \hookrightarrow \mathbf{Fun}(\mathbf{Span}(\mathbf{Fin}), C)$$

admit left adjoints. Moreover, the full subcategories $\mathbf{CMon}(C)$ and $\mathbf{CGrp}(C)$ inherit symmetric monoidal structures for which these left adjoints are symmetric monoidal Bousfield localizations of the Day convolution monoidal structure. Their tensor products preserve small colimits separately in both variables.

The free commutative monoid functor

$$F_C^{\mathbf{CMon}}: C \longrightarrow \mathbf{CMon}(C),$$

left adjoint to evaluation at $\langle 1 \rangle \in \mathbf{Span}(\mathbf{Fin})$, is symmetric monoidal.

Proof. The existence of the two left adjoints is the structured-object case of presentability recorded in Proposition 22.2.7.

The category $\mathbf{Span}(\mathbf{Fin})$ is small, so Corollary 16.2.8 and Proposition 22.5.5 equip $\mathbf{Fun}(\mathbf{Span}(\mathbf{Fin}), C)$ with its presentably symmetric monoidal Day convolution structure. It remains to apply the internal-hom criterion of Lemma 14.5.6. By Proposition 22.5.3, the category C is closed, and the Day convolution internal hom is computed by the end formula

$$\underline{\mathbf{Hom}}_{\text{Day}}(F, G)(S) \simeq \int_{T \in \mathbf{Span}(\mathbf{Fin})} \underline{\mathbf{Hom}}_C(F(T), G(S \times T)).$$

For every finite set T , the functor $- \times T$ preserves finite coproducts of finite sets, which are the finite products in $\mathbf{Span}(\mathbf{Fin})$. Consequently, if G preserves finite products, then so does the functor $S \mapsto G(S \times T)$. Internal homs and ends preserve limits, so $\underline{\mathbf{Hom}}_{\text{Day}}(F, G)$ again lies in $\mathbf{CMon}(C)$. If G is grouplike, the shear map of $\underline{\mathbf{Hom}}_{\text{Day}}(F, G)$ is obtained by applying the limit-preserving functor $\underline{\mathbf{Hom}}_{\text{Day}}(F, -)$ to the shear isomorphism of G , so $\underline{\mathbf{Hom}}_{\text{Day}}(F, G)$ is grouplike as well. Thus both local subcategories are closed under internal hom with arbitrary F , and Lemma 14.5.6 gives the claimed symmetric monoidal Bousfield localizations.

Part (3) of Proposition 14.5.3 shows that the localized tensor products preserve small colimits separately in both variables.

It remains to consider the free commutative monoid functor. Let $\iota: * \rightarrow \mathbf{Span}(\mathbf{Fin})$ select the monoidal unit $\langle 1 \rangle$ for the cartesian product of finite sets. Left Kan extension along ι is left adjoint to evaluation at $\langle 1 \rangle$,

$$\iota_!: C \simeq \mathbf{Fun}(*, C) \rightleftarrows \mathbf{Fun}(\mathbf{Span}(\mathbf{Fin}), C) : \iota^*.$$

The functor ι is symmetric monoidal, and Lemma 16.2.11 shows that $\iota_!$ is symmetric monoidal. The composite of $\iota_!$ with the symmetric monoidal reflection onto $\mathbf{CMon}(C)$ is left adjoint to evaluation at $\langle 1 \rangle$ on commutative monoids, hence is $F_C^{\mathbf{CMon}}$. It is therefore symmetric monoidal. $\square$

Proposition 16.4.2 (Commutative monoids and groups under adjunctions). *Let*

$$L: C \rightleftarrows D : R$$

be an adjunction between presentably symmetric monoidal ∞ -categories, and assume that L is symmetric monoidal. Postcomposition with R induces functors on commutative monoids and groups which admit symmetric monoidal left adjoints:

$$\tilde{L}: \mathbf{CMon}(C) \rightleftarrows \mathbf{CMon}(D) : R_* \quad \text{and} \quad \tilde{L}: \mathbf{CGrp}(C) \rightleftarrows \mathbf{CGrp}(D) : R_*.$$

If L preserves finite products, then the two functors $\tilde{L}$ are given by postcomposition with L .

Proof. The right adjoint R preserves finite products, so postcomposition with R preserves commutative monoids. It also preserves grouplike commutative monoids, since it carries shear isomorphisms to shear isomorphisms. This gives the two functors denoted by R_* in the statement.

Let E_C and E_D denote either the commutative-monoid categories or the commutative-group categories of C and D , respectively. Write Q_C and Q_D for the corresponding reflections of Proposition 16.4.1, and i_C and i_D for their fully faithful right adjoints. Postcomposition gives an adjunction

$$L^{\text{Span}}: \mathbf{Fun}(\mathbf{Span}(\mathbf{Fin}), C) \rightleftarrows \mathbf{Fun}(\mathbf{Span}(\mathbf{Fin}), D) : R^{\text{Span}}.$$

The left adjoint of R_* is the composite

$$\tilde{L} := Q_D L^{\text{Span}} i_C.$$

The adjunctions $Q_D \dashv i_D$ and $L^{\text{Span}} \dashv R^{\text{Span}}$, together with the full faithfulness of i_C , immediately give $\tilde{L} \dashv R_*$.

Since L is a left adjoint, it preserves small colimits, so Proposition 16.2.10 shows that L^{Span} is symmetric monoidal for the ambient Day convolution structures. Moreover, the two functors

$$\tilde{L} Q_C \quad \text{and} \quad Q_D L^{\text{Span}}$$

are left adjoint to the same functor $R^{\text{Span}} i_D = i_C R_*$. They are therefore naturally isomorphic. Since $Q_D L^{\text{Span}}$ is symmetric monoidal, the universal property of the symmetric monoidal Bousfield localization Q_C supplies $\tilde{L}$ with a unique symmetric monoidal refinement. Its right adjoint R_* acquires the canonical lax symmetric monoidal structure of Proposition 14.3.6.

If L preserves finite products, postcomposition with L preserves commutative monoids and groups. The object $L^{\text{Span}} i_C(A)$ is then already local for every A , so the localization map

$$L^{\text{Span}} i_C(A) \longrightarrow i_D Q_D L^{\text{Span}} i_C(A)$$

is an isomorphism. Thus $\tilde{L}$ is given by postcomposition with L , as claimed. $\square$

Lemma 16.4.3 (Bilinear maps). *Let C be a presentably symmetric monoidal ∞ -category. For $A, B, M \in \mathbf{CMon}(C)$ there is a natural equivalence*

$$\mathbf{Hom}_{\mathbf{CMon}(C)}(A \otimes B, M) \simeq \mathbf{Nat}_{\mathbf{Span}(\mathbf{Fin})^2}(A(-) \otimes_C B(-), M(- \times -)).$$

The same formula holds in $\mathbf{CGrp}(C)$ for commutative group objects. We refer to objects of the anima on the right as bilinear maps from A and B to M .

Proof. The tensor product $A \otimes B$ is the reflection of the ambient Day convolution $A \otimes_{\text{Day}} B$. Since M is local, maps from this reflection to M are the same as maps from $A \otimes_{\text{Day}} B$ to M . The universal property of Day convolution, in the form of Proposition 16.2.6, identifies the latter anima with the displayed anima of natural transformations. The argument for commutative groups is identical. $\square$

Example 16.4.4 (Abelian groups). Take $C = \text{Set}$ with its cartesian monoidal structure. Then $\text{CGrp}(\text{Set}) = \text{Ab}$, and the symmetric monoidal structure supplied by Proposition 16.4.1 is the usual tensor product of abelian groups. Indeed, regard abelian groups A , B and C as product-preserving functors $\text{Span}(\text{Fin}) \rightarrow \text{Set}$, so that $A(S) \cong A^S$, and similarly for B and C . By the universal property of Day convolution and the fact that C is local, maps from the localized Day convolution of A and B to C correspond to families of maps

$$A(S) \times B(T) \longrightarrow C(S \times T)$$

that are functorial in the finite sets S and T and their spans.

Evaluating such a family at $S = T = *$ gives a map $\beta: A \times B \rightarrow C$. Compatibility with the spans encoding addition shows that β is additive in each variable. Conversely, a bilinear map $\beta: A \times B \rightarrow C$ determines the family

$$((a_s)_{s \in S}, (b_t)_{t \in T}) \longmapsto (\beta(a_s, b_t))_{(s,t) \in S \times T},$$

and bilinearity is exactly what is needed for compatibility with composition of spans. We therefore obtain natural bijections

$$\text{Hom}_{\text{CGrp}(\text{Set})}(A \otimes B, C) \cong \{\text{Bilinear maps } A \times B \rightarrow C\} \cong \text{Hom}_{\text{Ab}}(A \otimes_{\mathbb{Z}} B, C).$$

By Yoneda, the tensor product furnished by the Day convolution localization is the usual tensor product $A \otimes_{\mathbb{Z}} B$.

Proposition 16.4.5 (Path components). *The adjunction*

$$\text{CGrp}(\text{An}) \xrightleftharpoons{\pi_0} \text{CGrp}(\text{Set}) = \text{Ab}$$

is symmetric monoidal on the left and lax symmetric monoidal on the right, where Ab is equipped with its usual tensor product.

Proof. The functor $\pi_0: \text{An} \rightarrow \text{Set}$ is a symmetric monoidal left adjoint for the cartesian monoidal structures and preserves finite products. Thus Proposition 16.4.2 applies. Finally, Example 16.4.4 identifies the symmetric monoidal structure on $\text{CGrp}(\text{Set}) = \text{Ab}$ with the usual tensor product. $\square$

For a presentably symmetric monoidal ∞ -category D , we now set

$$\text{CRig}(D) := \text{CAlg}(\text{CMon}(D)) \quad \text{and} \quad \text{CRing}(D) := \text{CAlg}(\text{CGrp}(D)),$$

where $\text{CMon}(D)$ and $\text{CGrp}(D)$ carry the symmetric monoidal structures of Proposition 16.4.1. We refer to their objects as *commutative semiring objects* and *commutative ring objects* in D , respectively. For $D = \text{Set}$, these recover the 1-categories of commutative semirings and commutative rings.

Proposition 16.4.6 (Maximal subgroupoids). *Passing to maximal subgroupoids defines a canonical lax symmetric monoidal functor*

$$(-)^{\simeq}: \text{CMon}(\text{Cat}_{\infty}) \longrightarrow \text{CMon}(\text{An}),$$

where both categories carry the symmetric monoidal structures of Proposition 16.4.1.

Proof. The inclusion $\text{An} \hookrightarrow \text{Cat}_{\infty}$ is a symmetric monoidal left adjoint for the cartesian monoidal structures, preserves finite products, and has right adjoint $(-)^{\simeq}$. The result is therefore the commutative-monoid instance of Proposition 16.4.2. $\square$

16.5 Spectra as reduced excisive functors

The construction of Sp in Chapter 4 used prespectra. For the symmetric monoidal structure, a different model is more convenient. We will consider functors from finite pointed animae which send the zero object to a terminal object and pushout squares to pullback squares. We will show that these reduced excisive functors give a second model for spectra, and that their inclusion into the full functor category admits a left adjoint. This realizes Sp as a Bousfield localization of a functor category on which Day convolution is available.

Definition 16.5.1. Let $F: C \rightarrow D$ be a functor between ∞ -categories.

- (1) If C admits pushouts, we say that F is *excisive* if it sends pushout squares in C to pullback squares in D .
- (2) If C admits a terminal object, we say that F is *reduced* if it sends the terminal object of C to a terminal object of D .

We write $\mathrm{Exc}(C, D) \subseteq \mathrm{Fun}(C, D)$ for the full subcategory of excisive functors, $\mathrm{Fun}_*(C, D) \subseteq \mathrm{Fun}(C, D)$ for the full subcategory of reduced functors, and $\mathrm{Exc}_*(C, D)$ for their intersection.

Notation 16.5.2. We write $\mathrm{An}_*^{\mathrm{fin}}$ for the small ∞ -category of finite pointed animae. Its universal property is given in Lemma 4.3.12.

Definition 16.5.3. Let C be an ∞ -category with finite limits. We denote by $\mathrm{Sp}^{\mathrm{exc}}(C)$ the full subcategory

$$\mathrm{Sp}^{\mathrm{exc}}(C) := \mathrm{Exc}_*(\mathrm{An}_*^{\mathrm{fin}}, C) \subseteq \mathrm{Fun}(\mathrm{An}_*^{\mathrm{fin}}, C)$$

spanned by the reduced, excisive functors from $\mathrm{An}_*^{\mathrm{fin}}$ to C . We denote by $\Omega^\infty: \mathrm{Sp}^{\mathrm{exc}}(C) \rightarrow C$ the evaluation at $S^0 \in \mathrm{An}_*^{\mathrm{fin}}$.

A reduced functor $\mathrm{An}_*^{\mathrm{fin}} \rightarrow C$ has a canonical lift to C_* , obtained from the maps $F(*) \rightarrow F(X)$; cf. Lemma 4.1.13. We will therefore freely use the equivalence

$$\mathrm{Fun}_*(\mathrm{An}_*^{\mathrm{fin}}, C) \simeq \mathrm{Fun}_*(\mathrm{An}_*^{\mathrm{fin}}, C_*),$$

and its restriction to reduced excisive functors.

The following criterion is the basic reason that reduced excisive functors behave like spectra.

Construction 16.5.4. Let C be a pointed ∞ -category with finite colimits, let D be a pointed ∞ -category with finite limits, and let $F: C \rightarrow D$ be a reduced functor. Applying F to the pushout square defining $\Sigma_C X$ gives a square

$$\begin{array}{ccc} F(X) & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & F(\Sigma_C X), \end{array}$$

and hence a canonical map

$$\eta_X: F(X) \rightarrow \Omega_D F(\Sigma_C X).$$

Proposition 16.5.5 (Excisiveness criterion). *Let C be a pointed ∞ -category with finite colimits and let D be a pointed ∞ -category with finite limits. A reduced functor $F: C \rightarrow D$ is excisive if and only if the map $\eta_X: F(X) \rightarrow \Omega_D F(\Sigma_C X)$ is an isomorphism for every object $X \in C$.*

Proof. If F is excisive, it sends the pushout square defining $\Sigma_C X$ to a pullback square, which identifies $F(X)$ with $\Omega_D F(\Sigma_C X)$.

Conversely, suppose all maps η_X are isomorphisms. For a commutative square Q as below, write

$$P_F(Q) := F(X) \times_{F(W)} F(Y)$$

and let $\alpha_F(Q): F(Z) \rightarrow P_F(Q)$ be the comparison map. Now let Q be a pushout square

$$\begin{array}{ccc} Z & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow \\ Y & \longrightarrow & W, \end{array}$$

The standard 3×3 diagram of cofibers contains a map $W \rightarrow \Sigma_C Z$ whose composites with $X \rightarrow W$ and $Y \rightarrow W$ are zero. Applying F therefore gives a morphism of cospans

$$\begin{array}{ccccc} F(X) & \longrightarrow & F(W) & \longleftarrow & F(Y) \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F(\Sigma_C Z) & \longleftarrow & 0. \end{array}$$

Taking pullbacks defines a map

$$\beta_F(Q): P_F(Q) \longrightarrow \Omega_D F(\Sigma_C Z).$$

Put $T(F) := \Omega_D F \Sigma_C$. The remaining faces of the 3×3 diagram identify the two indicated length-two composites in

$$\begin{array}{ccccccc} & & & & P_{\eta_F}(Q) & & \\ & & & & \curvearrowright & & \\ F(Z) & \xrightarrow{\alpha_F(Q)} & P_F(Q) & \xrightarrow{\beta_F(Q)} & T(F)(Z) & \xrightarrow{\alpha_{T(F)}(Q)} & P_{T(F)}(Q). \\ & \searrow & \eta_Z & \nearrow & & & \end{array}$$

The first identity is the suspension square defining η_Z . The second is obtained by taking pullbacks of the suspension squares defining η_X, η_W, η_Y ; here $P_{\eta_F}(Q)$ denotes the resulting map on pullbacks. The first length-two composite is an isomorphism by assumption, and the second is an isomorphism because it is induced on pullbacks by three isomorphisms. Hence the 2-out-of-6 property shows that $\alpha_F(Q)$ is an isomorphism. Thus F is excisive. $\square$

Exercise 16.5.6. Use the criterion to show that the following conditions are equivalent for a reduced functor $F: C \rightarrow D$ between stable ∞ -categories:

- (1) F is an exact functor;
- (2) F commutes with loop spaces;

(3) F commutes with suspensions.

Proposition 16.5.7. *Let C be a pointed ∞ -category with finite colimits and let D be an ∞ -category with finite limits. Then $\mathrm{Exc}_*(C, D)$ is stable.*

Proof. By replacing D with D_* , we may assume D is pointed. The ∞ -category $\mathrm{Exc}_*(C, D)$ has finite limits computed pointwise, and reduced excisive functors are closed under such limits. It is also pointed. By Theorem 4.2.2, it remains to show that its loop functor is an equivalence.

The loop functor is computed pointwise: $\Omega(F) = \Omega_D \circ F$. Define a shift functor by $\mathrm{Shift}(F) := F \circ \Sigma_C$. Since Σ_C preserves zero objects and pushout squares, $\mathrm{Shift}(F)$ is again reduced excisive. Applying an excisive functor F to the pushout square defining suspension gives natural isomorphisms

$$F \simeq \Omega_D F \Sigma_C = \Omega(\mathrm{Shift} F) = \mathrm{Shift}(\Omega F),$$

so Ω and Shift are inverse equivalences. $\square$

Lemma 16.5.8. *If C is stable, the evaluation functor $\Omega^\infty : \mathrm{Sp}^{\mathrm{exc}}(C) \rightarrow C$ is an equivalence.*

Proof. When C is stable, a reduced functor $\mathrm{An}_*^{\mathrm{fin}} \rightarrow C$ is excisive if and only if it preserves finite colimits. Hence

$$\mathrm{Sp}^{\mathrm{exc}}(C) = \mathrm{Exc}_*(\mathrm{An}_*^{\mathrm{fin}}, C) \simeq \mathrm{Fun}_*^{\mathrm{rex}}(\mathrm{An}_*^{\mathrm{fin}}, C).$$

Evaluation at S^0 is an equivalence by the universal property of $\mathrm{An}_*^{\mathrm{fin}}$ from Notation 16.5.2. $\square$

Corollary 16.5.9. *The ∞ -category $\mathrm{Sp}^{\mathrm{exc}}(C)$ is stable for every ∞ -category C with finite limits.*

Proof. Take the source category in Proposition 16.5.7 to be $\mathrm{An}_*^{\mathrm{fin}}$. $\square$

Proposition 16.5.10. *For every ∞ -category C with finite limits, the evaluation functor $\mathrm{ev}_{S^0} : \mathrm{Sp}^{\mathrm{exc}}(C) \rightarrow C$ exhibits $\mathrm{Sp}^{\mathrm{exc}}(C)$ as a stabilization of C .*

Proof. The functor ev_{S^0} is left exact, since limits in the functor category are computed pointwise and reduced excisive functors are closed under limits. Let D be stable. Since pushout squares and pullback squares agree in D , a functor $D \rightarrow C$ is left exact if and only if it is reduced and excisive. Thus it is enough to show that

$$\mathrm{Exc}_*(D, \mathrm{Sp}^{\mathrm{exc}}(C)) \rightarrow \mathrm{Exc}_*(D, C)$$

is an equivalence. Since reducedness and excisiveness are pointwise conditions, currying restricts to the relevant full subcategories and identifies the source with

$$\mathrm{Exc}_*(D, \mathrm{Exc}_*(\mathrm{An}_*^{\mathrm{fin}}, C)) \simeq \mathrm{Sp}^{\mathrm{exc}}(\mathrm{Exc}_*(D, C)).$$

The target $\mathrm{Exc}_*(D, C)$ is stable by Proposition 16.5.7, so evaluation at S^0 is an equivalence by Lemma 16.5.8. $\square$

Corollary 16.5.11. *There is a unique equivalence*

$$\mathrm{Sp} \simeq \mathrm{Sp}^{\mathrm{exc}}(\mathrm{An}) = \mathrm{Exc}_*(\mathrm{An}_*^{\mathrm{fin}}, \mathrm{An})$$

under which the two functors to An given by Ω^∞ and evaluation at S^0 agree.

Proof. By Propositions 4.3.16 and 16.5.10, both sides, equipped with the displayed functors to $\mathbf{An}$, are stabilizations of $\mathbf{An}$. The result follows from the uniqueness of stabilization. $\square$

Proposition 16.5.12 (Excisive approximation). *Let C be an ∞ -category that admits finite limits, finite colimits, and sequential colimits, and assume that the loop functor $\Omega: C_* \rightarrow C_*$ preserves sequential colimits. Then the two inclusions*

$$\mathrm{Sp}^{\mathrm{exc}}(C) = \mathrm{Exc}_*(\mathbf{An}_*^{\mathrm{fin}}, C) \hookrightarrow \mathrm{Fun}_*(\mathbf{An}_*^{\mathrm{fin}}, C) \quad \text{and} \quad \mathrm{Fun}_*(\mathbf{An}_*^{\mathrm{fin}}, C) \hookrightarrow \mathrm{Fun}(\mathbf{An}_*^{\mathrm{fin}}, C)$$

admit left adjoints

$$P_1: \mathrm{Fun}_*(\mathbf{An}_*^{\mathrm{fin}}, C) \rightarrow \mathrm{Exc}_*(\mathbf{An}_*^{\mathrm{fin}}, C) \quad \text{and} \quad (-)^{\mathrm{red}}: \mathrm{Fun}(\mathbf{An}_*^{\mathrm{fin}}, C) \rightarrow \mathrm{Fun}_*(\mathbf{An}_*^{\mathrm{fin}}, C).$$

Proof. For the reduction functor, let $F: \mathbf{An}_*^{\mathrm{fin}} \rightarrow C$ be arbitrary. Since $*$ is initial in $\mathbf{An}_*^{\mathrm{fin}}$, the unique natural transformation $\mathrm{const}_* \rightarrow \mathrm{id}$ induces $\mathrm{const}_{F(*)} \rightarrow F$. Define

$$F^{\mathrm{red}}(X) := * \sqcup_{F(*)} F(X).$$

This is reduced. If G is reduced, every natural transformation $F \rightarrow G$ vanishes on $F(*)$, and hence factors uniquely through $F \rightarrow F^{\mathrm{red}}$. Thus $(-)^{\mathrm{red}}$ is left adjoint to the inclusion of reduced functors.

For P_1 , we may replace C by C_* , which inherits finite and sequential colimits from C , so assume C is pointed. Write

$$T(F) := \Omega F \Sigma$$

for the endofunctor of $\mathrm{Fun}_*(\mathbf{An}_*^{\mathrm{fin}}, C)$, equipped with the natural transformation $\eta: \mathrm{id} \rightarrow T$ from Construction 16.5.4. We first record the coherence needed below. For a finite set S and $X \in \mathbf{An}_*^{\mathrm{fin}}$, let $K_S(X)$ be the colimit of the star-shaped diagram with central object X and one map $X \rightarrow 0$ for every element of S . Thus

$$K_\emptyset(X) = X, \quad K_{\{s\}}(X) = 0, \quad K_{\{0,1\}}(X) = \Sigma X.$$

Injectons of finite sets induce maps between these pointed cones. Since finite colimits commute with each other, interchanging the two colimit coordinates gives natural isomorphisms $K_S K_{S'} \simeq K_{S'} K_S$, coherently in S and S' . This simultaneous interchange is the coherence needed below; it does not require choosing an order in which to identify the two suspension coordinates. Moreover,

$$T(F)(X) \simeq \lim_{\emptyset \neq S \subseteq \{0,1\}} F(K_S(X)) \simeq F(0) \times_{F(\Sigma X)} F(0) \simeq \Omega F(\Sigma X).$$

Under this description, the maps $K_\emptyset(X) \rightarrow K_S(X)$ induce η_F . The isomorphisms which interchange K_S and $K_{S'}$ therefore identify the two natural transformations

$$T(\eta_F), \eta_{T(F)}: T(F) \longrightarrow T^2(F).$$

Now let $F: \mathbf{An}_*^{\mathrm{fin}} \rightarrow C$ be reduced. Iterate η to form

$$F \xrightarrow{\eta_F} T(F) \xrightarrow{\eta_{T(F)}} T^2(F) \longrightarrow \dots$$

and define

$$P_1(F) := \mathrm{colim}_{n \geq 0} T^n(F).$$

Thus $T^n(F) \simeq \Omega^n F \Sigma^n$, with the transition maps specified coherently by the pointed-cone construction.

The functor T preserves sequential colimits by assumption. Consequently,

$$T(P_1(F)) \simeq \operatorname{colim}_{n \geq 0} T^{n+1}(F),$$

and under this isomorphism $\eta_{P_1(F)}$ is the map from the defining telescope to its tail. The shift $n \mapsto n + 1$ is cofinal, so this map is an isomorphism. The excisiveness criterion now shows that $P_1(F)$ is excisive.

It remains to verify the universal property. The coherence $T(\eta_G) \simeq \eta_{T(G)}$ identifies the sequential diagram defining $P_1(T(G))$ with the tail of the diagram defining $P_1(G)$, compatibly with

$$P_1(\eta_G): P_1(G) \longrightarrow P_1(T(G)).$$

Thus $P_1(\eta_G)$ corresponds to the cofinal tail map and is an isomorphism. Moreover, P_1 preserves sequential colimits, since T does.

Let $u_F: F \rightarrow P_1(F)$ be the map into the telescope. Since P_1 preserves sequential colimits, the map

$$P_1(u_F): P_1(F) \longrightarrow P_1(P_1(F))$$

is the sequential colimit of the maps $P_1(F) \rightarrow P_1(T^n(F))$. Each of these is a composite of the isomorphisms $P_1(\eta_{T^k(F)})$, so $P_1(u_F)$ is an isomorphism. If G is already excisive, the excisiveness criterion shows that every map in the defining telescope for $P_1(G)$ is an isomorphism, so $u_G: G \rightarrow P_1(G)$ is an isomorphism. In particular, $u_{P_1(F)}$ is an isomorphism. By Proposition 21.8.9, the functor P_1 is left adjoint to the inclusion of reduced excisive functors. $\square$

Corollary 16.5.13. *The conclusion of Proposition 16.5.12 applies when $C = \mathbf{An}$ or $C = \mathbf{An}_*$.*

Proof. Both categories admit the required limits and colimits. Moreover, filtered colimits of anima $\mathbf{e}$ commute with finite limits, so their loop functors preserve sequential colimits. $\square$

Lemma 16.5.14. *The evaluation functor*

$$\operatorname{ev}_{S^0}: \operatorname{Exc}_*(\mathbf{An}_*^{\operatorname{fin}}, \mathbf{An}_*) \rightarrow \mathbf{An}_*$$

admits a left adjoint. It sends a pointed anima X to the excisive approximation of the smash product functor

$$P_1(X \wedge -): \mathbf{An}_*^{\operatorname{fin}} \rightarrow \mathbf{An}_*.$$

Proof. First consider evaluation at S^0 on the full functor category. Its left adjoint is the left Kan extension along $S^0: * \rightarrow \mathbf{An}_*^{\operatorname{fin}}$, which sends a pointed anima X to the functor

$$K \mapsto X \wedge \operatorname{fgt}(K)_+.$$

Its value at the zero object is X . Applying the reduction functor gives $K \mapsto X \wedge K$, since the cofiber of the canonical map $S^0 \rightarrow \operatorname{fgt}(K)_+$ is K . Thus $X \wedge -$ is left adjoint to evaluation on reduced functors. Composing with the excisive approximation P_1 , which exists by Corollary 16.5.13, gives the asserted left adjoint on reduced excisive functors. $\square$

16.6 The symmetric monoidal structure on $\mathbf{Sp}$

We will now combine the results from the previous sections to upgrade the tensor product $X \otimes Y$ of spectra from Section 4.4.3 to a symmetric monoidal structure $(\mathbf{Sp}, \otimes, \mathbb{S})$ on the ∞ -category of spectra.

Let us start by outlining the strategy. By Corollary 16.5.11, we have an equivalence $\mathbf{Sp} \simeq \mathbf{Exc}_*(\mathbf{An}_*^{\text{fin}}, \mathbf{An})$ between the ∞ -category of spectra and the ∞ -category of reduced excisive functors from finite pointed animae to animae. In particular, we may regard $\mathbf{Sp}$ as a full subcategory of the functor category $\mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An})$. By Proposition 16.2.7, this functor category admits the Day convolution monoidal structure. We will show that $\mathbf{Sp}$ is a symmetric monoidal Bousfield localization of it.

As the first step in this plan, we need to equip $\mathbf{An}_*^{\text{fin}}$ with a monoidal structure. The smash product on $\mathbf{An}_*$ from Definition 16.3.5 restricts to $\mathbf{An}_*^{\text{fin}}$ by Lemma 14.5.2. Indeed, for fixed $X \in \mathbf{An}_*^{\text{fin}}$, the functor $X \wedge -$ preserves finite colimits, so the fact that $\mathbf{An}_*^{\text{fin}}$ is generated under finite colimits by S^0 reduces closure under smash product to $X \wedge S^0 \xrightarrow{\sim} X$.

Combining everything, we can now construct the tensor product on $\mathbf{Sp}$:

Theorem 16.6.1. *The ∞ -category $\mathbf{Sp}$ admits a unique symmetric monoidal structure such that the reduced excisive approximation functor*

$$P_1^{\text{red}}: \mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An}) \rightarrow \mathbf{Exc}_*(\mathbf{An}_*^{\text{fin}}, \mathbf{An}) \simeq \mathbf{Sp}$$

is symmetric monoidal with respect to Day convolution on its source. This structure has the following properties:

- (1) *Its monoidal unit is the sphere spectrum $\mathbb{S}$, and the unreduced suspension spectrum functor $\mathbb{S}[-]: (\mathbf{An}, \times) \rightarrow (\mathbf{Sp}, \otimes)$ is strongly monoidal.*
- (2) *Its tensor product preserves small colimits separately in both variables.*
- (3) *Its underlying bifunctor agrees with the tensor product of spectra characterized in Proposition 4.4.16.*

Proof. Equip $\mathbf{An}_*^{\text{fin}}$ with the smash product monoidal structure from Definition 16.3.5, and equip $\mathbf{An}$ with the cartesian monoidal structure. The category $\mathbf{An}_*^{\text{fin}}$ is small by Notation 16.5.2. Corollary 16.2.12 therefore gives the Day convolution monoidal structure on $\mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An})$.

By Corollary 16.5.13, the inclusion

$$\iota: \mathbf{Sp} \simeq \mathbf{Exc}_*(\mathbf{An}_*^{\text{fin}}, \mathbf{An}) \hookrightarrow \mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An})$$

admits the left adjoint

$$P_1^{\text{red}} := P_1 \circ (-)^{\text{red}}.$$

We show that this Bousfield localization satisfies the internal-hom criterion of Lemma 14.5.6. By Proposition 16.2.13, the Day convolution monoidal structure on $\mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An})$ admits an internal hom given by

$$\underline{\text{Hom}}_{\mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, \mathbf{An})}(F, G)(X) \simeq \int_{Y \in \mathbf{An}_*^{\text{fin}}} \underline{\text{Hom}}_{\mathbf{An}}(F(Y), G(X \wedge Y)).$$

Here the internal hom in $\mathbf{An}$ agrees with the hom anima:

$$\underline{\mathrm{Hom}}_{\mathbf{An}}(A, B) \simeq \mathrm{Hom}_{\mathbf{An}}(A, B).$$

By Lemma 14.5.6, it now remains to show that this internal hom is reduced and excisive whenever G is.

We first check that it is reduced. If X is the zero object of $\mathbf{An}_*^{\mathrm{fin}}$, then so is $X \wedge Y$, hence $G(X \wedge Y)$ is terminal by reducedness of G . Since $\underline{\mathrm{Hom}}_{\mathbf{An}}(F(Y), -)$ preserves terminal anima and an end of terminal anima is terminal, this shows that the internal hom is reduced.

We now check that it is also excisive. Note that $- \wedge Y$ preserves pushouts, and so the functor $G(- \wedge Y): \mathbf{An}_*^{\mathrm{fin}} \rightarrow \mathbf{An}$ sends pushouts to pullbacks for all Y . The same then holds true for $\underline{\mathrm{Hom}}_{\mathbf{An}}(F(Y), G(- \wedge Y))$, and thus also for the end-expression since its defining limit commutes with pullbacks. We conclude that $\underline{\mathrm{Hom}}_{\mathrm{Fun}(\mathbf{An}_*^{\mathrm{fin}}, \mathbf{An})}(F, G)$ is excisive.

The internal-hom criterion, together with Proposition 14.5.3, now supplies the symmetric monoidal structure on the full subcategory of reduced excisive functors and the strong symmetric monoidal refinement of P_1^{red} . The ambient Day convolution structure is presentably symmetric monoidal by Proposition 22.5.5, so part (3) of Proposition 14.5.3 shows that the tensor product on $\mathbf{Sp}$ preserves small colimits separately in both variables.

To see uniqueness, suppose that another symmetric monoidal structure made P_1^{red} symmetric monoidal. For local objects X_i and Y , adjunction and symmetric monoidality give

$$\mathcal{M}_{\mathbf{Sp}}(\{X_i\}; Y) \simeq \mathcal{D}\mathrm{ay}(\mathcal{M}_{\mathbf{An}_*^{\mathrm{fin}}}, \mathcal{M}_{\mathbf{An}})(\{\iota(X_i)\}; \iota(Y)).$$

Its multimorphism operad is therefore the full suboperad of the ambient Day convolution operad on the local objects, which is determined by the localization. The remaining assertions are proved in Proposition 16.6.2 and Corollary 16.6.3. $\square$

Proposition 16.6.2. *The unreduced suspension spectrum functor $\mathbb{S}[-]: \mathbf{An} \rightarrow \mathbf{Sp}$ lifts to a strongly monoidal functor $\mathbb{S}[-]: (\mathbf{An}, \times) \rightarrow (\mathbf{Sp}, \otimes)$. In particular, the monoidal unit of $\mathbf{Sp}$ is the sphere spectrum $\mathbb{S}$.*

Proof. We will show that $\Omega^\infty: \mathbf{Sp} \rightarrow \mathbf{An}$ is lax symmetric monoidal and that it has a strong monoidal left adjoint.

Under the equivalence $\mathbf{Sp} \simeq \mathrm{Exc}_*(\mathbf{An}_*^{\mathrm{fin}}, \mathbf{An})$, we may identify Ω^∞ with the composite

$$\mathbf{Sp} \hookrightarrow \mathrm{Fun}(\mathbf{An}_*^{\mathrm{fin}}, \mathbf{An}) \xrightarrow{\mathrm{ev}_{S^0}} \mathbf{An}.$$

The first functor is lax symmetric monoidal by construction of the tensor product of $\mathbf{Sp}$. The second is lax symmetric monoidal by the precomposition statement of Proposition 16.2.10, applied to the symmetric monoidal functor $S^0: * \rightarrow \mathbf{An}_*^{\mathrm{fin}}$.

The left adjoint of the first functor is symmetric monoidal by construction of the tensor product on $\mathbf{Sp}$. Since $\mathbf{An}_*^{\mathrm{fin}}$ is small and the cartesian product of anima preserves small colimits separately in both variables, Lemma 16.2.11 shows that left Kan extension along $S^0: * \rightarrow \mathbf{An}_*^{\mathrm{fin}}$ is symmetric monoidal. The left adjoint $\mathbb{S}[-]$ of the displayed composite is therefore symmetric monoidal. $\square$

Corollary 16.6.3. *The tensor product functor $- \otimes -: \mathbf{Sp} \times \mathbf{Sp} \rightarrow \mathbf{Sp}$ defined by Theorem 16.6.1 agrees with the one characterized by Proposition 4.4.16.*

Proof. By parts (1) and (2) of Theorem 16.6.1, the tensor product preserves colimits in both variables, and the relation $\mathbb{S} \otimes \mathbb{S} \simeq \mathbb{S}$ holds since $\mathbb{S}$ is the monoidal unit. These are precisely the properties that determine the tensor product uniquely in Proposition 4.4.16, so the two agree. $\square$

16.6.1 The multiplicative infinite loop space machine

The Day convolution structures constructed above are compatible with the familiar passage from animae to spectra. We record the complete sequence, since it is the multiplicative input used for connective complex K-theory in Section 19.7.

Proposition 16.6.4. *The lax symmetric monoidal functors*

$$\mathrm{Sp} \xrightarrow{\Omega^\infty} \mathrm{CGrp}(\mathrm{An}) \hookrightarrow \mathrm{CMon}(\mathrm{An}) \xrightarrow{\mathrm{fgt}} \mathrm{An}_* \xrightarrow{\mathrm{fgt}} \mathrm{An}$$

admit strongly monoidal left adjoints:

$$(\mathrm{An}, \times) \xrightarrow{(-)_+} (\mathrm{An}_*, \wedge) \xrightarrow{F^{\mathrm{CMon}}} (\mathrm{CMon}(\mathrm{An}), \otimes) \xrightarrow{(-)^{\mathrm{grp}}} (\mathrm{CGrp}(\mathrm{An}), \otimes) \xrightarrow{\mathbf{B}^\infty} (\mathrm{Sp}, \otimes).$$

Here, $(-)_+$ adjoins a disjoint basepoint, F^{CMon} is the free commutative monoid functor, $(-)^{\mathrm{grp}}$ is group completion, and $\mathbf{B}^\infty$ is infinite delooping.

Proof. The functor

$$(\mathrm{An}, \times) \longrightarrow (\mathrm{Ar}(\mathrm{An}), \sqcap), \quad X \longmapsto (\emptyset \rightarrow X),$$

is symmetric monoidal: the pushout-product formula identifies its tensor comparison with $\emptyset \rightarrow X \times Y$, and its unit is $\emptyset \rightarrow *$. Its composite with the symmetric monoidal cofiber functor of Lemma 16.3.4 is $(-)_+$, so adjoining a basepoint is strongly monoidal.

The same conclusion for F^{CMon} follows from the bilinear universal property of Lemma 16.4.3. Indeed, for pointed animae X, Y and a commutative monoid M , there are natural equivalences

$$\begin{aligned} \mathrm{Hom}_{\mathrm{CMon}(\mathrm{An})}(F^{\mathrm{CMon}}(X) \otimes F^{\mathrm{CMon}}(Y), M) &\simeq \mathrm{Nat}_{\mathrm{Span}(\mathrm{Fin})^2}(F^{\mathrm{CMon}}(X)(-) \times F^{\mathrm{CMon}}(Y)(-), M(- \times -)) \\ &\simeq \mathrm{Hom}_{\mathrm{An}_*}(X \wedge Y, \mathrm{fgt}(M)) \\ &\simeq \mathrm{Hom}_{\mathrm{CMon}(\mathrm{An})}(F^{\mathrm{CMon}}(X \wedge Y), M). \end{aligned}$$

The middle equivalence is obtained by applying the free commutative monoid adjunction in each variable; the functors $S \mapsto M(S \times T)$ remain commutative monoids because $- \times T$ preserves the finite products of $\mathrm{Span}(\mathrm{Fin})$. The analogous nullary argument identifies the monoidal units. The group completion functor is strongly monoidal because it is the symmetric monoidal Bousfield localization of Proposition 16.4.1.

It remains to consider infinite delooping. Write $F^{\mathrm{CGrp}} := (-)^{\mathrm{grp}} F^{\mathrm{CMon}}$. The objects $F^{\mathrm{CGrp}}(X)$ generate $\mathrm{CGrp}(\mathrm{An})$ under colimits, and infinite delooping preserves colimits because it identifies $\mathrm{CGrp}(\mathrm{An})$ with the full subcategory $\mathrm{Sp}_{\geq 0} \subseteq \mathrm{Sp}$ of connective spectra. Under this identification,

$$\mathbf{B}^\infty F^{\mathrm{CGrp}}(X) \simeq \Sigma^\infty X.$$

Here $\Sigma^\infty X$ denotes the reduced suspension spectrum of the pointed anima X . Both functors

$$(G, H) \longmapsto \mathbf{B}^\infty(G \otimes H) \quad \text{and} \quad (G, H) \longmapsto \mathbf{B}^\infty G \otimes \mathbf{B}^\infty H$$

preserve colimits separately in G and H . On free generators, the canonical comparison is the strong monoidal comparison for reduced suspension spectra. This follows from Proposition 16.6.2: using the cofiber description of reduced suspension spectra from Remark 4.4.3, the pushout-product formula identifies this comparison with the isomorphism

$$\Sigma^\infty X \otimes \Sigma^\infty Y \simeq \Sigma^\infty (X \wedge Y).$$

The comparison is therefore an equivalence for all G and H . The same argument treats the unit and the coherence maps, proving that $\mathbf{B}^\infty$ is strongly monoidal. Finally, the right adjoints acquire the displayed lax symmetric monoidal structures by Proposition 14.3.6. $\square$

Corollary 16.6.5. *Infinite delooping induces a symmetric monoidal equivalence*

$$\mathbf{B}^\infty : (\mathrm{CGrp}(\mathrm{An}), \otimes) \xrightarrow{\simeq} (\mathrm{Sp}_{\geq 0}, \otimes).$$

Proof. The underlying functor is an equivalence by Theorem 5.4.5, and it is symmetric monoidal by Proposition 16.6.4. $\square$

16.6.2 Graded homotopy groups

The ordinary Day convolution tensor product on $\mathrm{Ab}^{\mathbb{Z}} = \mathrm{Fun}(\mathbb{Z}, \mathrm{Ab})$ is

$$(A \otimes B)_k = \bigoplus_{n+m=k} A_n \otimes B_m.$$

Its usual symmetry merely interchanges the two factors. For homotopy groups, the natural symmetry is instead the following signed variant.

Definition 16.6.6 (Koszul symmetry). Keeping the same underlying Day convolution monoidal structure on $\mathrm{Ab}^{\mathbb{Z}}$, we replace its ordinary symmetry by the *Koszul symmetry*, which on homogeneous elements is given by

$$A_n \otimes B_m \longrightarrow B_m \otimes A_n, \quad a \otimes b \longmapsto (-1)^{nm} b \otimes a.$$

We refer to the resulting symmetric monoidal category as the category of graded abelian groups with the Koszul symmetry.

Since $\mathrm{Ab}^{\mathbb{Z}}$ is a 1-category, it is enough to check the ordinary hexagon and involutivity axioms. These follow directly from the fact that $(n, m) \mapsto (-1)^{nm}$ is a symmetric bicharacter: bilinearity gives the two hexagon identities, while $(-1)^{nm}(-1)^{mn} = 1$ gives involutivity.

Observation 16.6.7 (Graded homotopy groups). The functor

$$\pi_* : \mathrm{Sp} \longrightarrow \mathrm{Ab}^{\mathbb{Z}}$$

admits a lax symmetric monoidal refinement when the target is equipped with the Koszul symmetry. Indeed, for all $n, m \in \mathbb{Z}$ the tensor product of representatives defines natural pairings

$$\pi_n(X) \otimes \pi_m(Y) \longrightarrow \pi_{n+m}(X \otimes Y), \quad [x] \otimes [y] \longmapsto [\mathbb{S}^{n+m} \simeq \mathbb{S}^n \otimes \mathbb{S}^m \xrightarrow{x \otimes y} X \otimes Y].$$

Associativity and unitality are inherited from the symmetric monoidal structure on $\mathbf{Sp}$. Interchanging x and y introduces the symmetry

$$\mathbb{S}^n \otimes \mathbb{S}^m \longrightarrow \mathbb{S}^m \otimes \mathbb{S}^n,$$

which acts by $(-1)^{nm}$ on $\mathbb{S}^{n+m}$. For $n, m \geq 0$, this is the sign of the block permutation interchanging n and m suspension coordinates; the general case follows by invertibility of suspension. This iterates the basic sign calculation of Warning 4.2.11. Thus the symmetry compatibility is exactly the Koszul rule. Associativity, unitality, and symmetry give all coherence conditions because the target is a 1-category, so the pairings assemble into the claimed lax symmetric monoidal structure.

With the ordinary, unsigned Day symmetry, the same construction is only lax monoidal. Consequently, the homotopy groups of an associative ring spectrum form a graded ring, while those of a commutative ring spectrum form a graded-commutative ring:

$$xy = (-1)^{nm}yx \quad (x \in \pi_n R, y \in \pi_m R).$$

Exercises

17 The envelope of an ∞ -operad

Recall from Section 12.3 that every colored operad $\mathcal{O}$ has an *envelope* $\text{Env}(\mathcal{O})$, a symmetric monoidal category equipped with a symmetric monoidal functor to $\mathbf{Fin}$ which captures the operad structure of $\mathcal{O}$. Also recall from Exercise 12.11 that the construction $\mathcal{O} \mapsto \text{Env}(\mathcal{O})$ is left adjoint to the assignment $C \mapsto \mathcal{M}_C$ of the multimorphism operad to a symmetric monoidal category C . In particular, $\mathcal{O}$ -algebras in C correspond to symmetric monoidal functors $A: \text{Env}(\mathcal{O}) \rightarrow C$.

The goal of this chapter is to introduce the analogous envelope construction for ∞ -operads, and show that there exists a similar adjunction

$$\text{Env}: \text{Op}_\infty \rightleftarrows \text{Cat}_\infty^\otimes : \mathcal{M}.$$

We will further show that the envelope construction induces a fully faithful functor $\text{Env}: \text{Op}_\infty \hookrightarrow (\text{Cat}_\infty^\otimes)_{/(\mathbf{Fin}, \amalg)}$ and characterize its essential image. Here $(\mathbf{Fin}, \amalg)$ is the category of finite sets, with the symmetric monoidal structure given by disjoint union.

The envelope $\text{Env}(\mathcal{O})$ of an ∞ -operad $\mathcal{O}$ is obtained by considering the functor $\mathcal{O}^\otimes \rightarrow \text{Span}(\mathbf{Fin})$ and ‘freely adding’ cocartesian lifts of all forward spans. Since $\mathcal{O}^\otimes$ already admits cocartesian lifts of backwards spans, the resulting functor to $\text{Span}(\mathbf{Fin})$ will be a cocartesian fibration, and one can show that its straightening $\text{Span}(\mathbf{Fin}) \rightarrow \text{Cat}_\infty$ preserves finite products, thus encoding a symmetric monoidal ∞ -category $\text{Env}(\mathcal{O})$.

Let us preview the resulting symmetric monoidal ∞ -category before developing the categorical machinery needed to construct it. Its underlying ∞ -category is the pullback

$$\text{Env}(\mathcal{O}) \simeq \mathbf{Fin} \times_{\text{Span}(\mathbf{Fin})} \mathcal{O}^\otimes,$$

where $\mathbf{Fin} \hookrightarrow \text{Span}(\mathbf{Fin})$ is the subcategory of forward spans. Thus the objects of $\text{Env}(\mathcal{O})$ are finite unordered tuples of colors, the tensor product is concatenation, and the hom anima are given by

$$\text{Hom}_{\text{Env}(\mathcal{O})}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J}) \simeq \coprod_{f: I \rightarrow J} \prod_{j \in J} \mathcal{O}(\{x_i\}_{i \in f^{-1}(j)}; y_j).$$

This formula, proved in Lemma 17.3.17, makes precise the sense in which the envelope retains exactly the operations of $\mathcal{O}$ while freely adjoining tensor products of objects.

It will be convenient to construct envelopes in a slightly more general setting, in particular encompassing Lurie’s definition of ∞ -operads as well. The same procedure will then produce an analogous functor $\text{Env}^{\text{Lurie}}: \text{Op}_\infty^{\text{Lurie}} \rightarrow \text{Cat}_\infty^\otimes$, which is compatible with the functor Env , in the sense that the

following diagram commutes:

$$\begin{array}{ccc}
 \mathrm{Op}_\infty & \xrightarrow{\mathrm{Env}} & (\mathrm{Cat}_\infty^\otimes)/(\mathrm{Fin}, \mathbb{I}) \\
 \downarrow -\times_{\mathrm{Span}(\mathrm{Fin})} \mathrm{Fin}_* & & \uparrow \mathrm{Env}^{\mathrm{Lurie}} \\
 \mathrm{Op}_\infty^{\mathrm{Lurie}} & &
 \end{array}$$

We will see that the functor $\mathrm{Env}^{\mathrm{Lurie}}$ is also fully faithful when sliced over $(\mathrm{Fin}, \mathbb{I})$, and that it has the same image as Env . In particular, the comparison functor $\mathrm{Op}_\infty \rightarrow \mathrm{Op}_\infty^{\mathrm{Lurie}}$ is an equivalence, making the span approach to ∞ -operads taken in this book compatible with the Lurie model used more commonly in the literature.

17.1 Factorization systems

We begin with the notion of a *factorization system* on an ∞ -category. The ∞ -categorical formulation is due to Joyal and generalizes the familiar decomposition of maps of sets into a surjection followed by an injection. Factorization systems are an important categorical structure in their own right, so we develop their basic theory in some detail. They also provide exactly the mechanism needed later in this chapter: an ∞ -operad already has cocartesian lifts over one class of a factorization system, and its envelope is obtained by freely adjoining lifts over the other class.

Orthogonality

Definition 17.1.1 (Orthogonality). Let C be an ∞ -category. A morphism $l: A \rightarrow B$ is said to be *left orthogonal* to a morphism $r: X \rightarrow Y$, written $l \perp r$, if the following square is a pullback square:

$$\begin{array}{ccc}
 \mathrm{Hom}_C(B, X) & \xrightarrow{- \circ l} & \mathrm{Hom}_C(A, X) \\
 r \circ - \downarrow & & \downarrow r \circ - \\
 \mathrm{Hom}_C(B, Y) & \xrightarrow{- \circ l} & \mathrm{Hom}_C(A, Y).
 \end{array}$$

Equivalently, for every solid commutative square

$$\begin{array}{ccc}
 A & \xrightarrow{\quad} & X \\
 l \downarrow & \nearrow & \downarrow r \\
 B & \xrightarrow{\quad} & Y,
 \end{array}$$

the anima of dashed fillers making both triangles commute is contractible. We then also say that r is *right orthogonal* to l .

Given a collection of morphisms S , we denote by $S^\perp$ the collection of all morphisms that are right orthogonal to every morphism in S , and by ${}^\perp S$ the collection of those that are left orthogonal to every morphism in S .

Observation 17.1.2. Every morphism $f: X \rightarrow Y$ satisfying $f \perp f$ is an isomorphism in C : the unique dashed filler in the following diagram defines an inverse to f :

$$\begin{array}{ccc} X & \longrightarrow & X \\ f \downarrow & \nearrow \text{dashed} & \downarrow f \\ Y & \longrightarrow & Y. \end{array}$$

These orthogonal classes satisfy a number of useful closure properties.

Proposition 17.1.3. *Let S be a collection of morphisms in an ∞ -category C .*

- (1) *The classes $S^\perp$ and ${}^\perp S$ are closed under composition and contain all equivalences. In particular, they are encoded by wide subcategories of C .*
- (2) *The classes $S^\perp$ and ${}^\perp S$ are closed under retracts.*
- (3) *The class $S^\perp$ is closed under base change (pullbacks).*
- (4) *The class ${}^\perp S$ is closed under cobase change (pushouts).*
- (5) *The class $S^\perp$ has the left cancellation property: if $g \circ f \in S^\perp$ and $g \in S^\perp$, then $f \in S^\perp$.*
- (6) *The class ${}^\perp S$ has the right cancellation property: if $g \circ f \in {}^\perp S$ and $f \in {}^\perp S$, then $g \in {}^\perp S$.*
- (7) *If C admits small limits, the class $S^\perp$ is closed under small limits in $\text{Ar}(C)$.*
- (8) *If C admits small colimits, the class ${}^\perp S$ is closed under small colimits in $\text{Ar}(C)$.*

Proof. These properties all follow from the definition of orthogonality in terms of hom animaes. The proofs of (1) and (2) are left to the reader. For (3), consider a pullback square in C where $r \in S^\perp$:

$$\begin{array}{ccc} X' & \longrightarrow & X \\ r' \downarrow & \lrcorner & \downarrow r \\ Y' & \longrightarrow & Y. \end{array}$$

For any $l: A \rightarrow B$ in S , we must show $l \perp r'$. This means showing that the left square in the diagram below is a pullback square of animaes:

$$\begin{array}{ccccc} \text{Hom}_C(B, X') & \longrightarrow & \text{Hom}_C(A, X') & \longrightarrow & \text{Hom}_C(A, X) \\ r' \circ - \downarrow & & \downarrow r' \circ - & & \downarrow r \circ - \\ \text{Hom}_C(B, Y') & \longrightarrow & \text{Hom}_C(A, Y') & \longrightarrow & \text{Hom}_C(A, Y). \end{array}$$

The right square is a pullback because $\text{Hom}_C(A, -)$ preserves pullbacks, so by the pasting lemma for pullback squares it remains to show that the outer rectangle is a pullback. But this holds as we may rewrite it as the pasting of the square obtained from the original pullback square by applying $\text{Hom}_C(B, -)$ and the pullback square exhibiting the orthogonality relation $l \perp r$. The argument for (4) is dual.

For (5), consider morphisms $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ in C , and assume that $g \circ f$ and g lie in $S^\perp$. To show that $f \in S^\perp$, let $l: A \rightarrow B$ be a morphism in S , and consider the following commutative diagram:

$$\begin{array}{ccccc} \mathrm{Hom}_C(B, X) & \xrightarrow{f \circ -} & \mathrm{Hom}_C(B, Y) & \xrightarrow{g \circ -} & \mathrm{Hom}_C(B, Z) \\ \downarrow - \circ l & & \downarrow - \circ l & & \downarrow - \circ l \\ \mathrm{Hom}_C(A, X) & \xrightarrow{f \circ -} & \mathrm{Hom}_C(A, Y) & \xrightarrow{g \circ -} & \mathrm{Hom}_C(A, Z). \end{array}$$

By assumption, the right and outer squares are pullback squares, hence by the pasting law so is the left square, as was to be shown. The argument for (6) is dual.

For (7), let $f: I \rightarrow \mathrm{Ar}(C)$ be a diagram where each morphism $f(i): X(i) \rightarrow Y(i)$ lies in $S^\perp$. Let $r: \lim_i X(i) \rightarrow \lim_i Y(i)$ be the limit morphism. For any $l: A \rightarrow B$ in S , we have a chain of equivalences:

$$\begin{aligned} \mathrm{Hom}_C(B, \lim_i X(i)) &\simeq \lim_i \mathrm{Hom}_C(B, X(i)) \\ \mathrm{Hom}_C(A, \lim_i X(i)) &\simeq \lim_i \mathrm{Hom}_C(A, X(i)) \\ \mathrm{Hom}_C(B, \lim_i Y(i)) &\simeq \lim_i \mathrm{Hom}_C(B, Y(i)) \\ \mathrm{Hom}_C(A, \lim_i Y(i)) &\simeq \lim_i \mathrm{Hom}_C(A, Y(i)). \end{aligned}$$

The square defining the orthogonality $l \perp r$ is the limit of the squares defining $l \perp f(i)$ for each $i \in I$. Since limits of pullback squares are pullback squares, the claim follows. The proof for (8) is dual. $\square$

Factorization systems

We now come to the central notion of this section.

Definition 17.1.4 (Factorization system). A *factorization system* on an ∞ -category C consists of a pair of wide subcategories $(\mathcal{L}, \mathcal{R})$ of C satisfying the following conditions:

- (1) *Orthogonality*: Every morphism $l \in \mathcal{L}$ is left orthogonal to every morphism $r \in \mathcal{R}$.
- (2) *Factorization*: Every morphism f in C admits a factorization $f \simeq r \circ l$, where $l \in \mathcal{L}$ and $r \in \mathcal{R}$.

Three fundamental examples of factorization systems are developed in Exercises 17.1 to 17.3 at the end of the chapter.

Proposition 17.1.5 ([ABFJ22, Lemma 3.1.9]). *If $(\mathcal{L}, \mathcal{R})$ is a factorization system on C , then $\mathcal{L} = {}^\perp \mathcal{R}$ and $\mathcal{R} = \mathcal{L}^\perp$.*

Proof. By symmetry, it suffices to prove $\mathcal{R} = \mathcal{L}^\perp$. The inclusion $\mathcal{R} \subseteq \mathcal{L}^\perp$ holds by definition. For the converse, let $r: X \rightarrow Y$ be a morphism in $\mathcal{L}^\perp$. By the factorization property, we can write $r \simeq r' \circ l$ for some $l: X \rightarrow Z$ in $\mathcal{L}$ and $r': Z \rightarrow Y$ in $\mathcal{R}$.

Since $l \in \mathcal{L}$ and $r \in \mathcal{L}^\perp$, we have $l \perp r$. This means there is a contractible anima of lifts in the commutative square

$$\begin{array}{ccc} X & \xrightarrow{\text{id}_X} & X \\ l \downarrow & \nearrow d & \downarrow r \\ Z & \xrightarrow{r'} & Y. \end{array}$$

Let $d: Z \rightarrow X$ be such a lift. The commutativity of the top triangle gives $d \circ l \simeq \text{id}_X$. To see that also $l \circ d \simeq \text{id}_Z$, consider the following lifting problem for $l \perp r'$:

$$\begin{array}{ccc} X & \xrightarrow{l} & Z \\ l \downarrow & & \downarrow r' \\ Z & \xrightarrow{r'} & Y. \end{array}$$

It is clear that $\text{id}_Z: Z \rightarrow Z$ is a filler. But also the morphism $l \circ d: Z \rightarrow Z$ is a filler, since $r' \circ (l \circ d) \simeq (r' \circ l) \circ d \simeq r \circ d \simeq r'$ and $(l \circ d) \circ l \simeq l \circ (d \circ l) \simeq l$. By uniqueness of fillers, we must have $l \circ d \simeq \text{id}_Z$.

We conclude that l is an isomorphism, hence in $\mathcal{R}$. Since $r \simeq r' \circ l$ and $\mathcal{R}$ is closed under composition, we conclude that also $r \in \mathcal{R}$. $\square$

Remark 17.1.6. By combining the previous proposition with Proposition 17.1.3, we conclude that the class $\mathcal{R}$ is automatically closed under retracts, base change, and left cancellation. Similarly, $\mathcal{L}$ is closed under retracts, cobase change, and right cancellation.

In the literature, the condition that $\mathcal{L}$ and $\mathcal{R}$ need to be closed under retracts is often taken as part of the definition of a factorization system, but the previous proposition shows this assumption is redundant.

In a factorization system, the factorization of a morphism is unique. To formulate a precise statement, we denote by $\text{Fact}_{\mathcal{L},\mathcal{R}}(C)$ the full subcategory of $\text{Fun}([2], C)$ spanned by diagrams $X \rightarrow Y \rightarrow Z$ where the morphism $X \rightarrow Y$ lies in $\mathcal{L}$ and the morphism $Y \rightarrow Z$ lies in $\mathcal{R}$. Restriction along the map $d_1: [1] \rightarrow [2]$ provides a ‘composition functor’ $-\circ -: \text{Fact}_{\mathcal{L},\mathcal{R}}(C) \rightarrow \text{Ar}(C)$.

Proposition 17.1.7 ([Lur09b, Proposition 5.2.8.17]). *Let $\mathcal{L}$ and $\mathcal{R}$ be wide subcategories of an ∞ -category C . Then the following two conditions are equivalent:*

- (1) *The pair $(\mathcal{L}, \mathcal{R})$ is a factorization system on C ;*
- (2) *The composition functor $-\circ -: \text{Fact}_{\mathcal{L},\mathcal{R}}(C) \rightarrow \text{Ar}(C)$ is an equivalence.*

Proof. Assume first that $(\mathcal{L}, \mathcal{R})$ is a factorization system. The existence of factorizations implies that $-\circ -$ is essentially surjective, so it remains to show it is also fully faithful. To this end, consider morphisms $l: X \rightarrow Z$ and $l': X' \rightarrow Z'$ in $\mathcal{L}$, and morphisms $r: Z \rightarrow Y$ and $r': Z' \rightarrow Y'$ in $\mathcal{R}$, and define $u := r \circ l$ and $u' := r' \circ l'$. We must show that the map

$$\text{Hom}_{\text{Fact}_{\mathcal{L},\mathcal{R}}(C)}((l, r), (l', r')) \rightarrow \text{Hom}_{\text{Ar}(C)}(u, u')$$

induced by composition in C is an equivalence of anima. Equivalently, we may show that the fiber over every morphism $u \rightarrow u'$ in $\text{Ar}(C)$ is contractible. Such a morphism consists of maps

$a: X \rightarrow X'$ and $b: Y \rightarrow Y'$ in C , together with a homotopy filling the outer rectangle in the following diagram:

$$\begin{array}{ccc} X & \xrightarrow{a} & X' \\ l \downarrow & & \downarrow l' \\ Z & \dashrightarrow & Z' \\ r \downarrow & & \downarrow r' \\ Y & \xrightarrow{b} & Y' \end{array}$$

Using the commutative-square and Segal axioms, we then compute that the fiber under consideration agrees with the fiber over $(l'a: X \rightarrow Z', br: Z \rightarrow Y')$ of the map

$$(l^*, r'_*): \text{Hom}_C(Z, Z') \rightarrow \text{Hom}_C(X, Z') \times_{\text{Hom}_C(X, Y')} \text{Hom}_C(Z, Y').$$

But this map is an equivalence, since $l \perp r'$.

The proof of the converse is similar: if the composition functor is an equivalence, then essential surjectivity provides the desired factorizations of morphisms in C , while full faithfulness applied to the objects $(l: A \rightarrow B, \text{id}_B: B \rightarrow B)$ and $(\text{id}_X: X \rightarrow X, r: X \rightarrow Y)$ of $\text{Fact}_{\mathcal{L}, \mathcal{R}}(C)$ provides the orthogonality condition. $\square$

Corollary 17.1.8. *Let $(\mathcal{L}, \mathcal{R})$ be a factorization system on C . Then for every morphism f of C , the fiber of $\text{Fact}_{\mathcal{L}, \mathcal{R}}(C) \rightarrow \text{Ar}(C)$ over f is contractible.* $\square$

Lemma 17.1.9. *Let $(\mathcal{L}, \mathcal{R})$ be a factorization system on C , and let $\text{Ar}_{\mathcal{R}}(C) \subseteq \text{Ar}(C)$ denote the full subcategory spanned by the morphisms in $\mathcal{R}$. Then the target functor $t: \text{Ar}_{\mathcal{R}}(C) \rightarrow C$ is a cocartesian fibration, with cocartesian morphisms given by those commutative squares*

$$\begin{array}{ccc} x & \xrightarrow{f} & x' \\ r \downarrow & & \downarrow r' \\ y & \xrightarrow{g} & y' \end{array}$$

such that $f \in \mathcal{L}$.

Proof. Given r and g , factor the composite gr as $r'f$ with $f \in \mathcal{L}$ and $r' \in \mathcal{R}$. This provides a lift of g of the displayed form. We show that every such lift is t -cocartesian.

Consider a third object $(r'': x'' \rightarrow y'')$ in $\text{Ar}_{\mathcal{R}}(C)$. We must show that the square

$$\begin{array}{ccc} \text{Hom}_{\text{Ar}_{\mathcal{R}}(C)}(r', r'') & \longrightarrow & \text{Hom}_{\text{Ar}_{\mathcal{R}}(C)}(r, r'') \\ t \downarrow & & \downarrow t \\ \text{Hom}_C(y', y'') & \xrightarrow{- \circ g} & \text{Hom}_C(y, y'') \end{array}$$

is a pullback square. Fix a morphism $g': y' \rightarrow y''$ and a morphism $(h, g'g): r \rightarrow r''$ over $g'g$. The fiber in question is the anima of maps $k: x' \rightarrow x''$ fitting into the diagram

$$\begin{array}{ccc} x & \xrightarrow{h} & x'' \\ f \downarrow & \nearrow k & \downarrow r'' \\ x' & \xrightarrow{g'r'} & y'' \end{array}$$

Indeed, the two required triangles say precisely that $kf \simeq h$ and $r''k \simeq g'r'$. This filler anima is contractible because $f \perp r''$. It follows that the displayed square of hom animae is a pullback, so the original morphism is t -cocartesian.

Conversely, every t -cocartesian lift of g with source r is isomorphic to one constructed by factoring gr . Its top morphism is therefore the composite of a morphism in $\mathcal{L}$ with an isomorphism, and hence lies in $\mathcal{L}$. $\square$

The inert-active factorization system on an ∞ -operad

In Remark 14.1.11, we saw that every morphism in the total category $O^\otimes$ of an ∞ -operad factors uniquely as an inert morphism followed by an active morphism. We will now make this precise, by showing that the inert and active morphisms define a factorization system on $O^\otimes$.

As our first step, we will show that every span category admits a canonical factorization system. Recall from Lemma 13.1.16 that for an adequate triple (C, C_L, C_R) , the ∞ -category $\text{Span}_{L,R}(C)$ contains C_R and C_L^{op} as wide subcategories. The morphisms in C_R are called *forwards maps*, while those in C_L^{op} are called *backwards maps*.

Proposition 17.1.10 (The factorization system on a span category, [HHLN23b, Proposition 4.9]). *Let (C, C_L, C_R) be an adequate triple. The pair $(\mathcal{L}, \mathcal{R}) = (C_L^{\text{op}}, C_R)$ forms a factorization system on $\text{Span}_{L,R}(C)$.*

Proof. Any span $X \xleftarrow{l} U \xrightarrow{r} Y$ can be factored as a backwards map followed by a forwards map:

$$X \xleftarrow{l} U \xrightarrow{\text{id}_U} U \quad \text{followed by} \quad U \xleftarrow{\text{id}_U} U \xrightarrow{r} Y.$$

It remains to check orthogonality. Let $l': B \rightarrow A$ be a backwards map (given by $l \in C_L$) and $r': X \rightarrow Y$ be a forwards map (given by $r \in C_R$). We need to show that the square of hom animae

$$\begin{array}{ccc} \text{Hom}_{\text{Span}_{L,R}(C)}(A, X) & \longrightarrow & \text{Hom}_{\text{Span}_{L,R}(C)}(A, Y) \\ \downarrow & & \downarrow \\ \text{Hom}_{\text{Span}_{L,R}(C)}(B, X) & \longrightarrow & \text{Hom}_{\text{Span}_{L,R}(C)}(B, Y) \end{array}$$

is a pullback. Using the formula for hom animae in a span category (see Lemma 13.1.13), this square is equivalent to

$$\begin{array}{ccc} ((C_L)_A)^{\simeq} \times_{C^{\simeq}} ((C_R)_X)^{\simeq} & \xrightarrow{\text{id} \times (r \circ -)} & ((C_L)_A)^{\simeq} \times_{C^{\simeq}} ((C_R)_Y)^{\simeq} \\ (l \circ -) \times \text{id} \downarrow & & \downarrow (l \circ -) \times \text{id} \\ ((C_L)_B)^{\simeq} \times_{C^{\simeq}} ((C_R)_X)^{\simeq} & \xrightarrow{\text{id} \times (r \circ -)} & ((C_L)_B)^{\simeq} \times_{C^{\simeq}} ((C_R)_Y)^{\simeq}. \end{array}$$

The square involving only C_L is degenerate, hence a pullback square, and similarly for the square involving C_R . We conclude that this square is a pullback square, finishing the proof. $\square$

Remark 17.1.11. The unique filler in a lifting problem in $\text{Span}_{L,R}(C)$ can be visualized directly. A commutative square with a backwards map on the left and a forwards map on the right corresponds

to diagram of spans in C as follows:

$$\begin{array}{ccccc}
 A & \longleftarrow & Z & \longrightarrow & X \\
 \uparrow l & & \uparrow & \lrcorner & \parallel \\
 B & \longleftarrow & Z' & \longrightarrow & X \\
 \parallel & & \downarrow \lrcorner & & \downarrow r \\
 B & \longleftarrow & Z'' & \longrightarrow & Y.
 \end{array}$$

Here the composite span is $A \leftarrow Z' \rightarrow Y$, computed both as the composite of the spans $A \xleftarrow{l} B \xrightarrow{\text{id}_B} B$ and $B \leftarrow Z'' \rightarrow Y$, as well as the composite of the spans $A \leftarrow Z \rightarrow X$ and $X \xleftarrow{\text{id}_X} X \xrightarrow{r} Y$. The unique filler span is then given by the span $B \leftarrow Z' \rightarrow X$.

To obtain a factorization system on $O^\otimes$ from the factorization system on $\text{Span}(\text{Fin})$, we will make use of the following result.

Lemma 17.1.12. *Let $(\mathcal{L}, \mathcal{R})$ be a factorization system on an ∞ -category C , and let $p: E \rightarrow C$ be a functor such that for every morphism $l: A \rightarrow B$ in $\mathcal{L}$ and every object $\tilde{A} \in E_A$, there exists a p -cocartesian lift $\tilde{A} \rightarrow \tilde{B}$ of l . Then E admits a factorization system $(\tilde{\mathcal{L}}, \tilde{\mathcal{R}})$, where $\tilde{\mathcal{L}}$ consists of the p -cocartesian morphisms over $\mathcal{L}$ and $\tilde{\mathcal{R}}$ consists of morphisms over $\mathcal{R}$.*

Proof. Both classes define wide subcategories of E : they contain all isomorphisms, and they are closed under composition because $\mathcal{L}$ and $\mathcal{R}$ are, and because composites of p -cocartesian morphisms are again p -cocartesian.

(1) For orthogonality, consider $\tilde{l}: \tilde{A} \rightarrow \tilde{B}$ in $\tilde{\mathcal{L}}$ and $\tilde{r}: \tilde{X} \rightarrow \tilde{Y}$ in $\tilde{\mathcal{R}}$. Let $l: A \rightarrow B$ and $r: X \rightarrow Y$ be their images in C . We need to show that the top square in the following commutative cube is a pullback square:

$$\begin{array}{ccccc}
 \text{Hom}_E(\tilde{B}, \tilde{X}) & \xrightarrow{\tilde{r} \circ -} & \text{Hom}_E(\tilde{B}, \tilde{Y}) & & \\
 \downarrow p & \searrow - \circ \tilde{l} & \downarrow p & \searrow - \circ \tilde{l} & \\
 & \text{Hom}_E(\tilde{A}, \tilde{X}) & \xrightarrow{\tilde{r} \circ -} & \text{Hom}_E(\tilde{A}, \tilde{Y}) & \\
 & \downarrow p & \downarrow p & & \\
 \text{Hom}_C(B, X) & \xrightarrow{r \circ -} & \text{Hom}_C(B, Y) & & \\
 & \searrow - \circ l & \downarrow p & \searrow - \circ l & \\
 & \text{Hom}_C(A, X) & \xrightarrow{r \circ -} & \text{Hom}_C(A, Y) &
 \end{array}$$

Since $\tilde{l}$ is p -cocartesian, the left and right squares are pullback squares, and since $l \perp r$ the bottom square is a pullback square. The claim thus follows from the pasting law of pullback squares.

(2) For factorization, consider a morphism $\tilde{f}: \tilde{X} \rightarrow \tilde{Z}$. We may factor its image $X \rightarrow Z$ in C as a composite of a morphism $l: X \rightarrow Y$ in $\mathcal{L}$ and a morphism $r: Y \rightarrow Z$ in $\mathcal{R}$. We may then take $\tilde{l}: \tilde{X} \rightarrow \tilde{Y}$ to be a p -cocartesian lift of l , which exists by assumption. The cocartesian property produces a unique morphism $\tilde{r}: \tilde{Y} \rightarrow \tilde{Z}$ over r through which $\tilde{f}$ factors, as desired. $\square$

Corollary 17.1.13. *Let O be an ∞ -operad. Then the inert and active morphisms define a factorization system on $O^\otimes$.*

Proof. This is an instance of Lemma 17.1.12 applied to the functor $p_O: O^\otimes \rightarrow \text{Span}(\text{Fin})$, where we equip the target with the factorization system from Proposition 17.1.10. $\square$

Combining Corollaries 17.1.8 and 17.1.13, we conclude that the anima of inert-active factorizations of every morphism in $O^\otimes$ is contractible. This makes precise the uniqueness assertion from Remark 14.1.11.

17.2 Freely adjoining cocartesian lifts

The envelope of an operad will be a special case of a construction which freely adjoins cocartesian lifts. Throughout the section, we fix an ∞ -category S and a wide subcategory $\mathcal{R} \subset S$. We will show that for any functor $p: E \rightarrow S$, there exists a functor $E_{\mathcal{R}}(p) \rightarrow S$ equipped with a map from E over S which universally adds cocartesian lifts over the morphisms in $\mathcal{R}$. In case $\mathcal{R}$ is part of a factorization system $(\mathcal{L}, \mathcal{R})$ on S and p already admits cocartesian lifts over the morphisms of $\mathcal{L}$, we will show that $E_{\mathcal{R}}(p)$ is in fact a cocartesian fibration over S . Finally, we show that after slicing over the target projection $t: \text{Ar}_{\mathcal{R}}(S) \rightarrow S$, the free functor becomes fully faithful.

Our exposition in this section closely follows that from Barkan–Haugsgeng–Steinebrunner [BHS22, Section 2]. The free cocartesian completion has previously appeared in work of Gepner–Haugsgeng–Nikolaus [GHN17, Section 4], and the variant for factorization systems has appeared in work of Ayala–Mazel-Gee–Rozenblyum [AFMR17, Proposition A.0.1] and Shah [Sha21, Theorem 3.6].

Definition 17.2.1 ($\mathcal{R}$ -cocartesian fibrations). Let S be an ∞ -category and $\mathcal{R} \subset S$ a wide subcategory. A functor $p: E \rightarrow S$ is called $\mathcal{R}$ -cocartesian if for every $e \in E$ and every $r: p(e) \rightarrow y'$ in $\mathcal{R}$ there exists a p -cocartesian morphism $\hat{r}_e: e \rightarrow e'$ lifting r . A functor $f: E \rightarrow E'$ over S between $\mathcal{R}$ -cocartesian fibrations is $\mathcal{R}$ -cocartesian if it preserves p -cocartesian morphisms over morphisms in $\mathcal{R}$. We denote by

$$(\text{Cat}_{\infty})_{/S}^{\mathcal{R}\text{-cocart}} \subset (\text{Cat}_{\infty})_{/S}$$

the (non-full) subcategory spanned by the $\mathcal{R}$ -cocartesian fibrations and $\mathcal{R}$ -cocartesian functors.

Construction 17.2.2 (Freely adjoining $\mathcal{R}$ -cocartesian morphisms). Using the notation $\text{Ar}_{\mathcal{R}}(S)$ from Lemma 17.1.9, for any $p: E \rightarrow S$ we define

$$E_{\mathcal{R}}(p) := E \times_{S, S} \text{Ar}_{\mathcal{R}}(S) \xrightarrow{t \circ \text{pr}_2} S.$$

This construction defines an endofunctor $E_{\mathcal{R}}: (\text{Cat}_{\infty})_{/S} \rightarrow (\text{Cat}_{\infty})_{/S}$.

Lemma 17.2.3. *For every functor $p: E \rightarrow S$, the functor $E_{\mathcal{R}}(p)$ is an $\mathcal{R}$ -cocartesian fibration, and for every functor $f: E \rightarrow E'$ over S with structure map $p': E' \rightarrow S$, the induced functor $E_{\mathcal{R}}(f): E_{\mathcal{R}}(p) \rightarrow E_{\mathcal{R}}(p')$ over S is an $\mathcal{R}$ -cocartesian functor.*

Proof. First note that the target functor $t: \text{Ar}_{\mathcal{R}}(S) \rightarrow S$ is an $\mathcal{R}$ -cocartesian fibration: given an object $(r: x \rightarrow y) \in \text{Ar}_{\mathcal{R}}(S)$ (i.e., a morphism in $\mathcal{R}$) and a morphism $r': y \rightarrow y'$ in $\mathcal{R}$, a t -cocartesian lift of r' starting in r is given by the commutative square

$$\begin{array}{ccc} x & \xlongequal{\quad} & x \\ r \downarrow & & \downarrow r'r \\ y & \xrightarrow{r'} & y'. \end{array}$$

In particular, we see that a morphism in $\text{Ar}_{\mathcal{R}}(S)$ over $r' \in \mathcal{R}$ is t -cocartesian if and only if its image under the source functor $s: \text{Ar}_{\mathcal{R}}(S) \rightarrow S$ is an isomorphism in S .

The general case is similar: given an object $(e, r: x \rightarrow y)$ in $E_{\mathcal{R}}(p)$ and a morphism $r': y \rightarrow y'$ in $\mathcal{R}$, an $\mathcal{R}$ -cocartesian lift is given by the map $(e, r: x \rightarrow y) \rightarrow (e, r'r: x \rightarrow y')$ in $E_{\mathcal{R}}(p)$. Indeed, the hom anima in the pullback $E_{\mathcal{R}}(p)$ is the corresponding pullback of the hom animae in E and $\text{Ar}_{\mathcal{R}}(S)$, and this morphism is the identity on the E -component. Its cocartesian property therefore follows from that of the displayed morphism in $\text{Ar}_{\mathcal{R}}(S)$.

Note that these morphisms leave the E -component fixed. In particular, it is clear that the functor $E_{\mathcal{R}}(f): E_{\mathcal{R}}(p) \rightarrow E_{\mathcal{R}}(p')$ over S is an $\mathcal{R}$ -cocartesian functor for every $f: E \rightarrow E'$ over S . $\square$

Since $\mathcal{R}$ contains every identity map, restriction along $[1] \rightarrow *$ induces a fully faithful functor $i: S \rightarrow \text{Ar}_{\mathcal{R}}(S)$, given by $x \mapsto \text{id}_x$. The target and source functors are respectively a left and a right adjoint to i :

$$t \dashv i \dashv s.$$

The unit $\text{id} \rightarrow i \circ t$ of the first adjunction and the counit $i \circ s \rightarrow \text{id}$ of the second are given at a morphism $r: x \rightarrow y$ by the squares

$$\begin{array}{ccc} x & \xrightarrow{r} & y \\ r \downarrow & & \parallel \\ y & \xrightarrow{\quad} & y \end{array} \quad \text{and} \quad \begin{array}{ccc} x & \xrightarrow{\quad} & x \\ \parallel & & \downarrow r \\ x & \xrightarrow{r} & y, \end{array}$$

respectively. Pulling back i also provides a fully faithful functor

$$i_p: E \simeq E \times_S S \xrightarrow{E \times_S i} E \times_S \text{Ar}_{\mathcal{R}}(S) = E_{\mathcal{R}}(p)$$

over S , for every $p: E \rightarrow S$. Note that this is natural in p , in the sense of defining a natural transformation $\text{id} \rightarrow E_{\mathcal{R}}$ of endofunctors of $(\text{Cat}_{\infty})/S$.

Lemma 17.2.4. *If $p: E \rightarrow S$ is an $\mathcal{R}$ -cocartesian fibration, the functor $i_p: E \rightarrow E_{\mathcal{R}}(p)$ admits a left adjoint $\pi_p: E_{\mathcal{R}}(p) \rightarrow E$, given on objects by sending a pair $(e, r: x \rightarrow y)$ to the target e' of the p -cocartesian lift $\hat{r}_e: e \rightarrow e'$ of r .*

Proof. By the pointwise criterion for left adjoints, it suffices to show that for a fixed object (e, r) of $E_{\mathcal{R}}(p)$, the object $e' \in E$ is a left adjoint object to (e, r) under the functor i_p . Note that there is a canonical map $(e, r) \rightarrow (e', \text{id}_y) = i_p(e')$ given by the following diagram:

$$\begin{array}{ccc} e & \xrightarrow{\hat{r}_e} & e' \\ \Downarrow & & \Downarrow \\ x & \xrightarrow{r} & y \\ r \downarrow & & \parallel \\ y & \xrightarrow{\quad} & y. \end{array}$$

We need to show that for every other object $e'' \in E$ with $z := p(e'') \in S$, restriction along this map induces an equivalence

$$\text{Hom}_E(e', e'') \rightarrow \text{Hom}_{E_{\mathcal{R}}(p)}((e', \text{id}_y), (e'', \text{id}_z)) \rightarrow \text{Hom}_{E_{\mathcal{R}}(p)}((e, r: x \rightarrow y), (e'', \text{id}_z)).$$

Equivalently, we may show that the fiber over every morphism $(e, r: x \rightarrow y) \rightarrow (e'', \text{id}_z)$ in $E_{\mathcal{R}}(p)$ is contractible. By unwinding definitions, this is a consequence of the fact that there is a unique map $e' \rightarrow e''$ whose composite with the p -cocartesian morphism $\hat{r}_e: e \rightarrow e'$ is the given map $e \rightarrow e''$, with the resulting commutative triangle living over the triangle $x \rightarrow y \rightarrow z$ in S . $\square$

Remark 17.2.5. The last argument in the previous proof may be pictured as finding unique dashed arrows in the following solid commutative diagram:

$$\begin{array}{ccccc}
 e & \xrightarrow{\hat{r}_e} & e' & \dashrightarrow & e'' \\
 \downarrow & & \downarrow & & \downarrow \\
 x & \xrightarrow{r} & y & \dashrightarrow & z \\
 \downarrow r & & \parallel & & \parallel \\
 y & \xrightarrow{\quad} & y & \dashrightarrow & z
 \end{array}$$

It is clear that two dashed maps $y \rightarrow z$ have to be the given map $y \rightarrow z$ at the bottom. The uniqueness of the remaining map $e' \rightarrow e''$ is then a consequence of $\hat{r}_e$ being p -cocartesian.

Lemma 17.2.6. *If $q: F \rightarrow S$ is an $\mathcal{R}$ -cocartesian fibration, the functor $\pi_q: E_{\mathcal{R}}(q) \rightarrow F$ preserves $\mathcal{R}$ -cocartesian morphisms. Moreover, given a morphism $g: F \rightarrow F'$ in $(\text{Cat}_{\infty})_{/S}^{\mathcal{R}\text{-cocart}}$, the Beck–Chevalley transformation*

$$\pi_{q'} E_{\mathcal{R}}(g) \xrightarrow{\eta} \pi_{q'} E_{\mathcal{R}}(g) i_q \pi_q \simeq \pi_{q'} i_{q'} g \pi_q \xrightarrow{\varepsilon} g \pi_q$$

is a natural isomorphism. In particular, the following diagram commutes:

$$\begin{array}{ccc}
 E_{\mathcal{R}}(q) & \xrightarrow{E_{\mathcal{R}}(g)} & E_{\mathcal{R}}(q') \\
 \pi_q \downarrow & & \downarrow \pi_{q'} \\
 F & \xrightarrow{g} & F'
 \end{array}$$

Proof. First consider an $\mathcal{R}$ -cocartesian morphism

$$(e, r: x \rightarrow y) \longrightarrow (e, r'r: x \rightarrow y')$$

in $E_{\mathcal{R}}(q)$. The functor π_q sends it to the induced morphism $r_! e \rightarrow (r'r)_! e$. Both the composite $e \rightarrow r_! e \rightarrow (r'r)_! e$ and its first factor are q -cocartesian, so the second factor is q -cocartesian by Lemma 23.1.14. Thus π_q preserves $\mathcal{R}$ -cocartesian morphisms.

It suffices to show that the transformation is a pointwise isomorphism for every individual object $(e, r: x \rightarrow y)$ in $E_{\mathcal{R}}(q)$. Let $\hat{r}_e: e \rightarrow e'$ be a q -cocartesian lift of r in F , and let $\hat{r}_{g(e)}: g(e) \rightarrow e''$ be a q -cocartesian lift of r in F' . The universal property of $\hat{r}_{g(e)}$ then induces a map $e'' \rightarrow g(e')$. But since g preserves $\mathcal{R}$ -cocartesian morphisms, the map $g(e) \rightarrow g(e')$ is already an $\mathcal{R}$ -cocartesian morphism, hence the map $e'' \rightarrow g(e')$ is an isomorphism in F' . Unwinding the definitions of the counit of the adjunction $\pi_{q'} \dashv i_{q'}$ reveals that this map is precisely the map $(\pi_{q'} E_{\mathcal{R}}(g))(e, r) \rightarrow (g \pi_q)(e, r)$, finishing the proof. $\square$

We are now ready to prove the universal property of the $\mathcal{R}$ -cocartesian fibration $E_{\mathcal{R}}(p)$:

Given functors $p: E \rightarrow S$ and $q: F \rightarrow S$, we write $\text{Fun}_{/S}(E, F)$ for the fiber of $\text{Fun}(E, F) \rightarrow \text{Fun}(E, S)$ over p . If both functors are $\mathcal{R}$ -cocartesian, we write $\text{Fun}_{/S}^{\mathcal{R}\text{-cocart}}(E, F)$ for the full subcategory spanned by the functors which preserve cocartesian morphisms over $\mathcal{R}$.

Proposition 17.2.7 (Free adjunction of $\mathcal{R}$ -cocartesian lifts). *The functor*

$$E_{\mathcal{R}}: (\text{Cat}_{\infty})_{/S} \rightarrow (\text{Cat}_{\infty})_{/S}^{\mathcal{R}\text{-cocart}}$$

from Lemma 17.2.3 is left adjoint to the forgetful functor, with unit given by $i_p: E \rightarrow E_{\mathcal{R}}(p)$ and counit given by $\pi_q: E_{\mathcal{R}}(q) \rightarrow F$. More precisely, given a functor $p: E \rightarrow S$ and an $\mathcal{R}$ -cocartesian fibration $q: F \rightarrow S$, precomposition with i_p induces an equivalence

$$\text{Fun}_{/S}^{\mathcal{R}\text{-cocart}}(E_{\mathcal{R}}(p), F) \xrightarrow{\sim} \text{Fun}_{/S}(E, F),$$

with inverse sending $f: E \rightarrow F$ to $\pi_q \circ E_{\mathcal{R}}(f)$.

Proof. Consider first a morphism $f: E \rightarrow F$ in $(\text{Cat}_{\infty})_{/S}$. We must show that the composite

$$E \xrightarrow{i_p} E_{\mathcal{R}}(p) \xrightarrow{E_{\mathcal{R}}(f)} E_{\mathcal{R}}(q) \xrightarrow{\pi_q} F$$

is naturally isomorphic to f . To this end, observe that by naturality of the map $i_p: E \rightarrow E_{\mathcal{R}}(p)$ in p , the composite of the first two maps is given by $E \xrightarrow{f} F \xrightarrow{i_q} E_{\mathcal{R}}(q)$, naturally in f . The desired equivalence is then induced by the counit map $\pi_q \circ i_q \xrightarrow{\sim} \text{id}_F$ of the adjunction $\pi_q \dashv i_q$, which is an equivalence by full faithfulness of i_q .

Next, consider a morphism $g: E_{\mathcal{R}}(p) \rightarrow F$ in $(\text{Cat}_{\infty})_{/S}^{\mathcal{R}\text{-cocart}}$. We must show that the composite

$$E_{\mathcal{R}}(p) \xrightarrow{E_{\mathcal{R}}(i_p)} E_{\mathcal{R}}(E_{\mathcal{R}}(p)) \xrightarrow{E_{\mathcal{R}}(g)} E_{\mathcal{R}}(q) \xrightarrow{\pi_q} F$$

is naturally isomorphic to g . By Lemma 17.2.6, the composite of the last two maps is naturally equivalent to the composite $E_{\mathcal{R}}(E_{\mathcal{R}}(p)) \xrightarrow{\pi_{E_{\mathcal{R}}(p)}} E_{\mathcal{R}}(p) \xrightarrow{g} F$; note that the natural isomorphism given there is natural in g . The map $\pi_{E_{\mathcal{R}}(p)}$ is given by composition in $\mathcal{R}$: it sends a tuple $((e, r: x \rightarrow y), r': y \rightarrow z)$ to $(e, r'r: x \rightarrow z)$. In particular, if $r = \text{id}_y$ is the identity, this results in $(e, r': y \rightarrow z)$. It follows that the composite $\pi_{E_{\mathcal{R}}(p)} \circ E_{\mathcal{R}}(i_p)$ is naturally equivalent to $\text{id}_{E_{\mathcal{R}}(p)}$. This finishes the proof of the second statement of the proposition.

The first statement immediately follows from the second one by passing to groupoid cores on both sides, so that the left-hand side becomes $\text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{R}\text{-cocart}}}(E_{\mathcal{R}}(p), F)$, while the right-hand side becomes $\text{Hom}_{(\text{Cat}_{\infty})_{/S}}(E, F)$. $\square$

In case the wide subcategory $\mathcal{R}$ is part of a factorization system $(\mathcal{L}, \mathcal{R})$ on S , the functor $E_{\mathcal{R}}$ turns $\mathcal{L}$ -cocartesian fibrations into cocartesian fibrations:

Proposition 17.2.8. *Let $(\mathcal{L}, \mathcal{R})$ be a factorization system on S . If $p: E \rightarrow S$ is $\mathcal{L}$ -cocartesian, then $E_{\mathcal{R}}(p) \rightarrow S$ is a cocartesian fibration, with cocartesian morphisms given by those diagrams*

$$\begin{array}{ccc} e & \xrightarrow{\hat{f}} & e' \\ \downarrow & & \downarrow \\ x & \xrightarrow{f} & x' \\ r \downarrow & & \downarrow r' \\ y & \xrightarrow{g} & y' \end{array}$$

such that f lies in $\mathcal{L}$ and $\hat{f}$ is a p -cocartesian morphism in E . Similarly, if $g: E \rightarrow E'$ is an $\mathcal{L}$ -cocartesian functor over S , the induced map $E_{\mathcal{R}}(g)$ is a cocartesian functor over S .

Proof. We start by showing that $E_{\mathcal{R}}(p)$ is a cocartesian fibration: given an object $(e, r: x \rightarrow y)$ in $E_{\mathcal{R}}(p)$ and a morphism $g: y \rightarrow y'$ in S , we will construct a cocartesian lift over g . To this end, consider the unique factorization of $gr: x \rightarrow y'$ as $r'f$, where $f: x \rightarrow x'$ is in $\mathcal{L}$ and $r': x' \rightarrow y'$ is in $\mathcal{R}$. Also consider a p -cocartesian lift $\hat{f}_e: e \rightarrow e'$ of f starting in e ; this exists by assumption on E . We claim that the resulting morphism $(\hat{f}, f, g): (e, r) \rightarrow (e', r')$ in $E_{\mathcal{R}}(p)$ is a cocartesian morphism with respect to the map $t \text{pr}_2: E_{\mathcal{R}}(p) \rightarrow S$. In other words, we must show that for every third object $(e'', r'': x'' \rightarrow y'')$, the outer square in the commutative diagram

$$\begin{array}{ccc} \text{Hom}_{E_{\mathcal{R}}(p)}((e', r'), (e'', r'')) & \xrightarrow{-\circ(\hat{f}, f, g)} & \text{Hom}_{E_{\mathcal{R}}(p)}((e, r), (e'', r'')) \\ \text{pr}_2 \downarrow & & \downarrow \text{pr}_2 \\ \text{Hom}_{\text{Ar}_{\mathcal{R}}(S)}(r', r'') & \xrightarrow{-\circ(f, g)} & \text{Hom}_{\text{Ar}_{\mathcal{R}}(S)}(r, r'') \\ t \downarrow & & \downarrow t \\ \text{Hom}_S(y', y'') & \xrightarrow{-\circ g} & \text{Hom}_S(y, y'') \end{array}$$

is a pullback square. But this follows from the pasting law of pullback squares: the top square is a pullback square since $\hat{f}_e$ is p -cocartesian, and the bottom square is a pullback square since the morphism $(f, g): r \rightarrow r'$ is t -cocartesian by Lemma 17.1.9.

This shows that $E_{\mathcal{R}}(p)$ is a cocartesian fibration. By uniqueness of cocartesian lifts, it also confirms the description of the cocartesian morphisms. The fact that $E_{\mathcal{R}}(g)$ preserves cocartesian morphisms over S is clear from this description. $\square$

Remark 17.2.9. Diagrammatically, the fact that the morphism displayed in the previous proposition is cocartesian boils down to finding unique dashed morphisms in the following commutative diagram:

$$\begin{array}{ccccc} e & \xrightarrow{f} & e' & \dashrightarrow & e'' \\ \downarrow & & \downarrow & & \downarrow \\ x & \xrightarrow{f} & x' & \dashrightarrow & x'' \\ r \downarrow & & \downarrow r' & & \downarrow r'' \\ y & \xrightarrow{g} & y' & \xrightarrow{g'} & y'' \end{array}$$

The uniqueness of $x' \rightarrow x''$ making the bottom diagram commute is a consequence of the factorization system, while the uniqueness of $e' \rightarrow e''$ making the top square commute holds by cocartesianness of $\hat{f}$.

Corollary 17.2.10. *The functor $E_{\mathcal{R}}$ restricts to a functor*

$$E_{\mathcal{R}}: (\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}} \longrightarrow \text{Cocart}(S)$$

which is left adjoint to the inclusion.

Proof. By Proposition 17.2.8, $E_{\mathcal{R}}$ sends $\mathcal{L}$ -cocartesian fibrations to cocartesian fibrations, showing that it restricts at the level of objects. The characterization of the cocartesian morphisms from Proposition 17.2.8 further shows that if $f: E \rightarrow E'$ preserves $\mathcal{L}$ -cocartesian morphisms, then $E_{\mathcal{R}}(f)$ preserves cocartesian morphisms, showing that $E_{\mathcal{R}}$ also restricts at the level of morphisms.

To show the resulting functor is left adjoint to the inclusion, we must show that the unit and counit of the adjunction from Proposition 17.2.7 lie in the appropriate subcategories. Given an $\mathcal{L}$ -cocartesian fibration $p: E \rightarrow S$, it is clear from the description of the cocartesian morphisms given in Proposition 17.2.8 that the unit $i_p: E \rightarrow E_{\mathcal{R}}(p)$ preserves $\mathcal{L}$ -cocartesian morphisms. Similarly, given a cocartesian fibration $q: F \rightarrow S$, the counit $\pi_q: E_{\mathcal{R}}(q) \rightarrow F$ preserves cocartesian morphisms: it sends a cocartesian morphism as displayed in Proposition 17.2.8 to the map $r_!e \rightarrow r'_!e'$, which is q -cocartesian by the right cancellation property of q -cocartesian morphisms (see Lemma 23.1.14). $\square$

To end the section, we will show that the functor $E_{\mathcal{R}}$ from the corollary becomes fully faithful when slicing it over the cocartesian fibration $t: \text{Ar}_{\mathcal{R}}(S) \rightarrow S$. Note that the identity functor $\text{id}_S: S \rightarrow S$ is an $\mathcal{L}$ -cocartesian fibration, and is in fact the terminal object of $(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}$. Also note that the cocartesian fibration $t: \text{Ar}_{\mathcal{R}}(S) \rightarrow S$ is the image of id_S under $E_{\mathcal{R}}$. As a result, $E_{\mathcal{R}}$ induces a functor on slice categories

$$E_{\mathcal{R}}: (\text{Cat}_{\infty})_{/\text{id}_S}^{\mathcal{L}\text{-cocart}} \simeq ((\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}})_{/S} \rightarrow \text{Cocart}(S)_{/\text{Ar}_{\mathcal{R}}(S)}. \quad (17.1)$$

Lemma 17.2.11. *The functor (17.1) is fully faithful.*

Proof. Consider two $\mathcal{L}$ -cocartesian fibrations $p: E \rightarrow S$ and $p': E' \rightarrow S$. We must show that the induced diagram

$$\begin{array}{ccc} \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, E') & \longrightarrow & \text{Hom}_{\text{Cocart}(S)}(E_{\mathcal{R}}(p), E_{\mathcal{R}}(p')) \\ \downarrow & & \downarrow \\ * \simeq \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, S) & \longrightarrow & \text{Hom}_{\text{Cocart}(S)}(E_{\mathcal{R}}(p), \text{Ar}_{\mathcal{R}}(S)) \end{array}$$

is a pullback square, where the horizontal maps are given by applying $E_{\mathcal{R}}$. By the universal property of $E_{\mathcal{R}}(p)$, we may identify this square with the square

$$\begin{array}{ccc} \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, E') & \longrightarrow & \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, E_{\mathcal{R}}(p')) \\ \downarrow & & \downarrow \\ \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, S) & \longrightarrow & \text{Hom}_{(\text{Cat}_{\infty})_{/S}^{\mathcal{L}\text{-cocart}}}(E, \text{Ar}_{\mathcal{R}}(S)), \end{array}$$

where now the horizontal maps are given by postcomposition with $i_{p'}: E' \rightarrow E_{\mathcal{R}}(p')$ and $i: S \rightarrow \text{Ar}_{\mathcal{R}}(S)$. But this is a simple consequence of the fact that the left square in the diagram

$$\begin{array}{ccccc} E' & \xrightarrow{i_{p'}} & E_{\mathcal{R}}(p') & \longrightarrow & E' \\ p' \downarrow & & \downarrow & \lrcorner & \downarrow p' \\ S & \xrightarrow{i} & \text{Ar}_{\mathcal{R}}(S) & \xrightarrow{s} & S \end{array}$$

is a pullback square by the pasting law, and that a functor into E' preserves $\mathcal{L}$ -cocartesian morphisms if and only if its composite with $i_{p'}$ does. $\square$

17.3 Operadic envelopes

We now construct the envelope of an ∞ -operad. As mentioned in the introduction to this chapter, it will be convenient to work in greater generality. Our approach is heavily inspired by [BHS22], in which the authors construct a very general envelope construction using a notion of *algebraic pattern*. To simplify the exposition, we will forgo some of the generality and only consider a special case of their setup.

Convention 17.3.1. Throughout this section, we fix an adequate triple $\mathcal{F} = (F, F_L, F_R)$ which satisfies the following three conditions:

- (1) The ∞ -category F is extensive, and the triple (F, F_L, F_R) is *weakly coextensive*, in the sense of Definition 13.3.1;
- (2) Morphisms in F_L satisfy left cancellation: given $l: I \rightarrow J$ and $l': J \rightarrow K$, if $l' \in F_L$ and $l'l \in F_L$ then also $l \in F_L$;
- (3) For every $I \in F$, the morphism $\emptyset \rightarrow I$ is contained in F_L .

Note that this in particular implies that F_L contains all inclusions $I_i \hookrightarrow \bigsqcup_{j=1}^n I_j$ for $I_1, \dots, I_n \in F$: by weak coextensivity, F_L is closed under finite coproducts, so the inclusion is the coproduct of id_{I_i} with the morphisms $\emptyset \rightarrow I_j$ for $j \neq i$.

Example 17.3.2. The main two examples are the adequate triples $(\text{Fin}, \text{Fin}, \text{Fin})$ and $(\text{Fin}, \text{Fin}_{\text{inj}}, \text{Fin})$, where $\text{Fin}_{\text{inj}} \subseteq \text{Fin}$ is the wide subcategory of injections.

These are the only two cases needed later in this book. The extra generality is nevertheless useful: it exhibits the envelope construction as a reusable feature of span patterns rather than a peculiarity of ordinary symmetric operads. The examples below indicate how the same method extends to settings arising in equivariant and global homotopy theory.

Example 17.3.3. More generally, let G be a finite group and let Fin_G be the category of finite G -sets. Then the adequate triples $(\text{Fin}_G, \text{Fin}_G, \text{Fin}_G)$ and $(\text{Fin}_G, (\text{Fin}_G)_{\text{inj}}, \text{Fin}_G)$ are examples.

Example 17.3.4. Let $\text{FinGrpd} \subseteq \text{Grpd} \subseteq \text{Cat}$ denote the full $(2,1)$ -category of finite groupoids. Under the fully faithful inclusion $\text{Grpd} \hookrightarrow \text{An}$, we may identify this with the full subcategory of animae spanned by objects of the form $\bigsqcup_{i=1}^n BG_i$, the finite disjoint unions of classifying animae of finite groups G_i . Then $(\text{FinGrpd}, \text{FinGrpd}, \text{FinGrpd})$ satisfies the conditions of Convention 17.3.1.

Definition 17.3.5 ($\mathcal{F}$ -monoid). Let $\mathcal{F} = (F, F_L, F_R)$ be an adequate triple satisfying the conditions of Convention 17.3.1.

- (1) Given an ∞ -category C with finite products, we say that a functor $M: F_L^{\text{op}} \rightarrow C$ satisfies the *Segal condition* if for objects $I_1, \dots, I_n \in F$, the maps $e_i: I_i \hookrightarrow \bigsqcup_{j=1}^n I_j$ induce an isomorphism

$$(e_i^*)_{i=1}^n: M(\bigsqcup_{j=1}^n I_j) \xrightarrow{\sim} \prod_{i=1}^n M(I_i).$$

- (2) For C with finite products, we define an $\mathcal{F}$ -monoid in C to be a functor

$$M: \text{Span}_{L,R}(F) \rightarrow C$$

such that $M|_{F_L^{\text{op}}}$ satisfies the Segal condition. We denote by

$$\text{Mon}_{\mathcal{F}}(C) \subseteq \text{Fun}(\text{Span}_{L,R}(F), C)$$

the full subcategory spanned by the $\mathcal{F}$ -monoids.

Remark 17.3.6. If F_L contains the fold maps $I \sqcup I \rightarrow I$, then the conditions of Convention 17.3.1 imply that (F, F_L, F_R) is weakly extensive. Indeed, finite coproducts and the required pullback squares follow from the extensivity of F , condition (2) in the definition of weak extensivity is also part of weak coextensivity, and condition (3) follows from the fold-map hypothesis together with part (3) of Convention 17.3.1. It follows that F_L admits finite coproducts, and a functor $F_L^{\text{op}} \rightarrow C$ satisfies the Segal condition if and only if it preserves finite products. Similarly, part (1) of Lemma 13.3.5 shows that $\text{Span}_{L,R}(F)$ admits finite products, and a functor $M: \text{Span}_{L,R}(F) \rightarrow C$ is an $\mathcal{F}$ -monoid if and only if it preserves finite products.

Example 17.3.7. For $\mathcal{F} = (\text{Fin}, \text{Fin}, \text{Fin})$, the $\mathcal{F}$ -monoids are precisely the commutative monoids from Definition 5.3.4.

For $\mathcal{F} = (\text{Fin}, \text{Fin}_{\text{inj}}, \text{Fin})$, the ∞ -category $\text{Span}_{\text{inj}, \text{all}}(\text{Fin})$ is equivalent to the (classical) category Fin_* of finite pointed sets, and the resulting $\mathcal{F}$ -monoids are precisely the commutative monoids as defined by Lurie [Lur17]. By Proposition 5.3.9, restriction along the inclusion induces an equivalence between these two definitions.

Definition 17.3.8. We define an $\mathcal{F}$ -monoidal ∞ -category to be an $\mathcal{F}$ -monoid in Cat_{∞} .

Remark 17.3.9. Using the straightening-unstraightening equivalence $\text{Un}^{\text{cc}}: \text{Fun}(\text{Span}_{L,R}(F), \text{Cat}_{\infty}) \xrightarrow{\sim} \text{Cocart}(\text{Span}_{L,R}(F))$, we may identify the category of $\mathcal{F}$ -monoidal ∞ -categories with a full subcategory of $\text{Cocart}(\text{Span}_{L,R}(F))$. We denote the unstraightening of an $\mathcal{F}$ -monoidal ∞ -category C by $p_C: C^{\otimes} \rightarrow \text{Span}_{L,R}(F)$.

In the following, we recall from Proposition 17.1.10 that the span category $\text{Span}_{L,R}(F)$ comes equipped with a canonical factorization system $(\mathcal{L}, \mathcal{R})$, where $\mathcal{L} = F_L^{\text{op}}$ consists of the backwards spans and $\mathcal{R} = F_R$ consists of the forwards spans.

Definition 17.3.10 ($\mathcal{F}$ -operad). Let $\mathcal{F} = (F, F_L, F_R)$ be an adequate triple satisfying the conditions of Convention 17.3.1. An $\mathcal{F}$ -operad is a pair $O = (O^{\otimes}, p_O)$ consisting of an ∞ -category $O^{\otimes}$ and a functor $p_O: O^{\otimes} \rightarrow \text{Span}_{L,R}(F)$ satisfying the following three conditions:

- (1) The functor p_O is an $\mathcal{L}$ -cocartesian fibration;
- (2) The straightening $F_L^{\text{op}} \rightarrow \text{Cat}_\infty$ of the resulting cocartesian fibration $O^\otimes|_{F_L^{\text{op}}} \rightarrow F_L^{\text{op}}$ satisfies the Segal condition: for all $J_1, \dots, J_n \in F$ we have

$$O^\otimes_{\sqcup_{i=1}^n J_i} \xrightarrow{\sim} \prod_{i=1}^n O^\otimes_{J_i}.$$

- (3) Consider objects $I, J_1, \dots, J_n \in F$, $X \in O_I^\otimes$ and $Y \in O_J^\otimes$ with $J := \sqcup_{i=1}^n J_i$. Let $Y \rightarrow Y_i$ be cocartesian lifts of the backwards maps $J \hookrightarrow J_i \xrightarrow{=} J_i$. Then the induced commutative square

$$\begin{array}{ccc} \text{Hom}_{O^\otimes}(X, Y) & \longrightarrow & \prod_{i=1}^n \text{Hom}_{O^\otimes}(X, Y_i) \\ p_O \downarrow & & \downarrow p_O \\ \text{Hom}_{\text{Span}_{L,R}(F)}(I, J) & \longrightarrow & \prod_{i=1}^n \text{Hom}_{\text{Span}_{L,R}(F)}(I, J_i) \end{array}$$

is a pullback square.

Given another $\mathcal{F}$ -operad $\mathcal{P} = (\mathcal{P}^\otimes, p_{\mathcal{P}})$, a *morphism of $\mathcal{F}$ -operads* is a $\mathcal{L}$ -cocartesian functor $f: O^\otimes \rightarrow \mathcal{P}^\otimes$ over $\text{Span}_{L,R}(F)$. We denote by

$$\text{Op}_{\mathcal{F}} \subseteq (\text{Cat}_\infty)^{\mathcal{L}\text{-cocart}}_{/\text{Span}_{L,R}(F)}$$

the full subcategory spanned by the $\mathcal{F}$ -operads.

Construction 17.3.11. We construct a functor

$$\mathcal{M}: \text{Mon}_{\mathcal{F}}(\text{Cat}_\infty) \rightarrow \text{Op}_{\mathcal{F}}.$$

Given an $\mathcal{F}$ -monoidal ∞ -category C , its unstraightening $p_C: C^\otimes \rightarrow \text{Span}_{L,R}(F)$ is an $\mathcal{F}$ -operad: conditions (1) and (2) are obvious, and (3) is proved just like in Lemma 14.2.4. Similarly, for an $\mathcal{F}$ -monoidal functor $C \rightarrow D$ (i.e. a morphism in $\text{Mon}_{\mathcal{F}}(\text{Cat}_\infty)$) the map on unstraightenings $C^\otimes \rightarrow D^\otimes$ is a cocartesian functor, hence in particular $\mathcal{L}$ -cocartesian. This shows that unstraightening defines the desired functor $\mathcal{M}$.

Proposition 17.3.12. *The functor $\mathcal{M}: \text{Mon}_{\mathcal{F}}(\text{Cat}_\infty) \rightarrow \text{Op}_{\mathcal{F}}$ admits a left adjoint*

$$\text{Env}_{\mathcal{F}}: \text{Op}_{\mathcal{F}} \rightarrow \text{Mon}_{\mathcal{F}}(\text{Cat}_\infty),$$

called the $\mathcal{F}$ -operadic envelope construction.

Proof. By definition, $\text{Op}_{\mathcal{F}}$ is a full subcategory of $(\text{Cat}_\infty)^{\mathcal{L}\text{-cocart}}_{/\text{Span}_{L,R}(F)}$. We will similarly identify $\text{Mon}_{\mathcal{F}}(\text{Cat}_\infty)$ with a full subcategory of $\text{Cocart}(\text{Span}_{L,R}(F))$, see Remark 17.3.9. Recall from Corollary 17.2.10 that the forgetful functor $\text{Cocart}(\text{Span}_{L,R}(F)) \hookrightarrow (\text{Cat}_\infty)^{\mathcal{L}\text{-cocart}}_{/\text{Span}_{L,R}(F)}$ admits a left adjoint

$$E_{\mathcal{R}}: (\text{Cat}_\infty)^{\mathcal{L}\text{-cocart}}_{/\text{Span}_{L,R}(F)} \rightarrow \text{Cocart}(\text{Span}_{L,R}(F)).$$

Since $\mathcal{M}$ is defined as the restriction of this forgetful functor, it remains to show that $E_{\mathcal{R}}$ restricts to a functor

$$\text{Env}_{\mathcal{F}}: \text{Op}_{\mathcal{F}} \rightarrow \text{Mon}_{\mathcal{F}}(\text{Cat}_\infty).$$

To this end, let $\mathcal{O}$ be an $\mathcal{F}$ -operad, and consider the cocartesian fibration $E_{\mathcal{R}}(p_{\mathcal{O}}) \rightarrow \text{Span}_{L,R}(F)$ from Proposition 17.2.8. We must show that its straightening $\text{Span}_{L,R}(F) \rightarrow \text{Cat}_{\infty}$ satisfies the Segal condition. The description of cocartesian morphisms in Proposition 17.2.8, applied to the factorization of a composite of spans, gives the following explicit description of this straightening:

- On objects, it sends $I \in F$ to the pullback ∞ -category $(F_R)_{/I} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes}$.
- Given a backwards morphism $g: J \leftarrow I$, the functor $(F_R)_{/J} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes} \rightarrow (F_R)_{/I} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes}$ sends a pair $(X, r: J' \rightarrow J)$ with $r \in F_R$ and $X \in \mathcal{O}_{J'}^{\otimes}$ to the pair $(Y, r': I' \rightarrow I)$, where r' is defined via the pullback square

$$\begin{array}{ccc} I' & \xrightarrow{f} & J' \\ r' \downarrow & \lrcorner & \downarrow r \\ I & \xrightarrow{g} & J, \end{array}$$

and where Y is the target of a $p_{\mathcal{O}}$ -cocartesian lift $\hat{f}_X: X \rightarrow Y$ of f .

- Given a forward morphism $g: I \rightarrow J$, the transport functor sends $(X, r: K \rightarrow I)$ to $(X, gr: K \rightarrow J)$.

Indeed, in the backwards case the displayed pullback is precisely the factorization of the composite span used in Proposition 17.2.8. In the forward case, the composite gr already belongs to F_R , so its factorization has left factor id_K and the cocartesian lift leaves X unchanged. Consider now a collection of objects $J_1, \dots, J_n$ and let $J := \bigsqcup_{i=1}^n J_i$. We must show that the map

$$(F_R)_{/J} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes} \rightarrow \prod_{i=1}^n (F_R)_{/J_i} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes}$$

is an equivalence.

Essential surjectivity: Consider an object $(X_i, r_i: I_i \rightarrow J_i)_{i=1}^n$ of the target. We define $I := \bigsqcup_{i=1}^n I_i$, let $r: I \rightarrow J$ be the map whose i -th component is r_i , and we let $X \in \mathcal{O}_I^{\otimes}$ correspond to the tuple $(X_i)_{i=1}^n$ under the equivalence $\mathcal{O}_I^{\otimes} \xrightarrow{\sim} \prod_{i=1}^n \mathcal{O}_{I_i}^{\otimes}$ from condition (2) of Definition 17.3.10. Then $(X, r: I \rightarrow J)$ is the desired preimage.

Full faithfulness: Consider two objects $(X, r: I \rightarrow J)$ and $(X', r': I' \rightarrow J)$ of $(F_R)_{/J} \times_{\text{Span}_{L,R}(F)} \mathcal{O}^{\otimes}$. We must show that the map

$$\text{Hom}_{(F_R)_{/J}}(I, I') \times_{\text{Hom}_{\text{Span}_{L,R}(F)}(I, I')} \text{Hom}_{\mathcal{O}^{\otimes}}(X, X') \rightarrow \prod_{i=1}^n \text{Hom}_{(F_R)_{/J_i}}(I_i, I'_i) \times_{\text{Hom}_{\text{Span}_{L,R}(F)}(I_i, I'_i)} \text{Hom}_{\mathcal{O}^{\otimes}}(X_i, X'_i)$$

is an equivalence. Extensivity of F gives an equivalence $F_{/J} \simeq \prod_i F_{/J_i}$. It restricts to the displayed slices of F_R : membership in F_R is preserved by base change along the inclusions $J_i \hookrightarrow J$ by adequacy, and is detected on the resulting components because F_R is closed under finite coproducts by weak coextensivity. Consequently $(F_R)_{/J} \rightarrow \prod_{i=1}^n (F_R)_{/J_i}$ is an equivalence, and hence there is a natural equivalence

$$\text{Hom}_{(F_R)_{/J}}(I, I') \simeq \prod_{i=1}^n \text{Hom}_{(F_R)_{/J_i}}(I_i, I'_i).$$

It thus remains to show that this map induces equivalences on fibers for every map $f \in \text{Hom}_{(F_R)/J}(I, I')$. Condition (3) of Definition 17.3.10 identifies the relevant fiber with the product of the fibers of $\text{Hom}_{\mathcal{O}^\otimes}(X, X'_i)$ over the spans $I \leftarrow I_i \rightarrow I'_i$. Cocartesian transport along the backwards spans $I \leftarrow I_i \xrightarrow{=} I_i$ then identifies these fibers with those of $\text{Hom}_{\mathcal{O}^\otimes}(X_i, X'_i)$ over the forward spans $I_i \rightarrow I'_i$. This is precisely the map on fibers displayed above, so it is an equivalence. $\square$

Let us explicitly record the main example of the $\mathcal{F}$ -operadic envelope construction:

Proposition 17.3.13. *The multimorphism operad functor $\mathcal{M}: \text{Cat}_\infty^\otimes \rightarrow \text{Op}_\infty$ admits a left adjoint*

$$\text{Env}: \text{Op}_\infty \rightarrow \text{Cat}_\infty^\otimes.$$

Proof. This is a special case of the previous proposition. For $\mathcal{F} = (\text{Fin}, \text{Fin}, \text{Fin})$, Proposition 14.1.7 identifies the objects of $\text{Op}_\mathcal{F}$ with the ∞ -operads from Definition 14.1.1, while Corollary 14.1.10 identifies the two notions of morphism. $\square$

Corollary 17.3.14. *Let $\mathcal{O}$ be an ∞ -operad and let C be a symmetric monoidal ∞ -category. There is a natural equivalence*

$$\text{Alg}_\mathcal{O}(C) \simeq \text{Fun}^\otimes(\text{Env}(\mathcal{O}), C).$$

In other words, an $\mathcal{O}$ -algebra in C is equivalently a symmetric monoidal functor from the envelope of $\mathcal{O}$ to C .

Proof. Let E be an ∞ -category. Using Lemma 14.2.12, the adjunction of Proposition 17.3.13, and currying, we obtain natural equivalences

$$\begin{aligned} \text{Hom}_{\text{Cat}_\infty}(E, \text{Alg}_\mathcal{O}(C)) &\simeq \text{Hom}_{\text{Op}_\infty}(\mathcal{O}, \mathcal{M}_{\text{Fun}(E, C)}) \\ &\simeq \text{Map}^\otimes(\text{Env}(\mathcal{O}), \text{Fun}(E, C)) \\ &\simeq \text{Hom}_{\text{Cat}_\infty}(E, \text{Fun}^\otimes(\text{Env}(\mathcal{O}), C)). \end{aligned}$$

Here $\text{Fun}(E, C)$ carries the pointwise symmetric monoidal structure. Since these equivalences are natural in E , the Yoneda lemma gives the claimed equivalence of ∞ -categories. $\square$

We refer to the symmetric monoidal ∞ -category $\text{Env}(\mathcal{O})$ as the *envelope* of the ∞ -operad $\mathcal{O}$. Note that its underlying ∞ -category sits in the following pullback square:

$$\begin{array}{ccc} \text{Env}(\mathcal{O}) & \xrightarrow{\quad} & \mathcal{O}^\otimes \\ \downarrow & \lrcorner & \downarrow p_\mathcal{O} \\ \text{Fin} & \hookrightarrow & \text{Span}(\text{Fin}). \end{array}$$

In particular, the objects of $\text{Env}(\mathcal{O})$ are unordered tuples $\{x_i\}_{i \in I}$ of colors of $\mathcal{O}$.

Lemma 17.3.15. *The monoidal unit of $\text{Env}(\mathcal{O})$ is the empty tuple, and its tensor product is given by concatenation of unordered tuples:*

$$\{x_i\}_{i \in I} \otimes \{y_j\}_{j \in J} \simeq \{x_i\}_{i \in I} \sqcup \{y_j\}_{j \in J}.$$

Proof. Under the Segal equivalence, the two tuples on the left determine the object of the fiber over $\langle 2 \rangle$ indexed by their disjoint union $I \sqcup J$. The tensor product is cocartesian transport of this object along the forward fold $\text{span } \langle 2 \rangle \xleftarrow{=} \langle 2 \rangle \rightarrow \langle 1 \rangle$. By the forward-transport calculation in the proof of Proposition 17.3.12, this transport leaves the object of $\mathcal{O}^\otimes$ unchanged and postcomposes its indexing map with the fold map. The result is the concatenated tuple. The same calculation for the unique forward map $\emptyset \rightarrow \langle 1 \rangle$ gives the empty tuple as the unit. $\square$

Lemma 17.3.16. *The envelope $\text{Env}(\text{Comm})$ of the commutative operad is the category of finite sets, with monoidal structure given by disjoint union.*

Proof. The total category of the terminal operad Comm is $\text{Span}(\text{Fin})$ itself. The pullback description of the underlying category of the envelope therefore gives $\text{Env}(\text{Comm}) \simeq \text{Fin}$, and Lemma 17.3.15 identifies the tensor product with disjoint union. $\square$

Lemma 17.3.17. *Let $\mathcal{O}$ be an ∞ -operad, and let $\{x_i\}_{i \in I}$ and $\{y_j\}_{j \in J}$ be objects of $\text{Env}(\mathcal{O})$. Then there is a natural equivalence*

$$\text{Hom}_{\text{Env}(\mathcal{O})}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J}) \simeq \coprod_{f: I \rightarrow J} \prod_{j \in J} \mathcal{O}(\{x_i\}_{i \in f^{-1}(j)}; y_j),$$

where the coproduct ranges over all maps of finite sets $f: I \rightarrow J$. In particular, for a single output color $y \in \mathcal{O}^\simeq$ there is a natural equivalence

$$\text{Hom}_{\text{Env}(\mathcal{O})}(\{x_i\}_{i \in I}, y) \simeq \mathcal{O}(\{x_i\}_{i \in I}; y).$$

Proof. As noted above, the underlying ∞ -category of $\text{Env}(\mathcal{O})$ is the pullback $\text{Fin} \times_{\text{Span}(\text{Fin})} \mathcal{O}^\otimes$ along the inclusion $\text{Fin} \hookrightarrow \text{Span}(\text{Fin})$ of the forward spans. Consequently, $\text{Hom}_{\text{Env}(\mathcal{O})}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J})$ is the pullback of

$$\text{Hom}_{\mathcal{O}^\otimes}(\{x_i\}_{i \in I}, \{y_j\}_{j \in J}) \longrightarrow \text{Hom}_{\text{Span}(\text{Fin})}(I, J)$$

and the subanima $\text{Hom}_{\text{Fin}}(I, J)$ of forward spans, which produces the coproduct over the maps $f: I \rightarrow J$. Fixing such an f , condition (3) of Definition 17.3.10 identifies the fiber over f with the product over $j \in J$ of the fibers of $\text{Hom}_{\mathcal{O}^\otimes}(\{x_i\}_{i \in I}, y_j)$ over the span $I \hookleftarrow f^{-1}(j) \rightarrow \langle 1 \rangle$. Cocartesian transport along the inert morphism $\{x_i\}_{i \in I} \rightarrow \{x_i\}_{i \in f^{-1}(j)}$ identifies the latter fiber with the multimorphism anima $\mathcal{O}(\{x_i\}_{i \in f^{-1}(j)}; y_j)$. The final statement is the case $J = \langle 1 \rangle$. $\square$

By functoriality, any morphism of ∞ -operads induces a symmetric monoidal functor on envelopes. In particular, the terminal morphism $\mathcal{O} \rightarrow \text{Comm}$ from any ∞ -operad $\mathcal{O}$ induces a symmetric monoidal functor $\text{Env}(\mathcal{O}) \rightarrow \text{Env}(\text{Comm}) \simeq (\text{Fin}, \amalg)$.

Proposition 17.3.18. *The induced functor*

$$\text{Env}: \text{Op}_\infty \rightarrow (\text{Cat}_\infty^\otimes)_{/(\text{Fin}, \amalg)}$$

is fully faithful. A symmetric monoidal functor $p: \mathcal{C} \rightarrow (\text{Fin}, \amalg)$ lies in the essential image if and only if it satisfies the following two conditions:

- (1) *For every finite set I , the tensor product in $\mathcal{C}$ defines an equivalence*

$$\prod_{i \in I} \mathcal{C}_{\{i\}} \xrightarrow{\sim} \mathcal{C}_I,$$

where $\mathcal{C}_I$ is the fiber of p over $I \in \text{Fin}$;

- (2) For every finite collection $(I_j)_{j \in J}$ of finite sets and every collection of objects $X_j \in p^{-1}(I_j)$ and $y_j \in p^{-1}(\{j\})$, the commutative square

$$\begin{array}{ccc} \prod_{j \in J} \mathrm{Hom}_C(X_j, y_j) & \xrightarrow{\otimes_{j \in J}} & \mathrm{Hom}_C(\bigotimes_{j \in J} X_j, \bigotimes_{j \in J} y_j) \\ \downarrow & & \downarrow p \\ * & \xrightarrow{\quad} & \mathrm{Hom}_{\mathrm{Fin}}(\bigsqcup_{j \in J} I_j, J) \end{array}$$

is a pullback square, where the bottom map hits the map $\bigsqcup_{j \in J} I_j \rightarrow \bigsqcup_{j \in J} * = J$ induced by the maps $I_j \rightarrow *$.

Proof. The full faithfulness is an instance of Lemma 17.2.11. To identify the essential image, we first show that the two conditions are always satisfied for $C = \mathrm{Env}(\mathcal{O})$. Condition (1) follows from the Segal condition on $\mathcal{O}$ and Lemma 17.3.15: the underlying map $\mathrm{Env}(\mathcal{O}) \rightarrow \mathrm{Fin}$ is simply the pullback of the map $\mathcal{O}^\otimes \rightarrow \mathrm{Span}(\mathrm{Fin})$. For condition (2), let us write $X := \bigotimes_{j \in J} X_j$ and $Y := \bigotimes_{j \in J} y_j$, where these two tensor products in $C = \mathrm{Env}(\mathcal{O})$ are computed as unordered concatenations in $\mathcal{O}^\otimes$. Then we have a pullback square

$$\begin{array}{ccc} \mathrm{Hom}_C(\bigotimes_{j \in J} X_j, \bigotimes_{j \in J} y_j) & \xrightarrow{\quad} & \mathrm{Hom}_{\mathcal{O}^\otimes}(X, Y) \\ p \downarrow & \lrcorner & \downarrow p_\otimes \\ \mathrm{Hom}_{\mathrm{Fin}}(\bigsqcup_{j \in J} I_j, J) & \xrightarrow{\quad} & \mathrm{Hom}_{\mathrm{Span}(\mathrm{Fin})}(\bigsqcup_{j \in J} I_j, J). \end{array}$$

The claim now follows from condition (3) of Definition 17.3.10.

It remains to show that any C satisfying (1) and (2) is of the form $\mathrm{Env}(\mathcal{O})$ for some ∞ -operad $\mathcal{O}$. We define $\mathcal{O}$ as the following pullback:

$$\begin{array}{ccc} \mathcal{O} & \xrightarrow{\quad} & \mathcal{M}_C \\ \downarrow & & \downarrow \mathcal{M}_p \\ \mathrm{Comm} & \xrightarrow{\quad} & \mathcal{M}_{(\mathrm{Fin}, \mathrm{II})}. \end{array}$$

The pullback is formed in $(\mathrm{Cat}_\infty)_{/\mathrm{Span}(\mathrm{Fin})}$. It is again an ∞ -operad: cocartesian lifts of backwards spans are formed componentwise, the Segal equivalences on fibers are preserved by pullback, and condition (3) is inherited by taking pullbacks of the corresponding squares. Here the bottom map is the unit of the adjunction $\mathrm{Env} \dashv \mathcal{M}$. The projection $\mathcal{O} \rightarrow \mathcal{M}_C$ induces a composite of symmetric monoidal functors

$$\mathrm{Env}(\mathcal{O}) \longrightarrow \mathrm{Env}(\mathcal{M}_C) \longrightarrow C,$$

where the second map is the counit of the adjunction. Equivalently, this is the symmetric monoidal functor corresponding under Corollary 17.3.14 to the projection $\mathcal{O} \rightarrow \mathcal{M}_C$. Naturality of the counit with respect to p shows that this composite is a functor over $(\mathrm{Fin}, \mathrm{II})$. We show that it is an equivalence.

The colors of $\mathcal{O}$ are precisely the objects of $C_{\langle 1 \rangle}$. Since every object of $\mathrm{Env}(\mathcal{O})$ is a tensor product of colors, essential surjectivity follows from condition (1) on C . For full faithfulness, first consider a target color $y \in C_{\langle 1 \rangle}$. We have equivalences:

$$\mathrm{Hom}_{\mathrm{Env}(\mathcal{O})}(\{x_i\}, y) \simeq \mathcal{O}(\{x_i\}; y) \simeq \mathrm{Hom}_C(\bigotimes_{i \in I} x_i, y)$$

The first is Lemma 17.3.17. For the second, the pullback defining $\mathcal{O}$ identifies its multimorphism anima with the fiber of

$$\mathrm{Hom}_C\left(\bigotimes_{i \in I} x_i, y\right) \longrightarrow \mathrm{Hom}_{\mathrm{Fin}}(I, \langle 1 \rangle)$$

over the unique map $I \rightarrow \langle 1 \rangle$; this fiber is the entire source because the target is contractible.

For a general target $\{y_j\}_{j \in J}$, Lemma 17.3.17 decomposes the hom anima in $\mathrm{Env}(\mathcal{O})$ as a coproduct indexed by maps $f: I \rightarrow J$, with the fiber over f equal to

$$\prod_{j \in J} \mathrm{Hom}_C\left(\bigotimes_{i \in f^{-1}(j)} x_i, y_j\right).$$

Condition (2) identifies this product with the fiber over f of

$$\mathrm{Hom}_C\left(\bigotimes_{i \in I} x_i, \bigotimes_{j \in J} y_j\right) \longrightarrow \mathrm{Hom}_{\mathrm{Fin}}(I, J).$$

Taking the coproduct over all f proves full faithfulness. $\square$

17.4 Comparison to Lurie's operads

As mentioned in Remark 12.4.6, our definition of ∞ -operads does not agree with Lurie's definition, which instead works with ∞ -categories over the category Fin_* of finite pointed sets. In this section, we recall Lurie's definition, and show that the ∞ -category $\mathrm{Op}_{\infty}^{\mathrm{Lurie}}$ of Lurie's ∞ -operads is equivalent to Op_{∞} .

Definition 17.4.1. Given a finite set I , we write $I_+ := I \sqcup *$ for the resulting finite pointed set. A morphism $f: I_+ \rightarrow J_+$ of finite pointed sets is called *inert* if on the preimage of J it restricts to a bijection $f^{-1}(J) \xrightarrow{\sim} J$. We say that f is *active* if $f^{-1}(*) = \{*\}$.

Given a finite set I , the *Segal map* $\rho_i: I_+ \rightarrow \{i\}_+$ for $i \in I$ is defined as the map that is the identity on $\{i\}$ and sends all other $j \in I \setminus \{i\}$ to the basepoint.

Remark 17.4.2. Recall from Lemma 5.3.8 that there is an equivalence

$$\mathrm{Span}_{\mathrm{inj}, \mathrm{all}}(\mathrm{Fin}) \simeq \mathrm{Fin}_*,$$

given on objects by sending I to I_+ and on morphisms by sending a span $I \xleftarrow{f} K \xrightarrow{g} J$ to the pointed map $h: I_+ \rightarrow J_+$ given by $h(i) = g(k)$ whenever $i = f(k)$ is in the image of f and $h(i) = *$ otherwise. Under this equivalence, the inert maps in Fin_* correspond to the backwards spans $I \xleftarrow{f} K \xrightarrow{\sim} K$, and the active maps in Fin_* correspond to the forward spans $I \xleftarrow{\sim} I \xrightarrow{g} J$. The Segal maps ρ_i correspond to the backwards spans

$$I \xleftarrow{\sim} \{i\} \xrightarrow{\sim} \{i\}.$$

We will write $\mathrm{Fin}_* \hookrightarrow \mathrm{Span}(\mathrm{Fin})$ for the inclusion of the subcategory $\mathrm{Span}_{\mathrm{inj}, \mathrm{all}}(\mathrm{Fin})$.

Definition 17.4.3. Let Fin_* be the category of finite pointed sets. An ∞ -operad in the sense of Lurie is a pair $(\mathcal{O}^{\otimes}, p_{\mathcal{O}})$ consisting of an ∞ -category $\mathcal{O}^{\otimes}$ equipped with a functor $p_{\mathcal{O}}: \mathcal{O}^{\otimes} \rightarrow \mathrm{Fin}_*$ satisfying the following conditions:

- (1) For any inert morphism $f: I_+ \rightarrow J_+$ and any object $X \in \mathcal{O}_I^\otimes$ there is a p_O -cocartesian lift $X \rightarrow Y$ in $\mathcal{O}^\otimes$.
- (2) Let X and Y be objects of $\mathcal{O}^\otimes$ and let $I_+ := p(X)$ and $J_+ := p(Y)$. For every $j \in J$, let $Y \rightarrow Y_j$ denote a p_O -cocartesian lift of the Segal map $\rho_j: J_+ \rightarrow \{j\}_+$. Then the induced commutative square

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{O}^\otimes}(X, Y) & \longrightarrow & \prod_{j \in J} \mathrm{Hom}_{\mathcal{O}^\otimes}(X, Y_j) \\ p_O \downarrow & & \downarrow p_O \\ \mathrm{Hom}_{\mathrm{Fin}_*}(I_+, J_+) & \longrightarrow & \prod_{j \in J} \mathrm{Hom}_{\mathrm{Fin}_*}(I_+, \{j\}_+) \end{array}$$

is a pullback square.

- (3) For every finite collection $\{X_j\}_{j \in J}$ of objects $X_j \in \mathcal{O}_{\{j\}}^\otimes$, there exists an object $X \in \mathcal{O}_J^\otimes$ and a collection of p_O -cocartesian morphisms $X \rightarrow X_j$ covering $\rho_j: J_+ \rightarrow \{j\}_+$.

We say a functor $f: \mathcal{O}^\otimes \rightarrow \mathcal{P}^\otimes$ over Fin_* is a *morphism of Lurie- ∞ -operads* if it preserves cocartesian morphisms over inerts. We let $\mathrm{Op}_\infty^{\mathrm{Lurie}} \subseteq (\mathrm{Cat}_\infty)_{/\mathrm{Fin}_*}$ denote the ∞ -category of the Lurie ∞ -operads and their morphisms.

Let $\mathcal{F}_{\mathrm{Lurie}} := (\mathrm{Fin}, \mathrm{Fin}_{\mathrm{inj}}, \mathrm{Fin})$, so that $\mathrm{Span}_{\mathrm{inj}, \mathrm{all}}(\mathrm{Fin}) \simeq \mathrm{Fin}_*$.

Lemma 17.4.4. *Under the equivalence $\mathrm{Span}_{\mathrm{inj}, \mathrm{all}}(\mathrm{Fin}) \simeq \mathrm{Fin}_*$, the $\mathcal{F}_{\mathrm{Lurie}}$ -operads of Definition 17.3.10 are precisely the ∞ -operads in the sense of Lurie, and the two notions of morphism agree. In particular, there is a canonical equivalence*

$$\mathrm{Op}_{\mathcal{F}_{\mathrm{Lurie}}} \simeq \mathrm{Op}_\infty^{\mathrm{Lurie}}.$$

Proof. Condition (1) in the two definitions is the same. Suppose first that $p_O: \mathcal{O}^\otimes \rightarrow \mathrm{Fin}_*$ is an ∞ -operad in the sense of Lurie. The functor

$$\mathcal{O}_J^\otimes \longrightarrow \prod_{j \in J} \mathcal{O}_{\{j\}}^\otimes$$

induced by cocartesian transport along the Segal maps is essentially surjective by condition (3) of Definition 17.4.3. It is fully faithful as well: condition (2), restricted to the fiber over id_{J_+} , identifies the hom anima in $\mathcal{O}_J^\otimes$ with the product of the hom animae in the fibers over the singletons, using cocartesian transport along the Segal maps on the sources. Thus condition (2) holds for the decomposition of J into singletons, and hence for every finite decomposition by grouping the singleton factors. Condition (3) follows similarly from condition (2): apply it to the singleton decomposition of J and to those of the summands J_i , and group the resulting factors according to $J = \bigsqcup_i J_i$.

Conversely, if $\mathcal{O}$ is an $\mathcal{F}_{\mathrm{Lurie}}$ -operad, condition (1) gives the required cocartesian lifts of inert maps. Specializing condition (3) to the decomposition of J into singletons gives condition (2) of Definition 17.4.3, while condition (2) gives condition (3). Finally, both notions of morphism require precisely the preservation of cocartesian morphisms over inert maps. $\square$

Construction 17.4.5. Using Lemma 17.4.4 and Proposition 5.3.9, we define the *Lurie operadic envelope*

$$\mathrm{Env}^{\mathrm{Lurie}} : \mathrm{Op}_{\infty}^{\mathrm{Lurie}} \longrightarrow \mathrm{Cat}_{\infty}^{\otimes}$$

as the functor $\mathrm{Env}_{\mathcal{F}_{\mathrm{Lurie}}}$ of Proposition 17.3.12, followed by the equivalence from $\mathcal{F}_{\mathrm{Lurie}}$ -monoidal ∞ -categories to symmetric monoidal ∞ -categories.

Lemma 17.4.6. *If $\mathcal{O} = (\mathcal{O}^{\otimes}, p_{\mathcal{O}})$ is an ∞ -operad in the sense of Definition 14.1.1, then its pullback $p_{\mathcal{P}} : \mathcal{P}^{\otimes} := \mathcal{O}^{\otimes} \times_{\mathrm{Span}(\mathrm{Fin})} \mathrm{Fin}_{*} \rightarrow \mathrm{Fin}_{*}$ is an ∞ -operad in the sense of Lurie. Moreover, forming pullbacks along the inclusion $\mathrm{Fin}_{*} \hookrightarrow \mathrm{Span}(\mathrm{Fin})$ induces a ‘forgetful functor’*

$$\mathrm{Op}_{\infty} \rightarrow \mathrm{Op}_{\infty}^{\mathrm{Lurie}}.$$

Proof. Any $p_{\mathcal{O}}$ -cocartesian lift of a backwards span $I \hookleftarrow J \xrightarrow{=} J$ will still be $p_{\mathcal{P}}$ -cocartesian, verifying condition (1). Condition (2) is immediate from Lemma 14.1.9. Condition (3) is immediate from condition (ii) in Proposition 14.1.7. The claim on morphisms is checked similarly. $\square$

Lemma 17.4.7. *Let*

$$u : \mathrm{Fin}_{*} \simeq \mathrm{Span}_{\mathrm{inj}, \mathrm{all}}(\mathrm{Fin}) \hookrightarrow \mathrm{Span}(\mathrm{Fin})$$

be the inclusion, and let $\mathcal{R}_{}$ and $\mathcal{R}$ denote the forward morphisms in Fin_{*} and $\mathrm{Span}(\mathrm{Fin})$, respectively. For every functor $p : E \rightarrow \mathrm{Span}(\mathrm{Fin})$, there is a natural equivalence of functors over Fin_{*}*

$$u^{*} E_{\mathcal{R}}(p) \simeq E_{\mathcal{R}_{*}}(u^{*} p).$$

Proof. By the pullback formula in Construction 17.2.2, it suffices to construct a natural equivalence

$$\mathrm{Fin}_{*} \times_{\mathrm{Span}(\mathrm{Fin}), t} \mathrm{Ar}_{\mathcal{R}}(\mathrm{Span}(\mathrm{Fin})) \simeq \mathrm{Ar}_{\mathcal{R}_{*}}(\mathrm{Fin}_{*})$$

over Fin_{*} . The objects on both sides are forward spans, equivalently maps of finite sets $r : I \rightarrow J$. A morphism on the left is a commutative square in $\mathrm{Span}(\mathrm{Fin})$

$$\begin{array}{ccc} I & \xrightarrow{f} & I' \\ r \downarrow & & \downarrow r' \\ J & \xrightarrow{g} & J' \end{array}$$

in which g belongs to Fin_{*} . The composite gr has injective backwards leg because this leg is obtained by pulling back the injective backwards leg of g along r . Since r' is forward, the backwards leg of $r'f$ is the backwards leg of f . The equivalence $r'f \simeq gr$ therefore shows that f also belongs to Fin_{*} . As $\mathrm{Fin}_{*} \hookrightarrow \mathrm{Span}(\mathrm{Fin})$ is a wide subcategory inclusion, its hom animae are unions of components, so the homotopy making the square commute also lies in Fin_{*} . Thus the displayed square is precisely a morphism in $\mathrm{Ar}_{\mathcal{R}_{*}}(\mathrm{Fin}_{*})$, which proves the claim. $\square$

Proposition 17.4.8 (cf. [BHS22, Theorem 5.1.1, Corollary 5.1.15]). *The forgetful functor $\mathrm{Op}_{\infty} \rightarrow \mathrm{Op}_{\infty}^{\mathrm{Lurie}}$ is an equivalence.*

Proof. By Lemma 17.4.4, we may work with $\mathcal{F}_{\mathrm{Lurie}}$ -operads. The proof of Proposition 17.3.18 applies to this span pattern as well. Indeed, full faithfulness is the general statement Lemma 17.2.11; the Segal and mapping-anima arguments use only conditions (2) and (3); and the forward-transport

calculation of Lemma 17.3.15 applies unchanged. After the equivalence of Proposition 5.3.9, we therefore obtain a fully faithful functor

$$\mathrm{Env}^{\mathrm{Lurie}}: \mathrm{Op}_{\infty}^{\mathrm{Lurie}} \longrightarrow (\mathrm{Cat}_{\infty}^{\otimes})/(\mathrm{Fin}, \mathrm{II})$$

with essential image characterized by the two conditions in Proposition 17.3.18.

Now let $u: \mathrm{Fin}_{*} \hookrightarrow \mathrm{Span}(\mathrm{Fin})$ be the inclusion and let $\mathcal{O}$ be an ∞ -operad over $\mathrm{Span}(\mathrm{Fin})$. Applying Lemma 17.4.7 to $p_{\mathcal{O}}$ gives a natural equivalence of $\mathcal{F}_{\mathrm{Lurie}}$ -monoidal ∞ -categories

$$\mathrm{Env}^{\mathrm{Lurie}}(u^{*}\mathcal{O}) \simeq u^{*}\mathrm{Env}(\mathcal{O}).$$

Under Proposition 5.3.9, the right-hand side corresponds to the original symmetric monoidal ∞ -category $\mathrm{Env}(\mathcal{O})$. The equivalence is compatible with the maps to $(\mathrm{Fin}, \mathrm{II})$ by naturality of the construction. We therefore obtain a naturally commutative diagram

$$\begin{array}{ccc} \mathrm{Op}_{\infty} & \xrightarrow{\mathrm{Env}} & (\mathrm{Cat}_{\infty}^{\otimes})/(\mathrm{Fin}, \mathrm{II}) \\ \downarrow -\times_{\mathrm{Span}(\mathrm{Fin})} \mathrm{Fin}_{*} & & \uparrow \mathrm{Env}^{\mathrm{Lurie}} \\ \mathrm{Op}_{\infty}^{\mathrm{Lurie}} & & \end{array}$$

Both diagonal functors are equivalences onto the same essential image. It follows by 2-out-of-3 that the left vertical functor is an equivalence, as desired. $\square$

As a first payoff of the comparison, we can now make precise the passage from classical colored operads to ∞ -operads. In Construction 12.4.1, we discussed informally how to associate an ∞ -operad to every colored operad. We can make this formal without choosing a nerve for the $(2,1)$ -category appearing there by first using Lurie's model and then applying the comparison from Proposition 17.4.8.

Proposition 17.4.9. *Every colored operad $\mathcal{O}$ canonically and functorially determines an ∞ -operad, also denoted $\mathcal{O}$.*

Proof. Define a classical 1-category $\mathcal{O}^{\otimes, \mathrm{Lurie}}$ as in Construction 12.4.1, with the additional requirement that in the underlying span $I \xleftarrow{f} K \xrightarrow{g} J$ the finite set K is an actual *subset* of I and f is its inclusion. Composition is defined as in Construction 12.4.1, taking as the pullback of spans the literal preimage: the composite of $(K \subseteq I, g: K \rightarrow J)$ with $(M \subseteq J, k: M \rightarrow L)$ has underlying subset $g^{-1}(M) \subseteq K \subseteq I$ and underlying map the restriction of $k \circ g$ to it. This is again a subset inclusion, and the resulting composition is strictly associative: composing with a third morphism $(N \subseteq L, l: N \rightarrow P)$ in either of the two possible orders yields the subset $g^{-1}(k^{-1}(N)) \subseteq I$, together with the restriction of $l \circ k \circ g$. The identity of $\{x_i\}_{i \in I}$ is given by $K = I$ and $g = \mathrm{id}_I$, labeled by identity multimorphisms. On the level of the multimorphism labels, associativity and unitality of this composition are precisely the associativity and unitality axioms of the colored operad $\mathcal{O}$. Thus $\mathcal{O}^{\otimes, \mathrm{Lurie}}$ is a classical 1-category. Sending $\{x_i\}_{i \in I}$ to I_{+} and a morphism $(K \subseteq I, g, \{\varphi_j\}_{j \in J})$ to the pointed map $I_{+} \rightarrow J_{+}$ that restricts to g on K and sends $I \setminus K$ to the basepoint defines a functor of 1-categories

$$p_{\mathcal{O}}: \mathcal{O}^{\otimes, \mathrm{Lurie}} \longrightarrow \mathrm{Fin}_{*}.$$

Its fiber over I_+ consists of the I -tuples of colors of O and the I -tuples of unary multimorphisms between them, i.e. it is the I -fold power of the category of colors and unary multimorphisms of O . For an inert morphism, the morphism obtained by restricting an I -tuple of colors and labeling the resulting unary operations by identities is p_O -cocartesian. These cocartesian lifts give the required equivalences from the fiber over I_+ to the product of the fibers over its elements.

It remains to verify the condition on morphism sets. Let $X = \{x_i\}_{i \in I}$ and $Y = \{y_j\}_{j \in J}$. For a pointed map $\alpha: I_+ \rightarrow J_+$, the set of morphisms $X \rightarrow Y$ over α is, by construction,

$$\prod_{j \in J} \mathcal{O}((x_i)_{i \in \alpha^{-1}(j)}; y_j).$$

The j -th factor is precisely the set of morphisms from X to y_j over the composite of α with the Segal map ρ_j . Consequently the square in condition (2) of Lurie's definition is a pullback. Thus p_O is an ∞ -operad in Lurie's sense. A morphism of colored operads acts on tuples and multimorphism labels and thereby induces a functor between the resulting categories over Fin_* . Applying an inverse to the equivalence in Proposition 17.4.8 therefore produces the claimed functor from colored operads to ∞ -operads over $\text{Span}(\text{Fin})$. $\square$

Corollary 17.4.10. *The colored operads $\mathcal{A}ssoc$, $\mathcal{L}Mod$, and Mod of Section 12.2, as well as the bimodule operad $\mathcal{B}Mod$ of Definition 19.2.1, canonically define ∞ -operads. Their multimorphism animae are the discrete animae associated with their classical multimorphism sets.*

Proof. Apply Proposition 17.4.9 to each of the indicated colored operads. The comparison with the span model is induced by restriction along $\text{Fin}_* \hookrightarrow \text{Span}(\text{Fin})$ and therefore preserves the fibers of hom animae over active maps. The displayed formula in the proof of Proposition 17.4.9 consequently identifies the resulting multimorphism animae with the discrete animae associated with the classical multimorphism sets. $\square$

Corollary 17.4.11. *For a colored operad O , the ∞ -categorical envelope of the associated ∞ -operad is equivalent to the category underlying its classical envelope from Definition 12.3.3, compatibly with the symmetric monoidal structures and the maps to $(\text{Fin}, \amalg)$.*

Proof. Both envelopes have the unordered tuples of colors as objects. The description of the classical morphism sets in Definition 12.3.3 agrees with the mapping-anima formula of Lemma 17.3.17, since the latter is discrete for a classical colored operad. These identifications respect composition, and both tensor products are given by concatenation of tuples. $\square$

The envelopes of the operads governing associative algebras and modules are also 1-categories. We record them for later use.

Example 17.4.12 (Envelopes for algebras and modules). All the following categories are symmetric monoidal under disjoint union.

- (1) The envelope $\text{Env}(\mathcal{A}ssoc)$ is the category Fin^{Alg} whose objects are finite sets and whose morphisms $S \rightarrow T$ are maps of finite sets together with a linear order on every fiber over $t \in T$.

- (2) The envelope $\text{Env}(\mathcal{L}\text{Mod})$ is the category Fin^{LMod} whose objects are pairs (S_A, S_M) of finite sets. A morphism $(S_A, S_M) \rightarrow (T_A, T_M)$ is a map

$$f: S_A \sqcup S_M \longrightarrow T_A \sqcup T_M$$

which restricts to a bijection $S_M \rightarrow T_M$, together with a linear order on $f^{-1}(t) \cap S_A$ for every $t \in T_A \sqcup T_M$.

- (3) The envelope $\text{Env}(\text{Mod})$ is the category Fin^{Mod} with the same objects and morphisms as in (2), but without the choices of linear orders.

These descriptions follow immediately from Lemma 17.3.17 and the multimorphism sets of $\mathcal{A}\text{ssoc}$, $\mathcal{L}\text{Mod}$, and Mod .

Exercises

Exercise 17.1. Let C be an ∞ -category. Show that the pairs $(C, C^\simeq)$ and $(C^\simeq, C)$ are both factorization systems on C .

Exercise 17.2. Let $C = \text{An}$ be the ∞ -category of (small) anima, let $\mathcal{L}$ be the class of essentially surjective morphisms, and let $\mathcal{R}$ be the class of monomorphisms. Show that the pair $(\mathcal{L}, \mathcal{R})$ is a factorization system on An .

Exercise 17.3. Let $C = \text{Cat}_\infty$ be the ∞ -category of (small) ∞ -categories, let $\mathcal{L}$ be the class of essentially surjective functors and let $\mathcal{R}$ be the class of fully faithful functors. Show that $(\mathcal{L}, \mathcal{R})$ is a factorization system on Cat_∞ .

Exercise 17.4. Using Corollary 15.2.21, show that the symmetric monoidal structure on $\text{Env}(\text{Comm}) \simeq (\text{Fin}, \amalg)$ from Lemma 17.3.16 is the cocartesian monoidal structure on Fin .

18 Operadic Day convolution and stabilization

In Chapter 16, we realized Sp as the ∞ -category of reduced excisive functors from finite pointed animae to animae and used Day convolution to construct its tensor product. This construction suggests a more general question. Recall from Section 4.3 that $\mathrm{Sp} = \mathrm{Sp}(\mathrm{An})$ is an instance of the stabilization $\mathrm{Sp}(C)$ of an ∞ -category C with finite limits, characterized as the cofree stable ∞ -category generated by C . If C carries a symmetric monoidal structure, when does $\mathrm{Sp}(C)$ inherit one as well?

Stabilization is the last of four parallel constructions: pointed objects C_* , commutative monoids $\mathrm{CMon}(C)$, commutative groups $\mathrm{CGrp}(C)$, and spectrum objects $\mathrm{Sp}(C)$. Following Nikolaus [Nik16], we treat all four uniformly. Their lax symmetric monoidal universal properties are most naturally formulated in the operadic setting. Thus, for an ∞ -operad $\mathcal{O}$ with sufficient (co)limits, we will construct:

- The pointed ∞ -operad $\mathcal{O}_*$ of pointed objects in $\mathcal{O}$;
- The semiadditive ∞ -operad $\mathrm{CMon}(\mathcal{O})$ of commutative monoid objects in $\mathcal{O}$;
- The additive ∞ -operad $\mathrm{CGrp}(\mathcal{O})$ of commutative group objects in $\mathcal{O}$;
- The stable ∞ -operad $\mathrm{Sp}(\mathcal{O})$ of spectrum objects in $\mathcal{O}$.

Each of these constructions comes with a universal property, exhibiting it as the *cofree* operad with the corresponding algebraic structure built from $\mathcal{O}$. These constructions always produce ∞ -operads, but even when $\mathcal{O} = \mathcal{M}_C$ they need not be represented by symmetric monoidal ∞ -categories. In the presentable setting they are represented by the corresponding presentably symmetric monoidal ∞ -categories; for stabilization, see Corollary 18.5.5.

Applying these four constructions to the cartesian symmetric monoidal ∞ -category $(\mathrm{An}, \times)$ recovers the symmetric monoidal structures on An_* , $\mathrm{CMon}(\mathrm{An})$, $\mathrm{CGrp}(\mathrm{An})$ and $\mathrm{Sp} = \mathrm{Sp}(\mathrm{An})$ constructed in Chapter 16.

The operadic results are genuinely more general than the presentable theory: they apply to non-presentable ∞ -operads, and the stabilization result characterizes lax symmetric monoidal functors whose underlying functors preserve finite limits. Readers interested only in presentably symmetric monoidal ∞ -categories may instead pass directly to Section 18.5, where the four constructions arise from smashing localizations of Pr^{L} . In that setting the operadic stabilization is represented by a symmetric monoidal structure, as we will explain in Corollary 18.5.5.

The chapter is organized as follows. We begin in Section 18.1 with the notions of operadic limit and colimit, which let us formulate the universal properties that the four constructions above should satisfy. We then construct the multiplicative Yoneda embedding in Section 18.2 and extend Day convolution to arbitrary target operads in Section 18.3. Combining these ingredients, Section 18.4 constructs the four cofree operads. The final Section 18.5 explains the common universal property of the resulting presentably symmetric monoidal ∞ -categories.

18.1 Operadic (co)limits

For our upcoming discussion of ∞ -operads with additional algebraic structure, we will need notions of limit and colimit that respect not only the underlying ∞ -category $\mathcal{O}_{\langle 1 \rangle}$ but also the multimorphisms of $\mathcal{O}$.

Definition 18.1.1. Let $\mathcal{O}$ be an ∞ -operad and let $F: I \rightarrow \mathcal{O}_{\langle 1 \rangle}$ be a functor.

- (1) A cone $\eta: \text{const}_y \rightarrow F$ in $\mathcal{O}_{\langle 1 \rangle}$ is called an *operadic limit* if for every $n \geq 0$ and all colors $x_1, \dots, x_n \in \mathcal{O}^\approx$ the induced map

$$\mathcal{O}((x_1, \dots, x_n); y) \rightarrow \lim_{i \in I} \mathcal{O}((x_1, \dots, x_n); F(i))$$

is an equivalence of anima.

- (2) A cocone $\varepsilon: F \rightarrow \text{const}_x$ in $\mathcal{O}_{\langle 1 \rangle}$ is called an *operadic colimit* if for every $n \geq 1$ and all colors $x_2, \dots, x_n, y \in \mathcal{O}^\approx$ the induced map

$$\mathcal{O}((x, x_2, \dots, x_n); y) \rightarrow \lim_{i \in I} \mathcal{O}(F(i), x_2, \dots, x_n; y)$$

is an equivalence of anima.

Remark 18.1.2. These conditions are direct operadic analogues of the mapping-anima characterizations of ordinary limits and colimits, with unary mapping anima replaced by multimorphism anima. Taking $n = 1$ shows that an operadic limit is a limit in $\mathcal{O}_{\langle 1 \rangle}$, and an operadic colimit is a colimit in $\mathcal{O}_{\langle 1 \rangle}$. We will therefore usually assume that the relevant (co)limits already exist in $\mathcal{O}_{\langle 1 \rangle}$ and only ask whether they are operadic.

For limits, there is also a useful reformulation through the envelope. Every object of $\text{Env}(\mathcal{O})$ is a tensor $x_1 \otimes \dots \otimes x_n$ of colors, and by Lemma 17.3.17 the multimorphism anima compute the mapping anima out of such tensors, $\mathcal{O}((x_1, \dots, x_n); y) \simeq \text{Hom}_{\text{Env}(\mathcal{O})}(x_1 \otimes \dots \otimes x_n, y)$. Testing against all colors $x_1, \dots, x_n$ therefore shows that a cone in $\mathcal{O}_{\langle 1 \rangle}$ is an operadic limit if and only if it becomes a limit cone in $\text{Env}(\mathcal{O})$ under the inclusion $\mathcal{O}_{\langle 1 \rangle} \hookrightarrow \text{Env}(\mathcal{O})$.

For colimits, the corresponding statement is only relative: after tensoring with any fixed colors $x_2, \dots, x_n$, the cocone is colimiting when tested by mapping into colors of $\mathcal{O}$. It need not be a colimit cocone in the whole envelope; in particular, an operadic initial object need not be initial there. Compare Example 18.1.7.

Example 18.1.3. If $\mathcal{O} = \mathcal{M}_C$ for a symmetric monoidal ∞ -category C , then any limit in C is automatically operadic. This is because the tensor product provides a natural equivalence

$$\mathcal{M}_C((x_1, \dots, x_n); -) \simeq \text{Hom}_C(x_1 \otimes \dots \otimes x_n, -),$$

and the functor $\text{Hom}_C(X, -)$ preserves limits for any object X .

Example 18.1.4. The claim from the previous example does not hold for general operads. For instance, consider the trivial operad $\mathcal{Triv}$. Its underlying ∞ -category is the terminal category $*$, which has all limits, and in particular a terminal object. However, for $n \neq 1$, we have $\mathcal{Triv}((x_1, \dots, x_n); -) = \emptyset$, so this terminal object is not an operadic terminal object.

Example 18.1.5. If $\mathcal{O} = \mathcal{M}_C$ for a symmetric monoidal ∞ -category C , then a colimit in C is operadic if and only if it is preserved by the functor $x \otimes - : C \rightarrow C$ for every object $x \in C$; the argument is similar to Example 18.1.3.

This brings us to the key definitions for this chapter, which are direct analogues of the corresponding notions for ∞ -categories:

Definition 18.1.6. An ∞ -operad $\mathcal{O}$ is called:

- (1) *Pointed* if $\mathcal{O}_{\langle 1 \rangle}$ has a zero object which is both operadic initial and operadic terminal.
- (2) *Semiadditive* if $\mathcal{O}_{\langle 1 \rangle}$ is semiadditive and all finite products and coproducts are operadic.
- (3) *Additive* if $\mathcal{O}_{\langle 1 \rangle}$ is additive and all finite products and coproducts are operadic.
- (4) *Stable* if $\mathcal{O}_{\langle 1 \rangle}$ is stable and all finite limits and colimits are operadic.

This allows us to define various subcategories of the ∞ -category $\mathbf{Op}_\infty$ of ∞ -operads:

- We define $\mathbf{Op}_\infty^* \subseteq \mathbf{Op}_\infty$ as the (non-full) subcategory spanned by ∞ -operads with an operadic terminal object and maps preserving them. We denote by $\mathbf{Op}_\infty^{\text{pt}} \subseteq \mathbf{Op}_\infty^*$ the full subcategory spanned by pointed ∞ -operads.
- We define $\mathbf{Op}_\infty^{\text{prod}} \subseteq \mathbf{Op}_\infty$ as the (non-full) subcategory spanned by ∞ -operads with finite operadic products and maps preserving them. We denote by $\mathbf{Op}_\infty^{\text{sadd}}, \mathbf{Op}_\infty^{\text{add}} \subseteq \mathbf{Op}_\infty^{\text{prod}}$ the full subcategories spanned by semiadditive and additive ∞ -operads, respectively.
- We define $\mathbf{Op}_\infty^{\text{lex}} \subseteq \mathbf{Op}_\infty$ as the (non-full) subcategory spanned by ∞ -operads with finite operadic limits and maps preserving them. We denote by $\mathbf{Op}_\infty^{\text{st}} \subseteq \mathbf{Op}_\infty^{\text{lex}}$ the full subcategory spanned by stable ∞ -operads.

For example, the operad $\mathcal{M}_{\text{Sp}}$ associated with the symmetric monoidal structure constructed in Section 16.6 is stable.

Example 18.1.7 (Pointed objects). Let C satisfy the hypotheses of Lemma 16.3.4, and equip C_* with its smash product monoidal structure. Its zero object is both operadic terminal and operadic initial, so $\mathcal{M}_{C_*}$ is a pointed ∞ -operad. Indeed, operadic terminality is automatic by Example 18.1.3, while operadic initiality says precisely that tensoring with the zero object gives the zero object.

Lemma 18.1.8 (Properties inherited by full suboperads). *Let $\mathcal{O} \subseteq \mathcal{P}$ be a full suboperad.*

- (1) *If $\mathcal{P}$ is pointed and $\mathcal{O}_{\langle 1 \rangle}$ contains its zero object, then $\mathcal{O}$ is pointed.*
- (2) *If $\mathcal{P}$ is semiadditive or additive and $\mathcal{O}_{\langle 1 \rangle}$ is closed under finite biproducts in $\mathcal{P}_{\langle 1 \rangle}$, then $\mathcal{O}$ is semiadditive or additive, respectively.*

(3) If $\mathcal{P}$ is stable and $\mathcal{O}_{\langle 1 \rangle}$ is closed under finite limits and finite colimits in $\mathcal{P}_{\langle 1 \rangle}$, then $\mathcal{O}$ is stable.

Proof. In each case the relevant (co)limits in $\mathcal{O}_{\langle 1 \rangle}$ are inherited from $\mathcal{P}_{\langle 1 \rangle}$. This gives the asserted property of the underlying ∞ -category. Since $\mathcal{P}$ has the corresponding operadic (co)limits and $\mathcal{O} \subseteq \mathcal{P}$ is full, the same multimorphism anima compute the operadic (co)limit conditions in $\mathcal{O}$. $\square$

The goal of the remainder of this chapter is to show that the four fully faithful inclusion functors

$$\begin{aligned} \mathrm{Op}_{\infty}^{\mathrm{pt}} &\hookrightarrow \mathrm{Op}_{\infty}^* \\ \mathrm{Op}_{\infty}^{\mathrm{sadd}} &\hookrightarrow \mathrm{Op}_{\infty}^{\mathrm{prod}} \\ \mathrm{Op}_{\infty}^{\mathrm{add}} &\hookrightarrow \mathrm{Op}_{\infty}^{\mathrm{prod}} \\ \mathrm{Op}_{\infty}^{\mathrm{st}} &\hookrightarrow \mathrm{Op}_{\infty}^{\mathrm{lex}} \end{aligned}$$

all admit right adjoints, which will correspond to ‘cofreely adding’ the respective algebraic structure to an ∞ -operad.

18.2 The multiplicative Yoneda embedding

We will now explain how to embed an arbitrary ∞ -operad into the multimorphism operad of a presheaf category. This is the multiplicative version of the Yoneda embedding, first established by Nikolaus [Nik16, Proposition 2.7].

Lemma 18.2.1. *Let D be a symmetric monoidal ∞ -category. Then the Hom-functor*

$$\mathrm{Hom}_D(-, -): D^{\mathrm{op}} \times D \rightarrow \mathbf{An}$$

admits a canonical lax symmetric monoidal structure, where $\mathbf{An}$ carries the cartesian monoidal structure.

Proof. The twisted arrow construction preserves finite products, hence preserves commutative monoid objects of $\mathbf{Cat}_{\infty}$. Thus the symmetric monoidal structure on D induces a symmetric monoidal structure on $\mathrm{Tw}(D)$. The source and target projections

$$s: \mathrm{Tw}(D) \rightarrow D^{\mathrm{op}}, \quad t: \mathrm{Tw}(D) \rightarrow D$$

are natural in D , and therefore assemble to a symmetric monoidal functor

$$(s, t): \mathrm{Tw}(D) \rightarrow D^{\mathrm{op}} \times D.$$

We claim that the induced functor on total categories $(s, t)^{\otimes}: \mathrm{Tw}(D)^{\otimes} \rightarrow (D^{\mathrm{op}} \times D)^{\otimes}$ is still a left fibration. Under straightening over $\mathrm{Span}(\mathbf{Fin})$, this functor is induced by the natural transformation whose value on a finite set I is the I -fold product

$$(s, t)^I: \mathrm{Tw}(D)^I \rightarrow (D^{\mathrm{op}} \times D)^I$$

of the usual source-target left fibration. Since left fibrations are stable under finite products, Exercise 23.2 shows that $(s, t)^{\otimes}$ is a left fibration. The fiber over an object $\{(x_i, y_i)\}_{i \in I}$ is the anima

$\prod_{i \in I} \text{Hom}_D(x_i, y_i)$, with transport induced by the symmetric monoidal structure on D and composition in D .

Straightening this left fibration gives a functor

$$H^\otimes: (D^{\text{op}} \times D)^\otimes \rightarrow \text{An}.$$

On the fiber over $\langle 1 \rangle$ this is the usual Hom-functor $\text{Hom}_D(-, -): D^{\text{op}} \times D \rightarrow \text{An}$. Moreover, the description of its fibers shows that $H^\otimes$ preserves finite products: products in $(D^{\text{op}} \times D)^\otimes$ are given by concatenating finite tuples, and $H^\otimes$ sends such a tuple to the corresponding product of hom animae. By the universal property of the cartesian monoidal structure on An from Theorem 15.3.11, this finite-product-preserving functor is precisely a lax symmetric monoidal structure on $\text{Hom}_D(-, -)$. $\square$

Proposition 18.2.2. *Let D be a small symmetric monoidal ∞ -category. Then the Yoneda embedding*

$$Y: D \rightarrow \text{PSh}(D) = \text{Fun}(D^{\text{op}}, \text{An})$$

admits a canonical strong symmetric monoidal structure, where $\text{PSh}(D)$ is equipped with the Day convolution monoidal structure.

Proof. By the universal property of Day convolution, a lax symmetric monoidal structure on Y is equivalent to a lax symmetric monoidal structure on the associated functor

$$D^{\text{op}} \times D \rightarrow \text{An}, \quad (d', d) \mapsto \text{Hom}_D(d', d).$$

This is provided by Lemma 18.2.1.

It remains to show that this lax structure is strong. Let $\{d_i\}_{i \in I}$ be a finite collection of objects of D . We must show that the induced map

$$\bigotimes_{i \in I}^{\text{Day}} Y(d_i) \rightarrow Y\left(\bigotimes_{i \in I} d_i\right)$$

is an isomorphism in $\text{PSh}(D)$. Testing against an arbitrary presheaf $H: D^{\text{op}} \rightarrow \text{An}$, the left-hand side represents natural transformations

$$\text{Nat}\left(\prod_{i \in I} Y(d_i)(-), H\left(\bigotimes_{i \in I} -\right)\right)$$

of functors $(D^I)^{\text{op}} \rightarrow \text{An}$ by the defining property of the Day convolution tensor product. Applying the Yoneda lemma for D^I identifies this anima with $H(\bigotimes_{i \in I} d_i)$, which is also $\text{Nat}(Y(\bigotimes_{i \in I} d_i), H)$. This proves the claim. $\square$

Proposition 18.2.3 (Multiplicative Yoneda embedding). *Let $\mathcal{O}$ be a small ∞ -operad and put $D := \text{Env}(\mathcal{O})$. The composite*

$$\mathcal{O} \xrightarrow{\eta_{\mathcal{O}}} \mathcal{M}_D \xrightarrow{M_Y} \mathcal{M}_{\text{PSh}(D)}$$

is fully faithful in the sense that it induces equivalences on all multimorphism animae. Moreover, if $\mathcal{O}$ admits operadic limits indexed by some ∞ -category I , then the induced functor on colors

$$\mathcal{O}_{\langle 1 \rangle} \rightarrow \text{PSh}(D)$$

preserves I -indexed limits.

Proof. Let $x_1, \dots, x_n, y$ be colors of $\mathcal{O}$. Since Y is strong symmetric monoidal by Proposition 18.2.2, we have natural equivalences

$$\begin{aligned} \mathcal{M}_{\text{PSh}(D)}(Yx_1, \dots, Yx_n; Yy) &\simeq \text{Hom}_{\text{PSh}(D)}(Yx_1 \otimes^{\text{Day}} \dots \otimes^{\text{Day}} Yx_n, Yy) \\ &\simeq \text{Hom}_{\text{PSh}(D)}(Y(x_1 \otimes \dots \otimes x_n), Yy) \\ &\simeq \text{Hom}_D(x_1 \otimes \dots \otimes x_n, y) \\ &\simeq \mathcal{O}(x_1, \dots, x_n; y). \end{aligned}$$

The first and last equivalences are induced by the unit $\eta_{\mathcal{O}}$ and agree with the identifications of Lemma 17.3.17. This proves full faithfulness. For the final claim, let $y \rightarrow F$ be an operadic limit cone in $\mathcal{O}_{\langle 1 \rangle}$. By Remark 18.1.2, its image in $D = \text{Env}(\mathcal{O})$ is an ordinary limit cone. Since the Yoneda embedding preserves limits, its image in $\text{PSh}(D)$ is again a limit cone. $\square$

Corollary 18.2.4. *Let $\mathcal{O}$ be a small ∞ -operad that admits finite operadic limits and put $D := \text{Env}(\mathcal{O})$. Then the color functor*

$$\mathcal{O}_{\langle 1 \rangle} \rightarrow \text{PSh}(D)$$

induced by the multiplicative Yoneda embedding is fully faithful, preserves finite limits, and has essential image closed under finite limits in $\text{PSh}(D)$. In particular, by Lemma 14.1.6, we may identify $\mathcal{O}$ with the full suboperad of $\mathcal{M}_{\text{PSh}(D)}$ spanned by this essential image. $\square$

18.3 Day convolution into arbitrary operads

The envelope allows us to construct Day convolution operads with arbitrary target operads. The key formal point is that every ∞ -operad embeds as a full suboperad of the multimorphism operad of its envelope, and that Day convolution restricts along full suboperads.

Lemma 18.3.1. *Let $i: \mathcal{O} \hookrightarrow \mathcal{Q}$ be a full suboperad. For every ∞ -operad $\mathcal{R}$, the induced functor*

$$\text{Fun}_{\text{Op}_{\infty}}(\mathcal{R}, \mathcal{O}) \rightarrow \text{Fun}_{\text{Op}_{\infty}}(\mathcal{R}, \mathcal{Q})$$

is fully faithful. Its essential image consists of those operad maps $\mathcal{R} \rightarrow \mathcal{Q}$ whose colors land in $\mathcal{O}_{\langle 1 \rangle} \subseteq \mathcal{Q}_{\langle 1 \rangle}$.

Proof. This is clear from the fact that the total category $\mathcal{O}^{\otimes}$ is the full subcategory of $\mathcal{Q}^{\otimes}$ spanned by those finite tuples whose entries are colors of $\mathcal{O}$. $\square$

Lemma 18.3.2. *Let $i: \mathcal{O} \hookrightarrow \mathcal{Q}$ be a full suboperad, let $\mathcal{C}$ be a symmetric monoidal ∞ -category, and assume that $\text{Day}(\mathcal{M}_{\mathcal{C}}, \mathcal{Q})$ exists. Let*

$$\mathcal{W} \subseteq \text{Day}(\mathcal{M}_{\mathcal{C}}, \mathcal{Q})$$

be the full suboperad spanned by those functors $\mathcal{C} \rightarrow \mathcal{Q}_{\langle 1 \rangle}$ whose values lie in $\mathcal{O}_{\langle 1 \rangle}$. Then $\mathcal{W}$ is a Day convolution operad $\text{Day}(\mathcal{M}_{\mathcal{C}}, \mathcal{O})$.

Proof. The evaluation map for $\text{Day}(\mathcal{M}_{\mathcal{C}}, \mathcal{Q})$ restricts to a map

$$\mathcal{W} \times \mathcal{M}_{\mathcal{C}} \rightarrow \mathcal{O},$$

because on colors it sends a pair (F, c) to $F(c)$. We verify the defining universal property. Let $\mathcal{R}$ be an arbitrary ∞ -operad. By Lemma 18.3.1, maps $\mathcal{R} \rightarrow \mathcal{W}$ are precisely maps $\mathcal{R} \rightarrow \mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{Q})$ whose color functors land in $\mathcal{O}_{\langle 1 \rangle}$. Transposing along the Day convolution adjunction for $\mathcal{Q}$, these are precisely maps

$$\mathcal{R} \times \mathcal{M}_C \rightarrow \mathcal{Q}$$

whose colors land in $\mathcal{O}_{\langle 1 \rangle}$. Applying Lemma 18.3.1 again, this is the same as giving a map $\mathcal{R} \times \mathcal{M}_C \rightarrow \mathcal{O}$. This identification is natural in $\mathcal{R}$, hence $\mathcal{W}$ satisfies the universal property of $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{O})$. $\square$

Theorem 18.3.3. *Let $\mathcal{C}$ be a symmetric monoidal ∞ -category. For every ∞ -operad $\mathcal{O}$, the Day convolution operad $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{O})$ exists.*

Proof. Put $\mathcal{D} := \text{Env}(\mathcal{O})$. By Lemma 17.3.17, the unit

$$\eta_{\mathcal{O}}: \mathcal{O} \rightarrow \mathcal{M}_{\mathcal{D}}$$

is fully faithful: its essential image is the full suboperad spanned by the singleton tuples of colors in $\mathcal{D}$. We may therefore identify $\mathcal{O}$ with this full suboperad by Lemma 14.1.6. The Day convolution operad $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_{\mathcal{D}})$ exists by Theorem 16.2.4, so Lemma 18.3.2 gives the result. $\square$

Remark 18.3.4. A relative version of this result was proved by Lurie [Lur17, Theorem 2.2.6.2]: if $f: \mathcal{O}' \rightarrow \mathcal{P}$ is a cocartesian fibration of ∞ -operads, then the pullback functor $f^*: (\text{Op}_{\infty})_{/\mathcal{P}} \rightarrow (\text{Op}_{\infty})_{/\mathcal{O}'}$ admits a right adjoint

$$\text{Nm}_{\mathcal{O}'/\mathcal{P}}: (\text{Op}_{\infty})_{/\mathcal{O}'} \rightarrow (\text{Op}_{\infty})_{/\mathcal{P}}.$$

By precomposing this functor with f^* , it then follows that the product functor $\mathcal{O}' \times_{\mathcal{P}} -: (\text{Op}_{\infty})_{/\mathcal{P}} \rightarrow (\text{Op}_{\infty})_{/\mathcal{P}}$ admits a right adjoint

$$\mathcal{D}\text{ay}_{/\mathcal{P}}(\mathcal{O}', -): (\text{Op}_{\infty})_{/\mathcal{P}} \rightarrow (\text{Op}_{\infty})_{/\mathcal{P}}.$$

If $\mathcal{P} = \text{Comm}$ is the terminal ∞ -operad and $\mathcal{O}' \simeq \mathcal{M}_C$ for some $\mathcal{C}$, this specializes to Theorem 18.3.3.

Theorem 18.3.5 (Multimorphism formula for Day convolution). *Let $\mathcal{C}$ be a small symmetric monoidal ∞ -category and let $\mathcal{O}$ be an ∞ -operad. For every finite set I and functors $F_i, G: \mathcal{C} \rightarrow \mathcal{O}_{\langle 1 \rangle}$, there is a natural equivalence*

$$\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{O})(\{F_i\}_{i \in I}; G) \simeq \int_{\{c_i\}_{i \in I} \in \mathcal{C}^I} \mathcal{O}(\{F_i(c_i)\}_{i \in I}; G(\bigotimes_C^I c_i)).$$

Proof. Put $\mathcal{D} := \text{Env}(\mathcal{O})$ and identify $\mathcal{O}$ with the full suboperad of $\mathcal{M}_{\mathcal{D}}$ spanned by the singleton tuples, as in the proof of Theorem 18.3.3. By Lemma 18.3.2, the operad $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{O})$ is the full suboperad of $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_{\mathcal{D}})$ spanned by the $\mathcal{O}_{\langle 1 \rangle}$ -valued functors. Therefore its multimorphism animae agree with the corresponding multimorphism animae in $\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{M}_{\mathcal{D}})$. By Proposition 16.2.6, we get

$$\mathcal{D}\text{ay}(\mathcal{M}_C, \mathcal{O})(\{F_i\}_{i \in I}; G) \simeq \text{Nat} \left(\bigotimes_D^I \circ \prod_{i \in I} F_i, G \circ \bigotimes_C^I \right).$$

Using the end formula for natural transformations from Proposition 23.6.5, the right-hand side is equivalent to

$$\int_{\{c_i\}_{i \in I} \in C^I} \mathrm{Hom}_D \left(\bigotimes_D^I F_i(c_i), G(\bigotimes_C^I c_i) \right).$$

By Lemma 17.3.17, the integrand is naturally equivalent to

$$\mathcal{O}(\{F_i(c_i)\}_{i \in I}; G(\bigotimes_C^I c_i)).$$

Taking ends preserves this equivalence of integrands, giving the desired formula. $\square$

18.4 Stabilization of ∞ -operads

In Section 18.1, we introduced various classes of ∞ -operads with additional algebraic structure: pointed, semiadditive, additive, and stable. The goal of this section is to show that each of the four fully faithful inclusions

$$\mathrm{Op}_\infty^{\mathrm{pt}} \hookrightarrow \mathrm{Op}_\infty^*, \quad \mathrm{Op}_\infty^{\mathrm{sadd}} \hookrightarrow \mathrm{Op}_\infty^{\mathrm{prod}}, \quad \mathrm{Op}_\infty^{\mathrm{add}} \hookrightarrow \mathrm{Op}_\infty^{\mathrm{prod}} \quad \text{and} \quad \mathrm{Op}_\infty^{\mathrm{st}} \hookrightarrow \mathrm{Op}_\infty^{\mathrm{lex}}$$

admits a right adjoint. These right adjoints can be thought of as *cofreely* adding the respective algebraic structure to an ∞ -operad. The constructions rely on the Day convolution machinery of Chapter 16 and its extension to arbitrary operadic targets in Section 18.3.

We treat the case of spectrum objects in detail; the pointed, semiadditive, and additive cases follow the same pattern and are treated more briefly at the end of the section.

18.4.1 Spectrum objects

Construction 18.4.1. Let $\mathcal{O}$ be an ∞ -operad that admits finite operadic limits. Consider the ∞ -category $\mathrm{An}_*^{\mathrm{fin}}$ equipped with the smash product monoidal structure from Definition 16.3.5. We define

$$\mathrm{Sp}(\mathcal{O}) \subseteq \mathrm{Day}(\mathcal{M}_{\mathrm{An}_*^{\mathrm{fin}}}, \mathcal{O})$$

to be the full suboperad whose colors are those functors $F: \mathrm{An}_*^{\mathrm{fin}} \rightarrow \mathcal{O}_{\langle 1 \rangle}$ that are reduced and excisive. Note that

$$\mathrm{Sp}(\mathcal{O})_{\langle 1 \rangle} \simeq \mathrm{Sp}^{\mathrm{exc}}(\mathcal{O}_{\langle 1 \rangle})$$

by construction; through Proposition 16.5.10, we identify this underlying ∞ -category with the abstract stabilization $\mathrm{Sp}(\mathcal{O}_{\langle 1 \rangle})$. The monoidal unit S^0 defines an operad map $\mathrm{Comm} \rightarrow \mathcal{M}_{\mathrm{An}_*^{\mathrm{fin}}}$, and restricting the evaluation map along it defines a morphism of ∞ -operads

$$\Omega^\infty: \mathrm{Sp}(\mathcal{O}) \rightarrow \mathcal{O}.$$

Lemma 18.4.2. *Let D be a small symmetric monoidal ∞ -category and equip $P := \mathrm{PSh}(D)$ with the Day convolution symmetric monoidal structure. Then $\mathrm{Sp}(\mathcal{M}_P)$ is a stable ∞ -operad.*

Proof. Put $A := \mathrm{An}_*^{\mathrm{fin}}$. The Day convolution structure on P identifies $\mathcal{M}_P$ with $\mathrm{Day}(\mathcal{M}_{D^{\mathrm{op}}}, \mathcal{M}_{\mathrm{An}})$. Thus, using the universal property of Day convolution twice, $\mathrm{Day}(\mathcal{M}_A, \mathcal{M}_P)$ identifies with

$$\mathrm{Day}(\mathcal{M}_{D^{\mathrm{op}}}, \mathrm{Day}(\mathcal{M}_A, \mathcal{M}_{\mathrm{An}})).$$

Under this identification, $\mathcal{S}p(\mathcal{M}_P)$ is obtained by restricting the inner $\mathcal{D}ay(\mathcal{M}_A, \mathcal{M}_{\text{An}})$ to reduced excisive functors, using Lemma 18.3.2. By Theorem 16.6.1 and the construction of the symmetric monoidal localization there, this restriction is precisely the multimorphism operad $\mathcal{M}_{\text{Sp}}$. Hence

$$\mathcal{S}p(\mathcal{M}_P) \simeq \mathcal{D}ay(\mathcal{M}_{D^{\text{op}}}, \mathcal{M}_{\text{Sp}}).$$

The latter is the multimorphism operad of $\text{Fun}(D^{\text{op}}, \text{Sp})$ with Day convolution. Since Sp is stable and its tensor product preserves colimits in each variable, so does the Day tensor product: for a fixed functor in all but one variable, it is computed as a left Kan extension of a pointwise tensor product, and both operations preserve colimits in the remaining variable. It follows that $\mathcal{S}p(\mathcal{M}_P)$ is stable. $\square$

Proposition 18.4.3. *Let $\mathcal{O}$ be an ∞ -operad that admits finite operadic limits. Then $\mathcal{S}p(\mathcal{O})$ is stable.*

Proof. Choose a universe in which $D := \text{Env}(\mathcal{O})$ is small, let $P := \text{PSh}(D)$ be the corresponding presheaf category, and suppress the universe decoration from the notation. Let

$$\iota: \mathcal{O} \hookrightarrow \mathcal{M}_P$$

be the multiplicative Yoneda embedding of Proposition 18.2.3. By Lemma 18.3.2, the operad $\mathcal{S}p(\mathcal{O})$ is the full suboperad of $\mathcal{S}p(\mathcal{M}_P)$ spanned by those reduced excisive functors $\text{An}_*^{\text{fin}} \rightarrow P$ whose values lie in the essential image of $\mathcal{O}_{\langle 1 \rangle} \rightarrow P$.

By Corollary 18.2.4, the essential image of $\mathcal{O}_{\langle 1 \rangle} \rightarrow P$ is closed under finite limits. Since ι is fully faithful and preserves finite limits, it also reflects them. Thus reducedness and excisiveness can be checked either in $\mathcal{O}_{\langle 1 \rangle}$ or after applying ι . Hence the essential image of the fully faithful functor

$$\text{Sp}^{\text{exc}}(\mathcal{O}_{\langle 1 \rangle}) \rightarrow \text{Sp}^{\text{exc}}(P)$$

is closed under finite limits, since limits of reduced excisive functors are computed pointwise. The source and target of this functor are stable by Proposition 16.5.7, and a finite-limit-preserving functor between stable ∞ -categories is exact. It follows that this essential image is closed under finite colimits as well. Since $\mathcal{S}p(\mathcal{M}_P)$ is stable by Lemma 18.4.2, the claim follows from Lemma 18.1.8. $\square$

18.4.2 The end calculation

The remaining point is to show that if $\mathcal{O}$ is already stable, then evaluation at S^0 is an equivalence of ∞ -operads between $\mathcal{S}p(\mathcal{O})$ and $\mathcal{O}$. The underlying categorical statement is Lemma 16.5.8; what is missing is full faithfulness on multimorphism animae. This is where the end formula from Theorem 18.3.5 enters.

Lemma 18.4.4. *Let $T: (\text{An}_*^{\text{fin}})^{\text{op}} \times \text{An}_*^{\text{fin}} \rightarrow \text{An}$ be a functor with the following properties:*

- (1) *For every $L \in \text{An}_*^{\text{fin}}$, the functor $T(-, L): (\text{An}_*^{\text{fin}})^{\text{op}} \rightarrow \text{An}$ sends finite colimits in An_*^{fin} to limits.*
- (2) *For every $K \in \text{An}_*^{\text{fin}}$, the functor $T(K, -): \text{An}_*^{\text{fin}} \rightarrow \text{An}$ is reduced and excisive.*

Then the projection from the end to the component at S^0 ,

$$\int_{K \in \text{An}_*^{\text{fin}}} T(K, K) \rightarrow T(S^0, S^0),$$

is an equivalence.

Proof. Let $A := \mathbf{An}_*^{\text{fin}}$ and let $j: A \hookrightarrow \mathbf{An}_*$ be the inclusion. Also let

$$W: \mathbf{An}_* \rightarrow \mathbf{PSh}(A), \quad X \mapsto \text{Hom}_{\mathbf{An}_*}(-, X)|_A$$

be the restricted Yoneda embedding from Lemma 22.3.9. By currying, the functor T determines a functor

$$\tilde{T}: A \rightarrow \mathbf{PSh}(A), \quad L \mapsto T(-, L).$$

By assumption (1) and Lemma 22.3.9, the functor $\tilde{T}$ factors uniquely as $W \circ T'$ for a functor $T': A \rightarrow \mathbf{An}_*$. Since W is fully faithful and preserves and reflects limits, and since limits in presheaf categories are computed pointwise, assumption (2) implies that T' is reduced and excisive.

Let $Y_A: A \rightarrow \mathbf{PSh}(A)$ denote the Yoneda embedding. Using the end formula for natural transformations, the Yoneda lemma, and the identity $W \circ j \simeq Y_A$, we obtain equivalences

$$\int_{K \in A} T(K, K) \simeq \text{Nat}(Y_A, \tilde{T}) \simeq \text{Nat}(j, T').$$

Since T' is reduced and excisive, Lemma 16.5.14 gives

$$\text{Nat}(j, T') \simeq \text{Nat}(P_1 j, T') \simeq \text{Hom}_{\mathbf{An}_*}(S^0, T'(S^0)).$$

Finally, by the definition of W we have

$$\text{Hom}_{\mathbf{An}_*}(S^0, T'(S^0)) \simeq W(T'(S^0))(S^0) \simeq T(S^0, S^0).$$

Tracing through the construction identifies this composite with projection from the end to the component at S^0 . $\square$

Proposition 18.4.5. *If $\mathcal{O}$ is stable, then the evaluation map*

$$\Omega^\infty: \mathcal{Sp}(\mathcal{O}) \rightarrow \mathcal{O}$$

is an equivalence of ∞ -operads.

Proof. On underlying ∞ -categories this is precisely Lemma 16.5.8. By Lemma 14.1.6, it remains to prove that Ω^∞ is fully faithful.

Let I be a finite set, and let $\{F_i: \mathbf{An}_*^{\text{fin}} \rightarrow \mathcal{O}_{\langle 1 \rangle}\}_{i \in I}$ and $G: \mathbf{An}_*^{\text{fin}} \rightarrow \mathcal{O}_{\langle 1 \rangle}$ be reduced excisive functors. By Theorem 18.3.5, the source of the induced map on multimorphism animae is

$$\int_{\{K_i\}_{i \in I} \in (\mathbf{An}_*^{\text{fin}})^I} \mathcal{O}(\{F_i(K_i)\}_{i \in I}; G(\bigwedge_{i \in I} K_i)).$$

By naturality of Theorem 18.3.5 in the evaluation operad map and the identification $\bigwedge_{i \in I} S^0 \simeq S^0$, the map induced by Ω^∞ is the projection from this end to the component where all K_i are equal to S^0 .

We prove a slightly stronger statement by induction on the cardinality of I : the same projection is an equivalence after adjoining any fixed finite tuple of spectator colors $z_1, \dots, z_m$ to the inputs of every multimorphism anima. For $I = \emptyset$ there is nothing to prove. Otherwise choose $i_0 \in I$ and put

$I' := I \setminus \{i_0\}$. By Fubini for ends, the relevant end is the end over $K \in \mathbf{An}_*^{\text{fin}}$ of $T(K, K)$, where for $K, L \in \mathbf{An}_*^{\text{fin}}$ we define

$$T(K, L) := \int_{\{K_i\}_{i \in I'} \in (\mathbf{An}_*^{\text{fin}})^{I'}} \mathcal{O}(z_1, \dots, z_m, F_{i_0}(K), \{F_i(K_i)\}_{i \in I'}; G(L \wedge \bigwedge_{i \in I'} K_i)).$$

We claim that T satisfies the hypotheses of Lemma 18.4.4. Since $\mathcal{O}$ is stable, a reduced excisive functor $\mathbf{An}_*^{\text{fin}} \rightarrow \mathcal{O}_{\langle 1 \rangle}$ preserves finite colimits: finite colimits in $\mathbf{An}_*^{\text{fin}}$ are generated by the initial object and pushouts, and a square in a stable ∞ -category is a pushout if and only if it is a pullback. Therefore F_{i_0} sends finite colimits in $\mathbf{An}_*^{\text{fin}}$ to finite operadic colimits in $\mathcal{O}$. Since finite operadic colimits in an input variable are detected by mapping out of them, $T(-, L)$ sends finite colimits to limits. Similarly, the functor $L \mapsto G(L \wedge \bigwedge_{i \in I'} K_i)$ is reduced and sends pushout squares to finite limits in $\mathcal{O}_{\langle 1 \rangle}$, and these limits are operadic. Thus $T(K, -)$ is reduced and excisive. Ends preserve the relevant limits throughout, and the fixed spectator colors do not affect either verification.

Applying Lemma 18.4.4 collapses the i_0 -variable to S^0 . The stronger induction hypothesis, with $F_{i_0}(S^0)$ adjoined to the spectator colors, now collapses the remaining variables and gives an equivalence from the displayed end to

$$\mathcal{O}(\{F_i(S^0)\}_{i \in I}; G(S^0)),$$

which is the multimorphism anima in $\mathcal{O}$ between the images under Ω^∞ . This proves full faithfulness. $\square$

Theorem 18.4.6 (Stabilization of operads). *Let $\mathcal{O}$ be an ∞ -operad that admits finite operadic limits. Then the ∞ -operad $Sp(\mathcal{O})$ is stable, and the resulting functor*

$$Sp: \mathbf{Op}_\infty^{\text{lex}} \rightarrow \mathbf{Op}_\infty^{\text{st}}$$

is right adjoint to the inclusion functor.

Proof. The first claim is Proposition 18.4.3. We now prove the adjunction. Postcomposition with a map in $\mathbf{Op}_\infty^{\text{lex}}$ preserves reduced excisive functors, since its color functor preserves finite limits. Thus the construction of $Sp(\mathcal{O})$ is functorial in $\mathcal{O} \in \mathbf{Op}_\infty^{\text{lex}}$, and evaluation at S^0 gives a natural transformation

$$\Omega^\infty: Sp \rightarrow \text{id}_{\mathbf{Op}_\infty^{\text{lex}}}.$$

By the dual of the recognition criterion for Bousfield localizations from Proposition 21.8.9, applied in $(\mathbf{Op}_\infty^{\text{lex}})^{\text{op}}$, it is enough to show that $\Omega_\mathcal{O}^\infty$ is an equivalence whenever $\mathcal{O}$ is stable, and that

$$Sp(\Omega_\mathcal{O}^\infty): Sp(Sp(\mathcal{O})) \rightarrow Sp(\mathcal{O})$$

is an equivalence for every $\mathcal{O} \in \mathbf{Op}_\infty^{\text{lex}}$. The first assertion is Proposition 18.4.5.

For the second assertion, observe that $Sp(Sp(\mathcal{O}))$ may be identified with the full suboperad of $\mathcal{Day}(\mathcal{M}_{\mathbf{An}_*^{\text{fin}} \times \mathbf{An}_*^{\text{fin}}}, \mathcal{O})$ spanned by bifunctors $\mathbf{An}_*^{\text{fin}} \times \mathbf{An}_*^{\text{fin}} \rightarrow \mathcal{O}_{\langle 1 \rangle}$ which are reduced and excisive in each variable. Under this identification, the map $Sp(\Omega_\mathcal{O}^\infty)$ evaluates in one variable at S^0 , while the map

$$\Omega_{Sp(\mathcal{O})}^\infty: Sp(Sp(\mathcal{O})) \rightarrow Sp(\mathcal{O})$$

evaluates in the other variable at S^0 . These two maps are conjugate by the symmetry of $\mathbf{An}_*^{\text{fin}} \times \mathbf{An}_*^{\text{fin}}$. Since $Sp(\mathcal{O})$ is stable by the first part of the theorem, Proposition 18.4.5 shows that $\Omega_{Sp(\mathcal{O})}^\infty$ is an equivalence. Hence $Sp(\Omega_\mathcal{O}^\infty)$ is an equivalence as well. $\square$

Recall from Definition 11.1.15 that a symmetric monoidal ∞ -category C is called *stably symmetric monoidal* if C is stable and the tensor product $- \otimes -: C \times C \rightarrow C$ is exact in both variables; equivalently, if $\mathcal{M}_C$ is a stable ∞ -operad.

Corollary 18.4.7. *Let C be a symmetric monoidal ∞ -category with finite limits, and assume that $\mathrm{Sp}(\mathcal{M}_C)$ is represented by a symmetric monoidal structure on $\mathrm{Sp}(C)$. Then $\Omega^\infty: \mathrm{Sp}(C) \rightarrow C$ admits a canonical lax symmetric monoidal structure. Furthermore, for every stably symmetric monoidal ∞ -category D , composition with Ω^∞ induces an equivalence of ∞ -categories*

$$\Omega^\infty \circ -: \mathrm{Fun}^{\otimes\text{-lax}, \text{lex}}(D, \mathrm{Sp}(C)) \xrightarrow{\sim} \mathrm{Fun}^{\otimes\text{-lax}, \text{lex}}(D, C),$$

where the superscript lex denotes the full subcategories of lax symmetric monoidal functors whose underlying functors preserve finite limits. On the left-hand side this is equivalently the condition that the underlying functor be exact.

Proof. Applying the adjunction of Theorem 18.4.6 to finite-operadic-limit-preserving maps $\mathcal{M}_D \rightarrow \mathcal{M}_C$ gives the asserted equivalence on maximal animae. The enhancement to an equivalence of ∞ -categories, including non-invertible lax monoidal natural transformations, is [Nik16, Corollary 4.13]; it is obtained by applying the functoriality of operadic stabilization to the corresponding operad-map categories. $\square$

Remark 18.4.8. The construction $\mathrm{Sp}(\mathcal{M}_C)$ always produces an ∞ -operad whose underlying ∞ -category is $\mathrm{Sp}(C)$, but this operad need not be the multimorphism operad of a symmetric monoidal ∞ -category. Thus the representability hypothesis in Corollary 18.4.7 is genuine. It does hold when C is presentably symmetric monoidal; this will follow from the mode-theoretic construction in Corollary 18.5.5.

18.4.3 Pointing and additivization

The pointed, semiadditive and additive cases work similarly.

Definition 18.4.9. If $\mathcal{O}$ is an ∞ -operad with an operadic terminal object, equip $[1]$ with the symmetric monoidal structure given by the minimum, as used in the proof of Lemma 16.3.3, and denote by

$$\mathcal{O}_* \subseteq \mathcal{D}\mathrm{ay}(\mathcal{M}_{[1]}, \mathcal{O})$$

the full suboperad whose colors are those functors $F: [1] \rightarrow \mathcal{O}_{\langle 1 \rangle}$ satisfying $F(0) \simeq *$. If in addition $\mathcal{O}$ has finite operadic products, equip $\mathrm{Span}(\mathrm{Fin})$ with the symmetric monoidal structure induced by the cartesian product of finite sets, and denote by

$$\mathrm{CGrp}(\mathcal{O}) \subseteq \mathrm{CMon}(\mathcal{O}) \subseteq \mathcal{D}\mathrm{ay}(\mathcal{M}_{\mathrm{Span}(\mathrm{Fin})}, \mathcal{O})$$

the full suboperads whose colors are those functors $F: \mathrm{Span}(\mathrm{Fin}) \rightarrow \mathcal{O}_{\langle 1 \rangle}$ that are commutative groups or commutative monoids in $\mathcal{O}_{\langle 1 \rangle}$, respectively.

Notice that this is not the cartesian monoidal structure on $\mathrm{Span}(\mathrm{Fin})$: its categorical products are disjoint unions by Lemma 13.3.5. The cartesian product of finite sets is the monoidal structure used in the Day convolution description of Proposition 16.4.1.

It is immediate from the definitions that the underlying ∞ -categories of these ∞ -operads are given by

$$(O_*)_{\langle 1 \rangle} \simeq (O_{\langle 1 \rangle})_*, \quad CMon(O)_{\langle 1 \rangle} \simeq CMon(O_{\langle 1 \rangle}) \quad \text{and} \quad CGrp(O)_{\langle 1 \rangle} \simeq CGrp(O_{\langle 1 \rangle}),$$

justifying the notation.

Lemma 18.4.10 (Day convolution on pointed objects). *Let C be an ∞ -category with finite products and finite colimits such that its cartesian product preserves finite colimits separately in both variables. Then there is a natural equivalence*

$$(\mathcal{M}_{(C, \times)})_* \simeq \mathcal{M}_{(C_*, \wedge)}.$$

Proof. The operad $(\mathcal{M}_{(C, \times)})_*$ is the full suboperad of the Day convolution operad on $\text{Ar}(C)$ spanned by the arrows with source the terminal object. The monoidal localization

$$\text{cofib}: \text{Ar}(C)^\square \longrightarrow (C_*, \wedge)$$

of Lemma 16.3.4 identifies this full suboperad with the multimorphism operad of its local objects. $\square$

Lemma 18.4.11. *If O is pointed, then evaluation at the target defines an equivalence of ∞ -operads*

$$\text{ev}_1: O_* \rightarrow O.$$

Proof. On underlying ∞ -categories, evaluation identifies the category of arrows from the zero object with $O_{\langle 1 \rangle}$. It remains to prove full faithfulness on multimorphism anima.

Let $\{F_i\}_{i \in I}$ and G be colors of O_* . By Theorem 18.3.5, their multimorphism anima is an end over $[1]^I$. We collapse its variables one at a time. After applying Fubini, the variable to be collapsed appears as the end of a bifunctor $T: [1]^{\text{op}} \times [1] \rightarrow \text{An}$, allowing arbitrary fixed spectator colors. This end is the pullback

$$T(0, 0) \times_{T(0, 1)} T(1, 1).$$

Both $T(0, 0)$ and $T(0, 1)$ are contractible because $F_i(0)$ is the zero object and hence operadic initial. Thus this pullback is equivalent to $T(1, 1)$. Fubini for ends and induction, with the values $F_i(1)$ already obtained treated as spectator colors, identify the original end with

$$O(\{F_i(1)\}_{i \in I}; G(1)).$$

By naturality of the end formula, this equivalence is the map induced by evaluation at 1. $\square$

For the semiadditive and additive cases below, the coherence of the universal properties is supplied by Nikolaus [Nik16, Theorem 5.7]; we include the argument that identifies the resulting operads and their underlying algebraic structure. The pointed case will be proved internally.

Theorem 18.4.12 (Additivization of operads, [Nik16, Theorem 5.7]). *Let O be an object in Op_∞^* or $\text{Op}_\infty^{\text{prod}}$ respectively. Then the operads O_* , $CMon(O)$ and $CGrp(O)$ are pointed, semiadditive, and additive, respectively. Moreover, the resulting functors*

$$(-)_*: \text{Op}_\infty^* \rightarrow \text{Op}_\infty^{\text{pt}}, \quad CMon: \text{Op}_\infty^{\text{prod}} \rightarrow \text{Op}_\infty^{\text{sadd}} \quad \text{and} \quad CGrp: \text{Op}_\infty^{\text{prod}} \rightarrow \text{Op}_\infty^{\text{add}}$$

are right adjoint to the respective inclusion functors.

Proof. We first prove the pointed case. Choose a universe in which $\text{Env}(\mathcal{O})$ is small and let $\mathcal{P}$ be the corresponding presheaf category. The multiplicative Yoneda embedding and Lemma 18.3.2 give full embeddings

$$\mathcal{O}_* \hookrightarrow (\mathcal{M}_{\text{Env}(\mathcal{O})})_* \hookrightarrow (\mathcal{M}_{\mathcal{P}})_*$$

where the last target is understood in the chosen universe. By Lemma 18.4.10, this last operad is the multimorphism operad of the pointed presheaf category and is therefore pointed by Example 18.1.7. The suboperad $\mathcal{O}_*$ contains its zero object, since the multiplicative Yoneda embedding preserves the operadic terminal object of $\mathcal{O}$. It follows from Lemma 18.1.8(1) that $\mathcal{O}_*$ is pointed.

Evaluation at the target gives a natural operad map $\mathcal{O}_* \rightarrow \mathcal{O}$. If $\mathcal{O}$ is pointed, this map is an equivalence by Lemma 18.4.11. Applying the pointed-object construction a second time gives the same conclusion because $\mathcal{O}_*$ is pointed. The dual of Proposition 21.8.9, applied in $(\text{Op}_\infty^*)^{\text{op}}$, therefore identifies $(-)_*$ as right adjoint to the inclusion of pointed operads.

For commutative monoids and groups, choose a universe in which $\mathcal{O}$ is small and apply the multiplicative Yoneda embedding to regard $\mathcal{O}$ as a full suboperad of the presheaf operad on $\text{Env}(\mathcal{O})$. In that presheaf category, Proposition 16.4.1 supplies the Day convolution operads of commutative monoids and commutative groups. Their underlying categories are semiadditive and additive by Proposition 5.3.19, and their tensor products preserve colimits separately in both variables. Hence their finite biproducts are operadic. These biproducts are computed pointwise. Since the multiplicative Yoneda image is closed under finite products, the full suboperads consisting of objects valued in that image are closed under these biproducts. They are precisely $\text{CMon}(\mathcal{O})$ and $\text{CGrp}(\mathcal{O})$, so Lemma 18.1.8(2) shows that the former is semiadditive and the latter additive.

If $\mathcal{P}$ is semiadditive, every color of $\mathcal{P}$ has a commutative-monoid structure supplied by its biproducts, compatibly with all multimorphisms because the biproducts are operadic. If $\mathcal{P}$ is additive, these structures are grouplike. The cited theorem supplies the coherent lifts through $\text{CMon}(\mathcal{O}) \rightarrow \mathcal{O}$ and $\text{CGrp}(\mathcal{O}) \rightarrow \mathcal{O}$ and identifies them with the right adjoints to the respective inclusions. $\square$

Corollary 18.4.13. *Applied to the cartesian multimorphism operad $\mathcal{M}_{(\text{An}, \times)}$, the four operadic constructions recover the symmetric monoidal structures of Chapter 16:*

$$\begin{aligned} (\mathcal{M}_{(\text{An}, \times)})_* &\simeq \mathcal{M}_{(\text{An}_*, \wedge)}, & \text{CMon}(\mathcal{M}_{(\text{An}, \times)}) &\simeq \mathcal{M}_{\text{CMon}(\text{An})}, \\ \text{CGrp}(\mathcal{M}_{(\text{An}, \times)}) &\simeq \mathcal{M}_{\text{CGrp}(\text{An})}, & \text{Sp}(\mathcal{M}_{(\text{An}, \times)}) &\simeq \mathcal{M}_{\text{Sp}}. \end{aligned}$$

Proof. The pointed case is Lemma 18.4.10, the semiadditive and additive cases are the symmetric monoidal localizations of Proposition 16.4.1, and the stable case is Theorem 16.6.1. $\square$

18.5 Modes and universal properties

The symmetric monoidal categories and transition functors constructed in Chapter 16 fit into a sequence of increasingly linear settings:

$$(\text{An}, \times) \xrightarrow{(-)_+} (\text{An}_*, \wedge) \xrightarrow{F^{\text{CMon}}} (\text{CMon}(\text{An}), \otimes) \xrightarrow{(-)^{\text{grp}}} (\text{CGrp}(\text{An}), \otimes) \xrightarrow{\mathbf{B}^\infty} (\text{Sp}, \otimes).$$

We have constructed the monoidal structures and transition functors directly. We now explain their common universal property. The language of *modes* packages pointedness, semiadditivity, additivity, and stability as instances of the same kind of structure.

Recall from Section 22.4 the symmetric monoidal ∞ -category Pr^{L} of presentable ∞ -categories and colimit-preserving functors. Its unit is An , and its tensor product is characterized by equivalences

$$\mathrm{Fun}^{\mathrm{L}}(C \otimes D, E) \simeq \mathrm{Fun}^{\mathrm{L}, \mathrm{L}}(C \times D, E),$$

where the right-hand side is the full subcategory of functors that preserve small colimits separately in both variables. By Remark 22.5.2, a commutative algebra object in Pr^{L} is equivalently a presentably symmetric monoidal ∞ -category.

Definition 18.5.1 (Mode). A *mode* is a commutative algebra $M \in \mathrm{CAlg}(\mathrm{Pr}^{\mathrm{L}})$ whose multiplication

$$M \otimes M \longrightarrow M$$

is an equivalence. A presentable ∞ -category C is called *M-local* if the unit of M induces an equivalence

$$C \simeq \mathrm{An} \otimes C \longrightarrow M \otimes C.$$

The terminology is taken from [CSY21, Section 5]. We only need the following basic formal properties of modes, which are instances of the general theory of idempotent commutative algebras.

Proposition 18.5.2 (Modes as properties, [Lur17, Propositions 4.8.2.4, 4.8.2.9 and 4.8.2.10]). *Let M be a mode.*

- (1) *A presentable ∞ -category admits an M -module structure if and only if it is M -local, and such a structure is unique.*
- (2) *Tensoring with M defines a Bousfield localization*

$$M \otimes -: \mathrm{Pr}^{\mathrm{L}} \longrightarrow (\mathrm{Pr}^{\mathrm{L}})^{M\text{-loc}},$$

where $(\mathrm{Pr}^{\mathrm{L}})^{M\text{-loc}}$ is the full subcategory of M -local presentable ∞ -categories.

- (3) *If D is M -local, then precomposition with the unit $C \rightarrow M \otimes C$ induces an equivalence*

$$\mathrm{Fun}^{\mathrm{L}}(M \otimes C, D) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{L}}(C, D).$$

Thus a mode classifies a property of presentable ∞ -categories which, whenever it exists, carries a unique coherent structure and can be imposed by a universal localization. The modes relevant here are the following.

Theorem 18.5.3 (The basic modes, [GGN15, Sections 3–5]). *The presentably symmetric monoidal ∞ -categories*

$$(\mathrm{An}_*, \wedge), \quad (\mathrm{CMon}(\mathrm{An}), \otimes), \quad (\mathrm{CGrp}(\mathrm{An}), \otimes), \quad (\mathrm{Sp}, \otimes)$$

are modes. For every presentable ∞ -category C , there are natural equivalences

$$\begin{aligned} \mathrm{An}_* \otimes C &\simeq C_*, & \mathrm{CMon}(\mathrm{An}) \otimes C &\simeq \mathrm{CMon}(C), \\ \mathrm{CGrp}(\mathrm{An}) \otimes C &\simeq \mathrm{CGrp}(C), & \mathrm{Sp} \otimes C &\simeq \mathrm{Sp}(C). \end{aligned}$$

Consequently, these four modes classify the properties recorded in Table 18.1. The functors in the sequence at the beginning of this section are morphisms of modes, meaning morphisms in $\mathrm{CAlg}(\mathrm{Pr}^{\mathrm{L}})$.

Mode	Local presentable ∞ -categories
$\mathbf{An}_*$	Pointed
$\mathbf{CMon}(\mathbf{An})$	Semiadditive
$\mathbf{CGrp}(\mathbf{An})$	Additive
$\mathbf{Sp}$	Stable

Table 18.1: The four basic modes of Theorem 18.5.3 and the properties of presentable ∞ -categories that they classify.

Proposition 18.5.4 (Multiplicative universal property of commutative monoids). *Let C and D be presentably symmetric monoidal ∞ -categories, and assume that D is semiadditive. Restriction along the free commutative monoid functor induces an equivalence*

$$\mathrm{Fun}^{\mathrm{L},\otimes}(\mathbf{CMon}(C), D) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{L},\otimes}(C, D)$$

of ∞ -categories.

Proof. Set $M := \mathbf{CMon}(\mathbf{An})$. By Theorem 18.5.3, M is a mode and there is a natural equivalence

$$M \otimes C \simeq \mathbf{CMon}(C).$$

Under this equivalence, the localization unit $C \simeq \mathbf{An} \otimes C \rightarrow M \otimes C$ is the free semiadditive functor $C \rightarrow \mathbf{CMon}(C)$. The symmetric monoidal refinement of this functor is the one constructed from Day convolution in Proposition 16.4.1.

Since D is semiadditive, it is M -local by Theorem 18.5.3. The localization $M \otimes -$ of Proposition 18.5.2 is induced by the idempotent commutative algebra M , so it restricts to a localization on commutative algebra objects of $\mathbf{Pr}^{\mathrm{L}}$. Its universal property gives an equivalence

$$\mathrm{Fun}^{\mathrm{L},\otimes}(M \otimes C, D) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{L},\otimes}(C, D).$$

Using the displayed identification of $M \otimes C$ with $\mathbf{CMon}(C)$ gives the result. Since the argument takes place in commutative algebra objects of $\mathbf{Pr}^{\mathrm{L}}$, it supplies the full symmetric monoidal coherence rather than only the individual tensor comparisons. $\square$

Corollary 18.5.5 (Presentable operadic stabilization, [Nik16, Proposition 4.9]). *Let C be a presentably symmetric monoidal ∞ -category. Then $\mathbf{Sp}(C)$ admits a canonical presentably symmetric monoidal structure, and the operadic stabilization $\mathbf{Sp}(\mathcal{M}_C)$ is represented by its multimorphism operad:*

$$\mathbf{Sp}(\mathcal{M}_C) \simeq \mathcal{M}_{\mathbf{Sp}(C)}.$$

In particular, the representability hypothesis in Corollary 18.4.7 is automatic in the presentable case.

Proof. The presentably symmetric monoidal structure on C is a commutative algebra object of $\mathbf{Pr}^{\mathrm{L}}$. Tensoring it with the commutative algebra $\mathbf{Sp}$ gives another commutative algebra

$$\mathbf{Sp} \otimes C \in \mathbf{CAlg}(\mathbf{Pr}^{\mathrm{L}}).$$

By Theorem 18.5.3, its underlying presentable ∞ -category is naturally equivalent to $\mathbf{Sp}(C)$. This equips $\mathbf{Sp}(C)$ with a presentably symmetric monoidal structure. The unit map $\mathbf{An} \rightarrow \mathbf{Sp}$ of the

mode induces a strongly monoidal stabilization functor $C \simeq \mathbf{An} \otimes C \rightarrow \mathbf{Sp} \otimes C$, whose right adjoint $\Omega^\infty: \mathbf{Sp}(C) \rightarrow C$ is lax symmetric monoidal.

Under the reduced-excise model of stabilization, the resulting multimorphism operad is the full suboperad of the Day convolution operad on $\mathbf{Fun}(\mathbf{An}_*^{\text{fin}}, C)$ spanned by the reduced excise functors. This is exactly the construction of $\mathcal{S}p(\mathcal{M}_C)$ from Section 18.4, giving the displayed equivalence. $\square$

The classification says, for example, that the free semiadditive presentable category generated by C is $\mathbf{CMon}(C)$, while its free stable presentable category is $\mathbf{Sp}(C)$. This is different from the cofree universal properties proved in Theorems 18.4.6 and 18.4.12. For example, operadic stabilization gives an equivalence

$$\mathbf{Fun}^{\otimes\text{-lax,lex}}(D, \mathbf{Sp}(C)) \simeq \mathbf{Fun}^{\otimes\text{-lax,lex}}(D, C)$$

for a stably symmetric monoidal D , whereas the mode $\mathbf{Sp}$ gives an equivalence

$$\mathbf{Fun}^{\mathbf{L}}(\mathbf{Sp}(C), D) \simeq \mathbf{Fun}^{\mathbf{L}}(C, D)$$

for a stable presentable D . The first is a mapping-into, cofree property for finite-limit-preserving functors and does not require presentability; the second is a mapping-out, free property for colimit-preserving functors and only concerns presentable categories. When the functors in question occur as adjoints, the two properties are closely related, but neither statement subsumes the other.

Applied to $C = \mathbf{An}$, the mode-theoretic classification gives the promised multiplicative mapping-out universal properties. Write $\mathbf{Fun}^{\mathbf{L},\otimes}(C, D)$ for the ∞ -category of colimit-preserving symmetric monoidal functors.

Corollary 18.5.6 (Multiplicative universal properties). *Let D be a presentably symmetric monoidal ∞ -category.*

(1) *If D is pointed, then restriction along $\mathbf{An} \rightarrow \mathbf{An}_*$ induces an equivalence*

$$\mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}_*, D) \xrightarrow{\sim} \mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}, D).$$

(2) *If D is semiadditive, then restriction along $\mathbf{An} \rightarrow \mathbf{CMon}(\mathbf{An})$ induces an equivalence*

$$\mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{CMon}(\mathbf{An}), D) \xrightarrow{\sim} \mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}, D).$$

(3) *If D is additive, then restriction along $\mathbf{An} \rightarrow \mathbf{CGrp}(\mathbf{An})$ induces an equivalence*

$$\mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{CGrp}(\mathbf{An}), D) \xrightarrow{\sim} \mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}, D).$$

(4) *If D is stable, then restriction along $\mathbf{An} \rightarrow \mathbf{Sp}$ induces an equivalence*

$$\mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{Sp}, D) \xrightarrow{\sim} \mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}, D).$$

Proof. Let M be any of the four modes in Theorem 18.5.3. If D is M -local, then the localization universal property of the idempotent commutative algebra M gives an equivalence

$$\mathbf{Fun}^{\mathbf{L},\otimes}(M, D) \xrightarrow{\sim} \mathbf{Fun}^{\mathbf{L},\otimes}(\mathbf{An}, D)$$

by restriction along the unit $\mathbf{An} \rightarrow M$. The four characterizations of M -local categories in Theorem 18.5.3 give the assertions. In the semiadditive case, this is also the specialization $C = \mathbf{An}$ of Proposition 18.5.4. $\square$

The tensor unit A_n carries the initial commutative algebra structure in $\mathbf{Pr}^L$, so the ∞ -category on the right in each case is contractible. Thus A_{n*} , $\mathbf{CMon}(A_n)$, $\mathbf{CGrp}(A_n)$, and $\mathbf{Sp}$ are initial among pointed, semiadditive, additive, and stable presentably symmetric monoidal ∞ -categories, respectively. In particular, the monoidal structures and transition functors constructed above are determined uniquely by these universal properties.

Exercises

19 Algebras and modules

The theory of algebras and modules is not specific to spectra, but makes sense in any suitable monoidal ∞ -category. Defining the objects requires no new work: an associative algebra is an $\mathcal{A}ssoc$ -algebra and a left module is an $\mathcal{L}Mod$ -algebra, so both are instances of notions we already have. The work lies in showing that they behave like their classical counterparts, and this is what we carry out here, in the generality required for the study of ring spectra in Part I (Chapter 8).

In Section 19.1, we introduce algebras and modules, construct limits in module categories, and establish higher-categorical analogues of basic constructions from classical algebra, such as free modules. In Section 19.2, we study the relative tensor product $M \otimes_A N$ as the composition of bimodules and deduce the resulting restriction and extension of scalars functors. We then briefly record the alternative simplicial models for algebras and modules in Section 19.3, both as a conceptual explanation of colimits of modules and as preparation for the cut diagrams used in Section 19.4. The latter section constructs endomorphism algebras and explains their universal relation to algebra actions on a fixed object.

The remaining sections specialize the theory and apply it. In Section 19.5 we pass to stable ∞ -categories and spectra, culminating in the monogenic Morita theorem, and Section 19.6 packages categories of algebraic objects by monads. Finally, Section 19.7 applies the coherent algebra developed in this part to construct connective complex K-theory as a commutative ring spectrum.

The stable and spectral consequences use the model-theoretic universal properties of Section 18.5. They and the final application to connective complex K-theory also use results established earlier in Part I.

The theory developed here is deep and often technically demanding. We will therefore frequently refer to Lurie’s foundational text [Lur17], where these topics are treated in exhaustive detail. Whenever we invoke a result formulated for Lurie’s model of ∞ -operads, we transport it through the equivalence of Proposition 17.4.8.

19.1 Foundations

In order to talk about modules over associative algebras, recall from Example 12.2.9 and Corollary 17.4.10 the ‘left module operad’ $\mathcal{L}Mod$ and its associated ∞ -operad. It comes equipped with

the following morphisms of ∞ -operads:

$$\begin{array}{ccccc} \mathcal{A}ssoc & \xleftarrow{\mathfrak{a}} & \mathcal{L}Mod & \xleftarrow{\mathfrak{m}} & \mathcal{T}riv \\ & \searrow & \downarrow & & \\ & & \mathcal{A}ssoc. & & \end{array}$$

We may describe their total ∞ -categories $\mathcal{A}ssoc^{\otimes}$ and $\mathcal{L}Mod^{\otimes}$ as follows:

- The objects of $\mathcal{A}ssoc^{\otimes}$ are finite sets I . A morphism from I to J consists of a span

$$I \xleftarrow{f} K \xrightarrow{g} J$$

of finite sets together with a collection of linear orders $(\preceq_j)_{j \in J}$ on the fibers $g^{-1}(j)$. Composition is given by composition of spans and the canonical ordering.

- The objects of $\mathcal{L}Mod^{\otimes}$ are pairs (I_A, I_M) of finite sets. A morphism from (I_A, I_M) to (J_A, J_M) is a span

$$I_A \sqcup I_M \xleftarrow{f} K \xrightarrow{g} J_A \sqcup J_M$$

together with a collection of linear orders $(\preceq_j)_{j \in J_A \sqcup J_M}$ on the fibers $g^{-1}(j)$, such that g induces a bijection between $f^{-1}(I_M)$ and J_M and, for each $j \in J_M$, the unique element of $g^{-1}(j) \cap f^{-1}(I_M)$ is maximal in the order $\preceq_j$.

The inclusion $\mathfrak{a}: \mathcal{A}ssoc^{\otimes} \hookrightarrow \mathcal{L}Mod^{\otimes}$ sends I to $(I, \emptyset)$, while the inclusion $\mathfrak{m}: \mathcal{T}riv^{\otimes} \hookrightarrow \mathcal{L}Mod^{\otimes}$ sends I to $(\emptyset, I)$. The map $\mathcal{L}Mod^{\otimes} \rightarrow \mathcal{A}ssoc^{\otimes}$ sends (I_A, I_M) to $I_A \sqcup I_M$.

Definition 19.1.1. Let C be a monoidal ∞ -category, regarded as $\mathcal{L}Mod$ -monoidal by pullback along the collapse map $\mathcal{L}Mod \rightarrow \mathcal{A}ssoc$. We denote by

$$\mathbf{LMod}(C) := \mathbf{Alg}_{\mathcal{L}Mod}(C)$$

the ∞ -category of $\mathcal{L}Mod$ -algebras in C . Restriction along $\mathfrak{a}$ defines a functor $\mathfrak{a}^*: \mathbf{LMod}(C) \rightarrow \mathbf{Alg}(C) := \mathbf{Alg}_{\mathcal{A}ssoc}(C)$. For an associative algebra $A \in \mathbf{Alg}(C)$ we define the ∞ -category $\mathbf{LMod}_A(C)$ of *left modules over A* as the fiber

$$\begin{array}{ccc} \mathbf{LMod}_A(C) & \longrightarrow & \mathbf{LMod}(C) \\ \downarrow & & \downarrow \mathfrak{a}^* \\ * & \xrightarrow{A} & \mathbf{Alg}(C). \end{array}$$

Remark 19.1.2. Because of the adjunction $\mathbf{Env}: \mathbf{Op}_{\infty} \rightleftarrows \mathbf{Cat}_{\infty}^{\otimes} : \mathcal{M}$, an associative algebra contains the same data as a symmetric monoidal functor $\mathbf{Env}(\mathcal{A}ssoc) \rightarrow C$. The concrete description $\mathbf{Env}(\mathcal{A}ssoc) \simeq \mathbf{Fin}^{\mathbf{Alg}}$ was given in Example 17.4.12. In particular, the maps $\langle 0 \rangle \rightarrow \langle 1 \rangle$ and $\langle 2 \rangle \rightarrow \langle 1 \rangle$ corresponding to the canonical ordering $\{1 \leq 2\}$ provide structure maps

$$e: \mathbb{1} \rightarrow A \quad \text{and} \quad m: A \otimes A \rightarrow A.$$

Furthermore, the following two diagrams commute:

$$\begin{array}{ccc}
 A \otimes \mathbb{1} & \xrightarrow{\text{id}_A \otimes e} & A \otimes A \\
 \searrow \simeq & & \downarrow m \\
 & & A
 \end{array}
 \quad
 \begin{array}{ccc}
 A \otimes A \otimes A & \xrightarrow{m \otimes \text{id}_A} & A \otimes A \\
 \text{id}_A \otimes m \downarrow & & \downarrow m \\
 A \otimes A & \xrightarrow{m} & A.
 \end{array}$$

Similarly, a module over A is given by a symmetric monoidal functor $\text{Env}(\mathcal{LMod}) \rightarrow C$ which on $\text{Env}(\mathcal{A}ssoc)$ restricts to A . The map $(\langle 1 \rangle, \langle 1 \rangle) \rightarrow (\emptyset, \langle 1 \rangle)$ in $\text{Env}(\mathcal{LMod})$ coming from the fold map $\langle 1 \rangle \sqcup \langle 1 \rangle \rightarrow \langle 1 \rangle$ gives rise to a map

$$\text{act}: A \otimes M \rightarrow M,$$

and the relations in $\text{Env}(\mathcal{LMod})$ ensure that the following two diagrams commute:

$$\begin{array}{ccc}
 \mathbb{1} \otimes M & \xrightarrow{e \otimes \text{id}_M} & A \otimes M \\
 \searrow \simeq & & \downarrow \text{act} \\
 & & M
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 A \otimes A \otimes M & \xrightarrow{\text{id}_A \otimes \text{act}} & A \otimes M \\
 m \otimes \text{id}_M \downarrow & & \downarrow \text{act} \\
 A \otimes M & \xrightarrow{\text{act}} & M.
 \end{array}$$

For future purposes, it will be convenient to introduce left modules in the more general setting of ∞ -categories that are ‘left-tensored’ over C .

Definition 19.1.3. Let C be a monoidal ∞ -category. A *left tensoring of an ∞ -category D over C* consists of an $\mathcal{LMod}$ -monoidal ∞ -category, given by a cocartesian fibration of ∞ -operads $p_D: D^\otimes \rightarrow \mathcal{LMod}^\otimes$, together with the following identifications:

- The underlying monoidal ∞ -category $\mathfrak{a}^*(p_D): \mathfrak{a}^*(D^\otimes) \rightarrow \mathcal{A}ssoc^\otimes$ is equivalent to C ;
- The underlying $\mathcal{Triv}$ -monoidal ∞ -category $\mathfrak{m}^*(p_D): \mathfrak{m}^*(D^\otimes) \rightarrow \mathcal{Triv}^\otimes$ corresponds to D under the equivalence $\text{Alg}_{\mathcal{Triv}}(\text{Cat}_\infty) \simeq \text{Cat}_\infty$.

We say that D is *left-tensored over C* if it is equipped with such a left tensoring.

Definition 19.1.4. Let D be left-tensored over C . Pullback along $\mathfrak{a}$ defines a forgetful functor $\mathfrak{a}^*: \text{Alg}_{\mathcal{LMod}}(D) \rightarrow \text{Alg}(C)$. Given an associative algebra $A \in \text{Alg}(C)$, we define the ∞ -category $\text{LMod}_A(D)$ of A -modules in D as the fiber of this functor over A .

Remark 19.1.5. By pulling back along the operad map $\mathcal{LMod} \rightarrow \mathcal{A}ssoc$, every monoidal ∞ -category C may be canonically regarded as left-tensored over itself. In this case, Definition 19.1.4 reduces to Definition 19.1.1.

Remark 19.1.6. All the definitions we just made also make sense for modules over *commutative* algebras, where we would replace the ∞ -operad $\mathcal{LMod}$ by its commutative version $\mathcal{Mod}$ from Example 12.2.8. It sits in a commutative diagram of ∞ -operads as follows:

$$\begin{array}{ccccc}
 \mathcal{A}ssoc & \longrightarrow & \mathcal{Comm} & & \\
 \downarrow & & \downarrow & & \\
 \mathcal{Triv} \hookrightarrow & \mathcal{LMod} & \longrightarrow & \mathcal{Mod} & \\
 \downarrow & & \downarrow & & \\
 \mathcal{A}ssoc & \longrightarrow & \mathcal{Comm}. & &
 \end{array}$$

In particular, restriction along the inclusion $Comm \hookrightarrow Mod$ defines a forgetful functor $Mod(C) := Alg_{Mod}(C) \rightarrow CAlg(C)$, allowing us to define an ∞ -category $Mod_A(C)$ for every commutative algebra A in C . It turns out this ∞ -category only depends on the underlying associative algebra of A :

Proposition 19.1.7. *Let C be a symmetric monoidal ∞ -category. Then the following commutative square is a pullback square:*

$$\begin{array}{ccc} Mod(C) & \xrightarrow{-|_{\mathcal{L}Mod}} & LMod(C) \\ \downarrow -|_{Comm} & & \downarrow -|_{\mathcal{A}ssoc} \\ CAlg(C) & \xrightarrow{-|_{\mathcal{A}ssoc}} & Alg(C). \end{array}$$

In particular, passing to vertical fibers over $R \in CAlg(C)$ induces an equivalence

$$Mod_R(C) \xrightarrow{\sim} LMod_{R'}(C),$$

where we denote the underlying associative algebra of R by R' .

Proof. This follows by combining Glasman [Gla14, Proposition 7] with [Lur17, Proposition 4.5.1.4]. More explicitly, the vertical functors are cartesian fibrations, the top comparison preserves cartesian morphisms because it does not change underlying objects, and the cited results identify it as an equivalence on every vertical fiber. Hence the square is a pullback. $\square$

19.1.1 Unit algebras and their modules

We will now show that the monoidal unit $\mathbb{1} \in C$ is always a commutative algebra, and modules over it are simply objects of C . In fact, we may do this more generally for *unital ∞ -operads*:

Definition 19.1.8. An ∞ -operad $\mathcal{O}$ is called *unital* if its total ∞ -category $\mathcal{O}^\otimes$ is pointed, i.e., if the terminal object (given by the empty tuple $\emptyset_{\mathcal{O}} := \{\}$) is also an initial object. Equivalently, $\mathcal{O}$ is unital if we have

$$\mathcal{O}(\{ \}; x) \simeq *$$

for every color $x \in \mathcal{O}^\simeq$.

Example 19.1.9. The ∞ -operads $Comm$, $\mathcal{A}ssoc$ and $\mathcal{E}_k$ are unital. The ∞ -operads $\mathcal{T}riv$, Mod and $\mathcal{L}Mod$ are not unital.

Definition 19.1.10. Let $\mathcal{O}$ be a unital ∞ -operad and let $C: \mathcal{O}^\otimes \rightarrow Cat_\infty$ be an $\mathcal{O}$ -monoidal ∞ -category. For every color $x \in \mathcal{O}^\simeq$, the unique multimorphism $e_x \in \mathcal{O}(\{ \}; x)$ defines a functor

$$\mathbb{1}_x: * \simeq C(\{ \}) \rightarrow C(\{x\}) = C_x.$$

We refer to the corresponding object $\mathbb{1}_x \in C_x$ as the *monoidal unit of C_x* .

Note that every $\mathcal{O}$ -algebra A comes equipped with ‘unit maps’ $e_x: \mathbb{1}_x \rightarrow A_x$ in C_x for every color $x \in \mathcal{O}^\simeq$: this map is the unique lift of the map $\{ \} \rightarrow \{A_x\}$ along the cocartesian morphism $\{ \} \rightarrow \{\mathbb{1}_x\}$. The following result by Lurie shows that, conversely, the monoidal units $\mathbb{1}_x \in C_x$ can always be assembled into an $\mathcal{O}$ -algebra $\mathbb{1}_C$ for which the maps $\mathbb{1}_x \rightarrow (\mathbb{1}_C)_x$ are isomorphisms:

Proposition 19.1.11 ([Lur17, Proposition 3.2.1.8]). *Let $\mathcal{O}$ be a unital ∞ -operad and let C be an $\mathcal{O}$ -monoidal ∞ -category. Then the ∞ -category $\mathrm{Alg}_{\mathcal{O}/\mathcal{O}}(C)$ of $\mathcal{O}$ -algebras in C admits an initial object $\mathbb{1}_C \in \mathrm{Alg}_{\mathcal{O}/\mathcal{O}}(C)$. Furthermore, an $\mathcal{O}$ -algebra is initial if and only if the unit maps $e_x: \mathbb{1}_x \rightarrow A_x$ are isomorphisms for all $x \in \mathcal{O}^\simeq$.*

The proof uses the theory of operadic left Kan extensions, which we do not develop here. We nevertheless record the short construction of $\mathbb{1}_C$ as an $\mathcal{O}$ -algebra of C from [HW21, Lemma II.45a]. It suffices to define a cocartesian functor

$$\begin{array}{ccc} \mathcal{O}^\otimes & \xrightarrow{\mathbb{1}_C} & C^\otimes \\ & \searrow & \downarrow p_C \\ & & \mathcal{O}^\otimes \end{array}$$

of cocartesian fibrations over $\mathcal{O}^\otimes$. By straightening-unstraightening, we may equivalently define a natural transformation of functors $\mathrm{const}_* \rightarrow \mathrm{Str}^{\mathrm{cc}}(p_C)$ of functors $\mathcal{O}^\otimes \rightarrow \mathrm{Cat}_\infty$. Since $\mathcal{O}^\otimes$ admits an initial object $\emptyset_{\mathcal{O}}$, there is a natural transformation $\mathrm{const}_{\emptyset_{\mathcal{O}}} \rightarrow \mathrm{id}_{\mathcal{O}^\otimes}$ of functors $\mathcal{O}^\otimes \rightarrow \mathcal{O}^\otimes$. As the fiber of p_C over $\emptyset_{\mathcal{O}}$ is the terminal ∞ -category, composing this transformation with $\mathrm{Str}^{\mathrm{cc}}(p_C)$ gives the desired map $\mathrm{const}_* \rightarrow \mathrm{Str}^{\mathrm{cc}}(p_C)$. It is clear from the construction that the maps $\mathbb{1}_x \rightarrow (\mathbb{1}_C)_x$ of the resulting $\mathcal{O}$ -algebra are isomorphisms.

Corollary 19.1.12 ([Lur17, Corollary 3.2.1.9]). *Let C be a symmetric monoidal ∞ -category. Then a commutative algebra R is initial in $\mathrm{CAlg}(C)$ if and only if the unit map $e: \mathbb{1} \rightarrow R$ is an isomorphism in C . In particular, the underlying object of the commutative algebra $\mathbb{1}_C \in \mathrm{CAlg}(C)$ from Proposition 19.1.11 is the monoidal unit of C . $\square$*

19.1.2 Free modules

Let A be an associative ring and let M_0 be an abelian group. The tensor product $M := A \otimes M_0$ admits the structure of a left A -module, given by

$$A \otimes M = A \otimes (A \otimes M_0) \simeq (A \otimes A) \otimes M_0 \xrightarrow{m \otimes M_0} A \otimes M_0 = M.$$

Moreover, M comes equipped with a morphism $\varphi: M_0 \simeq \mathbb{1} \otimes M_0 \xrightarrow{e \otimes M_0} A \otimes M_0 = M$ that exhibits M as the *free* A -module on M_0 : for every other left A -module N , composition with φ induces a bijection $\mathrm{Hom}_A(M, N) \xrightarrow{\sim} \mathrm{Hom}_{\mathrm{Ab}}(M_0, N)$, where $\mathrm{Hom}_A(M, N)$ is the set of A -module homomorphisms from M to N .

This admits the following ∞ -categorical refinement:

Proposition 19.1.13 ([Lur17, Proposition 4.2.4.2, Corollary 4.2.4.4]). *Let C be a monoidal ∞ -category and let D be an ∞ -category left tensored over C .*

(1) *The forgetful functor $\mathrm{LMod}(D) \xrightarrow{(\mathfrak{a}^*, \mathfrak{m}^*)} \mathrm{Alg}(C) \times D$ admits a left adjoint*

$$F: \mathrm{Alg}(C) \times D \rightarrow \mathrm{LMod}(D), \quad (A, M_0) \mapsto F_A(M_0)$$

satisfying $\mathfrak{a}^ F_A(M_0) \simeq A$ and $\mathfrak{m}^* F_A(M_0) \simeq A \otimes M_0$.*

- (2) For an associative algebra $A \in \text{Alg}(C)$, the forgetful functor $\mathfrak{m}^*: \text{LMod}_A(D) \rightarrow D$ admits a left adjoint

$$F_A: D \rightarrow \text{LMod}_A(D),$$

where the underlying object of $F_A(M_0)$ is $A \otimes M_0$.

As a consequence, we may deduce that modules over a trivial algebra are simply objects of D :

Corollary 19.1.14 ([Lur17, Proposition 4.2.4.9]). *Let C be a monoidal ∞ -category and let $A \in \text{Alg}(C)$ be an algebra object such that the unit map $e: \mathbb{1} \rightarrow A$ is an isomorphism in C . Then for every left C -tensored ∞ -category D the forgetful functor $\text{LMod}_A(D) \rightarrow D$ is an equivalence of ∞ -categories.*

Proof. By Proposition 19.1.13, the forgetful functor admits a left adjoint $F_A: D \rightarrow \text{LMod}_A(D)$. Since the forgetful functor is conservative, it suffices to show that the unit map $M_0 \rightarrow A \otimes M_0$ is an isomorphism. Since this map is given by tensoring the map $e: \mathbb{1} \rightarrow A$ by M_0 , this follows from the assumption on A . Indeed, the triangle identity then identifies the underlying counit with the inverse of the unit at the underlying object, and conservativity implies that the counit is also an isomorphism. $\square$

Remark 19.1.15. There is also a conceptual proof of Corollary 19.1.14. If $e: \mathbb{1} \rightarrow A$ is an isomorphism, then the monad $A \otimes -$ is isomorphic to the identity monad. Its category of modules is therefore equivalent to D . We have kept the direct proof above because it is shorter and does not require the monadicity package.

19.1.3 Limits and colimits of modules

Restriction of scalars can be organized into a cartesian fibration:

Proposition 19.1.16 ([Lur17, Corollary 4.2.3.2]). *Let D be left-tensored over a monoidal ∞ -category C . The restriction functor*

$$\alpha^*: \text{LMod}(D) \longrightarrow \text{Alg}(C)$$

is a cartesian fibration. A morphism in $\text{LMod}(D)$ is α^ -cartesian if and only if its underlying morphism in D is an isomorphism. Consequently, a morphism $f: A \rightarrow B$ of associative algebras induces a restriction-of-scalars functor*

$$f_*: \text{LMod}_B(D) \longrightarrow \text{LMod}_A(D)$$

which does not change the underlying object of D .

We can now deduce the limit statement for modules from the corresponding result for operadic algebras.

Corollary 19.1.17 ([Lur17, Corollaries 4.2.3.3 and 4.2.3.5]). *Let C be a monoidal ∞ -category, let D be left-tensored over C , and let $A \in \text{Alg}(C)$. Let $\mathfrak{m}^*: \text{LMod}_A(D) \rightarrow D$ be the forgetful functor, and let I be a small ∞ -category.*

- (1) If D admits I -indexed limits, then $\mathfrak{m}^*$ creates and preserves I -indexed limits.
- (2) If D admits I -indexed colimits and the functor $X \otimes -: D \rightarrow D$ preserves I -indexed colimits for every $X \in C$, then $\mathfrak{m}^*$ creates and preserves I -indexed colimits.

Proof. We prove (1). Equip $\text{Fun}(I, D)$ with the pointwise left tensoring over C . The identity functor of C and the constant-diagram functor $\text{const}: D \rightarrow \text{Fun}(I, D)$ assemble to an $\mathcal{L}\text{Mod}$ -monoidal functor between the corresponding left tensorings. Since const has right adjoint $\lim_I$, Proposition 14.4.7 equips the pair

$$(\text{id}_C, \lim_I)$$

with the right-adjoint lax $\mathcal{L}\text{Mod}$ -monoidal structure. Passing to $\mathcal{L}\text{Mod}$ -algebras and taking the fiber over A therefore gives an adjunction

$$\text{const}: \text{LMod}_A(D) \rightleftarrows \text{LMod}_A(\text{Fun}(I, D)) \simeq \text{Fun}(I, \text{LMod}_A(D)) : \lim_I.$$

The underlying object of the right adjoint is the limit in D . Thus $\mathfrak{m}^*$ preserves these limits. Since it detects isomorphisms, comparison with the constructed limit also shows that it creates them.

Part (2) is [Lur17, Corollary 4.2.3.5]. Its proof uses the simplicial model for modules to lift an operadic colimit of the underlying diagram; we briefly recall this model in Section 19.3. $\square$

19.2 Relative tensor products and change of algebras

If R is a classical commutative ring, then the category $\text{Mod}_R(\text{Ab})$ of R -modules comes equipped with a symmetric monoidal structure given by the *relative tensor product* $M \otimes_R N$ of two R -modules M and N . Moreover, if $f: R \rightarrow S$ is a morphism of commutative rings, then the relative tensor product $S \otimes_R M$ admits the structure of an S -module for every R -module M , resulting in a symmetric monoidal functor $f^*: \text{Mod}_R(\text{Ab}) \rightarrow \text{Mod}_S(\text{Ab})$ which is left adjoint to the forgetful functor $\text{Mod}_S(\text{Ab}) \rightarrow \text{Mod}_R(\text{Ab})$.

The goal of this section is to introduce this structure in an arbitrary symmetric monoidal ∞ -category that admits operadic geometric realizations. The organizing idea is that associative algebras are objects, bimodules are morphisms, and relative tensor products compose these morphisms. Associativity and unitality of the relative tensor product then express the coherence of this composition, while extension of scalars is composition with the regular bimodule associated to a morphism of algebras. We will make this perspective precise without constructing the ambient $(\infty, 2)$ -category explicitly.

19.2.1 Bimodules

Let A and B be ordinary associative algebras. Then we may speak of (A, B) -*bimodules*: objects M that are simultaneously a left A -module and a right B -module in a compatible way. This notion extends to the ∞ -categorical setting.

Definition 19.2.1. We define a colored operad $\mathcal{B}\text{Mod}$. It has three colors: $\mathfrak{a}_-$, $\mathfrak{a}_+$ and $\mathfrak{m}$. The multimorphism sets $\mathcal{B}\text{Mod}(\{X_i\}_{i \in I}; Y)$ are then given as follows:

- If $Y = \mathfrak{a}_-$, then this set is empty unless $X_i = \mathfrak{a}_-$ for all $i \in I$, in which case it is the set of linear orders on I ;
- If $Y = \mathfrak{a}_+$, then this set is empty unless $X_i = \mathfrak{a}_+$ for all $i \in I$, in which case it is the set of linear orders on I ;
- If $Y = \mathfrak{m}$, then this set is the set of linear orders $\{i_1 < i_2 < \cdots < i_n\}$ on I with the following property: there exists precisely one index $i_k \in I$ such that $X_{i_k} = \mathfrak{m}$, $X_{i_j} = \mathfrak{a}_-$ for $j < k$ and $X_{i_j} = \mathfrak{a}_+$ for $j > k$.

Composition in $\mathcal{B}Mod$ is given by composition of linear orders.

By Proposition 17.4.9, this colored operad canonically determines the ∞ -operad $\mathcal{B}Mod$ used below.

Observation 19.2.2. We identify $\mathcal{L}Mod$ with the full suboperad of $\mathcal{B}Mod$ spanned by the colors $\mathfrak{a}_-$ and $\mathfrak{m}$, and let $\mathcal{R}Mod \subseteq \mathcal{B}Mod$ denote the full suboperad spanned by $\mathfrak{m}$ and $\mathfrak{a}_+$. Thus $\mathcal{B}Mod$ contains two copies of $\mathcal{A}ssoc$, on $\mathfrak{a}_-$ and $\mathfrak{a}_+$, a copy of $\mathcal{T}riv$ on $\mathfrak{m}$, and the full suboperads $\mathcal{L}Mod$ and $\mathcal{R}Mod$. There is also an operad map

$$\mathcal{B}Mod \longrightarrow \mathcal{A}ssoc$$

which sends all three colors to $\mathfrak{a}$ and forgets the restrictions on the linear orders. Its restrictions to $\mathcal{L}Mod$ and $\mathcal{R}Mod$ are the usual collapse maps to $\mathcal{A}ssoc$.

Notation 19.2.3. For a monoidal ∞ -category C we write

$$\mathcal{R}Mod(C) := \mathcal{A}lg_{\mathcal{R}Mod}(C) \quad \text{and} \quad \mathcal{B}Mod(C) := \mathcal{A}lg_{\mathcal{B}Mod}(C).$$

Similarly, if A and B are associative algebras in C , we write

$$\mathcal{R}Mod_B(C) \quad \text{and} \quad {}_A\mathcal{B}Mod_B(C)$$

for the fibers over B and (A, B) of the two forgetful functors $\mathfrak{a}_+^*: \mathcal{R}Mod(C) \rightarrow \mathcal{A}lg(C)$ and $(\mathfrak{a}_-^*, \mathfrak{a}_+^*): \mathcal{B}Mod(C) \rightarrow \mathcal{A}lg(C) \times \mathcal{A}lg(C)$, respectively.

Observation 19.2.4. There is an operad map $\mathcal{B}Mod \rightarrow \mathcal{M}od$ which sends $\mathfrak{m}$ to $\mathfrak{m}$ and sends both $\mathfrak{a}_-$ and $\mathfrak{a}_+$ to $\mathfrak{a}$. Composing it with the inclusion $\mathcal{L}Mod \hookrightarrow \mathcal{B}Mod$ gives the canonical map $\mathcal{L}Mod \rightarrow \mathcal{M}od$, and similarly for $\mathcal{R}Mod$. It follows that for a commutative algebra A in a symmetric monoidal ∞ -category C , restriction along these operad maps defines forgetful functors

$$\begin{array}{ccccc} & & \mathcal{M}od_A(C) & & \\ & \swarrow \sim & \downarrow & \searrow \sim & \\ \mathcal{L}Mod_A(C) & \longleftarrow & {}_A\mathcal{B}Mod_A(C) & \longrightarrow & \mathcal{R}Mod_A(C). \end{array}$$

Here the left diagonal equivalence is the one from Proposition 19.1.7. The right one follows by applying Observation 19.2.5, since a commutative algebra is canonically isomorphic to its opposite algebra.

Observation 19.2.5. There are equivalences of ∞ -operads

$$\mathrm{rev}: \mathcal{A}ssoc \xrightarrow{\sim} \mathcal{A}ssoc, \quad \mathrm{rev}: \mathcal{L}Mod \xrightarrow{\sim} \mathcal{R}Mod \quad \text{and} \quad \mathrm{rev}: \mathcal{B}Mod \xrightarrow{\sim} \mathcal{B}Mod$$

given on objects by $\mathfrak{a} \mapsto \mathfrak{a}$, $\mathfrak{a}_- \mapsto \mathfrak{a}_+$, $\mathfrak{a}_+ \mapsto \mathfrak{a}_-$ and $\mathfrak{m} \mapsto \mathfrak{m}$, and on multimorphisms by reversing the order on the finite set. The map $\mathcal{A}ssoc \rightarrow \mathcal{C}omm$ coequalizes rev and the identity, so the symmetric monoidal structure on C canonically identifies the two induced monoidal structures. Consequently, restriction along rev defines an endofunctor $\mathrm{rev}^*: \mathrm{Alg}(C) \rightarrow \mathrm{Alg}(C)$. It sends an associative algebra A to its *opposite algebra*, denoted A^{rev} , which has the same underlying object but whose multiplication is given by

$$A \otimes A \xrightarrow{\sim} A \otimes A \xrightarrow{m_A} A,$$

where the first map is the swap map. For associative algebras A and B we thus get equivalences

$$\mathrm{LMod}_A(C) \xrightarrow{\sim} \mathrm{RMod}_{A^{\mathrm{rev}}}(C) \quad \text{and} \quad {}_A\mathrm{BMod}_B(C) \xrightarrow{\sim} {}_{B^{\mathrm{rev}}}\mathrm{BMod}_{A^{\mathrm{rev}}}(C).$$

Informally speaking, an (A, B) -bimodule M comes equipped with structure maps

$$A^{\otimes n} \otimes M \otimes B^{\otimes m} \rightarrow M$$

for all $n, m \geq 0$ that are compatible with the multiplication maps of A and B . By setting $m = 0$ or $n = 0$ we obtain a left A -module structure and a right B -module structure on M , respectively. These structures are compatible with each other in a very strong sense: the A -module structure on M can be upgraded to an A -module structure internal to the ∞ -category $\mathrm{RMod}_B(C)$ and vice versa:

Theorem 19.2.6 ([Lur17, Theorem 4.3.2.7]). *Let A and B be associative algebras in a monoidal ∞ -category C .*

- (1) *The ∞ -category $\mathrm{RMod}_B(C)$ is left tensored over C , and there is an equivalence*

$${}_A\mathrm{BMod}_B(C) \simeq \mathrm{LMod}_A(\mathrm{RMod}_B(C));$$

- (2) *The ∞ -category $\mathrm{LMod}_A(C)$ is right tensored over C , and there is an equivalence*

$${}_A\mathrm{BMod}_B(C) \simeq \mathrm{RMod}_B(\mathrm{LMod}_A(C)).$$

Corollary 19.2.7. *For associative algebras A and B in C , there are equivalences*

$${}_A\mathrm{BMod}_{\mathbb{1}}(C) \xrightarrow{\sim} \mathrm{LMod}_A(C) \quad \text{and} \quad {}_{\mathbb{1}}\mathrm{BMod}_B(C) \xrightarrow{\sim} \mathrm{RMod}_B(C).$$

Proof. In light of the equivalences $\mathrm{RMod}_{\mathbb{1}}(C) \xrightarrow{\sim} C$ and $\mathrm{LMod}_{\mathbb{1}}(C) \xrightarrow{\sim} C$ from Corollary 19.1.14, this follows directly from the previous theorem. $\square$

19.2.2 The relative tensor product

Let B be a discrete associative ring. A central construction in the theory of modules is the *relative tensor product*: Given a right B -module M and a left B -module N , their relative tensor product $M \otimes_B N$ is obtained as the quotient of $M \otimes N$ by identifying $mb \otimes n$ with $m \otimes bn$ for all $m \in M$, $n \in N$ and $b \in B$. In other words, $M \otimes_B N$ is the coequalizer of the following diagram:

$$M \otimes B \otimes N \rightrightarrows M \otimes N.$$

In the ∞ -categorical setting, this coequalizer needs to be replaced by the geometric realization of a simplicial diagram, called the *two-sided bar construction*. The construction is somewhat subtle, and is handled by Lurie in [Lur17, Construction 4.4.2.7]. We will first give the non-coherent description, and then indicate its coherent version in Remark 19.2.10 below.

Construction 19.2.8 (Bar construction). Let B be an associative algebra in a monoidal ∞ -category C , let M be a right B -module, and let N be a left B -module. The *two-sided bar construction* for M and N over B , denoted $\text{Bar}_B(M, N)_\bullet$, is a simplicial object in C described as follows:

- In degree $[n] \in \Delta^{\text{op}}$, it is given by $\text{Bar}_B(M, N)_n := M \otimes B^{\otimes n} \otimes N$;
- The face maps $d_i : \text{Bar}_B(M, N)_n \rightarrow \text{Bar}_B(M, N)_{n-1}$ are given by:
 - $d_0 = \text{act}_M \otimes \text{id}_{B^{\otimes(n-1)}} \otimes \text{id}_N : M \otimes B \otimes B^{\otimes(n-1)} \otimes N \rightarrow M \otimes B^{\otimes(n-1)} \otimes N$.
 - For $0 < i < n$, d_i is induced by the multiplication map $m : B \otimes B \rightarrow B$ on the i -th and $(i+1)$ -th factors of $B^{\otimes n}$.
 - $d_n = \text{id}_M \otimes \text{id}_{B^{\otimes(n-1)}} \otimes \text{act}_N : M \otimes B^{\otimes(n-1)} \otimes B \otimes N \rightarrow M \otimes B^{\otimes(n-1)} \otimes N$.
- The degeneracy maps $s_i : \text{Bar}_B(M, N)_n \rightarrow \text{Bar}_B(M, N)_{n+1}$ are induced by inserting the unit $e : \mathbb{1} \rightarrow B$ at the $(i+1)$ -th position.

Definition 19.2.9 (Relative tensor product). Let C be a monoidal ∞ -category. Assume that C admits geometric realizations and that the tensor product $- \otimes - : C \times C \rightarrow C$ preserves geometric realizations in both variables. For an associative algebra B , a right B -module M and a left B -module N , their *relative tensor product* is the object of C given by the geometric realization of the bar construction:

$$M \otimes_B N := |\text{Bar}_B(M, N)_\bullet| = \text{colim}_{[n] \in \Delta^{\text{op}}} \text{Bar}_B(M, N)_n.$$

More generally, let $A, B, D \in \text{Alg}(C)$. If $M \in {}_A\text{BMod}_B(C)$ and $N \in {}_B\text{BMod}_D(C)$, then the bar construction lifts to a simplicial object of ${}_A\text{BMod}_D(C)$. Geometric realizations of bimodules are created on underlying objects by [Lur17, Proposition 4.3.3.9], so its realization defines an (A, D) -bimodule

$$M \otimes_B N \in {}_A\text{BMod}_D(C).$$

Remark 19.2.10. To formally construct $\text{Bar}_B(M, N)_\bullet$, Lurie defines in [Lur17, Definition 4.4.1.1] an ∞ -operad $\mathcal{Tens}_2$ and shows in [Lur17, Proposition 4.4.1.11] that it sits in a pushout square of ∞ -operads as follows:

$$\begin{array}{ccc} \mathcal{A}ssoc & \xrightarrow{a_+} & \mathcal{B}Mod \\ a_- \downarrow & \lrcorner & \downarrow \\ \mathcal{B}Mod & \hookrightarrow & \mathcal{Tens}_2. \end{array}$$

In particular, for a monoidal ∞ -category C , the operad $\mathcal{Tens}_2$ encodes the data of algebras $A, B, D \in \text{Alg}(C)$, an (A, B) -bimodule M , and a (B, D) -bimodule N : there is a pullback square

$$\begin{array}{ccc} \text{Alg}_{\mathcal{Tens}_2}(C) & \longrightarrow & \text{BMod}(C) \\ \downarrow & \lrcorner & \downarrow \alpha_-^* \\ \text{BMod}(C) & \xrightarrow{\alpha_+^*} & \text{Alg}(C). \end{array}$$

For this reason, we denote the five colors of $\mathcal{Tens}_2$ by $\mathfrak{a}$, $\mathfrak{m}$, $\mathfrak{b}$, $\mathfrak{n}$ and $\mathfrak{c}$.

Lurie's construction supplies a functor

$$\text{BCut}: \Delta^{\text{op}} \longrightarrow \mathcal{Tens}_2^{\otimes}$$

which sends $[n]$ to the tuple $(\mathfrak{m}, \mathfrak{b}, \mathfrak{b}, \dots, \mathfrak{b}, \mathfrak{n})$ encoding $M \otimes B^{\otimes n} \otimes N$. It can be described as the pullback, in $\mathcal{Tens}_2^{\otimes}$, of the following cospan:

$$\begin{array}{ccc} & (\mathfrak{m}, \mathfrak{b}, \mathfrak{b}, \dots, \mathfrak{b}) & \\ & \downarrow & \\ (\mathfrak{b}, \mathfrak{b}, \dots, \mathfrak{b}, \mathfrak{n}) & \longrightarrow & (\mathfrak{b}, \mathfrak{b}, \dots, \mathfrak{b}). \end{array}$$

Both maps are inert, and their pullback is the indicated tuple. The compatibility of these pullbacks with the simplicial operators is part of [Lur17, Construction 4.4.2.7]; it may also be expressed using the cut functors discussed later in Section 19.3.

Now, given an (A, B) -bimodule M and a (B, D) -bimodule N , we may think of this data as defining a $\mathcal{Tens}_2$ -algebra in C , which in particular defines a functor $\mathcal{Tens}_2^{\otimes} \rightarrow C^{\otimes}$. Precomposing this with BCut gives a functor $\Delta^{\text{op}} \rightarrow C^{\otimes}$. One can check that this actually factors through the active morphisms. Using cocartesian transport, we can then turn this into the desired functor $\text{Bar}_B(M, N)_{\bullet}: \Delta^{\text{op}} \rightarrow C$.

Proposition 19.2.11. *Let C and D be monoidal ∞ -categories which admit geometric realizations and whose tensor products preserve geometric realizations separately in both variables. If $F: C \rightarrow D$ is a monoidal functor which preserves geometric realizations, then for every associative algebra $B \in \text{Alg}(C)$, right B -module M , and left B -module N , there is a natural isomorphism*

$$F(M) \otimes_{F(B)} F(N) \xrightarrow{\sim} F(M \otimes_B N).$$

Proof. The monoidal structure on F , together with the functorial construction of the coherent bar diagram from Remark 19.2.10, gives an isomorphism of simplicial objects

$$\text{Bar}_{F(B)}(F(M), F(N))_{\bullet} \xrightarrow{\sim} F(\text{Bar}_B(M, N)_{\bullet}).$$

Taking geometric realizations and using that F preserves them gives the asserted isomorphism. $\square$

The relative tensor product is therefore a composition law for bimodules. More precisely, it assembles into a functor

$$\text{BMod}(C) \times_{\text{Alg}(C)} \text{BMod}(C) \longrightarrow \text{BMod}(C),$$

where the fiber product matches the right algebra of the first bimodule with the left algebra of the second. This composition is coherently associative and unital. In particular, for bimodules ${}_A M_B$, ${}_B N_C$ and ${}_C P_D$ there is a natural isomorphism

$$(M \otimes_B N) \otimes_C P \simeq M \otimes_B (N \otimes_C P) \quad \in \quad {}_A \mathbf{BMod}_D(C).$$

For a bimodule ${}_A M_B$, the regular bimodules A and B act as units:

$$A \otimes_A M \simeq M \quad \text{and} \quad M \otimes_B B \simeq M \quad \in \quad {}_A \mathbf{BMod}_B(C).$$

These statements, including their coherent compatibility, are proved in [Lur17, Propositions 4.4.3.14 and 4.4.3.16]. They are the higher-categorical reason that the usual change-of-algebras formalism works exactly as in ordinary algebra.

Proposition 19.2.12 (Change of algebras). *Let C be a monoidal ∞ -category which admits geometric realizations and whose tensor product preserves geometric realizations separately in both variables. For every morphism $f : A \rightarrow B$ in $\mathbf{Alg}(C)$, the restriction functor from Proposition 19.1.16*

$$f_* : \mathbf{LMod}_B(C) \longrightarrow \mathbf{LMod}_A(C)$$

admits a left adjoint given by extension of scalars:

$$f^* := B \otimes_A - : \mathbf{LMod}_A(C) \longrightarrow \mathbf{LMod}_B(C).$$

Analogously, restriction from right B -modules to right A -modules admits the left adjoint $- \otimes_A B$.

Proof. The morphism f makes every left B -module into a left A -module and makes B into a (B, A) -bimodule. For a left A -module M and a left B -module N , an A -linear morphism $g : M \rightarrow f_* N$ induces the B -linear composite

$$B \otimes_A M \xrightarrow{\text{id}_B \otimes_A g} B \otimes_A f_* N \longrightarrow N,$$

where the final map is induced by the B -action on N . Conversely, precomposition with

$$M \xrightarrow{\sim} A \otimes_A M \xrightarrow{f \otimes_A \text{id}_M} B \otimes_A M$$

turns every B -linear morphism $B \otimes_A M \rightarrow N$ into an A -linear morphism $M \rightarrow f_* N$. These assignments arise from the displayed natural transformations, so they define maps between the corresponding mapping anima. Unitality and associativity of relative tensor products show that these maps are inverse. The argument for right modules is analogous. $\square$

If all the algebras involved are a fixed algebra A , the relative tensor product takes the form of a functor

$$- \otimes_A - : {}_A \mathbf{BMod}_A(C) \times {}_A \mathbf{BMod}_A(C) \rightarrow {}_A \mathbf{BMod}_A(C)$$

which satisfies associativity and unitality. Lurie shows that this assembles into a monoidal structure on ${}_A \mathbf{BMod}_A(C)$:

Proposition 19.2.13 ([Lur17, Proposition 4.4.3.12]). *Let C be a monoidal ∞ -category. Assume that C admits geometric realizations and that the tensor product $- \otimes - : C \times C \rightarrow C$ preserves geometric realizations in both variables. For an associative algebra A , there exists a monoidal structure on ${}_A \mathbf{BMod}_A(C)$ given by the relative tensor product over A .*

19.2.3 Modules over commutative algebras

Let R be a discrete commutative ring. Given R -modules M and N , the relative tensor product $M \otimes_R N$ is again an R -module, with the scalar multiplication given by $r(m \otimes n) = rm \otimes n = m \otimes rn$. In fact, the construction $(M, N) \mapsto M \otimes_R N$ equips the category $\text{Mod}_R(\text{Ab})$ of R -modules with a symmetric monoidal structure. The analogous statement also holds in the ∞ -categorical setting:

Theorem 19.2.14 ([Lur17, Theorem 4.5.2.1]). *Let C be a symmetric monoidal ∞ -category. Assume that C admits geometric realizations and that the tensor product of C preserves geometric realizations in both variables. Let R be a commutative algebra in C . Then:*

- (1) *The ∞ -category $\text{Mod}_R(C)$ admits a symmetric monoidal structure.*
- (2) *The functor $\text{Mod}_R(C) \rightarrow {}_R\text{BMod}_R(C)$ from Observation 19.2.4 refines to a monoidal functor, where the target has the monoidal structure from Proposition 19.2.13.*

In particular, the tensor product on $\text{Mod}_R(C)$ is given as the following composite:

$$\begin{aligned} \text{Mod}_R(C) \times \text{Mod}_R(C) &\longrightarrow {}_R\text{BMod}_R(C) \times {}_R\text{BMod}_R(C) \\ &\xrightarrow{-\otimes_R-} {}_R\text{BMod}_R(C) \longrightarrow \text{LMod}_R(C) \xrightarrow{\sim} \text{Mod}_R(C). \end{aligned}$$

Proposition 19.2.15 ([Lur17, Theorem 5.1.4.10 and Corollary 3.4.1.7]). *Let C be a symmetric monoidal ∞ -category. Assume that C admits geometric realizations and that the tensor product of C preserves geometric realizations in both variables. Let R be a commutative algebra in C . The forgetful functor $\text{CAlg}(\text{Mod}_R(C)) \rightarrow \text{CAlg}(C)$ lifts to an equivalence of ∞ -categories*

$$\text{CAlg}(\text{Mod}_R(C)) \xrightarrow{\sim} \text{CAlg}(C)_{R/}.$$

Warning 19.2.16. There is no analogous equivalence $\text{Alg}(\text{Mod}_R(C)) \simeq \text{Alg}(C)_{R/}$. Already for ordinary rings, the structure map from R to an associative algebra in $\text{Mod}_R(C)$ is central, whereas an arbitrary morphism from R to an associative algebra need not have central image; see also [Lur17, Warning 7.1.3.9].

Corollary 19.2.17 (Localization of a commutative base algebra). *Let $R \in \text{CAlg}(C)$ be as in Theorem 19.2.14, and let*

$$L: \text{Mod}_R(C) \rightleftarrows D: i$$

be a symmetric monoidal Bousfield localization as in Proposition 14.5.3. Then the local R -module $L(R)$ canonically underlies a commutative algebra in C equipped with a morphism $R \rightarrow L(R)$. It is initial among commutative algebras under R whose underlying R -modules are local.

Proof. The object R is the monoidal unit of $\text{Mod}_R(C)$ and hence the initial object of $\text{CAlg}(\text{Mod}_R(C))$. By Corollary 14.5.4, its reflection $L(R)$ is the initial object of $\text{CAlg}(D)$, and the essential image of

$$\text{CAlg}(D) \hookrightarrow \text{CAlg}(\text{Mod}_R(C))$$

consists of the commutative R -algebras with local underlying module. The equivalence $\text{CAlg}(\text{Mod}_R(C)) \simeq \text{CAlg}(C)_{R/}$ from Proposition 19.2.15 gives the result. $\square$

When C is presentably symmetric monoidal, the symmetric monoidal ∞ -category $\mathrm{Mod}_R(C)$ is again presentably symmetric monoidal: its underlying category is presentable by Propositions 19.1.7 and 22.5.5, and its relative tensor product preserves colimits separately by [Lur17, Corollary 4.4.2.15]. It is therefore closed by Proposition 22.5.3; in particular, each functor $- \otimes_R N$ admits an internal right adjoint $\underline{\mathrm{Hom}}_R(N, -)$.

19.2.4 Restriction and extension of scalars

Let $f: A \rightarrow B$ be a morphism of classical commutative rings. Every B -module M may be regarded as an A -module via the scalar multiplication $am := f(a)m$, resulting in a functor

$$f_*: \mathrm{Mod}_B(\mathrm{Ab}) \rightarrow \mathrm{Mod}_A(\mathrm{Ab})$$

called *restriction of scalars*¹⁹. Furthermore, this functor admits a left adjoint

$$f^*: \mathrm{Mod}_A(\mathrm{Ab}) \rightarrow \mathrm{Mod}_B(\mathrm{Ab})$$

sending an A -module to the relative tensor product $B \otimes_A M$ of B and M over A , where we regard B as a (B, A) -bimodule. Furthermore, this left adjoint is symmetric monoidal: there are natural isomorphisms of B -modules

$$B \otimes_A (M \otimes_A N) \xrightarrow{\sim} (B \otimes_A M) \otimes_B (B \otimes_A N)$$

for all A -modules M and N .

It turns out that this structure exists for commutative algebras in an arbitrary symmetric monoidal ∞ -category C .

Theorem 19.2.18 ([Lur17, Theorem 4.5.3.1]). *Let C be a symmetric monoidal ∞ -category. Assume that C admits geometric realizations and that the tensor product $\otimes: C \times C \rightarrow C$ preserves geometric realizations in both variables separately. Then there exists a cocartesian fibration*

$$p: \mathrm{Mod}(C)^\otimes \rightarrow \mathrm{CAlg}(C) \times \mathrm{Span}(\mathrm{Fin})$$

such that for every commutative algebra R the pullback along $\{R\} \times \mathrm{Span}(\mathrm{Fin}) \hookrightarrow \mathrm{CAlg}(C) \times \mathrm{Span}(\mathrm{Fin})$ is the cocartesian fibration

$$p_R: \mathrm{Mod}_R(C)^\otimes \rightarrow \mathrm{Span}(\mathrm{Fin})$$

encoding the symmetric monoidal structure on $\mathrm{Mod}_R(C)$ from Theorem 19.2.14.

By straightening-unstraightening, the cocartesian fibration p corresponds to a functor

$$\mathrm{Str}^{\mathrm{cc}}: \mathrm{CAlg}(C) \times \mathrm{Span}(\mathrm{Fin}) \rightarrow \mathrm{Cat}_\infty$$

which preserves finite products in the second variable. By currying, this may equivalently be encoded as a functor

$$\mathrm{CAlg}(C) \rightarrow \mathrm{Fun}^\times(\mathrm{Span}(\mathrm{Fin}), \mathrm{Cat}_\infty) = \mathrm{Cat}_\infty^\otimes$$

¹⁹While this assignment is *contravariant* in the ring homomorphism, the covariant notation f_* comes from the algebro-geometric perspective: f_* is geometrically thought of as the pushforward on quasi-coherent sheaves along the map $f: \mathrm{Spec}(B) \rightarrow \mathrm{Spec}(A)$.

from commutative algebras in C to symmetric monoidal ∞ -categories. In particular, every morphism $f: R \rightarrow S$ in $\mathbf{CAlg}(C)$ induces a symmetric monoidal functor

$$f^*: \mathbf{Mod}_R(C) \rightarrow \mathbf{Mod}_S(C).$$

When $R = \mathbb{1}$ is the monoidal unit, the resulting symmetric monoidal functor

$$C \simeq \mathbf{Mod}_{\mathbb{1}}(C) \rightarrow \mathbf{Mod}_S(C)$$

is the free module functor $F_S: C \rightarrow \mathbf{Mod}_S(C)$ from Proposition 19.1.13.

19.3 Simplicial models for algebras and modules

We briefly record alternative simplicial models for associative algebras and their modules. We will use their cut diagrams in the construction of endomorphism categories below. Lurie's proof of the colimit statement in Corollary 19.1.17 also uses these models to lift an operadic colimit of the underlying diagram to a colimit of modules.

19.3.1 Associative algebras as simplicial objects

For a nonempty finite linear order P , let $\text{Cut}(P)$ be its set of Dedekind cuts, namely the nontrivial partitions (P_0, P_1) such that $p_0 \leq p_1$ for all $p_0 \in P_0$ and $p_1 \in P_1$. Pullback of cuts along a map of finite linear orders defines a span of finite sets. The canonical ordering of the cuts refines this construction to a functor

$$\text{Cut}: \Delta^{\text{op}} \longrightarrow \mathcal{A}ssoc^{\otimes}.$$

Definition 19.3.1. Let C be a monoidal ∞ -category. An $\mathbb{A}_{\infty}$ -algebra in C is a functor

$$A: \Delta^{\text{op}} \longrightarrow C^{\otimes}$$

over $\text{Cut}: \Delta^{\text{op}} \rightarrow \mathcal{A}ssoc^{\otimes}$ which sends the dual of every interval inclusion

$$e_i: [1] \simeq \{i-1 \leq i\} \hookrightarrow [n]$$

to an inert morphism. We write

$$\mathbf{Alg}_{\mathbb{A}_{\infty}}(C) \subseteq \mathbf{Fun}_{/\mathcal{A}ssoc^{\otimes}}(\Delta^{\text{op}}, C^{\otimes})$$

for the full subcategory of such functors.

The inert condition identifies the components of $A_n \in C_n^{\otimes} \simeq C^n$ with A_1 . The resulting simplicial structure records the multiplication and all its coherences.

Theorem 19.3.2 ([Lur17, Proposition 4.1.3.19]). *Restriction along Cut induces an equivalence*

$$\text{Cut}^*: \mathbf{Alg}(C) \xrightarrow{\sim} \mathbf{Alg}_{\mathbb{A}_{\infty}}(C).$$

Applied to the cartesian monoidal ∞ -category $\mathbf{Cat}_{\infty}$, this identifies $\mathbb{A}_{\infty}$ -monoidal ∞ -categories with monoidal ∞ -categories.

19.3.2 Left modules as simplicial objects

Construction 19.3.3. Define

$$\text{LCut}: \Delta^{\text{op}} \times [1] \longrightarrow \mathcal{LMod}^{\otimes}$$

as the composite

$$\Delta^{\text{op}} \times [1] \xrightarrow{\text{Cut} \times [1]} \mathcal{A}ssoc^{\otimes} \times [1] \longrightarrow \mathcal{LMod}^{\otimes},$$

where the final functor sends $(I, 1)$ to $(I, \emptyset)$ and $(I, 0)$ to $(I, \{m\})$. The morphism from $(I, 0)$ to $(I, 1)$ is represented by the span

$$(I, \{m\}) \leftarrow (I, \emptyset) \xrightarrow{=} (I, \emptyset).$$

Definition 19.3.4. An $\mathbb{A}_{\infty}$ -module in C is a functor

$$F: \Delta^{\text{op}} \times [1] \longrightarrow C^{\otimes}$$

over LCut satisfying the following conditions:

- (1) The restriction to $\Delta^{\text{op}} \times \{1\}$ is an $\mathbb{A}_{\infty}$ -algebra.
- (2) Each morphism $F([n], 0) \rightarrow F([n], 1)$ is inert.
- (3) If $\alpha: [n] \rightarrow [m]$ is an interval inclusion with $\alpha(n) = m$, then $F(\alpha^{\text{op}}, \text{id}_0)$ is inert.

We write $\text{LMod}^{\mathbb{A}_{\infty}}(C)$ for the resulting full subcategory of $\text{Fun}_{/\mathcal{LMod}^{\otimes}}(\Delta^{\text{op}} \times [1], C^{\otimes})$.

Theorem 19.3.5 ([Lur17, Proposition 4.2.2.12]). *Restriction along LCut induces an equivalence*

$$\text{LCut}^*: \text{LMod}(C) \xrightarrow{\sim} \text{LMod}^{\mathbb{A}_{\infty}}(C).$$

For a module represented by F , write $A = F([1], 1)$ and $M = F([0], 0)$. The conditions identify

$$F([n], 0) \simeq (A, \dots, A, M),$$

so the simplicial structure records all the action maps $A^{\otimes n} \otimes M \rightarrow A^{\otimes m} \otimes M$. Applied to a diagram of A -modules, this model explains why a colimit of the underlying objects inherits an A -action when $A \otimes -$ preserves that colimit. In the proof cited in Corollary 19.1.17, this is made precise by taking the colimit in the functor category defining the simplicial model.

19.4 Endomorphism algebras

Let M be an abelian group, and let $\text{End}(M)$ denote the set of group homomorphisms from M to itself. We may equip $\text{End}(M)$ with the structure of an associative ring, where addition is given pointwise, and multiplication is given by composition of homomorphisms. Moreover, this ring admits the following universal property: for any other associative ring R , the data of a ring homomorphism $R \rightarrow \text{End}(M)$ is equivalent to the data of an R -module structure on M .

We will formulate the ∞ -categorical analogue in a slightly more general setting. Let C be a monoidal ∞ -category and let D be left-tensored over C . For an object $M \in D$, an object $E \in C$ equipped with a morphism $a: E \otimes M \rightarrow M$ is called an *endomorphism object of M* if composition with a induces an isomorphism

$$\mathrm{Hom}_C(X, E) \xrightarrow{\sim} \mathrm{Hom}_D(X \otimes M, M)$$

for every $X \in C$. We will explain why such an endomorphism object admits a preferred associative algebra structure and is universal among associative algebras acting on M . The necessary coherence is subtle, so we will only describe the idea of the construction and refer to Lurie [Lur17, Section 4.7.1] for the details.

Notation 19.4.1. In the ∞ -category $\mathcal{LMod}^\otimes$, we have the following two ‘canonical’ morphisms $(\{\mathfrak{a}\}, \{\mathfrak{m}\}) \rightarrow \{\mathfrak{m}\}$:

- The ‘projection map’ $\mathrm{pr}: (\{\mathfrak{a}\}, \{\mathfrak{m}\}) \rightarrow \{\mathfrak{m}\}$, given by the span

$$\{\mathfrak{a}\} \sqcup \{\mathfrak{m}\} \hookrightarrow \{\mathfrak{m}\} \xrightarrow{=} \{\mathfrak{m}\}.$$

- The ‘action map’ $\mathrm{act}: (\{\mathfrak{a}\}, \{\mathfrak{m}\}) \rightarrow \{\mathfrak{m}\}$, given by the span

$$\{\mathfrak{a}\} \sqcup \{\mathfrak{m}\} \xleftarrow{=} \{\mathfrak{a}\} \sqcup \{\mathfrak{m}\} \rightarrow \{\mathfrak{m}\}.$$

Construction 19.4.2 (cf. [Lur17, Definition 4.7.1.1]). Let $p_D: D^\otimes \rightarrow \mathcal{LMod}^\otimes$ exhibit D as left-tensored over C . An *enriched morphism* of D is a diagram

$$N \xleftarrow{\beta} X \xrightarrow{\alpha} M$$

in $D^\otimes$ satisfying the following conditions:

- (1) The morphism β is p_D -cocartesian and the image $p_D(\beta)$ in $\mathcal{LMod}^\otimes$ is the projection map $\mathrm{pr}: (\{\mathfrak{a}\}, \{\mathfrak{m}\}) \rightarrow \{\mathfrak{m}\}$;
- (2) The image $p_D(\alpha)$ in $\mathcal{LMod}^\otimes$ is the action map $\mathrm{act}: (\{\mathfrak{a}\}, \{\mathfrak{m}\}) \rightarrow \{\mathfrak{m}\}$.

We denote by

$$\mathrm{Ar}^{\mathrm{enr}}(D) \subseteq \mathrm{Fun}_{/\mathcal{LMod}^\otimes}(\ulcorner, D^\otimes)$$

the full subcategory spanned by the enriched morphisms of D , where $\ulcorner$ is the walking span of Definition 1.4.1. There are two forgetful functors

$$(s, t): \mathrm{Ar}^{\mathrm{enr}}(D) \rightarrow D \times D$$

given by $s(\alpha, \beta) := N$ and $t(\alpha, \beta) := M$.

Remark 19.4.3. Let us spell out the data contained in an enriched morphism. The fiber of $D^\otimes$ over $(\{\mathfrak{a}\}, \{\mathfrak{m}\})$ is equivalent to $C \times D$, hence we may write $X = (A, N')$ for some $A \in C$ and $N' \in D$. Condition (1) says that β induces an isomorphism $N' \xrightarrow{\sim} N$ in D . By condition (2), the morphism $\alpha: X \rightarrow M$ then corresponds to a morphism $A \otimes N \rightarrow M$ in D .

Definition 19.4.4. Let D be left-tensored over a monoidal ∞ -category C , and let $M \in D$ be an object. We define the *endomorphism category of M* as the fiber of (s, t) over (M, M) :

$$C[M] := \mathrm{Ar}^{\mathrm{enr}}(D) \times_{D \times D} \{(M, M)\}.$$

An object of $C[M]$ is equivalently a pair (A, η) consisting of an object $A \in C$ and a morphism $\eta: A \otimes M \rightarrow M$. For two such objects, the mapping anima is

$$\mathrm{Hom}_{C[M]}((X, \zeta), (A, \eta)) \simeq \mathrm{Hom}_C(X, A) \times_{\mathrm{Hom}_D(X \otimes M, M)} \{\zeta\},$$

where the map from $\mathrm{Hom}_C(X, A)$ is induced by η . Thus (A, η) is terminal precisely when composition with η induces an isomorphism

$$\mathrm{Hom}_C(X, A) \xrightarrow{\sim} \mathrm{Hom}_D(X \otimes M, M)$$

for every $X \in C$, which is exactly the endomorphism-object condition from the beginning of the section.

Construction 19.4.5 (Idea of the monoidal structure). We briefly indicate how Lurie equips $C[M]$ with a monoidal structure. For each $n \geq 0$, the twisted arrow poset $\mathrm{Tw}^r([n])$ from Definition 13.1.6 organizes the subintervals of $[n]$. Sending an interval $(i \leq j)$ to its length, and a subinterval inclusion $(k \leq l) \subseteq (i \leq j)$ to the dual of the translated interval inclusion $[l - k] \hookrightarrow [j - i]$, gives a diagram

$$\mathrm{Tw}^r([n]) \longrightarrow \Delta^{\mathrm{op}} \xrightarrow{(-, 0)} \Delta^{\mathrm{op}} \times [1] \xrightarrow{\mathrm{LCut}} \mathcal{LMod}^{\otimes}.$$

A suitable lift to $D^{\otimes}$ records a string of enriched morphisms

$$M_0 \longrightarrow M_1 \longrightarrow \cdots \longrightarrow M_n$$

together with all their coherent composites. Relative Kan extension makes these lifts functorial in $[n]$, and restriction to the consecutive subintervals satisfies the Segal condition. Taking the fiber over the constant string $(M, \dots, M)$ therefore produces an $\mathbb{A}_{\infty}$ -monoidal structure on $C[M]$. We refer to [Lur17, Definitions 4.7.1.5–4.7.1.6, Proposition 4.7.1.13] for the relative Kan extension and the verification of the Segal condition.

Under the comparison between $\mathbb{A}_{\infty}$ -monoidal categories and monoidal ∞ -categories obtained by applying Theorem 19.3.2 to Cat_{∞} , the resulting tensor product is given on objects by

$$(A, \eta) \otimes (A', \eta') = \left(A \otimes A', A \otimes A' \otimes M \xrightarrow{\mathrm{id}_A \otimes \eta'} A \otimes M \xrightarrow{\eta} M \right).$$

Theorem 19.4.6 ([Lur17, Proposition 4.7.1.30, Theorem 4.7.1.34]). *Let D be left-tensored over a monoidal ∞ -category C , and let $M \in D$. The endomorphism category $C[M]$ admits a monoidal structure for which the forgetful functor $C[M] \rightarrow C$ is monoidal. Moreover, there is an equivalence*

$$\mathrm{Alg}(C[M]) \xrightarrow{\sim} \mathrm{LMod}(D) \times_D \{M\}$$

compatible with the forgetful functors to $\mathrm{Alg}(C)$.

The equivalence in this theorem is obtained by combining the simplicial models for algebras and modules from Section 19.3 with evaluation of an enriched string on its longest interval. Lurie’s construction first replaces $C[M]$ by an equivalent auxiliary monoidal category defined using left Kan extensions; this is what ensures that evaluation on the longest interval retains all the coherent module data.

Corollary 19.4.7 ([Lur17, Corollary 4.7.1.40]). *Let D be left-tensored over a monoidal ∞ -category C , and let $M \in D$ admit an endomorphism object $\text{End}(M) \in C$. Then $\text{End}(M)$ admits a preferred structure of an associative algebra, and M admits a preferred structure of a left module over $\text{End}(M)$. Moreover, there is an equivalence of ∞ -categories making the following diagram commute:*

$$\begin{array}{ccc} \text{Alg}(C)_{/\text{End}(M)} & \xrightarrow{\simeq} & \text{LMod}(D) \times_D \{M\} \\ & \searrow \text{fgt} & \swarrow \text{fgt} \\ & \text{Alg}(C) & \end{array}$$

Proof. By definition, $\text{End}(M)$ is terminal in $C[M]$. The monoidal structure from Theorem 19.4.6 and Corollary 14.4.11 therefore give $\text{End}(M)$ a unique associative algebra structure in $C[M]$. Its image in C is an associative algebra, while its image under the equivalence of Theorem 19.4.6 is a left module structure on M .

The forgetful functor $\text{Alg}(C[M]) \rightarrow \text{Alg}(C)$ is a right fibration by [Lur17, Proposition 4.7.1.39]. Since its total category has the terminal object just constructed, this right fibration is represented by $\text{End}(M)$. This gives the displayed equivalence. $\square$

19.5 Stable and spectral consequences

We now specialize the preceding constructions to stable ∞ -categories and spectra. The stability of module categories is a direct consequence of the abstract limit and colimit results. Relative tensor products then identify colimit-preserving functors between categories of modules, while the endomorphism-algebra construction equips mapping spectra with their natural multiplicative structures. Together these results culminate in a symmetric monoidal form of the monogenic Morita theorem.

19.5.1 Module categories and bimodules

Corollary 19.5.1. *Let C be a monoidal ∞ -category whose underlying ∞ -category is stable and whose tensor product preserves finite colimits separately in both variables. For every associative algebra $A \in \text{Alg}(C)$, the ∞ -categories $\text{LMod}_A(C)$ and $\text{RMod}_A(C)$ are stable.*

Proof. By Corollary 19.1.17, the forgetful functor $\text{LMod}_A(C) \rightarrow C$ creates and preserves finite limits and colimits. Thus $\text{LMod}_A(C)$ has a zero object, and a commutative square in $\text{LMod}_A(C)$ is a pullback if and only if it is a pushout, since this may be checked in C . Hence $\text{LMod}_A(C)$ is stable. The proof for right modules is dual. $\square$

For an associative ring spectrum R , the relative tensor product gives a functor

$$- \otimes_R -: \text{RMod}_R \times_R \text{BMod}_R \longrightarrow \text{RMod}_R$$

which preserves colimits in both variables by [Lur17, Corollary 4.4.2.15]. Currying therefore gives a functor

$${}_R\mathbf{BMod}_R \longrightarrow \mathbf{Fun}^L(\mathbf{RMod}_R, \mathbf{RMod}_R).$$

Theorem 19.5.2 (Spectral Eilenberg–Watts theorem, [Lur17, Proposition 7.1.2.4]). *For every associative ring spectrum R , the functor*

$${}_R\mathbf{BMod}_R \longrightarrow \mathbf{Fun}^L(\mathbf{RMod}_R, \mathbf{RMod}_R)$$

is an equivalence.

Thus the spectral Eilenberg–Watts theorem is the morphism-level counterpart of Morita theory: bimodules define colimit-preserving functors between module categories by relative tensor product, and every such functor arises uniquely in this way. In particular, the relative tensor product describes not merely a distinguished class of functors, but all colimit-preserving endofunctors of $\mathbf{RMod}_R$.

For morphisms of commutative ring spectra, the symmetric monoidal extension-of-scalars functor fits into a triple of adjoints.

Proposition 19.5.3. *Let $f: R \rightarrow S$ be a morphism of commutative ring spectra.*

- (1) *The functor $f^*: \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$ admits a right adjoint*

$$f_*: \mathbf{Mod}_S \rightarrow \mathbf{Mod}_R.$$

For every S -module M , the underlying spectrum of $f_(M)$ is naturally isomorphic to the underlying spectrum of M .*

- (2) *The functor $f_*: \mathbf{Mod}_S \rightarrow \mathbf{Mod}_R$ admits a further right adjoint*

$$\underline{\mathrm{hom}}_R(S, -): \mathbf{Mod}_R \rightarrow \mathbf{Mod}_S$$

whose underlying R -module is the internal hom: it satisfies $f_ \underline{\mathrm{hom}}_R(S, -) \simeq \underline{\mathrm{Hom}}_R(S, -)$ as functors $\mathbf{Mod}_R \rightarrow \mathbf{Mod}_R$.*

Proof. For part (1), Theorem 22.2.5 reduces the existence of f_* to the fact that f^* preserves small colimits. This may be tested after postcomposing with the forgetful functor $\mathbf{Mod}_S \rightarrow \mathbf{Sp}$, where f^* is given by the relative tensor product $S \otimes_R -$.

To identify the underlying spectrum of $f_*(M)$, we may equivalently pass to left adjoints and show that for every $M_0 \in \mathbf{Sp}$ there is a natural isomorphism $f^*(F_R(M_0)) \simeq F_S(M_0)$. This follows from the functoriality of extension of scalars: F_R and F_S arise from the unit maps $\eta_R: \mathbb{S} \rightarrow R$ and $\eta_S: \mathbb{S} \rightarrow S$, and $\eta_S = f \circ \eta_R$.

For part (2), the adjoint functor theorem applies once more because f_* preserves colimits. This can be checked on underlying spectra, since restriction of scalars does not change them and the forgetful functors create colimits. The identification $f_* \underline{\mathrm{hom}}_R(S, -) \simeq \underline{\mathrm{Hom}}_R(S, -)$ may be checked after passing to left adjoints, where it is the identity

$$f_* f^* \simeq S \otimes_R -: \mathbf{Mod}_R \longrightarrow \mathbf{Mod}_R.$$

□

Corollary 19.5.4. *The restriction of the cocartesian fibration p from Theorem 19.2.18 to the fiber over $\langle 1 \rangle \in \text{Span}(\text{Fin})$,*

$$-|_{\text{Comm}} : \text{Mod}(\text{Sp}) \longrightarrow \text{CAlg}(\text{Sp}),$$

is also a cartesian fibration, hence straightens to a contravariant functor

$$\text{CAlg}(\text{Sp})^{\text{op}} \rightarrow \text{Cat}_{\infty}, \quad A \mapsto \text{Mod}_A, \quad f \mapsto f_*.$$

Proof. Recall from Lemma 23.1.20 that a cocartesian fibration $p : E \rightarrow C$ is also a cartesian fibration if and only if every induced functor between its fibers admits a right adjoint. For $-|_{\text{Comm}} : \text{Mod}(\text{Sp}) \rightarrow \text{CAlg}(\text{Sp})$, this is precisely Proposition 19.5.3(1). $\square$

19.5.2 Mapping spectra and Morita theory

Proposition 19.5.5 ([Lur17, Remark 7.1.2.2]). *Let D be a stable ∞ -category.*

- (1) *For every object $M \in D$, the mapping spectrum $\text{hom}_D(M, M)$ admits a preferred structure of an associative ring spectrum.*
- (2) *For every object $M \in D$, the mapping spectrum functor $\text{hom}_D(M, -) : D \rightarrow \text{Sp}$ admits a preferred lift*

$$\text{hom}_D(M, -) : D \rightarrow \text{RMod}_{\text{hom}_D(M, M)}.$$

Proof. We may assume that D is small by enlarging the ambient universe if necessary. Consider the fully faithful Yoneda embedding

$$D \hookrightarrow \widehat{D} := \text{Ind}(D).$$

The ∞ -category $\widehat{D}$ is presentable and stable, and the inclusion of D is exact; see Proposition 22.3.6 and [Lur17, Proposition 1.1.3.6]. In particular, it preserves mapping spectra. Being presentable and stable, $\widehat{D}$ is Sp -local, so Theorem 18.5.3 and Proposition 18.5.2 equip it with a unique Sp -module structure in Pr^{L} ; in particular $\widehat{D}$ is canonically left-tensored over Sp . The resulting action functor $- \otimes M : \text{Sp} \rightarrow \widehat{D}$ is colimit-preserving and satisfies $\mathbb{S} \otimes M \simeq M$, so by Theorem 4.4.14 it agrees with the tensoring of Proposition 4.4.9; its right adjoint is therefore $\text{hom}_{\widehat{D}}(M, -)$, and for spectra E there are natural isomorphisms

$$\text{Hom}_{\text{Sp}}(E, \text{hom}_D(M, M)) \xrightarrow{\sim} \text{Hom}_{\widehat{D}}(E \otimes M, M).$$

Consequently, $A := \text{hom}_D(M, M)$ is an endomorphism object of $M \in \widehat{D}$. Applying Corollary 19.4.7 gives A a preferred associative algebra structure and equips M with a preferred left A -module structure. This proves (1).

Using this module structure, we may form the two-sided bar construction in $\widehat{D}$, just as in Section 19.2; see also [Lur17, Construction 4.4.2.7]. Its geometric realization defines a colimit-preserving functor

$$F := - \otimes_A M : \text{RMod}_A \rightarrow \widehat{D}.$$

Since both categories are presentable, Theorem 22.2.5 gives a right adjoint

$$G : \widehat{D} \rightarrow \text{RMod}_A.$$

Let $U: \mathbf{RMod}_A \rightarrow \mathbf{Sp}$ denote the forgetful functor. For $K \in \mathbf{Sp}$ and $X \in \widehat{D}$, the free-right-module adjunction and the tensor–Hom adjunction give natural isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Sp}}(K, UG(X)) &\simeq \mathrm{Hom}_{\mathbf{RMod}_A}(K \otimes A, G(X)) \\ &\simeq \mathrm{Hom}_{\widehat{D}}((K \otimes A) \otimes_A M, X) \\ &\simeq \mathrm{Hom}_{\widehat{D}}(K \otimes M, X) \\ &\simeq \mathrm{Hom}_{\mathbf{Sp}}(K, \mathrm{hom}_{\widehat{D}}(M, X)). \end{aligned}$$

The Yoneda lemma therefore provides a natural isomorphism

$$UG(X) \simeq \mathrm{hom}_{\widehat{D}}(M, X).$$

Restricting G along $D \hookrightarrow \widehat{D}$ gives the lift in (2). $\square$

Theorem 19.5.6 (Monogenic Morita theorem, [Lur17, Theorem 7.1.2.1, Proposition 7.1.2.7]). *Let C be a presentably symmetric monoidal stable ∞ -category, and assume that its monoidal unit $\mathbb{1}$ is a compact generator. Then the endomorphism spectrum*

$$A := \mathrm{hom}_C(\mathbb{1}, \mathbb{1})$$

is canonically a commutative ring spectrum, and the enriched mapping-spectrum functor lifts to a symmetric monoidal equivalence

$$\Phi: C \xrightarrow{\simeq} \mathbf{Mod}_A(C), \quad X \mapsto \mathrm{hom}_C(\mathbb{1}, X).$$

Proof. By Corollary 18.5.6, the unit functor $- \otimes \mathbb{1}: \mathbf{Sp} \rightarrow C$ is the essentially unique colimit-preserving symmetric monoidal functor out of $\mathbf{Sp}$, and its right adjoint is the mapping-spectrum functor $\mathrm{hom}_C(\mathbb{1}, -)$ (Corollary 4.4.10). By Proposition 14.3.6 this right adjoint is canonically lax symmetric monoidal. In particular, it carries the commutative algebra $\mathbb{1} \in \mathbf{CAlg}(C)$ to the commutative algebra

$$A := \mathrm{hom}_C(\mathbb{1}, \mathbb{1}) \in \mathbf{CAlg}(\mathbf{Sp}).$$

Applying the functoriality of commutative module categories under lax symmetric monoidal functors gives a lax symmetric monoidal functor

$$\Phi: C \simeq \mathbf{Mod}_{\mathbb{1}}(C) \longrightarrow \mathbf{Mod}_A(C), \quad X \mapsto \mathrm{hom}_C(\mathbb{1}, X),$$

whose underlying A -module-valued functor is that of Proposition 19.5.5. Since $\mathbb{1}$ is compact, Φ preserves filtered colimits; being exact, it preserves finite colimits, and hence all colimits.

Using the right-module equivalence of Observation 19.2.4, we may identify $\mathbf{Mod}_A(C)$ with $\mathbf{RMod}_A(C)$. Under this identification, the two-sided bar construction used in the proof of Proposition 19.5.5 defines a colimit-preserving functor

$$\Psi: \mathbf{Mod}_A(C) \longrightarrow C, \quad M \mapsto M \otimes_A \mathbb{1},$$

where $\mathbb{1}$ carries the left A -module structure of Proposition 19.5.5. The same construction exhibits Ψ as a left adjoint to Φ . On the free module we have $\Psi(A) \simeq \mathbb{1}$, so the counit and unit of $\Psi \dashv \Phi$ restrict to isomorphisms on the generators: the counit at $\mathbb{1}$ is the canonical isomorphism $\Psi\Phi(\mathbb{1}) = A \otimes_A \mathbb{1} \simeq \mathbb{1}$, and the unit at A is the isomorphism $A \rightarrow \Phi\Psi(A) = \mathrm{hom}_C(\mathbb{1}, \mathbb{1}) = A$. Both

Φ and Ψ are exact and preserve colimits, so the full subcategories of C and of $\text{Mod}_A(C)$ on which the counit, respectively the unit, is an isomorphism are stable and closed under colimits. Since $\mathbb{1}$ is a compact generator of C and A is a compact generator of $\text{Mod}_A(C)$ (Example 22.3.10), these subcategories are everything. Hence Φ is an equivalence.

It remains to promote the lax symmetric monoidal structure on Φ to a strong one. Its unit comparison $A \rightarrow \Phi(\mathbb{1})$ is an isomorphism, and its multiplication comparison

$$\Phi(X) \otimes_A \Phi(Y) \longrightarrow \Phi(X \otimes Y)$$

is an isomorphism for $X = Y = \mathbb{1}$, where both sides are A . Fixing $X = \mathbb{1}$, the full subcategory of those Y for which the comparison is an isomorphism is stable, closed under colimits, and contains $\mathbb{1}$, hence is all of C . Running the same argument in the second variable, now for each fixed $Y \in C$, shows that the comparison is an isomorphism for all X and Y . Thus Φ is a symmetric monoidal equivalence. $\square$

19.6 Monads and monadicity

An adjunction $F \dashv G$ determines the endofunctor GF together with a unit and multiplication. In classical category theory these maps satisfy the monad identities strictly. For ∞ -categories, they must be accompanied by coherent higher homotopies, so it is preferable to package a monad as an algebra object rather than to list this structure by hand. The Barr–Beck theorem then recognizes when the target of the adjunction is recovered as the ∞ -category of modules over this monad.

Two ingredients enter. The coherent monad T associated to an adjunction $F \dashv G$, together with its Eilenberg–Moore category $\text{LMod}_T(C)$, is constructed in Lurie [Lur17, Section 4.7]; we import it. The recognition criterion is then internal to this book: it reduces to the comparison theorem Theorem 21.7.3, proved in Part III from split simplicial resolutions.

We first construct the monoidal structure under which monads are algebras, as an instance of the endomorphism-algebra construction of Section 19.4. Regard Cat_∞ as left-tensored over itself via the cartesian product, and fix the object C . By the currying axiom for functor categories (Axiom C.4), the functor category $\text{Fun}(C, C)$, equipped with the evaluation functor $\text{ev}: \text{Fun}(C, C) \times C \rightarrow C$, is an endomorphism object of C in $(\text{Cat}_\infty, \times)$: precomposition with ev induces an isomorphism

$$\text{Hom}_{\text{Cat}_\infty}(X, \text{Fun}(C, C)) \xrightarrow{\sim} \text{Hom}_{\text{Cat}_\infty}(X \times C, C)$$

for every $X \in \text{Cat}_\infty$. Applying Corollary 19.4.7 therefore equips $\text{Fun}(C, C)$ with a preferred associative algebra structure in $(\text{Cat}_\infty, \times)$. Under the simplicial comparisons of Theorems 19.3.2 and 19.3.5, this is a monoidal structure whose tensor product $S \otimes T \simeq S \circ T$ is given by composition, and the resulting left module structure exhibits C as left-tensored over $\text{Fun}(C, C)$ through evaluation. When C is not small we run this construction in a larger universe, exactly as in Proposition 19.5.5.

Definition 19.6.1 ([Lur17, Definition 4.7.0.1]). A *monad* on an ∞ -category C is an associative algebra object

$$T \in \text{Alg}(\text{Fun}(C, C))$$

for the composition monoidal structure. Its *Eilenberg–Moore category* is the ∞ -category

$$\text{LMod}_T(C)$$

of left T -module objects in C , formed using the evaluation left tensoring constructed above. It comes with a forgetful functor $U_T: \mathbf{LMod}_T(C) \rightarrow C$.

Informally, the algebra structure consists of a unit $\mathrm{id}_C \rightarrow T$ and a multiplication $T \circ T \rightarrow T$, together with all coherent associativity and unitality data. A left T -module is an object $c \in C$ with a coherent action $T(c) \rightarrow c$.

We record the part of Lurie's package needed below. Given an adjunction

$$F: C \rightleftarrows D : G,$$

the composite GF admits a preferred monad structure T , characterized as the endomorphism algebra of G . The resulting T -action on G determines a *comparison functor*

$$K_G: D \longrightarrow \mathbf{LMod}_T(C)$$

with $U_T K_G \simeq G$; here $\mathbf{Fun}(D, C)$ is left-tensored over $\mathbf{Fun}(C, C)$ by postcomposition, which is the action used to form the endomorphism algebra of G . See [Lur17, Definition 4.7.3.2, Proposition 4.7.3.3 and the discussion preceding Definition 4.7.3.4]. The forgetful functor U_T is conservative and admits a left adjoint F_T , whose underlying endofunctor is T . The mate

$$F_T \longrightarrow K_G F$$

is an isomorphism. Finally, every U_T -split simplicial object of $\mathbf{LMod}_T(C)$ admits a geometric realization which is preserved by U_T ; see [Lur17, Corollary 4.2.3.2, Corollary 4.2.4.8, Lemma 4.7.3.12].

Definition 19.6.2. A right adjoint $G: D \rightarrow C$ is *monadic* if its comparison functor $K_G: D \rightarrow \mathbf{LMod}_T(C)$ is an equivalence of ∞ -categories.

Theorem 19.6.3 (Barr–Beck monadicity, [Lur17, Theorem 4.7.3.5]). *Let $G: D \rightarrow C$ be a conservative functor admitting a left adjoint. Assume that D admits geometric realizations of G -split simplicial objects and that G preserves them. Then G is monadic.*

Proof. Let T be the monad associated to the adjunction and let $K_G: D \rightarrow \mathbf{LMod}_T(C)$ be the comparison functor. Apply Theorem 21.7.3 to G , the forgetful functor $U_T: \mathbf{LMod}_T(C) \rightarrow C$, and K_G . The identity $U_T K_G \simeq G$ has mate $F_T \rightarrow K_G F$, which is an isomorphism by the cited package. The same package supplies conservativity of U_T and the required geometric realizations on the Eilenberg–Moore side. All the hypotheses of the comparison theorem are therefore satisfied, so K_G is an equivalence. $\square$

Remark 19.6.4. Barr–Beck gives an alternative proof of the underlying equivalence in the monogenic Morita theorem, Theorem 19.5.6. In its notation, consider the adjunction

$$- \otimes \mathbb{1}: \mathbf{Sp} \rightleftarrows C : \mathrm{hom}_C(\mathbb{1}, -).$$

The right adjoint is conservative because $\mathbb{1}$ is a generator. It preserves all colimits: compactness of $\mathbb{1}$ gives preservation of filtered colimits, exactness gives preservation of finite colimits, and arbitrary coproducts are filtered colimits of finite coproducts. Barr–Beck therefore identifies C with the Eilenberg–Moore category of the associated monad on $\mathbf{Sp}$.

If $A = \text{hom}_C(\mathbb{1}, \mathbb{1})$, the canonical natural transformation

$$- \otimes A \longrightarrow \text{hom}_C(\mathbb{1}, - \otimes \mathbb{1})$$

is an isomorphism: both sides preserve colimits and it is an isomorphism on the sphere spectrum. This identification respects the monad structures by the construction of multiplication on the endomorphism spectrum A . Hence the Eilenberg–Moore category is $\text{Mod}_A(C)$, and its comparison functor is the functor Φ of Theorem 19.5.6. The proof given there additionally shows directly that this equivalence is symmetric monoidal.

The basic example is the forgetful functor from modules. The following result makes precise the idea that an A -module is an algebra for the operation of tensoring with A .

Proposition 19.6.5 (Monadicity of module categories). *Let C be a monoidal ∞ -category, let D be left-tensored over C , and let $A \in \text{Alg}(C)$. The forgetful functor*

$$U_A: \text{LMod}_A(D) \longrightarrow D$$

is monadic, and its associated monad is $A \otimes -: D \rightarrow D$, with unit and multiplication induced by those of A .

Proof. The left adjoint exists by Proposition 19.1.13, and its composite with U_A is $A \otimes -$. The forgetful functor is conservative by [Lur17, Corollary 4.2.3.2]. Moreover, every U_A -split simplicial object admits a geometric realization which is preserved by U_A ; this is [Lur17, Lemma 4.7.3.12]. The claim follows from Theorem 19.6.3. $\square$

Corollary 19.6.6 (Limits and colimits of Eilenberg–Moore categories). *Let T be a monad on an ∞ -category C , let $U_T: \text{LMod}_T(C) \rightarrow C$ be its forgetful functor, and let I be a small ∞ -category.*

- (1) *If C admits I -indexed limits, then U_T creates and preserves them.*
- (2) *If C admits I -indexed colimits and T preserves them, then U_T creates and preserves them.*

Proof. For (1), apply Corollary 19.1.17(1) to the evaluation left tensoring of C over $\text{Fun}(C, C)$.

Part (2) follows from [Lur17, Proposition 4.2.3.4], applied to the evaluation left tensoring of C over $\text{Fun}(C, C)$ and the algebra $T \in \text{Alg}(\text{Fun}(C, C))$. $\square$

Applied to the monad $A \otimes -$ of Proposition 19.6.5, part (2) shows that the forgetful functor $\text{LMod}_A(D) \rightarrow D$ creates any colimits which exist in D and are preserved by $A \otimes -$. For a fixed algebra this sharpens Corollary 19.1.17(2), whose more uniform hypothesis ensures the same conclusion simultaneously for every algebra in C .

Remark 19.6.7. The monadic description does not replace the cut constructions of Section 19.3. They provide the common simplicial language for the two-sided bar construction and for the endomorphism-category construction of Section 19.4.

19.7 Application: connective complex K-theory

We conclude by applying the coherent algebra developed in this part to connective complex K-theory. Direct sum and tensor product should make the moduli anima of complex vector bundles into a commutative semiring object. We first explain the general mechanism by which distributivity produces such semiring objects. We then construct the tensor-product structure on vector bundles explicitly in Lurie’s Fin_* -model, where the multilinear maps of bundles are naturally indexed by the fibers of pointed maps, and use the comparison of Proposition 17.4.8. Finally, we combine the resulting semiring structure with symmetric monoidal group completion to obtain the commutative ring spectrum structure on ku used in Chapter 9.

19.7.1 Semirings from distributivity

The tensor product $\otimes_{\text{coprod}}$ of Section 22.4 is characterized by functors which preserve finite coproducts separately in each variable. The following proposition relates this universal property directly to the Day convolution tensor product on commutative monoids.

Lemma 19.7.1. *Interpreted one universe higher, the ∞ -category $\text{Cat}_\infty^{\text{coprod}}$ is presentable and semi-additive. Its zero object is the terminal category, and the cartesian product $C \times D$ is also the coproduct of C and D . Moreover, the tensor product $\otimes_{\text{coprod}}$ makes $\text{Cat}_\infty^{\text{coprod}}$ presentably symmetric monoidal.*

Proof. Presentability is the finite-coproduct case of [Lur17, Lemma 4.8.4.2]. Finite products are computed in Cat_∞ and therefore given by cartesian products of ∞ -categories. The terminal category is also initial in $\text{Cat}_\infty^{\text{coprod}}$, since a finite-coproduct-preserving functor from it to C must select an initial object of C .

Let 0_C and 0_D denote initial objects. For every $E \in \text{Cat}_\infty^{\text{coprod}}$, restriction along the functors

$$C \longrightarrow C \times D, \quad c \longmapsto (c, 0_D), \quad \text{and} \quad D \longrightarrow C \times D, \quad d \longmapsto (0_C, d),$$

induces an equivalence

$$\text{Fun}^\Pi(C \times D, E) \xrightarrow{\sim} \text{Fun}^\Pi(C, E) \times \text{Fun}^\Pi(D, E).$$

Indeed, the coproduct functor $\amalg: E \times E \rightarrow E$, which is left adjoint to the diagonal and therefore preserves finite coproducts, supplies an inverse which sends a pair (F, G) to the composite

$$C \times D \xrightarrow{F \times G} E \times E \xrightarrow{\amalg} E.$$

The two composites are naturally isomorphic to the identity because every (c, d) is the coproduct of $(c, 0_D)$ and $(0_C, d)$ in $C \times D$. Thus $C \times D$ is also a coproduct in $\text{Cat}_\infty^{\text{coprod}}$, proving semiadditivity.

Finally, the defining universal property of $\otimes_{\text{coprod}}$ exhibits tensoring with a fixed object as a left adjoint. It therefore preserves colimits, which proves the final assertion. $\square$

Proposition 19.7.2 (Cocartesian semiring construction). *The functor*

$$\text{Cat}_\infty^{\text{coprod}} \longrightarrow \text{CMon}(\text{Cat}_\infty), \quad C \longmapsto (C, \amalg),$$

has a canonical lax symmetric monoidal refinement, where the source carries $\otimes_{\text{coprod}}$ and the target carries the symmetric monoidal structure of Proposition 16.4.1.

Proof. Write

$$U: \text{Cat}_\infty^{\text{coprod}} \longrightarrow \text{Cat}_\infty$$

for the forgetful functor. The multimorphism operad defining $\otimes_{\text{coprod}}$ is, by construction in Section 22.4, a suboperad of the cartesian multimorphism operad of Cat_∞ . This inclusion gives U a canonical lax symmetric monoidal structure.

By Lemma 19.7.1, the multimorphism operad of $\text{Cat}_\infty^{\text{coprod}}$ is semiadditive. Moreover, U preserves finite products, since they are computed by cartesian products of ∞ -categories on both sides. The adjunction of Theorem 18.4.12 therefore gives a unique lift

$$\begin{array}{ccc} & & \text{CMon}(\mathcal{M}_{(\text{Cat}_\infty, \times)}) \\ & \nearrow \text{dashed} & \downarrow \\ \mathcal{M}_{\text{Cat}_\infty^{\text{coprod}}} & \longrightarrow & \mathcal{M}_{(\text{Cat}_\infty, \times)}. \end{array}$$

By the definition of CMon and the Day convolution structure of Proposition 16.4.1, the upper-right operad is naturally equivalent to the multimorphism operad of $\text{CMon}(\text{Cat}_\infty)$:

$$\text{CMon}(\mathcal{M}_{(\text{Cat}_\infty, \times)}) \simeq \mathcal{M}_{\text{CMon}(\text{Cat}_\infty)}.$$

The lift consequently determines a lax symmetric monoidal functor from $\text{Cat}_\infty^{\text{coprod}}$ to $\text{CMon}(\text{Cat}_\infty)$. On an object C , it gives the commutative monoid structure induced by the biproducts in $\text{Cat}_\infty^{\text{coprod}}$. Its unit selects the initial object 0_C , while its addition is the codiagonal

$$C \times C \simeq C \oplus C \longrightarrow C.$$

Under the description of the coproduct in Lemma 19.7.1, this codiagonal is the coproduct functor $\amalg: C \times C \rightarrow C$. Thus the underlying functor sends C to $(C, \amalg)$, as claimed. $\square$

There is consequently a general mechanism that turns distributivity over finite coproducts into a commutative semiring object.

Proposition 19.7.3. *Let C be a small symmetric monoidal ∞ -category which admits finite coproducts. Assume that for every object $X \in C$, the functor*

$$X \otimes -: C \longrightarrow C$$

preserves finite coproducts. Then the cocartesian monoidal structure on C canonically refines C to an object of

$$\text{CRig}(\text{Cat}_\infty) = \text{CAlg}(\text{CMon}(\text{Cat}_\infty)).$$

Proof. The finite-coproduct specialization in Section 22.4 exhibits the given symmetric monoidal structure on C as a commutative algebra object of $(\text{Cat}_\infty^{\text{coprod}}, \otimes_{\text{coprod}})$. Applying the lax symmetric monoidal functor of Proposition 19.7.2 produces a commutative algebra in $\text{CMon}(\text{Cat}_\infty)$, whose underlying commutative monoid is the cocartesian monoidal structure $(C, \amalg)$. $\square$

Corollary 19.7.4. *Under the hypotheses of Proposition 19.7.3, the maximal subgroupoid $C^\simeq$ is canonically an object of $\text{CRig}(\text{An})$.*

Proof. Apply the lax symmetric monoidal maximal-subgroupoid functor of Proposition 16.4.6 to the commutative algebra supplied by Proposition 19.7.3. $\square$

19.7.2 The vector-bundle semiring

Let X be a topological space. We write $\text{Vect}^{\text{disc}}(X)$ for the 1-category whose objects are the finite-rank complex vector bundles over X and whose morphisms are the morphisms of vector bundles over X . The superscript disc indicates that we are not yet recording the topology on spaces of bundle maps. Since vector bundles are stable under base change along continuous maps, this gives a contravariant functor

$$\text{Vect}^{\text{disc}} : \text{Top}^{\text{op}} \rightarrow \text{Cat}.$$

We show that $\text{Vect}^{\text{disc}}(X)$ can be equipped with a symmetric monoidal structure via tensor product of complex vector bundles. We construct this structure explicitly using Lurie's equivalent model of symmetric monoidal categories over Fin_* .

Construction 19.7.5. For a finite pointed set S_+ , we write S for the complement of the basepoint. Define a 1-category

$$\text{Vect}^{\otimes, \text{disc}}(X)$$

equipped with a functor $p_X : \text{Vect}^{\otimes, \text{disc}}(X) \rightarrow \text{Fin}_*$ as follows. An object over S_+ is an S -indexed family $(V_s)_{s \in S}$ of vector bundles over X . A morphism over a pointed map $\alpha : S_+ \rightarrow T_+$ from $(V_s)_{s \in S}$ to $(W_t)_{t \in T}$ consists of, for every $t \in T$, a map over X

$$\varphi_t : \prod_{s \in \alpha^{-1}(t)} V_s \longrightarrow W_t$$

which is continuous and complex multilinear on each fiber. Here the product is the fiber product over X , and if $\alpha^{-1}(t) = \emptyset$ we interpret the source as X , so that φ_t is the same as a section of W_t .

Composition is substitution of multilinear maps. Thus, if $\beta : T_+ \rightarrow U_+$ is another pointed map and $\psi_u : \prod_{t \in \beta^{-1}(u)} W_t \rightarrow Z_u$ is the corresponding family of multilinear maps, the composite over $\beta\alpha$ is given by

$$(v_s)_{s \in (\beta\alpha)^{-1}(u)} \mapsto \psi_u \left(\left(\varphi_t((v_s)_{s \in \alpha^{-1}(t)}) \right)_{t \in \beta^{-1}(u)} \right).$$

This is again multilinear in all variables. The identity morphism over id_{S_+} is given by the identity maps $V_s \rightarrow V_s$.

Proposition 19.7.6. *The functor*

$$p_X : \text{Vect}^{\otimes, \text{disc}}(X) \rightarrow \text{Fin}_*$$

is a cocartesian fibration of 1-categories. It therefore defines a symmetric monoidal 1-category whose underlying category is $\text{Vect}^{\text{disc}}(X)$ and whose tensor product is the tensor product of vector bundles.

Proof. Let $\alpha : S_+ \rightarrow T_+$ be a pointed map and let $(V_s)_{s \in S}$ be an object over S_+ . For each $t \in T$, choose the tensor product

$$W_t := \bigotimes_{s \in \alpha^{-1}(t)} V_s,$$

with the convention that the tensor product over the empty set is the trivial line bundle $\underline{\mathbb{C}}$. Let

$$\mu_t : \prod_{s \in \alpha^{-1}(t)} V_s \rightarrow W_t$$

be the universal multilinear map. Local trivializations show that W_t is again a vector bundle and that μ_t is continuous. These maps define a morphism

$$(V_s)_{s \in S} \longrightarrow (W_t)_{t \in T}$$

over α .

We claim that this morphism is p_X -cocartesian. Let $\beta: T_+ \rightarrow U_+$ be another pointed map, and let $(Z_u)_{u \in U}$ be an object over U_+ . By definition, a morphism from $(W_t)_{t \in T}$ to $(Z_u)_{u \in U}$ over β is a family of multilinear maps

$$\prod_{t \in \beta^{-1}(u)} W_t \rightarrow Z_u.$$

Precomposition with the maps μ_t identifies this set with the set of multilinear maps

$$\prod_{s \in (\beta\alpha)^{-1}(u)} V_s \rightarrow Z_u,$$

for all $u \in U$, by the universal property of the tensor products W_t . This is precisely the required universal property of a cocartesian morphism.

The fiber over S_+ is visibly $\mathbf{Vect}^{\text{disc}}(X)^S$, so the Segal maps are isomorphisms. Thus p_X is a symmetric monoidal category in Lurie's model. Taking nerves gives the corresponding cocartesian fibration of ∞ -categories, and the equivalence of operad models from Proposition 17.4.8 carries it to the desired symmetric monoidal ∞ -category over $\mathbf{Span}(\mathbf{Fin})$. Its tensor product is obtained by pushing forward along the active map $\{1, 2\}_+ \rightarrow \{1\}_+$, hence is the usual tensor product of vector bundles. $\square$

The symmetric monoidal structure above interacts with the direct-sum operation in the expected way. The general semiring formalism of Section 16.4 packages all the resulting coherence.

Corollary 19.7.7. *For every topological space X , the groupoid*

$$\mathbf{Vect}^{\text{disc}}(X) \simeq$$

of finite-rank complex vector bundles and bundle isomorphisms is naturally an object of $\mathbf{CRig}(\mathbf{An})$. This structure is contravariantly functorial in X by pullback.

Proof. Direct sum gives finite coproducts in the small 1-category $\mathbf{Vect}^{\text{disc}}(X)$, with initial object the rank-zero bundle $\underline{0}$. Tensor product distributes over direct sums in each variable, so Corollary 19.7.4 applies to the symmetric monoidal structure of Proposition 19.7.6. This construction is natural in symmetric monoidal functors which preserve finite coproducts. Since pullback preserves direct sums, tensor products, and trivial bundles, it gives the asserted functoriality in X . $\square$

The 1-category $\mathbf{Vect}^{\text{disc}}(X)$ discards the natural compact-open topologies on its hom sets. Let us first explain how the familiar topological enrichment motivates the construction below. Suppose that X is compact Hausdorff. The spaces $\mathbf{Iso}_X(E, E')$ of bundle isomorphisms, with their compact-open topologies, form the hom spaces of a topologically enriched groupoid. Its homotopy coherent nerve, in the sense of Construction 24.3.2, remembers the underlying anima of each of these spaces of isomorphisms by Proposition 24.3.4.

The simplicial resolution used to construct this nerve has the same vector bundles as objects in every degree, while its morphisms in degree n are the continuous maps

$$\mathrm{Hom}_{\mathrm{Top}}(|\Delta^n|, \mathrm{Iso}_X(E, E')).$$

The total spaces of vector bundles over a compact Hausdorff space are locally compact Hausdorff, so the compact-open exponential law identifies this set with

$$\mathrm{Iso}_{X \times |\Delta^n|}(\mathrm{pr}_X^* E, \mathrm{pr}_X^* E').$$

Thus the groupoid in simplicial degree n is equivalent to the full subgroupoid of $\mathrm{Vect}^{\mathrm{disc}}(X \times |\Delta^n|)^{\simeq}$ spanned by the bundles pulled back from X . Every vector bundle over $X \times |\Delta^n|$ is isomorphic to one of this form by Theorem 9.1.4. Consequently the inclusion of this full subgroupoid is an equivalence. For compact Hausdorff X , the colimit below is therefore the standard simplicial model for the homotopy coherent nerve of the topologically enriched groupoid of vector bundles and bundle isomorphisms. We use the same formula as our definition for an arbitrary topological space.

Definition 19.7.8. The cosimplicial topological space $[n] \mapsto |\Delta^n|$ and pullback of vector bundles determine a simplicial object

$$[n] \longmapsto \mathrm{Vect}^{\mathrm{disc}}(X \times |\Delta^n|)^{\simeq}$$

in $\mathrm{CRig}(\mathrm{An})$. We define the *moduli anima of complex vector bundles over X* by

$$\mathrm{Vect}(X)^{\simeq} := \mathrm{colim}_{[n] \in \Delta^{\mathrm{op}}} \mathrm{Vect}^{\mathrm{disc}}(X \times |\Delta^n|)^{\simeq} \in \mathrm{CRig}(\mathrm{An}).$$

Thus points of $\mathrm{Vect}(X)^{\simeq}$ are represented by vector bundles over X , paths are represented by concordances over $X \times [0, 1]$, and the higher simplices encode higher concordances. For compact Hausdorff X , this models how the topology on the spaces of bundle isomorphisms enters the homotopy coherent nerve. The simplicial description also makes direct sums and tensor products available degreewise.

Proposition 19.7.9. *The construction of Definition 19.7.8 defines a functor*

$$\mathrm{Vect}(-)^{\simeq} : \mathrm{Top}^{\mathrm{op}} \longrightarrow \mathrm{CRig}(\mathrm{An}).$$

Its underlying anima is computed by the same colimit in An . If X is paracompact Hausdorff, the inclusion of the zeroth simplicial degree induces a natural bijection

$$\pi_0 \mathrm{Vect}^{\mathrm{disc}}(X)^{\simeq} \xrightarrow{\cong} \pi_0 \mathrm{Vect}(X)^{\simeq}.$$

Proof. Functoriality follows from pullback. The ∞ -category Δ^{op} is sifted by Exercise 21.6.9, and the forgetful functors

$$\mathrm{CRig}(\mathrm{An}) \longrightarrow \mathrm{CMon}(\mathrm{An}) \longrightarrow \mathrm{An}$$

create sifted colimits. For the second functor, this follows by regarding $\mathrm{CMon}(\mathrm{An})$ as the full subcategory of $\mathrm{Fun}(\mathrm{Span}(\mathrm{Fin}), \mathrm{An})$ spanned by the finite-product-preserving functors: pointwise sifted colimits preserve this condition by Lemma 21.6.11. For the first functor, use $\mathrm{CRig}(\mathrm{An}) = \mathrm{CAlg}(\mathrm{CMon}(\mathrm{An}))$, the fact that $\mathrm{CMon}(\mathrm{An})$ is presentably symmetric monoidal by Theorem 18.5.3, and [Lur17, Corollary 3.2.3.2]. This proves the assertion about the underlying anima.

It remains to identify its path components. Applying π_0 to the geometric realization gives the coequalizer of the two maps from simplicial degree 1 to degree 0. Every vector bundle over $X \times |\Delta^n|$

is isomorphic to the pullback of its restriction to a vertex: the product $X \times |\Delta^n|$ is again paracompact Hausdorff, and the identity is homotopic to the composite of the projection with the inclusion of a vertex, so this follows from Theorem 9.1.4. In particular, degree 0 surjects onto the coequalizer. The two endpoint restrictions of a vector bundle over $X \times [0, 1]$ are isomorphic by the same theorem, so the two maps in the coequalizer agree. The coequalizer is therefore $\pi_0 \text{Vect}^{\text{disc}}(X)^{\simeq}$. $\square$

Corollary 19.7.10. *The construction of Definition 19.7.8, applied to finite-rank real vector bundles, defines a functor*

$$\text{Vect}_{\mathbb{R}}(-)^{\simeq} : \text{Top}^{\text{op}} \longrightarrow \text{CRig}(\text{An})$$

whose underlying anima is computed by the corresponding colimit in An.

Proof. The constructions and proofs of Propositions 19.7.6 and 19.7.9 and Corollary 19.7.7 apply verbatim with $\mathbb{C}$ replaced by $\mathbb{R}$. $\square$

We now specialize to $X = \text{pt}$. Recall from Section 9.4.2 that the group completion of $\text{Vect}(\text{pt})^{\simeq}$ is the infinite loop anima underlying connective complex K-theory. The semiring structure above supplies its multiplication and all its coherences.

Proposition 19.7.11. *The connective complex K-theory spectrum ku admits a canonical structure of commutative ring spectrum.*

Proof. By Definition 19.7.8, we have

$$\text{Vect}(\text{pt})^{\simeq} \in \text{CAlg}(\text{CMon}(\text{An})).$$

By Proposition 16.6.4, the composite

$$\text{CMon}(\text{An}) \xrightarrow{(-)^{\text{grp}}} \text{CGrp}(\text{An}) \xrightarrow{\mathbf{B}^{\infty}} \text{Sp}$$

is symmetric monoidal. Applying it to the commutative algebra object $\text{Vect}(\text{pt})^{\simeq}$ gives a commutative algebra object of Sp , i.e. a commutative ring spectrum. Under Theorem 5.4.5, the functor $\mathbf{B}^{\infty} : \text{CGrp}(\text{An}) \rightarrow \text{Sp}_{\geq 0}$ is the inverse of the recognition equivalence, and $(-)^{\text{grp}}$ is the same group-completion left adjoint used in Section 9.4.2. Its underlying spectrum is therefore precisely ku . $\square$

Exercises

20 Monoidal Dwyer–Kan localizations and derived categories

In Section 14.5 we treated localizations that are *reflective*: the local objects form a full subcategory of C , and the monoidal structure is transported along the reflector. Derived categories are not presented in this way. The ∞ -category $D(R)$ arises from the 1-category $\text{Ch}(R)$ of chain complexes by inverting the quasi-isomorphisms, which is a Dwyer–Kan localization, and the reflective criteria do not apply to it. Supplying the missing monoidal statement is the purpose of this chapter.

The difficulty is that a symmetric monoidal structure on C is not extra structure on C alone, but a functor $C^{\otimes} \rightarrow \text{Span}(\text{Fin})$, so inverting a class W of morphisms of C is not by itself enough to say what should happen to the tensor product. The remedy is to invert morphisms upstairs as well: we localize $C^{\otimes}$ at a class $W^{\otimes}$ built from W , and give conditions under which the result is again a symmetric monoidal ∞ -category with the expected universal property.

Section 20.1 carries this out. The case where tensoring with any object preserves W is straightforward (Proposition 20.1.1); the substantial statement is Theorem 20.1.6, which only requires the monoidal structure to be *left derivable*, and which follows [NS18, Theorem A.7]. In Section 20.2 we apply it to unbounded chain complexes, obtaining both the derived tensor product on $D(R)$ and a lax symmetric monoidal refinement of the localization $\text{Ch}(R) \rightarrow D(R)$. We show that this tensor product preserves colimits, and record the compact presentation of $D(R)$ needed for the comparison with Eilenberg–MacLane modules in Corollary 8.3.4.

20.1 Monoidal Dwyer–Kan localizations

Let C be a symmetric monoidal ∞ -category and let W be a class of morphisms in C . The question addressed in this section is when the localization $C[W^{-1}]$ of C at W inherits a symmetric monoidal structure from C .

The situation is relatively straightforward if tensoring with any object preserves W :

Proposition 20.1.1 (Tensor-stable monoidal localization). *Let C be a symmetric monoidal ∞ -category, and let W be a class of morphisms in C such that, for every object $X \in C$, the functor $X \otimes -$ sends morphisms in W to morphisms in W .*

- (1) *The localization $C[W^{-1}]$ inherits a unique symmetric monoidal structure turning the localization functor $\gamma: C \rightarrow C[W^{-1}]$ into a symmetric monoidal functor.*

(2) For any other symmetric monoidal ∞ -category D , restriction along γ induces a fully faithful functor

$$\mathrm{Fun}^{\otimes}(C[W^{-1}], D) \hookrightarrow \mathrm{Fun}^{\otimes}(C, D)$$

whose essential image consists of those symmetric monoidal functors $C \rightarrow D$ which invert the morphisms in W .

Proof. We may first replace W by the smallest class of morphisms in C that contains W and is closed under isomorphisms, composition, and homotopy. This does not change the localization, since a functor out of C inverts the original class W if and only if it inverts this larger class. The enlarged class is still preserved by tensoring with an arbitrary object, since tensor products preserve isomorphisms, composition, and homotopies. Thus we may assume that W is a wide subcategory of C .

For (1), consider the full subcategory $\mathrm{RelCat}_{\infty} \subseteq \mathrm{Ar}(\mathrm{Cat}_{\infty})$ spanned by the *relative ∞ -categories*: those functors $W \hookrightarrow C$ which are simultaneously monomorphisms in Cat_{∞} and essentially surjective. (Such a choice of W corresponds to a *wide subcategory* of C .) We will generally denote objects of RelCat_{∞} as pairs (C, W) .

Observe that there is a fully faithful inclusion functor

$$i_{\mathrm{rel}}: \mathrm{Cat}_{\infty} \hookrightarrow \mathrm{RelCat}_{\infty}, \quad D \mapsto (D, D^{\simeq}).$$

Observe that for a relative ∞ -category (C, W) , the localization functor $\gamma: C \rightarrow C[W^{-1}]$ refines to a morphism of relative ∞ -categories $(C, W) \rightarrow i_{\mathrm{rel}}(C[W^{-1}])$. We claim that this map exhibits $C[W^{-1}]$ as a left adjoint object to (C, W) under i_{rel} . For every other ∞ -category D , precomposition with γ by definition induces a fully faithful inclusion of anima

$$\mathrm{Hom}_{\mathrm{Cat}_{\infty}}(C[W^{-1}], D) \rightarrow \mathrm{Hom}_{\mathrm{Cat}_{\infty}}(C, D)$$

whose image consists of the functors $C \rightarrow D$ that send W to isomorphisms. This image is precisely the image of $\mathrm{Hom}_{\mathrm{RelCat}_{\infty}}((C, W), i_{\mathrm{rel}}(D))$, proving the adjointness.

By the pointwise criterion for left adjoints (Lemma 21.1.4), the assignment $(C, W) \mapsto C[W^{-1}]$ assembles into a functor $\mathcal{L}: \mathrm{RelCat}_{\infty} \rightarrow \mathrm{Cat}_{\infty}$. Moreover, by Exercise 1.5.25, this functor preserves finite products, hence induces a functor on commutative monoids:

$$\mathcal{L}: \mathrm{CMon}(\mathrm{RelCat}_{\infty}) \rightarrow \mathrm{CMon}(\mathrm{Cat}_{\infty}).$$

Since also the right adjoint i_{rel} preserves finite products, the entire adjunction lifts to commutative monoids. Unwinding definitions, the condition on the pair (C, W) in the statement of the proposition precisely guarantees that the symmetric monoidal structure on C refines to an object $(C, W) \in \mathrm{CMon}(\mathrm{RelCat}_{\infty})$. The unit of the adjunction now takes the form of a symmetric monoidal functor $\gamma: C \rightarrow C[W^{-1}]$ refining the localization functor.

The adjunction on commutative monoids gives the claimed universal property on mapping anima. Applying the same argument with the pointwise symmetric monoidal categories $\mathrm{Fun}([n], D)$ for every $n \geq 0$, as in Definition 14.2.3, gives the asserted fully faithful functor between symmetric monoidal functor ∞ -categories. The uniqueness statement follows immediately. $\square$

Remark 20.1.2. In the situation of Proposition 14.5.3, we may consider the class W of L -equivalences in C . The assumption on L guarantees that also the condition of Proposition 20.1.1 is satisfied. In particular, Proposition 20.1.1 provides an alternative way of equipping the subcategory $D \subseteq C$ with a symmetric monoidal structure.

The criterion of Proposition 20.1.1 is often too strict in practice. For example, for chain complexes with quasi-isomorphisms, tensoring with a fixed complex preserves quasi-isomorphisms only under flatness hypotheses. We already encountered this phenomenon in Section 6.6, where we introduced the notion of a *derived functor*, and explained how to form the derived tensor product of chain complexes. The goal of the remainder of this section is to enhance the derived tensor product to a fully coherent symmetric monoidal structure. Our discussion closely follows that of [NS18, Appendix A].

Definition 20.1.3 (Left derivable monoidal structure). Let (C, W, I) be an ∞ -category with weak equivalences and cofibrations, and assume that C is given a symmetric monoidal structure. We call an object of C *tensor-cofibrant* if it belongs to the smallest full subcategory of C that contains all cofibrant objects and the monoidal unit and is closed under binary tensor products. We say that the monoidal structure is *left derivable* if for every finite set J , every J -indexed collection of cofibrant objects $\{X_j\}_{j \in J}$, every J -indexed collection of tensor-cofibrant objects $\{Y_j\}_{j \in J}$, and every collection of weak equivalences $f_j: X_j \rightarrow Y_j$, the induced map

$$\bigotimes_{j \in J} f_j: \bigotimes_{j \in J} X_j \rightarrow \bigotimes_{j \in J} Y_j$$

is again a weak equivalence.

Remark 20.1.4. In the model category literature, there exists a notion of *symmetric monoidal model category*. The underlying category with weak equivalences and cofibrations of such a symmetric monoidal model category is always left derivable. However, the condition of being left derivable is much weaker: we require only that tensoring preserve the relevant weak equivalences, with no general compatibility between tensor products and cofibrations.

A convenient sufficient criterion is the following. Assume that the monoidal unit is cofibrant, that the tensor product of two cofibrant objects is again cofibrant, and that for every cofibrant object Z the functor $- \otimes Z$ sends weak equivalences between cofibrant objects to weak equivalences. Then every tensor-cofibrant object is again cofibrant, and an induction on the cardinality of J shows that the monoidal structure is left derivable in the sense of the definition.

Construction 20.1.5. Let (C, W, I) be an ∞ -category with weak equivalences and cofibrations which comes equipped with a left derivable symmetric monoidal structure. For a finite set J , let W^J denote the collection of morphisms in $C^J \simeq C_J^\otimes$ given by J -tuples of weak equivalences, and let $W^\otimes$ be the collection of morphisms in $C^\otimes$ given by the union of the W^J . By construction, the cocartesian fibration $p_C: C^\otimes \rightarrow \text{Span}(\text{Fin})$ inverts all morphisms in $W^\otimes$, so the universal property of localizations produces a functor

$$p_{C[W^{-1}]}: C^\otimes[(W^\otimes)^{-1}] \rightarrow \text{Span}(\text{Fin}).$$

The localization functor $\gamma^\otimes: C^\otimes \rightarrow C^\otimes[(W^\otimes)^{-1}]$ then becomes a functor over $\text{Span}(\text{Fin})$.

Theorem 20.1.6 (cf. [NS18, Theorem A.7]). *Let (C, W, I) be an ∞ -category with weak equivalences and cofibrations, and assume that C comes equipped with a left derivable symmetric monoidal structure. Then the following statements hold:*

- (1) *The functor $p_{C[W^{-1}]}: C^{\otimes}[(W^{\otimes})^{-1}] \rightarrow \text{Span}(\text{Fin})$ defines a symmetric monoidal ∞ -category.*
- (2) *The localization functor $\gamma^{\otimes}: C^{\otimes} \rightarrow C^{\otimes}[(W^{\otimes})^{-1}]$ encodes a lax symmetric monoidal functor.*
- (3) *The underlying functor of $\gamma^{\otimes}$ exhibits its target as the localization of C at W , thus equipping $C[W^{-1}]$ with the structure of a symmetric monoidal ∞ -category.*
- (4) *For every symmetric monoidal ∞ -category D , the functor $\gamma^{\otimes}$ induces a fully faithful functor*

$$\text{Fun}^{\otimes\text{-lax}}(C[W^{-1}], D) \hookrightarrow \text{Fun}^{\otimes\text{-lax}}(C, D)$$

whose essential image consists of those lax symmetric monoidal functors $C \rightarrow D$ that invert the morphisms in W .

More informally, parts (1)-(3) of the theorem state that the localization $C[W^{-1}]$ is canonically symmetric monoidal, and that the localization functor $\gamma: C \rightarrow C[W^{-1}]$ canonically refines to a lax symmetric monoidal functor. Part (4) states that γ is in fact *initial* among lax symmetric monoidal functors $C \rightarrow D$ which invert the morphisms in W .

Before moving to the proof of this theorem, we record the bounded below case needed in Chapter 6.

Example 20.1.7 (Chain complexes). Let $\mathcal{A}$ be a symmetric monoidal abelian category with enough projectives. Assume that its monoidal unit is projective and that the tensor product of two projective objects is again projective. Equip the category $\text{Ch}^-(\mathcal{A})$ of bounded below chain complexes with the projective cofibration structure from Proposition 6.6.14 and the symmetric monoidal structure from Definition 6.6.22. We claim that this monoidal structure is left derivable. To see this, consider finitely many bounded below projective chain complexes $C_{\bullet}^1, \dots, C_{\bullet}^n$, finitely many tensor-cofibrant bounded below chain complexes $D_{\bullet}^1, \dots, D_{\bullet}^n$, and quasi-isomorphisms $f_i: C_{\bullet}^i \rightarrow D_{\bullet}^i$. By assumption on $\mathcal{A}$, each $D_{\bullet}^i$ is again a bounded below projective chain complex. Hence by Proposition 6.6.11, each f_i is actually a chain homotopy equivalence. Since the tensor product of chain maps preserves chain homotopies in each variable, it follows by induction on n that the tensor product

$$\bigotimes_{i=1}^n f_i: \bigotimes_{i=1}^n C_{\bullet}^i \rightarrow \bigotimes_{i=1}^n D_{\bullet}^i$$

is again a chain homotopy equivalence, and hence a quasi-isomorphism.

Thus Theorem 20.1.6 equips $\text{D}^-(\mathcal{A})$ with a symmetric monoidal structure and the localization $\text{Ch}^-(\mathcal{A}) \rightarrow \text{D}^-(\mathcal{A})$ with a lax symmetric monoidal refinement. For comparison with the construction in Chapter 6, Proposition 6.6.15 identifies $\text{D}^-(\mathcal{A})$ with the localization of the symmetric monoidal category $\text{Ch}_{\text{proj}}^-(\mathcal{A})$ at the quasi-isomorphisms. Since tensor product preserves quasi-isomorphisms between projective complexes in each variable, Proposition 20.1.1 gives a second description of the induced tensor product, namely the derived tensor product constructed in Definition 6.6.24.

In the remainder of this section, we explain the main ingredients for the proof of Theorem 20.1.6.

Recall from Definition 6.6.1 that a total left derived functor of $F: C \rightarrow D$ is a right Kan extension of F along the localization $\gamma: C \rightarrow C[W^{-1}]$. Such a derived functor $\mathbf{L}F$ is called *absolute* if for every functor $H: D \rightarrow E$, the composite $H \circ \mathbf{L}F$ is again a total left derived functor of $H \circ F$.

Example 20.1.8. Assume that $F: C \rightarrow D$ inverts morphisms in W , so that it induces a functor $\mathbf{L}F: C[W^{-1}] \rightarrow D$ by the universal property of localization. Then the natural isomorphism $\mathbf{L}F \circ \gamma \cong F$ exhibits $\mathbf{L}F$ as an absolute left derived functor of F .

Example 20.1.9. More generally, let (C, W, I) be an ∞ -category with weak equivalences and cofibrations, in the sense of Definition 2.2.4, and consider a functor $F: C \rightarrow D$ for which the restriction $F|_{C_c}: C_c \rightarrow D$ to the cofibrant objects inverts weak equivalences. It follows from Proposition 6.6.3 that F admits an absolute left derived functor $\mathbf{L}F: C[W^{-1}] \rightarrow D$, uniquely characterized by the existence of an isomorphism $\mathbf{L}F \circ \gamma \circ i \simeq F \circ i$ on the cofibrant objects.

Definition 20.1.10. Let $p: C \rightarrow S$ be a cocartesian fibration. Assume that for every object $s \in S$ we are given a class W_s of morphisms in the fiber $C_s := p^{-1}(s)$, and let $\gamma_s: C_s \rightarrow C_s[W_s^{-1}]$ denote the localization functor. We say that p is *left derivable with respect to* $\{W_s\}_{s \in S}$ if the following two conditions are satisfied:

- (1) For every morphism $f: s \rightarrow t$ in S , consider the cocartesian transport functor $f_!: C_s \rightarrow C_t$. Then $\gamma_t \circ f_!: C_s \rightarrow C_t[W_t^{-1}]$ admits an absolute left derived functor

$$\mathbf{L}f_!: C_s[W_s^{-1}] \rightarrow C_t[W_t^{-1}].$$

- (2) For two composable morphisms $s \xrightarrow{f} t \xrightarrow{g} u$ in S , the canonical comparison map constructed below

$$\mathbf{L}g_! \circ \mathbf{L}f_! \rightarrow \mathbf{L}(g \circ f)_!$$

is a natural isomorphism.

Observation 20.1.11. The comparison map in part (2) of Definition 20.1.10 is obtained from absoluteness. Write

$$\varepsilon_f: \mathbf{L}f_! \circ \gamma_s \Rightarrow \gamma_t \circ f_!, \quad \varepsilon_g: \mathbf{L}g_! \circ \gamma_t \Rightarrow \gamma_u \circ g_!$$

for the structure transformations. Since $\mathbf{L}f_!$ is absolute, the composite $\mathbf{L}g_! \circ \mathbf{L}f_!$ is a right Kan extension of $\mathbf{L}g_! \circ \gamma_t \circ f_!$ along γ_s . The composite transformation

$$\mathbf{L}g_! \circ \mathbf{L}f_! \circ \gamma_s \Rightarrow \mathbf{L}g_! \circ \gamma_t \circ f_! \Rightarrow \gamma_u \circ g_! \circ f_!$$

therefore induces, by the universal property of $\mathbf{L}(g \circ f)_!$, the canonical comparison

$$\mathbf{L}g_! \circ \mathbf{L}f_! \rightarrow \mathbf{L}(g \circ f)_!.$$

Example 20.1.12. In the situation of Definition 20.1.10, assume that every cocartesian transport functor $f_!: C_s \rightarrow C_t$ satisfies $f_!(W_s) \subseteq W_t$. Then p is left derivable with respect to $\{W_s\}_{s \in S}$. Indeed, the condition on $f_!$ shows that $\gamma_t \circ f_!$ uniquely factors through a functor $\mathbf{L}f_!: C_s[W_s^{-1}] \rightarrow C_t[W_t^{-1}]$, which by Example 20.1.8 is an absolute left derived functor. The second condition is then an immediate consequence of the universal property.

Example 20.1.13. If $p: C \rightarrow S$ is left derivable with respect to $\{W_s\}$, then for any functor $f: T \rightarrow S$ the pullback functor $C \times_S T \rightarrow T$ is left derivable with respect to $\{W_{f(t)}\}_{t \in T}$.

Proposition 20.1.14 ([NS18, Proposition A.14]). *Let $p: C \rightarrow S$ be a cocartesian fibration which is left derivable with respect to classes of morphisms $\{W_s\}_{s \in S}$. Let W denote the collection of morphisms in C given by the union of all W_s , and let $q: C[W^{-1}] \rightarrow S$ denote the unique factorization of p through the localization functor $\gamma: C \rightarrow C[W^{-1}]$. Then the following statements hold true:*

- (1) *The functor q is a cocartesian fibration;*
- (2) *For every object $s \in S$, the map on fibers*

$$\gamma_s: C_s \rightarrow C[W^{-1}]_s$$

induces an equivalence $C_s[W_s^{-1}] \xrightarrow{\sim} C[W^{-1}]_s$;

- (3) *Consider a morphism $f: s \rightarrow t$ in S . Under the equivalences from part (2), the transport functor*

$$C[W^{-1}]_s \rightarrow C[W^{-1}]_t$$

identifies with the total derived functor $\mathbf{L}f_!$ of $f_!: C_s \rightarrow C_t$.

Proof sketch. We refer to [NS18, Proposition A.14] for the proof and only indicate its structure. By Example 20.1.13, one may first restrict to the bases $[0]$ and $[1]$. The result is immediate over $[0]$. Over $[1]$, the cocartesian fibration is classified by a functor $F: C_0 \rightarrow C_1$. Its absolute left derived functor unstraightens to a cocartesian fibration with the desired fibers. The substantive point is that absoluteness identifies the mapping simplex of this derived functor with the localization of the mapping simplex of F , by the universal property of right Kan extension and the description in Example 23.2.4.

Finally, the total category of a cocartesian fibration commutes with colimits in the base, and localization commutes with colimits because it is a left adjoint on relative ∞ -categories. The class of bases for which the proposition holds is therefore closed under colimits. Since Cat_∞ is generated under colimits by $[0]$ and $[1]$ by Proposition 24.1.16, this proves the result for arbitrary S . $\square$

Proposition 20.1.15. *In the situation of Theorem 20.1.6, the cocartesian fibration $p_C: C^\otimes \rightarrow \text{Span}(\text{Fin})$ is left derivable with respect to the classes $\{W^I\}_{I \in \text{Fin}}$ on its fibers $C_I^\otimes \simeq C^I$.*

Proof. Fix a span $\alpha: I \xleftarrow{f} K \xrightarrow{g} J$ in $\text{Span}(\text{Fin})$. Under the identification $C_I^\otimes \simeq C^I$, the cocartesian transport functor along α is given by

$$\alpha_!: C^I \xrightarrow{f^*} C^K \xrightarrow{g_\otimes} C^J, \quad (x_i)_{i \in I} \mapsto \left(\bigotimes_{k \in g^{-1}(j)} x_{f(k)} \right)_{j \in J}.$$

The product category C^I again carries weak equivalences and cofibrations componentwise, by the finite-product version of Exercise 6.6.7. It therefore suffices, by Proposition 6.6.3, to show that $\alpha_!$ preserves componentwise weak equivalences between componentwise cofibrant objects.

This is clear for the restriction functor f^* . For the functor $g_\otimes$, let $\{u_k: x_k \rightarrow y_k\}_{k \in K}$ be a componentwise weak equivalence between componentwise cofibrant tuples. Then every object y_k is in particular tensor-cofibrant, and the j -th component of $g_\otimes(u)$ is the map

$$\bigotimes_{k \in g^{-1}(j)} x_k \xrightarrow{\bigotimes_{k \in g^{-1}(j)} u_k} \bigotimes_{k \in g^{-1}(j)} y_k,$$

which is a weak equivalence by the left derivability assumption. Thus $\alpha_!$ admits an absolute left derived functor

$$\mathbf{L}\alpha_!: C^I[(W^I)^{-1}] \rightarrow C^J[(W^J)^{-1}].$$

It remains to verify compatibility with composition. Consider composable spans

$$\alpha: I \xleftarrow{f} K \xrightarrow{g} J \quad \text{and} \quad \beta: J \xleftarrow{h} L \xrightarrow{k} M.$$

We show that the canonical comparison map

$$\mathbf{L}\beta_! \circ \mathbf{L}\alpha_! \rightarrow \mathbf{L}(\beta \circ \alpha)_!$$

is an isomorphism. Let $X = (X_i)_{i \in I}$ be componentwise cofibrant. Every component of $\alpha_!(X)$ is a finite tensor product of cofibrant objects, and hence is tensor-cofibrant. Choose any componentwise weak equivalence

$$q: P \xrightarrow{\sim} \alpha_!(X)$$

with P componentwise cofibrant. Such a map is obtained by factoring the morphism from the initial object to $\alpha_!(X)$ in C^J , and no functorial choice is required. By the defining property of total left derived functors, we have

$$(\mathbf{L}\beta_! \circ \mathbf{L}\alpha_!)(\gamma_I X) \simeq \mathbf{L}\beta_!(\gamma_J \alpha_!(X)) \simeq \gamma_M \beta_!(P).$$

The structure transformation for $\mathbf{L}\beta_!$ at $\alpha_!(X)$ is represented under this identification by the map

$$\gamma_M \beta_!(q): \gamma_M \beta_!(P) \longrightarrow \gamma_M \beta_! \alpha_!(X).$$

This map is an isomorphism. Indeed, restriction along h preserves weak equivalences, and each component after applying $k_\otimes$ is a tensor product of weak equivalences whose sources are cofibrant and whose targets are tensor-cofibrant; this is a weak equivalence by Definition 20.1.3. The composite structure transformation used to construct the comparison in Observation 20.1.11 is therefore an isomorphism at $\gamma_I X$. The structure transformation for $\mathbf{L}(\beta \circ \alpha)_!$ is also an isomorphism there, since X is componentwise cofibrant. Hence the comparison map is an isomorphism on the image of every componentwise cofibrant object of C^I . By part (1) of Corollary 2.2.9, every object of $C^I[(W^I)^{-1}]$ is isomorphic to the image of such an object, so the comparison map is an isomorphism everywhere. We conclude that p_C is left derivable with respect to $\{W^I\}_{I \in \mathbf{Fin}}$. $\square$

We may now finally give a proof of the main result of this section.

Proof of Theorem 20.1.6. Let us write

$$q: C^\otimes[(W^\otimes)^{-1}] \rightarrow \mathbf{Span}(\mathbf{Fin})$$

for the functor from Construction 20.1.5. By Proposition 20.1.15, the cocartesian fibration $p_C: C^\otimes \rightarrow \text{Span}(\text{Fin})$ is left derivable with respect to the classes $\{W^I\}_{I \in \text{Fin}}$ on its fibers. In particular, it follows from Proposition 20.1.14 that q is a cocartesian fibration. Moreover, for every finite set I , the induced map on fibers

$$C_I^\otimes = C^I \rightarrow C^\otimes[(W^\otimes)^{-1}]_I$$

exhibits the target as the localization $C^I[(W^I)^{-1}]$. Using Exercise 1.5.25, we obtain natural equivalences

$$C^\otimes[(W^\otimes)^{-1}]_I \simeq C^I[(W^I)^{-1}] \simeq (C[W^{-1}])^I.$$

For $i \in I$, let

$$\rho_i: I \hookrightarrow \{i\} \xrightarrow{=} \{i\}$$

be the corresponding inert span. Cocartesian transport along ρ_i in $C^\otimes$ is the projection $\text{pr}_i: C^I \rightarrow C$, which preserves componentwise weak equivalences. Its total left derived functor is therefore the induced projection

$$C^I[(W^I)^{-1}] \longrightarrow C[W^{-1}].$$

Part (3) of Proposition 20.1.14 identifies this with cocartesian transport along ρ_i for q . Consequently, under the equivalences above, the Segal map for the straightening of q is the equivalence

$$C^I[(W^I)^{-1}] \xrightarrow{\sim} (C[W^{-1}])^I.$$

The same statement for $I = \emptyset$ identifies the fiber with the terminal category. Thus the cocartesian straightening of q preserves finite products, and Lemma 14.2.4 shows that q defines a symmetric monoidal ∞ -category. This proves (1).

For (2), the structure transformation for each projection is invertible because the projection itself preserves weak equivalences. Thus $\gamma^\otimes$ carries cocartesian lifts of the spans ρ_i to cocartesian lifts. By part (iii) of Proposition 14.1.7, these lifts exhibit the product decompositions in the source and target total categories. It follows that $\gamma^\otimes$ preserves finite products. Thus it is a morphism of ∞ -operads, in the sense of Definition 14.1.1, and hence encodes a lax symmetric monoidal functor. This proves (2).

For (3), we evaluate part (2) of Proposition 20.1.14 on the one-point set $\langle 1 \rangle$. This identifies the underlying ∞ -category of the symmetric monoidal ∞ -category from (1) with the localization $C[W^{-1}]$, and under this equivalence the restriction of $\gamma^\otimes$ to the fiber over $\langle 1 \rangle$ is precisely the localization functor $\gamma: C \rightarrow C[W^{-1}]$. Thus the symmetric monoidal structure from (1) may be regarded as a symmetric monoidal structure on $C[W^{-1}]$ itself.

Finally, let D be a symmetric monoidal ∞ -category. Since $\gamma^\otimes: C^\otimes \rightarrow C^\otimes[(W^\otimes)^{-1}]$ is the localization of the total category at $W^\otimes$, precomposition with $\gamma^\otimes$ induces a fully faithful functor

$$\text{Fun}_{/\text{Span}(\text{Fin})}(C^\otimes[(W^\otimes)^{-1}], D^\otimes) \hookrightarrow \text{Fun}_{/\text{Span}(\text{Fin})}(C^\otimes, D^\otimes)$$

whose essential image consists of those functors over $\text{Span}(\text{Fin})$ that invert the morphisms in $W^\otimes$. In the sense of Definition 14.1.1, the lax symmetric monoidal functors are precisely the morphisms of ∞ -operads, i.e. the functors over $\text{Span}(\text{Fin})$ that preserve finite products. Since $\gamma^\otimes$ preserves finite products, precomposition with $\gamma^\otimes$ certainly sends lax symmetric monoidal functors to lax symmetric monoidal functors. Conversely, let $F: C^\otimes[(W^\otimes)^{-1}] \rightarrow D^\otimes$ be a functor over $\text{Span}(\text{Fin})$ such that $F \circ \gamma^\otimes$ preserves finite products. By part (2) of Proposition 20.1.14, the functor $\gamma^\otimes$ is essentially

surjective on each fiber, hence essentially surjective on the total category. Since it also preserves finite products, it follows that F preserves finite products as well. We conclude that the displayed fully faithful functor restricts to a fully faithful functor

$$\mathrm{Fun}^{\otimes\text{-lax}}(C[W^{-1}], D) \hookrightarrow \mathrm{Fun}^{\otimes\text{-lax}}(C, D)$$

whose essential image consists of those lax symmetric monoidal functors that invert the morphisms in W . Indeed, for a lax symmetric monoidal functor, inverting W on the fiber over $\langle 1 \rangle$ is equivalent to inverting $W^{\otimes}$ on every fiber, since the induced functor on the fiber over I is the product of the underlying functor on the fiber over $\langle 1 \rangle$. This proves (4). $\square$

20.2 The symmetric monoidal derived category

We finish this chapter by applying the preceding machinery to unbounded chain complexes. Recall from Theorem 6.1.29 that the derived ∞ -category of any abelian category is stable. The construction of the monoidal structure requires the following standard structure on unbounded chain complexes, which we use as a black box.

Proposition 20.2.1 (Unbounded projective cofibrations, [Lur17, Propositions 7.1.2.8 and 7.1.2.11]). *Let R be a commutative ring and let W be the class of quasi-isomorphisms in $\mathrm{Ch}(R)$. There is a class I_{proj} of projective cofibrations with the following properties:*

- (1) *The triple $(\mathrm{Ch}(R), W, I_{\mathrm{proj}})$ is a homotopy cocomplete ∞ -category with weak equivalences and cofibrations.*
- (2) *The unit $R[0]$ is cofibrant, and a finite tensor product of cofibrant complexes is again cofibrant.*
- (3) *If P is cofibrant, then $P \otimes_R -$ is homotopy cocontinuous and preserves quasi-isomorphisms.*

The bounded-below projective cofibrations of Proposition 6.6.14 admit a direct construction from ordinary projective resolutions. The unbounded statement is subtler, which is why we refer to Lurie [Lur17]. Its relevance here is that properties (2) and (3), together with Remark 20.1.4, give exactly the left derivability required in Definition 20.1.3.

Proposition 20.2.2 (The symmetric monoidal derived category). *Let R be a commutative ring. The derived ∞ -category*

$$\mathrm{D}(R) := \mathrm{Ch}(R)[\{\text{quasi-isomorphisms}\}^{-1}]$$

admits a canonical symmetric monoidal structure whose tensor product is the derived tensor product and whose unit is $R[0]$. The localization functor

$$\gamma: \mathrm{Ch}(R) \longrightarrow \mathrm{D}(R)$$

admits a canonical lax symmetric monoidal refinement. Moreover, the derived tensor product preserves small colimits separately in both variables.

Proof. Equip $\mathrm{Ch}(R)$ with the weak equivalences and projective cofibrations of Proposition 20.2.1. Properties (2) and (3) of that proposition show that the ordinary tensor product is left derivable. The

first two assertions therefore follow from Theorem 20.1.6. The description of cocartesian transport in part (3) of Proposition 20.1.14 identifies tensor product in the localization with the total left derived functor of ordinary tensor product, so this is the derived tensor product. Applying the same description to the unique span from $\emptyset$ to $\langle 1 \rangle$ identifies the monoidal unit with the image of the ordinary unit $R[0]$.

It remains to prove the assertion about colimits. By part (1) of Corollary 2.2.9, we may represent an object $X \in D(R)$ by a cofibrant complex P . For a complex Y , choose a cofibrant replacement $Q \xrightarrow{\sim} Y$. The preceding identification and part (3) of Proposition 20.2.1 give

$$X \otimes_R^L \gamma(Y) \simeq \gamma(P \otimes_R Q) \simeq \gamma(P \otimes_R Y).$$

Thus $X \otimes_R^L -$ is the functor on localizations induced by $P \otimes_R -$. The latter is homotopy cocontinuous by part (3) of Proposition 20.2.1. The dual form of Theorem 2.2.11 therefore shows that $X \otimes_R^L -$ preserves small colimits. By symmetry, the same holds in the other variable. $\square$

Proposition 20.2.3 (Bounded-below comparison). *Let R be a commutative ring. The full subcategory $D^-(R) \subseteq D(R)$ of bounded below objects is closed under the derived tensor product and contains the monoidal unit. It therefore inherits a symmetric monoidal structure from $D(R)$ for which the inclusion*

$$D^-(R) \hookrightarrow D(R)$$

is symmetric monoidal. This structure agrees with the symmetric monoidal structure on $D^-(R)$ constructed in Example 20.1.7.

Proof. The unit $R[0]$ is bounded below. Let $X, Y \in D^-(R)$, and choose bounded below projective resolutions $P \rightarrow X$ and $Q \rightarrow Y$. Since projective R -modules are flat, Lemma 6.6.26 shows that tensoring with either P or Q preserves quasi-isomorphisms.

Choose cofibrant replacements $U \xrightarrow{\sim} P$ and $V \xrightarrow{\sim} Q$ in the unbounded projective cofibration structure of Proposition 20.2.1. The composite

$$U \otimes_R V \longrightarrow U \otimes_R Q \longrightarrow P \otimes_R Q$$

is a quasi-isomorphism: this follows for the first map from part (3) of Proposition 20.2.1, and for the second from the preceding observation about Q . Hence the derived tensor product of X and Y in $D(R)$ is represented by the bounded below complex $P \otimes_R Q$. Thus $D^-(R)$ is closed under the ambient derived tensor product, and Lemma 14.5.2 gives the asserted restricted symmetric monoidal structure.

It remains to compare this restriction with the bounded-below construction. The inclusion $\text{Ch}^-(R) \hookrightarrow \text{Ch}(R)$ is symmetric monoidal. Composing it with the lax symmetric monoidal localization $\text{Ch}(R) \rightarrow D(R)$ and applying part (4) of Theorem 20.1.6 to the bounded-below localization produces a lax symmetric monoidal refinement of the inclusion $D^-(R) \hookrightarrow D(R)$. On bounded below projective resolutions P and Q , its binary structure map is represented by the composite quasi-isomorphism $U \otimes_R V \rightarrow P \otimes_R Q$ displayed above, and is therefore an isomorphism. Its unit map is likewise the identity of $R[0]$. The refinement is consequently symmetric monoidal, which identifies the bounded-below structure with the restriction of the unbounded one. $\square$

Corollary 20.2.4 (DGAs as derived algebras). *The lax symmetric monoidal localization of Proposition 20.2.2 induces functors*

$$\text{Alg}(\text{Ch}(R)) \longrightarrow \text{Alg}(D(R)), \quad \text{CAlg}(\text{Ch}(R)) \longrightarrow \text{CAlg}(D(R)).$$

Proof. Lax symmetric monoidal functors preserve associative and commutative algebra objects. $\square$

Thus every associative or commutative DGA determines an associative or commutative algebra object in $D(R)$, respectively.

We finally record the compact presentation of the derived category.

Corollary 20.2.5 (Perfect complexes). *The object $R[0]$ is a compact generator of $D(R)$. If*

$$\text{Perf}(R) := D(R)^\omega$$

denotes the full subcategory of compact objects, then $\text{Perf}(R)$ is the smallest thick subcategory of $D(R)$ containing $R[0]$. Its objects are precisely the retracts of bounded complexes of finitely generated free R -modules, and there is an equivalence of symmetric monoidal ∞ -categories

$$D(R) \simeq \text{Ind}(\text{Perf}(R)).$$

Here the symmetric monoidal structure on the right is the one supplied by Theorem 22.5.4.

Proof. Since R is projective as an R -module, Corollary 6.4.5 gives

$$\pi_n \text{hom}_{D(R)}(R[0], X) \cong H_n(X)$$

for every $X \in D(R)$. The canonical t-structure on $D(R)$ is compatible with filtered colimits by [Lur17, Proposition 1.3.5.21], so each homology functor preserves filtered colimits. Since homotopy groups of spectra preserve filtered colimits by Lemma 4.4.27, the displayed formula shows that $R[0]$ is compact. Its shifts jointly detect zero objects, since a complex is zero in $D(R)$ precisely when all its homology groups vanish. It follows from Proposition 22.3.4 that $R[0]$ is a compact generator.

By Proposition 22.3.4, the compact objects form the smallest thick subcategory containing $R[0]$; in particular, $\text{Perf}(R)$ is small. Bounded complexes of finitely generated free modules are obtained from shifts of $R[0]$ by finite cofiber sequences, using their finite filtrations by degree. Conversely, their retracts form a thick subcategory. This gives the asserted explicit description of $\text{Perf}(R)$.

The underlying equivalence $D(R) \simeq \text{Ind}(\text{Perf}(R))$ now follows from Proposition 22.3.7. The derived tensor product restricts to $\text{Perf}(R)$: this follows from its exactness separately and the description of $\text{Perf}(R)$ as the thick subcategory generated by the unit. The uniqueness assertion in Theorem 22.5.4 upgrades the equivalence to a symmetric monoidal equivalence. $\square$

These results provide the higher-algebra input for the comparison with Eilenberg–MacLane modules. Indeed, the Ind-presentation and Proposition 22.3.6 show that $D(R)$ is presentable. It is therefore a presentably symmetric monoidal stable ∞ -category with compact generator $R[0]$. Applying the monogenic Morita theorem, Theorem 19.5.6, and computing the endomorphism ring spectrum of $R[0]$ gives the symmetric monoidal equivalence

$$D(R) \simeq \text{Mod}_{HR}(\text{Sp}).$$

The computation and its interpretation in terms of Eilenberg–MacLane ring spectra are carried out in Corollary 8.3.4.

Exercises

Part III

Background on ∞ -category theory

Introduction to Part III

This part collects the ∞ -categorical background used throughout the book. The material is standard, but developing it at the point where it is first needed would repeatedly interrupt the main line of argument. These chapters are therefore meant to be consulted rather than read in order: whenever the main text appeals to one of their results, it forward-references the precise statement it uses.

Chapter 21 treats adjunctions between ∞ -categories, their equivalent characterizations, their interaction with (co)limits and Kan extensions, and Bousfield localizations. Chapter 22 develops presentable ∞ -categories through presentations by generators and relations, which is the form in which we use them to construct adjoints and localizations; it also treats compact generation and tensor products of categories with prescribed colimits. Chapter 23 treats cartesian and cocartesian fibrations together with the straightening/unstraightening correspondence, our main device for constructing functors into $\mathbf{Cat}_\infty$, such as the Hom functor $\mathrm{Hom}_C : C^{\mathrm{op}} \times C \rightarrow \mathbf{An}$. Finally, Chapter 24 presents ∞ -categories as complete Segal anima, which is how the span categories of Part II are constructed.

21 Adjunctions

Nearly every construction in this book is specified by a universal property, and in practice this means that it is produced by an adjoint functor: free algebras, localizations, Kan extensions and (co)limits are all of this form. Adjunctions are therefore the tool we reach for most often, and the goal of this chapter is to give a detailed treatment of them, together with the various related notions that the main text calls upon.

We start in Section 21.1 by recalling the definition and providing various equivalent characterizations. In Section 21.2 we discuss the relation between (co)limits and adjunctions, and deduce various useful consequences about limits and colimits. After introducing slice categories in Section 21.3, we discuss Kan extensions and their pointwise formulas in Section 21.4. In Section 21.5 we briefly introduce the notion of cofinality, and in Section 21.6 the filtered and sifted ∞ -categories, whose colimits commute with finite limits and finite products respectively. Section 21.7 proves a comparison theorem for categories of structured objects, the monad-free mechanism underlying the Barr–Beck–Lurie monadicity theorem. Finally, in Section 21.8 we introduce Bousfield localizations, give various examples, and provide a criterion for recognizing them.

21.1 Definition and characterizations

Recall from Definition 1.8.13 the definition of an adjunction between ∞ -categories:

Definition 21.1.1. Let $F: C \rightarrow D$ and $G: D \rightarrow C$ be two functors. An *adjunction* between C and D consists of a natural isomorphism

$$\mathrm{Hom}_D(F(-), -) \cong \mathrm{Hom}_C(-, G(-))$$

of functors $C^{\mathrm{op}} \times D \rightarrow \mathbf{An}$. We often write $F \dashv G$ to indicate that such an isomorphism has been given.

Proposition 21.1.2. Consider two functors $F: C \rightarrow D$ and $G: D \rightarrow C$. The following data are equivalent:

- (1) The data of an adjunction $F \dashv G$;
- (2) The data of a natural transformation $\eta: \mathrm{id}_C \rightarrow GF$, called the unit, such that for all objects X of C and Y of D the induced map

$$\mathrm{Hom}_D(FX, Y) \xrightarrow{G} \mathrm{Hom}_C(GFX, GY) \xrightarrow{- \circ \eta_X} \mathrm{Hom}_C(X, GY)$$

is an equivalence;

- (3) The data of a natural transformation $\varepsilon: FG \rightarrow \text{id}_D$, called the counit, such that for all objects X of C and Y of D the induced map

$$\text{Hom}_C(X, GY) \xrightarrow{F} \text{Hom}_D(FX, FGY) \xrightarrow{\varepsilon_Y \circ -} \text{Hom}_D(FX, Y)$$

is an equivalence;

- (4) The data of both a unit $\eta: \text{id} \rightarrow GF$ and a counit $\varepsilon: FG \rightarrow \text{id}$ together with commutative triangles as follows, called the triangle identities.²⁰

$$\begin{array}{ccc} & FGF & \\ F\eta \nearrow & & \searrow \varepsilon F \\ F & \xlongequal{\quad} & F \end{array} \quad \text{and} \quad \begin{array}{ccc} & GFG & \\ \eta G \nearrow & & \searrow G\varepsilon \\ G & \xlongequal{\quad} & G. \end{array}$$

Proof sketch. First assume (1), so that there is a natural equivalence $\text{Hom}_D(FX, Y) \simeq \text{Hom}_C(X, GY)$. If we fix $X \in C$, then by the Yoneda lemma the natural transformation $\text{Hom}_D(FX, -) \xrightarrow{\sim} \text{Hom}_D(X, G(-))$ corresponds to a map $\eta_X: X \rightarrow GFX$: we obtain this map by taking $Y = FX$ and letting η_X be the map that under this equivalence corresponds to the identity map $\text{id}_{FX}: FX \rightarrow FX$. Conversely, the map η_X determines the transformation as the composite

$$\text{Hom}_D(FX, -) \xrightarrow{G} \text{Hom}_C(GFX, G(-)) \xrightarrow{- \circ \eta_X} \text{Hom}_C(X, -).$$

The naturality of the equivalence in X corresponds to the naturality of $\eta: \text{id}_C \rightarrow GF$, showing the equivalence between (1) and (2).

The equivalence between (1) and (3) is proved similarly, where now the counit map $\varepsilon_Y: FGY \rightarrow Y$ corresponds to the identity map $\text{id}_{GY}: GY \rightarrow GY$.

To show that these conditions are also equivalent to (4), assume first that (1)-(3) are satisfied. For the first triangle identity, observe that by definition the map $\eta_X: X \rightarrow GFX$ is a preimage of $\text{id}_{FX}: FX \rightarrow FX$ under the equivalence

$$\text{Hom}_C(X, GFX) \xrightarrow{F} \text{Hom}_D(FX, FGF) \xrightarrow{\varepsilon_{FX} \circ -} \text{Hom}_D(FX, FX).$$

This precisely says that the composite $\varepsilon_{FX} \circ F\eta_X$ is equivalent to id_{FX} , naturally in X , which is the first triangle identity. The second triangle identity similarly follows from the fact that the morphism $\varepsilon_Y: FGY \rightarrow Y$ is a preimage of $\text{id}_{GY}: GY \rightarrow GY$ under the equivalence

$$\text{Hom}_D(FGY, Y) \xrightarrow{G} \text{Hom}_C(GFGY, GY) \xrightarrow{- \circ \eta_{GY}} \text{Hom}_C(GY, GY).$$

Finally, assume that (4) is satisfied. An easy check shows that in this case the map defined in (2) is inverse to the map defined in (3), producing the desired natural equivalence $\text{Hom}_D(FX, Y) \simeq \text{Hom}_C(X, GY)$. $\square$

It turns out that we may detect left/right functors ‘objectwise’:

Definition 21.1.3. Let $F: C \rightarrow D$ be a functor and let Y be an object of D .

²⁰The details aren’t super relevant here, but strictly speaking the data of an adjunction should only contain either the first or the second triangle identity, while for the other triangle we only demand the *existence*; see e.g. [Lur26, Tag 02F4].

- (1) An object Z of C is called a *right adjoint object to Y under F* if it comes equipped with a map $\varepsilon_Y: FZ \rightarrow Y$ such that for every other object X of C the induced map

$$\mathrm{Hom}_C(X, Z) \xrightarrow{F} \mathrm{Hom}_D(FX, FZ) \xrightarrow{\varepsilon_Y \circ -} \mathrm{Hom}_D(FX, Y)$$

is an equivalence of anima.

- (2) Dually, Z is called a *left adjoint object to Y under F* if it comes equipped with a map $\eta_Y: Y \rightarrow FZ$ such that for every other object X of C the induced map

$$\mathrm{Hom}_C(Z, X) \xrightarrow{F} \mathrm{Hom}_D(FZ, FX) \xrightarrow{- \circ \eta_Y} \mathrm{Hom}_D(Y, FX)$$

is an equivalence of anima.

Lemma 21.1.4. *Consider a functor $F: C \rightarrow D$ and let $D_R \subseteq D$ be a full subcategory. Assume that for every $Y \in D_R$ there exists a right adjoint object Z to Y under F . Then these right adjoint objects assemble into a functor $G: D_R \rightarrow C$ which comes equipped with a natural isomorphism*

$$\mathrm{Hom}_D(F(-), -) \cong \mathrm{Hom}_C(-, G(-))$$

of functors $C^{\mathrm{op}} \times D_R \rightarrow \mathbf{An}$.

Dually, if $D_L \subseteq D$ is a subcategory such that every $Y \in D_L$ admits a left adjoint object under F , then these left adjoint objects assemble into a functor $L: D_L \rightarrow C$ that comes equipped with a natural isomorphism

$$\mathrm{Hom}_C(L(-), -) \cong \mathrm{Hom}_D(-, F(-))$$

of functors $D_L^{\mathrm{op}} \times C \rightarrow \mathbf{An}$.

Proof. We will prove the claim for right adjoints; the claim for left adjoints is dual. The assumption on F is that for every Y in D_R the functor $\mathrm{Hom}_D(F(-), Y): C^{\mathrm{op}} \rightarrow \mathbf{An}$ is representable, in the sense that it lies in the image of the Yoneda embedding $Y: C \hookrightarrow \mathrm{Fun}(C^{\mathrm{op}}, \mathbf{An})$. Since the Yoneda embedding is fully faithful, this means there exists a functor

$$G: D_R \rightarrow C$$

making the following diagram commute:

$$\begin{array}{ccc} & & C \\ & \nearrow G & \downarrow Y \\ D_R & \xrightarrow{Y \mapsto \mathrm{Hom}_D(F(-), Y)} & \mathrm{Fun}(C^{\mathrm{op}}, \mathbf{An}). \end{array}$$

Here the bottom functor is the one obtained from currying the functor $\mathrm{Hom}_D(F(-), -): C^{\mathrm{op}} \times D_R \rightarrow \mathbf{An}$. Since $Y(X) = \mathrm{Hom}_C(-, X)$, this commutative triangle precisely encodes a natural isomorphism

$$\mathrm{Hom}_D(F(-), -) \cong \mathrm{Hom}_C(-, G(-))$$

of functors $C^{\mathrm{op}} \times D_R \rightarrow \mathbf{An}$, as desired. $\square$

Corollary 21.1.5. *A functor $F: C \rightarrow D$ admits a right adjoint if and only if every object of D admits a right adjoint object under F . Dually, F admits a left adjoint if and only if every object of D admits a left adjoint object under F .*

Proof. We again only prove the claim about right adjoints. If a right adjoint $G: D \rightarrow C$ exists, then by Proposition 21.1.2 we see that the object $Z := GY$ together with the counit map $\varepsilon_Y: FGY \rightarrow Y$ provides a right adjoint object to Y under F . Conversely, if all the right adjoint objects exists, then we may apply Lemma 21.1.4 with $D = D_R$ to obtain a functor $G: D \rightarrow C$ together with a natural isomorphism

$$\mathrm{Hom}_C(F(-), -) \cong \mathrm{Hom}_D(-, G(-)),$$

which means that G is a right adjoint to F . □

Every adjunction automatically induces new adjunctions at the level of functor categories:

Lemma 21.1.6. *Let $F \dashv G$ be an adjunction of ∞ -categories. Then:*

(1) *For every ∞ -category E the functors*

$$G^*: \mathrm{Fun}(C, E) \rightleftarrows \mathrm{Fun}(D, E) : F^*$$

form an adjunction, with unit and counit given by

$$G^*F^* = (FG)^* \xrightarrow{\varepsilon^*} \mathrm{id}_D^* = \mathrm{id}_{\mathrm{Fun}(D, E)} \quad \text{and} \quad \mathrm{id}_{\mathrm{Fun}(C, E)} = \mathrm{id}_C^* \xrightarrow{\eta^*} (GF)^* = F^*G^*.$$

(2) *For every ∞ -category E the functors*

$$F_*: \mathrm{Fun}(E, C) \rightleftarrows \mathrm{Fun}(E, D) : G_*$$

form an adjunction, with unit and counit given by

$$F_*G_* = (FG)_* \xrightarrow{\varepsilon_*} (\mathrm{id}_D)_* = \mathrm{id}_{\mathrm{Fun}(D, E)} \quad \text{and} \quad \mathrm{id}_{\mathrm{Fun}(C, E)} = (\mathrm{id}_C)_* \xrightarrow{\eta_*} (GF)_* = G_*F_*.$$

Proof. The triangle identities for ε^* and η^* , resp. ε_* and η_* , follow immediately from the triangle identities for ε and η . We leave the details to the reader. □

Every adjunction induces an equivalence on geometric realizations:

Lemma 21.1.7. *Let $F: C \rightleftarrows D : G$ be an adjunction of ∞ -categories. Then the induced maps $\|F\|: \|C\| \rightarrow \|D\|$ and $\|G\|: \|D\| \rightarrow \|C\|$ are inverse equivalences.*

Proof. Thinking of the counit $\varepsilon: FG \rightarrow \mathrm{id}_D$ as a functor $[1] \times D \rightarrow D$, it induces another functor

$$[1] \times \|D\| \rightarrow \|[1]\| \times \|D\| \xrightarrow{1.5.25} \|[1] \times D\| \xrightarrow{\|\varepsilon\|} \|D\|,$$

i.e. a natural transformation $\|F\|\|G\| = \|FG\| \rightarrow \|\mathrm{id}_D\| = \mathrm{id}_{\|D\|}$. Since $\|D\|$ is an anima, this natural transformation is automatically a natural isomorphism by Axiom J. Similarly, the unit $\eta: \mathrm{id}_C \rightarrow GF$ induces a natural transformation $\|\eta\|: \mathrm{id}_{\|C\|} \rightarrow \|G\|\|F\|$. This shows that $\|G\|$ is inverse to $\|F\|$, as desired. □

21.2 (Co)limits and adjunctions

Just like for classical categories, we may describe the existence of (co)limits in terms of adjunctions:

Lemma 21.2.1. *Let I and C be ∞ -categories and let $F: I \rightarrow C$ be a functor.*

- (1) *An object W in C equipped with a cone $\varepsilon_F: \text{const}_W \rightarrow F$ is a limit of F in C , in the sense of Definition 1.7.1, if and only if ε_F exhibits W as a right adjoint object to F under the functor $\text{const}: C \rightarrow \text{Fun}(I, C)$, in the sense of Definition 21.1.3.*
- (2) *Dually, an object W equipped with a cocone $\eta_F: F \rightarrow \text{const}_W$ is a colimit of F in C if and only if it exhibits W as a left adjoint object to F under $\text{const}: C \rightarrow \text{Fun}(I, C)$.*

Proof. This is simply a matter of unwinding definitions: the condition in (1) that W is a right adjoint object to F under const means that for every other object X of C the composite

$$\text{Hom}_C(X, W) \xrightarrow{\text{const}} \text{Nat}(\text{const}_X, \text{const}_W) \xrightarrow{\varepsilon} \text{Nat}(\text{const}_X, F)$$

is an equivalence, where we recall that we write $\text{Nat}(-, -)$ for the hom anima in $\text{Fun}(I, C)$. But this is precisely the universal property of the limit from Definition 1.7.1! $\square$

Corollary 21.2.2. *Let I and C be ∞ -categories. Then C admits all I -indexed limits if and only if the functor*

$$\text{const}: C \rightarrow \text{Fun}(I, C), \quad W \mapsto \text{const}_W$$

admits a right adjoint

$$\lim_I: \text{Fun}(I, C) \rightarrow C.$$

Dually, C admits all I -indexed colimits if and only if $\text{const}: C \rightarrow \text{Fun}(I, C)$ admits a left adjoint

$$\text{colim}_I: \text{Fun}(I, C) \rightarrow C.$$

Proof. In light of Lemma 21.2.1 this is an immediate consequence of the pointwise criterion for adjoints from Corollary 21.1.5. $\square$

As a particular consequence of the previous corollary, we see that limit cones and colimit cocones are fully functorial in the functor $F: I \rightarrow C$: the transformations

$$\text{const}_{\lim F} \rightarrow F \quad \text{and} \quad F \rightarrow \text{const}_{\text{colim } F}$$

are given by the counit and unit, respectively, of the adjunctions $\text{const} \dashv \lim_I$ and $\text{colim}_I \dashv \text{const}$.

Remark 21.2.3. For an object x of C , it follows from Corollary 21.2.2 that x is terminal if and only if the functor

$$x: * \rightarrow C$$

is right adjoint to the functor $p_C: C \rightarrow *$. Dually, x is initial if and only if it is *left* adjoint to $p_C: C \rightarrow *$.

Lemma 21.2.4. *Let i be an initial object of an ∞ -category I . Then every ∞ -category C admits I -indexed limits, and the limit functor is given by evaluation at i :*

$$\lim_I = \text{ev}_i : \text{Fun}(I, C) \rightarrow C.$$

Dually, if i is terminal in I then C admits I -indexed colimits which are given by evaluating at i .

Proof. By Corollary 21.2.2 it suffices to show that ev_i defines a right adjoint to $\text{const} : C \rightarrow \text{Fun}(I, C)$. This adjunction may be obtained by considering the adjunction $p_I : I \rightarrow * : i$ and passing to functor categories using Lemma 21.1.6.

(An alternative proof would be to cite Lemma 21.5.4.) $\square$

Lemma 21.2.5. *Let $F : C \rightarrow D$ be a functor and let I be an ∞ -category such that both C and D admit I -indexed limits. Then F preserves I -indexed limits if and only if the ‘canonical’ natural transformation*

$$F(\lim_I(-)) \rightarrow \lim_I(F_*(-))$$

of functors $\text{Fun}(I, C) \rightarrow D$ is a natural isomorphism.

Proof. Consider the following commutative diagram:

$$\begin{array}{ccc} C & \xrightarrow{F} & D \\ \text{const} \downarrow & & \downarrow \text{const} \\ \text{Fun}(I, C) & \xrightarrow{F_*} & \text{Fun}(I, D). \end{array}$$

The ‘canonical’ transformation above is the one that under the adjunction $\text{const} \dashv \lim_I$ corresponds to the transformation

$$\text{const}_{F(\lim_I(-))} \cong F_*(\text{const}_{\lim_I(-)}) \rightarrow F_*(-)$$

induced by the counit $\text{const}_{\lim_I(X)} \rightarrow X$. By Axiom J we may check that this map is an isomorphism for every fixed diagram $X : I \rightarrow C$. But now we observe that the map $F(\lim_I(X)) \rightarrow \lim_I(F_*(-))$ is an isomorphism if and only if the associated cone $\text{const}_{F(\lim_I(X))} \rightarrow F_*(X) = F \circ X$ is a limit cone, giving the claim. $\square$

Lemma 21.2.6. *Let $F : C \rightleftarrows D : G$ be an adjunction of ∞ -categories. Then F preserves all colimits that exist in C and G preserves all limits that exist in D .*

Proof. We prove that F preserves colimits; the proof for G preserving limits is dual. Let $X : I \rightarrow C$ be a diagram that admits a colimit in C , and let $\eta : X \rightarrow \text{const}_{\text{colim}_I X}$ denote the associated colimit cone. We need to show that the cocone

$$F(\eta) : F \circ X \rightarrow F(\text{const}_{\text{colim}_I X}) = \text{const}_{F(\text{colim}_I X)}$$

is a colimit cone in D . By definition, this means that for every other object W in D the map

$$\text{Hom}_D(F(\text{const}_{\text{colim}_I X}), W) \rightarrow \text{Nat}(F \circ X, \text{const}_W)$$

is an equivalence of anima. But under the adjunction $F \dashv G$ and the induced adjunction $F \circ - \dashv G \circ -$ from Lemma 21.1.6, this map is equivalent to the map

$$\text{Hom}_C(\text{const}_{\text{colim}_I X}, G(W)) \rightarrow \text{Nat}(X, G \circ \text{const}_W) = \text{Nat}(X, \text{const}_{G(W)}).$$

Since the latter map is an equivalence by the universal property of the colimit $\operatorname{colim}_I X$, this finishes the proof. $\square$

As a consequence, we see that limits commute with limits and that colimits commute with colimits.

Corollary 21.2.7. *If C admits I -indexed limits, then $\lim_I: \operatorname{Fun}(I, C) \rightarrow C$ preserves all limits that exist in $\operatorname{Fun}(I, C)$. If C admits I -indexed colimits, then $\operatorname{colim}_I: \operatorname{Fun}(I, C) \rightarrow C$ preserves all colimits that exist in $\operatorname{Fun}(I, C)$.*

Lemma 21.2.8. *Consider ∞ -categories C and I , and assume that C admits all I -indexed limits (resp. colimits).*

- (1) *For every ∞ -category E , the functor category $\operatorname{Fun}(E, C)$ admits I -indexed limits (resp. colimits).*
- (2) *For every functor $E' \rightarrow E$, the restriction functor $\operatorname{Fun}(E, C) \rightarrow \operatorname{Fun}(E', C)$ preserves I -indexed limits (resp. colimits).*
- (3) *In particular, for every object X of E the functor*

$$\operatorname{ev}_X: \operatorname{Fun}(E, C) \rightarrow C$$

preserves I -indexed limits (resp. colimits).

Proof. We treat the case for limits; the case for colimits is dual. By Corollary 21.2.2 there is an adjunction

$$\operatorname{const}: C \rightleftarrows \operatorname{Fun}(I, C) : \lim_I.$$

By Lemma 21.1.6 this adjunction induces an adjunction on functor categories of the form

$$\operatorname{Fun}(E, C) \rightleftarrows \operatorname{Fun}(E, \operatorname{Fun}(I, C)) \simeq \operatorname{Fun}(I, \operatorname{Fun}(E, C)).$$

By applying Corollary 21.2.2 another time this says that $\operatorname{Fun}(E, C)$ admits I -indexed limits.

Observe that the limit-functor for $\operatorname{Fun}(E, C)$ is given as the composite

$$\operatorname{Fun}(I, \operatorname{Fun}(E, C)) \simeq \operatorname{Fun}(E, \operatorname{Fun}(I, C)) \xrightarrow{(\lim_I)_*} \operatorname{Fun}(E, C).$$

This description makes it clear that precomposition with any functor $E' \rightarrow E$ preserves I -limits. $\square$

Since the ∞ -category $\mathbf{An}$ admits small limits and colimits, this immediately implies:

Corollary 21.2.9. *Let C be an ∞ -category. The presheaf category $\mathbf{PSh}(C)$ admits all small limits and colimits. The evaluation functors $\operatorname{ev}_x: \mathbf{PSh}(C) \rightarrow \mathbf{An}$ preserve all small limits and colimits. $\square$*

As another consequence, we get the following result on colimits along product categories:

Lemma 21.2.10. *Let I, J and C be ∞ -categories such that C admits all I -indexed and J -indexed colimits. Then C admits all $(I \times J)$ -indexed colimits, and for a functor $F: I \times J \rightarrow C$ we have*

$$\operatorname{colim}_{(i,j) \in I \times J} F(i, j) \cong \operatorname{colim}_{i \in I} \operatorname{colim}_{j \in J} F(i, j).$$

Proof. The adjunction $\text{colim}_J : \text{Fun}(J, C) \rightleftarrows C : \text{const}$ induces an adjunction

$$\text{Fun}(I \times J, C) \simeq \text{Fun}(I, \text{Fun}(J, C)) \rightleftarrows \text{Fun}(I, C)$$

by applying Lemma 21.1.6. Composing this with the adjunction $\text{colim}_I : \text{Fun}(I, C) \rightleftarrows C : \text{const}$ then gives the result. $\square$

A fact which will frequently be useful is that colimits along an indexing category I which itself may be written as a colimit may be computed as an iterated colimit:

Theorem 21.2.11. *Let J be a small ∞ -category and consider a J -indexed family of small ∞ -categories $I_\bullet : J \rightarrow \text{Cat}_\infty$. Define $I := \text{colim}_j I_j$, and let $\varepsilon : I_\bullet \rightarrow \text{const}_I$ denote the colimit cone, giving functors $\varepsilon_j : I_j \rightarrow I$. Let C be an ∞ -category and let $F : I \rightarrow C$ be a functor.*

(1) *Assume that the restricted diagram*

$$F|_{I_j} : I_j \xrightarrow{\varepsilon_j} I \xrightarrow{F} C$$

admits a colimit for every $j \in J$. Then these colimits assemble into a functor

$$F' : J \rightarrow C, \quad j \mapsto \text{colim}(F|_{I_j} : I_j \rightarrow C).$$

(2) *In this case, the functor F admits a colimit in C if and only if the functor F' admits a colimit in C , and then we have*

$$\text{colim}_{i \in I} F(i) \cong \text{colim}_{j \in J} \text{colim}(F|_{I_j}).$$

Dually, if each $F|_{I_j}$ has a limit, then we have $\lim_J F \cong \lim_{j \in J^{\text{op}}} \lim_{I_j} F|_{I_j}$ as soon as either side exists.

Proof. **TO DO.** $\square$

21.3 Slice categories

A useful categorical construction, which we will need in the next two sections, is that of the (*relative*) *slice category*.

Definition 21.3.1 (Slice category). Let C be an ∞ -category and let x be an object in C . We define the *slice categories* $C_{/x}$ and $C_{x/}$, also known as the *over category* and the *under category*, respectively, via the following two pullback squares:

$$\begin{array}{ccc} C_{/x} & \longrightarrow & \text{Ar}(C) \\ \downarrow & \lrcorner & \downarrow t \\ * & \xrightarrow{x} & C \end{array} \quad \text{and} \quad \begin{array}{ccc} C_{x/} & \longrightarrow & \text{Ar}(C) \\ \downarrow & \lrcorner & \downarrow s \\ * & \xrightarrow{x} & C. \end{array}$$

Note that an object of $C_{/x}$ is a pair $(y, f: y \rightarrow x)$ of an object y equipped with a morphism to x , and dually objects of $C_{x/}$ are pairs $(y, f: x \rightarrow y)$. As a result of Proposition 1.4.4 morphisms $(y, f) \rightarrow (y', f')$ in $C_{/x}$ may equivalently be encoded by commutative triangles of the form

$$\begin{array}{ccc} y & \xrightarrow{g} & y' \\ & \searrow f & \swarrow f' \\ & x, & \end{array}$$

and dually for morphisms in $C_{x/}$.

We denote by

$$s: C_{/x} \rightarrow C \quad \text{and} \quad t: C_{x/} \rightarrow C$$

the assignments $s(y, f: y \rightarrow x) = y$ and $t(y, f: x \rightarrow y) = y$, i.e. we first include the slice into the arrow category $\text{Ar}(C)$ and then apply the source/target functor.

Lemma 21.3.2. *Consider an object x of C and consider two objects (y, f) and (y', f') of the slice category $C_{/x}$. Then the hom anima in $C_{/x}$ sits in a pullback square of the form*

$$\begin{array}{ccc} \text{Hom}_{C_{/x}}((y, f), (y', f')) & \longrightarrow & \text{Hom}_C(y, y') \\ \downarrow & \lrcorner & \downarrow f' \circ - \\ * & \xrightarrow{f} & \text{Hom}_C(y, x). \end{array}$$

A dual formula holds for the hom anima of $C_{x/}$.

Proof. By Exercise 1.5, the hom anima in $C_{/x}$ may be computed as the following pullback:

$$\begin{array}{ccc} \text{Hom}_{C_{/x}}((y, f), (y', f')) & \longrightarrow & \text{Nat}(f: y \rightarrow x, f': y \rightarrow x) \\ \downarrow & \lrcorner & \downarrow \\ * & \xrightarrow{\text{id}_x} & \text{Hom}_C(x, x). \end{array}$$

Using the equivalence $\text{Fun}([1] \times [1], C) \simeq \text{Fun}([2], C) \times_{\text{Ar}(C)} \text{Fun}([2], C)$ from the commutative square axiom, the top right hom anima in $\text{Ar}(C)$ sits in a pullback square as follows:

$$\begin{array}{ccc} \text{Nat}(f: y \rightarrow x, f': y \rightarrow x) & \longrightarrow & \text{Hom}_C(y, y') \\ \downarrow & \lrcorner & \downarrow f' \circ - \\ \text{Hom}_C(x, x) & \xrightarrow{- \circ f} & \text{Hom}(y, x). \end{array}$$

The claim thus follows by pasting pullback squares. $\square$

Corollary 21.3.3. *Let x be an object of C . Then the slice category $C_{/x}$ admits a terminal object (x, id_x) , while $C_{x/}$ admits an initial object (x, id_x) .*

Proof. The claim for $C_{x/}$ is immediate from Lemma 21.3.2, since for every $(y, f) \in C_{/x}$ the induced map $\text{id}_x \circ -: \text{Hom}_C(y, x) \rightarrow \text{Hom}_C(y, x)$ is an equivalence, hence its fiber over f is contractible. The claim for $C_{/x}$ is dual. $\square$

Lemma 21.3.4. *Let x be an object in an ∞ -category C .*

- (1) *The object x is terminal in C if and only if the forgetful functor $s: C_{/x} \rightarrow C$ is an equivalence;*
- (2) *The object x is initial in C if and only if the forgetful functor $t: C_{x/} \rightarrow C$ is an equivalence.*

Proof. We only prove (1); the case of (2) is dual. If $C_{/x} \rightarrow C$ is an equivalence, then it follows from Corollary 21.3.3 that x is terminal in x , as it is the image under this equivalence of the terminal object $(x, \text{id}_x) \in C_{/x}$. So assume that x is a terminal object. By Remark 21.2.3, this means that the functor $x: * \rightarrow C$ is a right adjoint to $p_C: * \rightarrow C$, and in particular there is a natural transformation $\varepsilon: \text{id}_C \rightarrow \text{const}_x$ satisfying the triangle identity. We may now construct a functor

$$\varphi: C \rightarrow C_{/x}, \quad y \mapsto (y, \varepsilon_y: y \rightarrow x).$$

More precisely, we may think of ε as a functor $\bar{\varepsilon}: C \rightarrow \text{Ar}(C)$, and since the target of ε is const_x we see that $\bar{\varepsilon}$ canonically factors through the slice $C_{/x}$. It is also clear from this definition that we have $s \circ \varphi \cong \text{id}_C$. It remains to show that we also have $\varphi \circ s \cong \text{id}_{C_{/x}}$.

The functor $\bar{\varepsilon}: C \rightarrow \text{Ar}(C)$ induces a functor $\bar{\varepsilon}_*: \text{Ar}(C) \rightarrow \text{Fun}([1] \times [1], C)$, which informally speaking sends a morphism $f: y \rightarrow z$ to the ‘naturality square’ induced by ε :

$$\begin{array}{ccc} y & \xrightarrow{\varepsilon_y} & x \\ f \downarrow & & \parallel \\ z & \xrightarrow{\varepsilon_z} & x. \end{array}$$

Note that for $z = x$, it follows from the triangle identities for the adjunction that the map $\varepsilon_x: x \rightarrow x$ is the identity map. So restricting the naturality square to objects of the form $(y, f) \in C_{/x}$, this square takes the form

$$\begin{array}{ccc} y & \xrightarrow{\varepsilon_y} & x \\ f \downarrow & & \parallel \\ x & \xlongequal{\quad} & x. \end{array}$$

This shows that the object (y, f) of $C_{/x}$ is naturally isomorphic to $(y, \varepsilon_y: y \rightarrow x)$, producing the desired natural isomorphism $\varphi \circ s \cong \text{id}_{C_{/x}}$. $\square$

Definition 21.3.5 (Relative slice category). For a functor $F: C \rightarrow D$ and an object d of D , we define the *relative slice categories* $C_{/d}$ and $C_{d/}$ via the following two pullback squares:

$$\begin{array}{ccc} C_{/d} & \longrightarrow & D_{/d} \\ s \downarrow & \lrcorner & \downarrow s \\ C & \xrightarrow{F} & D \end{array} \quad \text{and} \quad \begin{array}{ccc} C_{d/} & \longrightarrow & D_{d/} \\ t \downarrow & \lrcorner & \downarrow t \\ C & \xrightarrow{F} & D \end{array}$$

Objects of $C_{/d}$ consist of pairs $(y, f: F(y) \rightarrow d)$, and objects of $C_{d/}$ consist of pairs $(y, f: d \rightarrow F(y))$.

Remark 21.3.6. These relative slice categories are sometimes also denoted by F/d and d/F , which makes their dependencies on F more transparent.

21.4 Kan extensions

Definition 21.4.1. Let I, J and C be ∞ -categories and let $\varphi: I \rightarrow J$ be a functor.

- (1) Given a functor $F: I \rightarrow C$, a *left Kan extension of F along φ* is a left adjoint object to F under the restriction functor $\varphi^*: \text{Fun}(J, C) \rightarrow \text{Fun}(I, C)$. More explicitly, it is a functor $\varphi_!(F): J \rightarrow C$ equipped with a natural transformation $\eta_F: F \rightarrow \varphi^* \varphi_!(F)$ such that for every other functor $G: J \rightarrow C$ the composite

$$\text{Nat}(\varphi_!(F), G) \xrightarrow{\varphi^*} \text{Nat}(\varphi^* \varphi_!(F), \varphi^* G) \xrightarrow{- \circ \eta_F} \text{Nat}(F, \varphi^* G)$$

is an equivalence.

- (2) Dually, a *right Kan extension of F along φ* is a right adjoint object to F under φ^* , i.e. a functor $\varphi_*(F): J \rightarrow C$ equipped with a transformation $\eta_F: \varphi^* \varphi_*(F) \rightarrow F$ inducing equivalences

$$\text{Nat}(G, \varphi_*(F)) \xrightarrow{\sim} \text{Nat}(\varphi^*(G), F)$$

for all $G: J \rightarrow C$.

If every functor $F: I \rightarrow C$ admits a left Kan extension along φ , then by Corollary 21.1.5 these left Kan extensions assemble into a left adjoint

$$\varphi_!: \text{Fun}(I, C) \rightarrow \text{Fun}(J, C)$$

to the restriction functor $\varphi^*: \text{Fun}(J, C) \rightarrow \text{Fun}(I, C)$, called the functor of *left Kan extension along φ* . Dually, if every functor F admits a *right Kan extension along φ* then φ^* admits a right adjoint

$$\varphi_*: \text{Fun}(I, C) \rightarrow \text{Fun}(J, C)$$

called *right Kan extension along φ* .

Observation 21.4.2. Note that the left Kan extension of a functor $F: I \rightarrow C$ along the functor $p_I: I \rightarrow *$ is precisely a colimit of F . Dually, a right Kan extension of F along p_C is a limit of F .

The most useful and common form of Kan extensions are the *pointwise Kan extensions*: those that can be computed using explicit pointwise formulas. In practice, the only Kan extensions one cares about are the pointwise Kan extensions.

Theorem 21.4.3. Let $F: I \rightarrow C$ and $\varphi: I \rightarrow J$ be functors.

- (1) Assume that for every object j of J the composite functor

$$I_{/j} \xrightarrow{s} I \xrightarrow{F} C$$

admits a colimit in C . Then these colimits assemble into a functor

$$\varphi_!(F): J \rightarrow C.$$

Furthermore, the maps

$$F(i) \simeq \operatorname{colim}_{i' \in I_{/i}} F(i') \rightarrow \operatorname{colim}_{i' \in I_{/\varphi(i)}} F(i') = \varphi_!(F)(\varphi(i)) = \varphi^* \varphi_!(F)(i)$$

assemble into a natural transformation $\eta_F: F \rightarrow \varphi^* \varphi_!(F)$ exhibiting $\varphi_!(F)$ as a left Kan extension of F along φ .

(2) Dually, if for every j in J the composite

$$I_{j/} \xrightarrow{t} I \xrightarrow{F} C$$

admits a limit in C , then these limits assemble into functor

$$\varphi_*(F): J \rightarrow C$$

which is a right Kan extension to F .

Proof. **TO DO.** Unfortunately we do not have the techniques yet to be able to prove this result. I may or may not type up a proof of this result in Chapter 23 in the future. $\square$

Definition 21.4.4. If the condition in (1) is satisfied, we say that F admits a pointwise left Kan extension along φ , and if the condition in (2) is satisfied we say that F admits a pointwise right Kan extension along φ .

In general, the Kan extension of F might not actually be an ‘extension’, in the sense that the unit map $F \rightarrow \varphi^*(\varphi_! F)$ and counit map $\varphi^*(\varphi_* F) \rightarrow F$ may not be invertible. However, this is the case whenever φ is fully faithful, explaining the terminology. For this we need the following alternative characterization of fully faithful functors:

Lemma 21.4.5. Let $F: C \rightarrow D$ be a functor. Then the following conditions are equivalent:

- (1) The functor F is fully faithful;
- (2) For every object y of C the commutative square

$$\begin{array}{ccc} C_{/y} & \xrightarrow{F} & C_{/Fy} \\ s \downarrow & & \downarrow s \\ C & \xrightarrow{F} & D \end{array}$$

is a pullback square;

- (3) For every object x of C the commutative square

$$\begin{array}{ccc} C_{y/} & \xrightarrow{F} & C_{Fx/} \\ t \downarrow & & \downarrow t \\ C & \xrightarrow{F} & D \end{array}$$

is a pullback square.

Proof sketch. We prove the equivalence between (1) and (2); the equivalence between (1) and (3) is dual. The two vertical functors in the commutative square in (2) are *right fibrations*, in a sense to be explained below in Chapter 23. One can prove that a functor between two right fibrations is an equivalence if and only if it induces equivalences on all fibers. Since the induced map on fibers is precisely the map $\mathrm{Hom}_C(x, y) \rightarrow \mathrm{Hom}_C(Fx, Fy)$, it follows that condition (2) is equivalent to full faithfulness of F . $\square$

Lemma 21.4.6. *Assume that $\varphi: I \hookrightarrow J$ is a fully faithful functor, and let $F: I \rightarrow C$ be a functor.*

- (1) *Assume that F admits a pointwise left Kan extension $\varphi_!(F)$ along φ . Then the unit map $\eta_F: F \rightarrow \varphi^* \varphi_!(F)$ is a natural isomorphism.*
- (2) *Dually, if F admits a pointwise right Kan extension $\varphi_*(F)$ along φ then the counit $\varepsilon_F: \varphi^* \varphi_*(F) \rightarrow F$ is a natural isomorphism.*

Proof. We prove (1); the proof of (2) is dual. By Axiom J we may check that η_F is a pointwise isomorphism. For an object i of I , the map $F(i) \rightarrow \varphi^* \varphi_!(F)$ is induced by the map of slices

$$I_{/i} \rightarrow I_{/\varphi(i)}, \quad (i', f: i' \rightarrow i) \mapsto (i', F(f): F(i') \rightarrow F(i)).$$

But since φ is fully faithful, this functor is an equivalence by Lemma 21.4.5, and it follows that $\eta_F(i)$ is an isomorphism. $\square$

21.5 Cofinal functors

We will briefly discuss the useful notion of *initial/final functors*. The following result is taken as a black box:

Theorem 21.5.1 (Joyal’s version of Quillen’s Theorem A, [Lur09b, Theorem 4.1.3.1]). *For a functor $\alpha: I \rightarrow J$, the following two conditions are equivalent:*

- (1) *A functor $F: J \rightarrow C$ has a colimit if and only if the functor $F \circ \alpha: I \rightarrow C$ has a colimit, and in this case the canonical map*

$$\mathrm{colim}_I (F \circ \alpha) \rightarrow \mathrm{colim}_J F$$

is an isomorphism in C ;

- (2) *For every object j of J , the relative slice category $I_{j/} = I \times_J J_{j/}$ is weakly contractible, meaning that its geometric realization $\|I_{j/}\|$ is contractible.*

The dual result is also true, where in (1) we use limits and in (2) we use the under categories $I_{j/}$.

Definition 21.5.2. A functor α satisfying the conditions of the theorem is called *final*. We say that α is *initial* if $\alpha^{\mathrm{op}}: I^{\mathrm{op}} \rightarrow J^{\mathrm{op}}$ is final (i.e. each relative slice $I_{j/}$ is contractible, or equivalently limits of functors $F: J \rightarrow C$ may be computed as limits of $F \circ \alpha: I \rightarrow C$).

Warning 21.5.3. The terminology regarding final functors is extremely inconsistent in the literature. Because of historic reasons, final functors are also called *cofinal* functors, where the ‘co’ roughly means ‘jointly final’ and has nothing to do with the usual meaning of ‘co’ in category theory. Nevertheless, some authors use the words ‘final’ and ‘cofinal’ for the two notions from Definition 21.5.2, and there is no consensus as to which of the two words should refer to which of the two notions.

We hope that the convention in Definition 21.5.2 is easy to remember in light of the following lemma:

Lemma 21.5.4. *Consider an object j of an ∞ -category J . Then $j: * \rightarrow C$ is an initial functor if and only if j is an initial object, and it is a final functor if and only if j is a final (i.e. terminal) object of J .*

Proof. Given an object x of C , unwinding definitions reveals that the relative slice category of the functor $j: * \rightarrow C$ is equivalent to the hom anima $\mathrm{Hom}_C(j, x)$. Since this is already an anima, it is equivalent to its geometric realization, and hence it is weakly contractible if and only if it is contractible. So we see that j is an initial functor if and only if the hom anima $\mathrm{Hom}_C(j, x)$ is contractible for every object x of C , which by is the case by definition precisely if j is an initial object. $\square$

A common way of showing that the relative slice categories $I_{j/}$ or $I_{/j}$ are weakly contractible is by showing they admit an initial or terminal object:

Lemma 21.5.5. *Let C be an ∞ -category that admits an initial (resp. terminal) object $x: * \rightarrow C$. Then C is weakly contractible: $\|C\| \simeq *$.*

Proof. By Remark 21.2.3 we have an adjunction $x: * \rightleftarrows C: p_C$, which by Lemma 21.1.7 induces an equivalence $* \simeq \|*\| \simeq \|C\|$. $\square$

Corollary 21.5.6. *If a functor $\alpha: I \rightarrow J$ admits a left adjoint, then α is a final functor. Dually, if it admits a right adjoint it is an initial functor.*

Proof. We need to show that for every object j the relative slice category $I_{j/}$ is weakly contractible. Let $\beta: J \rightarrow I$ be a left adjoint to α . Then the relative slice category $I_{j/}$ is equivalent to the slice category $I_{\beta(j)/}$ of I . But this admits an initial object given by $(\beta(j), \mathrm{id}_{\beta(j)})$, hence is weakly contractible by Lemma 21.5.5. $\square$

21.6 Filtered and sifted ∞ -categories

We introduce the notions of filtered and sifted ∞ -categories, and record their basic properties.

Definition 21.6.1. An ∞ -category K is called *filtered* if for every finite diagram $F: I \rightarrow K$ in K there exists an object $W \in K$ with a cocone $F \rightarrow \mathrm{const}_W$; equivalently, F extends to a functor $\tilde{F}: I^\triangleright \rightarrow K$.

Example 21.6.2. Every directed poset is a filtered ∞ -category. In particular, the poset $(\mathbb{N}, \leq)$ of natural numbers is filtered.

More generally, any category which is filtered in the classical sense is filtered as an ∞ -category.

The following important characterization of filtered ∞ -categories will be treated as a black box:

Theorem 21.6.3 ([Lur09b, Proposition 5.3.3.3]). *An ∞ -category K is filtered if and only if the functor $\operatorname{colim}_K : \operatorname{Fun}(K, \operatorname{An}) \rightarrow \operatorname{An}$ preserves finite limits.*

Let us list various consequences of this characterization:

Corollary 21.6.4. *A filtered colimit $\operatorname{colim}_K *$ of terminal anima is again terminal.*

Proof. A terminal anima is a limit over the finite category $\emptyset$. □

Corollary 21.6.5. *The functor $\Omega : \operatorname{An}_* \rightarrow \operatorname{An}_*$ preserves filtered colimits.*

Proof. By construction, this functor is given as the composite

$$\operatorname{An}_* \simeq \operatorname{An}_* \times_{\operatorname{An}_*} \operatorname{An}_* \hookrightarrow \operatorname{Fun}([1], \operatorname{An}_*) \times_{\operatorname{An}_*} \operatorname{Fun}([1], \operatorname{An}_*) \simeq \operatorname{Fun}(\sqcup, \operatorname{An}_*) \xrightarrow{\lim} \operatorname{An}_*.$$

The last functor preserves filtered colimits by Proposition 21.6.3. The inclusion $\operatorname{An}_* \simeq (\operatorname{An}_*)_* \rightarrow \operatorname{Fun}([1], \operatorname{An}_*)$ preserves filtered colimits by Corollary 21.6.4. □

Corollary 21.6.6. *For every $n \geq 0$, the functor $\pi_n : \operatorname{An}_* \rightarrow \operatorname{Set}$ preserves filtered colimits.*

Proof. We may factor this functor as follows:

$$\operatorname{An}_* \xrightarrow{\Omega^n} \operatorname{An}_* \xrightarrow{\operatorname{fgt}} \operatorname{An} \xrightarrow{\pi_0} \operatorname{Set}.$$

Since each of these functors preserves filtered colimits, the claim follows. □

Lemma 21.6.7. *Every filtered ∞ -category K is weakly contractible.*

Proof. For every ∞ -category I , its geometric realization may be computed as the colimit in An of the constant I -indexed diagram with value $*$. Indeed, for every anima X there are natural isomorphisms

$$\begin{aligned} \operatorname{Hom}_{\operatorname{An}}(\operatorname{colim}_{i \in I} *, X) &\simeq \operatorname{Nat}(\operatorname{const}_*, \operatorname{const}_X) \\ &\simeq \operatorname{Fun}(I, X) \\ &\simeq \operatorname{Hom}_{\operatorname{An}}(\|I\|, X). \end{aligned}$$

Indeed, such a natural transformation chooses a point of X for every object of I , a path between the corresponding points for every morphism of I , and all the higher coherence data; this is precisely a functor $I \rightarrow X$. The last isomorphism follows from the universal property of the localization $I \rightarrow \|I\|$. The Yoneda lemma therefore gives

$$\|I\| \simeq \operatorname{colim}_{i \in I} *.$$

Applying this to the filtered ∞ -category K , we conclude from Corollary 21.6.4 that

$$\|K\| \simeq \operatorname{colim}_{k \in K} * \simeq *.$$

□

Definition 21.6.8. An ∞ -category I is called *sifted* if it is nonempty and the diagonal functor $\Delta: I \rightarrow I \times I$ is final.

Exercise 21.6.9. The goal of this exercise is to show that the opposite of the simplex category Δ^{op} is sifted, or equivalently that the diagonal functor $\Delta \rightarrow \Delta \times \Delta$ is initial: for $[n], [m] \in \Delta$ the relative slice category $\Delta \times_{\Delta \times \Delta} (\Delta \times \Delta)_{/([n],[m])}$ is weakly contractible.

- (1) Show that this relative slice category is equivalent to the full subcategory $\Delta_{/[n] \times [m]}$ of $\text{Poset}_{/[n] \times [m]}$ consisting of morphisms of posets of the form $\varphi: [k] \rightarrow [n] \times [m]$.
- (2) Consider the subcategory $C_{n,m} \subseteq \Delta_{/[n] \times [m]}$ consisting of all objects, but only those morphisms $\psi: [k] \rightarrow [l]$ over $[n] \times [m]$ satisfying $\psi(k) = l$. Construct two functors

$$f: C_{n,m} \hookrightarrow \Delta_{/[n] \times [m]} \quad \text{and} \quad g: \Delta_{/[n] \times [m]} \hookrightarrow C_{n,m}$$

together with natural transformations $fg \rightarrow \text{id}$ and $gf \rightarrow \text{id}$. Conclude that f and g induce equivalences on geometric realizations.

- (3) Construct two functors

$$h: C_{n,m} \rightarrow [n] \times [m] \quad \text{and} \quad k: [n] \times [m] \rightarrow C_{n,m},$$

together with an identification $hk = \text{id}$ and a natural transformation $kh \rightarrow \text{id}$. Conclude that h and k induce equivalences on geometric realizations.

- (4) Show that $\|C \times D\| \simeq \|C\| \times \|D\|$ for ∞ -categories C and D ;
- (5) Show that $[n]$ and $[m]$ are both weakly contractible.
- (6) Combine these points to conclude that Δ^{op} is sifted.

Lemma 21.6.10. *Every filtered ∞ -category is sifted.*

Proof. Let K be a filtered ∞ -category. We need to show that for every object $(k, l) \in K \times K$ the relative slice category $K \times_{K \times K} (K \times K)_{(k,l)}/$ is weakly contractible. We will show that this relative slice category is in fact a filtered ∞ -category, so that Lemma 21.6.7 implies the claim. To this end, consider a finite ∞ -category I and a functor $F: I \rightarrow K \times_{K \times K} (K \times K)_{(k,l)}/$. We may equivalently encode this diagram as a functor $P \rightarrow I$, where P is defined as the following pushout of ∞ -categories:

$$\begin{array}{ccc} I \sqcup I & \hookrightarrow & (I \sqcup I)^{\triangleleft} \\ \downarrow \nabla & & \downarrow \\ I & \xrightarrow{\quad r \quad} & P. \end{array}$$

Note that P is again finite, so that by filteredness of K we obtain an extension $P^{\triangleright} \rightarrow K$. This extension then translates back into an extension $\tilde{F}$ of F , finishing the proof. $\square$

Lemma 21.6.11. *Let C be an ∞ -category that admits finite products, and assume that for every object $X \in C$ the functor $X \times -: C \rightarrow C$ preserves colimits. Then the product functor $- \times -: C \times C \rightarrow C$ preserves sifted colimits.*

Proof. Let I be a sifted ∞ -category and let $X, Y: I \rightarrow C$ be two functors. We need to show that the map

$$\operatorname{colim}_{i \in I} (X_i \times Y_i) \rightarrow (\operatorname{colim}_{i \in I} X_i) \times (\operatorname{colim}_{i \in I} Y_i)$$

is an equivalence. This follows from the following computation:

$$\begin{aligned} \operatorname{colim}_{i \in I} (X_i \times Y_i) &\stackrel{(1)}{\simeq} \operatorname{colim}_{(i,j) \in I \times I} (X_i \times Y_j) \\ &\stackrel{(2)}{\simeq} \operatorname{colim}_{i \in I} \operatorname{colim}_{j \in I} (X_i \times Y_j) \\ &\stackrel{(3)}{\simeq} \operatorname{colim}_{i \in I} (X_i \times \operatorname{colim}_{j \in I} Y_j) \\ &\stackrel{(4)}{\simeq} (\operatorname{colim}_{i \in I} X_i) \times (\operatorname{colim}_{j \in I} Y_j). \end{aligned}$$

Here equivalence (1) holds because the diagonal functor $I \rightarrow I \times I, i \mapsto (i, i)$ is final by assumption on I . The equivalence (2) is an instance of Lemma 21.2.10. The equivalences (3) and (4) hold because the product in C preserves colimits in each variable. $\square$

Lemma 21.6.12. *The geometric realization functor $|-|: \mathbf{sAn} \rightarrow \mathbf{An}$ preserves finite products.*

Proof. By definition, this functor is $\operatorname{colim}_{[n] \in \Delta^{\text{op}}}: \mathbf{Fun}(\Delta^{\text{op}}, \mathbf{An}) \rightarrow \mathbf{An}$, and the claim is equivalent to the statement that the product functor $- \times -: \mathbf{An} \times \mathbf{An} \rightarrow \mathbf{An}$ preserves Δ^{op} -indexed colimits. The ∞ -category Δ^{op} is sifted by Exercise 21.6.9. Furthermore, for an anima X the functor $X \times -: \mathbf{An} \rightarrow \mathbf{An}$ admits a right adjoint $\operatorname{Hom}_{\mathbf{An}}(X, -): \mathbf{An} \rightarrow \mathbf{An}$, hence preserves all colimits. This verifies the assumptions of Lemma 21.6.11, proving the claim. $\square$

21.7 A comparison theorem

The argument below is the monad-free comparison mechanism underlying the Barr–Beck–Lurie monadicity theorem. Suppose that two categories of structured objects come with forgetful functors to the same ∞ -category. To compare them, it is often enough to compare their free objects: every object can be reconstructed from free ones by a canonical simplicial resolution. We isolate this mechanism here without introducing the structured objects classified by monads.

We first recall the class of simplicial objects for which the relevant geometric realizations exist formally. Let $\Delta_{-\infty}$ be the category whose objects are the symbols $[n]$ for $n \geq -1$ and whose morphisms $[m] \rightarrow [n]$ are the order-preserving maps

$$\{-\infty\} \sqcup [m] \longrightarrow \{-\infty\} \sqcup [n]$$

which preserve $-\infty$. There are inclusions $\Delta \subseteq \Delta_+ \subseteq \Delta_{-\infty}$, where a morphism belongs to Δ_+ precisely when the inverse image of $-\infty$ is $\{-\infty\}$.

Definition 21.7.1 ([Lur17, Definition 4.7.2.2]). An augmented simplicial object $X: \Delta_+^{\text{op}} \rightarrow C$ is *split* if it extends to a functor $\Delta_{-\infty}^{\text{op}} \rightarrow C$. A simplicial object is *split* if it extends to a split augmented simplicial object.

Let $G: D \rightarrow C$ be a functor. A simplicial object $X_\bullet: \Delta^{\text{op}} \rightarrow D$ is *G-split* if $GX_\bullet$ is split in C .

Lemma 21.7.2 (Absolute realization of split simplicial objects). *Every split augmented simplicial object is a colimit diagram. In particular, every split simplicial object admits a geometric realization, and every functor preserves this realization.*

Proof. Reversing the order on the finite ordered sets identifies an extension to $\Delta_{-\infty}^{\text{op}}$ with the extra-degeneracy structure of Definition 5.2.3. The first claim is therefore Lemma 5.2.4. Applying any functor preserves the extension to $\Delta_{-\infty}^{\text{op}}$, so the same lemma computes the image of the colimit cocone. Thus the colimit is absolute. $\square$

The following theorem is the monad-free form of [Lur17, Corollary 4.7.3.16]. The bar-resolution argument in its proof is also the engine of the proof of the ∞ -categorical Barr–Beck theorem [Lur17, Theorem 4.7.3.5, Proposition 4.7.3.14].

Theorem 21.7.3 (Comparison). *Let $G : D \rightarrow C$ and $G' : D' \rightarrow C$ be functors admitting left adjoints F and F' , respectively. Let $K : D \rightarrow D'$ be equipped with a natural isomorphism*

$$\theta : G'K \xrightarrow{\sim} G$$

whose mate $\theta^b : F' \rightarrow KF$ is an isomorphism. Assume that:

- (1) *The functors G and G' are conservative;*
- (2) *The ∞ -category D admits geometric realizations of G -split simplicial objects, and G preserves them;*
- (3) *The ∞ -category D' admits geometric realizations of G' -split simplicial objects, and G' preserves them.*

Then K is an equivalence of ∞ -categories.

The strategy is to resolve every object functorially by objects in the image of the relevant left adjoint. The mate condition identifies the two classes of free objects, while the hypotheses on split simplicial objects ensure that both categories can reconstruct arbitrary objects from these resolutions.

Proof. Write $\eta : \text{id}_C \rightarrow GF$ and $\varepsilon : FG \rightarrow \text{id}_D$ for the unit and counit of the first adjunction. For $d \in D$, its *bar resolution* is the augmented simplicial object

$$B_{\bullet}(d) \longrightarrow d, \quad B_n(d) := (FG)^{n+1}d.$$

Its face maps are obtained by applying the counit ε in one of the $n + 1$ occurrences of FG , and its degeneracy maps by inserting $F\eta G$ between two consecutive occurrences. Naturality and the triangle identities give the simplicial identities. After applying G , the augmented simplicial object has terms

$$GB_n(d) \simeq (GF)^{n+1}Gd.$$

The unit η supplies the extra degeneracies, by inserting a new initial occurrence of GF . Hence $GB_{\bullet}(d) \rightarrow Gd$ is split. By assumption, $B_{\bullet}(d)$ admits a geometric realization in D , and the augmentation induces a morphism

$$|B_{\bullet}(d)| \longrightarrow d.$$

After applying G , this becomes the colimit map of the split augmented simplicial object $GB_\bullet(d) \rightarrow Gd$, which is an isomorphism by Lemma 21.7.2. Since G is conservative, the displayed morphism is an isomorphism. Thus every object of D is the geometric realization of a simplicial object whose terms are free objects Fc .

We next observe that K preserves the geometric realizations under consideration. Let $X_\bullet$ be a G -split simplicial object of D . The isomorphism $G'KX_\bullet \simeq GX_\bullet$ shows that $KX_\bullet$ is G' -split, so both geometric realizations exist. Apply G' to the canonical comparison morphism

$$|KX_\bullet| \longrightarrow K|X_\bullet|.$$

Using the preservation hypotheses and $G'K \simeq G$, its image identifies with the identity of $|GX_\bullet|$. It is therefore an isomorphism. Conservativity of G' now shows that the comparison morphism itself is an isomorphism.

We prove that K is fully faithful. For $c \in C$ and $d \in D$, adjunction and the mate isomorphism give natural isomorphisms

$$\begin{aligned} \mathrm{Hom}_D(Fc, d) &\simeq \mathrm{Hom}_C(c, Gd) \\ &\simeq \mathrm{Hom}_C(c, G'Kd) \\ &\simeq \mathrm{Hom}_{D'}(F'c, Kd) \\ &\simeq \mathrm{Hom}_{D'}(KFc, Kd). \end{aligned}$$

By the definition of the mate, their composite is the map induced by K . Since every $B_n(d)$ is free, resolving the source and using preservation of the realization gives, for all $d, e \in D$,

$$\begin{aligned} \mathrm{Hom}_D(d, e) &\simeq \lim_{[n] \in \Delta} \mathrm{Hom}_D(B_n(d), e) \\ &\simeq \lim_{[n] \in \Delta} \mathrm{Hom}_{D'}(KB_n(d), Ke) \\ &\simeq \mathrm{Hom}_{D'}(Kd, Ke). \end{aligned}$$

Thus K is fully faithful.

It remains to prove essential surjectivity. Form the analogous bar resolution $B'_\bullet(d') \rightarrow d'$ for an object $d' \in D'$. Every term has the form $F'c$ and is therefore in the essential image of K , because $F' \simeq KF$. Since K is fully faithful, the entire simplicial diagram lifts to a simplicial object $Y_\bullet$ of D with $KY_\bullet \simeq B'_\bullet(d')$. The simplicial object $Y_\bullet$ is G -split, since

$$GY_\bullet \simeq G'KY_\bullet \simeq G'B'_\bullet(d')$$

is split. Consequently, $Y_\bullet$ admits a geometric realization and K preserves it. Applying the bar-resolution computation in D' gives

$$K|Y_\bullet| \simeq |KY_\bullet| \simeq |B'_\bullet(d')| \simeq d'.$$

Hence every object of D' lies in the essential image of K . □

21.8 Bousfield localizations

We introduce the notion of a Bousfield localization, and prove a useful characterization of them in Proposition 21.8.9.

Definition 21.8.1. A functor $L: C \rightarrow D$ is called a *Bousfield localization* if it admits a fully faithful right adjoint $R: D \hookrightarrow C$. In this case, we also call D a *reflective subcategory of C* .

A *Bousfield colocalization* is instead a functor $C \rightarrow D$ that admits a fully faithful *left* adjoint.

Example 21.8.2 (Rationalization). The inclusion $\text{Vect}_{\mathbb{Q}} \hookrightarrow \text{Ab}$ of rational vector spaces into abelian groups admits a left adjoint

$$\mathbb{Q} \otimes_{\mathbb{Z}} -: \text{Ab} \rightarrow \text{Vect}_{\mathbb{Q}},$$

called the *rationalization functor*. It is a Bousfield localization.

Example 21.8.3 (Abelianization). The inclusion $\text{Ab} \hookrightarrow \text{Grp}$ of abelian groups into all groups admits a left adjoint

$$(-)^{\text{ab}}: \text{Grp} \rightarrow \text{Ab},$$

given by the *abelianization functor*.

Example 21.8.4 (Group completion). The inclusion $\text{Grp} \hookrightarrow \text{Mon}$ of groups into monoids admits a left adjoint

$$(-)^{\text{grp}}: \text{Mon} \rightarrow \text{Grp}$$

called the (discrete) *group completion functor*.

Example 21.8.5 (Path components). The inclusion $\text{Set} \hookrightarrow \text{An}$ admits a left adjoint

$$\pi_0: \text{An} \rightarrow \text{Set}$$

given by the set of path components.

Example 21.8.6 (Geometric realization). The inclusion $\text{An} \hookrightarrow \text{Cat}_{\infty}$ has a left adjoint

$$\| - \|: \text{Cat}_{\infty} \rightarrow \text{An}$$

given by the geometric realization functor. It also admits a right adjoint

$$(-)^{\simeq}: \text{Cat}_{\infty} \rightarrow \text{An}$$

given by the groupoid core functor.

Example 21.8.7 (Homotopy category). The inclusion $\text{Cat} \hookrightarrow \text{Cat}_{\infty}$ has a left adjoint

$$\text{Ho}(-): \text{Cat}_{\infty} \rightarrow \text{Cat}_1$$

given by the homotopy category functor.

The terminology is justified by the fact that every Bousfield (co)localization is a localization in the sense of Definition 1.5.19:

Lemma 21.8.8. *Let $L: E \rightarrow C$ be a Bousfield localization, and let $W := L^{-1}(C^{\simeq})$ be the class of morphisms in E inverted by L . Then L exhibits C as a localization of E at W . The same conclusion holds if L is a Bousfield colocalization.*

Proof. The adjunction $L \vdash R$ induces for any ∞ -category D an adjunction

$$R^*: \text{Fun}(C, D) \rightleftarrows \text{Fun}(E, D) : L^*$$

on functor categories. The counit $LR \xrightarrow{\sim} \text{id}$ induces the counit $R^*L^* = (LR)^* \xrightarrow{\sim} \text{id}^* = \text{id}$, hence L^* is fully faithful. To identify the essential image of L^* , note that a functor $F: E \rightarrow D$ is in the essential image if and only if the unit $F \rightarrow L^*R^*F = F \circ R \circ L$ is a natural isomorphism. This happens if and only if F sends the unit $\eta_X: X \rightarrow R(L(X))$ in E to an isomorphism in D for every $X \in E$. Since $L(\eta_X): L(X) \rightarrow L(R(L(X)))$ is an isomorphism, this is clearly implied by F inverting the maps in $W = L^{-1}(C^\simeq)$. Conversely, if F inverts each of the maps η_X , then for every morphism $f: X \rightarrow X'$ in $W = L^{-1}(C^\simeq)$ it inverts three out of the four maps in the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & X' \\ \eta_X \downarrow & & \downarrow \eta_{X'} \\ RLX & \xrightarrow{LRf} & RLX', \end{array}$$

and so by 2-out-of-3 it also inverts f . This shows that the essential image of L^* consists of the functors F that invert W , as desired.

The proof for Bousfield colocalizations is dual. □

Here is a useful criterion for checking that a full subcategory is a reflective subcategory:

Proposition 21.8.9 (cf. [Lur09b, Proposition 5.2.7.4]). *Let $L: C \rightarrow C$ be a functor equipped with a natural transformation $\eta: \text{id}_C \Rightarrow L$ such that both maps*

$$\eta_{Lx}: Lx \rightarrow LLx \quad \text{and} \quad L\eta_x: Lx \rightarrow LLx$$

are isomorphisms for all $x \in C$. Then the functor $L: C \rightarrow \text{im}(L)$ is left adjoint to the inclusion $\text{im}(L) \hookrightarrow C$, with unit η . In particular, $L: C \rightarrow \text{im}(L)$ is a Bousfield localisation.

Proof. We must show that for objects x and y in C , precomposition with $\eta_x: x \rightarrow Lx$ induces an equivalence of anima

$$- \circ \eta_x: \text{Hom}_C(Lx, Ly) \xrightarrow{\sim} \text{Hom}_C(x, Ly).$$

We will do this by showing separately that it admits a left inverse and a right inverse.

We start by constructing a left inverse. By assumption on η , the map $\eta_{Ly}: Ly \rightarrow LLy$ is an isomorphism for every y , hence we may form the following composite:

$$\text{Hom}_C(x, Ly) \xrightarrow{L} \text{Hom}_C(Lx, LLy) \xrightarrow{\eta_{Ly}^{-1} \circ -} \text{Hom}_C(Lx, Ly).$$

The fact this is a left inverse follows from the following commutative diagram:

$$\begin{array}{ccccc} \text{Hom}_C(x, Ly) & \xrightarrow{L} & \text{Hom}_C(Lx, LLy) & & \\ & \searrow \eta_{Ly} \circ - & \swarrow - \circ \eta_x & \searrow \eta_{Ly}^{-1} \circ - & \\ & \text{Hom}_C(x, LLy) & & \text{Hom}_C(Lx, Ly) & \\ & \searrow \eta_{Ly}^{-1} \circ - & & \swarrow - \circ \eta_x & \\ & & \text{Hom}_C(x, Ly) & & \end{array}$$

Here the upper left triangle commutes since for any morphism $f: x \rightarrow Ly$ the transformation η provides a naturality square

$$\begin{array}{ccc} x & \xrightarrow{f} & Ly \\ \eta_x \downarrow & & \downarrow \eta_{Ly} \\ Lx & \xrightarrow{Lf} & LLy \end{array}$$

which is fully functorial in $f \in \text{Hom}_C(x, Ly)$.

We now turn to the construction of a right inverse. Using the assumption that the map $L\eta_y: Ly \rightarrow LLy$ is an isomorphism, we may consider the following composite:

$$\text{Hom}_C(x, Ly) \xrightarrow{L} \text{Hom}_C(Lx, LLy) \xrightarrow{(L\eta_y)^{-1} \circ -} \text{Hom}_C(Lx, Ly).$$

This is a right inverse by the following commutative diagram:

$$\begin{array}{ccccc} & & \text{Hom}_C(x, Ly) & & \\ & \nearrow^{-\circ \eta_x} & & \searrow^L & \\ \text{Hom}_C(Lx, Ly) & \xrightarrow{L} & \text{Hom}_C(LLx, LLy) & \xrightarrow{-\circ L\eta_x} & \text{Hom}_C(Lx, LLy) \\ & \searrow & \nearrow^{L\eta_y \circ -} & & \downarrow (L\eta_y)^{-1} \circ - \\ & & \text{Hom}_C(Lx, Ly) & \xlongequal{\quad} & \text{Hom}_C(Lx, Ly) \end{array}$$

The top triangle commutes since the functor L preserves composition. The triangle in the middle commutes since for any morphism $g: Lx \rightarrow Ly$, the transformation η provides a naturality square

$$\begin{array}{ccc} Lx & \xrightarrow{g} & Ly \\ L(\eta_x) \downarrow & & \downarrow L(\eta_y) \\ LLx & \xrightarrow{Lg} & LLy \end{array}$$

which is fully functorial in $g \in \text{Hom}_C(Lx, Ly)$. The bottom right triangle clearly commutes. This finishes the proof. $\square$

Exercises

22 Presentable ∞ -categories

The large ∞ -categories that occur in stable homotopy theory are usually built from small data. Presheaf categories are the basic examples: they are large enough to admit all small colimits, but every presheaf is assembled from the small collection of representable presheaves. Presentable ∞ -categories are obtained from presheaf categories by imposing a small collection of relations.

The usual treatment of presentability uses regular cardinals, accessible categories, and accessible functors. We will not develop that language here. Instead, we take the presentation by generators and relations as our definition. We prove the elementary localization and compactness arguments that fit naturally with this presentation and record the remaining consequences needed elsewhere in the book as black boxes. This is equivalent to the standard definition; see [Lur09b, Theorem 5.5.1.1 and Proposition 5.5.4.15]. We freely use the Yoneda lemma and the elementary presheaf formalism from Section 1.8.

22.1 Presheaves and free cocompletions

The free-cocompletion theorem from Theorem 1.8.9 has a useful variant in which one freely adds only a prescribed collection of colimits.

Definition 22.1.1. Let $\mathcal{K}$ be a collection of small ∞ -categories and let C be a small ∞ -category. We denote by

$$\mathrm{PSh}^{\mathcal{K}}(C) \subseteq \mathrm{PSh}(C)$$

the smallest full subcategory of $\mathrm{PSh}(C)$ that contains the representable presheaves and is closed under I -indexed colimits for all $I \in \mathcal{K}$.

Proposition 22.1.2 (Lurie [Lur09b, Proposition 5.3.6.2]). *Let $\mathcal{K}$ be a collection of small ∞ -categories. For every small ∞ -category C and every (possibly large) ∞ -category D with I -indexed colimits for all $I \in \mathcal{K}$, restriction along the Yoneda embedding $Y: C \hookrightarrow \mathrm{PSh}^{\mathcal{K}}(C)$ induces an equivalence*

$$\mathrm{Fun}_{\mathcal{K}}(\mathrm{PSh}^{\mathcal{K}}(C), D) \xrightarrow{\sim} \mathrm{Fun}(C, D),$$

where the left-hand side denotes the full subcategory spanned by the functors $\mathrm{PSh}^{\mathcal{K}}(C) \rightarrow D$ that preserve I -indexed colimits for all $I \in \mathcal{K}$.

22.2 Presentable categories and adjoint functors

We first make precise what it means to impose a small collection of relations on a presheaf category. Let E be an ∞ -category and let S be a collection of morphisms in E . An object $X \in E$ is called *S -local* if for every morphism $f: A \rightarrow B$ in S , precomposition with f induces an equivalence

$$\mathrm{Hom}_E(B, X) \xrightarrow{\sim} \mathrm{Hom}_E(A, X).$$

We write $E^{S\text{-loc}} \subseteq E$ for the full subcategory of S -local objects.

Definition 22.2.1. An ∞ -category D is called *presentable* if there exist a small ∞ -category C and a small collection S of morphisms in $\mathrm{PSh}(C)$ such that the inclusion

$$\mathrm{PSh}(C)^{S\text{-loc}} \hookrightarrow \mathrm{PSh}(C)$$

admits a left adjoint and D is equivalent to $\mathrm{PSh}(C)^{S\text{-loc}}$.

Thus a presentation of D consists of a small collection of generators, encoded by C , and a small collection of relations, encoded by S . The presentation is not part of the structure on D , and is very far from unique. The existence of the reflector may in fact be omitted from the definition: any small collection of morphisms in a presheaf category generates a Bousfield localization. We include it to keep the definition directly tied to the localization formalism of Section 21.8.

Theorem 22.2.2 (Basic constructions on presentable categories, [Lur09b, Theorem 5.5.1.1 and Propositions 5.5.3.6, 5.5.3.10–5.5.3.11 and 5.5.4.15]). *The following statements hold.*

- (1) *For every small ∞ -category C , the presheaf category $\mathrm{PSh}(C)$ is presentable. In particular, $\mathrm{An} \simeq \mathrm{PSh}(*)$ is presentable.*
- (2) *If C is presentable and K is small, then the functor category $\mathrm{Fun}(K, C)$ is presentable.*
- (3) *If C is presentable and $X \in C$, then the slice categories $C_{/X}$ and $C_{X/}$ are presentable.*
- (4) *If C is presentable and S is a small collection of morphisms in C , then the full subcategory $C^{S\text{-loc}}$ is presentable and its inclusion into C admits a left adjoint.*

We will also use the following immediate consequence of the definition: every presentable ∞ -category C admits a small collection $\mathcal{G}$ of objects such that the functors $\mathrm{Hom}_C(G, -)$, for $G \in \mathcal{G}$, jointly detect isomorphisms. Indeed, in any presentation of C as a localization of a presheaf category, one may take the images under the reflector of the representable presheaves. The adjunction and the Yoneda lemma then reduce detection to the fact that isomorphisms of presheaves are detected objectwise.

We write Pr^{L} for the ∞ -category whose objects are presentable ∞ -categories and whose morphisms are the colimit-preserving functors. By the adjoint functor theorem below, these are precisely the left adjoint functors. Similarly, we write Pr^{R} for the ∞ -category of presentable ∞ -categories and right adjoint functors. Passing between left and right adjoints gives an equivalence

$$(\mathrm{Pr}^{\mathrm{L}})^{\mathrm{op}} \simeq \mathrm{Pr}^{\mathrm{R}}.$$

Their objects are large ∞ -categories, so all statements about Pr^{L} and Pr^{R} are interpreted one universe higher, in the enlarged universe $\widehat{\mathrm{Cat}}_{\infty}$ from Remark 1.8.1.

Theorem 22.2.3 (Limits of presentable categories, [Lur09b, Proposition 5.5.3.13 and Theorem 5.5.3.18]). *The ∞ -categories Pr^{L} and Pr^{R} admit small limits. Moreover, the inclusion functors*

$$\mathrm{Pr}^{\mathrm{L}} \longrightarrow \widehat{\mathrm{Cat}}_{\infty} \quad \text{and} \quad \mathrm{Pr}^{\mathrm{R}} \longrightarrow \widehat{\mathrm{Cat}}_{\infty}$$

preserve small limits.

Corollary 22.2.4 (Presentability of kernels). *Let $F: C \rightarrow D$ be a colimit-preserving functor between pointed presentable ∞ -categories. Then the full subcategory*

$$\ker(F) := \{X \in C \mid F(X) \simeq 0\} \subseteq C$$

is presentable, and its inclusion into C preserves small colimits.

Proof. The functor $* \rightarrow D$ selecting the zero object is a left adjoint, since the zero object is initial. The pullback

$$\begin{array}{ccc} \ker(F) & \longrightarrow & C \\ \downarrow & & \downarrow F \\ * & \xrightarrow{0} & D \end{array}$$

may therefore be formed in Pr^{L} . By Theorem 22.2.3, its underlying ∞ -category is the pullback in $\widehat{\mathrm{Cat}}_{\infty}$, which identifies it with the displayed full subcategory. The projection $\ker(F) \rightarrow C$ is a morphism in Pr^{L} , so it preserves small colimits. $\square$

The main reason presentable categories are useful is that functors out of them have adjoints as soon as they preserve colimits. This is the form of the adjoint functor theorem used throughout the book.

Theorem 22.2.5 (Adjoint functor theorem, [Lur09b, Corollary 5.5.2.9]). *Let C and D be presentable ∞ -categories. A functor $F: C \rightarrow D$ admits a right adjoint if and only if it preserves small colimits.*

There is a dual-looking criterion for the existence of a left adjoint, but preservation of limits by itself is not enough: one must also impose a size condition on the functor. We will not need its general formulation. The special cases involving structured objects that occur later are included in Proposition 22.2.7 below.

Corollary 22.2.6 (Representability, [Lur09b, Proposition 5.5.2.2]). *Let C be a presentable ∞ -category. A functor $F: C^{\mathrm{op}} \rightarrow \mathbf{An}$ is representable if and only if it preserves small limits.*

Proposition 22.2.7 (Presentability of structured objects). *Let C be a presentable ∞ -category.*

(1) *The ∞ -categories $\mathrm{CMon}(C)$ and $\mathrm{CGrp}(C)$ are presentable, and the inclusions*

$$\mathrm{CGrp}(C) \hookrightarrow \mathrm{CMon}(C) \hookrightarrow \mathrm{Fun}(\mathrm{Span}(\mathrm{Fin}), C)$$

admit left adjoints.

(2) *If C is stable and $\mathcal{G}$ is a small collection of objects of C , then the smallest full stable subcategory $\mathrm{Loc}(\mathcal{G}) \subseteq C$ that contains $\mathcal{G}$ and is closed under small colimits is presentable. Its inclusion into C preserves small colimits and consequently admits a right adjoint.*

Proof. For (1), consider the functor category $\text{Fun}(\text{Span}(\text{Fin}), C)$, which is presentable by Theorem 22.2.2. Let $\mathcal{G}$ be a small collection of objects of C whose corepresentable functors jointly detect isomorphisms. For every finite set S , evaluation at S admits a left adjoint

$$\gamma_S: C \rightarrow \text{Fun}(\text{Span}(\text{Fin}), C),$$

given by left Kan extension along the inclusion $S: * \rightarrow \text{Span}(\text{Fin})$. Consider now the small collection of morphisms in $\text{Fun}(\text{Span}(\text{Fin}), C)$ consisting, for every object $G \in \mathcal{G}$ and all $S, T \in \text{Fin}$, of the maps

$$\emptyset \longrightarrow \gamma_\emptyset(G) \quad \text{and} \quad \gamma_S(G) \sqcup \gamma_T(G) \longrightarrow \gamma_{S \sqcup T}(G).$$

The second morphism is induced by the two product projections $S \sqcup T \rightarrow S, T$ in $\text{Span}(\text{Fin})$. By the adjunctions defining the functors γ_S , a functor $F \in \text{Fun}(\text{Span}(\text{Fin}), C)$ is local with respect to these morphisms precisely when the maps

$$\text{Hom}_C(G, F(\emptyset)) \longrightarrow *$$

and

$$\text{Hom}_C(G, F(S \sqcup T)) \longrightarrow \text{Hom}_C(G, F(S)) \times \text{Hom}_C(G, F(T))$$

are equivalences for every $G \in \mathcal{G}$ and all $S, T \in \text{Fin}$. Since these objects jointly detect isomorphisms, this says exactly that F preserves finite products. Thus the local objects are precisely the objects of $\text{CMon}(C)$. It follows from Theorem 22.2.2(4) that $\text{CMon}(C)$ is presentable and that its inclusion into $\text{Fun}(\text{Span}(\text{Fin}), C)$ admits a left adjoint.

Let $\sigma: \langle 2 \rangle \rightarrow \langle 2 \rangle$ be the morphism in $\text{Span}(\text{Fin})$ represented by the shear matrix

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

For every G in the detecting collection, it induces an endomorphism

$$\sigma^*: \gamma_{\langle 2 \rangle}(G) \longrightarrow \gamma_{\langle 2 \rangle}(G).$$

Locality with respect to these endomorphisms says that $F(\sigma)$ is an isomorphism. For a product-preserving functor F , the map $F(\sigma)$ is the shear map of the commutative monoid F . Consequently, adjoining the morphisms σ^* to the preceding localizing collection cuts out precisely $\text{CGrp}(C) \subseteq \text{Fun}(\text{Span}(\text{Fin}), C)$. This proves that $\text{CGrp}(C)$ is presentable and reflective in $\text{Fun}(\text{Span}(\text{Fin}), C)$. The reflector

$$\text{Fun}(\text{Span}(\text{Fin}), C) \longrightarrow \text{CGrp}(C)$$

restricts to a left adjoint to the inclusion $\text{CGrp}(C) \hookrightarrow \text{CMon}(C)$.

For (2), apply [Lur17, Proposition 1.4.4.11(2)] to the small collection of all shifts $G[n]$ with $G \in \mathcal{G}$ and $n \in \mathbb{Z}$. In a stable ∞ -category, the smallest full subcategory containing these objects and closed under extensions and small colimits is precisely $\text{Loc}(\mathcal{G})$. It is therefore presentable. Its inclusion into C preserves small colimits by definition and hence admits a right adjoint by Theorem 22.2.5. $\square$

22.3 Compact objects and Ind-categories

We now specialize to finiteness controlled by ordinary filtered colimits. This is sufficient for all applications in the book, so we will not need the more general notion of κ -compactness.

Definition 22.3.1 (Compact object). Let C be an ∞ -category with filtered colimits. An object $X \in C$ is called *compact* if the functor $\mathrm{Hom}_C(X, -): C \rightarrow \mathbf{An}$ preserves filtered colimits. We write $C^\omega \subseteq C$ for the full subcategory of compact objects.

Corollary 22.3.2. *If C is a stable ∞ -category with filtered colimits, then C^ω is a thick subcategory of C .*

Definition 22.3.3. An ∞ -category C is called *compactly generated* if it admits small colimits, the subcategory C^ω is small, and every object of C is a filtered colimit of compact objects.

If C is stable, a small collection $\mathcal{G}$ of compact objects is called a *collection of compact generators* if the smallest localizing subcategory of C containing $\mathcal{G}$ is C itself. Here a localizing subcategory is a full stable subcategory closed under small colimits.

Proposition 22.3.4 (Compact generation criterion, cf. Hovey, Palmieri, and Strickland [HPS97, Corollary 2.3.12]). *Let C be a stable ∞ -category with small colimits and let $\mathcal{G}$ be a small collection of compact objects. The following conditions are equivalent:*

(1) *The collection $\mathcal{G}$ generates C .*

(2) *The functors*

$$\mathrm{Hom}_C(G[n], -): C \rightarrow \mathbf{An}, \quad G \in \mathcal{G}, n \in \mathbb{Z},$$

are jointly conservative.

(3) *Every object of C is a filtered colimit of objects in the smallest thick subcategory of C containing $\mathcal{G}$.*

If these conditions hold, then C is compactly generated and its compact objects form the smallest thick subcategory of C containing $\mathcal{G}$.

Proof. Let $C^{\mathcal{G}\text{-fin}} \subseteq C$ denote the smallest thick subcategory containing $\mathcal{G}$, and call its objects $\mathcal{G}$ -finite. Since $\mathcal{G}$ is small, the subcategory $C^{\mathcal{G}\text{-fin}}$ is small: it is obtained from $\mathcal{G}$ by iterating shifts, finite colimits, and retracts. Moreover, every $\mathcal{G}$ -finite object is compact by Corollary 22.3.2.

Condition (3) implies condition (1), since every localizing subcategory containing $\mathcal{G}$ contains all $\mathcal{G}$ -finite objects and their filtered colimits. We next prove that condition (1) implies condition (2). Since C is stable, it suffices to show that the displayed functors jointly detect zero objects. Suppose that $X \in C$ satisfies

$$\mathrm{Hom}_C(G[n], X) \simeq *$$

for every $G \in \mathcal{G}$ and $n \in \mathbb{Z}$. The full subcategory of objects $Y \in C$ satisfying $\mathrm{Hom}_C(Y[n], X) \simeq *$ for every $n \in \mathbb{Z}$ is localizing and contains $\mathcal{G}$, hence is all of C . Taking $Y = X$ shows that the identity of X is nullhomotopic, so $X \simeq 0$.

It remains to prove that condition (2) implies condition (3). Fix an object $X \in C$, and let $I_X \subseteq C/X$ be the full subcategory spanned by the morphisms $Y \rightarrow X$ for which Y is $\mathcal{G}$ -finite. This is a small

∞ -category, and it is closed under finite colimits in $C_{/X}$. It is therefore filtered. The tautological maps $Y \rightarrow X$ induce a morphism

$$\varphi: \operatorname{colim}_{(Y \rightarrow X) \in I_X} Y \longrightarrow X.$$

We claim that φ is an isomorphism. By condition (2), it is enough to show that mapping out of every shift of every object of $\mathcal{G}$ sends φ to an equivalence. We will prove the stronger statement that

$$\varphi \circ -: \operatorname{Hom}_C(Y', \operatorname{colim}_{(Y \rightarrow X) \in I_X} Y) \longrightarrow \operatorname{Hom}_C(Y', X)$$

is an equivalence for every $\mathcal{G}$ -finite object Y' .

Since Y' is compact, the source is the filtered colimit of the anima $\operatorname{Hom}_C(Y', Y)$. Both sides are grouplike E_∞ -anima under addition of morphisms, and the displayed map preserves this structure. Translation therefore identifies the homotopy groups at every component with those based at the zero morphism, so it suffices to show that the induced map is an isomorphism on π_0 and on all homotopy groups based at zero. By stability, we have

$$\pi_k \operatorname{Hom}_C(Y', Z) \cong \pi_0 \operatorname{Hom}_C(Y'[k], Z)$$

for every $k \geq 0$, and $Y'[k]$ is again $\mathcal{G}$ -finite. We may therefore restrict attention to π_0 . We must show that the map

$$\operatorname{colim}_{(Y \rightarrow X) \in I_X} \pi_0 \operatorname{Hom}_C(Y', Y) \longrightarrow \pi_0 \operatorname{Hom}_C(Y', X)$$

is an isomorphism.

Surjectivity follows directly from the definition of I_X . Indeed, a morphism $f: Y' \rightarrow X$ defines an object (Y', f) of I_X , and the class of f is represented at this stage by $\operatorname{id}_{Y'}$. For injectivity, suppose that a class in the kernel is represented at a stage (Y, f) by a morphism $g: Y' \rightarrow Y$. Thus the composite $fg: Y' \rightarrow X$ is nullhomotopic. After choosing such a nullhomotopy, f factors through the cofiber $Y'' := \operatorname{cofib}(g)$, giving a morphism $h: Y'' \rightarrow X$. The object Y'' is $\mathcal{G}$ -finite, so (Y'', h) belongs to I_X . At this stage the class of g becomes zero, since the composite $Y' \rightarrow Y \rightarrow Y''$ is nullhomotopic. This proves injectivity, and hence proves that φ is an isomorphism. Condition (3) follows.

We have already seen that every $\mathcal{G}$ -finite object is compact. Conversely, if X is compact, then the preceding filtered-colimit presentation of X shows that the identity of X factors through some $\mathcal{G}$ -finite object $Y \rightarrow X$ in I_X . Thus X is a retract of Y , and is therefore $\mathcal{G}$ -finite. Consequently,

$$C^\omega = C^{\mathcal{G}\text{-fin}}.$$

The right-hand side is small, and condition (3) expresses every object of C as a filtered colimit of its objects. Hence C is compactly generated. $\square$

The preceding statement is often the most convenient way to prove compact generation. Notice the distinction between the two closure operations: the generators produce all objects under small colimits, but the compact objects are obtained from the generators using only finite colimits, shifts, and retracts.

Definition 22.3.5 (Ind-completion). Let C be a small ∞ -category. Its *Ind-completion* is the full subcategory

$$\mathrm{Ind}(C) \subseteq \mathrm{PSh}(C)$$

spanned by the filtered colimits of representable presheaves. The Yoneda embedding factors through a fully faithful functor $C \hookrightarrow \mathrm{Ind}(C)$.

Thus $\mathrm{Ind}(C)$ is the partial free cocompletion from Proposition 22.1.2 obtained by freely adjoining filtered colimits. In particular, if D admits filtered colimits, restriction along the Yoneda embedding induces an equivalence

$$\mathrm{Fun}_\omega(\mathrm{Ind}(C), D) \xrightarrow{\sim} \mathrm{Fun}(C, D),$$

where the left-hand side denotes the full subcategory of functors preserving filtered colimits.

Proposition 22.3.6 ([Lur09b, Corollary 4.2.3.11, Propositions 5.3.2.9, 5.3.5.10 and 5.3.5.14, and Theorem 5.5.1.1]). *If C is a small ∞ -category with finite colimits, then $\mathrm{Ind}(C)$ admits small colimits and is presentable. Moreover, the inclusion $C \hookrightarrow \mathrm{Ind}(C)$ preserves finite colimits, and the inclusion*

$$\mathrm{Ind}(C) \hookrightarrow \mathrm{PSh}(C)$$

identifies $\mathrm{Ind}(C)$ with the full subcategory spanned by the presheaves $C^{\mathrm{op}} \rightarrow \mathbf{An}$ which send finite colimits in C to limits in $\mathbf{An}$.

Proposition 22.3.7 (Compactly generated categories as Ind-completions, [Lur09b, Propositions 5.5.7.8 and 5.5.7.10]). *Let D be an ∞ -category with small colimits and assume that D^ω is small. The following are equivalent:*

- (1) *Every object of D is a filtered colimit of compact objects.*
- (2) *The restricted Yoneda functor induces an equivalence*

$$D \xrightarrow{\sim} \mathrm{Ind}(D^\omega).$$

Under these conditions, a colimit-preserving functor out of D is uniquely determined by its restriction to D^ω .

Remark 22.3.8 (Compact presentations). A presentable ∞ -category D is compactly generated if and only if it admits a presentation

$$D \simeq \mathrm{PSh}(C)^{S\text{-loc}}$$

in which C is small and S is a small collection of morphisms between compact objects of $\mathrm{PSh}(C)$. We will not use this characterization; see [Lur09b, Corollary 5.5.7.3 and Proposition 5.5.7.8].

Lemma 22.3.9 (Animae as Ind-completions). *The restricted Yoneda functors induce equivalences*

$$\mathbf{An} \xrightarrow{\sim} \mathrm{Ind}(\mathbf{An}^{\mathrm{fin}}) \quad \text{and} \quad \mathbf{An}_* \xrightarrow{\sim} \mathrm{Ind}(\mathbf{An}_*^{\mathrm{fin}}).$$

Proof. The first equivalence follows immediately from Propositions 4.3.11 and 22.3.6. For the second, Lemma 4.1.13 gives

$$\mathrm{Fun}^{\mathrm{lex}}((\mathbf{An}_*^{\mathrm{fin}})^{\mathrm{op}}, \mathbf{An}) \xrightarrow{\sim} \mathrm{Fun}^{\mathrm{lex}}((\mathbf{An}_*^{\mathrm{fin}})^{\mathrm{op}}, \mathbf{An}_*),$$

and the right-hand side is equivalent to $\mathbf{An}_*$ by Lemma 4.3.12. In both cases the inverse to evaluation is the restricted Yoneda functor, so the result follows from Proposition 22.3.6. $\square$

Example 22.3.10. The categories $\mathbf{An}$, $\mathbf{An}_*$, and $\mathbf{Sp}$ are compactly generated. For animae and pointed animae this follows from Lemma 22.3.9. The sphere spectrum is a compact generator of $\mathbf{Sp}$: compactness follows because $\mathrm{Hom}_{\mathbf{Sp}}(\mathbb{S}, -) \simeq \Omega^\infty$ preserves filtered colimits by Lemma 4.3.21, while its shifts jointly detect isomorphisms by Corollary 4.4.26; now apply Proposition 22.3.4.

For every associative ring spectrum A , the free module A is a compact generator of $\mathbf{LMod}_A$ by Lemma 8.1.8; for every commutative ring R , the complex $R[0]$ is a compact generator of $\mathbf{D}(R)$ by Corollary 20.2.5. Consequently,

$$\mathbf{Sp} \simeq \mathrm{Ind}(\mathbf{Sp}^\omega), \quad \mathbf{LMod}_A \simeq \mathrm{Ind}(\mathrm{Perf}(A)), \quad \mathbf{D}(R) \simeq \mathrm{Ind}(\mathrm{Perf}(R)).$$

The Eilenberg–MacLane comparison of Corollary 8.3.4 identifies the last presentation with the corresponding presentation of $\mathbf{Mod}_{HR}$.

22.4 Tensor products of presentable ∞ -categories

In a context of ∞ -categories admitting colimits, one frequently works with bifunctors $C \times D \rightarrow E$ that are bilinear, in the sense that they preserve colimits in both arguments separately. There exists a universal such bifunctor, which is denoted by $C \times D \rightarrow C \otimes_{\mathrm{colim}} D$, and is characterized by the fact that bilinear bifunctors correspond to colimit-preserving functors $C \otimes_{\mathrm{colim}} D \rightarrow E$. The goal of this section is to explain how to turn this into a symmetric monoidal structure on the (very large) ∞ -category of ∞ -categories with colimits. It is convenient to allow for an arbitrary class $\mathcal{K}$ of indexing categories. We specialize to tensor products of presentable ∞ -categories afterwards.

For a collection $\mathcal{K}$ of small ∞ -categories, write

$$\widehat{\mathrm{Cat}}_{\infty}^{\mathcal{K}\text{-colim}} \subseteq \widehat{\mathrm{Cat}}_{\infty}$$

for the ∞ -category whose objects are the possibly large ∞ -categories admitting I -indexed colimits for every $I \in \mathcal{K}$, and whose morphisms preserve all such colimits. More generally, for $C_1, \dots, C_n, D$ admitting these colimits, we say that a functor

$$F: C_1 \times \dots \times C_n \longrightarrow D$$

$\mathcal{K}$ -multilinear if it preserves I -indexed colimits in each variable separately for every $I \in \mathcal{K}$.

Lemma 22.4.1. *There is a unique suboperad*

$$\mathcal{O}_{\mathcal{K}\text{-colim}} \subseteq \mathcal{O}_{\widehat{\mathrm{Cat}}_{\infty}}^{\times}$$

of the cartesian monoidal structure on $\widehat{\mathrm{Cat}}_{\infty}$ whose colors are the objects of $\widehat{\mathrm{Cat}}_{\infty}^{\mathcal{K}\text{-colim}}$, and its multimorphisms are the $\mathcal{K}$ -multilinear functors.

Proof. The two conditions specify a unique subcategory

$$\mathcal{O}_{\mathcal{K}\text{-colim}}^{\otimes} \subseteq \widehat{\mathrm{Cat}}_{\infty}^{\times}$$

of the total category of the cartesian monoidal structure, and we need to verify this defines a suboperad. Identity functors preserve the prescribed colimits. Operadic composites of $\mathcal{K}$ -multilinear

functors are again $\mathcal{K}$ -multilinear: when one input varies, it occurs in a single inner operation, and preservation of its colimit may first be passed through that operation and then through the relevant variable of the outer operation. The cocartesian lift of a backwards span in the cartesian operad merely reindexes a tuple of colors, and all its component operations are identity functors. It therefore belongs to the suboperad. Its cocartesian universal property restricts to the suboperad because both sides of the relevant mapping-anima equivalence impose the same componentwise $\mathcal{K}$ -multilinearity condition. Finally, the Segal decomposition of a tuple into its individual colors restricts to the suboperad, since $\mathcal{K}$ -multilinearity of a functor with tuple-valued target can be checked componentwise. The characterization in Proposition 14.1.7 now applies. $\square$

The non-formal point is that the multimorphisms in this operad are corepresentable.

Theorem 22.4.2 (Tensor products of categories with prescribed colimits, [Lur17, Corollary 4.8.1.4 and Variant 4.8.1.11]). *The ∞ -operad $\mathcal{O}_{\mathcal{K}\text{-colim}}$ is the multimorphism operad of a symmetric monoidal ∞ -category*

$$(\widehat{\text{Cat}}_{\infty}^{\mathcal{K}\text{-colim}}, \otimes_{\mathcal{K}}).$$

Remark 22.4.3. Thus, for every finite collection $C_1, \dots, C_n$ of its objects, there is an object

$$C_1 \otimes_{\mathcal{K}} \cdots \otimes_{\mathcal{K}} C_n$$

and a $\mathcal{K}$ -multilinear functor from $C_1 \times \cdots \times C_n$ to this object such that composition induces an equivalence

$$\text{Fun}_{\mathcal{K}}(C_1 \otimes_{\mathcal{K}} \cdots \otimes_{\mathcal{K}} C_n, D) \xrightarrow{\sim} \text{Fun}_{\mathcal{K}, \dots, \mathcal{K}}(C_1 \times \cdots \times C_n, D).$$

Here the right-hand side is the full subcategory spanned by the $\mathcal{K}$ -multilinear functors. If each C_i is small, then $C_1 \otimes_{\mathcal{K}} \cdots \otimes_{\mathcal{K}} C_n$ is small as well.

Suppose first that $\mathcal{K}$ consists of the finite discrete categories. Then $\widehat{\text{Cat}}_{\infty}^{\mathcal{K}\text{-colim}}$ is the ∞ -category $\widehat{\text{Cat}}_{\infty}^{\text{coprod}}$ of possibly large ∞ -categories with finite coproducts and finite-coproduct-preserving functors. We write the resulting symmetric monoidal structure as $(\widehat{\text{Cat}}_{\infty}^{\text{coprod}}, \otimes_{\text{coprod}})$; its tensor product is characterized by finite-coproduct-preserving multilinear functors. The full subcategory $\text{Cat}_{\infty}^{\text{coprod}}$ on the small colors is closed under this tensor product. In particular, a commutative algebra in $(\text{Cat}_{\infty}^{\text{coprod}}, \otimes_{\text{coprod}})$ is precisely a small symmetric monoidal ∞ -category whose tensor product preserves finite coproducts separately in each variable.

At the other extreme, let $\mathcal{K}$ consist of all small ∞ -categories, denote the resulting ∞ -category by $\widehat{\text{Cat}}_{\infty}^{\text{colim}}$, and write $\otimes$ for this specialization of $\otimes_{\mathcal{K}}$. Its colors are the cocomplete ∞ -categories and its morphisms are the colimit-preserving functors. The full subcategory $\text{Pr}^{\text{L}} \subseteq \widehat{\text{Cat}}_{\infty}^{\text{colim}}$ on the presentable colors contains the unit $\mathbf{An}$ and is closed under the tensor product by [Lur17, Proposition 4.8.1.15]. It is therefore a symmetric monoidal subcategory

$$(\text{Pr}^{\text{L}}, \otimes) \subseteq (\widehat{\text{Cat}}_{\infty}^{\text{colim}}, \otimes),$$

with tensor product characterized by

$$\text{Fun}^{\text{L}}(C \otimes D, E) \simeq \text{Fun}^{\text{L}, \text{L}}(C \times D, E).$$

This is the monoidal structure used in Section 18.5.

22.5 Presentably symmetric monoidal categories

Definition 22.5.1. A symmetric monoidal ∞ -category $(C, \otimes)$ is called *presentably symmetric monoidal* if C is presentable and the tensor product $- \otimes -: C \times C \rightarrow C$ preserves small colimits separately in both variables.

Remark 22.5.2. The suboperad description in Section 22.4 identifies presentably symmetric monoidal ∞ -categories with commutative algebra objects of $(\mathrm{Pr}^{\mathrm{L}}, \otimes)$. Indeed, a commutative algebra object is an operad map

$$\mathrm{Comm} \longrightarrow \mathcal{M}_{(\mathrm{Pr}^{\mathrm{L}}, \otimes)} \subseteq \mathcal{O}_{\mathrm{Cat}_{\infty}}^{\times}.$$

After forgetting the displayed factorization, this is a symmetric monoidal ∞ -category $(C, \otimes_C)$. Its underlying ∞ -category is presentable, and each of its multimorphism functors $C^n \rightarrow C$ preserves small colimits separately in every variable because the operad map lands in $\mathcal{M}_{(\mathrm{Pr}^{\mathrm{L}}, \otimes)}$. In particular, its binary tensor product has the property required in Definition 22.5.1.

Conversely, let $(C, \otimes_C)$ be presentably symmetric monoidal. Its symmetric monoidal structure gives an operad map $\mathrm{Comm} \rightarrow \mathcal{O}_{\mathrm{Cat}_{\infty}}^{\times}$. The binary tensor product preserves small colimits separately by assumption, and hence so do all iterated tensor products. The operad map therefore factors through $\mathcal{M}_{(\mathrm{Pr}^{\mathrm{L}}, \otimes)}$. The same factorization criterion for morphisms identifies maps in $\mathrm{CAlg}(\mathrm{Pr}^{\mathrm{L}})$ with colimit-preserving symmetric monoidal functors. This gives the claimed identification of ∞ -categories.

Proposition 22.5.3. *Every presentably symmetric monoidal ∞ -category is closed: for every object $X \in C$, the functor $X \otimes -: C \rightarrow C$ admits a right adjoint $\underline{\mathrm{Hom}}_C(X, -)$.*

Proof. This follows immediately from Theorem 22.2.5. □

The construction of Ind-categories is compatible with symmetric monoidal structures. We state the version for stable categories, where preservation of finite colimits is equivalent to exactness.

Theorem 22.5.4 (Symmetric monoidal Ind-completion, [Lur17, Corollary 4.8.1.14]). *Let C be a small stable symmetric monoidal ∞ -category, and assume that the tensor product on C is exact separately in both variables. Then $\mathrm{Ind}(C)$ admits a unique symmetric monoidal structure with the following properties:*

- (1) *The Yoneda embedding $C \hookrightarrow \mathrm{Ind}(C)$ is symmetric monoidal.*
- (2) *The tensor product on $\mathrm{Ind}(C)$ preserves colimits separately in both variables.*

For filtered-colimit presentations $X \simeq \mathrm{colim}_i X_i$ and $Y \simeq \mathrm{colim}_j Y_j$ with $X_i, Y_j \in C$, its underlying tensor product is characterized by

$$X \otimes Y \simeq \mathrm{colim}_{(i,j)} (X_i \otimes Y_j).$$

Two further closure properties are used repeatedly in Part II.

Proposition 22.5.5. *The following statements hold.*

- (1) *If C is a small symmetric monoidal ∞ -category and D is presentably symmetric monoidal, then the Day convolution structure on $\mathrm{Fun}(C, D)$ is presentably symmetric monoidal.*

(2) If C is presentably symmetric monoidal and $A \in \text{Alg}(C)$, then $\text{LMod}_A(C)$ and $\text{RMod}_A(C)$ are presentable. Their forgetful functors to C create limits and small colimits.

Proof. For (1), the underlying functor category $\text{Fun}(C, D)$ is presentable by Theorem 22.2.2, and Corollary 16.2.8 equips it with the Day convolution monoidal structure. The category D is complete because it is presentable, and it is closed by Proposition 22.5.3. Hence Proposition 16.2.13 shows that the Day convolution structure is closed. Its tensor product therefore preserves small colimits separately in both variables.

Part (2) is [Lur17, Corollary 4.2.3.7]. □

Exercises

23 Cocartesian fibrations and (un)straightening

An important notion in ∞ -category theory is that of a *cocartesian fibration*. Roughly speaking, a cocartesian fibration is a functor $p: E \rightarrow C$ whose fibers $E_x := E \times_C \{x\}$ vary covariantly in x , in the sense that for every morphism $f: x \rightarrow y$ in C we obtain a *cocartesian transport functor* $f_!: E_x \rightarrow E_y$. These functors are fully coherent in f , in the sense that they assemble into a functor $\mathrm{Str}(p): C \rightarrow \mathrm{Cat}_\infty$, given on objects by sending x to E_x and on morphisms by sending f to $f_!$. This provides a powerful way to construct functors into Cat_∞ , known as the *straightening/unstraightening correspondence*. For example, it will allow us to formally define the Hom-functor $\mathrm{Hom}_C: C^{\mathrm{op}} \times C \rightarrow \mathrm{An}$ of an ∞ -category C .

23.1 Cartesian and cocartesian fibrations

We start with some basic definitions regarding (co)cartesian fibrations.

Definition 23.1.1. Consider a functor $p: E \rightarrow C$.

- (1) A morphism $\varphi: e \rightarrow e'$ in E is called *p-cocartesian* if, for every other object e'' in E , the commutative square

$$\begin{array}{ccc} \mathrm{Hom}_E(e', e'') & \xrightarrow{- \circ \varphi} & \mathrm{Hom}_E(e, e'') \\ p \downarrow & & \downarrow p \\ \mathrm{Hom}_C(pe', pe'') & \xrightarrow{- \circ p(\varphi)} & \mathrm{Hom}_C(pe, pe'') \end{array}$$

is a pullback square. Informally, this means that for every solid diagram of the form

$$\begin{array}{ccc} & e' & \\ \varphi \nearrow & & \searrow \exists! \\ e & \longrightarrow & e'' \end{array} \quad \xrightarrow{p} \quad \begin{array}{ccc} & pe' & \\ p\varphi \nearrow & & \searrow \\ pe & \longrightarrow & pe'' \end{array}$$

in E for which its image in C has been completed to a triangle, there exists a unique map $e' \rightarrow e''$ that forms a commutative triangle in E and projects to the given one in C .

- (2) We say that p is a *cocartesian fibration* if for every morphism $f: x \rightarrow y$ in C and every object $e \in E_x$ there exists a *p-cocartesian* morphism $\varphi: e \rightarrow e'$ satisfying $p(\varphi) = f$. We refer to such a morphism φ as a *p-cocartesian lift* of f .

- (3) Let $p: E \rightarrow C$ and $p': E' \rightarrow C$ be two cocartesian fibrations. A functor $F: E \rightarrow E'$ over C , i.e., a commutative triangle

$$\begin{array}{ccc} E & \xrightarrow{F} & E' \\ p \searrow & & \swarrow p' \\ & C, & \end{array}$$

is called *cocartesian over C* if it sends p -cocartesian morphisms in E to p' -cocartesian morphisms in E' .

- (4) If C is a small ∞ -category, we denote by

$$\mathrm{Cocart}(C) \subseteq (\mathrm{Cat}_\infty)_{/C}$$

the non-full subcategory spanned by the cocartesian fibrations $p: E \rightarrow C$ and the cocartesian functors between them.

The cocartesian lift $\varphi: e \rightarrow e'$ is functorial in the pair (f, e) . To show this, the following reformulation of p -cocartesian morphisms is useful:

Exercise 23.1.2. Let $p: E \rightarrow C$ be a functor. Consider the functor

$$\mathrm{Ar}(E) \rightarrow \mathrm{Ar}(C) \times_{s,C,p} E, \quad (\varphi: e \rightarrow e') \mapsto (p(\varphi): p(e) \rightarrow p(e'), e).$$

Then a p -cocartesian lift of a morphism $f: x \rightarrow y$ starting in e is precisely the same thing as a left adjoint object to (f, e) under this functor.

Notation 23.1.3. Given an object $e \in E_x$, we may sometimes denote a cocartesian lift of $f: x \rightarrow y$ by $\varphi: e \rightarrow f_!e$, and refer to $f_!e$ as the *cocartesian transport of e along f* . This is well-defined, since left adjoint objects are unique whenever they exist.

Proposition 23.1.4 (Functoriality of partial cocartesian lifts). *Let $p: E \rightarrow C$ be a functor, and let $W \subseteq \mathrm{Ar}(C)$ be a subcategory. Suppose that for every $(f, e) \in W \times_{s,C,p} E$ there exists a p -cocartesian lift of f starting in e . Then these lifts assemble into a functor*

$$\mathrm{lift}_W: W \times_{s,C,p} E \longrightarrow \mathrm{Ar}(E)$$

More precisely, this functor factors through

$$\mathrm{Ar}(E) \times_{\mathrm{Ar}(C) \times_{s,C,p} E} (W \times_{s,C,p} E),$$

and the resulting functor is left adjoint to the projection from this pullback to $W \times_{s,C,p} E$.

Proof. By the previous exercise, the required cocartesian lifts are precisely the left adjoint objects under the restricted functor

$$\mathrm{Ar}(E) \times_{\mathrm{Ar}(C) \times_{s,C,p} E} (W \times_{s,C,p} E) \longrightarrow W \times_{s,C,p} E.$$

The claim is therefore an immediate consequence of the pointwise criterion for adjunctions from Lemma 21.1.4. $\square$

Corollary 23.1.5. *Let $p: E \rightarrow C$ be a cocartesian fibration. Then the cocartesian lifts assemble into a left adjoint*

$$\text{lift}: \text{Ar}(C) \times_{s,C,p} E \rightarrow \text{Ar}(E), \quad (f, e) \mapsto (\varphi: e \rightarrow f_!)$$

to the functor from the previous exercise.

Proof. Apply Proposition 23.1.4 with $W = \text{Ar}(C)$. □

Corollary 23.1.6. *Let $p: E \rightarrow C$ be a cocartesian fibration, and let $f: x \rightarrow y$ be a morphism in C . Then cocartesian transport assembles into a functor $f_!: E_x \rightarrow E_y$.*

Proof. We may obtain $f_!$ by restricting the functor lift from the previous corollary. □

We may also deduce the uniqueness of cocartesian lifts:

Exercise 23.1.7. For a functor $p: E \rightarrow C$, consider the full subcategory

$$\text{Ar}^{\text{cocart}}(E) \subseteq \text{Ar}(E)$$

spanned by the p -cocartesian morphisms. Show that the functor

$$\text{Ar}^{\text{cocart}}(E) \rightarrow \text{Ar}(C) \times_{s,C,p} E, \quad (\varphi: e \rightarrow e') \mapsto (p\varphi, e)$$

is fully faithful. Deduce that p is a cocartesian fibration if and only if this functor is an equivalence.

The definition of cocartesian fibrations may be dualized:

Definition 23.1.8. For a functor $p: E \rightarrow C$, we say that a morphism $\varphi: e \rightarrow e'$ in E is p -cartesian if for every third object e'' in E the commutative square

$$\begin{array}{ccc} \text{Hom}_E(e'', e) & \xrightarrow{\varphi \circ -} & \text{Hom}_E(e'', e') \\ p \downarrow & & \downarrow p \\ \text{Hom}_C(pe'', pe) & \xrightarrow{p\varphi \circ -} & \text{Hom}_C(pe'', pe') \end{array}$$

is a pullback square. We say p is a *cartesian fibration* if for every morphism $f: x \rightarrow y$ and every $e' \in E_y$ there exists a p -cartesian lift $\varphi: e \rightarrow e'$ of f . We similarly obtain a notion of a *cartesian functor over C* , and this leads to a (non-full) subcategory

$$\text{Cart}(C) \subseteq (\text{Cat}_\infty)_{/C}.$$

Remark 23.1.9. Note that a functor $p: E \rightarrow C$ is a cartesian fibration if and only if its opposite $p^{\text{op}}: E^{\text{op}} \rightarrow C^{\text{op}}$ is a cocartesian fibration. All properties about cocartesian fibrations have dual analogues for cartesian fibrations; we will generally not spell these out explicitly.

Let us go through a basic example of a (co)cartesian fibration to see these definitions in action:

Example 23.1.10. Let C be an ∞ -category, let $E := \text{Ar}(C)$ be its arrow category, and consider the target functor $p := t: \text{Ar}(C) \rightarrow C$. An object of E is given by a morphism in C , which for emphasis we will write vertically: $\begin{pmatrix} X \\ \downarrow \\ Y \end{pmatrix}$. A morphism $\varphi: \begin{pmatrix} X \\ \downarrow \\ Y \end{pmatrix} \rightarrow \begin{pmatrix} X' \\ \downarrow \\ Y' \end{pmatrix}$ in E is given by a commutative square in C of the form

$$\begin{array}{ccc} X & \xrightarrow{g} & X' \\ \downarrow & & \downarrow \\ Y & \xrightarrow{f} & Y' \end{array}$$

We claim:

- (1) The morphism φ in E is p -cocartesian if and only if g is an equivalence. In particular, t is always a cocartesian fibration: given $e = \begin{pmatrix} X \\ \downarrow \\ Y \end{pmatrix}$ and $f: p(\begin{pmatrix} X \\ \downarrow \\ Y \end{pmatrix}) = Y \rightarrow Y'$ we may always form the square with $X' := X$ and $g = \text{id}_X$.
- (2) The morphism φ in E is p -cartesian if and only if the square is a pullback square (a.k.a. a *cartesian square*). In particular, t is a cartesian fibration if and only if C admits pullbacks.

I will leave it to the reader to spell out a detailed proof of these claims, see Exercise 23.3. But let's at least see what these claims are heuristically saying.

Claim (1) says that for every third object $\begin{pmatrix} X'' \\ \downarrow \\ Y'' \end{pmatrix}$ of E and every solid diagram below, there exists a unique dashed arrow making the diagram commute:

$$\begin{array}{ccccc} & & X' & & \\ & \nearrow g & \downarrow & \searrow & \\ X & \xrightarrow{\quad} & X'' & & \\ \downarrow & & \downarrow & & \downarrow \\ Y & \xrightarrow{f} & Y' & \searrow & Y'' \\ & & & \nearrow & \end{array}$$

If g is an equivalence, we may invert it and it is clear that the dashed filler is unique. Conversely, if φ is p -cocartesian, then by taking $X'' = X$ we obtain a morphism $g': X' \rightarrow X$ satisfying $g'g \cong \text{id}_X$, and by taking $X'' = X'$ we then use uniqueness to get that $gg' = \text{id}_{X'}$. (Alternatively, we use that there always exists a p -cocartesian lift with g an equivalence so the claim follows by uniqueness of p -cocartesian lifts.)

Claim (2) says that for every third object $\begin{pmatrix} X'' \\ \downarrow \\ Y'' \end{pmatrix}$ of E and every solid diagram below, there exists a unique dashed arrow making the diagram commute:

$$\begin{array}{ccccc} & & X & & \\ & \nearrow & \downarrow & \searrow g & \\ X'' & \xrightarrow{\quad} & X' & & \\ \downarrow & & \downarrow & & \downarrow \\ Y'' & \xrightarrow{\quad} & Y' & & \\ & & \nearrow f & & \end{array}$$

This is a reformulation of the condition that the original square is a pullback square.

Example 23.1.11. The forgetful functor $\text{VectBund} \rightarrow \text{Top}$, $(X, E) \mapsto X$ is a cartesian fibration. See Exercise 23.4 for details.

Lemma 23.1.12. *Show that cocartesian fibrations are closed under composition and pullback.*

Proof. This is left as an exercise, see Exercises 23.5 and 23.6 □

Exercise 23.1.13. Show that $C \rightarrow *$ is both a cartesian and cocartesian fibration. Deduce that any projection functor $\text{pr}_D: C \times D \rightarrow D$ is both a cartesian and cocartesian fibration, and that a morphism $(f: c \rightarrow c', g: d \rightarrow d')$ is pr_D -cocartesian (or pr_D -cartesian) if and only if f is an isomorphism.

Lemma 23.1.14 (Right cancellation of cocartesian morphisms). *Let $p: E \rightarrow C$ be a functor and let $f: x \rightarrow y$ and $g: y \rightarrow z$ be morphisms in E such that f is p -cocartesian. Then g is p -cocartesian if and only if gf is p -cocartesian.*

Proof. For every other object $w \in E$, we consider the following commutative diagram:

$$\begin{array}{ccccc} \text{Hom}_E(z, w) & \xrightarrow{- \circ g} & \text{Hom}_E(y, w) & \xrightarrow{- \circ f} & \text{Hom}_E(x, w) \\ p \downarrow & & p \downarrow p & & \downarrow p \\ \text{Hom}_C(pz, pw) & \xrightarrow{- \circ p(g)} & \text{Hom}_C(py, pw) & \xrightarrow{- \circ p(f)} & \text{Hom}_C(px, pw). \end{array}$$

Since f is p -cocartesian, the right square is a pullback square. It follows from the pasting law for pullback squares that the left square is a pullback if and only if the outer square is a pullback. This proves the claim. □

Exercise 23.1.15. Let $p: E \rightarrow C$ be a functor and let $\varphi: e \rightarrow e'$ be a p -cocartesian morphism. Then φ is an isomorphism if and only if $p(\varphi)$ is an isomorphism.

23.1.1 Left and right fibrations

Proposition 23.1.16. *Let $p: E \rightarrow C$ be a cocartesian fibration. Then the following three conditions are equivalent:*

- (1) *The functor p is conservative, i.e., a morphism f in E is an isomorphism as soon as $p(f)$ is an isomorphism in C ;*
- (2) *For each $x \in C$, the fiber $E_x := \{x\} \times_C E$ is an anima;*
- (3) *Every morphism in E is p -cocartesian.*

Proof. It is clear that (1) implies (2): every morphism in the fiber E_x is sent to the identity morphism id_x , hence is an isomorphism by conservativity of p .

Assuming (2), consider a morphism $\psi: e \rightarrow e''$ in E . We need to show that ψ is p -cocartesian. Since p is a cocartesian fibration, there exists a p -cocartesian morphism $\varphi: e \rightarrow e'$ lifting $p(\psi)$. Since φ is p -cocartesian, there then exists a unique morphism $\xi: e' \rightarrow e''$ fitting in the following commutative diagram lifting the degenerate one in C :

$$\begin{array}{ccc} & e' & \\ \varphi \nearrow & & \searrow \xi \\ e & \xrightarrow{\psi} & e'' \end{array} \quad \xrightarrow{p} \quad \begin{array}{ccc} & pe'' & \\ p\psi \nearrow & & \searrow \\ pe & \xrightarrow{p\psi} & pe'' \end{array}$$

But then the morphism ξ lies in the fiber $E_{pe''}$, which is an anima, so ξ is an isomorphism. In particular ξ is p -cocartesian, and hence so is $\psi = \xi \circ \varphi$.

Finally, assume that (3) holds, and consider a morphism $\varphi: e \rightarrow e'$ in E such that $p\varphi$ is invertible. We need to show that φ is invertible. By assumption, there is an inverse $g: pe' \rightarrow pe$ of $p\varphi$. We may then find a p -cocartesian lift $\psi: e' \rightarrow e''$ of g . But then $\varphi \circ \psi$ and $\psi \circ \varphi$ are cocartesian lifts of the respective identity maps on pe and pe' . By uniqueness of cocartesian lifts (Exercise 23.1.7) these composites are isomorphisms. It follows that φ is an isomorphism with inverse ψ . $\square$

Definition 23.1.17. A functor $p: E \rightarrow C$ is called a *left fibration* if it is a cocartesian fibration satisfying the equivalent conditions (1), (2) and (3) from the proposition. This gives rise to a full subcategory

$$\mathbf{LFib}(C) \subseteq \mathbf{Cocart}(C).$$

Note that $\mathbf{LFib}(C)$ is also a full subcategory of $(\mathbf{Cat}_\infty)_/C$ (in contrast to $\mathbf{Cocart}(C)$).

Dually, p is called a *right fibration* if it is a cartesian fibration which satisfies the equivalent conditions (1), (2) and (3'): Every morphism in E is cartesian. We write $\mathbf{RFib}(C) \subseteq (\mathbf{Cat}_\infty)_/C$ for the resulting full subcategory.

Lemma 23.1.18. *Left fibrations are closed under composition and pullback.*

Proof. Follows from Lemma 23.1.12. $\square$

Proposition 23.1.19. *Let C be an ∞ -category. For every object $x \in C$, the functor $t: C_{x/} \rightarrow C$ sending a morphism $x \rightarrow y$ to its target y is a left fibration. Dually, the source functor $s: C_{/x} \rightarrow C$ is a right fibration.*

Proof. We prove the case for $t: C_{x/} \rightarrow C$; the other case is dual. Consider a morphism in $C_{x/}$, which takes the form of a commutative triangle

$$\begin{array}{ccc} & x & \\ g \swarrow & & \searrow h \\ y & \xrightarrow{f} & z. \end{array}$$

To show that this morphism is cocartesian with respect to the target functor $t: C_{x/} \rightarrow C$, we have to show that for every third object $x \rightarrow w$, the top square in the following commutative diagram is a pullback square:

$$\begin{array}{ccccc} \mathrm{Hom}_{C_{x/}}(z, w) & \xrightarrow{-\circ f} & \mathrm{Hom}_{C_{x/}}(y, w) & & \\ \downarrow & \searrow t & \downarrow & \searrow t & \\ \mathrm{Hom}_C(z, w) & \xrightarrow{-\circ f} & \mathrm{Hom}_C(y, w) & & \\ \downarrow & \searrow -\circ h & \downarrow & \searrow -\circ g & \\ * & \xrightarrow{\quad} & * & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ \mathrm{Hom}_C(x, w) & \xrightarrow{\quad} & \mathrm{Hom}_C(x, w) & & \end{array}$$

But this follows from the pasting law for pullback squares, since the bottom square and left/right faces are pullback squares. $\square$

23.1.2 Adjunctions

Lemma 23.1.20. *Let $p: E \rightarrow C$ be a cocartesian fibration. Then the following conditions are equivalent:*

- (1) *The functor p is also a cartesian fibration.*
- (2) *For every morphism $f: X \rightarrow Y$ in C , the cocartesian transport functor $f_!: E_X \rightarrow E_Y$ admits a right adjoint $f^*: E_Y \rightarrow E_X$.*

Proof. First assume (2). We need to show that for every morphism $f: X \rightarrow Y$ and any lift $\tilde{Y} \in E_Y$ of Y , there exists a p -cartesian lift $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$. To this end, we define $\tilde{X} := f^*\tilde{Y} \in E_X$. There exists a p -cocartesian lift $\tilde{X} \rightarrow f_!\tilde{X}$ in E , and by composing this with the counit $\varepsilon: f_!\tilde{X} = f_!f^*\tilde{Y} \rightarrow \tilde{Y}$ we obtain a morphism $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$. To show this is p -cartesian, consider a third object $\tilde{Z}$. We must show that the square

$$\begin{array}{ccc} \mathrm{Hom}_E(\tilde{Z}, \tilde{X}) & \xrightarrow{\tilde{f} \circ -} & \mathrm{Hom}_E(\tilde{Z}, \tilde{Y}) \\ p \downarrow & & \downarrow p \\ \mathrm{Hom}_C(Z, X) & \xrightarrow{f \circ -} & \mathrm{Hom}_C(Z, Y) \end{array}$$

is a pullback square. We may check this on fibers over any morphism $g: Z \rightarrow X$, which we denote by

$$\tilde{f} \circ - : \mathrm{Hom}_E^g(\tilde{Z}, \tilde{X}) \rightarrow \mathrm{Hom}_E^{fg}(\tilde{Z}, \tilde{Y}).$$

By picking a cocartesian lift $\tilde{g}: \tilde{Z} \rightarrow g_!\tilde{Z}$, and using the universal property to replace $\tilde{Z}$ by $g_!\tilde{Z}$, we may reduce to the case $Z = X$ and $g = \mathrm{id}_X$. In this case, we must show that the top composite in the following commutative diagram is an equivalence:

$$\begin{array}{ccccc} \mathrm{Hom}_E^{\mathrm{id}_X}(\tilde{Z}, \tilde{X}) & \longrightarrow & \mathrm{Hom}_E^f(\tilde{Z}, f_!\tilde{X}) & \xrightarrow{\varepsilon \circ -} & \mathrm{Hom}_E^f(\tilde{Z}, \tilde{Y}) \\ & \searrow f_! & \uparrow \sim & & \uparrow \sim \\ & & \mathrm{Hom}_E^{\mathrm{id}_Y}(f_!\tilde{Z}, f_!\tilde{X}) & \xrightarrow{\varepsilon \circ -} & \mathrm{Hom}_E^{\mathrm{id}_Y}(f_!\tilde{Z}, \tilde{Y}). \end{array}$$

Here the two vertical equivalences are given by precomposition with the p -cocartesian morphism $\tilde{Z} \rightarrow f_!\tilde{Z}$, the right square commutes by associativity of composition, and the left square commutes by the functoriality of cocartesian lifts (Corollary 23.1.5). We may thus equivalently show that the bottom composite is an equivalence. But this composite is precisely the adjunction equivalence

$$\mathrm{Hom}_{E_X}(\tilde{Z}, f^*\tilde{Y}) \xrightarrow{\sim} \mathrm{Hom}_{E_Y}(f_!\tilde{Z}, \tilde{Y}).$$

This shows that (2) $\Rightarrow$ (1). Conversely, if p is a cartesian fibration, then for an object $\tilde{Y} \in E_Y$ we obtain a p -cartesian lift $f^*\tilde{Y} \rightarrow \tilde{Y}$ in E . This map factors uniquely through the cocartesian morphism $f^*\tilde{Y} \rightarrow f_!f^*\tilde{Y}$, resulting in a morphism $\varepsilon: f_!f^*\tilde{Y} \rightarrow \tilde{Y}$. We claim that this exhibits $f^*\tilde{Y}$ as a right-adjoint object to $\tilde{Y}$ under the functor $f_!$, i.e. that for every $\tilde{Z} \in E_X$ the composite

$$\mathrm{Hom}_{E_X}(\tilde{Z}, f^*\tilde{Y}) \xrightarrow{f_!} \mathrm{Hom}_{E_Y}(f_!\tilde{Z}, f_!f^*\tilde{Y}) \xrightarrow{\varepsilon \circ -} \mathrm{Hom}_{E_Y}(f_!\tilde{Z}, \tilde{Y})$$

is an equivalence. Reversing the logic used for the other implication, this is a consequence of the fact that the morphism $f^*\tilde{Y} \rightarrow \tilde{Y}$ is p -cartesian. $\square$

Corollary 23.1.21. *Let $F: C \rightarrow D$ be a functor between small ∞ -categories, encoded by a morphism $[1] \rightarrow \text{Cat}_\infty$. Let $E \rightarrow [1]$ be the cocartesian unstraightening of this functor. Then F admits a right adjoint if and only if the cocartesian fibration $E \rightarrow [1]$ is also a cartesian fibration.* $\square$

23.2 Straightening/unstraightening

Let $p: E \rightarrow C$ be a cocartesian fibration. For every object x of C , we may consider the fiber E_x of p over x . For a morphism $f: x \rightarrow y$ in C , there is a *cocartesian transport* functor $f_!: E_x \rightarrow E_y$. Furthermore, given another morphism $g: y \rightarrow z$, there is a natural isomorphism

$$g_! \circ f_! \cong (g \circ f)_!: E_x \rightarrow E_z.$$

Morally speaking, this construction should assemble into a functor

$$\text{Str}^{\text{cc}}(p): C \rightarrow \text{Cat}_\infty, \quad x \mapsto E_x, \quad f \mapsto f_!.$$

This is precisely the content of the straightening/unstraightening correspondence:

Theorem 23.2.1 (Straightening/Unstraightening, Lurie [Lur09b], Hebestreit, Heuts, and Ruit [HHR21]). *Let C be a small ∞ -category. Then there exist equivalences of ∞ -categories*

$$\text{Str}^{\text{cc}}: \text{Cocart}(C) \xrightarrow{\sim} \text{Fun}(C, \text{Cat}_\infty)$$

and

$$\text{Str}^{\text{ct}}: \text{Cart}(C) \xrightarrow{\sim} \text{Fun}(C^{\text{op}}, \text{Cat}_\infty).$$

Furthermore, these equivalences are natural²¹ in C , in the sense that for every functor $F: C \rightarrow D$ the following two squares commute:

$$\begin{array}{ccc} \text{Cocart}(D) & \xrightarrow{\text{Str}^{\text{cc}}} & \text{Fun}(D, \text{Cat}_\infty) \\ F^* \downarrow & & \downarrow - \circ F \\ \text{Cocart}(C) & \xrightarrow{\text{Str}^{\text{cc}}} & \text{Fun}(C, \text{Cat}_\infty) \end{array} \quad \text{and} \quad \begin{array}{ccc} \text{Cart}(D) & \xrightarrow{\text{Str}^{\text{ct}}} & \text{Fun}(D^{\text{op}}, \text{Cat}_\infty) \\ F^* \downarrow & & \downarrow - \circ F^{\text{op}} \\ \text{Cart}(C) & \xrightarrow{\text{Str}^{\text{ct}}} & \text{Fun}(C^{\text{op}}, \text{Cat}_\infty). \end{array}$$

Finally, the composites

$$\begin{aligned} \text{Cat}_\infty &\simeq \text{Cocart}(*) \xrightarrow{\text{Str}^{\text{cc}}} \text{Fun}(*, \text{Cat}_\infty) \simeq \text{Cat}_\infty \\ \text{Cat}_\infty &\simeq \text{Cart}(*) \xrightarrow{\text{Str}^{\text{ct}}} \text{Fun}(*, \text{Cat}_\infty) \simeq \text{Cat}_\infty \end{aligned}$$

are (isomorphic to) the identity functors.

This is one of the most fundamental and most important results in the theory of ∞ -categories. The theorem was first proved in the model of quasicategories by Jacob Lurie; the proof has since been substantially simplified by [HHR21]. Since this result is so fundamental to the theory, it may also be taken as an axiom, which is what we will do in this book.

²¹See Remark 23.2.10 for a more refined version of the naturality.

Given a cocartesian fibration $p: E \rightarrow C$, we refer to the functor $\mathrm{Str}^{\mathrm{cc}}(p): C \rightarrow \mathrm{Cat}_\infty$ as its *(cocartesian) straightening*, and similarly for $\mathrm{Str}^{\mathrm{ct}}(p)$. The inverses of $\mathrm{Str}^{\mathrm{cc}}$ and $\mathrm{Str}^{\mathrm{ct}}$ are denoted by

$$\mathrm{Un}^{\mathrm{cc}}: \mathrm{Fun}(C, \mathrm{Cat}_\infty) \xrightarrow{\sim} \mathrm{Cocart}(C), \quad \text{and} \quad \mathrm{Un}^{\mathrm{ct}}: \mathrm{Fun}(C^{\mathrm{op}}, \mathrm{Cat}_\infty) \xrightarrow{\sim} \mathrm{Cart}(C).$$

Given a functor $F: C \rightarrow \mathrm{Cat}_\infty$, we refer to the cocartesian fibration $\mathrm{Un}^{\mathrm{cc}}(F) \rightarrow C$ as its *cocartesian unstraightening*, and to the cartesian fibration $\mathrm{Un}^{\mathrm{ct}}(F)$ as the *cartesian unstraightening*.

We will use the following compatibility of unstraightening with functor categories. Here $\mathrm{const}: C \rightarrow \mathrm{Fun}(I, C)$ denotes the constant-diagram functor.

Proposition 23.2.2. *Let $p: E \rightarrow C$ be a cocartesian fibration classified by a functor $F: C \rightarrow \mathrm{Cat}_\infty$, and let I be a small ∞ -category. The composite*

$$C \xrightarrow{F} \mathrm{Cat}_\infty \xrightarrow{\mathrm{Fun}(I, -)} \mathrm{Cat}_\infty$$

is classified by the cocartesian fibration

$$C \times_{\mathrm{Fun}(I, C)} \mathrm{Fun}(I, E) \longrightarrow C,$$

where the map $\mathrm{Fun}(I, E) \rightarrow \mathrm{Fun}(I, C)$ is induced by p and the map from C sends an object to its constant I -diagram.

Proof. Cocartesian lifts for $\mathrm{Fun}(I, E) \rightarrow \mathrm{Fun}(I, C)$ are formed pointwise, so this map is a cocartesian fibration. The displayed pullback is therefore a cocartesian fibration by Lemma 23.1.12. Its fiber over $x \in C$ is

$$\mathrm{Fun}(I, E) \times_{\mathrm{Fun}(I, C)} \{\mathrm{const}_x\} \simeq \mathrm{Fun}(I, E_x) \simeq \mathrm{Fun}(I, F(x)).$$

Under these equivalences, cocartesian transport along a morphism $f: x \rightarrow y$ is postcomposition with the transport functor $f_!: E_x \rightarrow E_y$. Its straightening is therefore the stated composite. $\square$

Example 23.2.3. When $C = [0]$, the last statement of the theorem implies that for a cocartesian fibration $p: E \rightarrow C$ the ∞ -categories $\mathrm{Str}^{\mathrm{cc}}(p)(x)$ for $x \in C$ are precisely the fibers of p .

Example 23.2.4. An illustrative example is (un)straightening over the walking morphism $C = [1]$. Consider first a cocartesian fibration $p: E \rightarrow [1]$, with E small, and let E_0 and E_1 denote its fibers over 0 and 1, respectively. Cocartesian transport along the canonical morphism in $[1]$ results in a functor $F: E_0 \rightarrow E_1$, which corresponds to a morphism in Cat_∞ , i.e. a functor $[1] \rightarrow \mathrm{Cat}_\infty$. This is $\mathrm{Str}^{\mathrm{cc}}(p)$.

Conversely, consider a functor $F: E_0 \rightarrow E_1$ between small ∞ -categories. Regarding this as a morphism in Cat_∞ , its unstraightening takes the form of a cocartesian fibration $E \rightarrow [1]$. Since the fibers over 0 and 1 are E_0 and E_1 , there are inclusions $i_0: E_0 \hookrightarrow E$ and $i_1: E_1 \hookrightarrow E$. Moreover, by choosing coherent choices of cocartesian lifts, there is a natural transformation $\alpha: i_0 \Rightarrow i_1 \circ F$ of functors $E_0 \rightarrow E$. All in all, we have constructed a commutative square in Cat_∞ as follows:

$$\begin{array}{ccc} E_0 \times \{1\} & \hookrightarrow & E_0 \times [1] \\ & & \downarrow \alpha \\ E_1 & \xrightarrow{i_1} & E. \end{array}$$

It can be shown that this square is a pushout square in Cat_∞ , see for example [Lur09b, Section 3.2.2].

Let us now state some immediate consequences of the theorem.

Corollary 23.2.5. *The straightening equivalences Str^{cc} and Str^{ct} restrict to equivalences*

$$\text{Str}^{\text{cc}}: \text{LFib}(C) \xrightarrow{\sim} \text{Fun}(C, \text{An}) \quad \text{and} \quad \text{Str}^{\text{ct}}: \text{RFib}(C) \xrightarrow{\sim} \text{Fun}(C^{\text{op}}, \text{An}).$$

Proof. Given a cocartesian fibration $p: E \rightarrow C$, the individual values of $\text{Str}^{\text{cc}}(p)$ are precisely the fibers of p . In particular, $\text{Str}^{\text{cc}}(p)$ lands in the subcategory $\text{An} \subseteq \text{Cat}_{\infty}$ if and only if all fibers are anima. By Proposition 23.1.16, this is equivalent to p being a left fibration, proving the first equivalence. The dual discussion provides the right equivalence. $\square$

Corollary 23.2.6. *Let X be an anima. Then there is an equivalence of ∞ -categories*

$$\text{Str}: \text{An}/X \xrightarrow{\sim} \text{Fun}(X, \text{An}).$$

Proof. If C is an anima, then every functor $E \rightarrow C$ is both a cocartesian and a cartesian fibration. Moreover, it is a left/right fibration if and only if also E is an anima. The claim now becomes an instance of Corollary 23.2.5. $\square$

Notation 23.2.7. We will denote the inverses to the straightening functors Str^{cc} and Str^{ct} by

$$\text{Un}^{\text{cc}}: \text{Fun}(C, \text{Cat}_{\infty}) \xrightarrow{\sim} \text{Cocart}(C) \quad \text{and} \quad \text{Un}^{\text{ct}}: \text{Fun}(C^{\text{op}}, \text{Cat}_{\infty}) \xrightarrow{\sim} \text{Cart}(C),$$

and refer to these as (cocartesian/cartesian) *unstraightening*. Given a functor $F: C \rightarrow \text{Cat}_{\infty}$, we may describe its unstraightening $\text{Un}^{\text{cc}}(F)$ informally as follows:

- An object in $\text{Un}^{\text{cc}}(F)$ is a pair (x, a) , where x is an object of C and a is an object of $F(x)$;
- A morphism in $\text{Un}^{\text{cc}}(F)$ from (x, a) to (y, b) is a pair (f, φ) , where $f: x \rightarrow y$ is a morphism in C and $\varphi: f_!(a) \rightarrow b$ is a morphism in $F(y)$. Here we write $f_!$ for the functor $F(f): F(x) \rightarrow F(y)$ induced by f using the functoriality of F .
- Composition of $(f, \varphi): (x, a) \rightarrow (y, b)$ and $(g, \psi): (y, b) \rightarrow (z, c)$ is defined as $(g \circ f, \psi \circ g_!(\varphi))$.

The functor $\text{Un}^{\text{cc}}(F) \rightarrow C$ is given by $(x, a) \mapsto x$ and $(f, \varphi) \mapsto f$. A dual description holds for $\text{Un}^{\text{ct}}(F)$.

Lemma 23.2.8 (Products in cocartesian unstraightenings). *Let $q: E \rightarrow B$ be a cocartesian fibration, where B admits finite products. If the straightening*

$$\text{Str}^{\text{cc}}(q): B \rightarrow \text{Cat}_{\infty}$$

preserves finite products, then E admits finite products and q preserves them. More precisely, let $X_a \in E_{b_a}$ be a finite family of objects and let $b := \prod_a b_a$. Under the equivalence

$$E_b \simeq \prod_a E_{b_a},$$

let $X \in E_b$ correspond to the tuple $(X_a)_a$. Then the q -cocartesian lifts $X \rightarrow X_a$ of the product projections $b \rightarrow b_a$ exhibit X as the product $\prod_a X_a$ in E .

Proof. Let $Z \in E$ and write $c := q(Z)$. The asserted product cone induces a commutative square

$$\begin{array}{ccc} \mathrm{Hom}_E(Z, X) & \longrightarrow & \prod_a \mathrm{Hom}_E(Z, X_a) \\ \downarrow & & \downarrow \\ \mathrm{Hom}_B(c, b) & \xrightarrow{\sim} & \prod_a \mathrm{Hom}_B(c, b_a). \end{array}$$

For a morphism $f: c \rightarrow b$, the map induced on the fibers over f and its components identifies with

$$\mathrm{Hom}_{E_b}(f_! Z, X) \longrightarrow \prod_a \mathrm{Hom}_{E_{b_a}}((\mathrm{pr}_a f)_! Z, X_a).$$

This is an equivalence because, under the product equivalence $E_b \simeq \prod_a E_{b_a}$, the objects $f_! Z$ and X correspond to the tuples $((\mathrm{pr}_a f)_! Z)_a$ and $(X_a)_a$, respectively. Thus the top map in the square is an equivalence, proving that $X \simeq \prod_a X_a$. The image of this product cone under q is the chosen product cone in B , so q preserves finite products. $\square$

Corollary 23.2.9 (Coproducts in cartesian unstraightenings). *Let $q: E \rightarrow B$ be a cartesian fibration, where B admits finite coproducts. If the straightening*

$$\mathrm{Str}^{\mathrm{ct}}(q): B^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}$$

preserves finite products, then E admits finite coproducts and q preserves them. More precisely, if $X_a \in E_{b_a}$ and $X \in E_{\coprod_a b_a}$ corresponds to $(X_a)_a$ under the equivalence

$$E_{\coprod_a b_a} \simeq \prod_a E_{b_a},$$

then the q -cartesian lifts $X_a \rightarrow X$ of the coproduct inclusions exhibit X as the coproduct $\coprod_a X_a$ in E .

Proof. Apply Lemma 23.2.8 to $q^{\mathrm{op}}: E^{\mathrm{op}} \rightarrow B^{\mathrm{op}}$. $\square$

Remark 23.2.10. In Theorem 23.2.1 we only formulated a ‘morphismwise’ version of functoriality of the straightening equivalences. Let us remark on how to formulate a fully functorial version.

For this, we will need to assume the existence of an even bigger universe $\widehat{\mathrm{Cat}}_{\infty}$ of large ∞ -categories, and assume that Cat_{∞} is an object of $\widehat{\mathrm{Cat}}_{\infty}$, cf. Remark 1.8.1. Applying cartesian straightening to the cartesian fibration $t: \mathrm{Ar}(\mathrm{Cat}_{\infty}) \rightarrow \mathrm{Cat}_{\infty}$ results in a contravariant functor

$$\mathrm{Cat}_{\infty}^{\mathrm{op}} \rightarrow \widehat{\mathrm{Cat}}_{\infty}, \quad C \mapsto (\mathrm{Cat}_{\infty})_{/C}, \quad (F: C \rightarrow D) \mapsto (F^*: (\mathrm{Cat}_{\infty})_{/D} \rightarrow (\mathrm{Cat}_{\infty})_{/C}).$$

Since cocartesian fibrations are closed under pullback, this restricts to a functor

$$\mathrm{Cocart}(-): \mathrm{Cat}_{\infty}^{\mathrm{op}} \rightarrow \widehat{\mathrm{Cat}}_{\infty}.$$

Similarly, the assignment $C \mapsto \mathrm{Fun}(C, \mathrm{Cat}_{\infty})$ defines a functor

$$\mathrm{Fun}(-, \mathrm{Cat}_{\infty}): \mathrm{Cat}_{\infty}^{\mathrm{op}} \rightarrow \widehat{\mathrm{Cat}}_{\infty},$$

where functoriality is given by precomposition. The fully coherent version of the straightening equivalence would now ask for a natural equivalence

$$\mathrm{Str}^{\mathrm{cc}}: \mathrm{Cocart}(-) \xrightarrow{\sim} \mathrm{Fun}(-, \mathrm{Cat}_{\infty})$$

of functors $\mathrm{Cat}_{\infty}^{\mathrm{op}} \rightarrow \widehat{\mathrm{Cat}}_{\infty}$.

Remark 23.2.11. Recall that if two functors $F: C \rightarrow D$ and $G: D \rightarrow E$ admit right adjoints F^R and G^R , then GF admits a right adjoint $F^R G^R$. Using straightening/unstraightening, we can define a highly coherent version of this functoriality of passing to right adjoints. Consider the wide subcategories

$$\mathrm{Cat}_{\infty}^{\mathrm{L}} \subseteq \mathrm{Cat}_{\infty} \quad \text{and} \quad \mathrm{Cat}_{\infty}^{\mathrm{R}} \subseteq \mathrm{Cat}_{\infty}$$

denote the wide subcategories spanned by the left adjoints and right adjoints, respectively. We claim that passing to right adjoints defines a functor

$$(-)^{\mathrm{R}}: (\mathrm{Cat}_{\infty}^{\mathrm{L}})^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}^{\mathrm{R}}.$$

To this end, choose a bigger universe $\widehat{\mathrm{Cat}}_{\infty}$ containing Cat_{∞} as an object. The inclusion $\mathrm{Cat}_{\infty}^{\mathrm{L}} \hookrightarrow \widehat{\mathrm{Cat}}_{\infty}$ can then be straightened to a cocartesian fibration $E \rightarrow \mathrm{Cat}_{\infty}^{\mathrm{L}}$. Given a morphism $F: C \rightarrow D$ in $\mathrm{Cat}_{\infty}^{\mathrm{L}}$, the induced cocartesian transport map $E_C \rightarrow E_D$ on fibers is precisely the functor F itself. Since by assumption F admits a right adjoint, it follows from Lemma 23.1.20 that this cocartesian fibration is also a cartesian fibration, hence straightens to a functor $(-)^{\mathrm{R}}: (\mathrm{Cat}_{\infty}^{\mathrm{L}})^{\mathrm{op}} \rightarrow \widehat{\mathrm{Cat}}_{\infty}$. By checking on objects and morphisms, this factors through $\mathrm{Cat}_{\infty}^{\mathrm{R}}$.

23.3 Fiberwise criteria

We obtain the following useful consequence of straightening-unstraightening:

Theorem 23.3.1. *Consider a cocartesian functor*

$$\begin{array}{ccc} E & \xrightarrow{F} & E' \\ & \searrow p & \swarrow p' \\ & C, & \end{array}$$

where p and p' are cocartesian fibrations between small ∞ -categories. Then F is an equivalence of ∞ -categories if and only if for every object x of C , the induced functor on fibers $F_x: E_x \rightarrow E'_x$ is an equivalence.

Proof. Under the equivalence $\mathrm{Str}^{\mathrm{cc}}: \mathrm{Cocart}(C) \xrightarrow{\sim} \mathrm{Fun}(C, \mathrm{Cat}_{\infty})$, F corresponds to a natural transformation $\mathrm{Str}^{\mathrm{cc}}(F): \mathrm{Str}^{\mathrm{cc}}(p) \rightarrow \mathrm{Str}^{\mathrm{cc}}(p')$, and this natural transformation is a natural equivalence if and only if it is a pointwise equivalence. $\square$

Corollary 23.3.2. *A cocartesian fibration $p: E \rightarrow C$ is an equivalence if and only if all of its fibers are contractible.*

Proof. Apply the previous theorem to $E' = C$. $\square$

Corollary 23.3.3. *A morphism of anima $p: X \rightarrow Y$ is an equivalence if and only if all its fibers are contractible.*

Proof. Since X is an anima, every morphism in X is an isomorphism, and in particular p -cocartesian. It follows that p is a left fibration, and thus the claim holds by the previous corollary. $\square$

Proposition 23.3.4 (Fiberwise initial objects in cartesian fibrations, [Lur09b, Proposition 2.4.4.9]). *Let $p: E \rightarrow C$ be a cartesian fibration whose fibers admit initial objects. Then the fiberwise initial objects assemble uniquely into a fully faithful left-adjoint section*

$$s: C \rightarrow E$$

of p . Dually, the fiberwise terminal objects of a cocartesian fibration assemble uniquely into a fully faithful right-adjoint section.

Proof. For every $x \in C$, choose an initial object $e_x \in E_x$. We claim that e_x , together with the identity morphism $x \rightarrow p(e_x) = x$, is a left adjoint object to x under p . Indeed, for any $e \in E$ consider the map

$$\mathrm{Hom}_E(e_x, e) \longrightarrow \mathrm{Hom}_C(x, p(e)).$$

For a morphism $f: x \rightarrow p(e)$, choose a p -cartesian lift $f^*e \rightarrow e$. The defining property of this lift identifies the fiber of the displayed map over f with

$$\mathrm{Hom}_{E_x}(e_x, f^*e),$$

which is contractible because e_x is initial. It follows from Corollary 23.3.3 that $\mathrm{Hom}_E(e_x, e) \rightarrow \mathrm{Hom}_C(x, p(e))$ is an equivalence. This proves the claim.

The pointwise criterion for adjunctions from Lemma 21.1.4 now assembles the objects e_x and the identity morphisms $x \rightarrow p(e_x)$ into a left adjoint $s: C \rightarrow E$ whose unit $\mathrm{id}_C \rightarrow ps$ is a natural isomorphism. Hence s is a fully faithful section of p . Any other section selecting initial objects in the fibers is a left adjoint to p by the same argument, and is therefore unique. The final statement follows by duality. $\square$

Lemma 23.3.5. *Let $p: E \rightarrow C$ be a functor. Then a morphism $\varphi: e \rightarrow e'$ in E is p -cocartesian if and only if the commutative square*

$$\begin{array}{ccc} E_{e'/|} & \xrightarrow{-\circ\varphi} & E_{e/|} \\ p \downarrow & & \downarrow p \\ C_{pe'/|} & \xrightarrow{-\circ p\varphi} & C_{pe/|} \end{array}$$

is a pullback square. A dual criterion holds for p -cartesian morphisms.

Proof. The horizontal maps are left fibrations, hence in particular cocartesian fibrations. It follows from closure under base change that also the map from the pullback to $E_{e/|}$ is a cocartesian fibration. By Theorem 23.3.1, the functor $E_{e'/|} \rightarrow E_{e/|} \times_{C_{pe/|}} C_{pe'/|}$ is an equivalence if and only if this holds fiberwise over $E_{e/|}$, which means that for every morphism $\psi: e \rightarrow e''$ the map

$$\mathrm{Hom}_{E_{e/|}}(\varphi, \psi) \rightarrow \mathrm{Hom}_{C_{pe/|}}(p\varphi, p\psi)$$

is an equivalence. Since this is a morphism of anima, this may be checked fiberwise, where it precisely becomes the definition of cocartesianness for φ . $\square$

Lemma 23.3.6. *Let $q: E \rightarrow D$ and $p: D \rightarrow C$ be functors and assume that p is a left fibration. Then q is a left fibration if and only if pq is a left fibration.*

Proof. As a consequence of Lemma 23.3.5, we see that a morphism $\varphi: e \rightarrow e'$ in E is q -cocartesian if and only if it is pq -cocartesian: in the commutative diagram

$$\begin{array}{ccc} E_{e'/|} & \xrightarrow{-\circ\varphi} & E_{e/|} \\ q \downarrow & & \downarrow q \\ D_{qe'/|} & \xrightarrow{-\circ q\varphi} & D_{qe/|} \\ p \downarrow & & \downarrow p \\ C_{pqe'/|} & \xrightarrow{-\circ pq\varphi} & C_{pqe/|}, \end{array}$$

the bottom square is a pullback square, hence the top square is a pullback square if and only if the bottom one is. It follows that every morphism in E is q -cocartesian if and only if every morphism is pq -cocartesian. The claim follows. $\square$

Proposition 23.3.7. *For a functor $p: E \rightarrow C$, the following conditions are equivalent:*

- (1) *The functor p is a left fibration;*
- (2) *For every object e of E , the induced functor $p: E_{e/|} \rightarrow C_{pe/|}$ is an equivalence;*
- (3) *The commutative square*

$$\begin{array}{ccc} \text{Ar}(E) & \xrightarrow{p_*} & \text{Ar}(C) \\ s \downarrow & & \downarrow s \\ E & \xrightarrow{p} & C \end{array}$$

is a pullback square.

Proof. For (1) $\Rightarrow$ (2), assume that p is a left fibration. Since $E_{e/|} \rightarrow E$ and $C_{pe/|} \rightarrow C$ are left fibrations as well, it follows from Lemma 23.3.6 that also $E_{e/|} \rightarrow C_{pe/|}$ is a left fibration, hence it is an equivalence if and only if each of its fibers is contractible. But the fiber over $f \in C_{pe/|}$ is the anima $L(e, f)$ of p -cocartesian lifts of f starting in e , hence this is contractible by Exercise 23.1.7.

For (2) $\Rightarrow$ (1), first note that it follows from (2) that every morphism $pe \rightarrow c'$ has a lift to some morphism $e \rightarrow e'$. It remains to show that every morphism $\varphi: e \rightarrow e'$ in E is p -cocartesian. But this is a direct consequence of Lemma 23.3.5: the two vertical maps in the square are equivalences, hence the square is a pullback square.

For (2) $\iff$ (3), note that the square is a pullback square if and only if the horizontal map in the following diagram is an equivalence:

$$\begin{array}{ccc} \text{Ar}(E) & \xrightarrow{(s, p_*)} & E \times_C \text{Ar}(C) \\ s \searrow & & \swarrow \text{pr} \\ & E. & \end{array}$$

By the dual of Example 23.1.10, the two vertical functors are cartesian fibrations. In light of Theorem 23.3.1, the horizontal functor is an equivalence if and only if it induces equivalences on

fibers over each object $e \in E$. Unwinding definitions, these functors on fibers are precisely the functors $E_{e/} \rightarrow C_{pe/}$ from condition (2). $\square$

23.4 The Hom-functor

Last semester, we had taken it as a black box that every ∞ -category C admits a Hom-functor

$$\mathrm{Hom}_C : C^{\mathrm{op}} \times C \rightarrow \mathrm{An}.$$

Using the straightening/unstraightening correspondence, we will now finally construct this functor.

Lemma 23.4.1. *Let C be an ∞ -category. Then the functor*

$$\begin{array}{ccc} \mathrm{Ar}(C) & \xrightarrow{(s,t)} & C \times C \\ & \searrow t \quad \swarrow \mathrm{pr}_2 & \\ & C & \end{array}$$

is a cocartesian functor over C .

Proof. Recall from Example 23.1.10 that the target functor $t : \mathrm{Ar}(C) \rightarrow C$ is a cocartesian fibration whose cocartesian morphisms are those squares inverted by $s : \mathrm{Ar}(C) \rightarrow C$. The projection functor pr_2 is cocartesian by Exercise 23.1.13, and its cocartesian morphisms are those pairs $(g : X \rightarrow X', f : Y \rightarrow Y')$ such that g is an isomorphism. It follows immediately from these descriptions that (s, t) is a cocartesian functor. $\square$

Construction 23.4.2 (The Hom-functor). Let C be a small ∞ -category. We construct a functor $\mathrm{Hom}_C : C^{\mathrm{op}} \times C \rightarrow \mathrm{An}$.

By Lemma 23.4.1, the functor $(s, t) : \mathrm{Ar}(C) \rightarrow C \times C$ defines a morphism in $\mathrm{Cocart}(C)$. Applying the cocartesian straightening functor, we thus obtain a morphism

$$\mathrm{Str}^{\mathrm{cc}}(t) \rightarrow \mathrm{Str}^{\mathrm{cc}}(\mathrm{pr}_2)$$

in $\mathrm{Fun}(C, \mathrm{Cat}_\infty)$. Now, note that pr_2 is the pullback along $p_C : C \rightarrow *$ of the map $p_C : C \rightarrow *$, and it follows from the naturality of straightening that $\mathrm{Str}^{\mathrm{cc}}(\mathrm{pr}_2)$ is given by the composite

$$C \xrightarrow{p_C} * \xrightarrow{\mathrm{Str}(p_C)} \mathrm{Cat}_\infty.$$

Since $\mathrm{Str}(p_C)$ is simply the map picking out the object $C \in \mathrm{Cat}_\infty$, this means that $\mathrm{Str}^{\mathrm{cc}}(\mathrm{pr}_2)$ is the constant functor $\mathrm{const}_C : C \rightarrow \mathrm{Cat}_\infty$. In particular, the map $\mathrm{Str}^{\mathrm{cc}}(t) \rightarrow \mathrm{Str}^{\mathrm{cc}}(\mathrm{pr}_2) \cong \mathrm{const}_C$ gives rise to an object

$$\mathrm{Str}^{\mathrm{cc}}(t) \in \mathrm{Fun}(C, \mathrm{Cat}_\infty)_{/\mathrm{const}_C} \simeq \mathrm{Fun}(C, (\mathrm{Cat}_\infty)_/C),$$

i.e., we may treat it as a functor $C \rightarrow (\mathrm{Cat}_\infty)_/C$. We now claim that this functor factors through the full subcategory $\mathrm{RFib}(C) \subseteq (\mathrm{Cat}_\infty)_/C$. We may check this on objects: we have to show that

for every object $x \in C$, the functor $\mathrm{Str}^{\mathrm{cc}}(t)(x) \rightarrow \mathrm{Str}^{\mathrm{cc}}(\mathrm{pr}_2)(x)$ is a right fibration. By naturality of $\mathrm{Str}^{\mathrm{cc}}(-)$, this evaluation is computed by forming the induced map on fibers over x :

$$\begin{array}{ccccc} C_{/x} & \xrightarrow{s} & C & \xrightarrow{\quad} & C \times C \\ & \searrow & \swarrow & \nearrow (s,t) & \swarrow \\ & & \mathrm{Ar}(C) & & \\ & \searrow & \swarrow & \nearrow & \swarrow \\ & & * & \xrightarrow{x} & C. \end{array}$$

But this functor $s: C_{/x} \rightarrow C$ is indeed a right fibration by Proposition 23.1.19. We thus obtain a functor

$$C \rightarrow \mathrm{RFib}(C) \xrightarrow[\sim]{\mathrm{Str}^{\mathrm{ct}}} \mathrm{Fun}(C^{\mathrm{op}}, \mathrm{An}).$$

Uncurrying then gives the desired functor $\mathrm{Hom}_C: C^{\mathrm{op}} \times C \rightarrow \mathrm{An}$.

Remark 23.4.3. There is an asymmetry in our definition of Hom_C : we could also have considered the *cartesian* functor

$$\begin{array}{ccc} \mathrm{Ar}(C) & \xrightarrow{(s,t)} & C \times C \\ & \searrow s & \swarrow \mathrm{pr}_1 \\ & & C \end{array}$$

to get a functor $C^{\mathrm{op}} \rightarrow \mathrm{LFib}(C) \simeq \mathrm{Fun}(C, \mathrm{An})$. The resulting two functors can be shown to agree with each other: more generally, for *bifibrations* $E \rightarrow C \times D$ the two ways of constructing a functor $C^{\mathrm{op}} \times D \rightarrow \mathrm{An}$ agree, see [HHLN23b, Proposition 6.18].

23.5 Limits and colimits of ∞ -categories

As a consequence of straightening/unstraightening, we obtain explicit formulas for limits and colimits in Cat_{∞} .

Construction 23.5.1. Let $p: E \rightarrow I$ be a cocartesian fibration.

- We write $E[\mathrm{cc}^{-1}]$ for the localization of E at the collection of p -cocartesian morphisms.
- We write $\Gamma_I(E)$ for the ∞ -category of *sections* of p , defined via the following pullback square:

$$\begin{array}{ccc} \Gamma_I(E) & \longrightarrow & \mathrm{Fun}(I, E) \\ \downarrow & \lrcorner & \downarrow p_* \\ * & \xrightarrow{\mathrm{id}_I} & \mathrm{Fun}(I, I). \end{array}$$

- We write $\Gamma_I^{\mathrm{cc}}(E)$ for the full subcategory of $\Gamma_I(E)$ spanned by the *cocartesian sections*: those sections

$$\begin{array}{ccc} I & \xrightarrow{s} & E \\ & \searrow & \swarrow p \\ & & I \end{array}$$

that are cocartesian functors over I (i.e., that send arbitrary morphisms in I to p -cocartesian morphisms in E).

Proposition 23.5.2. *Let I be a small ∞ -category and let $F: I \rightarrow \text{Cat}_\infty$ be a functor.*

(1) *The functor F admits a colimit in Cat_∞ , which can be computed by the formula*

$$\text{colim}_I F(i) \simeq \text{Un}^{\text{cc}}(F)[\text{cc}^{-1}].$$

(2) *The functor F admits a limit in Cat_∞ , which can be computed by the formula*

$$\lim_I F(i) \simeq \Gamma_I^{\text{cc}}(\text{Un}^{\text{cc}}(F)).$$

Proof. (1) We need to show that there exists a natural transformation $\eta_F: F \rightarrow \text{const}_{\text{Un}^{\text{cc}}(F)[\text{cc}^{-1}]}$ of functors $I \rightarrow \text{Cat}_\infty$ which exhibits $\text{Un}^{\text{cc}}(F)[\text{cc}^{-1}]$ as a left adjoint object to F under the functor

$$\text{const}: \text{Cat}_\infty \rightarrow \text{Fun}(I, \text{Cat}_\infty).$$

By straightening/unstraightening, this functor agrees with the functor

$$\text{Cat}_\infty \simeq \text{Cocart}(\ast) \xrightarrow{p_I^\ast} \text{Cocart}(I), \quad C \mapsto (\text{pr}_I: C \times I \rightarrow I)$$

given by pullback along $p_I: I \rightarrow \ast$. A transformation of the form η_F corresponds to a cocartesian functor over I of the form

$$\begin{array}{ccc} \text{Un}^{\text{cc}}(F) & \xrightarrow{\bar{\eta}} & \text{Un}^{\text{cc}}(F)[\text{cc}^{-1}] \times I \\ & \searrow p_F & \swarrow \text{pr}_I \\ & I & \end{array}$$

We define $\bar{\eta} := (\gamma, p_F)$, where $\gamma: \text{Un}^{\text{cc}}(F) \rightarrow \text{Un}^{\text{cc}}(F)[\text{cc}^{-1}]$ is the localization functor. Since the cocartesian morphisms of pr_I are those whose component in $\text{Un}^{\text{cc}}(F)[\text{cc}^{-1}]$ is an equivalence (see Exercise 23.1.13), $\bar{\eta}$ is indeed a cocartesian functor over I . To show that it exhibits $\text{Un}^{\text{cc}}(F)[\text{cc}^{-1}]$ as a left adjoint object, we must show that for any other ∞ -category C the induced map

$$\text{Hom}_{\text{Cat}_\infty}(\text{Un}^{\text{cc}}(F)[\text{cc}^{-1}], C) \xrightarrow{(- \circ \bar{\eta}) \circ p_I^\ast} \text{Hom}_{\text{Cocart}(I)}(\text{Un}^{\text{cc}}(F), C \times I) \quad (23.1)$$

is an equivalence. For this, we compare this map with the composite

$$\text{Hom}_{\text{Cat}_\infty}(\text{Un}^{\text{cc}}(F), C) \xrightarrow{(- \circ (\text{id}, p_F)) \circ p_I^\ast} \text{Hom}_{(\text{Cat}_\infty)_I}(\text{Un}^{\text{cc}}(F), C \times I),$$

which is an equivalence due to the fact that the forgetful functor $(\text{Cat}_\infty)_I \rightarrow \text{Cat}_\infty$ is left adjoint to $p_I^\ast: \text{Cat}_\infty \rightarrow (\text{Cat}_\infty)_I$. But then also the composite (23.1) is a composite: its source is the full subanima of $\text{Hom}_{\text{Cat}_\infty}(\text{Un}^{\text{cc}}(F), C)$ spanned by those functors $\varphi: \text{Un}^{\text{cc}}(F) \rightarrow C$ that invert the cocartesian morphisms, which is equivalent to the condition that the functor $(\varphi, p_F): \text{Un}^{\text{cc}}(F) \rightarrow C \times I$ preserves cocartesian morphisms. This finishes the proof for colimits.

(2) Similar to the case of colimits, we may translate the problem into finding a cocartesian functor

$$\begin{array}{ccc} \Gamma_I^{\text{cc}}(\text{Un}^{\text{cc}}(F)) \times I & \xrightarrow{\bar{\varepsilon}} & \text{Un}^{\text{cc}}(F) \\ & \searrow \text{pr}_I & \swarrow p_F \\ & I & \end{array}$$

over I that exhibits $\Gamma_I^{\text{cc}}(\text{Un}^{\text{cc}}(F))$ as a right adjoint to $\text{Un}^{\text{cc}}(F)$ under the pullback functor $p_I^*: \text{Cat}_\infty \rightarrow \text{Cocart}(I)$. By definition of $\Gamma_I(\text{Un}^{\text{cc}}(F))$, the evaluation functor $\text{Fun}(I, \text{Un}^{\text{cc}}(F)) \times I \rightarrow \text{Un}^{\text{cc}}(F)$ restricts to a functor

$$\begin{array}{ccc} \Gamma_I(\text{Un}^{\text{cc}}(F)) \times I & \xrightarrow{\text{ev}} & \text{Un}^{\text{cc}}(F) \\ & \searrow \text{pr}_I \quad \swarrow p_F & \\ & C & \end{array}$$

over C . We claim that this exhibits $\Gamma_I(\text{Un}^{\text{cc}}(F))$ as a right adjoint object to $\text{Un}^{\text{cc}}(F)$ under the functor $\text{Cat}_\infty \rightarrow (\text{Cat}_\infty)_I$: for every other ∞ -category C we have

$$\begin{aligned} \text{Hom}_{\text{Cat}_\infty}(C, \Gamma_I(\text{Un}^{\text{cc}}(F))) &\simeq \text{fib}(\text{Hom}_{\text{Cat}_\infty}(C, \text{Fun}(I, \text{Un}^{\text{cc}}(F))) \rightarrow \text{Hom}_{\text{Cat}_\infty}(C, \text{Fun}(I, I))) \\ &\simeq \text{fib}(\text{Hom}_{\text{Cat}_\infty}(C \times I, \text{Un}^{\text{cc}}(F)) \rightarrow \text{Hom}_{\text{Cat}_\infty}(C \times I, I)) \\ &\simeq \text{Hom}_{(\text{Cat}_\infty)_I}(C \times I, \text{Un}^{\text{cc}}(F)). \end{aligned}$$

By defining $\bar{e}$ as the restriction of the evaluation functor, we may immediately deduce that also the functor

$$\text{Hom}_{\text{Cat}_\infty}(C, \Gamma_I^{\text{cc}}(\text{Un}^{\text{cc}}(F))) \xrightarrow{(\bar{e} \circ -) \circ p_I^*} \text{Hom}_{\text{Cocart}(I)}(C \times I, \text{Un}^{\text{cc}}(F))$$

is an equivalence: a functor $\varphi: C \rightarrow \Gamma_I(\text{Un}^{\text{cc}}(F))$ lands in the subcategory of cocartesian sections if and only if the associated functor $C \times I \rightarrow \text{Un}^{\text{cc}}(F)$ over I is a cocartesian functor. This finishes the proof for limits. $\square$

Corollary 23.5.3. *Let I be a small ∞ -category and let $F: I \rightarrow \text{An}$ be functor.*

(1) *The functor F admits a colimit in An , which can be computed by the formula*

$$\text{colim}_I F(i) \simeq \|\text{Un}^{\text{cc}}(F)\|.$$

(2) *The functor F admits a limit in An , which can be computed by the formula*

$$\lim_I F(i) \simeq \Gamma_I(\text{Un}^{\text{cc}}(F)).$$

Proof. We compute the limit and colimit in Cat_∞ and observe that it is still in An . For (1) this is clear: all morphisms in $\text{Un}^{\text{cc}}(F)$ are cocartesian, see Proposition 23.1.16, hence inverting all cocartesian morphisms gives an anima. For (2), any section $s: I \rightarrow \text{Un}^{\text{cc}}(F)$ of p_F is cocartesian, again since all morphisms are cocartesian, so $\Gamma_I^{\text{cc}}(\text{Un}^{\text{cc}}(F)) = \Gamma_I(\text{Un}^{\text{cc}}(F))$. Note that this is already an anima: any morphism of sections is invertible since all fibers of $p_F: \text{Un}^{\text{cc}}(F) \rightarrow I$ are anima. $\square$

We will now investigate the behavior of limits of ∞ -categories in more detail.

Proposition 23.5.4 (Limits of adjunctions). *Let K be a small ∞ -category and consider functors $C_\bullet, D_\bullet: K \rightarrow \text{Cat}_\infty$. Let $F_\bullet: C_\bullet \rightarrow D_\bullet$ be a natural transformation satisfying the following two conditions:*

- (1) *(Pointwise adjoints) The functor $F_k: C_k \rightarrow D_k$ admits a right adjoint $G_k: D_k \rightarrow C_k$ for every $k \in K$;*

(2) (*Beck-Chevalley condition*) For every morphism $f: k \rightarrow k'$ in K , the Beck-Chevalley transformation $\text{BC}: f_* G_k \rightarrow G_{k'} f_*$, defined as the composite

$$f_* G_k \xrightarrow{\text{unit}} G_{k'} F_{k'} f_* G_k \cong G_{k'} f_* F_k G_k \xrightarrow{\text{counit}} G_{k'} f_*,$$

is a natural isomorphism.

Then the functors G_k assemble into a natural transformation $G_\bullet: D_\bullet \rightarrow C_\bullet$, and the units and counits assemble into natural transformations $\varepsilon_\bullet: F_\bullet \circ G_\bullet \rightarrow \text{id}_\bullet$ and $\eta_\bullet: \text{id}_\bullet \rightarrow G_\bullet \circ F_\bullet$ that satisfy the triangle identities.

As a result, the induced functors

$$F := \lim_k F_k: \lim_k C_k \rightarrow \lim_k D_k \quad \text{and} \quad G := \lim_k G_k: \lim_k D_k \rightarrow \lim_k C_k$$

are adjoint to each other.

Proof. We may interpret the transformation $F_\bullet$ as a functor $K \rightarrow \text{Fun}([1], \text{Cat}_\infty)$, $k \mapsto (F_k: C_k \rightarrow D_k)$. Using the equivalence $\text{Fun}([1], \text{Cat}_\infty) \simeq \text{CoCart}([1])$, each F_k corresponds to a cocartesian fibration $E_k \rightarrow [1]$. The assumption that F_k admits a right adjoint G_k implies that $E_k \rightarrow [1]$ is also a cartesian fibration; see Corollary 23.1.21. The Beck-Chevalley condition is precisely the condition that the induced functor $E_k \rightarrow E_{k'}$ for $k \rightarrow k'$ in K is a cartesian functor over $[1]$. So conditions (1) and (2) guarantee that the composite $K \rightarrow \text{Fun}([1], \text{Cat}_\infty) \simeq \text{CoCart}([1]) \subseteq (\text{Cat}_\infty)_{/[1]}$ lands in the subcategory $\text{Cart}([1]) \simeq \text{Fun}([1]^{\text{op}}, \text{Cat}_\infty) \simeq \text{Fun}([1], \text{Cat}_\infty)$. This results in a natural transformation $G_\bullet: D_\bullet \rightarrow C_\bullet$ which is pointwise for $k \in K$ given by forming the right adjoint G_k of F_k .

The constructions of $\varepsilon_\bullet$ and $\eta_\bullet$ is similar. Since formation of cartesian/cocartesian lifts is functorial in the morphism (see Corollary 23.1.5), there is a functorial way to extract from a cartesian fibration $E \rightarrow [1]$ the cartesian lifts $\text{can}^* x \rightarrow x$ of the canonical morphism $\text{can}: 0 \rightarrow 1$ in $[1]$. This morphism in turn factors functorially through the cocartesian lift $\text{can}^* x \rightarrow \text{can}_! \text{can}^* x$, providing a morphism $\text{can}_! \text{can}^* x \rightarrow x$ which is functorial in the pair $(E, x \in E_1)$. An analogous construction gives $\eta_\bullet$, and the triangle identities. We leave the details to the interested reader.

For the final claim, passing to limits produces the functors F and G , and natural transformations $\varepsilon: FG \rightarrow \text{id}$ and $\eta: \text{id} \rightarrow GF$ satisfying the triangle identities. This shows that G is right adjoint to F . $\square$

Corollary 23.5.5. *Let I and K be a small ∞ -category and let $C_\bullet: K \rightarrow \text{Cat}_\infty$ be a K -indexed diagram of small ∞ -categories. Assume that C_k admits I -indexed limits (resp. colimits) for all $k \in K$, and that the functor $C_k \rightarrow C_{k'}$ preserves I -indexed limits (resp. colimits) for every morphism $k \rightarrow k'$ in K . Then the limit $C := \lim_{k \in K} C_k$ in Cat_∞ also admits I -indexed limits (resp. colimits) and each functor $C \rightarrow C_k$ preserves them.*

Proof. We prove the claim about limits; the claim for colimits is dual. By the description of limits in terms of an adjunction on functor categories from Corollary 21.2.2, we need to show that the constant diagram functor $\lim_{k \in K} C_k \rightarrow \text{Fun}(I, \lim_{k \in K} C_k)$ admits a right adjoint. Since $\text{Fun}(I, -): \text{Cat}_\infty \rightarrow \text{Cat}_\infty$ is right adjoint to $I \times -$, it preserves limits, giving an equivalence

$$\text{Fun}(I, \lim_{k \in K} C_k) \simeq \lim_{k \in K} \text{Fun}(I, C_k).$$

The resulting map $\lim_{k \in K} C_k \rightarrow \lim_{k \in K} \text{Fun}(I, C_k)$ is a limit of the constant diagram functors $C_k \rightarrow \text{Fun}(I, C_k)$. The conditions on the C_k then precisely amount to saying that each of these functors admit right adjoints, and that the Beck-Chevalley conditions are satisfied, so the claim follows from the previous proposition. $\square$

23.6 Hom animae in functor categories

We may use the results from the previous section to compute the hom animae in a functor category $\text{Fun}(C, D)$. For this, we introduce the notion of an *end* in the ∞ -categorical setting.

Definition 23.6.1 (Twisted arrow category). Let C be an ∞ -category. We define its *twisted arrow category* $\text{Tw}(C)$ as the source of the left fibration whose cocartesian straightening is the Hom-functor $\text{Hom}_C: C^{\text{op}} \times C \rightarrow \text{An}$:

$$\text{Tw}(C) := \text{Un}^{\text{cc}}(\text{Hom}_C: C^{\text{op}} \times C \rightarrow \text{An}).$$

In particular, it comes equipped with a left fibration $(s, t): \text{Tw}(C) \rightarrow C^{\text{op}} \times C$ whose fibers are the hom animae of C :

$$\begin{array}{ccc} \text{Hom}_C(x, y) & \longrightarrow & \text{Tw}(C) \\ \downarrow & \lrcorner & \downarrow (s, t) \\ * & \xrightarrow{(x, y)} & C^{\text{op}} \times C. \end{array}$$

Unwinding the description of the unstraightening construction, we see that $\text{Tw}(C)$ may be described as follows:

- The objects of $\text{Tw}(C)$ are morphisms in C , which we will display as $\begin{pmatrix} X \\ \downarrow \varphi \\ Y \end{pmatrix}$. We have $s\left(\begin{pmatrix} X \\ \downarrow \varphi \\ Y \end{pmatrix}\right) = X$ and $t\left(\begin{pmatrix} X \\ \downarrow \varphi \\ Y \end{pmatrix}\right) = Y$.
- A morphism $\begin{pmatrix} f \\ g \end{pmatrix}: \begin{pmatrix} X \\ \downarrow \varphi \\ Y \end{pmatrix} \rightarrow \begin{pmatrix} X' \\ \downarrow \varphi' \\ Y' \end{pmatrix}$ in $\text{Tw}(C)$ is a triple (f, g, α) consisting of a morphism $f: X \rightarrow X'$ in C^{op} (i.e., $f: X' \rightarrow X$ in C), a morphism $g: Y \rightarrow Y'$ in C , and a morphism $\alpha: (f, g)! \begin{pmatrix} X \\ \downarrow \varphi \\ Y \end{pmatrix} = \begin{pmatrix} X' \\ \downarrow \varphi' \\ Y' \end{pmatrix}$ in the fiber $\text{Tw}(C)_{(X', Y')} = \text{Hom}_C(X', Y')$. We will display this data by means of diagrams of the form

$$\begin{array}{ccc} X & \xleftarrow{f} & X' \\ \varphi \downarrow & & \downarrow \varphi' \\ Y & \xrightarrow{g} & Y'. \end{array}$$

Note that this is *not* a morphism in the usual arrow category $\text{Ar}(C)$, since f points in the opposite direction.

Definition 23.6.2 (End). Let C be an ∞ -category and consider a functor $F: C^{\text{op}} \times C \rightarrow D$. The *end* of F , denoted by $\int_{c \in C} F(c, c)$ if it exists, is the limit of the composite functor

$$\text{Tw}(C) \xrightarrow{(s, t)} C^{\text{op}} \times C \xrightarrow{F} D.$$

Remark 23.6.3. If C is small and D admits small limits, the end of F always exists. Furthermore, if $D \rightarrow D'$ is a functor preserving small limits, then it also preserves ends.

Lemma 23.6.4 (Fubini rule for ends). *Let C and C' be ∞ -categories and let $F: C^{\text{op}} \times C \times C'^{\text{op}} \times C' \rightarrow D$ be a functor. Then there is an equivalence*

$$\int_{c \in C} \int_{c' \in C'} F(c, c, c', c') \simeq \int_{(c, c') \in C \times C'} F(c, c, c', c').$$

Proof. Since a limit over a product category may be computed as an iterated limit, it will suffice to produce an equivalence

$$\text{Tw}(C \times C') \simeq \text{Tw}(C) \times \text{Tw}(C')$$

of left fibrations over $C^{\text{op}} \times C \times C'^{\text{op}} \times C'$. By straightening-unstraightening, it will suffice to produce a natural isomorphism between their unstraightenings:

$$(x, y, x', y') \mapsto \text{Hom}_{C \times C'}((x, x'), (y, y')) \simeq \text{Hom}_C(x, y) \times \text{Hom}_{C'}(x', y').$$

Unwinding the definition of the hom functors, this boils down to the canonical equivalence

$$\text{Ar}(C \times C') \xrightarrow{\sim} \text{Ar}(C) \times \text{Ar}(C')$$

over $C \times C \times C' \times C'$, coming from the fact that $\text{Ar}(-) = \text{Fun}([1], -)$ preserves products. $\square$

Proposition 23.6.5 (Hom animae in functor categories). *Let $F, G: C \rightarrow D$ be functors. Then we have the following end-formula for the anima of natural transformations from F to G :*

$$\text{Nat}(F, G) \simeq \int_{x \in C} \text{Hom}_D(F(x), G(x)).$$

Here the right-hand side is the end of the composite functor $C^{\text{op}} \times C \xrightarrow{F^{\text{op}} \times G} D^{\text{op}} \times D \xrightarrow{\text{Hom}_D} \text{An}$.

Proof. By definition, the end on the right-hand side is a certain limit in An over the twisted arrow category $\text{Tw}(C)$. As shown in Corollary 23.5.3, this limit is equivalent to the anima of sections of the unstraightening of the composite $\text{Tw}(C) \rightarrow \text{An}$. By definition, the unstraightening of Hom_D is the twisted arrow category $\text{Tw}(D) \rightarrow D^{\text{op}} \times D$. The unstraightening of the composite $\text{Tw}(C) \rightarrow \text{An}$ is thus given by pulling back $\text{Tw}(D)$ along $F^{\text{op}} \times G$ and (s, t) :

$$\begin{array}{ccccccc} P' & \xrightarrow{\quad} & P & \xrightarrow{\quad} & \text{Tw}(D) = \text{Un}(\text{Hom}_D) \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow (s, t) \\ \text{Tw}(C) & \xrightarrow{(s, t)} & C^{\text{op}} \times C & \xrightarrow{F^{\text{op}} \times G} & D^{\text{op}} \times D & \xrightarrow{\text{Hom}_D} & \text{An}. \end{array}$$

From the left pullback square it in particular follows that the sections of the map $P' \rightarrow \text{Tw}(C)$ are equivalent to maps $\text{Tw}(C) \rightarrow P$ over $C^{\text{op}} \times C$. Since both sides define left fibrations over $C^{\text{op}} \times C$, the anima of such maps over $C^{\text{op}} \times C$ is equivalent to the anima of natural transformations between their straightenings, which for $\text{Tw}(C)$ is Hom_C and for P is $\text{Hom}_D(F(-), G(-))$. Let us summarize what we have so far:

$$\begin{aligned} \int_{x \in C} \text{Hom}_D(F(x), G(x)) &= \lim \left(\text{Tw}(C) \xrightarrow{(s, t)} C^{\text{op}} \times C \xrightarrow{F^{\text{op}} \times G} D^{\text{op}} \times D \xrightarrow{\text{Hom}_D} \text{An} \right) \\ &\simeq \Gamma_{\text{Tw}(C)}(P') = \text{Hom}_{\text{LFib}(\text{Tw}(C))}(\text{Tw}(C), P') \\ &\simeq \text{Hom}_{\text{LFib}(C^{\text{op}} \times C)}(\text{Tw}(C), P) \\ &\simeq \text{Hom}_{\text{Fun}(C^{\text{op}} \times C, \text{An})}(\text{Hom}_C, \text{Hom}_D \circ (F^{\text{op}} \times G)). \end{aligned}$$

Now, under the equivalence $\text{Fun}(C^{\text{op}} \times C, \text{An}) \simeq \text{Fun}(C, \text{PSh}(C))$, the functor Hom_C corresponds to the Yoneda embedding $Y^C: C \hookrightarrow \text{Fun}(C^{\text{op}}, \text{An})$, while $\text{Hom}_D \circ (F^{\text{op}} \times G)$ corresponds to the composite

$$C \xrightarrow{G} D \xrightarrow{Y^D} \text{PSh}(D) \xrightarrow{F^*} \text{PSh}(C).$$

The hom anima we need to compute is thus given by

$$\text{Hom}_{\text{Fun}(C, \text{PSh}(C))}(Y^C, F^* \circ Y^D \circ G).$$

The functor $F^*: \text{PSh}(D) \rightarrow \text{PSh}(C)$ is restriction along $F^{\text{op}}: C^{\text{op}} \rightarrow D^{\text{op}}$. Since $\text{PSh}(D)$ is cocomplete, Theorem 21.4.3 gives it a left adjoint $F_!$, defined by left Kan extension along F^{op} . The Yoneda lemma identifies $F_!(Y^C(c))$ with $Y^D(F(c))$, naturally in c , so that $F_!$ fits in a commutative diagram

$$\begin{array}{ccc} C & \xrightarrow{F} & D \\ Y^C \downarrow & & \downarrow Y^D \\ \text{PSh}(C) & \xrightarrow{F_!} & \text{PSh}(D). \end{array}$$

This provides equivalences

$$\begin{aligned} \text{Hom}_{\text{Fun}(C, \text{PSh}(C))}(Y^C, F^* \circ Y^D \circ G) &\simeq \text{Hom}_{\text{Fun}(C, \text{PSh}(D))}(F_! \circ Y^C, Y^D \circ G) \\ &\simeq \text{Hom}_{\text{Fun}(C, \text{PSh}(D))}(Y^D \circ F, Y^D \circ G) \\ &\simeq \text{Hom}_{\text{Fun}(C, D)}(F, G) = \text{Nat}(F, G), \end{aligned}$$

where the last equivalence follows from full faithfulness of the Yoneda embedding Y^D . $\square$

Proposition 23.6.6. *Let $F: I \rightarrow C$ be a functor of ∞ -categories. A natural transformation $\eta: F \rightarrow \text{const}_x$ exhibits $x \in C$ as the colimit of F if and only if the natural map*

$$\eta^*: \text{Hom}_C(x, y) \xrightarrow{\sim} \lim_{i \in I^{\text{op}}} \text{Hom}_C(F(i), y)$$

obtained by applying $\text{Hom}_C(-, y)$ to η is an equivalence for all $y \in C$. A similar assertion holds for limits.

Proof. Consider the following commutative diagram:

$$\begin{array}{ccccc} \text{Hom}_C(x, y) & \xrightarrow{\text{const}} & \text{Nat}(\text{const}_x, \text{const}_y) & \xrightarrow{- \circ \eta} & \text{Nat}(F, \text{const}_y) \\ & \searrow \eta^* & & & \downarrow \\ & & & & \lim_{i \in I^{\text{op}}} \text{Hom}_C(F(i), y); \end{array}$$

here the right vertical map is induced by the evaluation functors $\text{ev}_i: \text{Fun}(I, C) \rightarrow C$. By definition, x is a colimit of F if and only if the top map is an equivalence for all y , hence it will suffice to show that the right vertical map is an equivalence. By Proposition 23.6.5, the top right is equivalent to $\lim_{(i \rightarrow i') \in \text{Tw}(I)} \text{Hom}_C(F(i), y)$. It thus remains to show that the forgetful functor $s: \text{Tw}(I) \rightarrow I^{\text{op}}$ is final. By Theorem 21.5.1, it suffices to show that the relative slices are weakly contractible. But these relative slices are weakly equivalent [explain]²² to the actual slices $I_{i/}$, which have initial objects, so the claim follows from Lemma 21.5.5. $\square$

²²There is an adjunction $\text{Tw}(I)_{i/} \rightleftarrows I_{i/}$, and we have Lemma 21.1.7

Corollary 23.6.7. *The functors $Y_C : C \hookrightarrow \text{Fun}(C^{\text{op}}, \text{An})$ and $Y_{C^{\text{op}}} : C^{\text{op}} \hookrightarrow \text{Fun}(C, \text{An})$ preserve limits.*

Proof. The statement for $Y_{C^{\text{op}}}$ is immediate from Proposition 23.6.6. The other one is proved dually, or may be recovered by replacing C by C^{op} . $\square$

23.7 Descent and ∞ -topoi

An important property that distinguishes the ∞ -category An from other ∞ -categories is a property called *descent*. We have already encountered this property in the proof of the Recognition Principle (Theorem 5.2.5). In this section, we give a self-contained treatment.

Definition 23.7.1 (Descent). Let I be a small ∞ -category, and let T be an ∞ -category with pullbacks and I -indexed colimits. We say that T *satisfies descent for I -indexed colimits* if the functor

$$T^{\text{op}} \rightarrow \text{Cat}, \quad X \mapsto T/X$$

preserves I -indexed limits.

Definition 23.7.2 (∞ -Topos). An ∞ -topos is a presentable ∞ -category satisfying descent for all small colimits.

Let us unwind Definition 23.7.1. Given a small ∞ -category I and an I -indexed diagram $X_\bullet : I \rightarrow T$, set $X := \text{colim}_i X_i$. The descent condition demands that the canonical functor $T/X \rightarrow \lim_i T/X_i$, obtained from the base change functors along the maps $X_i \rightarrow X$, is an equivalence.

Theorem 23.7.3. *The ∞ -category An is an ∞ -topos.*

Proof. We need to show that for any functor $X_\bullet : I \rightarrow \text{An}$, with colimit $X_\infty := \text{colim}_i X_i$, the functor $\text{An}/X_\infty \rightarrow \lim_i \text{An}/X_i$ is an equivalence. By Corollary 23.2.6, we have an equivalence $\text{An}/Y \xrightarrow{\sim} \text{Fun}(Y, \text{An})$ which is natural in Y . So, the claim is equivalent to showing that the map

$$\text{Fun}(X_\infty, \text{An}) \rightarrow \lim_i \text{Fun}(X_i, \text{An})$$

is an equivalence. But this follows from the fact that $\text{Fun}(-, \text{An}) : \text{An}^{\text{op}} \rightarrow \text{Cat}$ sends colimits in An to limits in Cat . $\square$

To understand the descent condition better, we provide an alternative description of the limit $\lim_i T/X_i$ in terms of cartesian natural transformations.

Definition 23.7.4 (Cartesian natural transformation). Given two functors $F, G : I \rightarrow T$, a natural transformation $\alpha : F \rightarrow G$ is called *cartesian* if for every morphism $i \rightarrow j$ in I the naturality square

$$\begin{array}{ccc} F(i) & \xrightarrow{\alpha(i)} & G(i) \\ \downarrow & \lrcorner & \downarrow \\ F(j) & \xrightarrow{\alpha(j)} & G(j) \end{array}$$

is a pullback square. We denote by $\text{Fun}^{\text{cart}}(I, T) \subseteq \text{Fun}(I, T)$ the wide subcategory spanned by the cartesian natural transformations.

Given a diagram $X_\bullet: I \rightarrow T$, evaluation at $i \in I$ determines a functor $\text{ev}_i: \text{Fun}^{\text{cart}}(I, T)_{/X_\bullet} \rightarrow T_{/X_i}$. Furthermore, by definition of cartesianness these evaluation functors are compatible with base change, resulting in a comparison functor $\text{Fun}^{\text{cart}}(I, T)_{/X_\bullet} \rightarrow \lim_{i \in I^{\text{op}}} T_{/X_i}$.

Lemma 23.7.5. *Let C be an ∞ -category with pullbacks and let $X_\bullet: I \rightarrow C$ be a functor. Then the functor $\text{Fun}^{\text{cart}}(I, C)_{/X_\bullet} \rightarrow \lim_{i \in I^{\text{op}}} C_{/X_i}$ is an equivalence.*

More informally, objects of the limit are given by I -indexed diagrams $Y_\bullet$ in C equipped with a cartesian natural transformation $Y_\bullet \rightarrow X_\bullet$.

Proof. Recall that a limit of a diagram $I^{\text{op}} \rightarrow \text{Cat}$ of ∞ -categories may be computed as the ∞ -category of cartesian sections of the cartesian unstraightening of the diagram. By definition, the functor $C_{/-}: C^{\text{op}} \rightarrow \text{Cat}$ is the cartesian straightening of the target map $t: \text{Ar}(C) \rightarrow C$, so the unstraightening of $C_{/-}$ is t . It follows that the unstraightening of the functor $i \mapsto C_{/X_i}$ is the base change of t along $X_\bullet$:

$$\begin{array}{ccc} C/X_\bullet & \longrightarrow & \text{Ar}(C) \\ \downarrow & \lrcorner & \downarrow t \\ I & \xrightarrow{X_\bullet} & C. \end{array}$$

The ∞ -category of sections $I \rightarrow C/X_\bullet$ of this map is equivalent to the ∞ -category of maps $I \rightarrow \text{Ar}(C)$ whose target projection is $X_\bullet: I \rightarrow C$, which in turn is equivalent to the slice- ∞ -category $\text{Fun}(I, C)_{/X_\bullet}$. It thus remains to show that a section $s: I \rightarrow C/X_\bullet$ is cartesian precisely if the corresponding natural transformation $Y_\bullet \rightarrow X_\bullet$ is a cartesian natural transformation. But this is an immediate consequence of the fact that the t -cartesian morphisms in $\text{Ar}(C)$ are precisely the pullback squares in C . $\square$

With this alternative description of $\lim_i T_{/X_i}$ at hand, we can show that the functor $T_{/X} \rightarrow \lim_i T_{/X_i}$ admits a left adjoint:

Lemma 23.7.6. *Let T be an ∞ -category with pullbacks and I -indexed colimits, and let $X = \text{colim}_i X_i$ be a colimit in T . Then the functor $T_{/X} \rightarrow \lim_i T_{/X_i}$ admits a left adjoint*

$$\text{colim}: \lim_i T_{/X_i} \rightarrow T_{/X}$$

sending a cartesian transformation $Y_\bullet \rightarrow X_\bullet$ to its colimit $\text{colim}_i Y_i \rightarrow \text{colim}_i X_i$.

Proof. The colimit functor $\text{colim}: \text{Fun}(I, T) \rightarrow T$ admits a right adjoint $\text{const}: T \rightarrow \text{Fun}(I, T)$ sending an object Y to the constant functor on Y . By slicing this adjunction over an arbitrary diagram $X_\bullet \in \text{Fun}(I, T)$, with colimit denoted X , we obtain another adjunction

$$\text{colim}: \text{Fun}(I, T)_{/X_\bullet} \rightleftarrows T_{/X},$$

see [Lur09b, Proposition 5.2.5.1]. The right adjoint in this adjunction is given by the composite $T_{/X} \xrightarrow{\text{const}} \text{Fun}(I, T)_{/\text{const } X} \rightarrow \text{Fun}(I, T)_{/X_\bullet}$, with the second map given by base change along the colimit cocone $X_\bullet \rightarrow \text{const}_X$. We now make the following two observations:

- The ∞ -category $\text{Fun}(I, T)_{/X_\bullet}$ contains the limit $\lim_{i \in I^{\text{op}}} C_{/X_i} \simeq \text{Fun}^{\text{cart}}(I, T)_{/X_\bullet}$ as a full subcategory: given transformations $Y_\bullet \rightarrow Z_\bullet \rightarrow X_\bullet$, if both $Z_\bullet \rightarrow X_\bullet$ and $Y_\bullet \rightarrow X_\bullet$ are cartesian transformations, then so is $Y_\bullet \rightarrow Z_\bullet$ by the pasting law for pullback squares.

- Since any transformation $\text{const } Y \rightarrow \text{const } X$ between constant diagrams is cartesian, and cartesian transformations are closed under base change, the right adjoint $T_{/X} \rightarrow \text{Fun}(I, T)_{/X_\bullet}$ lands in this full subcategory.

It follows that the adjunction restricts to an adjunction $\lim_i T_{/X_i} \rightleftarrows T_{/X}$, as desired. $\square$

We can now characterize when the functor $T_{/X} \rightarrow \lim_i T_{/X_i}$ is an equivalence. By Lemma 23.7.6, this happens precisely when the unit and counit of the adjunction are natural isomorphisms.

Lemma 23.7.7. *Let T be an ∞ -category with pullbacks and I -indexed colimits, and let $X_\bullet : I \rightarrow T$ be a diagram with colimit $X := \text{colim}_i X_i$. Then T satisfies descent for I -indexed colimits if and only if the following two conditions hold:*

- (1) *For any map $Y \rightarrow X$, the canonical map $\text{colim}_i (X_i \times_X Y) \rightarrow Y$ is an isomorphism.*
- (2) *Given a cartesian transformation $Y_\bullet \rightarrow X_\bullet$, each map $Y_i \rightarrow (\text{colim}_j Y_j) \times_X X_i$ is an isomorphism.*

Proof. By definition, T satisfies descent for I -indexed colimits if and only if the canonical functor $T_{/X} \rightarrow \lim_i T_{/X_i}$ is an equivalence. By Lemma 23.7.6, this functor admits a left adjoint given by sending a cartesian transformation $Y_\bullet \rightarrow X_\bullet$ to its colimit. The functor is an equivalence if and only if both the unit and counit of this adjunction are isomorphisms. The counit at $Y \rightarrow X$ is precisely the map in (1), while the unit at a cartesian transformation $Y_\bullet \rightarrow X_\bullet$ consists of the maps in (2). $\square$

23.7.1 Examples of descent

Let us illustrate the descent condition by giving some examples.

Example 23.7.8 (Strictly initial object). Let T be an ∞ -category with an initial object. Then the initial object $\emptyset$ satisfies descent if and only if it is *strictly initial*, meaning that any morphism $X \rightarrow \emptyset$ is an isomorphism.

Indeed, by definition $\emptyset$ satisfies descent if and only if the functor $T_{/\emptyset} \rightarrow *$ is an equivalence. This functor admits a fully faithful left adjoint $* \rightarrow T_{/\emptyset}$ hitting the identity $\text{id}_\emptyset$. For the functor to be an equivalence, the counit $\emptyset \rightarrow X$ must be an isomorphism for every $X \in T_{/\emptyset}$, which is precisely the strict initiality condition.

Example 23.7.9 (Disjoint coproducts). For an ∞ -category T with arbitrary coproducts, we say that *coproducts are disjoint* if for all $X, Y \in T$ the diagram

$$\begin{array}{ccc} \emptyset & \longrightarrow & Y \\ \downarrow & & \downarrow \\ X & \longrightarrow & X \sqcup Y \end{array}$$

is a pullback square. We claim that arbitrary coproducts in T satisfy descent if and only if they are universal and T has disjoint coproducts.

By Lemma 23.7.7, coproducts satisfy descent if and only if for every collection of maps $\{Y_i \rightarrow X_i\}_{i \in I}$ the commutative square

$$\begin{array}{ccc} Y_j & \longrightarrow & Y := \coprod_{i \in I} Y_i \\ \downarrow & & \downarrow \\ X_j & \longrightarrow & X := \coprod_{i \in I} X_i \end{array}$$

is a pullback square for every $j \in I$. To see that this reduces to disjointness, consider $X' := \coprod_{i \in I \setminus \{j\}} X_i$ and $Y' := \coprod_{i \in I \setminus \{j\}} Y_i$. We may write $X = X_j \sqcup X'$ and $Y = Y_j \sqcup Y'$. Invoking universality of coproducts once more shows that the condition reduces to disjointness of binary coproducts.

Example 23.7.10 (Grothendieck construction). Let T be an ∞ -topos. Given an anima $A \in \mathbf{An}$, we obtain an equivalence

$$\mathrm{Fun}(A, T) = \lim_A T_{/*} \simeq T_{/\mathrm{colim}_A *}.$$

We refer to this equivalence as the *Grothendieck construction*.

23.8 Exponentiability of cartesian fibrations

We will need one further formal property of cartesian fibrations. Recall that a morphism $p: E \rightarrow B$ in an ∞ -category with finite products is called *exponentiable* if the functor $- \times_B E$ admits a right adjoint. In the present context we will only use the following consequence, which says that base change along a cartesian fibration behaves like tensoring with a fixed object.

Proposition 23.8.1 (Exponentiability of cartesian fibrations, [Lur17, Appendix B.3]). *Let $p: E \rightarrow B$ be a cartesian fibration. Then the functor*

$$- \times_B E: (\mathrm{Cat}_\infty)_{/B} \rightarrow (\mathrm{Cat}_\infty)_{/B}$$

preserves colimits. Dually, if p is a cocartesian fibration, then $- \times_B E$ preserves colimits as well.

Proof. This is Lurie’s flatness theorem for cartesian fibrations; see also [AF20]. $\square$

Remark 23.8.2. The proposition is a genuine property of cartesian fibrations, not a formal property of arbitrary maps. For example, let $\ell: [1] \rightarrow [2]$ be the long edge. The category $[2]$ is the pushout of the spine

$$[1] \sqcup_{[0]} [1],$$

where the two copies of $[1]$ meet in the middle vertex. Pulling this pushout back along ℓ gives two discrete vertices, whereas the pullback of $[2]$ along ℓ is $[1]$. Thus $- \times_{[2]} [1]$ does not preserve this pushout.

Theorem 23.8.3 (Currying for cartesian fibrations). *Let $p: E \rightarrow B$ be a cartesian fibration and let C be an ∞ -category. Write*

$$A := \mathrm{Str}^{\mathrm{ct}}(p): B^{\mathrm{op}} \rightarrow \mathrm{Cat}_\infty.$$

Thus, for a morphism $\beta: b \rightarrow b'$ in B , the induced functor is the cartesian transport functor $\beta^: E_{b'} \rightarrow E_b$. Let*

$$H: B \rightarrow \mathrm{Cat}_\infty, \quad b \mapsto \mathrm{Fun}(E_b, C), \quad \beta \mapsto (\beta^*)^*$$

and let $q: Q := \text{Un}^{\text{cc}}(H) \rightarrow B$ be its cocartesian unstraightening. Then there exists a functor

$$\text{ev}: Q \times_B E \rightarrow C$$

which restricts on the fiber over $b \in B$ to the usual evaluation functor

$$\text{Fun}(E_b, C) \times E_b \rightarrow C$$

and which has the following universal property: for every object $D \rightarrow B$ of $(\text{Cat}_\infty)_/B$, the composite

$$\text{Fun}_/B(D, Q) \xrightarrow{- \times_B E} \text{Fun}_/B(D \times_B E, Q \times_B E) \xrightarrow{\text{ev} \circ -} \text{Fun}(D \times_B E, C)$$

is an equivalence of ∞ -categories.

Proof. This is the dual of Gepner, Haugseng, and Nikolaus [GHN17, Proposition 7.3]. That result identifies the internal hom obtained by exponentiating along a cocartesian fibration as the cartesian unstraightening of the fiberwise functor categories. Passing to opposite categories gives the present statement for a cartesian fibration: the right adjoint to $- \times_B E$ is the cocartesian unstraightening of

$$b \mapsto \text{Fun}(E_b, C),$$

with transport given by precomposition with cartesian transport in E . Its counit is the displayed evaluation functor, and the adjunction gives the asserted equivalence. $\square$

Exercises

Exercise 23.1. Let $p: D \rightarrow C$ and $q: E \rightarrow D$ be two cocartesian fibrations.

(1) Show that $pq: E \rightarrow C$ is again a cocartesian fibration.

(2) Show that the functor p is a cocartesian functor over C .

Exercise 23.2. Let S be a small ∞ -category, and let $\alpha: F \Rightarrow G$ be a natural transformation between functors $F, G: S \rightarrow \text{Cat}_\infty$. Assume that for every object $s \in S$ the functor

$$\alpha_s: F(s) \rightarrow G(s)$$

is a left fibration. Show that the induced functor on cocartesian unstraightenings

$$\text{Un}^{\text{cc}}(\alpha): \text{Un}^{\text{cc}}(F) \rightarrow \text{Un}^{\text{cc}}(G)$$

is a left fibration. More explicitly, identify the fiber over an object $(s, g) \in \text{Un}^{\text{cc}}(G)$ with the fiber of α_s over g , and describe the transport along a morphism $(s, g) \rightarrow (t, g')$ in $\text{Un}^{\text{cc}}(G)$.

Exercise 23.3. Let C be an ∞ -category and consider $p := t: \text{Ar}(C) \rightarrow C$.

(1) Show that a commutative square

$$\begin{array}{ccc} X & \longrightarrow & X' \\ \downarrow & & \downarrow \\ Y & \xrightarrow{f} & Y' \end{array}$$

is p -cocartesian (when regarded as a morphism in $\text{Ar}(C)$ from $\begin{pmatrix} X \\ \downarrow \\ Y \end{pmatrix}$ to $\begin{pmatrix} X' \\ \downarrow \\ Y' \end{pmatrix}$) if and only if the map $X \rightarrow X'$ is an isomorphism.

- (2) Conclude that p is always a cocartesian fibration.
- (3) Show that a commutative square like the one above is p -cartesian if and only if it is a pullback square in C . (Since pullback squares are also called ‘cartesian squares’, this explains the name ‘cartesian morphisms’).
- (4) Conclude that p is a cartesian fibration if and only if C admits pullbacks.

Dualize this statement by showing that s is always a cartesian fibration and that it is additionally a cocartesian fibration if and only if C admits pushouts.

Exercise 23.4. Consider the category $\mathbf{VectBund}$ whose objects are pairs (X, E) consisting of a topological space X and a real vector bundle $p: E \rightarrow X$, and whose morphisms are commutative squares

$$\begin{array}{ccc} E & \xrightarrow{F} & E' \\ p \downarrow & & \downarrow p' \\ X & \xrightarrow{f} & X' \end{array}$$

of topological spaces such that F induces $\mathbb{R}$ -linear maps $E_x \rightarrow E'_{f(x)}$ on fibers. Show that the forgetful functor $\mathbf{VectBund} \rightarrow \mathbf{Top}, (X, E) \mapsto X$ is a cartesian fibration.

Exercise 23.5. Let $p: D \rightarrow C$ and $q: E \rightarrow D$ be two cocartesian fibrations.

- (1) Show that $pq: E \rightarrow C$ is again a cocartesian fibration.
- (2) Show that the functor q is a cocartesian functor over C .

Exercise 23.6. Consider a pullback square of ∞ -categories of the form

$$\begin{array}{ccc} E' & \xrightarrow{F} & E \\ p' \downarrow & \lrcorner & \downarrow p \\ C' & \longrightarrow & C, \end{array}$$

where p is a cocartesian fibration. Show that p' is a cocartesian fibration as well, and a morphism $f: x \rightarrow y$ in E' is a p' -cocartesian morphism if and only if $G(f)$ is a p -cocartesian morphism.

Exercise 23.7. For a functor $p: E \rightarrow C$, consider the full subcategory

$$\{p\text{-cocart}\} \subseteq \mathbf{Ar}(E)$$

spanned by the p -cocartesian morphisms.

- (1) Show that the functor

$$\{p\text{-cocart}\} \rightarrow E \times_{p, C, s} \mathbf{Ar}(C), \quad (\varphi: e \rightarrow e') \mapsto (e, p\varphi)$$

is fully faithful.

- (2) Deduce that p is a cocartesian fibration if and only if this functor is an equivalence.

- (3) Use this equivalence to show that for every morphism $f: x \rightarrow y$, the assignment $e \mapsto f_! e$ from Notation 23.1.3 can be enhanced to a cocartesian transport functor

$$f_!: E_x \rightarrow E_y.$$

Exercise 23.8. A morphism $f: Y \rightarrow X$ of anima is called a *covering* if all its fibers are sets, i.e., if all their higher homotopy groups vanish. Letting $\text{Cov}(X) \subseteq \text{An}/_X$ denote the full subcategory spanned by the coverings, deduce that the straightening equivalence induces an equivalence

$$\text{Cov}(X) \xrightarrow{\sim} \text{Fun}(\Pi_1(X), \text{Set})$$

between the category of covering spaces over X and the category of functors $\Pi_1(X) \rightarrow \text{Set}$. Explain how this provides a high-tech version of the classical classification of covering spaces.

Exercise 23.9. Let I and C ∞ -categories, and let $F = \text{const}_C: I \rightarrow \text{Cat}_\infty$ be the constant functor with value C .

- Show that the colimit of F is the product $C \times \|I\|$, where $\|I\| := C[\text{all}^{-1}]$ is the geometric realization of I (obtained by inverting arbitrary morphisms).
- Show that the limit of F is $\text{Fun}(\|I\|, C)$.
- Conclude that for an anima X we have $\text{colim}_X * \simeq X$.

Exercise 23.10. Use the formula for colimits in Cat_∞ to compute the pushout in Cat_∞ of the following diagram:

$$\begin{array}{ccc} * & \xrightarrow{0} & [1] \\ \downarrow 1 & & \\ [1] & & \end{array}$$

(You may take for granted that for a 1-category I the unstraightening of a functor $I \rightarrow \text{Cat}_\infty$ that happens to land in the subcategory $\text{Cat}_1 \subseteq \text{Cat}_\infty$ of 1-categories is again a 1-category, computed by the Grothendieck construction.)

24 Complete Segal animae

Constructing ∞ -categories is more subtle than constructing classical categories: it does not suffice to define composition of morphisms by hand and check unitality and associativity, due to the fact that there are higher coherences that should be encoded. In most cases, the ∞ -category one is interested in can be built up from a few elementary constructions, such as products, pullbacks, and functor categories. However, in certain cases, like that for ∞ -categories of *spans* (see Chapter 13), this is not immediately possible.

In this chapter, we will introduce an alternative method for constructing ∞ -categories, originally due to Rezk [Rez01]: by writing down a simplicial anima $\Delta^{\text{op}} \rightarrow \mathbf{An}$ satisfying certain *Segal* and *Rezk* conditions. Given an ∞ -category C , the assignment $[n] \mapsto \text{Map}([n], C)$ can be made functorial in $[n] \in \Delta^{\text{op}}$, resulting in a simplicial anima $N(C): \Delta^{\text{op}} \rightarrow \mathbf{An}$ called the *nerve* of C . It turns out that this assignment defines a fully faithful functor $N: \text{Cat}_{\infty} \rightarrow \mathbf{sAn}$ from the ∞ -category of small ∞ -categories to the ∞ -category of simplicial animae. The image of this functor consists of those simplicial animae that satisfy the Segal and Rezk conditions, which are called *complete Segal animae*. A precise formulation of this result is given in Theorem 24.1.2 below.

This theorem is powerful since it lets us manipulate or construct ∞ -categories at the level of simplicial animae. We recall the definition of complete Segal animae in Section 24.1, and use the theorem in Section 24.2 to draw some consequences regarding filtered colimits and geometric realizations of ∞ -categories. The online version collects additional proofs at the end of the chapter.

24.1 ∞ -Categories as complete Segal animae

We will now introduce the notion of complete Segal animae, which are simplicial animae that satisfy certain conditions. The motivating example of a complete Segal anima is the nerve of an ∞ -category:

Definition 24.1.1 (Nerve). The *nerve* of an ∞ -category C is the simplicial anima $N(C): \Delta^{\text{op}} \rightarrow \mathbf{An}$ defined by $N(C)_n := \text{Map}([n], C)$, i.e. as the composite

$$\Delta^{\text{op}} \hookrightarrow \text{Cat}^{\text{op}} \hookrightarrow \text{Cat}_{\infty}^{\text{op}} \xrightarrow{\text{Hom}_{\text{Cat}_{\infty}}(-, C)} \mathbf{An}.$$

This construction is functorial in C and thus defines a functor $N: \text{Cat}_{\infty} \rightarrow \mathbf{sAn}$.

We may think of points of $\text{Map}([n], C)$ as ‘strings of n composable morphisms in C ’:

- For $n = 0$, $\text{Map}([0], C) = \text{Fun}(*, C)^{\simeq} \simeq C^{\simeq}$ is just the groupoid core of C ;
- For $n = 1$, $\text{Map}([1], C)$ is the anima of morphisms in C ;

- For $n = 2$, $\text{Map}([2], C)$ is the anima of commutative triangles in C , which by the Segal axiom, Proposition 1.4.5, is equivalent to $\text{Map}([1], C) \times_{C^\approx} \text{Map}([1], C)$;
- More generally, the ‘higher version’ of the Segal condition inductively implies that for every $n \geq 0$ we have

$$\text{Map}([n], C) \xrightarrow{\sim} \text{Map}([1], C) \times_{C^\approx} \text{Map}([1], C) \times_{C^\approx} \cdots \times_{C^\approx} \text{Map}([1], C)$$

for all n .

Note that all the relevant information of C is captured by its nerve: the objects are captured by the anima $N(C)_0 = C^\approx$, the hom anima $\text{Hom}_C(x, y)$ may be recovered as the fiber over (x, y) of the map $(d_1, d_0): N(C)_1 \rightarrow N(C)_0 \times N(C)_0$, and composition in C is captured by $N(C)_2$. It thus seems plausible that C may be recovered from $N(C)$. In fact, we have the following much stronger statement:

Theorem 24.1.2 ([JT07, Section 4], [Lur09a, Corollary 4.3.16], [HS23]). *The nerve functor*

$$N: \text{Cat}_\infty \hookrightarrow \text{sAn}$$

is fully faithful. Its image consists of the complete Segal anima (see below). In particular, the nerve functor induces an equivalence

$$N: \text{Cat}_\infty \xrightarrow{\sim} \text{CSeg}(\text{An})$$

between Cat_∞ and the ∞ -category of complete Segal anima.

Most of the basic aspects of an ∞ -category C , like its composition of morphisms and its definition of isomorphisms, can more generally be phrased in terms of simplicial anima. We will now spell out the details.

Definition 24.1.3 (Segal anima). A simplicial anima $X: \Delta^{\text{op}} \rightarrow \text{An}$ is called a *Segal anima* if the Segal condition is satisfied: For every $n \geq 1$, the inclusion maps $e_i: [1] \cong \{i-1 \leq i\} \hookrightarrow [n]$ induce an isomorphism of anima

$$(e_1^*, \dots, e_n^*): X_n \xrightarrow{\sim} X_1 \times_{X_0} X_1 \times_{X_0} \cdots \times_{X_0} X_1.$$

We write $\text{Seg}(\text{An}) \subseteq \text{sAn} = \text{Fun}(\Delta^{\text{op}}, \text{An})$ for the full subcategory of Segal anima.

Observe that the nerve $N(C)$ of an ∞ -category C is always a Segal anima: this is essentially the content of the Segal Axiom 1.4.5.

One may show, just like we did for ∞ -categories in Section 1.4.2, that a Segal anima X comes equipped with a composition map

$$- \circ -: X_1 \times_{X_0} X_1 \xrightarrow{d_1} X_1$$

and that composition is unital and associative.

Convention 24.1.4. Given a Segal anima X , we refer to 0-simplices $x \in X_0$ as *objects* of X , and to 1-simplices $f \in X_1$ as *morphisms* of X . We use the notation $f: x \rightarrow y$ when $d_1(f) \simeq x$ and $d_0(f) \simeq y$. Given objects $x, y \in X_0$, we define the *hom anima* in X as the following pullback:

$$\begin{array}{ccc} \mathrm{Hom}_X(x, y) & \longrightarrow & X_1 \\ \downarrow & \lrcorner & \downarrow (d_1, d_0) \\ * & \xrightarrow{(x, y)} & X_0 \times X_0. \end{array}$$

We will sometimes refer to 2-simplices as *homotopies*.

Definition 24.1.5 (Dwyer-Kan equivalences). A morphism $f: X \rightarrow Y$ of Segal anima is called a *Dwyer-Kan equivalence* if the following two conditions are satisfied:

- (1) (Essential surjectivity): For every object $y \in Y_0$ there exists an object $x \in X_0$ satisfying $f(x) \simeq y$ in Y_0 .
- (2) (Fully faithfulness): For objects $x, x' \in X_0$, the induced map $f: \mathrm{Hom}_X(x, x') \rightarrow \mathrm{Hom}_Y(fx, fx')$ is an equivalence of anima.

Remark 24.1.6. Note that a functor $F: C \rightarrow D$ between ∞ -categories is essentially surjective (resp. fully faithful) precisely if the induced map on nerves $N(F): N(C) \rightarrow N(D)$ is essentially surjective (resp. fully faithful). In particular, F is an equivalence if and only if $N(F)$ is a Dwyer-Kan equivalence.

Definition 24.1.7. Let $X \in \mathrm{Seg}(\mathrm{An})$ be a Segal anima. A morphism $f: x \rightarrow y$ in X is called an *isomorphism* if there exist 2-simplices $\sigma, \tau \in X_2$ satisfying the relations $d_0(\sigma) \simeq f$, $d_1(\sigma) \simeq \mathrm{id}_x$, $d_2(\tau) \simeq f$ and $d_1(\tau) \simeq \mathrm{id}_y$. In other words: f is an isomorphism if there exist morphisms $g, h: y \rightarrow x$ together with homotopies $gf \simeq \mathrm{id}_x$ and $fg \simeq \mathrm{id}_y$.

We write $X_1^\sim \subseteq X_1$ for the full subanima spanned by the isomorphisms.

Definition 24.1.8 (Complete Segal anima, Rezk [Rez01]). A Segal anima $X: \Delta^{\mathrm{op}} \rightarrow \mathrm{An}$ is called *complete*, or is said to satisfy the *Rezk condition*, if the map

$$X_0 \rightarrow X_1^\sim, \quad x \mapsto \mathrm{id}_x$$

is an equivalence of anima. We denote the full subcategory of complete Segal anima by $\mathrm{CSeg}(\mathrm{An}) \subseteq \mathrm{Seg}(\mathrm{An})$.

The nerve $N(C)$ of an ∞ -category is complete by Axiom F. In particular, the nerve functor $N: \mathrm{Cat}_\infty \rightarrow \mathrm{sAn}$ factors through $\mathrm{CSeg}(\mathrm{An}) \subseteq \mathrm{sAn}$.

Additional proofs and details for the results below are given in the online version.

Lemma 24.1.9 ([HS23, Lemma 2.3]). *Let $\varphi: X \rightarrow Y$ be a morphism of complete Segal anima. Then φ is an equivalence if and only if it is a Dwyer-Kan equivalence.*

We will now assign an ∞ -category to every simplicial anima.

Lemma 24.1.10. *The nerve functor admits a left adjoint $\mathrm{ac}: \mathrm{sAn} \rightarrow \mathrm{Cat}_\infty$.*

Definition 24.1.11. Given a simplicial anima X , we refer to the ∞ -category $\mathrm{ac}(X)$ as the *associated category*.

Since the functor ac is defined in terms of left Kan extensions, the pointwise formula from Theorem 21.4.3 allows us to compute it in terms of colimits in Cat_∞ , which in turn admit an explicit description by Proposition 23.5.2. The resulting expression can then be simplified somewhat further to obtain a concrete expression for $\mathrm{ac}(X)$ as a certain localization, as explained in [HS23, Corollary 3.8]. This explicit expression can then be used to prove the following important result:

Proposition 24.1.12 (Hebestreit and Steinebrunner [HS23, Corollary 3.15, Lemma 4.1]). *For every Segal anima X , the unit map $\eta: X \rightarrow N(\mathrm{ac}(X))$ is a Dwyer-Kan equivalence.*

We refer to [HS23] for the details of the argument. Together with the results above, this supplies the main theorem stated at the start of the section.

Corollary 24.1.13. *The associated category functor $\mathrm{ac}: \mathrm{Seg}(\mathrm{An}) \rightarrow \mathrm{Cat}_\infty$ is a Bousfield localization. A morphism $\varphi: X \rightarrow Y$ of Segal anima is inverted by $\mathrm{ac}(-)$ if and only if it is a Dwyer-Kan equivalence.*

Corollary 24.1.14. *The inclusions $\mathrm{CSeg}(\mathrm{An}) \hookrightarrow \mathrm{sAn}$ and $\mathrm{CSeg}(\mathrm{An}) \hookrightarrow \mathrm{Seg}(\mathrm{An})$ admit left adjoints given by $X \mapsto N(\mathrm{ac}(X))$.*

Notation 24.1.15. The left adjoint to $\mathrm{CSeg}(\mathrm{An}) \hookrightarrow \mathrm{Seg}(\mathrm{An})$ is often called the *completion functor* or *Rezk completion* and is denoted by

$$\widehat{(-)}: \mathrm{Seg}(\mathrm{An}) \rightarrow \mathrm{CSeg}(\mathrm{An}).$$

The following is a useful consequence of the identification of Cat_∞ with the ∞ -category of complete Segal anima:

Proposition 24.1.16. *The ∞ -category Cat_∞ is generated under colimits by $[0]$ and $[1]$.*

24.2 Filtered colimits and geometric realizations of categories

The description of ∞ -categories as complete Segal anima allows us to deduce some useful statements about the interaction of finite limits and filtered colimits in Cat_∞ . It will also allow us to clarify the terminology regarding the phrase ‘geometric realization’, which we have used in two different contexts so far.

Lemma 24.2.1. *The subcategory $\mathrm{CSeg}(\mathrm{An}) \subseteq \mathrm{sAn}$ is closed under limits and filtered colimits.*

Proof. Recall that limits and colimits in sAn are computed pointwise. The conditions defining the subcategory $\mathrm{CSeg}(\mathrm{An}) \subseteq \mathrm{sAn}$ are expressed in terms of finite limits. Since finite limits commute with both limits and filtered colimits²³ in An , the claim follows. $\square$

²³See e.g. [Hau25, Section 9.9] for a ‘synthetic’ proof of the fact that finite limits commute with filtered colimits in An .

Corollary 24.2.2. *The nerve functor $N: \text{Cat}_\infty \hookrightarrow \text{sAn}$ preserves filtered colimits. In particular, finite limits commute with filtered colimits in Cat_∞ .*

Proof. The first claim is immediate from Lemma 24.2.1 and the equivalence $N: \text{Cat}_\infty \xrightarrow{\sim} \text{CSeg}(\text{An})$. The second claim follows since limits commute with filtered colimits in An and hence in sAn . $\square$

Corollary 24.2.3. *For every finite category I , the functor $\text{Hom}_{\text{Cat}_\infty}(I, -): \text{Cat}_\infty \rightarrow \text{An}$ preserves filtered colimits. In particular the core functor $(-)^{\simeq}: \text{Cat}_\infty \rightarrow \text{An}$ preserves filtered colimits.*

Proof. By definition, the subcategory $\text{Cat}_\infty^{\text{fin}} \subseteq \text{Cat}_\infty$ is the smallest subcategory closed under pushouts which contains $\emptyset$, $[0]$ and $[1]$. Since filtered colimits commute with pullbacks in An , the class I of ∞ -categories for which the claim holds is closed under pushouts. It thus remains to show the claim for $I = \emptyset, [0], [1]$. The claim for $\emptyset$ is clear, since $\text{Hom}_{\text{Cat}_\infty}(\emptyset, C)$ is terminal for each C and filtered colimits of terminal objects are terminal. The claim for $[0]$ and $[1]$ follows from the previous lemma, since $\text{Hom}_{\text{Cat}_\infty}([n], -)$ agrees with the composite $\text{Cat}_\infty \xrightarrow{N} \text{sAn} \xrightarrow{\text{ev}_n} \text{An}$.

The last claim is the special case $I = [0]$. $\square$

Corollary 24.2.4. *The ∞ -category Cat_∞ is compactly generated. Its compact objects are precisely the retracts of finite ∞ -categories.*

Proof. Every finite ∞ -category is compact by Corollary 24.2.3. By Proposition 24.1.16, every small ∞ -category is a small colimit of copies of $[0]$ and $[1]$. Writing such a colimit as a filtered colimit of its finite subcolimits expresses every small ∞ -category as a filtered colimit of finite ∞ -categories.

It remains to identify all compact objects. If C is compact, then applying compactness to a filtered-colimit presentation of C by finite ∞ -categories shows that id_C factors through one of its finite stages. Thus C is a retract of a finite ∞ -category. Conversely, every such retract is compact. The subcategory of compact objects is therefore the idempotent completion of the small ∞ -category $\text{Cat}_\infty^{\text{fin}}$, and is consequently small. $\square$

Next, we will address the terminology regarding the phrase ‘geometric realization’. Recall that we have previously used this phrase for two a priori unrelated constructions:

- For an ∞ -category C , the geometric realization $\|C\|$ is obtained from C by inverting all the morphisms in C :

$$\|C\| := C[W^{-1}], \quad W := \text{Map}([1], C);$$

- For a simplicial anima $X: \Delta^{\text{op}} \rightarrow \text{An}$, its geometric realization is its colimit:

$$|X| := \text{colim}_{[n] \in \Delta^{\text{op}}} X_n.$$

The relation between these two notions is explained by Theorem 24.1.2:

Lemma 24.2.5. *The geometric realization $\|C\|$ of an ∞ -category C is naturally equivalent to the geometric realization $|N(C)|$ of its nerve.*

Proof. The colimit-functor is by definition a left adjoint to the constant simplicial anima functor:

$$|-| : \mathbf{sAn} \rightleftarrows \mathbf{An} : \text{const}.$$

For any anima X , the constant simplicial anima $\text{const}_X : \Delta^{\text{op}} \rightarrow \mathbf{An}$ is a complete Segal anima (equivalent to the nerve $N(X)$ of X), and hence this adjunction restricts to an adjunction

$$|-| : \mathbf{CS}(\mathbf{An}) \rightleftarrows \mathbf{An} : \text{const}.$$

Under the equivalence $\mathbf{Cat}_\infty \simeq \mathbf{CS}(\mathbf{An})$ from Theorem 24.1.2, the constant functor corresponds to the inclusion $\mathbf{An} \hookrightarrow \mathbf{Cat}_\infty$, and by uniqueness of adjoints this implies the claim. $\square$

24.3 Homotopy coherent nerves

In a classical category, the collection of morphisms between two objects forms a *set*. In contrast, we have seen in Definition 1.4.17 that for ∞ -categories the morphisms between objects instead form an *anima*. As a first approximation, we may thus think of an ∞ -category as a ‘category enriched in animae’. While a formal implementation of enrichments in animae would already require a well-developed theory of ∞ -categories²⁴, we may sidestep this problem by working instead with enrichment in the 1-category of *spaces*: every topological space X admits an underlying anima $\Pi_\infty(X)$, and the functor $\Pi_\infty(-) : \mathbf{Top} \rightarrow \mathbf{An}$ preserves finite products. In this case, classical enriched category theory provides a well-defined notion of a topologically enriched category (see Definition 24.3.1), of which classical topology provides plenty of examples. It was shown by Bergner [Ber07] that the theory of topologically enriched categories provides a reasonable model for the theory of ∞ -categories.

The goal of this section is to explain how to assign an ∞ -category $N^{\text{top}}(D)$ to every topologically enriched category D , called the *homotopy coherent nerve* of D . The standard way to make this construction precise is via quasicategories, cf. Example 1.6.14. In this section, we provide an alternative approach to homotopy coherent nerves that avoids explicit models by quasicategories, fitting it in the model-agnostic approach to higher category theory taken in this book.

Definition 24.3.1. A *topologically enriched category* is a classical 1-category D together with a chosen topology on each of the hom sets $\text{Hom}_D(X, Y)$ for $X, Y \in D$, satisfying the property that the composition maps $\text{Hom}_D(Y, Z) \times \text{Hom}_D(X, Y) \rightarrow \text{Hom}_D(X, Z)$ are continuous with respect to these topologies. We denote by $\text{Hom}_D^{\text{top}}(X, Y) \in \mathbf{Top}$ the hom set in C equipped with the chosen topology.

Construction 24.3.2 (Homotopy coherent nerve). Consider a topologically enriched category D . We will define an ∞ -category $N^{\text{top}}(D)$, called the *homotopy coherent nerve* of D .

For every $[n] \in \Delta^{\text{op}}$, consider the (classical) 1-category D_n defined as follows:

- The objects of D_n are the objects of D ;
- The morphisms $X \rightarrow Y$ of D_n are the continuous maps $|\Delta^n| \rightarrow \text{Hom}_D^{\text{top}}(X, Y)$;
- The identity in D_n is the constant map $|\Delta^n| \rightarrow \text{Hom}_D^{\text{top}}(X, X)$ with value id_X ;

²⁴See [Hei23, Lemma 6.18] for a precise statement.

- The composition in D_n of morphisms $f: |\Delta^n| \rightarrow \mathrm{Hom}_D^{\mathrm{top}}(X, Y)$ and $g: |\Delta^n| \rightarrow \mathrm{Hom}_D^{\mathrm{top}}(Y, Z)$ is the composite

$$|\Delta^n| \xrightarrow{\Delta} |\Delta^n| \times |\Delta^n| \xrightarrow{g \times f} \mathrm{Hom}_D^{\mathrm{top}}(Y, Z) \times \mathrm{Hom}_D^{\mathrm{top}}(X, Y) \xrightarrow{\circ} \mathrm{Hom}_D^{\mathrm{top}}(X, Z).$$

Since $|\Delta^n|$ is functorial in $[n] \in \mathbf{\Delta}$ (see Example 1.6.8) we see that the assignment $[n] \mapsto D_n$ determines a functor

$$D_\bullet: \mathbf{\Delta}^{\mathrm{op}} \rightarrow \mathrm{Cat}_1 \hookrightarrow \mathrm{Cat}_\infty.$$

We then define the ∞ -category $N^{\mathrm{top}}(D)$ as the colimit of this functor:

$$N^{\mathrm{top}}(D) := \mathrm{colim}(D_\bullet: \mathbf{\Delta}^{\mathrm{op}} \rightarrow \mathrm{Cat}_\infty).$$

Note that D_0 is the underlying category of D . Since it admits a canonical functor to $N^{\mathrm{top}}(D)$, we see that every object of D gives rise to an object of $N^{\mathrm{top}}(D)$, and in fact all objects are of this form.

Remark 24.3.3. The fact that we refer to $N^{\mathrm{top}}(D)$ as the ‘homotopy coherent nerve’ is somewhat abusive: this terminology is often used instead for the simplicial set $N^\Delta(\mathbf{C})$ associated to a simplicially enriched category, cf. Example 1.6.14. The two constructions are closely related: given a topologically enriched category D , we may obtain a simplicially enriched category $\mathbf{C}$ by taking $\mathrm{Hom}_{\mathbf{C}}^\Delta(X, Y) := \mathrm{Sing}(\mathrm{Hom}_D^{\mathrm{top}}(X, Y))$, i.e. we apply the singular simplicial complex construction from Example 1.6.8 to each of the hom spaces of D . Then the underlying ∞ -category of the quasi-category $N^\Delta(\mathbf{C})$ is equivalent to $N^{\mathrm{top}}(D)$. The proof of this fact is quite technical: it relies on the fact that there is a model structure (due to Bergner [Ber07]) on the category of (small) simplicially enriched categories whose localization at the weak equivalences is Cat_∞ . A reference for this claim in the literature may be found in Gepner and Meier [GM23, Corollary B.4].

Proposition 24.3.4. *For objects X and Y of D , the hom anima between X and Y in $N^{\mathrm{top}}(D)$ is equivalent to the underlying anima of the topological space $\mathrm{Hom}_D^{\mathrm{top}}(X, Y)$:*

$$\mathrm{Hom}_{N^{\mathrm{top}}(D)}(X, Y) \simeq \Pi_\infty(\mathrm{Hom}_D^{\mathrm{top}}(X, Y)).$$

The proof is given in Section 24.3.2 below.

24.3.1 Proofs concerning complete Segal animae (online only)

In this online-only subsection, we discuss the proofs of several statements made above. We first compare equivalences of complete Segal animae with Dwyer-Kan equivalences. We then construct the associated category functor and use it to prove that complete Segal animae model ∞ -categories. Finally, we deduce the localization results and the generation of Cat_∞ by $[0]$ and $[1]$.

We start by showing that the equivalences between complete Segal animae are precisely the Dwyer-Kan equivalences. The key point is that completeness promotes essential surjectivity to an equivalence on the animae of objects.

Proof of Lemma 24.1.9. It is clear that every equivalence is a Dwyer-Kan equivalence. For the converse, assume φ is a Dwyer-Kan equivalence. We start by showing that φ is *conservative*,

meaning that the following square is a pullback square:

$$\begin{array}{ccc} X_1^\sim & \hookrightarrow & X_1 \\ \downarrow & & \downarrow \varphi_1 \\ Y_1^\sim & \hookrightarrow & Y_1. \end{array}$$

Let $f: x \rightarrow y$ be a morphism in X_1 and assume that its image $\varphi(f): \varphi(x) \rightarrow \varphi(y)$ is an isomorphism in Y . We have to show that already f itself is contained in $X_1^\sim$. To produce a left inverse, pick a left inverse $g': \varphi(y) \rightarrow \varphi(x)$ of $\varphi(f)$. By assumption on φ , the map $\text{Hom}_X(y, x) \xrightarrow{\sim} \text{Hom}_Y(\varphi y, \varphi x)$ is an equivalence, and in particular there exists a preimage $g: y \rightarrow x$ of g' . We claim that $gf \simeq \text{id}_x$. But since $\varphi(gf) \simeq \varphi(g)\varphi(f) \simeq g'\varphi(f) \simeq \text{id}_{\varphi(x)} \simeq \varphi(\text{id}_x)$, this follows from the equivalence $\varphi: \text{Hom}_X(x, x) \xrightarrow{\sim} \text{Hom}_Y(\varphi x, \varphi x)$. Thus g is a left inverse of f . Since $\varphi(g) = g'$ is invertible in Y , we may apply the argument to g to conclude that also g admits some left inverse h , so that g is invertible in X and hence so is f . This finishes the proof of conservativity of φ .

To show that φ is an equivalence, consider the following commutative square:

$$\begin{array}{ccccccc} X_0 & \xrightarrow[\sim]{s_0} & X_1^\sim & \hookrightarrow & X_1 & \xrightarrow{(d_1, d_0)} & X_0 \times X_0 \\ \varphi_0 \downarrow & & \downarrow \varphi_1 & & \downarrow \varphi_1 & & \downarrow \varphi_0 \times \varphi_0 \\ Y_0 & \xrightarrow[\sim]{s_0} & Y_1^\sim & \hookrightarrow & Y_1 & \xrightarrow{(d_1, d_0)} & Y_0 \times Y_0. \end{array}$$

The horizontal maps in the left-hand square are equivalences by the completeness of X and Y , and so the left square is a pullback square. The middle square is a pullback square by conservativity of φ . The right-hand square is a pullback square, since on horizontal fibers it induces the maps $\text{Hom}_X(x, y) \rightarrow \text{Hom}_Y(\varphi x, \varphi y)$, which are equivalences by full faithfulness of φ . It follows that the outer rectangle is a pullback square, and hence that $\varphi_0: X_0 \rightarrow Y_0$ is a monomorphism of animae. Since it is also surjective on objects, it follows that φ_0 is an equivalence. The right-hand pullback square then shows that also $\varphi_1: X_1 \rightarrow Y_1$ is an equivalence and using the Segal condition we conclude that $\varphi_n: X_n \rightarrow Y_n$ is an equivalence for all n . $\square$

We next construct the associated category functor. The construction itself is forced by the requirement that it preserve colimits, while its adjunction with the nerve follows by checking the desired mapping property on representable simplicial animae.

Proof of Lemma 24.1.10. Since Cat_∞ admits all colimits, Theorem 1.8.9 guarantees that the inclusion $\Delta \hookrightarrow \text{Cat}_\infty$ uniquely extends to a colimit-preserving functor $\text{ac}: \text{sAn} = \text{PSh}(\Delta) \rightarrow \text{Cat}_\infty$. To show that this is a left adjoint to the nerve functor, we must produce a natural isomorphism

$$\text{Hom}_{\text{Cat}_\infty}(\text{ac}(X), C) \cong \text{Hom}_{\text{sAn}}(X, N(C))$$

for every simplicial anima X and every ∞ -category C . Regarding both sides as functors $\text{sAn}^{\text{op}} \rightarrow \text{Fun}(\text{Cat}_\infty, \text{An})$, it follows from Proposition 1.8.6 that both are limit-preserving, hence by another application of Theorem 1.8.9 it suffices to construct such a natural isomorphism after restriction along the Yoneda embedding $\Delta^\bullet: \Delta \hookrightarrow \text{sAn}$. But in this case we may use the composite

$$\text{Hom}_{\text{Cat}_\infty}(\text{ac}(\Delta^n), C) \cong \text{Hom}_{\text{Cat}_\infty}([n], C) = N(C)_n \cong \text{Hom}_{\text{sAn}}(\Delta^n, N(C)),$$

where the last step uses the Yoneda lemma. $\square$

We can now prove that the nerve identifies ∞ -categories with complete Segal anima. The comparison with Dwyer-Kan equivalences shows that the completion map is an equivalence whenever its source is already complete.

Proof of Theorem 24.1.2. Since the nerve functor lands in $\mathbf{CSeg}(\mathbf{An})$, the adjunction $\mathbf{ac} \dashv N$ induces an adjunction

$$\mathbf{ac}: \mathbf{CSeg}(\mathbf{An}) \rightleftarrows \mathbf{Cat}_\infty : N.$$

We have to show that both the unit and the counit are equivalences. For the unit, let X be a complete Segal anima. The unit map $\eta: X \rightarrow N(\mathbf{ac}(X))$ is a Dwyer-Kan equivalence by Proposition 24.1.12 and thus an equivalence by Lemma 24.1.9, since both sides are complete Segal anima. For the counit, let C be an ∞ -category. To show that $\varepsilon: \mathbf{ac}(N(C)) \rightarrow C$ is an equivalence, it suffices by Remark 24.1.6 to show that $N(\varepsilon): N(\mathbf{ac}(N(C))) \rightarrow N(C)$ is an equivalence. But this is a consequence of the triangle identity and the fact that $\eta_{N(C)}: N(C) \rightarrow N(\mathbf{ac}(N(C)))$ is an equivalence. This finishes the proof. $\square$

This equivalence also describes the associated category functor as a localization: it inverts exactly the Dwyer-Kan equivalences.

Proof of Corollary 24.1.13. The first statement is a consequence of the full faithfulness of the nerve functor $N: \mathbf{Cat}_\infty \hookrightarrow \mathbf{Seg}(\mathbf{An})$ from Theorem 24.1.2. For the second statement, consider the following commutative square:

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & Y \\ \simeq \downarrow \eta_X & & \eta_Y \downarrow \simeq \\ N(\mathbf{ac}(X)) & \xrightarrow{N(\mathbf{ac}(\varphi))} & N(\mathbf{ac}(Y)). \end{array}$$

The vertical morphisms are Dwyer-Kan equivalences by Proposition 24.1.12. Since Dwyer-Kan equivalences satisfy the two-out-of-three property, it follows that φ is a Dwyer-Kan equivalence if and only if $N(\mathbf{ac}(\varphi))$ is. The latter is precisely saying that $\mathbf{ac}(\varphi): \mathbf{ac}(X) \rightarrow \mathbf{ac}(Y)$ is an equivalence by Remark 24.1.6. $\square$

Composing this localization with the nerve gives the completion functor, and hence the left adjoint to each of the two inclusions of complete Segal anima considered above.

Proof of Corollary 24.1.14. Given a complete Segal anima Y , we have

$$\begin{aligned} \mathrm{Hom}_{\mathbf{CSeg}(\mathbf{An})}(N(\mathbf{ac}(X)), Y) &\simeq \mathrm{Hom}_{\mathbf{Cat}_\infty}(\mathbf{ac}(X), \mathbf{ac}(Y)) \\ &\simeq \mathrm{Hom}_{\mathbf{sAn}}(X, N(\mathbf{ac}(Y))) \\ &\simeq \mathrm{Hom}_{\mathbf{sAn}}(X, Y), \end{aligned}$$

where the first isomorphism holds by the equivalence $\mathbf{Cat}_\infty \simeq \mathbf{CSeg}(\mathbf{An})$, the second comes from the adjunction $\mathbf{ac} \dashv N$, and the third is induced by $Y \xrightarrow{\sim} N(\mathbf{ac}(Y))$. $\square$

We conclude by applying the localization description to generators. The representable simplicial anima generate all simplicial anima under colimits, and their associated categories can in turn be assembled from $[0]$ and $[1]$.

Proof of Proposition 24.1.16. The presheaf category $\mathbf{sAn} = \mathbf{Fun}(\Delta^{\mathrm{op}}, \mathbf{An})$ is generated by the representable presheaves by Theorem 1.8.9. It follows that $\mathbf{Cat}_\infty$ is generated under colimits by the images of these representable presheaves under the localization functor $\mathrm{ac}: \mathbf{sAn} \rightarrow \mathbf{Cat}_\infty$, which are the posets $[n]$. The claim now follows, since by the Segal condition each $[n]$ can be written as an iterated pushout of copies of $[1]$ along $[0]$. $\square$

24.3.2 The hom animae of a homotopy coherent nerve (online only)

In order for our definition of $N^{\mathrm{top}}(D)$ to be reasonable, we should check that for objects $X, Y \in D$, the hom anima between X and Y in $N^{\mathrm{top}}(D)$ is given by taking the topological space $\mathrm{Hom}_D^{\mathrm{top}}(X, Y)$ and passing to its underlying anima. This computation is somewhat intricate: since Δ^{op} is not filtered, we cannot use the results from Section 24.2 about filtered colimits of ∞ -categories. Indeed, the subcategory $\mathrm{Seg}(\mathbf{An})$ of $\mathbf{sAn}$ is not closed under arbitrary sifted colimits. Nevertheless, we will now see that under suitable assumptions on the diagram, a sifted colimit of Segal animae is again a Segal anima, giving us control over the hom animae in this colimit. I thank Maxime Ramzi for useful discussions.

We will start with some preliminary results regarding pullbacks in $\mathbf{An}$.

Lemma 24.3.5. *Consider a commutative square*

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

of animae. Assume Z is connected and let $z \in Z$ be a choice of basepoint. Then the square is a pullback square if and only if the induced square of fibers over z

$$\begin{array}{ccc} W_z & \longrightarrow & X_z \\ \downarrow & & \downarrow \\ Y_z & \longrightarrow & * \end{array}$$

is a pullback square (so that W_z is the product of X_z and Y_z).

Proof. By the Recognition Principle, Theorem 5.2.1, we may identify Z with the classifying anima $\mathbf{B}\Omega(Z, z)$ of its loop group $\Omega(Z, z)$:

$$Z \simeq \mathrm{colim} \left(\dots \rightrightarrows \Omega(Z, z)^2 \rightrightarrows \Omega(Z, z) \rightrightarrows * \right).$$

As a consequence of descent for colimits (Theorem 5.2.5) it will thus suffice to show that the original commutative square becomes a pullback square after pulling it back along each map $\Omega(Z, z)^n \rightarrow Z$ for $n \geq 0$. Since each of these maps factors through $z: * \rightarrow Z$, it follows from the pasting law of pullbacks that this condition reduces to the case $n = 0$, i.e. the condition that the square becomes a pullback after taking fibers over $z \in Z$. But this holds by assumption. $\square$

Proposition 24.3.6. *Let I be a sifted ∞ -category with initial object i_0 , and consider a pullback square*

$$\begin{array}{ccc} W & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow f \\ Y & \xrightarrow{g} & Z \end{array}$$

in the ∞ -category $\mathrm{Fun}(I, \mathbf{An})$, and assume that the composite $I \xrightarrow{Z} \mathbf{An} \xrightarrow{\pi_0} \mathbf{Set}$ is a constant functor. Then the commutative square

$$\begin{array}{ccc} \mathrm{colim}_{i \in I} W(i) & \longrightarrow & \mathrm{colim}_{i \in I} X(i) \\ \downarrow & \lrcorner & \downarrow f \\ \mathrm{colim}_{i \in I} Y(i) & \xrightarrow{g} & \mathrm{colim}_{i \in I} Z(i) \end{array}$$

is a pullback square in $\mathbf{An}$.

Proof. Case 1: We will first prove the case where each $Z(i)$ is connected and where $Y = \mathrm{const}_*$ is the terminal object in $\mathrm{Fun}(I, \mathbf{An})$. For each $i \in I$, we treat the map $g(i): * = Y(i) \rightarrow Z(i)$ as a basepoint z_i of the connected anima $Z(i)$, and we may consider the Čech nerve $\overline{\Omega}Z(i)$ of the map $z_i: * \rightarrow Z(i)$:

$$\overline{\Omega}Z(i) = \left(\dots \rightrightarrows \Omega Z(i)^2 \rightrightarrows \Omega Z(i) \rightrightarrows * \right).$$

We may define another simplicial object $\overline{\Omega}Z(i) \ltimes W(i)$ via the following pullback squares:

$$\begin{array}{ccc} (\overline{\Omega}Z(i) \ltimes W(i))_k & \longrightarrow & X(i) \\ \downarrow & \lrcorner & \downarrow f(i) \\ (\Omega Z(i))^k & \longrightarrow & Z(i). \end{array}$$

More formally, $\overline{\Omega}Z(i) \ltimes W(i)$ is the pullback of $\overline{\Omega}Z(i)$ and $\mathrm{const}_{X(i)}$ over $\mathrm{const}_{Z(i)}$ in $\mathrm{Fun}(\Delta^{\mathrm{op}}, \mathbf{An})$. It is an immediate consequence of the pasting law of pullback squares that the resulting transformation $\overline{\Omega}Z(i) \ltimes W(i) \rightarrow \overline{\Omega}Z(i)$ is cartesian, in the sense that all commutative squares that comprise it are cartesian. Also note that $(\overline{\Omega}Z(i) \ltimes W(i))_0 \simeq W(i)$ and, more generally,

$$(\overline{\Omega}Z(i) \ltimes W(i))_k \simeq (\Omega Z(i))^k \times W(i)$$

for all $[k] \in \Delta$. We may think of the simplicial object $\overline{\Omega}Z(i) \ltimes W(i)$ as encoding an action of the group $\overline{\Omega}Z(i)$ on the anima $W(i)$, which also explains the choice of notation.

By the Recognition Principle (Theorem 5.2.1), we may recover $Z(i)$ from $\overline{\Omega}Z(i)$ by forming the classifying anima. Similarly, it follows from descent for colimits (Theorem 5.2.5) that $X(i)$ is the colimit of the simplicial object $\overline{\Omega}Z(i) \ltimes W(i)$:

$$Z(i) \simeq \mathbf{B}\overline{\Omega}Z(i), \quad X(i) \simeq \mathbf{B}(\overline{\Omega}Z(i) \ltimes W(i)).$$

We may think of the right-hand side as the homotopy quotient of $W(i)$ by the group object $\overline{\Omega}Z(i)$.

Since all the constructions we did are functorial in the anima $Z(i)$ and $X(i)$, these constructions assemble into a transformation $\overline{\Omega}Z \ltimes W \rightarrow \overline{\Omega}Z$ of functors $I \rightarrow \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathbf{An})$. Passing to colimits,

we thus obtain a natural transformation $\text{colim}_i (\overline{\Omega}Z(i) \ltimes W(i)) \rightarrow \text{colim}_i \overline{\Omega}Z(i)$ of simplicial anima. We claim that this transformation is still a cartesian transformation. Indeed, by the pasting law for pullback squares we may reduce to the case of a morphism $\varphi: [0] \rightarrow [k]$ in Δ . In this case, we need to show that the square

$$\begin{array}{ccc} \text{colim}_i ((\Omega Z(i))^k \times W(i)) & \longrightarrow & \text{colim}_i W(i) \\ \downarrow & & \downarrow \\ \text{colim}_i (\Omega Z(i))^k & \longrightarrow & \text{colim}_i * \end{array}$$

is a pullback square. This is a consequence of the fact that the sifted colimits commute with finite products in $\mathbf{An}$, see Lemma 21.6.11. As a consequence of descent for colimits (Theorem 5.2.5) it then follows that also the square

$$\begin{array}{ccc} \text{colim}_i W(i) & \longrightarrow & \mathbf{B} \text{colim}_i (\overline{\Omega}Z(i) \ltimes W(i)) \\ \downarrow & & \downarrow \\ \text{colim}_i * \simeq * & \longrightarrow & \mathbf{B} \text{colim}_i \overline{\Omega}Z(i) \end{array}$$

is a pullback square. But since colimits commute with each other, the right vertical map is precisely the map $\text{colim}_{i \in I} X(i) \rightarrow \text{colim}_{i \in I} Z(i)$, and so this square is precisely the square we wished to show is a pullback square.

Case 2: We will now do the case where each $Z(i)$ is still connected, but where Y (and X) are allowed to be arbitrary.

Since I by assumption has an initial object i_0 , it follows from Lemma 21.2.4 that $Z(i_0) \simeq \lim_{i \in I} Z(i)$, and in particular the counit of the adjunction provides a map $\text{const}(Z(i_0)) \rightarrow Z$ in $\text{Fun}(I, \mathbf{An})$. Fixing a basepoint $z_0 \in Z(i_0)$, we thus obtain a map $z: \text{const}_* \rightarrow Z$ in $\text{Fun}(I, \mathbf{An})$, providing a coherent family of basepoints $z_i \in Z(i)$.

Consider now the following commutative diagram:

$$\begin{array}{ccccc} \text{colim}_i W(i)_{z_i} & \xrightarrow{\quad} & \text{colim}_i X(i)_{z_i} & & \\ \downarrow & \searrow & \downarrow & \searrow & \\ & \text{colim}_i W(i) & \xrightarrow{\quad} & \text{colim}_i X(i) & \\ \downarrow & \downarrow & \downarrow & \downarrow & \\ \text{colim}_i Y(i)_{z_i} & \xrightarrow{\quad} & \text{colim}_i * \simeq * & \xrightarrow{\quad} & \text{colim}_i Z(i) \\ & \searrow & \searrow & \searrow & \\ & \text{colim}_i Y(i) & \xrightarrow{\quad} & \text{colim}_i Z(i) & \end{array}$$

We wish to show that the front square is a pullback square. By Case 1, we know that the bottom, top, left and right squares are pullback squares, and in particular the back square is obtained from the front square by passing to fibers over $z: * \rightarrow \text{colim}_i Z(i)$. Since $\text{colim}_i Z(i)$ is connected, it thus suffices by Lemma 24.3.5 to show that the back square is a pullback square. But this is again a consequence of the fact that sifted colimits commute with products in $\mathbf{An}$.

Step 3: We now address the general case. Let S be a set satisfying $\pi_0(Z) = \text{const}_S$. The given pullback square in $\text{Fun}(I, \mathbf{An})$ then lives over const_S , hence we may regard it as a pullback square

in $\text{Fun}(I, \text{An}/_S) \simeq \text{Fun}(I, \text{An})^S$. By treating each component separately, we may reduce to the case $S = *$, which is precisely Case 2. $\square$

We may now formulate and prove the result about sifted colimits of Segal anima:

Proposition 24.3.7. *Let I be a sifted ∞ -category with an initial object, and let $X: I \rightarrow \text{sAn}$ be a diagram of ∞ -categories. Assume that the following two conditions are satisfied:*

- (1) *Each simplicial anima $X(i)$ satisfies the Segal condition*
- (2) *The composite*

$$I \xrightarrow{X} \text{sAn} \xrightarrow{\text{ev}_0} \text{An} \xrightarrow{\pi_0} \text{Set}$$

is a constant functor.

Then also the simplicial anima $\text{colim}_{i \in I} X(i)$ satisfies the Segal condition.

Proof. By induction it suffices to show that for all $n \geq 1$, the commutative square

$$\begin{array}{ccc} \text{colim}_{i \in I} X(i)_n & \longrightarrow & \text{colim}_{i \in I} X(i)_1 \\ \downarrow & & \downarrow \\ \text{colim}_{i \in I} X(i)_{n-1} & \longrightarrow & \text{colim}_{i \in I} X(i)_0 \end{array}$$

is a pullback square. This is an immediate consequence of Proposition 24.3.6. $\square$

As a consequence of these results, we may now compute the hom anima in sifted colimits of ∞ -categories.

Corollary 24.3.8. *Let I be a sifted ∞ -category with an initial object, and let $C: I \rightarrow \text{Cat}_\infty$ be a diagram of small ∞ -categories. Assume that for each map $i \rightarrow j$ in I the functor $C(i) \rightarrow C(j)$ induces a bijection $\pi_0 C(i)^\simeq \rightarrow \pi_0 C(j)^\simeq$ on isomorphism classes of objects. Then, given families of objects $x, y: \text{const}_* \rightarrow C$ of the categories $C(i)$, there is an equivalence*

$$\text{colim}_{i \in I} \text{Hom}_{C(i)}(x_i, y_i) \xrightarrow{\sim} \text{Hom}_{\text{colim}_i C(i)}(x, y).$$

Proof. Recall the fully faithful nerve functor $N: \text{Cat}_\infty \hookrightarrow \text{Seg}(\text{An})$ from Chapter 24, which admits a left adjoint $\text{ac}: \text{Seg}(\text{An}) \rightarrow \text{Cat}_\infty$. It follows that the colimit of C in Cat_∞ may be computed as the ∞ -category $\text{ac}(\text{colim}_i N(C(i)))$ associated to the colimit in sAn of the nerves of each of the ∞ -categories $C(i)$. By Proposition 24.3.7, this colimit in sAn is again a Segal anima. It thus follows from Proposition 24.1.12 that the map $\text{colim}_i N(C(i)) \rightarrow N(\text{colim}_i C(i))$ is a Dwyer-Kan equivalence, so that the hom anima of $\text{colim}_i C(i)$ may be computed as the hom anima of $\text{colim}_i N(C(i))$. It thus remains to show that the commutative square

$$\begin{array}{ccc} \text{colim}_i \text{Hom}_{C(i)}(x_i, y_i) & \longrightarrow & \text{colim}_i N(C(i))_1 \\ \downarrow & & \downarrow (d_1, d_0) \\ * \simeq \text{colim}_i * & \longrightarrow & N(C(i))_0 \times N(C(i))_0 \end{array}$$

is a pullback square. But this is an instance of Proposition 24.3.6. $\square$

Proof of Proposition 24.3.4. Since Δ^{op} is sifted, we may apply Corollary 24.3.8 to compute the hom anima in the colimit $N^{\text{top}}(D)$: we get

$$\begin{aligned}
\text{Hom}_{N^{\text{top}}(D)}(X, Y) &\simeq \text{colim}_{[n] \in \Delta^{\text{op}}} \text{Hom}_{D_n}(X, Y) \\
&\simeq \text{colim}_{[n] \in \Delta^{\text{op}}} \text{Hom}_{\text{Top}}(|\Delta^n|, \text{Hom}_D^{\text{top}}(X, Y)) \\
&\simeq \text{colim}_{[n] \in \Delta^{\text{op}}} \text{Sing}(\text{Hom}_D^{\text{top}}(X, Y))_n \\
&= \Pi_{\infty}(\text{Hom}_D^{\text{top}}(X, Y)). \quad \square
\end{aligned}$$

Exercises

Exercise 24.1. Show that composition in a Segal anima is unital and associative.

Exercise 24.2. Consider the functor $\Delta \rightarrow \Delta$ which sends $[n]$ to the poset $[n]^{\text{op}} \star [n] \cong [2n+1]$ that has $2(n+1)$ elements $\bar{n}, \overline{n-1}, \dots, \bar{1}, \bar{0}, 0, 1, \dots, n$ with partial order given by

$$\bar{n} \leq \overline{n-1} \leq \dots \leq \bar{1} \leq \bar{0} \leq 0 \leq 1 \leq \dots \leq n.$$

- Show that this assignment extends to a well-defined functor $D: \Delta \rightarrow \Delta$;
- Show that for every ∞ -category C , the simplicial anima $\text{NTw}(C) \in \text{sAn}$ defined as the composite

$$\Delta^{\text{op}} \xrightarrow{D^{\text{op}}} \Delta^{\text{op}} \hookrightarrow \text{Cat}_{\infty}^{\text{op}} \xrightarrow{\text{Hom}_{\text{Cat}_{\infty}}(-, C)} \text{An}$$

is a complete Segal anima, hence is equivalent to the nerve $N(\text{Tw}(C))$ of some ∞ -category $\text{Tw}(C)$.

- Produce natural transformations $s: [n]^{\text{op}} \hookrightarrow [n]^{\text{op}} \star [n]$ and $t: [n] \hookrightarrow [n]^{\text{op}} \star [n]$ given by $s(i) = \bar{i}$ and $t(i) = i$. Deduce that there are functors

$$s: \text{Tw}(C) \rightarrow C^{\text{op}} \quad \text{and} \quad t: \text{Tw}(C) \rightarrow C.$$

- Show that the fiber of $(s, t): \text{Tw}(C) \rightarrow C^{\text{op}} \times C$ over (x, y) is equivalent to $\text{Hom}_C(x, y)$.

In the following two exercises, you may freely use that filtered colimits commute with finite limits in An .

Exercise 24.3. Show that the subcategory $\text{CSeg}(\text{An}) \subseteq \text{sAn}$ is closed under limits and filtered colimits. Deduce that the nerve functor $N: \text{Cat}_{\infty} \hookrightarrow \text{sAn}$ preserves filtered colimits and that finite limits commute with filtered colimits in Cat_{∞} .

Exercise 24.4. Show that, for every finite category I , the functor $\text{Hom}_{\text{Cat}_{\infty}}(I, -): \text{Cat}_{\infty} \rightarrow \text{An}$ preserves filtered colimits.

Applying this exercise to $I = *$ in particular shows that the core functor $(-)^{\simeq}: \text{Cat}_{\infty} \rightarrow \text{An}$ preserves filtered colimits.

Part IV

Solutions and supplementary proofs

Introduction to Part IV

This online-only part collects solutions and supplementary proofs whose inclusion in the main text would obscure its mathematical progression or take up disproportionate space. The organization and scope are provisional. We begin with the technical proofs underlying the axiomatic development of ∞ -categories in Chapter 1; later versions may include solutions or supplementary arguments from other chapters as well.

25 Solutions to Chapter 1

The material is arranged by dependency, which need not coincide exactly with the order in which the corresponding statements occur in Chapter 1. We include enough detail to make the dependency chain explicit. In particular, a proof may use results stated later in Chapter 1, but every such forward dependency is recorded.

25.1 Basic constructions of ∞ -categories

Solution 25.1.1 (Internal universal properties). We prove Proposition 1.3.24: the canonical comparisons involving terminal and initial categories, products, coproducts, pullbacks, and currying are equivalences.

The point is to construct the inverse comparison functors using only products, coproducts, pullbacks, and currying. In particular, we will not assemble functors pointwise.

For products, postcomposition with the projections $\text{pr}_i: \prod_i C_i \rightarrow C_i$, followed by the product construction, defines the comparison functor

$$\Phi: \text{Fun}\left(T, \prod_{i \in I} C_i\right) \longrightarrow \prod_{i \in I} \text{Fun}(T, C_i).$$

To construct an inverse, first form, for every i , the functor

$$\left(\prod_{j \in I} \text{Fun}(T, C_j)\right) \times T \xrightarrow{\text{pr}_i \times \text{id}_T} \text{Fun}(T, C_i) \times T \xrightarrow{\text{ev}_i} C_i.$$

The product axiom assembles these functors into a functor

$$\left(\prod_{i \in I} \text{Fun}(T, C_i)\right) \times T \longrightarrow \prod_{i \in I} C_i,$$

which we curry to obtain the desired inverse to Φ . Indeed, after composing one way, projecting to the i -th factor, and uncurrying, we recover the displayed evaluation functor, which is the uncurrying of the i -th projection. After composing the other way and uncurrying, postcomposition with every pr_i gives evaluation followed by pr_i . The uniqueness clauses for products and functor categories therefore give both inverse relations. When I is empty, the same argument gives $\text{Fun}(T, *) \xrightarrow{\sim} *$.

The coproduct case requires one additional step, since the coproduct axiom cannot be applied directly to a functor whose source is a product. For every i , consider the diagram

$$\begin{array}{ccccc} X \times C_i & \longrightarrow & X \times \bigsqcup_{j \in I} C_j & \xrightarrow{\text{pr}_X} & X \\ \downarrow & & \downarrow & & \parallel \\ C_i & \xrightarrow{\text{incl}_i} & \bigsqcup_{j \in I} C_j & \longrightarrow & *. \end{array}$$

The right square and the outer rectangle are the pullback squares which express the corresponding products as pullbacks over $*$. Hence the left square is a pullback by Lemma 1.3.15. Part (2) of Axiom C.3, applied to the projection $X \times \bigsqcup_i C_i \rightarrow \bigsqcup_i C_i$, now gives an equivalence

$$\bigsqcup_{i \in I} (X \times C_i) \xrightarrow{\sim} X \times \bigsqcup_{i \in I} C_i,$$

induced on the i -th summand by $(\text{pr}_X, \text{incl}_i \text{pr}_{C_i})$.

For every i , evaluation gives a functor

$$\left(\prod_{j \in I} \text{Fun}(C_j, T) \right) \times C_i \xrightarrow{\text{pr}_i \times \text{id}_{C_i}} \text{Fun}(C_i, T) \times C_i \xrightarrow{\text{ev}_i} T.$$

The coproduct axiom assembles these into a functor

$$\bigsqcup_{i \in I} \left(\left(\prod_{j \in I} \text{Fun}(C_j, T) \right) \times C_i \right) \longrightarrow T.$$

Transporting this functor across the preceding equivalence and currying gives a functor

$$\prod_{i \in I} \text{Fun}(C_i, T) \longrightarrow \text{Fun} \left(\bigsqcup_{i \in I} C_i, T \right).$$

It is inverse to restriction along the inclusions. In one direction, project to the i -th factor and uncurry; in the other, uncurry and precompose with the equivalence

$$\bigsqcup_{i \in I} \left(\text{Fun} \left(\bigsqcup_j C_j, T \right) \times C_i \right) \xrightarrow{\sim} \text{Fun} \left(\bigsqcup_j C_j, T \right) \times \bigsqcup_{i \in I} C_i.$$

The resulting functors agree on every summand with the appropriate evaluation functor. The uniqueness clauses for products, coproducts, and functor categories give both inverse relations. The empty-index case gives $\text{Fun}(\emptyset, T) \xrightarrow{\sim} *$.

Evaluation at the unique object defines a functor $\text{Fun}(*, T) \rightarrow T$. Its inverse is obtained by currying the projection $T \times * \rightarrow T$. The product axiom shows that this projection is an equivalence, and the defining property of currying gives one inverse relation. For the other, uncurry both endofunctors of $\text{Fun}(*, T)$: both become naturally isomorphic to evaluation. Thus $\text{Fun}(*, T) \xrightarrow{\sim} T$.

Now let $f: C \rightarrow E$ and $g: D \rightarrow E$. To construct an inverse to the canonical functor

$$\text{Fun}(T, C \times_E D) \longrightarrow \text{Fun}(T, C) \times_{\text{Fun}(T, E)} \text{Fun}(T, D),$$

consider the two evaluation functors

$$\begin{aligned} (\text{Fun}(T, C) \times_{\text{Fun}(T, E)} \text{Fun}(T, D)) \times T &\longrightarrow \text{Fun}(T, C) \times T \xrightarrow{\text{ev}_C} C, \\ (\text{Fun}(T, C) \times_{\text{Fun}(T, E)} \text{Fun}(T, D)) \times T &\longrightarrow \text{Fun}(T, D) \times T \xrightarrow{\text{ev}_D} D. \end{aligned}$$

The structural natural isomorphism of the pullback, after uncurrying, identifies their composites to E . The pullback axiom therefore assembles them into a functor to $C \times_E D$, and currying gives the desired inverse. The two inverse relations follow by uncurrying and comparing the projections to C and D , including their compatibility over E , using the uniqueness clauses for pullbacks and functor categories.

Finally, the functor

$$\text{Fun}(T, \text{Fun}(C, D)) \longrightarrow \text{Fun}(T \times C, D)$$

is obtained by currying the composite

$$\text{Fun}(T, \text{Fun}(C, D)) \times T \times C \xrightarrow{\text{ev} \times \text{id}_C} \text{Fun}(C, D) \times C \xrightarrow{\text{ev}} D.$$

For its inverse, curry the evaluation functor $\text{Fun}(T \times C, D) \times T \times C \rightarrow D$ first in C and then in T . Repeatedly uncurrying either composite gives the corresponding evaluation functor, so the uniqueness clause of the functor-category axiom gives both inverse relations.

25.2 Animaes and groupoid cores

Solution 25.2.1 (The mapping property of the groupoid core). We prove Exercise 1.4.13: for an anima X , postcomposition with $\gamma_C: C^\simeq \rightarrow C$ induces an equivalence $\text{Map}(X, C^\simeq) \simeq \text{Map}(X, C)$.

Restrict evaluation along the core functor to obtain

$$\text{Map}(X, C) \times X \longrightarrow C.$$

The source is an anima, since products of animaes are animaes, so Axiom E provides a factorization through $C^\simeq$. Currying this factorization gives a functor

$$\text{Map}(X, C) \longrightarrow \text{Fun}(X, C^\simeq).$$

Applying the groupoid core axiom once more, now to the target $\text{Fun}(X, C^\simeq)$, gives a functor

$$R: \text{Map}(X, C) \longrightarrow \text{Map}(X, C^\simeq).$$

Let

$$\gamma_*: \text{Map}(X, C^\simeq) \longrightarrow \text{Map}(X, C)$$

be induced by postcomposition with γ_C . To compare $\gamma_* R$ with the identity, compose both with the core functor $\text{Map}(X, C) \rightarrow \text{Fun}(X, C)$ and uncurry. The resulting functors $\text{Map}(X, C) \times X \rightarrow C$ are naturally isomorphic by the chosen factorization of evaluation through $C^\simeq$. The lifting clause of the groupoid core axiom therefore gives a natural isomorphism $\gamma_* R \cong \text{id}$.

For the other composite, compose $R\gamma_*$ and the identity with the core functor $\text{Map}(X, C^\simeq) \rightarrow \text{Fun}(X, C^\simeq)$ and uncurry. This gives two functors

$$\text{Map}(X, C^\simeq) \times X \longrightarrow C^\simeq.$$

After postcomposition with γ_C , both are naturally isomorphic to the composite of evaluation with γ_C . Since their source is an anima, the lifting clause for γ_C identifies them. Currying and applying the lifting clause once more, now for the core of $\text{Fun}(X, C^\simeq)$, gives $R\gamma_* \cong \text{id}$. Thus R is an inverse to $\gamma_*.X$

Solution 25.2.2 (Groupoid cores preserve pullbacks). We prove Exercise 1.4.14: for functors $C \rightarrow E \leftarrow D$, the canonical comparison $(C \times_E D)^\simeq \rightarrow C^\simeq \times_{E^\simeq} D^\simeq$ is an equivalence.

The projections from $C \times_E D$, restricted to its core, factor through $C^\simeq$ and $D^\simeq$. The structural natural isomorphism over E lifts to one over $E^\simeq$ by Axiom E. The pullback universal property therefore gives a functor

$$(C \times_E D)^\simeq \longrightarrow C^\simeq \times_{E^\simeq} D^\simeq.$$

Conversely, $C^\simeq \times_{E^\simeq} D^\simeq$ is an anima: an inverse to any of its morphisms is constructed from the inverses of its two projections, and the pullback uniqueness clause supplies the inverse relations. Its projections, followed by the core functors to C and D , are compatible over E . They therefore induce a functor to $C \times_E D$, which factors through $(C \times_E D)^\simeq$ by the groupoid-core axiom.

After composing the first resulting composite with the projections to $C^\simeq$ and $D^\simeq$, one obtains the corresponding identity projections. Hence the pullback uniqueness clause identifies this composite with the identity of $C^\simeq \times_{E^\simeq} D^\simeq$. For the other composite, first compose with the core functor $(C \times_E D)^\simeq \rightarrow C \times_E D$. Its two projections agree with those of the identity, including their compatibility over E , so pullback uniqueness gives a natural isomorphism after this composition. The lifting clause of the groupoid-core axiom then gives a natural isomorphism before it. Thus the two functors are inverse equivalences.

25.3 Commutative diagrams and composition

Solution 25.3.1 (Triangles as squares with an identity edge). We prove Proposition 1.4.4: precomposition with p_0 and p_2 identifies triangles with squares whose left and right column, respectively, is an identity.

We first treat the pullback square involving p_0 . It is the outer rectangle in

$$\begin{array}{ccc} \text{Fun}([2], C) & \xrightarrow{p_0^*} & \text{Fun}([1] \times [1], C) \\ d_1^* \downarrow & & \downarrow j_2^* \\ \text{Fun}([1], C) & \xrightarrow{s_0^*} & \text{Fun}([2], C) \\ \text{ev}_0 \downarrow & & \downarrow d_2^* \\ C & \xrightarrow{(p_{[1]})^*} & \text{Fun}([1], C). \end{array}$$

Here $j_2 = (s_1, s_0): [2] \rightarrow [1] \times [1]$, and the composite down the right side is restriction to the left column. Thus this factorization is an equality of the restriction functors appearing in the statement, up to the specified simplicial isomorphisms.

The upper square is a pullback. Indeed, consider

$$\begin{array}{ccccc}
\mathrm{Fun}([2], C) & \xrightarrow{p_0^*} & \mathrm{Fun}([1] \times [1], C) & \xrightarrow{j_0^*} & \mathrm{Fun}([2], C) \\
d_1^* \downarrow & & \downarrow j_2^* & & \downarrow d_1^* \\
\mathrm{Fun}([1], C) & \xrightarrow{s_0^*} & \mathrm{Fun}([2], C) & \xrightarrow{d_1^*} & \mathrm{Fun}([1], C).
\end{array}$$

The commutative-square case of Axiom D says that the square of indexing categories obtained by gluing the two copies of $[2]$ along $[1]$ is a pushout. By the definition of a pushout of ∞ -categories, applying $\mathrm{Fun}(-, C)$ gives precisely the right-hand pullback square above. For the outer rectangle, contravariant functoriality of restriction and the relations $p_0 \circ j_0 \cong \mathrm{id}_{[2]}$ and $s_0 \circ d_1 \cong \mathrm{id}_{[1]}$ identify both horizontal composites with identity functors. Its comparison functor is therefore an equivalence, so the outer rectangle is a pullback. The upper square is now a pullback by Lemma 1.3.15.

For the lower square, use the diagram

$$\begin{array}{ccccc}
\mathrm{Fun}([1], C) & \xrightarrow{s_0^*} & \mathrm{Fun}([2], C) & \xrightarrow{d_0^*} & \mathrm{Fun}([1], C) \\
\mathrm{ev}_0 \downarrow & & d_2^* \downarrow & & \downarrow \mathrm{ev}_0 \\
C & \xrightarrow{(p_{[1]})^*} & \mathrm{Fun}([1], C) & \xrightarrow{\mathrm{ev}_1} & C.
\end{array}$$

The right square is the pullback square obtained by applying $\mathrm{Fun}(-, C)$ to the case $n = 2$ of the first pushout in Axiom D, namely the Segal pullback square. In the outer rectangle, the relations $s_0 \circ d_0 \cong \mathrm{id}_{[1]}$ and $\mathrm{ev}_1 \circ (p_{[1]})^* \cong \mathrm{id}_C$ again identify both horizontal composites with identity functors. Hence the outer rectangle, and therefore the lower square by Lemma 1.3.15, is a pullback. Pasting the upper and lower squares proves the first claim.

For the second square, repeat the construction with the reflected simplicial identities. Explicitly, interchange p_0 with p_2 , j_0 with j_2 , d_0 with d_2 , s_0 with s_1 , and ev_0 with ev_2 . The same two pushout squares from Axiom D and the same pasting argument then prove the second claim.

25.4 Isomorphisms and subcategories

Solution 25.4.1 (Isomorphisms form a subcategory of arrows). We prove Proposition 1.5.6: the projection $\pi_{\mathrm{Iso}}: \mathrm{Iso}(C) \rightarrow \mathrm{Ar}(C)$ is a monomorphism. Equivalently, we must show that its diagonal $\mathrm{Iso}(C) \rightarrow \mathrm{Iso}(C) \times_{\mathrm{Ar}(C)} \mathrm{Iso}(C)$ is an equivalence.

The strategy is to show that the first projection $\mathrm{Iso}(C) \times_{\mathrm{Ar}(C)} \mathrm{Iso}(C) \rightarrow \mathrm{Iso}(C)$ is an equivalence. Composition with the middle isomorphism gives equivalences on the left- and right-inverse data; a pullback square combines these into the desired equivalence of the first projection. Since the diagonal is a section of that projection, it is then an equivalence by 2-out-of-3.

Write

$$r, m, l: \mathrm{Iso}(C) \longrightarrow \mathrm{Ar}(C)$$

for the first, middle, and third projections. Thus $m = \pi_{\mathrm{Iso}}$, while the defining pullback for $\mathrm{Iso}(C)$ supplies natural isomorphisms

$$m \circ r \cong (p_{[1]})^* \circ t \circ m, \quad l \circ m \cong (p_{[1]})^* \circ s \circ m.$$

Here and below, expressions such as $m \circ r$ mean the composite formed by Construction 1.4.7. Associativity and unitality give a natural isomorphism of functors $r \cong l$: explicitly,

$$l \cong l \circ (m \circ r) \cong (l \circ m) \circ r \cong r.$$

The composition functor and the pullback axiom now define functors

$$\begin{aligned} L: \text{Ar}(C) \times_{t,C,s} \text{Iso}(C) &\longrightarrow \text{Ar}(C) \times_{t,C,t} \text{Iso}(C), & (g, u) &\longmapsto (m(u) \circ g, u), \\ R: \text{Iso}(C) \times_{t,C,s} \text{Ar}(C) &\longrightarrow \text{Iso}(C) \times_{s,C,s} \text{Ar}(C), & (u, h) &\longmapsto (u, h \circ m(u)). \end{aligned}$$

These formulas are shorthand for actual functors: for example, the two components of L are the composite of the projections with the composition functor, and the projection to $\text{Iso}(C)$; the source and target identities provide the compatibility needed to apply the pullback axiom.

Define functors in the opposite directions, again using composition and the pullback axiom, by

$$L^{-1}(k, u) := (r(u) \circ k, u), \quad R^{-1}(u, k) := (u, k \circ l(u)).$$

For L , one composite reduces by associativity to $(m \circ r) \circ k \cong k$, while the other reduces to $(r \circ m) \circ g \cong (l \circ m) \circ g \cong g$. For R , the corresponding reductions are

$$(k \circ l) \circ m \cong k \circ (l \circ m) \cong k \quad \text{and} \quad (h \circ m) \circ l \cong h \circ (m \circ r) \cong h,$$

where the second calculation again uses $l \cong r$. Since all the displayed isomorphisms are isomorphisms of functors, this proves that L and R are equivalences, without choosing inverses objectwise.

We next construct the pullback square used in the argument. Every arrow in the following diagram is induced from projection functors, the composition functor, and the pullback axiom:

$$\begin{array}{ccc} \text{Iso}(C) \times_{\text{Ar}(C)} \text{Iso}(C) & \longrightarrow & \text{Ar}(C) \times_{t,C,s} \text{Iso}(C) \times_{t,C,s} \text{Ar}(C) \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow (g,f,h) \mapsto (f \circ g, f, h \circ f) \\ \text{Iso}(C) & \xrightarrow{f \mapsto (\text{id}_y, f, \text{id}_x)} & \text{Ar}(C) \times_{t,C,t} \text{Iso}(C) \times_{s,C,s} \text{Ar}(C). \end{array}$$

More precisely, the top functor has components $(r \text{ pr}_1, \text{pr}_1, l \text{ pr}_2)$, where the defining pullback of $\text{Iso}(C) \times_{\text{Ar}(C)} \text{Iso}(C)$ identifies $m \text{ pr}_1$ with $m \text{ pr}_2$. The bottom functor is induced by the constant-arrow functors at the source and target of m . The defining isomorphisms for $m \circ r$ and $l \circ m$ make the square commute.

To verify the pullback claim formally, form the pullback of the right vertical functor along the bottom functor and regard it as the parameter category. Its projections provide a middle isomorphism structure u and arrows g and h , together with compatible natural isomorphisms

$$m(u) \circ g \cong \text{id}_{t(m(u))}, \quad h \circ m(u) \cong \text{id}_{s(m(u))}.$$

Applying the pullback universal property that defines $\text{Iso}(C)$ to the triples

$$(g, m(u), l(u)) \quad \text{and} \quad (r(u), m(u), h)$$

gives two functors to $\text{Iso}(C)$ with the same middle projection, and hence a functor to $\text{Iso}(C) \times_{\text{Ar}(C)} \text{Iso}(C)$. Conversely, the top functor gives a functor in the other direction. After projecting to each

arrow-category and $\text{Iso}(C)$ factor, both composites are naturally isomorphic to the identity, using the same natural uniqueness calculation for inverse arrows when necessary. The uniqueness clauses for the defining pullbacks therefore show that these two functors are inverse equivalences. This proves that the displayed square is a pullback square.

The right vertical functor is the fiber product over $\text{Iso}(C)$ of L and R . The pullback axiom applied to L^{-1} and R^{-1} constructs an inverse, with the inverse relations supplied by its uniqueness clause. Hence the right vertical functor is an equivalence. Its pullback pr_1 is therefore an equivalence as well: an inverse is induced by an inverse of the right vertical functor together with the bottom arrow, and the pullback uniqueness clause gives the inverse relations. Since $\text{pr}_1 \circ \Delta_{\pi_{\text{Iso}}} \cong \text{id}_{\text{Iso}(C)}$, the diagonal $\Delta_{\pi_{\text{Iso}}}$ is an equivalence by 2-out-of-3. This says precisely that $\pi_{\text{Iso}}: \text{Iso}(C) \rightarrow \text{Ar}(C)$ is a monomorphism.

Solution 25.4.2 (The action of a functor on morphisms). We prove Exercise 1.5.7: the two vertical functors in Axiom H.1 are well defined and fit into the displayed commutative square.

For any ∞ -category D , curry the composite

$$\text{Map}(T, D) \times \text{Map}([1], T) \times [1] \longrightarrow \text{Map}(T, D) \times T \longrightarrow D,$$

where the first arrow evaluates the second factor and the second arrow evaluates the first. This constructs a functor

$$\text{Map}(T, D) \times \text{Map}([1], T) \longrightarrow \text{Fun}([1], D).$$

Its source is an anima, since products of animae are animae, so the groupoid-core axiom factors it through $\text{Map}([1], D)$. Currying gives a functor

$$\text{Map}(T, D) \longrightarrow \text{Fun}(\text{Map}([1], T), \text{Map}([1], D)).$$

The source is again an anima, so a second application of the groupoid-core axiom gives

$$a_D: \text{Map}(T, D) \longrightarrow \text{Map}(\text{Map}([1], T), \text{Map}([1], D)).$$

Taking $D = A$ and $D = C$ produces the two vertical functors in the axiom.

It remains to make the square commute. Postcompose a_A with the functor induced by $\text{Map}([1], i)$, and compare it with a_C after postcomposition by $i: A \rightarrow C$. Include both resulting functors into

$$\text{Fun}(\text{Map}([1], T), \text{Fun}([1], C))$$

and uncurry. Both become the functor

$$\text{Map}(T, A) \times \text{Map}([1], T) \longrightarrow \text{Fun}([1], C), \quad (F, u) \longmapsto i \circ F \circ u,$$

up to the preferred associator for composition. Since the source is an anima, the lifting clause first lifts this associator to an isomorphism between the two uncurried functors with target $\text{Map}([1], C)$. After currying, a second application of the lifting clause, now for the core of $\text{Fun}(\text{Map}([1], T), \text{Map}([1], C))$, gives the required natural isomorphism between the two composites. Hence the square in Axiom H.1 is well defined and commutative.

Solution 25.4.3 (Full subcategories from collections of objects). We prove Proposition 1.5.11: the subcategory $\langle M_\Gamma \rangle_C \hookrightarrow C$ is full and has core equivalent to Γ over $C^\simeq$.

Write $A := \langle M_\Gamma \rangle_C$. By Axiom H.1, applied with $T = *$, the square comparing objects of A and C with their identity arrows is a pullback. Here we use the equivalences $\text{Map}(*, A) \simeq A^\simeq$ and $\text{Map}([1], *) \simeq *$ from Proposition 1.3.24 and Exercise 1.4.13. The subcategory axiom identifies $\text{Map}([1], A)$ with M_Γ . Pasting this square with the pullback defining M_Γ gives a pullback square

$$\begin{array}{ccc} A^\simeq & \longrightarrow & C^\simeq \\ \downarrow & & \downarrow \Delta \\ \Gamma \times \Gamma & \longrightarrow & C^\simeq \times C^\simeq. \end{array}$$

Consequently,

$$A^\simeq \simeq \Gamma \times_{C^\simeq} \Gamma \simeq \Gamma.$$

The last equivalence is the diagonal equivalence characterizing the monomorphism $\Gamma \rightarrow C^\simeq$. All comparison functors are induced by the projections in the three defining pullbacks, so this equivalence lies over $C^\simeq$.

It remains to verify the full-subcategory pullback for a general ∞ -category T . Combining Axiom H.1 with the defining pullback of M_Γ says that a functor $T \rightarrow C$ factors through A precisely when the endpoint functor

$$\text{Map}([1], T) \longrightarrow C^\simeq \times C^\simeq$$

factors through $\Gamma \times \Gamma$. This condition is equivalent, as a pullback condition on mapping anima, to requiring that the object functor $T^\simeq \rightarrow C^\simeq$ factor through Γ . Indeed, a lift $T^\simeq \rightarrow \Gamma$ gives the endpoint lift by precomposition with $s, t: \text{Map}([1], T) \rightarrow T^\simeq$. Conversely, restrict an endpoint lift along the constant-arrow functor $T^\simeq \rightarrow \text{Map}([1], T)$ and project to the first copy of Γ . The first composite is the identity because source followed by the constant-arrow functor is the identity. For the other composite, project the two endpoint lifts to either copy of Γ . Their composites with the monomorphism $\Gamma \rightarrow C^\simeq$ are naturally isomorphic, so the uniqueness property of a monomorphism identifies them. The product uniqueness clause then identifies the two endpoint lifts. Thus the two constructions are inverse. We therefore obtain a pullback square

$$\begin{array}{ccc} \text{Map}(T, A) & \longrightarrow & \text{Map}(T, C) \\ \downarrow & & \downarrow \\ \text{Map}(T^\simeq, \Gamma) & \longrightarrow & \text{Map}(T^\simeq, C^\simeq). \end{array}$$

Using $A^\simeq \simeq \Gamma$, this is precisely the square required in Definition 1.5.9.

Solution 25.4.4 (Functor categories of morphism-inverting functors). We prove Exercise 1.5.18: the functors $C \rightarrow D$ which invert W form a collection of objects of $\text{Fun}(C, D)$ and therefore span a full subcategory $\text{Fun}^W(C, D)$.

The action-on-morphisms functor of Exercise 1.5.7, followed by restriction along $W \hookrightarrow \text{Map}([1], C)$, gives a functor

$$\text{Map}(C, D) \longrightarrow \text{Map}(W, \text{Map}([1], D)).$$

Thus this is an actual functor constructed from composition, groupoid-core factorization, and currying, rather than merely an objectwise assignment.

We first verify that the functor

$$\mathrm{Iso}(D)^{\simeq} \longrightarrow \mathrm{Map}([1], D)$$

induced by π_{Iso} is a monomorphism. By Exercise 1.4.14, taking groupoid cores preserves the pullback which defines the diagonal of π_{Iso} . Hence Proposition 1.5.6 implies that the displayed functor has an equivalent diagonal.

The same argument, together with Proposition 1.3.24, gives

$$\mathrm{Map}(W, A \times_B E) \simeq \mathrm{Map}(W, A) \times_{\mathrm{Map}(W, B)} \mathrm{Map}(W, E).$$

Consequently, applying $\mathrm{Map}(W, -)$ preserves monomorphisms. In particular, the lower horizontal functor in the pullback square

$$\begin{array}{ccc} \mathrm{Map}^W(C, D) & \hookrightarrow & \mathrm{Map}(C, D) \\ \downarrow & \lrcorner & \downarrow \\ \mathrm{Map}(W, \mathrm{Iso}(D)^{\simeq}) & \xrightarrow{(\pi_{\mathrm{Iso}})_*} & \mathrm{Map}(W, \mathrm{Map}([1], D)) \end{array}$$

is a monomorphism. Its base change $\mathrm{Map}^W(C, D) \hookrightarrow \mathrm{Map}(C, D)$ is again a monomorphism: its diagonal is obtained by base change from the diagonal of the lower horizontal functor.

An object of $\mathrm{Map}^W(C, D)$ is precisely a functor $F: C \rightarrow D$ whose action $W \rightarrow \mathrm{Map}([1], D)$ factors through $\mathrm{Iso}(D)^{\simeq}$. Factorization through a monomorphism of animae may be checked on objects, so this says exactly that $F(f)$ is invertible for every f contained in W . Hence $\mathrm{Map}^W(C, D) \hookrightarrow \mathrm{Map}(C, D)$ is the collection of objects consisting of the functors which invert W . Applying Proposition 1.5.11 constructs the required full subcategory $\mathrm{Fun}^W(C, D) \subseteq \mathrm{Fun}(C, D)$.

25.5 Hom animae

Solution 25.5.1 (Hom animae are animae). We prove Proposition 1.4.25: the ∞ -category $\mathrm{Hom}_C(x, y)$ defined by the endpoint pullback is an anima.

Define $C_{/y}$ by the pullback square

$$\begin{array}{ccc} C_{/y} & \longrightarrow & \mathrm{Ar}(C) \\ \downarrow & & \downarrow t \\ * & \xrightarrow{y} & C. \end{array}$$

Let $\pi: C_{/y} \rightarrow C$ be the composite of the top functor with source evaluation $s: \mathrm{Ar}(C) \rightarrow C$. Consider the commutative diagram

$$\begin{array}{ccccc} \mathrm{Hom}_C(x, y) & \longrightarrow & C_{/y} & \longrightarrow & \mathrm{Ar}(C) \\ \downarrow & & \downarrow \pi & & \downarrow (s, t) \\ * & \xrightarrow{x} & C & \xrightarrow{(\mathrm{id}_C, \mathrm{const}_y)} & C \times C \\ & & \downarrow p_C & & \downarrow \mathrm{pr}_2 \\ & & * & \xrightarrow{y} & C. \end{array}$$

The lower-right square is a pullback by the product universal property, while the outer rectangle formed by the two right-hand squares is the pullback defining $C_{/y}$. Hence the upper-right square is

a pullback by Lemma 1.3.15. The outer rectangle formed by the two upper squares is the pullback defining $\text{Hom}_C(x, y)$. A second application of the pasting lemma therefore shows that the upper-left square is a pullback. Consequently, there is a preferred equivalence

$$\text{Hom}_C(x, y) \xrightarrow{\sim} * \times_{x, C, \pi} C/y.$$

The right-hand triangle encoding in Proposition 1.4.4 gives a pullback square

$$\begin{array}{ccc} \text{Ar}(C/y) & \xrightarrow{\pi_*} & \text{Ar}(C) \\ t \downarrow & & \downarrow t \\ C/y & \xrightarrow{\pi} & C. \end{array}$$

Here is the formal identification. Applying $\text{Fun}([1], -)$ to the pullback defining C/y is legitimate by Proposition 1.3.24; it expresses an arrow of C/y as a commutative square in C whose right edge is the identity of y . The cited triangle encoding identifies such squares with triangles. Under this equivalence, restriction to the target object of the arrow in C/y and restriction to the left edge of the square give the two vertical maps displayed above. Thus the comparison functor from $\text{Ar}(C/y)$ to the indicated pullback is an equivalence.

Apply $\text{Ar}(-) = \text{Fun}([1], -)$ to the pullback expressing $\text{Hom}_C(x, y)$ as the fiber of π over x . By Proposition 1.3.24, this gives

$$\text{Ar}(\text{Hom}_C(x, y)) \xrightarrow{\sim} \text{Ar}(*) \times_{\text{Ar}(C)} \text{Ar}(C/y).$$

The preceding pullback square and the already proved equivalence $\text{Ar}(*) \xrightarrow{\sim} *$ therefore give, by pullback pasting, a chain of equivalences

$$\text{Ar}(\text{Hom}_C(x, y)) \xrightarrow{\sim} \text{Ar}(*) \times_{\text{Ar}(C)} (C/y \times_C \text{Ar}(C)) \xrightarrow{\sim} * \times_C C/y \xrightarrow{\sim} \text{Hom}_C(x, y).$$

Consequently, the composite equivalence $\text{Ar}(\text{Hom}_C(x, y)) \rightarrow \text{Hom}_C(x, y)$ is the target functor. Since its composite with the constant-arrow functor is the identity, the constant-arrow functor

$$(p_{[1]})^*: \text{Hom}_C(x, y) \longrightarrow \text{Ar}(\text{Hom}_C(x, y))$$

is an equivalence by 2-out-of-3.

On the other hand, the Rezk axiom applied to $\text{Hom}_C(x, y)$ says that the functor

$$i: \text{Hom}_C(x, y) \rightarrow \text{Iso}(\text{Hom}_C(x, y))$$

is an equivalence. Since $(p_{[1]})^* \cong \pi_{\text{Iso}} \circ i$, another application of 2-out-of-3 shows that

$$\pi_{\text{Iso}}: \text{Iso}(\text{Hom}_C(x, y)) \rightarrow \text{Ar}(\text{Hom}_C(x, y))$$

is an equivalence. Therefore every morphism in $\text{Hom}_C(x, y)$ is isomorphic in its arrow category to the middle arrow of an object of $\text{Iso}(\text{Hom}_C(x, y))$. If r , m and l are the three arrows in such an object, then associativity and unitality give

$$l \cong l \circ (m \circ r) \cong (l \circ m) \circ r \cong r.$$

Thus either inverse may be used on both sides, so m is invertible in the sense of Definition 1.4.10. Transporting an inverse and its two triangles across the isomorphism in the arrow category shows that the original morphism is invertible. Hence $\text{Hom}_C(x, y)$ is an anima.

Bibliography

- [AB19] Omar Antolín-Camarena and Tobias Barthel. “A simple universal property of Thom ring spectra”. *J. Topol.* 12.1 (2019), pp. 56–78. DOI: 10.1112/topo.12084.
- [ABFJ22] Mathieu Anel, Georg Biedermann, Eric Finster, and André Joyal. “Left-exact localizations of ∞ -topoi. I: Higher sheaves”. *Adv. Math.* 400 (2022). Id/No 108268, p. 64.
- [ABS64] Michael F. Atiyah, Raoul Bott, and A. Shapiro. “Clifford modules”. *Topology* 3 (1964), pp. 3–38.
- [Ada58] J. F. Adams. “On the structure and applications of the Steenrod algebra”. *Comment. Math. Helv.* 32 (1958), pp. 180–214.
- [Ada74] J. F. Adams. *Stable homotopy and generalised homology*. Chicago Lectures in Mathematics. Chicago - London: The University of Chicago Press. X, 373 p. £ 3.00 (1974). 1974.
- [AF20] David Ayala and John Francis. “Fibrations of ∞ -categories”. *High. Struct.* 4.1 (2020), pp. 168–265.
- [AFMR17] David Ayala, John Francis, Aaron Mazel-Gee, and Nick Rozenblyum. *Factorization homology of enriched ∞ -categories*. Preprint, arXiv:1710.06414. 2017. URL: <https://arxiv.org/abs/1710.06414>.
- [AG14] Benjamin Antieau and David Gepner. “Brauer groups and étale cohomology in derived algebraic geometry”. English. *Geom. Topol.* 18.2 (2014), pp. 1149–1244. ISSN: 1465-3060. DOI: 10.2140/gt.2014.18.1149.
- [AH59] Michael F. Atiyah and Friedrich Hirzebruch. “Riemann-Roch theorems for differentiable manifolds”. *Bull. Am. Math. Soc.* 65 (1959), pp. 276–281.
- [And+14] Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk. “An ∞ -categorical approach to R -line bundles, R -module Thom spectra, and twisted R -homology”. *J. Topol.* 7.3 (2014), pp. 869–893.
- [Ati61a] Michael F. Atiyah. “Bordism and cobordism”. *Proc. Camb. Philos. Soc.* 57 (1961), pp. 200–208.
- [Ati61b] Michael F. Atiyah. “Thom complexes”. *Proc. Lond. Math. Soc.* (3) 11 (1961), pp. 291–310.
- [Bar17] Clark Barwick. “Spectral Mackey functors and equivariant algebraic K -theory. I.” *Adv. Math.* 304 (2017), pp. 646–727.

- [BBD82] A. A. Beilinson, J. Bernstein, and Pierre Deligne. *Perverse sheaves*. French. Astérisque 100, 172 p. (1982). 1982.
- [Ber07] Julia E. Bergner. “A model category structure on the category of simplicial categories”. English. *Trans. Am. Math. Soc.* 359.5 (2007), pp. 2043–2058. ISSN: 0002-9947. DOI: 10.1090/S0002-9947-06-03987-0.
- [BG99] James C. Becker and Daniel Henry Gottlieb. “A history of duality in algebraic topology”. *History of topology*. Amsterdam: Elsevier, 1999, pp. 725–745.
- [BH21] Tom Bachmann and Marc Hoyois. *Norms in motivic homotopy theory*. Vol. 425. Astérisque. Paris: Société Mathématique de France (SMF), 2021.
- [BHS22] Shaul Barkan, Rune Haugseng, and Jan Steinebrunner. *Envelopes for Algebraic Patterns*. Preprint, arXiv:2208.07183. 2022. URL: <https://arxiv.org/abs/2208.07183>.
- [Bou79] A. K. Bousfield. “The localization of spectra with respect to homology”. *Topology* 18 (1979), pp. 257–281.
- [Bro62] Edgar H. Brown. “Cohomology theories”. *Ann. Math. (2)* 75 (1962), pp. 467–484.
- [BS22] Shaul Barkan and Jan Steinebrunner. *The equifibered approach to ∞ -properads*. Preprint, arXiv:2211.02576. 2022. URL: <https://arxiv.org/abs/2211.02576>.
- [BV73] J. M. Boardman and R. M. Vogt. *Homotopy invariant algebraic structures on topological spaces*. Vol. 347. Lect. Notes Math. Springer, Cham, 1973.
- [CCNW26] Denis-Charles Cisinski, Bastiaan Cnossen, Kim Nguyen, and Tashi Walde. “Synthetic category theory” (2026). URL: <https://drive.google.com/file/d/1lKaq7watGGl3xvjqw9qHjm6SDPFJ2-0o/view>.
- [CH21] Hongyi Chu and Rune Haugseng. “Homotopy-coherent algebra via Segal conditions”. *Adv. Math.* 385 (2021). Id/No 107733, p. 95.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*. Vol. 180. Camb. Stud. Adv. Math. Cambridge: Cambridge University Press, 2019.
- [CLL25] Bastiaan Cnossen, Tobias Lenz, and Sil Linskens. *Universality of span 2-categories and the construction of 6-functor formalisms*. Preprint, arXiv:2505.19192. 2025. URL: <https://arxiv.org/abs/2505.19192>.
- [CM13] Denis-Charles Cisinski and Ieke Moerdijk. “Dendroidal Segal spaces and ∞ -operads”. *J. Topol.* 6.3 (2013), pp. 675–704.
- [CSY21] Shachar Carmeli, Tomer M. Schlank, and Lior Yanovski. “Ambidexterity and height”. *Adv. Math.* 385 (2021), p. 90.
- [Day71] Brian Day. “Construction of biclosed categories”. *Bull. Aust. Math. Soc.* 5 (1971), pp. 139–140.
- [DHKS04] William G. Dwyer, Philip S. Hirschhorn, Daniel M. Kan, and Jeffrey H. Smith. *Homotopy limit functors on model categories and homotopical categories*. Vol. 113. Math. Surv. Monogr. Providence, RI: American Mathematical Society (AMS), 2004.
- [EKMM97] A. D. Elmendorf, Igor Kříž, Michael A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory. With an appendix by M. Cole*. Vol. 47. Math. Surv. Monogr. Providence, RI: American Mathematical Society, 1997.

- [Fre37] H. Freudenthal. “Über die Klassen der Sphärenabbildungen. I. Große Dimensionen.” *Compos. Math.* 5 (1937), pp. 299–314.
- [Gep20] David Gepner. “An introduction to higher categorical algebra”. *Handbook of homotopy theory*. Boca Raton, FL: CRC Press, 2020, pp. 487–548.
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus. “Universality of multiplicative infinite loop space machines”. *Algebr. Geom. Topol.* 15.6 (2015), pp. 3107–3153.
- [GHK22] David Gepner, Rune Haugseng, and Joachim Kock. “ ∞ -operads as analytic monads”. *Int. Math. Res. Not.* 2022.16 (2022), pp. 12516–12624.
- [GHN17] David Gepner, Rune Haugseng, and Thomas Nikolaus. “Lax colimits and free fibrations in ∞ -categories”. *Doc. Math.* 22 (2017), pp. 1225–1266.
- [Gla14] Saul Glasman. *A note on commutative algebras and their modules in quasicategories*. Preprint, arXiv:1409.3506. 2014. URL: <https://arxiv.org/abs/1409.3506>.
- [Gla16] Saul Glasman. “Day convolution for ∞ -categories”. *Math. Res. Lett.* 23.5 (2016), pp. 1369–1385.
- [GM23] David Gepner and Lennart Meier. “On equivariant topological modular forms”. *Compos. Math.* 159.12 (2023), pp. 2638–2693.
- [Gro16] Moritz Groth. “Characterizations of abstract stable homotopy theories”. *arXiv preprint arXiv:1602.07632* (2016).
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge: Cambridge University Press, 2002. ISBN: 0-521-79540-0.
- [Hat03] Allen Hatcher. “Vector bundles and K-theory”. Available online at <https://pi.math.cornell.edu/~hatcher/> (2003).
- [Hau22] Rune Haugseng. “ ∞ -operads via symmetric sequences”. *Math. Z.* 301 (2022), pp. 115–171. DOI: 10.1007/s00209-021-02881-w. arXiv: 1708.09632.
- [Hau23a] Rune Haugseng. “An allegedly somewhat friendly introduction to ∞ -operads” (2023).
- [Hau23b] Rune Haugseng. *From analytic monads to ∞ -operads through Lawvere theories*. Preprint, arXiv:2301.01199. 2023. URL: <https://arxiv.org/abs/2301.01199>.
- [Hau25] Rune Haugseng. *Yet another introduction to ∞ -categories*. Available on the author’s homepage. 2025. URL: https://runegha.folk.ntnu.no/naiveocat_web.pdf.
- [Hei23] Hadrian Heine. “An equivalence between enriched ∞ -categories and ∞ -categories with weak action”. *Adv. Math.* 417 (2023), p. 140.
- [HHLN23a] Rune Haugseng, Fabian Hebestreit, Sil Linskens, and Joost Nuiten. “Lax monoidal adjunctions, two-variable fibrations and the calculus of mates”. *Proc. Lond. Math. Soc.* (3) 127.4 (2023), pp. 889–957.
- [HHLN23b] Rune Haugseng, Fabian Hebestreit, Sil Linskens, and Joost Nuiten. “Two-variable fibrations, factorisation systems and ∞ -categories of spans”. *Forum Math. Sigma* 11 (2023). Id/No e111, p. 70.
- [HHR21] Fabian Hebestreit, Gijs Heuts, and Jaco Ruit. *A short proof of the straightening theorem*. Preprint, arXiv:2111.00069. 2021. URL: <https://arxiv.org/abs/2111.00069>.

- [Hin20] Vladimir Hinich. “Yoneda lemma for enriched ∞ -categories”. *Adv. Math.* 367 (2020). Id/No 107129, p. 119.
- [Hir15] Philip S. Hirschhorn. *Functorial CW-approximation*. Preprint, arXiv:1508.01944. 2015. URL: <https://arxiv.org/abs/1508.01944>.
- [HM22] Gijs Heuts and Ieke Moerdijk. *Simplicial and dendroidal homotopy theory*. Vol. 75. *Ergeb. Math. Grenzgeb.*, 3. Folge. Cham: Springer, 2022. ISBN: 978-3-031-10446-6; 978-3-031-10447-3.
- [HM24a] Claudius Heyer and Lucas Mann. *6-Functor Formalisms and Smooth Representations*. Preprint, arXiv:2410.13038. 2024. URL: <https://arxiv.org/abs/2410.13038>.
- [HM24b] Vladimir Hinich and Ieke Moerdijk. “On the equivalence of Lurie’s ∞ -operads and dendroidal ∞ -operads”. *Journal of Topology* 17.4, e70003 (2024). DOI: 10.1112/topo.70003.
- [Hov99] Mark Hovey. *Model categories*. Vol. 63. *Math. Surv. Monogr.* Providence, RI: American Mathematical Society, 1999. ISBN: 0-8218-1359-5.
- [HPS97] Mark Hovey, John H. Palmieri, and Neil P. Strickland. *Axiomatic stable homotopy theory*. Vol. 610. *Mem. Am. Math. Soc.* Providence, RI: American Mathematical Society (AMS), 1997.
- [HS23] Fabian Hebestreit and Jan Steinebrunner. *A short proof that Rezk’s nerve is fully faithful*. Preprint, arXiv:2312.09889. 2023. URL: <https://arxiv.org/abs/2312.09889>.
- [HSS00] Mark Hovey, Brooke Shipley, and Jeff Smith. “Symmetric spectra”. *J. Am. Math. Soc.* 13.1 (2000), pp. 149–208.
- [Hus93] Dale H. Husemoller. *Fibre bundles*. 3rd ed. Vol. 20. *Grad. Texts Math.* Berlin: Springer-Verlag, 1993.
- [HW21] Fabian Hebestreit and Ferdinand Wagner. *Algebraic and Hermitian K-Theory*. 2021.
- [Joy08] André Joyal. “Notes on quasi-categories”. *preprint* (2008).
- [JT07] André Joyal and Myles Tierney. “Quasi-categories vs Segal spaces”. *Categories in algebra, geometry and mathematical physics. Conference and workshop in honor of Ross Street’s 60th birthday, Sydney and Canberra, Australia, July 11–16/July 18–21, 2005*. Providence, RI: American Mathematical Society (AMS), 2007, pp. 277–326.
- [Lan21] Markus Land. *Introduction to infinity-categories*. *Compact Textb. Math.* Cham: Birkhäuser, 2021.
- [Lim59] Elon L. Lima. “The Spanier-Whitehead duality in new homotopy categories”. *Summa Brasil. Math.* 4 (1959), pp. 91–147.
- [LMS86] L. G. jun. Lewis, J. P. May, and M. Steinberger. *Equivariant stable homotopy theory. With contributions by J. E. McClure*. Vol. 1213. *Lect. Notes Math.* Springer, Cham, 1986.
- [Lur09a] Jacob Lurie. *(Infinity, 2)-categories and the Goodwillie calculus I*. Preprint, arXiv:0905.0462. 2009. URL: <https://arxiv.org/abs/0905.0462>.
- [Lur09b] Jacob Lurie. *Higher topos theory*. Vol. 170. *Ann. Math. Stud.* Princeton, NJ: Princeton University Press, 2009.

- [Lur17] Jacob Lurie. “Higher algebra”. <https://www.math.ias.edu/~lurie/papers/HA.pdf> (2017).
- [Lur26] Jacob Lurie. “Kerodon”. *online textbook, available at <https://kerodon.net>* (2026).
- [May72] J. P. May. *The geometry of iterated loop spaces*. Vol. 271. Lect. Notes Math. Springer, Cham, 1972.
- [May77] J. Peter May. *E_∞ ring spaces and E_∞ ring spectra. With contributions by Frank Quinn, Nigel Ray, and Jorgen Tornehave*. English. Vol. 577. Lect. Notes Math. Springer, Cham, 1977. doi: 10.1007/BFb0097608.
- [May99a] J. P. May. *A concise course in algebraic topology*. Chicago, IL: University of Chicago Press, 1999.
- [May99b] J. P. May. “Stable algebraic topology, 1945–1966”. *History of topology*. Amsterdam: Elsevier, 1999, pp. 665–723.
- [Mil59] John Milnor. “On spaces having the homotopy type of a CW-complex”. *Transactions of the American Mathematical Society* 90.2 (1959), pp. 272–280.
- [Mil63] John Milnor. *Morse theory*. Vol. 51. Annals of Mathematics Studies. Princeton, NJ: Princeton University Press, 1963.
- [MS60] John W. Milnor and Edwin H. Spanier. “Two remarks on fiber homotopy type”. *Pac. J. Math.* 10 (1960), pp. 585–590.
- [MS74] John W. Milnor and James D. Stasheff. *Characteristic classes*. Vol. 76. Ann. Math. Stud. Princeton University Press, Princeton, NJ, 1974.
- [MW07] Ieke Moerdijk and Ittay Weiss. “Dendroidal sets”. *Algebr. Geom. Topol.* 7 (2007), pp. 1441–1470.
- [Nar21] Denis Nardin. *Introduction to stable homotopy theory*. 2021.
- [Nik16] Thomas Nikolaus. *Stable ∞ -operads and the multiplicative Yoneda lemma*. Preprint, arXiv:1608.02901. 2016. URL: <https://arxiv.org/abs/1608.02901>.
- [Nik17] Thomas Nikolaus. “The group completion theorem via localizations of ring spectra”. *Available on author’s webpage* (2017).
- [NS18] Thomas Nikolaus and Peter Scholze. “On topological cyclic homology”. *Acta Math.* 221.2 (2018), pp. 203–409.
- [Pon59] Lev S. Pontryagin. “Smooth manifolds and their applications in homotopy theory”. *Am. Math. Soc. Transl. (2)* 11 (1959). Translated from the Russian original, Trudy Mat. Inst. Steklov 45 (1955), pp. 1–114.
- [Qui67] D. G. Quillen. *Homotopical algebra*. Vol. 43. Lect. Notes Math. Springer, Cham, 1967.
- [Rav84] Douglas C. Ravenel. “Localization with respect to certain periodic homology theories”. *Am. J. Math.* 106 (1984), pp. 351–414.
- [Rav86] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres*. Vol. 121. Pure Appl. Math., Academic Press. Academic Press, New York, NY, 1986.
- [Rav92] Douglas C. Ravenel. *Nilpotence and periodicity in stable homotopy theory*. Vol. 128. Ann. Math. Stud. Princeton, NJ: Princeton University Press, 1992.

- [Rez01] Charles Rezk. “A model for the homotopy theory of homotopy theory”. *Trans. Am. Math. Soc.* 353.3 (2001), pp. 973–1007.
- [Rie14] Emily Riehl. *Categorical homotopy theory*. Vol. 24. New Math. Monogr. Cambridge: Cambridge University Press, 2014.
- [Sch25] Peter Scholze. *Six-Functor Formalisms*. Preprint, arXiv:2510.26269. 2025. URL: <https://arxiv.org/abs/2510.26269>.
- [Seg68] Graeme Segal. “Classifying spaces and spectral sequences”. *Publ. Math., Inst. Hautes Étud. Sci.* 34 (1968), pp. 105–112.
- [Ser53] Jean-Pierre Serre. “Groupes d’homotopie et classes de groupes abéliens”. *Ann. Math.* (2) 58 (1953), pp. 258–294.
- [Sha21] Jay Shah. *Parametrized higher category theory II: Universal constructions*. Preprint, arXiv:2109.11954. 2021. URL: <https://arxiv.org/abs/2109.11954>.
- [Sta63] James Dillon Stasheff. “Homotopy associativity of H-spaces. II”. *Transactions of the American Mathematical Society* 108.2 (1963), pp. 293–312.
- [Sto68] Robert E. Stong. *Notes on cobordism theory*. Vol. 7. Mathematical Notes. Princeton, NJ: Princeton University Press, 1968.
- [SW53] Edwin H. Spanier and J. H. C. Whitehead. “A first approximation to homotopy theory”. *Proc. Natl. Acad. Sci. USA* 39 (1953), pp. 655–660.
- [Swi75] Robert M. Switzer. *Algebraic topology - homotopy and homology*. Vol. 212. Grundlehren Math. Wiss. Springer, Cham, 1975.
- [tDie08] Tammo tom Dieck. *Algebraic topology*. EMS Textb. Math. Zürich: European Mathematical Society (EMS), 2008.
- [Tho54] René Thom. “Quelques propriétés globales des variétés différentiables”. *Comment. Math. Helv.* 28 (1954), pp. 17–86.
- [Vol25] Marco Volpe. “Verdier duality on conically smooth stratified spaces”. *Algebr. Geom. Topol.* 25.2 (2025), pp. 919–950.
- [Wei94] Charles A. Weibel. *An introduction to homological algebra*. Vol. 38. Camb. Stud. Adv. Math. Cambridge University Press, 1994.
- [Wei99] Charles A. Weibel. “History of homological algebra”. *History of topology*. Amsterdam: Elsevier, 1999, pp. 797–836.
- [Whi62] George W. Whitehead. “Generalized homology theories”. *Trans. Am. Math. Soc.* 102 (1962), pp. 227–283.
- [Win25] Christoph Winges. *Hinich’s model for Day convolution revisited*. Preprint, arXiv:2506.06025. 2025. URL: <https://arxiv.org/abs/2506.06025>.

Index of notation

$X \oplus Y$, 114	$\text{fib}(g)$, 106
$X \otimes Y$, 134	Fin , 153
$[n]$, 113	Fin_* , 471
$\mathbb{1}$, 243	$\text{Fun}(C, D)$, 24
$\text{Alg}_O(C)$, 337, 381	$\text{Fun}^\otimes(C, D)$, 243
An , 49	$\text{Gr}_k(\mathbb{C}^n)$, 275
An_* , 77	$\text{Grp}(C)$, 143
$\text{Ar}(C)$, 29	HA , 168
$\mathcal{A}ssoc$, 336	$\text{Ho}(C)$, 53
$\mathbf{B}X$, 146	$\text{Hom}_C(x, y)$, 32
$\text{BMod}(C)$, 501	$\text{hom}_C(X, Y)$, 131
C^{op} , 35	$\underline{\text{Hom}}(X, Z)$, 320
C_* , 106	$\text{Ind}(C)$, 567
$C^\simeq$, 31	$\mathcal{K}(\mathcal{A})$, 185
$C[W^{-1}]$, 39	KU , 290
$\text{CAlg}(C)$, 491	ku , 288
$\text{Cart}(C)$, 574	$\text{LFib}(C)$, 577
Cat_∞ , 49	$\mathcal{L}ie$, 336
$\text{Cat}_\infty^{\text{st}}$, 112	$\lim_k^1 A_k$, 138
$\text{CGrp}(C)$, 155	$\text{LMod}_A(C)$, 497
$\text{Ch}(\mathcal{A})$, 176	$\mathcal{L}Mod$, 341
$\text{CMon}(C)$, 155	$\mathcal{M}_C$, 379
$\text{Cocart}(C)$, 572	MO , 305
$\text{cofib}(f)$, 106	$\text{Mod}_A(C)$, 501
$Comm$, 336	Mod , 340
$D(\mathcal{A})$, 178	$\text{Mon}(C)$, 142
$\mathcal{D}ay$, 424	MU , 307
Δ^n , 41	$O_{\langle 1 \rangle}$, 352
Δ , 41	$O^\otimes$, 349
End , 337	ΩX , 106
$\text{Env}(O)$, 343	Ω^∞ , 129
Ext , 200	

Op_∞ , 369
 $\pi_0(X)$, 53
 Π_∞ , 61
 $\pi_k(X)$, 135
 $\mathrm{Pic}(C)$, 320
 Pr^L , 562
 $\mathrm{PSh}(C)$, 51

 $\mathrm{RFib}(C)$, 577

 $\mathbb{S}/p$, 236
 $\mathbb{S}/p^\infty$, 236
 S^n , 81
 $\mathbb{S}$, 130
 ΣX , 106
 Σ^∞ , 129
 Σ_+^∞ , 129
 Sp , 129
 $\mathrm{Sp}_\mathbb{Q}$, 235
 $\mathrm{Sp}(C)$, 118
 $\mathrm{Span}(\mathrm{Fin})$, 153
 St , 580

 $\mathrm{Th}(E)$, 296
 Tor , 206
 $\mathcal{T}riv$, 336
 $\mathrm{Tw}(C)$, 591

 $\mathrm{Un}^{\mathrm{cc}}$, 580

 $\mathrm{Vect}(X)$, 273

Index

- 1-category, **53**
- A_∞ -algebra, **509**
- A_∞ -module, **510**
- abelian category, 109, **175**
- abelianization functor, **558**
- active morphism, **349**
- additive ∞ -category, **114**
- additive operad, **479**
- adequate triple, **354**
- adjointable
 - left, **399**
 - right, **400**
- adjunction, 52, **539**
- algebra over an operad, **337**, 381
- algebraic pattern, **464**
- anima, **12**
 - of colors, **370**
 - of factorizations, **373**
 - of multimorphisms, **370**
 - of objects, **19**
- arithmetic fracture square, **235**
- arrow category, **29**
- associated category, **604**
- associative algebra, **342**
- associative operad, **336**
- associative ring spectrum, **246**
- Atiyah duality, **328**
- attaching map, **93**
- augmented simplex category, **144**
- augmented simplicial object, **144**
- bar resolution, **556**
- base space, **278**
- Beck–Chevalley condition, **398**
- Beck–Chevalley fibration, **400**
- bifiber sequence, **109**
- bifibration, **587**
- bilinear map, **438**
- bimodule, **501**
- bordant, **308**
- B -bordism cycle, **308**
- Bott class, **286**
- Bott map, **290**
- Bott periodicity, **286**, 290, 291
- bounded, **199**
 - above, **199**
 - below, **199**
 - derived category, **199**
- Bousfield
 - class, **241**
 - colocalization, **558**
 - localization, **228**
- Brown representability, **92**, 100
- canonical bundle, **276**
- cartesian
 - fibration, **574**
 - functor, **574**
 - monoidal structure, 243, **398**
 - morphism, **574**
 - natural transformation, **594**
- Čech nerve functor, **144**
- cell complex, **70**
- cellular
 - approximation theorem, **94**
 - boundary map, **96**
 - chain complex, **96**
 - homology, **96**
- chain
 - complex, **176**
 - homotopy, 12, **177**
 - homotopy equivalence, **177**

- map, **176**
- Chern character, **293**
- class of backwards morphisms, **355**
- class of cofibrations, **67**
- class of fibrations, **66**
- class of forward morphisms, **355**
- classifying anima, **146**
- classifying space, **276**
- closed symmetric monoidal ∞ -category, **320**
- clutching function, **274**
- cocartesian
 - fibration, **572**
 - monoidal structure, 243, **398**
 - morphism, **572**
 - operad, **398**
 - section, **587**
 - square, **26**
 - transport, **573**
- cocone, **44**
- coconnective, **190**
- coevaluation map, **318**
- cofiber, 78, **106**
 - sequence, 78, **106**
- cofibrant, **67**
 - replacement, **210**
- cofibration, **67**
- cofree object, **477**
- cogroup object, **82**, 140
- cokernel, **175**
- colimit, **44**
 - cocone, **44**
 - diagram, **45**
- collared, **74**
- colored operad, **338**
- colors of an operad, **338**
- commutative
 - algebra, **244**
 - group, **155**
 - ∞ -operad, **373**
 - monoid, **155**
 - operad, **336**
 - ring object, **439**
 - ring spectrum, **246**
 - semiring, **294**
 - semiring object, **439**
- square, **27**
- compact generators, **565**
- compact object, **565**
- compactly generated, **565**
- comparison functor
 - monadic, **518**
- complete Segal anima, 354, **603**
- complete Segal space, 34
- completeness condition, **34**
- complex
 - bordism spectrum, **307**
 - K-theory, **272**
 - connective, **288**
 - periodic, **290**
 - vector bundle, **272**
- composite of morphisms, **29**
- composition functor, **29**
- composition map of an operad, **336**
- concentrated in a degree, **177**
- cone, **44**
- cone of a space, **65**
- connected anima, **81**
- connective, **165**, 190
- connectivity-preserving, **247**
- conservative functor, **607**
- contractible
 - ∞ -category, **20**
 - anima, **14**
 - chain complex, **212**
- contraction, **20**
- coproduct, **20**
- coPuppe sequence, **89**
- cospan, **27**
- cotensoring functor, **162**
- counit of an adjunction, **540**
- cup product, **242**
- currying, **24**
- CW-complex, **70**
- CW-structure on an anima, **93**
- Day convolution, 243, **424**
- décalage, **148**
- degeneracy map, **18**
- degree of a map, **97**
- degreewise projective, **212**
- degreewise split monomorphism, **185**

- delooping, **141**
- derived
 - category, **12**
 - functor, **209**
 - total left, **209**
 - total right, **210**
 - ∞ -category, **178**
 - limit
 - first, **138**
 - p -completion groups, **238**
 - tensor product, **216**
- descent, **149**
- DGA, **252**
- differential graded algebra, *see* DGA
- dimension axiom, **88**
- direct sum of vector bundles, **273**
- disjoint union, *see* coproduct
- distinguished triangle, **109**
- double mapping cylinder, **63**
- dual object, **318**
- duality datum, **318**
- dualizable, **318**
- Dwyer–Kan equivalence, **603**

- $\mathcal{E}_0$ -operad, **373**
- E_1 -monoid, **142**
- E -acyclic, **229**
- E -cohomology
 - reduced, **91**
- E -equivalence, **229**
- E -fundamental class, **330**
- E -homology
 - reduced, **137**
- $\mathcal{E}_k$ -algebra, **342**
- E -local, **229**
- E -localization, **230**
- $\mathcal{E}_n$ -monoid, **170**
- Eilenberg–MacLane
 - anima, **184**
 - ring spectrum, **253**
 - spectrum, **168, 184**
- Eilenberg–Moore category, **517**
- empty ∞ -operad, **377**
- empty anima, **13**
- end of a functor, **591**
- endomorphism
 - category, **512**
 - object, **511**
 - operad, **337**
- enriched morphism, **511**
- envelope of an operad, **343**
- equivalence of ∞ -categories, **17**
- equivalence of anima, **13**
- essentially surjective functor, **39**
- evaluation functor, **24**
- evaluation map of a duality, **318**
- exact
 - functor, **109**
 - sequence, **109**
 - sequence in an abelian category, **176**
 - sequence of pointed sets, **78**
 - square, **112**
 - triangle, **109**
- excisive functor, **440**
- exponentiable fibration, **597**
- Ext-group, **200, 264**
- extensive, **364**
- external direct sum, **297**
- extra degeneracies, **148**

- $\mathcal{F}$ -monoid, **465**
- $\mathcal{F}$ -monoidal ∞ -category, **465**
- $\mathcal{F}$ -operad, **465**
- face map, **18**
- factorization system, **373, 451**
- fiber
 - of a functor, **22**
 - of a morphism, **78, 106**
 - sequence, **78, 106**
- fiber product, **21**
- fiberwise one-point compactification, **296**
- fibrant, **67**
- fibration, **67**
- filtered ∞ -category, **552**
- final functor, **551**
- finite ∞ -category, **116**
- finite CW-complex, **94**
- finite-dimensional CW-complex, **94**
- finitely generated module, **265**
- flat
 - module, **266**
 - object, **205**

- resolution, **208**
- free cocompletion, **51**
- free group functor, **171**
- free module, **265**
- full subcategory inclusion, **38**
- full suboperad, **393**
- fully faithful functor, **38**
- fully faithful morphism of ∞ -operads, **370**
- functor, **15**
- functor category, **24**
- fundamental ∞ -groupoid, **13**

- G*-principal bundle, **277**
- G*-split, **555**
- generalized (co)homology theory, **88**
- geometric realization
 - of a simplicial object, **48**
 - of a simplicial set, **43**
 - of an ∞ -category, **40**
- graded abelian group, **88**
- Grassmannian
 - infinite, **275**
- Grothendieck construction, **597**
- group completion, **166**
- group completion theorem, **169**
- group object, **143**
- grouplike, **143**
- groupoid core, **13, 31**

- heart of a *t*-structure, **189**
- hom anima, **32**
- Hom functor, **50**
- hom spectrum, **131**
- homology of a chain complex, **177**
- homotopical *n*-sphere, **81**
- homotopy
 - category, **53**
 - cocomplete, **69**
 - cocontinuous, **69**
 - cofiber, **65**
 - coherent diagram, **11**
 - coherent nerve, **606**
 - commutative square, **64**
 - complete, **69**
 - continuous, **69**
 - equivalence, **11**
 - fiber, **66**
 - group object, **195**
 - group of a spectrum, **135**
 - group of an anima, **81**
 - hypothesis, **61**
 - pullback, **63**
 - pushout, **63**
 - ring spectrum, **242**
- homotopy category of chain complexes, **185**
- horizontal composite, **16**

- identity
 - morphism, **19**
 - operation, **336**
 - span, **353**
- image of a morphism, **176**
- Ind-completion, **567**
- inert morphism, **171, 349**
- infinite loop space, **165**
- $(\infty, 1)$ -category, **14**
- ∞ -category, **14**
 - with weak equivalences and cofibrations, **67**
 - with weak equivalences and fibrations, **67**
- ∞ -groupoid, **31**
- ∞ -operad, **349**
 - in the sense of Lurie, **471**
- ∞ -topos, **594**
- initial functor, **551**
- initial object, **46**
- input colors, **338**
- internal duality functor, **321**
- internal hom, **320**
 - of chain complexes, **218**
- inverse functor, **17**
- inversion map, **83**
- invertible morphism, **31**
- invertible object, **320**
- invertible spectrum, **301**
- isomorphism, **31**

- J-homomorphism, **303**

- k*-coconnective, **180**
- k*-connective, **180**
- k*-horn, **41**

- K-theory group, **283**
 - reduced, **284**
- Kan complex, **42**
- Kan extension
 - pointwise, **549**
- kernel, **175**
- Koszul symmetry, **448**
- Künneth theorem, **223**
 - algebraic, **221**
- L*-equivalence, **393**
- large ∞ -category, **50**
- lax monoidal functor, **391**
- lax structure maps, **382**
- lax symmetric monoidal functor, **243**, **382**
- left
 - adjoint object, **230**, **541**
 - bounded, **199**
 - complete t-structure, **198**
 - deformation, **211**
 - derivable, **528**
 - exact ∞ -category, **117**
 - exact functor, **68**, **117**
 - fibration, **577**
 - Kan extension, **51**, **549**
 - module, **342**
 - module operad, **341**
 - Ore condition, **259**
 - orthogonal, **451**
 - separated t-structure, **198**
 - t-exact, **190**
 - tensoring, **497**
- Lie operad, **336**
- limit, **44**
 - cone, **44**
 - diagram, **45**
- little cubes operad, **337**
- local system of lines, **310**
- localization, **39**
- localization of a ring spectrum, **257**
- locally trivial bundle, **278**
- loop
 - anima, **79**
 - group, **145**
 - object, **106**
 - space, **66**
- Lurie operadic envelope, **473**
- M*-local, **491**
- mapping
 - anima, **31**
 - cone, **177**
 - cylinder, **73**, **181**
 - path space, **75**
 - spectrum, *see* hom spectrum
 - telescope, **137**
- Mayer–Vietoris property, **99**
- mode, **491**
- model category, **66**
- module operad, **340**
- module over an algebra, **342**
- moduli anima of vector bundles, **302**
- monad, **517**
- monadic functor, **518**
- monoid object, **142**
- monoid over an operad, **389**
- monoidal ∞ -category, **389**
- monoidal unit, **243**
- monomorphism, **36**
- Moore spectrum, **236**
- morphism, **18**
 - of adequate triples, **355**
 - of colored operads, **339**
 - of ∞ -operads, **369**
 - of spectra, **90**
 - of vector bundles, **273**
- multicategory, *see* colored operad
- multimorphism, **338**
- multimorphism operad, **379**
- multiplication of a monoid, **142**
- n*-ary operation, **335**
- n*-cells, **93**
- n*-connected, **165**
- n*-fold group, **169**
- n*-fold monoid, **169**
- n*-isomorphism, **16**
- n*-simplex, **41**
- n*-universal, **100**
- natural isomorphism, **15**
- natural transformation, **33**
- NDR-pair, **74**

- neighborhood deformation retract pair, *see* NDR-pair
- nerve, **43**
- neutral element, **142**
- non-colored operad, **336**
- normal B -structure, **307**
- null map, **77**, **105**
- nullhomotopy, **78**, **106**
- nullsequence, **77**, **106**

- $\mathcal{O}$ -monoidal ∞ -category, **389**
- object, **18**
- one-point anima, **13**
- operad, **336**
- operadic colimit, **478**
- operadic envelope, **466**
- operadic limit, **478**
- oplax symmetric monoidal functor, **382**
- opposite
 - algebra, **503**
 - category, **35**
 - symmetric monoidal ∞ -category, **378**
 - t -structure, **191**
- ordinary cohomology, **187**
- ordinary homology, **187**
- orthofibration, **427**
- output color, **338**
- over category, **546**

- p -complete, **236**
- p -completion, **235**
- P -divisible, **232**
- p -local, **231**
- p -localization, **232**
- P -power torsion, **233**
- paracompact, **273**
- partition of unity, **273**
- path components, set of, **53**
- perfect module, **246**
- Picard anima, **320**
- Poincaré duality, **330**
- pointed
 - anima, **77**
 - functor, **105**
 - ∞ -category, **105**
 - objects, ∞ -category of, **106**
 - operad, **336**, **479**
 - pointwise symmetric monoidal structure, **243**
 - Pontryagin–Thom collapse map, **297**
 - Pontryagin–Thom theorem, **309**
 - Postnikov t -structure, **190**
 - preadditivity, **115**
 - presentable ∞ -category, **562**
 - presentably symmetric monoidal, **570**
 - presheaf, **51**
 - category, **51**
 - representable, **51**
 - prespectrum, **119**
 - product, **13**, **20**
 - projective
 - module, **265**
 - object, **199**
 - resolution, **202**
 - pullback, **21**
 - square, **23**
 - pushout, **26**
 - square, **26**

 - quasi-isomorphism, **12**, **177**
 - quasicategory, **40**, **42**

 - $\mathcal{R}$ -cocartesian fibration, **458**
 - R -line, **310**
 - R -orientation, **310**
 - rank of a vector bundle, **273**
 - rational equivalence, **229**
 - rational spectrum, **235**
 - rationalization, **232**
 - recognition principle
 - for infinite loop spaces, **141**
 - for loop spaces, **141**
 - reduced
 - chain complex, **187**
 - cochain complex, **187**
 - cohomology theory, **89**
 - functor, **440**
 - homology theory, **88**
 - suspension spectrum, **129**
 - reflective subcategory, **526**
 - regular module, **246**
 - relative
 - adjoint, **383**

- adjunction, **383**
- cell complex, **70**
- ∞ -category, **527**
- slice category, **548**
- tensor product, **501**
- representable functor, **99**
- restriction of scalars, **508**
- Rezk completion, **604**
- Rezk condition, **603**
- right
 - adjoint object, **541**
 - bounded, **199**
 - complete t-structure, **198**
 - exact, **117**
 - fibration, **577**
 - Kan extension, **549**
 - orthogonal, **451**
 - separated t-structure, **198**
 - t-exact, **190**
- S -local, **255**
- S -nilpotent, **258**
- section of a fibration, **587**
- Segal
 - anima, **602**
 - axiom, **29**
 - condition, **142**
 - map, **142**
- semiadditive ∞ -category, **114**
- semiadditive operad, **479**
- semiorthogonal decomposition, **263**
- Serre fibration, **74**
- shear map, **143**
- shift functor, **113**, **177**
- short exact sequence, **109**, **175**
- sifted ∞ -category, **554**
- simplex category, **41**
- simplicial identities, **41**
- simplicial object, **41**
- singular chain complex, **92**
- singular simplicial complex, **42**
- skeleton, **93**
- slice category, **546**
- small ∞ -category, **49**
- small anima, **49**
- smash product, **80**
- smashing localization, **234**
- span, **27**, **353**
 - category, **353**
 - category of finite sets, **153**
- Spanier–Whitehead dual, **323**
- spectra, ∞ -category of, **129**
- spectrification functor, **125**
- spectrum, **90**, **129**
- spectrum objects, ∞ -category of, **119**
- sphere spectrum, **130**
- spherical fibration, **299**
 - stable, **300**
- split short exact sequence, **115**
- splitting lemma, **115**
- stabilization, **118**
- stable
 - Grassmannian, **282**
 - homotopy groups, **136**
 - ∞ -category, **109**
 - ∞ -category of small stable ∞ -categories, **112**
 - normal bundle, **299**
 - operad, **479**
 - tangent bundle
 - negative, **324**
- stably symmetric monoidal, **322**
- standard t-structure, **190**
- Stiefel manifold, **275**
- straightening, **580**
- strictly initial, **596**
- strongly dualizable, **321**
- structure group, **278**
- structure maps of a spectrum, **90**
- subcategory, **36**
- subcategory inclusion, **37**
- suspension, **79**, **106**
- suspension isomorphism, **88**
- swap map, **80**
- symmetric monoidal
 - adjunction, **386**
 - functor, **243**
 - ∞ -category, **243**
 - model category, **528**
- t-exact, **190**
- t-flat, **205**

- t-projective, **199**
- t-structure, **189**
- telescope, **168**
- tensor product
 - derived, **216**
 - of chain complexes, **216**
 - of spectra, **134**
 - of vector bundles, **273**
 - relative, **501**
- tensor-cofibrant, **528**
- terminal object, **46**
- thick subcategory, **246**
- Thom
 - anima, **299**
 - class, **310**
 - space, **296**
 - spectrum, **301**
- Thom isomorphism, **311, 312**
- topologically enriched category, **606**
- Tor-amplitude, **270**
- Tor-group, **206, 266**
- total space, **278**
- totalization, **48**
- triangle identities, **540**
- trivial
 - fibration, **67**
 - ∞ -operad, **377**
 - operad, **336**
- trivialization, **310**
- twisted arrow category, **591**
- twisted suspension, **297**
- two-sided bar construction, **504**

- uncurrying, **24**
- under category, **546**
- underlying ∞ -category of an operad, **369**
- underlying anima
 - of a spectrum, **129**
 - of a topological space, **13**
- underlying category of an operad, **352**
- unfurling, **398**
- uniqueness, **14**
- unit of a monoid, **142**
- unit of an adjunction, **539**
- unital ∞ -operad, **498**
- universal cocartesian fibration, **49**
- universal coefficient theorem
 - algebraic, **222**
 - for cohomology, **224**
 - for homology, **224**
- universal element, **100**
- universe of animae, **49**
- universe of ∞ -categories, **49**
- unordered concatenation, **343**
- unordered tuple, **343**
- unoriented bordism spectrum, **305**
- unreduced suspension spectrum, **129**
- unstraightening, **580**

- virtual vector bundle, **303**

- weak
 - dual, **321**
 - equivalence, **67**
 - homotopy equivalence, **62**
 - homotopy invariant, **61**
- weakly coextensive, **364**
- weakly extensive, **364**
- wedge, **79**
 - axiom, **99**
- well-pointed, **80**
- Whitehead theorem, **71**
 - for animae, **82**
- Whitney embedding theorem, **324**
- wide subcategory, **354**

- Yoneda embedding, **51**
- Yoneda extension, **201**
- Yoneda lemma, **51**
 - stable, **131**

- zero object, **77, 105**