

(Geometric Singular) Perturbation Theory by Arjen Obelmann

1

perturbation: $\exists \epsilon: 0 < \epsilon \ll 1$

often in the neighborhood of a bifurcation

natural small quantity

- in fluidodynamics viscosity

ex: air at the boundary layer of a plane wing

Reaction-diffusion equations:
$$\begin{cases} v_t = \epsilon^2 v_{xx} + g(u, v) = 0 \\ u_t = u_{xx} + f(u, v) = 0 \end{cases}$$

regular setting (ODE)

$$\begin{cases} \dot{x} = f(x, t; \epsilon) \\ x(0) = a(\epsilon) \end{cases} \quad \text{in } \mathbb{R}^n$$

assume sufficient smoothness

expand $f(x, \epsilon) = f_0(x) + \epsilon f_1(x) + \dots$

$$x(t) = x_0(t) + \epsilon x_1(t) + \dots$$

$$a(\epsilon) = a_0 + \epsilon a_1 + \dots$$

$$\begin{aligned} \frac{d}{dt} x &= \dot{x}_0 + \epsilon \dot{x}_1 + \dots = f_0(x) + \epsilon f_1(x) + \dots \\ &= f_0(x_0 + \epsilon x_1 + \dots) + \epsilon f_1(x_0 + \epsilon x_1 + \dots) + \dots \\ &= f_0(x_0) + \frac{\partial f_0}{\partial x}(x_0) \cdot \epsilon x_1 + \dots + \epsilon f_1(x_0) + \epsilon^2 \frac{\partial f_1}{\partial x}(x_0) \cdot x_1 + \dots \end{aligned}$$

$$\mathcal{O}(1): \begin{cases} \dot{x}_0 = f_0(x_0) \\ x_0(0) = a_0 \end{cases} \quad \text{non-linear}$$

$$\mathcal{O}(\epsilon): \begin{cases} \dot{x}_1 = \frac{\partial f_0}{\partial x}(x_0) \cdot x_1 + f_1(x_0) \\ x_1(0) = a_1 \end{cases} \quad \begin{matrix} \text{linear} \\ + \text{inhomogeneous} \end{matrix}$$

up to $\epsilon^N, N \geq 1$ $x(t) = x_0(t) + \dots + \epsilon^N x_N(t) + \dots$

Poincaré $\forall T > 0 \exists \epsilon_0 > 0$ and $C > 0$ s.t. $\|x - (x_0 + \dots + \epsilon^N x_N)\| < C \epsilon^{N+1} \quad \forall t < T$

Thus 'validity' only up to

$$T = \mathcal{O}(1) \quad \text{w.r.t. } \epsilon$$


Ex. • $\ddot{x} + \epsilon \dot{x} + x = 0$
friction

$O(1)$ it $\sim \frac{1}{\epsilon}$

2

• $\ddot{x} + (1-\epsilon)x = 0$

the perturbed system lags behind!



perturbed


perturbation theory: this approach with a higher T

SINGULAR PERTURBATION

"the $\epsilon=0$ limit systems differ significantly from the $\epsilon \neq 0$ system"

Ex. $\epsilon \dot{x} = x \xrightarrow{\epsilon \downarrow 0} 0 = x$

RELATION
 $x(t) = Ce^{\frac{1}{\epsilon}t}$ $x=$



in the $\dot{x} = f(x, t; \epsilon)$ setting

SP $\iff f(x, t; \epsilon)$ is NOT smooth as function of ϵ , for $\epsilon \rightarrow 0$

$\dot{x} = \frac{1}{\epsilon} x$

$f(x, \epsilon) = \frac{x}{\epsilon}$

\rightarrow that's why it doesn't follow Poincaré

Ex. $\begin{cases} \epsilon \ddot{u} + \dot{u} - u = 0 \\ u(0) = u(1) = 0 \end{cases} \iff \frac{d}{dx}$
B.V.P.
works for $\epsilon \neq 0$

$\epsilon = 0 \quad \dot{u} = u \quad u(t) = Ce^x$

$u(x) = e^{\lambda x}$ $\epsilon \lambda^2 + \lambda - 1 = 0$

$\lambda_{\pm} = \frac{-1 \pm \sqrt{1+4\epsilon}}{2\epsilon}$

$1+2\epsilon + O(\epsilon^2)$

$\lambda_+ = 1 + O(\epsilon)$


$\lambda_- = -\frac{1}{\epsilon} + O(\epsilon)$

at leading order after B.C.

$u(x) = \frac{e-1}{e} e^{-\frac{1}{\epsilon}x} + \frac{1}{e} e^x$

only "non-zero" for x close to 0

for $\epsilon \downarrow 0$, what about the B.V.?



$$\begin{cases} \dot{u} = v \\ \dot{v} = \frac{1}{\varepsilon}(u-v) \end{cases} \quad x \in [0, 1]$$

$$\rightarrow \frac{d}{dt} \begin{pmatrix} u \\ v \end{pmatrix} = \underbrace{\begin{pmatrix} 0 & 1 \\ \frac{1}{\varepsilon} & -\frac{1}{\varepsilon} \end{pmatrix}}_{\text{not smooth}} \begin{pmatrix} u \\ v \end{pmatrix}$$

3

or rescaling time

$$\begin{cases} u' = \varepsilon v \\ v' = (u-v) \end{cases} \quad \text{smooth BUT } x \in [0, \frac{1}{\varepsilon}]$$

outside the Poincaré time scale

eigenvalues


$$\lambda_+ = 1 + O(\varepsilon)$$

$$\lambda_- = -\frac{1}{\varepsilon} + O(1)$$

$$\vec{v}_+ = \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \text{h.o.t.}$$

$$\vec{v}_- =$$

saddle point



General (*)

$$\begin{cases} \dot{x} = f(x, y; \varepsilon) \\ \dot{y} = \varepsilon g(x, y; \varepsilon) \end{cases}$$

FAST SCALING

$$x \in \mathbb{R}^k$$

$$y \in \mathbb{R}^l$$

$$\varepsilon \in \mathbb{R}^n = \mathbb{R}^{k+l}$$

background

$$\begin{cases} u_{xx} + F(u, v) = 0 \\ \varepsilon^2 v_{xx} + G(u, v) = 0 \end{cases} \rightarrow \begin{cases} u_x = p \\ p_x = -F(u, v) \\ \varepsilon v_x = q \\ \varepsilon q_x = -g(u, v) \end{cases}$$

$x \rightarrow X$ rescaling

$$\begin{cases} u_X = \varepsilon p \\ p_X = -\varepsilon F(u, v) \\ v_X = q \\ q_X = -g(u, v) \end{cases}$$


prototype

$$\begin{cases} \dot{x}_1 = x_2 \\ \dot{x}_2 = x_1 - x_1^2 + \varepsilon h(x_1, x_2, y) \\ \dot{y} = \varepsilon g(x_1, x_2, y) \end{cases}$$

(*) Assume I have critical points P_1, P_2 ($P_1 = P_2$)
 can I construct a HOMOCLINIC or HETEROCLINIC orbits?

$\hookrightarrow f(x, y) = g(x, y) = 0$

unbounded domain,
no help from
Poincaré.



(*) rescale 'time' $\rightarrow \begin{cases} \epsilon x' = f(x, y, \epsilon) \\ y' = g(x, y, \epsilon) \end{cases}$
 SLOW SCALING

$\Leftrightarrow \begin{cases} \dot{x} = f(x, y, \epsilon) \\ \dot{y} = g(x, y, \epsilon) \cdot \epsilon \end{cases}$
 FAST SCALING

$\downarrow \epsilon \rightarrow 0$
 Manifold $\mathcal{M}_0 \rightarrow \begin{cases} f(x, y, 0) = 0 \\ y' = g(x, y, 0) \end{cases}$
 SLOW REDUCED LIMIT
 l-dim dynamical system on \mathcal{M}_0



$\downarrow \epsilon \rightarrow 0$
 $\begin{cases} \dot{x} = f(x, y, 0) \\ \dot{y} = 0 \end{cases}$
 FAST REDUCED LIMIT
 $y = y_0$ parameter
 and $\dot{x} = f(x, y_0, 0)$
 l-parameter dyn. sys.

NOTE: \mathcal{M}_0 corresponds to the collection of critical points of the fast reduced limit

Example on the prototype case


$\begin{cases} \ddot{x}_1 = x_2 \\ \ddot{x}_2 = x_1 - x_1^2 + \epsilon h(x_1, x_2, y) \\ \dot{y} = \epsilon g(x_1, x_2, y) \end{cases} \rightarrow \begin{cases} \epsilon \dot{x}_1' = x_2 \\ \epsilon \dot{x}_2' = x_1 - x_1^2 + \epsilon h(x_1, x_2, y) \\ y' = g(x_1, x_2, y) \end{cases}$
 $\downarrow \epsilon \rightarrow 0$

FRL $\begin{cases} \ddot{x}_1 = x_1 - x_1^2 \\ \dot{y} = 0 \end{cases}$





SRL $\begin{cases} 0 = x_2 \\ 0 = x_1 - x_1^2 \\ y' = g(x_1, x_2, y) \\ y' = \begin{cases} g(0, 0, y) \\ g(1, 0, y) \end{cases} \end{cases}$
 if it has 2 critical points, let's assume

$\mathcal{M}_{0,1}$ and $\mathcal{M}_{0,2}$
 $\begin{cases} x_2 = 0 \\ x_1 = 0, +1 \end{cases}$




THEOREM Fenichel 1

If \mathcal{M}_0 corresponds to critical points of FRL that have eigenvalues λ with $\text{Re}(\lambda) \neq 0$, then \mathcal{M}_0 persists as invariant manifold \mathcal{M}_ϵ for the full system. at leading order



THEOREM Fenichel 2

\mathcal{M}_0 has stable and unstable manifolds $W^s(\mathcal{M}_0)$ and $W^u(\mathcal{M}_0)$ if \mathcal{M}_0 is normally hyperbolic ($\text{Re}(\lambda) \neq 0$) then also $W^s(\mathcal{M}_\epsilon)$ and $W^u(\mathcal{M}_\epsilon)$.



Holnikov theory

Heteroclinic $p_1 \rightarrow p_2$ \longrightarrow γ_* homoclinic to \mathcal{M}_ϵ

email: Anjem

Some references:

6

"Geometric singular perturbation theory in biological practice"
by Geertje Hek, 2010, J. Math. Biol.

"An Introduction To Geometric Methods and Dynamical
Systems Theory for Singular Perturbation Theory"
by Tasso J. Kaper, 1999, Proceeds of Symposia in
Applied Mathematics

"Geometric Singular Perturbation Theory"
by Christopher K.R.T. Jones [BOOK]