1 SchemeIBPME

1.1 Setup() \rightarrow (mpk, msk)

```
q \leftarrow \|\mathbb{G}\|

generate g \in \mathbb{G}_1 randomly

generate h \in \mathbb{G}_1 randomly

generate x, \alpha \in \mathbb{Z}_r randomly

H_1 : \{0,1\}^* \to \mathbb{G}_1

H_2 : \{0,1\}^* \to \mathbb{G}_1

H_3 : \{0,1\}^* \to \mathbb{G}_1

H_4 : \{0,1\}^* \to \mathbb{G}_1

H_5 : \{0,1\}^* \to \mathbb{G}_1

H_6 : \{0,1\}^* \to \mathbb{G}_1

H_7 : \{0,1\}^* \to \mathbb{G}_1

y \leftarrow g^x

mpk \leftarrow (G,G_T,q,g,e,h,H_1,H_2,H_3,H_4,H_5,H_6,H_7,y)

msk \leftarrow (x,\alpha)

return (mpk,msk)
```

$1.2 \quad \mathrm{DKGen}(id_R) ightarrow dk_{id_R}$

```
\begin{aligned} dk_{id_R,1} &\leftarrow H_1(id_R)^x \\ dk_{id_R,1} &\leftarrow H_1(id_R)^\alpha \\ dk_{id_R} &\leftarrow (dk_{id_R,1}, dk_{id_R,2}) \\ \mathbf{return} & dk_{id_R} \end{aligned}
```

1.3 $\mathrm{EKGen}(id_S) ightarrow ek_{id_S}$

```
ek_{id_S} \leftarrow H_2(id_S)^{\alpha}

return ek_{id_S}
```

1.4 ReEKGen $(ek_{id_2}, dk_{id_2}, id_1, id_2, id_3) ightarrow rk$

```
generate N \in \{0,1\}^{\lambda} randomly
generate \bar{x} \in \mathbb{Z}_r randomly
rk_1 \leftarrow g^{\bar{x}}
rk_2 \leftarrow dk_{id_2,1}h^{\bar{x}}H_6(e(y,H_1(id_3))^{\bar{x}})
K \leftarrow e(ek_{id_2},H_1(id_3))
rk_3 \leftarrow e(H_2(id_1),H_7(K||id_2||id_3||N) \cdot dk_{id_2,2})
rk \leftarrow (N,rk_1,rk_2,rk_3)
return rk
```

1.5 $\operatorname{Enc}(\boldsymbol{ek_{id_1}}, \boldsymbol{id_2}, m) \rightarrow \boldsymbol{ct}$

```
generate \sigma \in \mathbb{G}_1 randomly
generate \eta \in \mathbb{G}_T randomly
r \leftarrow H_3(m||\sigma||\eta)
ct_1 \leftarrow h^r
ct_2 \leftarrow g^r
ct_3 \leftarrow (m||\sigma) \oplus H_4(e(y, H_1(id_2))^r) \oplus H_4(\eta)
```

```
ct_4 \leftarrow \eta \cdot e(ek_{id_1}, H_1(id_2))
ct_5 \leftarrow H_5(ct_1||ct_2||ct_3||ct_4)^r
ct \leftarrow (ct_1, ct_2, ct_3, ct_4, ct_5)
return ct
```

1.6 ReEnc $(ct, rk) \rightarrow ct'$

If
$$e(ct_1, g) = e(h, ct_2) \wedge e(ct_1, H_5(ct_1||ct_2||ct_3||ct_4)) = e(h, ct_5)$$
 then $ct'_4 \leftarrow \frac{ct_4}{rk_3}$ $ct_7 \leftarrow \frac{e(rk_2, ct_2)}{e(ct_1, rk_1)}$ $ct' \leftarrow (ct_2, ct_3, ct'_4, ct_6, ct_7, N)$ else $ct' \leftarrow \bot$ return ct'

1.7 $\operatorname{Dec}_1(\operatorname{dk}_{\operatorname{id}_2},\operatorname{id}_1,\operatorname{ct}) \to m$

```
If e(ct_1,g) = e(h,ct_2) \wedge e(ct_1,H_5(ct_1||ct_2||ct_3||ct_4)) = e(h,ct_5):

V \leftarrow e(dk_{id_2,2},H_2(id_1))

\eta' \leftarrow \frac{ct_4}{V}

r \leftarrow H_3((ct_3 \oplus H_4(e(dk_{id_2,1})) \oplus H_4(\eta'))||\eta')

If g^r = ct_2:

return m
```

1.8 $\operatorname{Dec}_2(d\mathbf{k}_{id_3}, i\mathbf{d}_1, i\mathbf{d}_2, c\mathbf{t}') \to m$

```
V \leftarrow e(dk_{id_{3},2}, H_{2}(id_{2}))
\eta' \leftarrow ct'_{4} \cdot e(H_{2}(id_{1}), H_{7}(V||id_{2}||id_{3}||N))
R \leftarrow \frac{ct_{7}}{e(H_{6}(e(dk_{id_{3},1},ct_{6}),ct_{2})}
r \leftarrow H_{3}((ct_{3} \oplus H_{4}(R) \oplus H_{4}(\eta'))||\eta')
If g^{r} = ct_{2}:
return m
```