1 SchemeIBPME

This scheme is applicable to symmetric and asymmetric groups of prime orders.

1.1 Setup() \rightarrow (mpk, msk)

```
q \leftarrow \|\mathbb{G}\|
 g \leftarrow 1_{\mathbb{G}_1}
 \hat{g} \leftarrow 1_{\mathbb{G}_2}
 generate s, \alpha, \beta_0, \beta_1 \in \mathbb{Z}_r randomly
 g_1 \leftarrow g^{\alpha}
 f \leftarrow g^{\beta_0}
 \hat{f} \leftarrow \hat{g}^{\beta_0}
 h \leftarrow g^{\beta_1}

\hat{h} \leftarrow \hat{g}^{\beta_1} 

H : \mathbb{G}_T \to \mathbb{Z}_r

 H_1:\{0,1\}^*\to\mathbb{G}_1
 H_2: \{0,1\}^* \to \mathbb{G}_2
H_3: \mathbb{G}_T \to \mathbb{Z}_r
H_4: \{0,1\}^{\lambda} \times \mathbb{G}_T^2 \times \mathbb{G}_1^2 \to \{0,1\}^{\lambda}
H_5: \{0,1\}^{\lambda} \times \mathbb{G}_T^2 \times \mathbb{G}_1^2 \to \{0,1\}^{\lambda}
H_6: \mathbb{G}_T \to \{0,1\}^{3\lambda}
 H_7: \mathbb{G}_T \to \{0,1\}^{2\lambda}
 mpk \leftarrow (g, \hat{g}, g_1, f, h, \hat{f}, \hat{h}, H, H_1, H_2, H_3, H_4, H_5, H_6, H_7)
 msk \leftarrow (s, \alpha)
 return (mpk, msk)
```

1.2 SKGen $(\sigma) \rightarrow ek_{\sigma}$

```
ek_{\sigma} \leftarrow H_1(\sigma)^s

return ek_{\sigma}
```

1.3 $\operatorname{RKGen}(\rho) \to dk_{\rho}$

```
d_1 \leftarrow H_2(\rho)^s
d_2 \leftarrow H_2(\rho)^\alpha
dk_\rho \leftarrow (d_1, d_2)
return dk_\rho
```

1.4 $PKGen(dk_{\rho}, \sigma) \rightarrow pdk_{\rho, \sigma}$

```
generate y \leftarrow \mathbb{Z}_r randomly \eta \leftarrow e(H_1(\sigma), d_1)

y_1 \leftarrow d_2^{H_3(\eta)} (\hat{f}\hat{h}^{H(\eta)})^y

y_2 \leftarrow \hat{g}^y

pdk_{(\rho,\sigma)} \leftarrow (y_1, y_2)

return pdk_{(\rho,\sigma)}
```

1.5 $\operatorname{Enc}(\boldsymbol{ek_{\sigma}}, \boldsymbol{id_{2}}, m) \rightarrow \boldsymbol{ct}$

```
generate r \in \mathbb{Z}_r randomly \eta \leftarrow e(ek_{\sigma}, H_2(\rho)) K_R \leftarrow e(g_1, H_2(\rho))^{r \cdot H_3(\eta)} C_1 \leftarrow g^r C_2 \leftarrow (fh^{H(\eta)})^r K_C \leftarrow H_4(m, \eta, K_R) Y \leftarrow H_5(m, K_C, K_R, C_1, C_2) C_3 \leftarrow (m||K_C||Y) \oplus H_6(K_R) C \leftarrow (C_1, C_2, C_3) return C
```

1.6 $\operatorname{ProxyDec}(pdk, C) \to CT$

```
\begin{split} &K_R \leftarrow e(C_1,y_1)/e(C_2,y_2)\\ &m||K_C||Y \leftarrow C_3 \oplus H_6(K_R)\\ &\textbf{if }Y = H_5(m,K_C,K_R,C_1,C_2)\textbf{then}\\ &CT_1 \leftarrow C_1\\ &CT_2 \leftarrow (m||K_C) \oplus H_7(K_R)\\ &CT \leftarrow (CT_1,CT_2)\\ &\textbf{else}\\ &CT \leftarrow \bot\\ &\textbf{end if}\\ &\textbf{return }CT \end{split}
```

1.7 $\mathbf{Dec}_1(\mathbf{dk}_{\rho}, \sigma, C) \to m$

```
\begin{split} \eta &\leftarrow e(H_1(\sigma), d_1) \\ K_R &\leftarrow e(C_1, d_2^{H_3(\eta)}) \\ m||K_C||Y &\leftarrow C_3 \oplus H_6(K_R) \\ \text{if } K_C &\neq H_4(m, \eta, K_R) \lor Y \neq H_5(m, K_C, K_R, C_1, C_2) \text{then} \\ m &\leftarrow \perp \\ \text{end if} \\ \text{return } m \end{split}
```

1.8 $\operatorname{Dec}_1(\operatorname{dk}_{\rho}, \sigma, \operatorname{CT}) \to m'$

```
\begin{split} \eta &\leftarrow e(H_1(\sigma), d_1) \\ K_R &\leftarrow e(C_1, d_2^{H_3(\eta)}) \\ m||K_C &\leftarrow CT_2 \oplus H_7(K_R) \\ \text{if } K_C \neq H_4(m, \eta, K_R) \text{then} \\ m &\leftarrow \perp \\ \text{end if} \\ \text{return } m \end{split}
```