

TER NOTEBOOK

towards Tate's Thesis

Contents

1	Riemann's zeta function	2
1.1	Riemann's idea I: analytical continuation of zeta function	2
1.2	Riemann's idea II: relation with theta function	5
1.3	Value of Riemann's Zeta function	8
2	L-function	10
2.1	Some preparation	10
2.2	Functional equation of L-function	10
3	Dedekind zeta function	14
3.1	Lattice	14
3.2	Functional Equation	15
4	Hecke L-function	17
4.1	Größencharakter	17
4.2	Ideal number system	22
4.3	Primitive character and Gauss sum	23
5	Tate's Thesis	24
5.1	Adele	24

1 Riemann's zeta function

1.1 Riemann's idea I: analytical continuation of zeta function

As the core object in number theory, it is formally defined as a meromorphic function on a complex plane by Riemann in 1859, in his famous-9-page paper. **This section is the reconstruction of the zeta function along the Riemann's original ideas, it refers to pages 1 and 2 of the original paper.**

$$\sum_{n \geq 1} \frac{1}{n^s} = \prod_p \frac{1}{1 - p^{-s}} \quad (1)$$

This is Euler's product formula proved by Euler (around 1740), with the basic ideas of geometric series

$$\frac{1}{1 - p^{-s}} = 1 + p^{-s} + p^{-2s} + \dots$$

using unique factorization to build a basic correspondence between terms. The formal proof of the formula holds true in half-plane $\Re(s) > 1$ is given by Dirichlet, it is important that the analytic formula makes sense and plays a core role in the analytic number theory. As a student of Dirichlet, Riemann knew the analytic results about the series exactly, and in his era, complex analysis was an advanced branch and analytic continuation was developed completely, hence Riemann tried to treat Euler's formula 1 as an analytic function instead of a formal identity:

$$\zeta(s) = \sum_{n \geq 1} \frac{1}{n^s}, \quad \Re(s) > 1$$

A similar object is the Gamma function, which was called "factorial function" in the age of Gauss and Euler, and Euler noticed an identity of integration:

$$n! = \int_0^\infty e^{-x} x^n dx, \quad n = 1, 2, 3, \dots \quad (2)$$

which motivates Gauss (1813) to introduce the notation

$$\Pi(s) = \int_0^\infty e^{-x} x^s dx, \quad \Re(s) > -1$$

By the modern notation, $\Pi(s - 1) = \Gamma(s)$. Before Riemann developed the zeta function, some important functional equations of the Gamma function were already known, especially the following results:

$$\Gamma(s + 1) = s\Gamma(s) \quad (3)$$

$$\Gamma(s)\Gamma(1 - s) = \frac{\pi}{\sin(\pi s)} \quad (4)$$

$$\Gamma(s)\Gamma\left(s + \frac{1}{2}\right) = 2^{1-2s} \sqrt{\pi} \Gamma(2s) \quad (5)$$

the duplication formula was developed by Legendre (1809), many results about the Gamma function were developed by Euler and Gauss, Riemann surely knew these and he gave the analytic continuation of the zeta function by the following integration:

$$\int_0^\infty e^{-nx} x^{s-1} dx = \frac{\Gamma(s)}{n^s} \quad (6)$$

hence it is natural to consider the summation, it is not difficult to conclude that

$$\int_0^\infty \frac{x^{s-1}}{e^x - 1} dx = \Gamma(s)\zeta(s), \quad \Re(s) > 1 \quad (7)$$

Then Riemann considered the contour integral

$$\int_{\mathcal{H}} \frac{(-z)^{s-1}}{e^z - 1} dz$$

where the contour $\mathcal{H}$ can be drawn as following:




Figure 1: Henkel contour

This type of contour is called **the Henkel contour**, it was systematically studied by Hermann Hankel, who was the student of Riemann and influenced by him greatly. For the sake of convenience, we denote $\delta = \frac{\varepsilon}{\sqrt{2}}$ and $T = \sqrt{R^2 - \varepsilon^2}$, and then we can write

$$\gamma_+(x) = x - i\delta, \quad (-\gamma_-)(x) = x + i\delta$$

for $x \in [\delta, T]$. Notice that $\mathcal{H}$ has negative orientation, and $f : z \mapsto \frac{(-z)^{s-1}}{e^z - 1}$ has simple poles in the imaginary axis, so we choose ε sufficiently small such that $B(0, \varepsilon)$ only contains the pole at zero, then by the residual theorem we have

$$\int_{\mathcal{H}} \frac{(-z)^{s-1}}{e^z - 1} dz = -2\pi i \sum_{|k| \in \mathbb{Z} \cap (0, R)} \text{Res}(f, 2\pi i k) \quad (8)$$

Hence we should evaluate the left hand side. For the small circle, we can estimate that

$$\left| \int_{C_\varepsilon^+} f(z) dz \right| \leq \int_{\pi/4}^{3\pi/4} \frac{\varepsilon^{\Re(s)-1}}{e^{\varepsilon \cos \theta} - 1} d\theta \xrightarrow{\varepsilon \rightarrow 0} 0$$

and for the middle two lines, we take the principal branch of the logarithm Log and write

$$(-z)^{s-1} = e^{(s-1) \text{Log}(-z)}$$

the identity makes sense when $z \notin [0, +\infty)$, hence f is exactly well-defined on the contour, and it allows us to calculate

$$\int_{\gamma_+ \cup \gamma_-} f(z) dz = \int_{\delta}^T \frac{e^{(s-1)\text{Log}(-x+i\delta)}}{e^{x-i\delta} - 1} dx - \int_{\delta}^T \frac{e^{(s-1)\text{Log}(-x-i\delta)}}{e^{x+i\delta} - 1} dx$$

By argument principle, for any $z \in \mathbb{C} - \mathbb{R}_-$ we have

$$\text{Log } |z| = \ln |z| + i \cdot \arg(z)$$

where $\arg$ is a continuous function with value in $(-\pi, \pi)$, which allows us to take limit

$$\text{Log}(-x + i\delta) = \ln \sqrt{x^2 + \delta^2} + i \cdot \arg(-x + i\delta) \xrightarrow{\delta \rightarrow 0} \ln x + i\pi$$

Hence we can evaluate

$$\int_{\gamma_+ \cup \gamma_-} f(z) dz \xrightarrow{\delta \rightarrow 0} (e^{-i\pi s} - e^{i\pi s}) \int_0^T \frac{x^{s-1}}{e^x - 1} dx$$

in particular, when R tends to infinity, we can apply the integration (7) here

$$\int_{\gamma_+ \cup \gamma_-} f(z) dz \xrightarrow{\delta \rightarrow 0, R \rightarrow \infty} -2i \cdot \sin(\pi s) \Gamma(s) \zeta(s) \quad (9)$$

Finally we consider the large circle,

$$\begin{aligned} \left| \int_{C_R^-} f(z) dz \right| &\leq \int_{0+c}^{2\pi-c} \frac{R^{\Re(s)-1}}{e^{R \cos \theta} - 1} d\theta \\ &\xrightarrow{R \rightarrow \infty, \cos(\theta) \neq 0} 0 \end{aligned}$$

In conclusion, by taking limits in (8), and setting (9) into it, we have

$$\begin{aligned} 2 \sin(\pi s) \Gamma(s) \zeta(s) &= 2\pi \sum_{k \in \mathbb{Z} \setminus \{0\}} \text{Res}(f, 2\pi i k) \\ &= 2\pi \sum_{k \in \mathbb{Z} \setminus \{0\}} (-2\pi i k)^{s-1} \\ &= 2 \sin\left(\frac{\pi s}{2}\right) (2\pi)^s \sum_{n \geq 1} \frac{1}{n^{1-s}} \end{aligned}$$

which is the equation given in Riemann's paper, but before the equation he had stated that zeta function is actually meromorphic in the whole complex plane with only one simple pole at $s = 1$, reason is that we just consider the integral on the path $\mathcal{H} - C_R^-$, and we can conclude

$$2 \sin(\pi s) \Gamma(s) \zeta(s) = i \int_{\mathcal{H} - C_R^-} \frac{(-x)^{s-1}}{e^x - 1} dx$$

the right hand side is a entire function and we can get the analytical continuation of zeta. Therefore, Riemann's euqation can be rewritten as

$$\sin(\pi s) \Gamma(s) \zeta(s) = \sin\left(\frac{\pi s}{2}\right) (2\pi)^s \zeta(1-s), \quad s \neq 0, 1 \quad (10)$$

many modern books pointed out that the proof of Riemann is not rigorous enough, but it is not the core of the paper, Riemann noticed the symmetry of the zeta function and gave the functional equation

$$\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s) = \pi^{\frac{1-s}{2}}\Gamma(\frac{1-s}{2})\zeta(1-s)$$

Proof. The short paper do not give the proof, but with a hint of using the properties of Gamma function, so it can be seen as one of "exercises left to the reader" in this paper. Substituting $\sin(\pi s)$ in (10) by the reflection formula (4), we have

$$\zeta(s) = 2^s \pi^{s-1} \sin(\frac{\pi s}{2}) \Gamma(1-s) \zeta(1-s)$$

and substitute $\sin(\frac{\pi s}{2})$ by the duplication formula (5), we have

$$\Gamma(\frac{s}{2})\zeta(s) = \pi^{s-\frac{1}{2}}\Gamma(\frac{1-s}{2})\zeta(1-s)$$

then it is not difficult to rearrange the equation to the desired form. □

1.2 Riemann's idea II: relation with theta function

We can find that zeta function is not so perfectly symmetric as Gamma function, the functional equation has a complicated form with other factor involved, so Riemann introduced the function we talked above, in the sense of modern notation:

$$Z(s) = \pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s)$$

it is so-called **completed zeta function**, with the functional equation $Z(s) = Z(1-s)$. Riemann further considered the integration representation of $Z(s)$ by some work on previous integral (6) and (7), the idea is really straightforward: by just consider the term n^2 we have

$$\frac{\Gamma(s)}{n^{2s}} = \int_0^\infty e^{-n^2 x} x^{s-1} dx$$

we expect the appearance of some factor about π , then we change variable to the right side by setting $x = \pi t$, hence

$$\frac{\Gamma(s)}{n^{2s}} = \pi^s \int_0^\infty e^{-n^2 \pi t} t^{s-1} dt$$

without talking about the interchange of integration, we rearrange and sum over n we can get

$$Z(s) = \int_0^\infty \left(\sum_{n \geq 1} e^{-n^2 \pi t} \right) t^{\frac{s}{2}-1} dt$$

Riemann noticed that the summation is closely related to the theta function, and hence he cited the formula of Jacobi to get the analytical continuation of $Z(s)$ and define the xi-function, which is the core object in his following discussion. We stop here for the moment, the page 3 is so rich that we need to rewrite it carefully in modern notation, the reason is as following:

- Maybe Riemann do not realize that he actually gave the analytic continuation of zeta function again, because completed zeta function

Definition 1. The **Jacobi theta function** is defined as

$$\theta(z) = \sum_{n \in \mathbb{Z}} e^{\pi i n^2 z}$$

it is holomorphic in the upper half-plane $\mathbb{H} = \{z \in \mathbb{C} : \Im(z) > 0\}$.

It is not difficult to see that

$$|e^{\pi i n^2 (x+iy)}| = |e^{-\pi n^2 y}|$$

so the series convergence absolutely and uniformly on any compact subset $\mathbb{H}$, so θ is exactly holomorphic, and it has a important functional equation, which was first proved by Jacobi (1829), and Riemann cited it in his paper without proof.

Lemma 2. (Jacobi) For any $z \in \mathbb{H}$, we have

$$\theta(-1/z) = \sqrt{z/i} \cdot \theta(z)$$

In particular, when $z = it$ for $t > 0$, i.e. on the positive imaginary axis, θ function has a deep relation with the completed zeta function:

Proposition 3. For $\Re(s) > 1$, completed zeta function is holomorphic and

$$Z(s) = \frac{1}{2} \int_0^\infty (\theta(it) - 1) t^{\frac{s}{2}-1} dt$$

Proof. ζ and Γ are both holomorphic on $\Re(s) > 1$, and $s \mapsto \pi i^{-\frac{s}{2}}$ is entire, so Z is holomorphic on $\Re(s) > 1$. By definition of θ function, we have

$$\frac{\theta(it) - 1}{2} = \sum_{n \geq 1} e^{-n^2 \pi t}$$

which allows us to rewrite

$$\frac{1}{2} \int_0^\infty (\theta(it) - 1) t^{\frac{s}{2}-1} dt = \int_0^\infty \left(\sum_{n \geq 1} e^{-n^2 \pi t} \right) t^{\frac{s}{2}-1} dt$$

and if we admits the integration representation (7) and calculate as above we can get

$$Z(s) = \sum_{n \geq 1} \int_0^\infty e^{-\pi n^2 t} t^{\frac{s}{2}-1} dt$$

for $\Re(s) > 1$, so we should verify the interchange of summation and integration.

$$\left| \sum_{n \geq 1} e^{-\pi n^2 t} t^{s/2-1} \right| \leq \sum_{n \geq 1} e^{-\pi n t} t^a = \frac{t^a}{e^{\pi t} - 1} \sim_{t \rightarrow \infty} t^a e^{-\pi t}$$

with $a = \Re(s)/2 - 1 > -1/2$, so the integration converges at infinity, and by Jacobi's formula we can estimate the integration near zero:

$$\theta(it) = (t)^{-1/2} \theta(i/t) = t^{-1/2} \sum_{k \in \mathbb{Z}} e^{-\pi k^2/t} \xrightarrow[t \rightarrow 0]{} t^{-1/2}$$

which allows us to get a global bound function for some constant $C > 0$:

$$|\sum_{n \geq 1} e^{-\pi n^2 t} t^{s/2-1}| \leq g(t) = \begin{cases} C t^a e^{-\pi t} & t \geq 1 \\ C t^{(a-1/2)} & 0 < t < 1 \end{cases}$$

and g is measurable and integrable on $(0, \infty)$, hence by Dominated Convergence Theorem we can interchange the summation and integration. $\square$

with the integral representation, completed zeta function can extend meromorphically to the whole complex plane like what we do to Gamma function, for the sake of convenience, Riemann introduced the auxiliary function:

$$\psi(t) = \frac{\theta(it) - 1}{2}, \quad t > 0$$

by the Jacobi's functional equation, we can get the reflection formula

$$t^{1/2}(2\psi(t) + 1) = 2\psi(1/t) + 1, \quad t > 0 \quad (11)$$

with that we can get the following result which is roughly sketched in Riemann's paper:

Theorem 4. The completed zeta function Z has a analytic continuation to complex plane $\mathbb{C}$, and

- it only has simple poles at $s = 0$ and $s = 1$;
- residue number: $\text{Res}(Z, 0) = -1$ and $\text{Res}(Z, 1) = 1$;
- it satisfies the functional equation $Z(s) = Z(1-s)$ for $s \neq 0, 1$.

Proof. The proof is based on the integral representation on $\Re(s) > 1$, we divide the integration into two parts and change variable

$$\begin{aligned} Z(s) &= \int_0^1 \psi(t) t^{\frac{s}{2}-1} dt + \int_1^\infty \psi(t) t^{\frac{s}{2}-1} dt \\ &= \int_1^\infty \psi(1/u) u^{-\frac{s}{2}-1} du + \int_1^\infty \psi(t) t^{\frac{s}{2}-1} dt \end{aligned}$$

then apply the formula (11) to the first term, we have

$$\begin{aligned} Z(s) &= \frac{1}{2} \int_1^\infty t^{-s/2-1/2} dt - \frac{1}{2} \int_1^\infty t^{s/2-1} dt + \int_1^\infty \psi(t) (t^{s/2-1} + t^{-s/2-1/2}) dt \\ &= \frac{1}{s-1} - \frac{1}{s} + \int_1^\infty \psi(t) (t^{s/2-1} + t^{-s/2-1/2}) dt \\ &= \frac{1}{s-1} - \frac{1}{s} + F(s) + F(1-s) \end{aligned}$$

where $F(s) = \int_1^\infty \psi(t) t^{s/2-1} dt$ is an entire function, which can be proved by theorem of convergence dominated (wait to complete...). Hence we can get the analytic continuation of Z to the whole complex plane, with only simple poles at $s = 0$ and $s = 1$ left, and the residue number can be calculated by the formula above. Finally, the functional equation when $s \mapsto 1-s$. $\square$

Corollary 5. Riemann zeta function ζ has a analytic continuation to the whole complex plane except for a simple pole at $s = 1$ with residue number 1, with functional equation

$$\zeta(1-s) = 2(2\pi)^s \cos(\pi s/2) \Gamma(s) \zeta(s) \quad s \neq 1, 0, -1, -2, \dots$$

Proof. By completed zeta function, we have

$$\zeta(s) = \frac{Z(s)}{\pi^{-s/2} \Gamma\left(\frac{s}{2}\right)}$$

notice that Z and Γ have both simple poles at $s = 0$, so ζ is actually holomorphic at $s = 0$, with only simple pole at $s = 1$ left by $Z(s)$, that because Γ has no zeros and $s \mapsto \pi^{-s/2}$ is entire. $\square$

Remark 6. The method of analytic continuation via integral representation can be generalized, it is so called Mellin transformation, an important and natural integration transform: Let f be a positive real-valued function, then its **Mellin transform** is defined as

$$\mathcal{M}(f, s) = \int_0^\infty f(t) t^{s-1} dt$$

here completed zeta function is just the Mellin transform of ψ function.

1.3 Value of Riemann's Zeta function

This section is to give a basic description of the distribution of values of Riemann zeta function, mainly focus on the value of $\mathbb{Z}$, and then the result of Hadamard about the critical strip.

Firstly, the value at the positive odd integer of zeta function is hard and few results about that we can know. In 1978, Apéry proved that $\zeta(3)$ is irrational; In 2001, Rivoal proved that there exists infinite irrational number in $\{\zeta(2k+1) \mid k \in \mathbb{N}\}$, and he gave the lower bound of the cardinal (as the scale of $\log k$); Another folklore conjecture about the irrationality is that $\zeta(2k+1)$ is algebraical independent over $\mathbb{Q}$, it is the corollary of the famous **Grothendieck's period conjecture**, which is still open.

The simplest case is at the negative even integer, which is concluded in Riemann's paper as a small comments: $\zeta(-2k) = 0$ for any $k \in \mathbb{N}$.

Proof. The proof is based on the functional equation of zeta function, it refers to corollary 5, just take $1-s = -2k$, for $k \in \mathbb{N}$; then we can find that $\cos(\pi s/2) = 0$, and we can conclude the result. A little remark here is that the method can not be extended to the positive odd integer, the reason is that Γ is only well-defined on the positive integer, which shows that the lack of information is the difficulty of the question at positive odd integer. $\square$

The value at negative odd integers and at positive even integers are the same story, that is clear by the functional equation, if we set $1-s = -2k-1$ for $k \in \mathbb{N}$, then we can conclude

$$\zeta(-2k-1) = 2(2\pi)^{2k+4} (-1)^{k+1} (2k+1)! \zeta(2k+2)$$

so our attention will focus on the values of positive even integers firstly. Indeed the values of zeta function at $\mathbb{Z}$ has a deep relation with Bernoulli's number, formally we define:

Definition 7. The function $z \mapsto \frac{z}{e^z - 1}$ is holomorphic at the voisinage of zero, with the expansion of the series:

$$\frac{z}{e^z - 1} = \sum_{n \geq 0} \frac{B_n}{n!} z^n$$

the n -th **Bernoulli's number** is defined as B_n , with respect to the coefficient of the series.

The original definition of Bernoulli's number is given by Jacob Bernoulli in his book "Ars Conjectandi" (1713), he defined the number by the summation of powers:

$$\sum_{k=1}^m k^n = \frac{1}{n+1} \sum_{j=0}^n \binom{n+1}{j} B_j \cdot m^{n+1-j}$$

for any $n, m \in \mathbb{N}$, and the two definitions can be shown to be equivalent by the method of **generating function**. The definition of power series is easy for us to calculate the values of Bernoulli's number, for Example

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_3 = 0, \quad B_4 = -\frac{1}{30} \quad \dots$$

Lemma 8. The Bernoulli's number is rational, and $B_{2k+3} = 0$ for any $k \in \mathbb{N}$.

Proof. Observe that

$$\frac{ze^z}{e^z - 1} = \frac{-z}{e^{-z} - 1}$$

which allows us to write

$$z = \frac{ze^z}{e^z - 1} - \frac{z}{e^z - 1} = \sum_{n \geq 0} ((-1)^n - 1) \frac{B_n}{n!} \cdot z^n$$

at the voisinage of zero, hence by comparing the coefficients we can conclude that $B_0 = 1$ and $B_n = 0$ for any odd integer $n \geq 3$. To prove the rationality, by the properties of analytical function we know

$$B_n = n! \frac{d^n}{dz^n} \left(\frac{z}{e^z - 1} \right) \Big|_{z=0}$$

the derivatives can be proved to be a rational number by induction on n , hence the result holds true. $\square$

With the notation of the Bernoulli's number, the values can be written as a neat form

Proposition 9. Let $k \in \mathbb{N}$, then

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2(2k)!} \cdot B_{2k}$$

$$\zeta(-2k+1) = \frac{B_{2k}}{-2k}$$

which shows that the values at negative odd integers are rational numbers, while the values at positive even integers are transcendental numbers (rational number $\times$ the factor π).

Proof. One way to prove the result is to use the formula of cotangent function

$$\pi z \cot(\pi z) = 1 + \sum_{n=1}^{\infty} \frac{2z^2}{z^2 - n^2} \quad z \in \mathbb{C} - \mathbb{Z}$$

using geometric series we can rewrite the formula

$$\begin{aligned} \pi z \cot(\pi z) &= 1 + 2z^2 \sum_{n \geq 1} \frac{-1}{n^2} \cdot \frac{1}{1 - z^2/n^2} \\ &= 1 - 2z^2 \sum_{n \geq 1} \frac{1}{n^2} \sum_{k \geq 0} \left(\frac{z^2}{n^2} \right)^k \\ &= 1 - 2 \sum_{k \geq 1} \zeta(2k) z^{2k} \end{aligned}$$

here the formula holds true for $|z| < 1$, and the last equality holds by Fubini's theorem, so we get a power series expansion at zeros. To finish the proof, we write the trigonometric function in terms of exponential function, then we can calculate

$$z \cot(z) = \frac{2iz}{e^{2iz} - 1} + iz = 1 + \sum_{k=1}^{\infty} (-1)^k \frac{2^{2k}}{(2k)!} B_{2k} \cdot z^{2k}$$

which allows us to get another power series expansion at zeros, if we replace z by πz and compare the two expansions, then we can conclude the first formula. The second formula can be concluded by the functional equation of zeta function. $\square$

2 L-function

2.1 Some preparation

2.2 Functional equation of L-function

Review the functional equation of Riemann zeta function and its analytical continuation, we can conclude the method by two steps: Firstly, we find an integral representation of zeta function by the Gamma function, and rearrange the integration to get the integral representation of completed zeta function, it gives a relation with theta function; Secondly, we use the functional equation of theta function to rewrite the integral, which allows us to get the functional equation of completed zeta function, and hence the functional equation of zeta function.

For any Dirichlet character χ modulo q , there exists a conductor q' which divides q such that

$$\chi = \chi' \cdot 1$$

with χ' a primitive Dirichlet character modulo q' , and 1 the trivial character modulo q . Hence for any L-function on χ , we can write

$$L(s, \chi) = L(s, \chi') \cdot \prod_{p|q} (1 - \chi(p)p^{-s})$$

the last term defines a entire function, so the analytic continuation will be reduced to the case of primitive character.

The method is so natural that it can be generalized to the L-function, and we start from the integral

$$\pi^{-\frac{s}{2}} \Gamma(\frac{s}{2}) \frac{1}{n^s} = \int_0^\infty e^{-\pi n^2 t} t^{\frac{s}{2}-1} dt \quad \Re(s) > 1$$

if we sum over n , that will be what we do in the last section, but now we want to sum over n with **some primitive Dirichlet character** χ modulo q , and we expect the auxillary function is a certain arithmetic function, with at least the information of the character (periodic q).

$$(q/\pi)^{s/2} \Gamma(\frac{s}{2}) \frac{\chi(n)}{n^s} = \frac{1}{\sqrt{q}} \int_0^\infty [\chi(n) e^{-\pi n^2 t/p}] t^{\frac{s}{2}-1} dt \quad (12)$$

what we do is to change the variable by setting $t \rightarrow t/q$, hence we can sum over n and get

$$(q/\pi)^{s/2} \Gamma(\frac{s}{2}) L(s, \chi) = \frac{1}{\sqrt{q}} \int_0^\infty [\sum_{n \geq 1} \chi(n) e^{-\pi n^2 t/q}] t^{\frac{s}{2}-1} dt$$

we can extend the sum to the whole integer, but it depends on the values of $\chi(-1)$. If $\chi(-1) = 1$, then we can write

$$\sum_{n \geq 1} \chi(n) e^{-\pi n^2 t/q} = \frac{1}{2} \sum_{n \in \mathbb{Z}} \chi(n) e^{-\pi n^2 t/q}$$

and if $\chi(-1) = -1$ we find that the right term above is zero, it makes no sense, hence we should modify the integration (12) a little to avoid the problem, we can introduce a translation on s by setting $s \rightarrow s+1$, then we can write

$$(q/\pi)^{(s+1)/2} \Gamma(\frac{s+1}{2}) \frac{\chi(n)}{n^s} = \frac{1}{\sqrt{q}} \int_0^\infty [n \chi(n) e^{-\pi n^2 t/p}] t^{\frac{s-1}{2}} dt \quad (13)$$

then the sum over n allows us to get

$$\sum_{n \geq 1} n \chi(n) e^{-\pi n^2 t/q} = \frac{1}{2} \sum_{n \in \mathbb{Z}} n \chi(n) e^{-\pi n^2 t/q}$$

It is the basic idea of the proof, similarly we introduce the **auxillary function**

$$\psi(t, \chi) = \sum_{n \in \mathbb{Z}} n^{\mathfrak{v}} \chi(n) e^{-\pi n^2 t/q}$$

with $\mathfrak{v} = \frac{1-\chi(-1)}{2}$, then we can deduce the functional equation via Poisson's formula.

Lemma 10. let $t > 0$, then ψ satisfies the equation

$$\tau(\bar{\chi}) \psi(t, \chi) = (i/t)^{\mathfrak{v}} \sqrt{q/t} \cdot \psi(\bar{\chi}, 1/t)$$

where $\tau(\chi) = G(1, \chi)$ as the Gauss sum of χ .

Proof. We consider the $f(x) = e^{-\pi x^2 t}$ for some $t > 0$, and then we can calculate

$$\hat{f}(s) = \frac{1}{\sqrt{t}} f\left(\frac{s}{t}\right) = \frac{1}{\sqrt{t}} e^{-\pi s^2/t}$$

hence by poisson's formula we have

$$\sum_{n \in \mathbb{Z}} e^{-\pi(n+a)^2 t} = \frac{1}{\sqrt{t}} \sum_{n \in \mathbb{Z}} e^{-\pi n^2/t} e^{2\pi i n a} \quad (14)$$

now we consider χ is a arithmetic function on $\mathbb{Z}$ with period q , then we can write

$$\chi(n) = \frac{\tau(\chi)}{q} \sum_{m=1}^q \bar{\chi}(m) e^{2\pi i m n/q}$$

by the primitive character $\overline{\tau(\chi)} = \chi(-1)\tau(\bar{\chi})$, we have

$$\tau(\bar{\chi})\chi(n) = \sum_{m=1}^q \bar{\chi}(m) e^{2\pi i m n/q}$$

hence we can calculate

$$\begin{aligned} \tau(\bar{\chi})\psi(t, \chi) &= \sum_{n \in \mathbb{Z}} n^{\mathfrak{v}} \tau(\bar{\chi})\chi(n) e^{-\pi n^2 t/q} \\ &= \sum_{m=1}^q \bar{\chi}(m) \sum_{n \in \mathbb{Z}} n^{\mathfrak{v}} e^{-\pi n^2 t/q} e^{2\pi i m n/q} \end{aligned}$$

when $\chi(-1) = 1$, i.e. $\mathfrak{v} = 0$, we can apply the Poisson's formula (14) to get

$$\begin{aligned} \tau(\bar{\chi})\psi(t, \chi) &= \sqrt{q/t} \cdot \sum_{m=1}^q \bar{\chi}(m) \cdot \sum_{n \in \mathbb{Z}} e^{-\pi(qn+m)^2/qt} \\ &= \sqrt{q/t} \cdot \sum_{l \in \mathbb{Z}} \bar{\chi}(l) e^{-\pi l^2/t} \\ &= \sqrt{q/t} \cdot \psi(\bar{\chi}, 1/t) \end{aligned}$$

when $\chi(-1) = -1$, i.e. $\mathfrak{v} = 1$, we can modify the poisson's formula (14) by taking derivative of $f(x)$, then we have

$$it^{3/2} \sum_{n \in \mathbb{Z}} (n+a) e^{-\pi(n+a)^2 t} = \sum_{n \in \mathbb{Z}} n e^{-\pi n^2/t} e^{2\pi i n a}$$

then apply it we can get

$$\begin{aligned} \tau(\bar{\chi})\psi(t, \chi) &= \sum_{m=1}^q \bar{\chi}(m) \cdot i(q/t)^{3/2} \sum_{n \in \mathbb{Z}} \left(n + \frac{m}{q}\right) e^{-\pi(qn+m)^2/qt} \\ &= (i/t) \sqrt{q/t} \cdot \sum_{l \in \mathbb{Z}} l \cdot \bar{\chi}(l) e^{-\pi l^2/t} \\ &= (i/t) \sqrt{q/t} \cdot \psi(\bar{\chi}, 1/t) \end{aligned}$$

□

With the functional equation of ψ , it is sufficient to get the functional equation of L function, firstly we define the **completed L function** by

$$\Lambda(s, \chi) = (q/\pi)^{\frac{s+\mathfrak{v}}{2}} \Gamma\left(\frac{s+\mathfrak{v}}{2}\right) L(s, \chi)$$

by (12) and (13), we can write down the integral representation

$$\Lambda(s, \chi) = \frac{1}{2\sqrt{q}} \int_0^\infty \psi(t, \chi) t^{\frac{s+\mathfrak{v}}{2}-1} dt$$

when $\Re(s) > 1$, and then by the functional equation of ψ , we can get

Theorem 11. For any **non-principal** primitive Dirichlet character χ modulo q , the completed L function $\Lambda(s, \chi)$ has an analytic continuation to the whole complex plane, and it satisfies the functional equation

$$\Lambda(1-s, \bar{\chi}) = \frac{i^{\mathfrak{v}} \sqrt{q}}{\tau(\chi)} \Lambda(s, \chi)$$

Proof. when $\chi(-1) = 1$, i.e. $\mathfrak{v} = 0$, we have

$$\begin{aligned} 2\sqrt{q}\Lambda(s, \chi) &= \int_0^1 \psi(t, \chi) t^{\frac{s}{2}-1} dt + \int_1^\infty \psi(t, \chi) t^{\frac{s}{2}-1} dt \\ &= \int_1^\infty \psi(1/u, \chi) u^{-\frac{s}{2}-1} du + \int_1^\infty \psi(t, \chi) t^{\frac{s}{2}-1} dt \\ &= \frac{\tau(\chi)}{\sqrt{q}} \int_1^\infty \psi(u, \bar{\chi}) u^{-\frac{s-1}{2}} du + \int_1^\infty \psi(t, \chi) t^{\frac{s}{2}-1} dt \end{aligned}$$

notice that in this case we have $\overline{\tau(\chi)} = \tau(\bar{\chi})$, so we can write

$$\begin{aligned} 2\sqrt{q}\Lambda(s, \chi) &= \frac{\tau(\chi)}{\sqrt{q}} \left[\int_1^\infty \psi(u, \bar{\chi}) u^{-\frac{s-1}{2}} du + \frac{\tau(\bar{\chi})}{\sqrt{q}} \int_1^\infty \psi(t, \bar{\chi}) t^{-\frac{s-1}{2}-1} dt \right] \\ &= \frac{\tau(\chi)}{\sqrt{q}} \cdot 2\sqrt{q}\Lambda(1-s, \bar{\chi}) \end{aligned}$$

Finally, when $\chi(-1) = -1$, i.e. $\mathfrak{v} = 1$, we have

$$\begin{aligned} 2\sqrt{q}\Lambda(s, \chi) &= \int_0^1 \psi(t, \chi) t^{\frac{s-1}{2}} dt + \int_1^\infty \psi(t, \chi) t^{\frac{s-1}{2}} dt \\ &= \int_1^\infty \psi(1/u, \chi) u^{-\frac{s+3}{2}} du + \int_1^\infty \psi(t, \chi) t^{\frac{s-1}{2}} dt \\ &= \frac{\tau(\chi)}{i\sqrt{q}} \int_1^\infty \psi(u, \bar{\chi}) u^{-\frac{s}{2}} du + \int_1^\infty \psi(t, \chi) t^{\frac{s-1}{2}} dt \end{aligned}$$

and notice that in this case we have $\overline{\tau(\chi)} = -\tau(\bar{\chi})$, so we can write

$$\begin{aligned} 2\sqrt{q}\Lambda(s, \chi) &= \frac{\tau(\chi)}{i\sqrt{q}} \left[\int_1^\infty \psi(u, \bar{\chi}) u^{-\frac{s}{2}} du + \frac{\tau(\bar{\chi})}{i\sqrt{q}} \int_1^\infty \psi(t, \bar{\chi}) t^{\frac{s-1}{2}} dt \right] \\ &= \frac{\tau(\chi)}{i\sqrt{q}} \cdot 2\sqrt{q}\Lambda(1-s, \bar{\chi}) \end{aligned}$$

□

similarly, we can conclude that the

3 Dedekind zeta function

3.1 Lattice

Definition 12. Let $(V, \langle \cdot, \cdot \rangle)$ be a finite-dimensional inner product space over $\mathbb{R}$:

- A **lattice** L in V is a discrete subgroup of V .

- A lattice L is called **complete** if it is a free $\mathbb{Z}$ -module of rank $n = \dim V$ with basis $\{v_1, \dots, v_n\}$, its fundamental domain is defined as

$$P = \left\{ \sum_i^n c_i v_i \mid c_i \in [0, 1) \right\}$$

- For any lattice $L \subset V$, the dual lattice under the inner product is defined as

$$L^* = \{v \in V \mid \langle v, w \rangle \in \mathbb{Z}, \forall w \in L\}$$

Notice that V/L is compact if and only if L is complete, hence we can define the **volume** of V/L by

$$\text{Vol}(V/L) = m(P) = \sqrt{\det G} \cdot l(P)$$

where m is unique Haar measure induced by the inner product while l is the usual Lebesgue measure, and G is the Gram matrix with entries $\langle v_i, v_j \rangle$.

Lemma 13.

$$\text{Vol}(V/L) \cdot \text{Vol}(V/L^*) = 1$$

Proof. □

Theorem 14 (Poisson's formula for lattice). Let L be a complete lattice in V , then for any Schwartz function f on V , we have

$$\sum_{v \in L} f(v) = \frac{1}{\text{vol}(V/L)} \sum_{v^* \in L^*} \hat{f}(v^*)$$

Proof. □

Corollary 15. Let A be a positive-definite symmetric matrix, and $L \subset V$ be a complete lattice, then

$$\sum_{v \in L} e^{-\pi \cdot v^t A v} = \frac{\det A^{-1/2}}{\text{vol}(V/L)} \sum_{v \in L^*} e^{-\pi \cdot v^t A^{-1} v}$$

In particular, if $A = \text{diag}(c_1, \dots, c_n)$ and $L = \mathbb{Z}^n$, then

$$\sum_{v \in \mathbb{Z}^n} e^{-\pi \sum_i c_i v_i^2} = \frac{1}{\sqrt{c_1 \cdots c_n}} \sum_{v \in \mathbb{Z}^n} e^{-\pi \sum_i v_i^2 / c_i}$$

Proof. □

Now we consider the case of number field, suppose that $k/\mathbb{Q}$ is a number field with degree N , and we embed k into $\mathbb{R}^N$ by

$$\sigma : k \rightarrow \mathbb{R}^s \times \mathbb{C}^t, \quad x \mapsto (\sigma_i(x))_i$$

view $\mathbb{C}$ as a real vector space and then we can define inner product by trace

$$\langle \sigma(x), \sigma(y) \rangle = \text{Tr}_{k/\mathbb{Q}}(xy)$$

Notice that $\mathcal{O}_k$ is a free $\mathbb{Z}$ -module of rank N , so $\sigma(\mathcal{O}_k)$ is a complete lattice, and it motivates us to define its dual lattice.

Definition 16. The inverse different of k is a fractional ideal defined by

$$\mathfrak{d}^{-1} = \{x \in k \mid \text{Tr}_{k/\mathbb{Q}}(x\mathcal{O}_k) \subset \mathbb{Z}\}$$

Lemma 17. Along the notion above,

1. $\sigma(\mathfrak{d}^{-1})$ is the dual lattice of $\sigma(\mathcal{O}_k)$ under the inner product defined by trace.
2. $\mathfrak{d} := (\mathfrak{d}^{-1})^{-1}$ is an ideal of $\mathcal{O}_k$, and it is called the **different** of k .
3. The norm of different is the discriminant of the field $N(\mathfrak{d}) = |d_k|$
4. For any ideal $\mathfrak{a}$, we set $\mathfrak{a}^\vee = \mathfrak{d}^{-1}\mathfrak{a}^{-1}$, then $\sigma(\mathfrak{a}^\vee)$ is the dual lattice of $\sigma(\mathfrak{a})$.

Proposition 18. Let $\mathfrak{a}$ be an fractional ideal of k , and define

$$\Theta(c, \mathfrak{a}) = \sum_{z \in \mathfrak{a}} \exp(-\pi \sum_{c_i})$$

where $\|\cdot\|$ is the norm defined by trace paring, then we have the equation

$$\Theta(z, \mathfrak{a}) = \frac{z^{-N/2}}{N(\mathfrak{a})\sqrt{|d_k|}} \Theta(1/z, \mathfrak{a}^\vee)$$

3.2 Functional Equation

Suppose that $k/\mathbb{Q}$ is a number field with degree N , we make the convention for the notion:

1. r and t are the number of real and complex embedding of k respectively, so $N = r + 2t$;
2. $C_k = J_k/P_k$ is the ideal class group of k , and $h_k = |C_k|$ is the class number of k ;
3. w_k is the number of roots of unity in k , r_k is the regulator of k , and d_k is the discriminant of k .

For any ideal class $R \in C_k$, we take an representative $\mathfrak{a}$ of R^{-1} , then for any integral ideal $\mathfrak{b} \in R$ we have a well-defined applicaion to P_k :

$$\mathfrak{b} \mapsto \mathfrak{a}\mathfrak{b} = (x)$$

notice that $\mathfrak{b} \subset \mathcal{O}_k$ as an integral ideal, so $x \in \mathfrak{a}^*$. Conversely, for any $x \in \mathfrak{a}^*$ we can set $\mathfrak{b} = a^{-1}(x)$, then $\mathfrak{b} \subset \mathcal{O}_k^*$ implies $\mathfrak{b}$ is exactly a integral ideal, so the application is surjective to $\mathfrak{a}^*$. For a bijection, we consider an equivalence relation on $\mathfrak{a}^*$ by setting $x \sim x'$ if and only if $x' = ux$ for some unit u , i.e. two elements are associative, it is equivalent to the the action of $\mathcal{O}_k^*$ on $\mathfrak{a}^*$, hence we can get a bijection

$$\{\text{The integral ideal of } R\} \longleftrightarrow \mathfrak{a}^*/\mathcal{O}_k^*$$

the right term is the orbit space of the action. Hence we can rewrite the Dedekind zeta function

$$\zeta_k(s) = \sum_{R \in C_k} \zeta(s, R)$$

as the sum of function on ideal classes.

$$\zeta(s, R) = N(\mathfrak{a})^s \sum_{x \in \mathfrak{a}^*/\mathcal{O}_k^*} \frac{1}{|N(x)|^s} = (d_{\mathfrak{a}}/d_k)^{1/2} \sum \dots$$

where $\mathfrak{a}$ is the representative of R^{-1} , Hence it motivates us to find the functional equation of zeta function on ideal class.

We consider the minkowski space $\mathbb{R}^r \times \mathbb{C}^t$, then in it we can denote the norm as following

$$|N(x)| = \prod_v |x_v|^{a_v}$$

with $x_v = \sigma_v(x)$ and v runs over embeddings modulo conjugation, and

$$a_v = \begin{cases} 1 & \text{if } \sigma_v \text{ is a real embedding} \\ 2 & \text{if } \sigma_v \text{ is a complex embedding} \end{cases}$$

Proposition 19. With the notion above, we set

$$A = 2^{-t} d_k^{-1/2} \pi^{-N/2}$$

and let

$$\zeta(s, \infty) = A^s \Gamma(s/2)^r \Gamma(s)^t$$

then define the completed ideal class zeta function

$$\Lambda(s, R) = \zeta(s, \infty) \zeta(s, R)$$

Then $\Lambda(s, R)$ attains an analytical continuation on complex plane with only simple poles at $s = 0$ and $s = 1$, and it satisfies a functional equation for any $s \neq 0, 1$

$$\Lambda(1-s, R^{-1}) = \Lambda(s, R)$$

Proof. (1) Representation of function: We consider the haar measure on $\mathbb{R}_{>0}$ and definition of Gamma function, it sufficient to prove that

$$\frac{\Gamma(s/2)}{a^s} = \int_{\mathbb{R}_{>0}} e^{-a^2 y} y^{s/2} \frac{dy}{y}$$

Applying this formula to $a^2 = \dots$, it is enough to get

$$(\pi^{-1/2} d_k^{1/2N} N(\mathfrak{a})^{1/N})^s = \int_{\mathbb{R}_{>0}} \exp(-\pi d_{\mathfrak{a}}^{-1/N} |x_v|^2 y) y^{s/2} \frac{dy}{y} \quad (15)$$

For real parts we will multiply (15) r times, while for complex embedding, we should let $s \rightarrow 2s$, and change variable in integral such that $y \mapsto 2y$. The first operation is because we will only multiply complex parts t times while $N = r + 2t$; The second operation is necessary, because we expect a fourier transform with respects to the inner product in Minkowski space, hence for complex parts $|x_v|$ we have

$$(2^{-1} \pi^{-1} d_k^{1/N} N(\mathfrak{a})^{2/N})^s = \int_{\mathbb{R}_{>0}} \exp(-\pi d_{\mathfrak{a}}^{-1/N} \cdot 2|x_v|^2 y) y^s \frac{dy}{y} \quad (16)$$

As we mentioned, consider (15) $\times r + (16) \times t$ and sum over $\mathfrak{a}^*/\mathcal{O}_k^*$ we have

$$\Lambda(s, R) = \int_G \sum_x \exp(-\pi d_{\mathfrak{a}}^{1/N} \cdot \langle x, y \rangle_m) \|y\|^{s/2} d^*y$$

where $G = \mathbb{R}_{>0}^{s+t}$ as a locally compact group under the restricted product with Harr measure such that

$$d^*y = \prod_v \frac{dy_v}{y_v}$$

(2) Decomposition of integration: We now consider the spherical decomposition of group $G = \mathbb{R}_{>0} \times G_0$ with

$$G_0 = \{x \in G \mid \|x\| = 1\}$$

i.e. we set $y = l^{1/N} c$ with $l = \|y\|$, then we have

$$\Lambda(s, R) = \int_{\mathbb{R}_{>0}} \int_{G_0} \exp(-\pi d_{\mathfrak{a}}^{1/N} \cdot \langle x, l^{1/N} c \rangle) d^*c \quad t^{s/2} \frac{dt}{t}$$

□

4 Hecke L-function

4.1 Größencharakter

As we have done for Riemann zeta function, we hope to consider dedekind zeta function with character, so it is natural to consider the character

$$\chi_{\mathfrak{m}} : (\mathcal{O}_k/\mathfrak{m})^* \rightarrow \mathbb{C}^*$$

where $\mathfrak{m}$ is an ideal in $\mathcal{O}_k$. Notice that $\mathcal{O}_k/\mathfrak{m}$ is finite, so the character is finite onto $\mathbb{S}^1$ exactly. However, ring of integers $\mathcal{O}_k$ is not always a UFD, so it is natural to consider the character on ideals.

Firstly set some notation here:

- $J(\mathfrak{m}) = \{\mathfrak{a} \in J_k \mid v_{\mathfrak{p}}(\mathfrak{a}) = 0, \forall \mathfrak{p} \mid \mathfrak{m}\}$ fractional ideals coprime with $\mathfrak{m}$
- $P(\mathfrak{m}) = J(\mathfrak{m}) \cap P_k$ principal ideals coprime with $\mathfrak{m}$
- $\iota : k^* \rightarrow P_k, x \mapsto (x)$ is a surjective group homomorphism with kernel $\mathcal{O}_k^*$ the group of units.
- $k(\mathfrak{m}) = \iota^{-1}(P(\mathfrak{m}))$ the pre-image, it is a subgroup of k^*

The first things is to translate the additive group structure to multiplicative group structure, from elements to ideals

To determine which elements are inversible in $\mathcal{O}_k/\mathfrak{m}$, we can write

$$\begin{aligned} \bar{a} \in (\mathcal{O}_k/\mathfrak{m})^* &\iff \exists b \in \mathcal{O}_k, ab = 1 \pmod{\mathfrak{m}} \\ &\implies 1 = ab + (ab - 1) \in (a) + \mathfrak{m} \\ &\implies (a) + \mathfrak{m} = \mathcal{O}_k \end{aligned}$$

conversely, if $(a) + \mathfrak{m} = \mathcal{O}_k$, then there exists $x \in \mathfrak{m}$ and $b \in \mathcal{O}_k$ such that $1 = ab + x$, so $ab - 1 = x = 0 \pmod{\mathfrak{m}}$, hence we have

$$\mathcal{O}_k(\mathfrak{m}) = \{a \in \mathcal{O}_k \mid (a) + \mathfrak{m} = \mathcal{O}_k\} = k(\mathfrak{m}) \cap \mathcal{O}_k$$

where $\mathcal{O}_k(\mathfrak{m})$ is exactly the set of elements which are invertible in $\mathcal{O}_k/\mathfrak{m}$, so it restricts us to consider characters on $k(\mathfrak{m})$.

Then we rewrite the residue class, for any $a, b \in k(\mathfrak{m})$

$$\begin{aligned} a = b \pmod{\mathfrak{m}} &\iff a - b \in \mathfrak{m} \\ &\iff \mathfrak{m} \mid (a - b) \\ &\iff v_{\mathfrak{p}}(a - b) \geq v_{\mathfrak{p}}(\mathfrak{m}), \forall \mathfrak{p} \mid \mathfrak{m} \\ &\iff a - b \in S \end{aligned}$$

where $S = \{x \in k(\mathfrak{m}) \mid v_{\mathfrak{p}}(x) \geq v_{\mathfrak{p}}(\mathfrak{m}), \forall \mathfrak{p} \mid \mathfrak{m}\}$ as the set of elements $x = 0 \pmod{\mathfrak{m}}$. We can investigate the structure of S to find that

- $a, b \in S$ then $ab \in S$: $v_{\mathfrak{p}}(ab) = v_{\mathfrak{p}}(a) + v_{\mathfrak{p}}(b) \geq 2v_{\mathfrak{p}}(\mathfrak{m}) \geq v_{\mathfrak{p}}(\mathfrak{m})$.
- S is not a subgroup since for any $a \in S$, $v_{\mathfrak{p}}(a^{-1}) = -v_{\mathfrak{p}}(a) \leq 0$.
- $aS = S$ for any $a \in k(\mathfrak{m})$: for any $s \in S$, we have

$$v_{\mathfrak{p}}(as) = v_{\mathfrak{p}}(a) + v_{\mathfrak{p}}(s) \geq 0 + v_{\mathfrak{p}}(\mathfrak{m})$$

where $\mathfrak{p}|\mathfrak{m}$, so it is enough to get $as \in S$. Conversely, $a^{-1} \in k(\mathfrak{m})$ since $k(\mathfrak{m})$ is a group, hence for any $s \in S$,

$$v_{\mathfrak{p}}(s) = v_{\mathfrak{p}}(a^{-1}) + v_{\mathfrak{p}}(as) = v_{\mathfrak{p}}(as)$$

which implies $as \in S$.

Hence S is not a good choice to rewrite the residue class, but by the final point above, we can keep going on

$$\begin{aligned} a = b \pmod{\mathfrak{m}} &\iff a \in b + S = b + bS = b(1 + S) \\ &\iff a/b \in 1 + S \end{aligned}$$

so here the relation given by addition turns to a relation given by multiplication, and hence we can define

$$\begin{aligned} k_{\mathfrak{m}} = 1 + S &= \{x \in k(\mathfrak{m}) \mid v_{\mathfrak{p}}(x - 1) \geq v_{\mathfrak{p}}(\mathfrak{m}), \forall \mathfrak{p}(\mathfrak{m})\} \\ &= \{x \in k(\mathfrak{m}) \mid x = 1 \pmod{\mathfrak{m}}\} \end{aligned}$$

and then we can conclude the following things:

Lemma 20. $k_{\mathfrak{m}}$ is a subgroup of $k(\mathfrak{m})$, and there exists an isomorphism

$$(\mathcal{O}_k/\mathfrak{m})^* \rightarrow k(\mathfrak{m})/k_{\mathfrak{m}}, \quad x + \mathfrak{m} \mapsto x \cdot k_{\mathfrak{m}}$$

hence any character $\chi_{\mathfrak{m}}$ on $(\mathcal{O}_k/\mathfrak{m})^*$ can be lifted to a character f on $k(\mathfrak{m})$ with $f|_{k_{\mathfrak{m}}} = 1$; Conversely, any such character can be pushed down to a character on $(\mathcal{O}_k/\mathfrak{m})^*$.

Now we consider the principal ideals by natural map:

$$\iota : k(\mathfrak{m}) \rightarrow P(\mathfrak{m}), \quad x \mapsto (x) = x\mathcal{O}_k$$

then the kernel is $\mathcal{O}_k^* \cap k(\mathfrak{m}) = \mathcal{O}_k^*$ such that we can consider the following diagram

$$\begin{array}{ccc} k(\mathfrak{m}) & \xrightarrow{\iota} & P(\mathfrak{m}) \\ \downarrow & & \downarrow \pi \\ k(\mathfrak{m})/\mathcal{O}_k^* \cdot k_{\mathfrak{m}} & \xrightarrow{\sim} & P(\mathfrak{m})/P_{\mathfrak{m}} \end{array}$$

here the kernel is calculated as following

$$\begin{aligned} \ker(\pi \circ \iota) &= \{x \in k(\mathfrak{m}) \mid (x) \in P_{\mathfrak{m}}\} \\ &= \{x \in k(\mathfrak{m}) \mid \exists u \in \mathcal{O}_k^*, ux \in k_{\mathfrak{m}}\} \\ &= \mathcal{O}_k^* \cdot k_{\mathfrak{m}} \end{aligned}$$

Then we consider the infinite places, it is necessary to handle unit here.

We set the minkowski's space $k_{\infty}^* = (\mathbb{R}^*)^r \times (\mathbb{C}^*)^s$, and then

$$k^* \xhookrightarrow{i} k_{\infty}^* \xrightarrow{\chi_{\infty}} \mathbb{C}^*$$

we consider the continuous character here, its form can be determined as following

Lemma 21. Along the notation

1. Any continous character $\mathbb{R}^* \rightarrow \mathbb{C}^*$ is of the form

$$\chi(t) = \chi(\text{sign}(t)) \cdot |t|^\lambda$$

with $\lambda \in \mathbb{C}$ and $\text{sign}(t) = \frac{t}{|t|} = \pm 1$.

2. Any continous character $\mathbb{C}^* \rightarrow \mathbb{C}^*$ is of the form

$$\chi(z) = \left(\frac{z}{|z|}\right)^n \cdot |z|^\lambda$$

with $\lambda \in \mathbb{C}$ and $n \in \mathbb{Z}$.

3. Any continous character $\chi_\infty : k_\infty^* \rightarrow \mathbb{C}^*$ is of the form

$$\chi_\infty(x) = w(x) \cdot \prod_v |x_v|^{it_v}$$

where $t_v \in \mathbb{R}$ while w is a finite character on the group of units $U_\infty = \{x \in k_\infty^* \mid |x_v| = 1, \forall v\}$.

Proof.

□

The reason why we consider the character on infinite places is as following: for a character $\chi_{\mathfrak{m}} : (\mathcal{O}_k/\mathfrak{m})^* \rightarrow \mathbb{S}^1$, we consider its induced character $f : k(\mathfrak{m}) \rightarrow \mathbb{S}^1$ with $f|_{k_{\mathfrak{m}}} = 1$. It motivates us to define as following

$$\chi : P(\mathfrak{m}) \rightarrow \mathbb{S}^1, \quad (a) \mapsto f(a)$$

however, here are some problems:

- Any $\mathfrak{a} \in J(\mathfrak{m})$, then $\mathfrak{a}^h = (a) \in P(\mathfrak{m})$ with h here the class number, so we can try to extend χ on $P(\mathfrak{m})$ by writing

$$\chi(\mathfrak{a})^h = \chi((a)) = f(a)$$

but the extension is not unique since we can choose any h -th root of $f(a)$ to be $\chi(\mathfrak{a})$.

- Above map we defined is not well-defined: if $(a) = (b) = (ua)$ for some unit u , then we have

$$\chi((a)) = \chi((ua)) \iff f(u) = 1$$

but f can be non-trivial on units, so problems comes out.

- To solve the problem, we try to invite the character on infinite places satisfying

$$\chi((a)) = f(a) \cdot \chi_\infty(i(a)) \quad \forall a \in k(\mathfrak{m})$$

in this case if $u \in \mathcal{O}_k^*$

$$1 = f(u) \cdot \chi_\infty(i(u))$$

and if $a \in k_{\mathfrak{m}}$ such that $f(a) = 1$

$$\chi((a)) = \chi_\infty(i(a))$$

this gives the restrictions on χ , or a commute digram as following

$$\begin{array}{ccc} k_{\mathfrak{m}} & \xrightarrow{\iota} & P(\mathfrak{m}) \\ \downarrow & & \downarrow \chi \\ k_{\infty}^* & \xrightarrow{\chi_{\infty}} & \mathbb{S}^1 \end{array}$$

Theorem 22 (Weak apporximation theorem).

Let $v_1, \dots, v_m$ be pairwise inequivalent **absolute values** of the field k , and $a_1, \dots, a_m \in k$ be any elements, then for any $\epsilon > 0$, there exists an element $x \in k$ such that

$$v_i(x - a_i) < \epsilon \quad \text{for } i = 1, \dots, m$$

Proof. notice that for any i , there exists an element y_i such that

$$v_j(y_i) \begin{cases} > 1 & j = i \\ < 1 & j \neq i \end{cases}$$

hence we can consider the sequence $x_n^{(i)} = \frac{y_i^n}{1+y_i^n}$, then

$$v_j(x_n^{(i)}) \xrightarrow{n \rightarrow \infty} \begin{cases} 1 & j = i \\ 0 & j \neq i \end{cases}$$

Hence we can consider $x_n = \sum_{k=1}^m a_k x_n^{(k)}$, then for any i we have

$$\begin{aligned} v_i(x_n - a_i) &= v_i(a_i)v_i(1 - x_n^{(i)}) + \sum_{k \neq i} v_i(a_k)v_i(x_n^{(k)}) \\ &\xrightarrow{n \rightarrow \infty} v_i(a_i) \cdot 0 + 0 = 0 \end{aligned}$$

so we can choose a suitable n for any $\epsilon > 0$ such that $v_i(x_n - a_i) < \epsilon$ for any i . □

Corollary 23. $k_{\mathfrak{m}}$ is dense in k_{∞}^* under the embedding

$$i : k \hookrightarrow k_{\infty}, \quad x \mapsto (x_{\sigma})_{\sigma \in \Sigma_o}$$

where Σ_o is the set of ditinct embeddings up to conjuation, and $x_{\sigma} = \sigma(x)$ is the image of x under embedding σ .

Proof. Let $S = S_{\mathfrak{m}} \cup S_{\infty}$ be a set of absolute values on k , where $S_{\mathfrak{m}}$ is the set of non-archmedean absolute values given by prime ideals diving $\mathfrak{m}$, while S_{∞} is the set of archemedean absolute values given by $v(x) = |\sigma(x)|$ for any $\sigma \in \Sigma_o$. Hence we choose $a_v = 1$ if $v \in S_{\mathfrak{m}}$, and $a_v \in k^*$ arbitrary if $v \in S_{\infty}$, apply the weak approximation theorem, we can get an element $x \in k$ such that $v(x - 1) < 1$ for any $v \in S_{\mathfrak{m}}$, and $v(x - a_v) < \epsilon$ for any $v \in S_{\infty}$. The first condition implies that $x \in k_{\mathfrak{m}}$, while the second condition implies that $i(x)$ is close to $(a_v)_v$ in k_{∞}^* , hence we can conclude that $i(k_{\mathfrak{m}}) \subset i(k^*)$ dense, and then we can conclude that $k_{\mathfrak{m}}$ is dense in k_{∞}^* since $i(k)$ is dense in k_{∞} . □

With enough preparation, we can define the character defined by Hecke (1920) in his original paper as following

Definition 24. Let $\mathfrak{m}$ be an ideal of $\mathcal{O}_k$, a **Größencharakter modulo $\mathfrak{m}$** is a character $\chi : J(\mathfrak{m}) \rightarrow \mathbb{S}^1$ for which there exists a pair of (continuous) characters

$$\chi_{\mathfrak{m}} : (\mathcal{O}_k/\mathfrak{m})^* \rightarrow \mathbb{S}^1, \quad \chi_{\infty} : k_{\infty}^* \rightarrow \mathbb{S}^1$$

such that

$$\chi((a)) = \chi_{\mathfrak{m}}(\bar{a}) \cdot \chi_{\infty}(i(a))$$

for any $a \in k(\mathfrak{m}) \cap \mathcal{O}_k$.

Remark 25. Here are several explanations:

- $\chi_{\mathfrak{m}}$ can be identified as a character $f : k(\mathfrak{m}) \rightarrow \mathbb{S}^1$ as before, what we do here is to translate the relation before back to $\chi_{\mathfrak{m}}$.
- Review that $\chi((a)) = \chi_{\infty}(i(a))$ for any $a \in k_{\mathfrak{m}}$, the condition of continuity uniquely determines χ_{∞} since embedding $k_{\mathfrak{m}}$ is dense in k_{∞}^* .
- Conversely, we can determine χ by the values on $P(\mathfrak{m})$, but it is not unique.

4.2 Ideal number system

In Hecke's paper, he uses the notation of ideal numbers to make sum on ideals can be written as sum of elements, that is what we done above in Dedekind zeta function to make the functional equation.

Definition 26. For a number field k , the **ideal number system** of k is a multiplicative abelian group $\hat{k}^*$ containing k^* such that the following diagram commutes and the rows are exact sequences

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \mathcal{O}_k^* & \hookrightarrow & k^* & \longrightarrow & P_k & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \mathcal{O}_k^* & \longrightarrow & \hat{k}^* & \longrightarrow & J_k & \longrightarrow & 1 \end{array}$$

The construction of the system can be done as following:

By the theory of Minkowski, there exists a prime ideal $\mathfrak{p}$ in each class in class group J_k/P_k , so for any class we choose exactly one prime ideal $\mathfrak{p}$ as the representative, then there exists a divisor d of class number h_k such that $\mathfrak{p}^d = (a)$ for some $a \in \mathcal{O}_k$, hence we define $\hat{p} \in \hat{k}^*$ as an ideal number of $\mathfrak{p}$ which satisfies $(\hat{p}) = \mathfrak{p}$, immediately $\hat{p}^d = ua$ for some unit u .

Then we can define $\hat{k}^*$ as a group generated by k^* and finite ideal numbers $\hat{p}$ of all classes (notice that in principal class, $\hat{p}$ is just in k^*). Now if we take any fractional ideals $\mathfrak{b}$, it will be in some class $[\mathfrak{p}]$, then there exists $x \in k^*$ such that

$$\mathfrak{b} = (x)\mathfrak{p} = (x)(\hat{p})$$

anyway some problem here ...

some properties of the system:

- By two exact sequences, we have $k^*/\mathcal{O}_k^* \cong P_k$ and $\hat{k}^*/\mathcal{O}_k^* \cong J_k$, hence by isomorphism theorem, we can conclude an isomorphism for class group

$$\hat{k}^*/k^* \cong J_k/P_k$$

- The norm of elements can be extends to ideal numbers as following:

$$\mathcal{N}(\hat{a}) = N((\hat{a}))$$

i.e. make use of the absolute norm of ideals, clearly if we restrict the norm on k^* , it will be the algebraic norm $N_{k/\mathbb{Q}}$.

- Let $\hat{a} \in k^*$, we call it an ideal integer if $(\hat{a})$ is an integral ideal, and we define $\hat{\mathcal{O}}$ as the set of all ideal integers, it is a semi-group (may not have a) under multiplication

4.3 Primitive character and Gauss sum

5 Tate's Thesis

5.1 Adele Ring