


# 2WBB0 Calculus for BCS

Dr. Björn Baumeier



Technische Universiteit  
Eindhoven  
University of Technology

# Why do we (=you) need Calculus?



# Why do we (=you) need Calculus?

December 21, 1968? Shorter clip


# Why do we (=you) need Calculus?

December 21, 1968? Shorter clip




# Why do we (=you) need Calculus?

December 21, 1968? Shorter clip



## Why do we (=you) need Calculus?

December 21, 1968?




# Why do we (=you) need Calculus?

December 21, 1968?



# Why do we (=you) need Calculus?

November 16, 2022




# Why do we (=you) need Calculus?

April 26, 2023

## THE GLITCH THAT BROUGHT DOWN JAPAN'S LUNAR LANDER

by: Matthew Carlson

37 Comments



June 3, 2023



# Why do we (=you) need Calculus?

April 26, 2023

When a computer crashes, it usually doesn't leave debris. But when a computer happens to be descending towards the lunar surface and glitches out, that's a very different story. Turns out that's what happened on April 26th, as the Japanese Hakuto-R Lunar lander made its mark on the Moon...by crashing into it. [Scott Manley] dove in to try and understand the software bug that caused an otherwise flawless mission to go splat.

The lander began the descent sequence as expected at 100 km above the surface. However, as it descended, the altitude sensor reported the altitude as much lower than it was. It thought it was at zero altitude once it reached about 5 km above the surface. Confused by the fact it hadn't yet detected physical contact with the surface, the craft continued to slowly descend until it ran out of fuel and plunged to the surface.

Ultimately it all came down to sensor fusion. The lander merges several noisy sensors, such as accelerometers, gyroscopes, and radar, into one cohesive source of truth. The craft passed over a particularly large cliff that caused the radar altimeter to suddenly spike up 3 km. Like good filtering software, the craft reasons that the sensor must be getting spurious data and filters it out. It was now just estimating its altitude by looking at its acceleration. As anyone who has tried to track an object through space using just gyros and accelerometers alone can attest, errors accumulate, and suddenly you're not where you think you are.

# Why do we (=you) need Calculus?

There are 20 people in the room. They all need to say good bye to each other, and shake hands. How many handshakes?

# Why do we (=you) need Calculus?

There are 20 people in the room. They all need to say good bye to each other, and shake hands. How many handshakes?

## Without Calculus

```
Readln('how many people?', X);
cnt := 0;
for i := 1 to X do
    for j := 1 to X do
        if j>i then inc(cnt);
Writeln(cnt);
```

# Why do we (=you) need Calculus?

There are 20 people in the room. They all need to say good bye to each other, and shake hands. How many handshakes?

## Without Calculus

```
Readln('how many people?', X);
cnt := 0;
for i := 1 to X do
    for j := 1 to X do
        if j>i then inc(cnt);
Writeln(cnt);
```

## With Calculus

```
Readln('how many people?', X);
Writeln(X*(X-1)/2);
```

# Why do we (=you) need Calculus?

There are 20 people in the room. They all need to say good bye to each other, and shake hands. How many handshakes?

## Without Calculus

```
Readln('how many people?', X);
cnt := 0;
for i := 1 to X do
    for j := 1 to X do
        if j > i then inc(cnt);
Writeln(cnt);
```

## With Calculus

```
Readln('how many people?', X);
Writeln(X*(X-1)/2);
```

- ▶ For 20 people, nobody can tell the difference.

# Why do we (=you) need Calculus?

There are 20 people in the room. They all need to say good bye to each other, and shake hands. How many handshakes?

## Without Calculus

```
Readln('how many people?', X);
cnt := 0;
for i := 1 to X do
    for j := 1 to X do
        if j>i then inc(cnt);
Writeln(cnt);
```

## With Calculus

```
Readln('how many people?', X);
Writeln(X*(X-1)/2);
```

- ▶ For 20 people, nobody can tell the difference.
- ▶ For one million people in the room, you get to do  $10^{14}$  loops. You need a 64 bit machine for storing the counter. At 3 GHz, that takes 166 seconds (way more in real life). With Calculus: instant!

# Calculus in Computer Science

## For video game programming.

Calculus, when taught alongside Physics, gives you the background necessary to understand how objects can interact and move based on forces. Even if you don't ever write your own game engine, you'll be a better game programmer if you have a mastery of calculus.

## For scientific applications.

A lot of computer scientists work alongside scientists like biologists, physicists, and chemists. CS people help them solve their data-analysis problems using large clusters of computers, and anytime you have to understand how equations work, it is hard to escape calculus.

## For machine learning.

Basically machine learning is mathematics meets computer science meets data. This is one of the hottest topics in CS right now, but you have to know your mathematics to succeed with ML.

## In research.

Researchers such as Professors and PhD students use mathematics when writing papers. If you want to one day pursue an advanced degree, you'll have to decipher these papers and then write some of your own. I would not recommend starting out a career in research without at least a basic command over calculus.


# Analyze a Computer Algorithm

Imagine you are a programmer of an automatic, autonomous guidance system (think of cars, rockets). You have an algorithm that takes data collected from a number of input sources (sensors) and processes it to adjust the control of the vehicle. It takes 50ms to obtain the adjustment after the acquisition of data (reaction time).

Now, the engineers plan to double the number of sensors to improve the guidance. The maximal acceptable reaction time for the vehicle is 150ms.

**Can your algorithm deal with double the amount of data within the 150ms?**

# Different scaling of (parts of complex) algorithms



## Example: Bubble Sort

Sorting an unordered list of five numbers by swapping adjacent elements that are out of place:

17	12	8	3	7
12	17	8	3	7
12	8	17	3	7
12	8	3	17	7
12	8	3	7	17

How many comparisons?  $n - 1$

## Example: Bubble Sort

Now, do it again

12	8	3	7	17
8	12	3	7	17
8	3	12	7	17
8	3	7	12	17

How many comparisons?  $n - 2$

And again:

8	3	7	12	17
3	8	7	12	17
3	7	8	12	17

How many comparisons?  $n - 3$

## Example: Bubble Sort

Finally:

3   7   8   12   17

How many comparisons?  $n - 4$

So all in all:

$$(n - 1) + (n - 2) + (n - 3) + \dots + 1 = \frac{n(n - 1)}{2}$$

# Ranking Different Algorithms

- ▶ How to compare different algorithms?
- ▶ Two algorithms

Algorithm 1  $\rightarrow f(n) = n$

Algorithm 2  $\rightarrow g(n) = n^2$

- ▶ Which one is "faster"?
- ▶ Algorithm 1, because it grows at a slower rate with  $n$

# Ranking Different Algorithms

- ▶ How to formulate this more generally?  
→ Limits

- ▶
$$\lim_{n \rightarrow \infty} \frac{f}{g} = 0 \quad g \text{ grows faster}$$

$$\lim_{n \rightarrow \infty} \frac{f}{g} = c \quad g \text{ grows same as } f$$

$$\lim_{n \rightarrow \infty} \frac{f}{g} = \infty \quad g \text{ grows slower}$$

# Limits not always straightforward

- ▶  $f(n) = n \log(n)$

- ▶  $g(n) = n^{3/2}$

- ▶

$$\lim_{n \rightarrow \infty} \frac{n^{3/2}}{n \log(n)} = ?$$


- ▶ To evaluate we will need:

- ▶ derivatives
- ▶ Taylor series/polynomials
- ▶ Exponential functions and Logarithms
- ▶ rational functions
- ▶ ...

# Overview

- ▶ W1: Numbers, Functions, (In-)equalities, Polynomials, Rational functions
- ▶ W2: Trigonometric functions/Vectors in 2 and 3 dimensions
- ▶ W3: Limits, continuity, differentiation I
- ▶ W4: Differentiation II and Inverse functions
- ▶ W5: Exponential function and logarithm/Taylor series/Limits II
- ▶ W6: Integration/Integration techniques I
- ▶ W7: Integration techniques II/1st order differential equations
- ▶ W8: Exam and W1 – W7 summary (no new material)

Shall we begin?



# Topics in Week 1

- ▶ Real numbers and the real line
- ▶ Graphs of quadratic equations
- ▶ Polynomials and rational functions
- ▶ Powers and roots
- ▶ Cartesian coordinates in the plane
- ▶ More functions and their graphs
- ▶ Combining functions to make new functions

# Real numbers

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$$

↑  
is a subset of

- $\mathbb{N}$  natural numbers:  $0, 1, 2, \dots$  (sometimes without 0)
- $\mathbb{Z}$  integer numbers:  $\dots, -2, -1, 0, 1, 2, \dots$
- $\mathbb{Q}$  rational numbers = fractions:  $\frac{p}{q}$  with  $p, q \in \mathbb{Z}$ ,  $q \neq 0$
- $\mathbb{R}$  real numbers:  $\mathbb{Q}$  and  $\pi$ ,  $\sqrt{2}$ , etc
- $\mathbb{C}$  complex numbers:  $2 + 3i$ , with  $i \cdot i = i^2 = -1$  (not in curriculum)

# Decimal Expansion

For rational numbers, the “decimal expansion” has a repeating pattern, for irrational numbers, this is not the case:

$$\frac{1}{2} = 0.50000|0|\dots$$

$$\frac{1}{3} = 0.33333|3|\dots$$

$$\frac{1}{11} =$$

# Decimal Expansion

For rational numbers, the “decimal expansion” has a repeating pattern, for irrational numbers, this is not the case:

$$\frac{1}{2} = 0.50000|0|\dots$$

$$\frac{1}{3} = 0.33333|3|\dots$$

$$\frac{1}{11} = 0.0909|09|\dots$$

$$\frac{1}{7} =$$

# Decimal Expansion

For rational numbers, the “decimal expansion” has a repeating pattern, for irrational numbers, this is not the case:

$$\frac{1}{2} = 0.50000|0|\dots$$

$$\frac{1}{3} = 0.33333|3|\dots$$

$$\frac{1}{11} = 0.0909|09|\dots$$

$$\frac{1}{7} = 0.142857|142857|\dots$$

$$\pi =$$

# Decimal Expansion

For rational numbers, the “decimal expansion” has a repeating pattern, for irrational numbers, this is not the case:

$$\frac{1}{2} = 0.50000|0|\dots$$

$$\frac{1}{3} = 0.33333|3|\dots$$

$$\frac{1}{11} = 0.0909|09|\dots$$

$$\frac{1}{7} = 0.142857|142857|\dots$$

$$\pi = 3.141592653589793\dots$$

# Intervals

Notation for intervals:

- ▶  $[ , ]$ : end points are part of the interval
- ▶  $( , )$ : end points are NOT part of the interval

interval	alternative notation	type	type
$(0, 1)$	$0 < x < 1$	open	finite
$[0, 1)$	$0 \leq x < 1$	half open	finite
$(0, 1]$	$0 < x \leq 1$	half open	finite
$[0, 1]$	$0 \leq x \leq 1$	closed	finite
$[1, \infty)$	$1 \leq x < \infty$		infinite
$(-\infty, 3)$	$-\infty < x < 3$		infinite
$(-\infty, \infty)$	$-\infty < x < \infty$		infinite

# Symbols

$\cup$	union
$\cap$	intersection
$\in$	element of
$\vee$	or
$\wedge$	and
$(A) \implies (B)$	from (A) follows (B)
$(A) \Leftrightarrow (B)$	from (A) follows (B) and from (B) follows (A) in other words: they are equivalent (A) "if and only if" (B) (iff)

For two sets  $A$  and  $B$ :

- ▶  $A - B$
- ▶  $A \setminus B$

represents all  $x \in A$  for which  $x \notin B$ .

# Polynomials

A polynomial is a “many term” construct

The degree (order) of a polynomial is the highest occurring power

Ex:  $p(x) = x^3 - 2x^2 + 1$  is of degree 3

Ex:  $p(x) = 0 \cdot x^3 - 2x^2 + 1$  is of degree 2

Ex:  $p(x) = (x + 1)^2 - (x - 1)^2$  is of degree 1

Ex:  $p(x) = 3 = 3 \cdot x^0$  is of degree 0

Computers can only do

- ▶ additions/subtractions
- ▶ multiplications

All other operations, such as division or calculation of a root have been performed using only these basis operations – and then cost 3 to 5 times more calculational time

# Frequently appearing polynomials

Degree	Name	Example
0	constant	$p(x) = 2$
1	linear	$p(x) = 2x + 3$
2	quadratic	$p(x) = 5x^2 + 4x + 5$
3	cubic	$p(x) = 2x^3 + 1$

Equations with polynomials of degree 1 always have a solution:

Ex:  $4x - 3 = 0$

# Frequently appearing polynomials

Degree	Name	Example
0	constant	$p(x) = 2$
1	linear	$p(x) = 2x + 3$
2	quadratic	$p(x) = 5x^2 + 4x + 5$
3	cubic	$p(x) = 2x^3 + 1$

Equations with polynomials of degree 1 always have a solution:

Ex:  $4x - 3 = 0 \implies 4x = 3 \implies x = \frac{3}{4}$

Theorem:

Equations of polynomials of degree  $n$  have a maximum of  $n$  real solutions

# Equations with a polynomial of degree 2

Quadratic functions are more than algebraic curiosities – they are widely used in science, business, and engineering:

- ▶ trajectories of water jets in a fountain or of a bouncing ball
- ▶ parabolic reflectors that form the base of satellite dishes and car headlights
- ▶ forecast business profit and loss
- ▶ data interpolation

Equations of quadratic polynomials can always be written into the general form:

$$ax^2 + bx + c = 0$$

# Equations with a polynomial of degree 2

Equations of quadratic polynomials can always be written into the general form:

$$ax^2 + bx + c = 0$$

Our task is to find the solutions (or zeros) of this equation.

3 different ways to find the solution:

- ▶ completing the square
- ▶ abc-formula (quadratic formula)
- ▶ “Sum-product”-formula

Use the way that you like best **for this purpose** but learn the concept of **completing the square!** It will be useful later.

# Idea behind Completing the Square

Ex: :  $x^2 = 3$

# Idea behind Completing the Square

Ex: :  $x^2 = 3 \Leftrightarrow x = \pm\sqrt{3}$

# Idea behind Completing the Square

Ex: :  $x^2 = 3 \Leftrightarrow x = \pm\sqrt{3}$

This is easy (and also an equation with a quadratic polynomial)!

So, in general, if we have an equation

$$(\text{something})^2 = \text{something else}$$

we can simply take the square root on both sides.

# Idea behind Completing the Square

Ex:  $x^2 = 3 \Leftrightarrow x = \pm\sqrt{3}$

This is easy (and also an equation with a quadratic polynomial)!

So, in general, if we have an equation

$$(\text{something})^2 = \text{something else}$$

we can simply take the square root on both sides.

Ex:  $(x - 1)^2 = 9$

## Idea behind Completing the Square

Ex:  $x^2 = 3 \Leftrightarrow x = \pm\sqrt{3}$

This is easy (and also an equation with a quadratic polynomial)!

So, in general, if we have an equation

$$(\text{something})^2 = \text{something else}$$

we can simply take the square root on both sides.

Ex:  $(x - 1)^2 = 9 \Leftrightarrow x - 1 = \pm 3 \Leftrightarrow x = 1 \pm 3$

Rewrite a quadratic equation to a form with a complete square on the left-hand side

$$(x \pm R)^2 = T$$

and take the square root on both sides, if possible.

# The ONE basic trick for completing the square

Remember the binomial formulas:

$$\begin{aligned}(n+m)^2 &= n^2 + 2nm + m^2 \\(n-m)^2 &= n^2 - 2nm + m^2\end{aligned}$$

and that one can also use them from right to left!

# The ONE basic trick for completing the square

Remember the binomial formulas:

$$\begin{aligned}(n + m)^2 &= n^2 + 2nm + m^2 \\(n - m)^2 &= n^2 - 2nm + m^2\end{aligned}$$

and that one can also use them from right to left!

Ex:  $x^2 - 4x + 4 = (x - 2)^2$

# The ONE basic trick for completing the square

Remember the binomial formulas:

$$\begin{aligned}(n+m)^2 &= n^2 + 2nm + m^2 \\(n-m)^2 &= n^2 - 2nm + m^2\end{aligned}$$

and that one can also use them from right to left!

Ex:  $x^2 - 4x + 4 = (x - 2)^2$

Ex:  $4x^2 + 12x + 9 = (\underbrace{2x}_{=n})^2 + \underbrace{2}_{\text{binom. form.}} \cdot \underbrace{2x}_{=n} \cdot \underbrace{3}_{=m} + (\underbrace{3}_{=m})^2 = (2x + 3)^2$

Those examples are complete squares. They can be rewritten directly into one of the two binomial formulas.

# The ONE basic trick for completing the square

Remember the binomial formulas:

$$\begin{aligned}(n+m)^2 &= n^2 + 2nm + m^2 \\(n-m)^2 &= n^2 - 2nm + m^2\end{aligned}$$

and that one can also use them from right to left!

$$\text{Ex: } x^2 - 4x + 4 = (x - 2)^2$$

$$\text{Ex: } 4x^2 + 12x + 9 = (\underbrace{2x}_{=n})^2 + \underbrace{2}_{\text{binom. form.}} \cdot \underbrace{2x}_{=n} \cdot \underbrace{3}_{=m} + (\underbrace{3}_{=m})^2 = (2x + 3)^2$$

Those examples are complete squares. They can be rewritten directly into one of the two binomial formulas.

But what about  $x^2 - 4x + 3$ ? We are missing  $+1$  for a complete square!

# The ONE basic trick for completing the square

Remember the binomial formulas:

$$\begin{aligned}(n+m)^2 &= n^2 + 2nm + m^2 \\(n-m)^2 &= n^2 - 2nm + m^2\end{aligned}$$

and that one can also use them from right to left!

$$\text{Ex: } x^2 - 4x + 4 = (x - 2)^2$$

$$\text{Ex: } 4x^2 + 12x + 9 = (\underbrace{2x}_{=n})^2 + \underbrace{2}_{\text{binom. form.}} \cdot \underbrace{2x}_{=n} \cdot \underbrace{3}_{=m} + (\underbrace{3}_{=m})^2 = (2x + 3)^2$$

Those examples are complete squares. They can be rewritten directly into one of the two binomial formulas.

But what about  $x^2 - 4x + 3$ ? We are missing +1 for a complete square!

$$x^2 - 4x + 3 = x^2 - 4x + 3 + 0 = x^2 - 4x + 3 + 1 - 1 = x^2 - 4x + 4 - 1 = (x - 2)^2 - 1$$

# Completing the square to solve quadratic equations

Solve:

$$ax^2 + bx + c = 0$$

- 1 Identify  $n$  and  $m$  in the expression.

- ▶  $n$  is the square root of the term with  $x^2$   
 $ax^2 \rightarrow n = \sqrt{ax}$
- ▶  $m$  is the linear term in  $x$ , divided by  $2n$   
 $m = bx/(2 \cdot \sqrt{ax}) = b/(2\sqrt{a})$

- 2 Use binomial formula

$$(n + m)^2 = (\sqrt{ax} + \frac{b}{2\sqrt{a}})^2 = ax^2 + bx + \frac{b^2}{4a}$$

- 3 what is missing/too much

$$ax^2 + bx + c = ax^2 + bx + \frac{b^2}{4a} - \frac{b^2}{4a} + c = (\sqrt{ax} + \frac{b}{2\sqrt{a}})^2 - \frac{b^2}{4a} + c = 0$$

- 4 isolate the square, take roots, simplify

$$(\sqrt{ax} + \frac{b}{2\sqrt{a}})^2 = \frac{b^2}{4a} - c \Leftrightarrow \sqrt{ax} + \frac{b}{2\sqrt{a}} = \pm \frac{\sqrt{b^2 - 4ac}}{2\sqrt{a}}$$

$$\Leftrightarrow x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$ax^2 + bx + c = 0$  via abc-formula (I)

2nd degree polynomial  $ax^2 + bx + c = 0$  with  $a \neq 0$ :

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

How many solutions are possible?

Discriminant:  $D = b^2 - 4ac$

$D > 0$



$ax^2 + bx + c = 0$  via abc-formula (I)

2nd degree polynomial  $ax^2 + bx + c = 0$  with  $a \neq 0$ :

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

How many solutions are possible?

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$



$ax^2 + bx + c = 0$  via abc-formula (I)

2nd degree polynomial  $ax^2 + bx + c = 0$  with  $a \neq 0$ :

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

How many solutions are possible?

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$  1 double solution (2 solutions  $x_1 = x_2$ )

$D < 0$



$ax^2 + bx + c = 0$  via abc-formula (I)

2nd degree polynomial  $ax^2 + bx + c = 0$  with  $a \neq 0$ :

$$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

How many solutions are possible?

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$  1 double solution (2 solutions  $x_1 = x_2$ )

$D < 0$  no (real) solution

The zeros determine the factoring of the polynomial:

$$ax^2 + bx + c = (x - x_1)(x - x_2).$$

Pay attention to the minus sign!

## $ax^2 + bx + c = 0$ via abc-formula (II)

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$  1 double solution (2 solutions  $x_1 = x_2$ )

$D < 0$  no (real) solution

Ex:  $x^2 + 1 = 0$

## $ax^2 + bx + c = 0$ via abc-formula (II)

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$  1 double solution (2 solutions  $x_1 = x_2$ )

$D < 0$  no (real) solution

Ex:  $x^2 + 1 = 0 \implies x^2 = -1 \implies$  no solution

indeed:  $D = 0^2 - 4 \cdot 1 \cdot 1 = -4 < 0 \implies$  no solution

Ex:  $x^2 + 2x + 1 = 0$

## $ax^2 + bx + c = 0$ via abc-formula (II)

Discriminant:  $D = b^2 - 4ac$

$D > 0$  2 different solutions ( $x_1 \neq x_2$ )

$D = 0$  1 double solution (2 solutions  $x_1 = x_2$ )

$D < 0$  no (real) solution

Ex:  $x^2 + 1 = 0 \implies x^2 = -1 \implies$  no solution

indeed:  $D = 0^2 - 4 \cdot 1 \cdot 1 = -4 < 0 \implies$  no solution

Ex:  $x^2 + 2x + 1 = 0 : D = 2^2 - 4 \cdot 1 \cdot 1 = 0 \implies$  double solution

indeed:  $x^2 + 2x + 1 = 0 \implies (x + 1)^2 = 0$

$\implies x + 1 = 0$  or  $x + 1 = 0 \implies$  double solution  $x = -1$

$ax^2 + bx + c = 0$  via “Sum-product”

Set  $a = 1$ :

$$x^2 + \underbrace{(r+s)x}_{=b(\text{sum})} + \underbrace{rs}_{=c(\text{product})} = (x+r)(x+s)$$

Ex:  $x^2 + 5x + 6 =$

$ax^2 + bx + c = 0$  via “Sum-product”

Set  $a = 1$ :

$$x^2 + \underbrace{(r+s)x}_{=b(\text{sum})} + \underbrace{rs}_{=c(\text{product})} = (x+r)(x+s)$$

Ex:  $x^2 + 5x + 6 = (x+2)(x+3)$

Because  $b = 5 = 2 + 3$  and  $c = 6 = 2 \cdot 3$

Ex:  $x^2 - 9 = (x-3)(x+3)$

Attention!  $x^2 = 9 \not\Rightarrow x = 3$ , but :

$$\begin{aligned} x^2 = 9 &\implies x^2 - 9 = 0 \implies (x-3)(x+3) = 0 \implies \\ x - 3 = 0 \text{ or } x + 3 = 0 &\implies x = 3 \quad \text{or} \quad x = -3 \end{aligned}$$

## Examples: Quadratic Equations

- 1 Solve:  $x^2 - 3x + 2 = 0$
- 2 Factorize:  $x^2 = 2$
- 3 Solve:  $x^2 + 2 = 0$
- 4 Complete the square:  $x^2 + 3x + \frac{1}{4}$
- 5 Complete the square:  $4x^2 + 12x + 1$

# Equations with a 3rd degree polynomial

- 1 Guess first zero. Try out first  $a = 0, 1, -1, 2, -2, \dots$
- 2 Divide by factor  $x - a$
- 3 The rest must be 0 (if not, you made a mistake)
- 4 Find solutions of the remaining 2nd degree polynomial equation, if they exist

Ex:  $x^3 + 2x^2 - x - 2 = 0$

Try:

# Equations with a 3rd degree polynomial

- 1 Guess first zero. Try out first  $a = 0, 1, -1, 2, -2, \dots$
- 2 Divide by factor  $x - a$
- 3 The rest must be 0 (if not, you made a mistake)
- 4 Find solutions of the remaining 2nd degree polynomial equation, if they exist

Ex:  $x^3 + 2x^2 - x - 2 = 0$

Try:  $x = 1$  is a solution

Then is  $x^3 + 2x^2 - x - 2 = (x - 1) \cdot (\text{2nd degree polynomial})$

$$x^3 + 2x^2 - x - 2 = (x - 1) \cdot (\text{2nd degree polynomial})$$

To determine the 2nd order polynomial, divide out (see later):

$$\frac{x^3 + 2x^2 - x - 2}{x - 1} = x^2 + 3x + 2$$

And then  $x^2 + 3x + 2 =$

$$x^3 + 2x^2 - x - 2 = (x - 1) \cdot (\text{2nd degree polynomial})$$

To determine the 2nd order polynomial, divide out (see later):

$$\frac{x^3 + 2x^2 - x - 2}{x - 1} = x^2 + 3x + 2$$

And then  $x^2 + 3x + 2 = (x + 2)(x + 1)$  via abc-formula

Conclusion  $x^3 + 2x^2 - x - 2 = (x - 1)(x + 2)(x + 1)$

Thus solution of  $x^3 + 2x^2 - x - 2 = 0$ :

$$x^3 + 2x^2 - x - 2 = (x - 1) \cdot (\text{2nd degree polynomial})$$

To determine the 2nd order polynomial, divide out (see later):

$$\frac{x^3 + 2x^2 - x - 2}{x - 1} = x^2 + 3x + 2$$

And then  $x^2 + 3x + 2 = (x + 2)(x + 1)$  via abc-formula

Conclusion  $x^3 + 2x^2 - x - 2 = (x - 1)(x + 2)(x + 1)$

Thus solution of  $x^3 + 2x^2 - x - 2 = 0$ :  $x = 1$  or  $x = -1$  or  $x = -2$

# Equations with 3rd, 4th, 5th, ... -degree polynomials

Principle: guess, divide, guess, etc, until 2nd-degree remains

- 1 First, have a look if there is not a special easier case,  
as  $p(x) = 0$  with  $p(x) = x^3 - 8$  or  $p(x) = x^4 + 2x^2 + 1$
- 2 If not, guess a zero, say  $x = a$ ,  $a = 0, 1, -1, 2, -2, \dots$
- 3 Divide out by Long Division:  $p(x)/(x - a)$   
(here **must** have rest 0),  
this yields a new polynomial  $q(x)$
- 4 If the degree of  $q(x)$  is still larger than 2,  
repeat the above
- 5 If the degree of  $q(x)$  is equal to 2,  
try with sum-product technique or the abc-formula

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

4th degree but 2nd degree in  $x^2$ ! Rename  $p = x^2$

It holds that  $(x^2)^2 = x^2 \cdot x^2 = x^{2+2} = x^4$

$$p^2 - 50p + 49 = 0$$

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

4th degree but 2nd degree in  $x^2$ ! Rename  $p = x^2$

It holds that  $(x^2)^2 = x^2 \cdot x^2 = x^{2+2} = x^4$

$$\begin{aligned} p^2 - 50p + 49 &= 0 \implies (p - 49)(p - 1) = 0 \implies p = 49 \text{ or } p = 1 \\ \stackrel{p=x^2}{\implies} \implies x^2 &= 49 \text{ or } x^2 = 1 \implies x = \pm 7 \text{ or } x = \pm 1 \end{aligned}$$

Higher degree but only products of lower degrees

$$x^4(x - 2)^2 = 0$$

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

4th degree but 2nd degree in  $x^2$ ! Rename  $p = x^2$

It holds that  $(x^2)^2 = x^2 \cdot x^2 = x^{2+2} = x^4$

$$\begin{aligned} p^2 - 50p + 49 &= 0 \implies (p - 49)(p - 1) = 0 \implies p = 49 \text{ or } p = 1 \\ \stackrel{p=x^2}{\implies} \implies x^2 &= 49 \text{ or } x^2 = 1 \implies x = \pm 7 \text{ or } x = \pm 1 \end{aligned}$$

Higher degree but only products of lower degrees

$$x^4(x - 2)^2 = 0 \implies x^4 = 0 \text{ or } (x - 2)^2 = 0$$

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

4th degree but 2nd degree in  $x^2$ ! Rename  $p = x^2$

It holds that  $(x^2)^2 = x^2 \cdot x^2 = x^{2+2} = x^4$

$$\begin{aligned} p^2 - 50p + 49 &= 0 \implies (p - 49)(p - 1) = 0 \implies p = 49 \text{ or } p = 1 \\ \stackrel{p=x^2}{\implies} \implies x^2 &= 49 \text{ or } x^2 = 1 \implies x = \pm 7 \text{ or } x = \pm 1 \end{aligned}$$

Higher degree but only products of lower degrees

$$\begin{aligned} x^4(x - 2)^2 &= 0 \implies x^4 = 0 \text{ or } (x - 2)^2 = 0 \\ \implies x &= 0 \text{ or } x - 2 = 0 \end{aligned}$$

## Degree 4, 5, . . .: Special cases

Ex:  $x^4 - 50x^2 + 49 = 0$

4th degree but 2nd degree in  $x^2$ ! Rename  $p = x^2$

It holds that  $(x^2)^2 = x^2 \cdot x^2 = x^{2+2} = x^4$

$$\begin{aligned} p^2 - 50p + 49 &= 0 \implies (p - 49)(p - 1) = 0 \implies p = 49 \text{ or } p = 1 \\ \stackrel{p=x^2}{\implies} \implies x^2 &= 49 \text{ or } x^2 = 1 \implies x = \pm 7 \text{ or } x = \pm 1 \end{aligned}$$

Higher degree but only products of lower degrees

$$\begin{aligned} x^4(x - 2)^2 &= 0 \implies x^4 = 0 \text{ or } (x - 2)^2 = 0 \\ \implies x &= 0 \text{ or } x - 2 = 0 \implies x = 0 \text{ or } x = 2 \end{aligned}$$

(respectively 4-fold and 2-fold)

# Factoring and roots (zeros) of polynomials

Writing a polynomial as a product of two or more polynomials is called factoring.

Ex:  $x^3 - 3x^2 - 2x + 6 = (x - 3)(x^2 - 2)$

The polynomials  $x - 3$  and  $x^2 - 2$  are factors with degree 1 and 2.

Factoring breaks up a complicated polynomial into easier, lower degree pieces.

# Factoring and roots (zeros) of polynomials

Writing a polynomial as a product of two or more polynomials is called factoring.

Ex:  $x^3 - 3x^2 - 2x + 6 = (x - 3)(x^2 - 2)$

The polynomials  $x - 3$  and  $x^2 - 2$  are factors with degree 1 and 2.

Factoring breaks up a complicated polynomial into easier, lower degree pieces.

In the example, we can do more:

Ex:  $x^3 - 3x^2 - 2x + 6 = (x - 3)(x^2 - 2) = (x - 3)(x - \sqrt{2})(x + \sqrt{2})$

This polynomial is factored into three **linear** polynomials. We can't do any better, the polynomial is **factored completely**.

Knowing all roots of a polynomial  $\Leftrightarrow$  Knowing the factoring of a polynomial

# Polynomials: When to let factors stand as they are?

When a polynomial consists of one term of factors:

$$p(x) = (x + 1)^3 \text{ or } p(x) = (x - 1)(x + 2)$$

Most of the times, it is useful to let these stand, because:

- ▶ it is easy, no extra effort
- ▶ you see the zeros at once
- ▶ sometimes this is easier to differentiate/integrate (later in Calculus)  
(eg derivative or primitive of  $(x + 1)^{40}$ )
- ▶ useful for limits (eg  $\lim_{x \rightarrow 1^+} (x - 1)^3$  etc)

# Polynomials: When to write factors out?

When a polynomial consists of several terms (of factors), eg:

$$p(x) = (x + 1)^3 - (x - 1)^3$$

Most of the times, it is useful to write these out, because:

- ▶ Terms can cancel each other out, eg:

$$(x + 1)^3 - (x - 1)^3 = 6x^2 + 2 \text{ shows to be of degree 2 (not 3)}$$

## Examples: Higher order polynomial equations

- 1 Solve  $3x^4 = 9x^2$
- 2 Solve  $x^3 + 9x^2 + 26x + 24 = 0$
- 3 Solve  $x^4 - 26x^2 + 25 = 0$
- 4 What are all solutions to  $(x^2 - 1)^4(x + 3)^7 = 0$
- 5 Which of these polynomials has roots  $x = 1 \vee x = 2 \vee x = 3/4$ 
  - a)  $p(x) = (x + 1)(x + 2)(x + 3/4)$
  - b)  $p(x) = (x - 1)(x - 2)(x - 3/4)$

# Inequalities with polynomials

Finding solutions to inequalities of the kind

$$p(x) > d \quad p(x) \geq d \quad p(x) < d \quad p(x) \leq d$$

is done in the following steps:

- 1 Replace  $<$ ,  $\leq$ ,  $>$  or  $\geq$  by  $=$
- 2 Get a 0 to the right hand side
- 3 Mark the solutions (zeros) on the  $x$ -axis
- 4 Determine the signs for each interval (e.g., by filling in an arbitrary point)
- 5 Determine the solution of the inequality

Simple example: solve  $x^2 > 9$

# Inequalities with polynomials

Finding solutions to inequalities of the kind

$$p(x) > d \quad p(x) \geq d \quad p(x) < d \quad p(x) \leq d$$

is done in the following steps:

- 1 Replace  $<$ ,  $\leq$ ,  $>$  or  $\geq$  by  $=$
- 2 Get a 0 to the right hand side
- 3 Mark the solutions (zeros) on the  $x$ -axis
- 4 Determine the signs for each interval (e.g., by filling in an arbitrary point)
- 5 Determine the solution of the inequality

Simple example: solve  $x^2 > 9 \iff x \in (-\infty, -3) \cup (3, \infty)$

# Absolute value


$$|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

“Distance to 0”. Properties:

- ▶  $|x| = |-x|$
- ▶  $|xy| = |x| \cdot |y|$
- ▶  $|x + y| \leq |x| + |y|$
- ▶  $|x - a|$  “distance to  $a$ ”

For example:

- ▶  $|(-2)(-4)| = |-2| \cdot |-4|$
- ▶  $|2 + 4| = |2| + |4|, \quad |2 - 4| < |2| + |-4|$



# Absolute values: Examples

Attention!

$$(\sqrt{a})^2 = a \text{ but } \sqrt{a^2} = |a|$$

Ex:  $\sqrt{(-3)^2} = 3$

Only if you know that  $a \geq 0$ , then  $\sqrt{a^2} = |a| = a$ , eg

$$\sqrt{x^4 + 2x^2 + 1} = \sqrt{(x^2 + 1)^2} = |x^2 + 1| = x^2 + 1$$

The last step only because  $x^2 + 1$  always  $\geq 0$  (even  $\geq 1$ )

# Equations and inequalities with absolute values

## 1 Equations

split into the two parts and solve separately

## 2 Inequalities

- ▶ Just as for polynomial inequalities: replace  $<$  or  $\leq$  by  $=$
- ▶ Solve and mark solutions on the  $x$ -axis with 0s
- ▶ Determine sign by filling in arbitrary points within the intervals
- ▶ Determine the final solution

# Fractions

**Addition:** make denominators same, then you can add

$$\frac{1}{p} + \frac{1}{q} = \frac{q}{pq} + \frac{p}{pq} = \frac{p+q}{pq}$$

Up to “smallest common multiple”:

$$\frac{1}{4} + \frac{1}{12} = \frac{3}{12} + \frac{1}{12} = \frac{4}{12} = \frac{1}{3} \quad \frac{1}{6} + \frac{1}{8} = \frac{4}{24} + \frac{3}{24} = \frac{7}{24}$$

**Multiplication:** Multiply numerators and denominators

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$$

**Simplify:** divide numerator and denominator by the same factor

$$\frac{144}{30} = \frac{72}{15} = \frac{24}{5}$$

$$\frac{2x+6}{4} = \frac{x+3}{2} = \frac{1}{2}x + \frac{3}{2} \quad \text{but} \quad \frac{4}{2x+6} = \frac{2}{x+3} \neq \frac{2}{x} + \frac{2}{3} !$$

**Division** by a fraction is multiplication with the inverse

$$\frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \Big/ \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{ad}{bc}$$

# Equations with rational functions

Rational functions =  $\frac{\text{polynomial}}{\text{polynomial}}$        $R(x) = \frac{P(x)}{Q(x)}$

Two strategies:

(1) Crosswise multiplication

(2) Bring everything on the same denominator

$$(1) : \frac{3}{x+1} = \frac{2}{x+2}$$

# Equations with rational functions

Rational functions =  $\frac{\text{polynomial}}{\text{polynomial}}$        $R(x) = \frac{P(x)}{Q(x)}$

Two strategies:

(1) Crosswise multiplication

(2) Bring everything on the same denominator

$$(1) : \frac{3}{x+1} = \frac{2}{x+2} \implies 3(x+2) = 2(x+1) \text{ and } x \neq -1 \text{ and } x \neq -2$$

# Equations with rational functions

Rational functions =  $\frac{\text{polynomial}}{\text{polynomial}}$        $R(x) = \frac{P(x)}{Q(x)}$

Two strategies:

(1) Crosswise multiplication

(2) Bring everything on the same denominator

$$(1) : \frac{3}{x+1} = \frac{2}{x+2} \implies 3(x+2) = 2(x+1) \text{ and } x \neq -1 \text{ and } x \neq -2$$
$$\implies x = -4 \quad (\text{and } x \neq -1 \text{ and } x \neq -2)$$

(however the condition  $x \neq -1$  and  $x \neq -2$  is thus not relevant)

$$(2) : \frac{3(x+2)}{(x+1)(x+2)} - \frac{2(x+1)}{(x+1)(x+2)} = 0 \implies \frac{x+4}{(x+1)(x+2)} = 0$$

Thus numerator = 0:  $x = -4$

and denominator  $\neq 0$ :  $x \neq -1$  and  $x \neq -2$ , but that does not apply here

# Division with rest

Rewrite a fraction into an integer + “something with a numerator smaller than the denominator”:  $\frac{7}{4} = 1 + \frac{3}{4}$ :

This also works with rational functions:

$$\frac{\text{polynomial}}{\text{polynomial}} = \text{polynomial} + \frac{\text{polynomial P}}{\text{polynomial Q}}$$

so that the degree of P < degree of Q.

Previous example:  $x = 1$  was zero of  $x^3 + 2x^2 - x - 2$  and

$$\frac{x^3 + 2x^2 - x - 2}{x - 1} = x^2 + 3x + 2$$

with rest-term  $\frac{\text{polynomial P}}{\text{polynomial Q}}$  equal to zero.

# Division with rest, different formulation

Target: Write “rational function = polynomial + rational function” so that the degree of numerator < degree of denominator

Ex:

$$\frac{x^3 + 2}{x^2 + x}$$

Procedure: The same as for long division for finding zeros.  
However, the rest does not have to be 0.

# Long division with rest

Target: rational function = polynomial + other rational function  
while the degree of the numerator < degree of the denominator

$$x^2 + x \quad / \quad x^3 \quad \quad \quad +2 \quad \backslash$$

Conclusion

$$\frac{x^3 + 2}{x^2 + x} = x - 1 + \frac{x + 2}{x^2 + x}$$

Indeed: here degree of numerator is smaller than the degree of the denominator

Somewhat analog to  $\frac{7}{4} = 1 + \frac{3}{4}$ :

integer number + “something with numerator smaller than denominator”

# Long division with rest

Target: rational function = polynomial + other rational function  
while the degree of the numerator < degree of the denominator

$$\begin{array}{r} x^2 + x \quad / \quad x^3 \quad \quad \quad +2 \quad \backslash \quad x \\ x \cdot \quad (x^2 + x) \rightarrow x^3 + x^2 \end{array}$$

Conclusion

$$\frac{x^3 + 2}{x^2 + x} = x - 1 + \frac{x + 2}{x^2 + x}$$

Indeed: here degree of numerator is smaller than the degree of the denominator

Somewhat analog to  $\frac{7}{4} = 1 + \frac{3}{4}$ :

integer number + “something with numerator smaller than denominator”

# Long division with rest

Target: rational function = polynomial + other rational function  
while the degree of the numerator < degree of the denominator

$$\begin{array}{r} x^2 + x \quad / \quad x^3 \quad \quad +2 \quad \backslash \quad x \\ x \cdot \quad (x^2 + x) \rightarrow x^3 + x^2 \\ \hline -x^2 \quad \quad \quad +2 \end{array}$$

Conclusion

$$\frac{x^3 + 2}{x^2 + x} = x - 1 + \frac{x + 2}{x^2 + x}$$

Indeed: here degree of numerator is smaller than the degree of the denominator

Somewhat analog to  $\frac{7}{4} = 1 + \frac{3}{4}$ :

integer number + “something with numerator smaller than denominator”

## Long division with rest

Target: rational function = polynomial + other rational function  
while the degree of the numerator < degree of the denominator

$$\begin{array}{r}
 x^2 + x \quad / \quad x^3 \quad +2 \quad \backslash \quad \textcolor{blue}{x} - 1 \\
 \times \cdot \quad (x^2 + x) \rightarrow x^3 + x^2 \\
 \hline
 \end{array}$$

## Conclusion

$$\frac{x^3 + 2}{x^2 + x} = x - 1 + \frac{x + 2}{x^2 + x}$$

Indeed: here degree of numerator is smaller than the degree of the denominator

Somewhat analog to  $\frac{7}{4} = 1 + \frac{3}{4}$ :

integer number + “something with numerator smaller than denominator”

# Long division with rest

Target: rational function = polynomial + other rational function  
while the degree of the numerator < degree of the denominator

$$\begin{array}{r} x^2 + x \quad / \quad x^3 \quad \quad +2 \quad \backslash \quad x - 1 \\ x \cdot \quad (x^2 + x) \rightarrow x^3 + x^2 \\ \hline -x^2 \quad +2 \\ -1 \cdot \quad (x^2 + x) \rightarrow \quad -x^2 - x \\ \hline \quad \quad \quad x + 2 \end{array}$$

Conclusion

$$\frac{x^3 + 2}{x^2 + x} = x - 1 + \frac{x + 2}{x^2 + x}$$

Indeed: here degree of numerator is smaller than the degree of the denominator

Somewhat analog to  $\frac{7}{4} = 1 + \frac{3}{4}$ :

integer number + “something with numerator smaller than denominator”

# Inequalities, abs. values, rational funct.

1  $x^2 - 3x \geq -2$

2  $\frac{x+1}{x} \geq 2$

3 Simplify  $\sqrt{x^4 + 6x^2 + 9}$

4  $|2x - 5| < 6$

# Roots $\sqrt[n]{a}$

$$x^2 = a \implies x = \pm\sqrt{a}$$

$$x^3 = a \implies x = \sqrt[3]{a}$$

$$x^4 = a \implies x = \pm\sqrt[4]{a}$$

$$x^5 = a \implies x = \sqrt[5]{a}$$


here must be  $a \geq 0$

here  $a$  can be everything ( $a \in \mathbb{R}$ )

here must be  $a \geq 0$

here  $a$  can be everything ( $a \in \mathbb{R}$ )

etc; these are standard results that you can use in exams



Ex:  $x^3 = 8$  thus  $x^3 - 8 = 0$ : you could say right away  $x = 2$

but you could also solve this using the recipe for higher order polynomials

- ▶ guess solution:  $x = 2$
- ▶ long division:  $\frac{x^3-8}{x-2} = x^2 + 2x + 4$
- ▶ the discriminant of  $x^2 + 2x^2 + 4 = -12 < 0$   
thus no further solutions

## $n$ -th power roots: examples

$$x^3 = 27$$

## $n$ -th power roots: examples

$$x^3 = 27 \quad x = \sqrt[3]{27} =$$

## $n$ -th power roots: examples

$$\begin{aligned}x^3 &= 27 \\x^3 &= -27\end{aligned}$$

$$x = \sqrt[3]{27} = 3$$

## $n$ -th power roots: examples

$$x^3 = 27 \quad x = \sqrt[3]{27} = 3$$

$$x^3 = -27 \quad x = \sqrt[3]{-27} = -3$$

$$x^4 = 4$$

## $n$ -th power roots: examples

$$x^3 = 27 \quad x = \sqrt[3]{27} = 3$$

$$x^3 = -27 \quad x = \sqrt[3]{-27} = -3$$

$$x^4 = 4 \quad x = \pm \sqrt[4]{4} =$$

## $n$ -th power roots: examples

$$x^3 = 27$$

$$x = \sqrt[3]{27} = 3$$

$$x^3 = -27$$

$$x = \sqrt[3]{-27} = -3$$

$$x^4 = 4$$

$$x = \pm \sqrt[4]{4} = \pm \sqrt[2]{\sqrt[2]{4}} = \pm \sqrt[2]{2} = \pm \sqrt{2}$$

$$x^4 = -4$$

different notation:  $\pm 4^{\frac{1}{4}} = \pm (4^{\frac{1}{2}})^{\frac{1}{2}} = \pm 2^{\frac{1}{2}}$

## $n$ -th power roots: examples

$$x^3 = 27$$

$$x = \sqrt[3]{27} = 3$$

$$x^3 = -27$$

$$x = \sqrt[3]{-27} = -3$$

$$x^4 = 4$$

$$x = \pm \sqrt[4]{4} = \pm \sqrt[2]{\sqrt[2]{4}} = \pm \sqrt[2]{2} = \pm \sqrt{2}$$

$$x^4 = -4$$

different notation:  $\pm 4^{\frac{1}{4}} = \pm (4^{\frac{1}{2}})^{\frac{1}{2}} = \pm 2^{\frac{1}{2}}$

no solutions

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} = 5 \cdot 3\sqrt{5} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} = 5 \cdot 3\sqrt{5} = 15\sqrt{5}$$

Remove the root from the denominator:

$$\frac{1}{\sqrt{2}} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} = 5 \cdot 3\sqrt{5} = 15\sqrt{5}$$

Remove the root from the denominator:

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2} = \frac{1}{2}\sqrt{2}$$

"Root trick": Make use of  $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) = a - b$ :

$$\frac{1}{\sqrt{a} - 2} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} = 5 \cdot 3\sqrt{5} = 15\sqrt{5}$$

Remove the root from the denominator:

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2} = \frac{1}{2}\sqrt{2}$$

"Root trick": Make use of  $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) = a - b$ :

$$\frac{1}{\sqrt{a} - 2} = \frac{1}{\sqrt{a} - 2} \cdot \frac{\sqrt{a} + 2}{\sqrt{a} + 2} = \frac{\sqrt{a} + 2}{(\sqrt{a})^2 - 2^2} = \frac{\sqrt{a} + 2}{a - 4}$$

$$\frac{1}{\sqrt{a} + \sqrt{b}} =$$

# Working with roots

Basic property (definition):  $\sqrt{x} \cdot \sqrt{x} = x$  ( $x \geq 0$ )

Simplification in steps: decompose into factors:

$$\sqrt{1125} = \sqrt{5 \cdot 225} = \sqrt{5 \cdot 5 \cdot 45} = 5\sqrt{45} = 5\sqrt{9 \cdot 5} = 5 \cdot 3\sqrt{5} = 15\sqrt{5}$$

Remove the root from the denominator:

$$\frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{2} = \frac{1}{2}\sqrt{2}$$

"Root trick": Make use of  $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) = a - b$ :

$$\frac{1}{\sqrt{a} - 2} = \frac{1}{\sqrt{a} - 2} \cdot \frac{\sqrt{a} + 2}{\sqrt{a} + 2} = \frac{\sqrt{a} + 2}{(\sqrt{a})^2 - 2^2} = \frac{\sqrt{a} + 2}{a - 4}$$

$$\frac{1}{\sqrt{a} + \sqrt{b}} = \frac{1}{\sqrt{a} + \sqrt{b}} \cdot \frac{\sqrt{a} - \sqrt{b}}{\sqrt{a} - \sqrt{b}} = \frac{\sqrt{a} - \sqrt{b}}{(\sqrt{a})^2 - (\sqrt{b})^2} = \frac{\sqrt{a} - \sqrt{b}}{a - b}$$

= and  $<$ ,  $\geq$  with roots

You have to pay attention to one thing: by taking a square you can come to solutions that cannot exist  $\implies$  extra check

$$\sqrt{4x + 1} = x - 1$$

= and  $<$ ,  $\geq$  with roots

You have to pay attention to one thing: by taking a square you can come to solutions that cannot exist  $\implies$  extra check

$$\begin{aligned}\sqrt{4x+1} = x - 1 &\implies 4x + 1 = (x - 1)^2 \implies x^2 - 6x = 0 \\ &\implies x(x - 6) = 0 \implies x = 0 \text{ or } x = 6\end{aligned}$$

Check:

## = and <, $\geq$ with roots

You have to pay attention to one thing: by taking a square you can come to solutions that cannot exist  $\Rightarrow$  extra check

$$\begin{aligned}\sqrt{4x+1} = x - 1 &\Rightarrow 4x + 1 = (x - 1)^2 \Rightarrow x^2 - 6x = 0 \\ &\Rightarrow x(x - 6) = 0 \Rightarrow x = 0 \text{ or } x = 6\end{aligned}$$

Check:  $x = 0$  is no solution,  $x = 6$  is

- ▶ Equations: solve by taking squares (evt several times)  
check answers always at the end
- ▶ Inequalities: solve the equality  
also pay attention that for the solutions (intervals) all arguments under the roots have to be  $\geq 0$

# Algebraic skills: frequently made mistakes

**FAIL:**  $\sqrt{x+y} = \sqrt{x} + \sqrt{y}$

**GOOD:**  $\sqrt{xy} = \sqrt{x}\sqrt{y}$  (if  $x \geq 0, y \geq 0$ )

**FAIL:**  $x(x-2) = 1 \implies x = 1 \text{ or } x - 2 = 1$

**GOOD:**  $x(x-2) = 0 \implies x = 0 \text{ or } x - 2 = 0$

**FAIL:**  $x^2 = 3x \implies x = 3$

**GOOD:**  $x^2 = 3x \implies x = 3 \text{ or } x = 0$  because  $x(x-3) = 0$

**FAIL:**  $\sqrt{x^2} = x$

**GOOD:**  $\sqrt{x^2} = |x|$

**FAIL:**  $x^2 = a \implies x = \sqrt{a}$

**GOOD:**  $x^2 = a \implies x = \pm\sqrt{a}$


**FAIL:**  $\frac{2}{x+3} = \frac{2}{x} + \frac{2}{3}$

**GOOD:**  $\frac{x+3}{2} = \frac{x}{2} + \frac{3}{2}$

# The plane $\mathbb{R}^2$ , distance, lines

$\mathbb{R}^2$ : notation for a 2-dimensional space

Distance  $d$  between 2 points  $(x_1, y_1) = (-1, 1)$  adn  $(x_2, y_2) = (-2, 3)$



- $\Delta x = x_2 - x_1 = -2 - (-1) = -1$ ,     $\Delta y = y_2 - y_1 = 3 - 1 = 2$
- $d = \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{1 + 4} = \sqrt{5}$     (Pythagoras! [A B12])

Slope  $m$  of the line between the points  $(-1, 1)$  and  $(-2, 3)$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x} = \frac{2}{-1} = -2 \quad [\text{A P.2 E7}]$$

# Lines (I)

The line through 2 points  $(x_1, y_1)$  and  $(x_2, y_2)$

**two-point equation:**  $(y - y_1)\Delta x = (x - x_1)\Delta y$

(Remember:  $\Delta x = x_2 - x_1$  and  $\Delta y = y_2 - y_1$ ).

**point-slope equation:**  $y = m(x - x_1) + y_1$

Formula is correct because filling in yields:

- ▶  $(x_1, y_1)$ :  $0 \cdot \Delta x = 0 \cdot \Delta y$
- ▶  $(x_2, y_2)$ :  $(y_2 - y_1)\Delta x = (x_2 - x_1)\Delta y \implies \Delta y \Delta x = \Delta x \Delta y$

## Lines (II)

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$\begin{aligned}(y - y_1) \underbrace{\Delta x}_{=-1} &= (x - x_1) \underbrace{\Delta y}_{=2} \implies (y - 1) \cdot -1 = (x + 1) \cdot 2 \\ &\implies (y - 1) = -2(x + 1) \\ &\implies y = -2x - 1\end{aligned}$$

because  $\Delta x = x_2 - x_1 = -2 - (-1) = -1$ ,  $\Delta y = y_2 - y_1 = 3 - 1 = 2$ .  
Straight lines thus have the form  $ax + by = c$ . Special cases:

- ▶  $a = 0$ : horizontal line
- ▶  $b = 0$ : vertical line

Ex: Set point  $(-2, 3)$

Horizontal line through the point has the eq.:

## Lines (II)

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$\begin{aligned}(y - y_1) \underbrace{\Delta x}_{=-1} &= (x - x_1) \underbrace{\Delta y}_{=2} \implies (y - 1) \cdot -1 = (x + 1) \cdot 2 \\ &\implies (y - 1) = -2(x + 1) \\ &\implies y = -2x - 1\end{aligned}$$

because  $\Delta x = x_2 - x_1 = -2 - (-1) = -1$ ,  $\Delta y = y_2 - y_1 = 3 - 1 = 2$ .  
Straight lines thus have the form  $ax + by = c$ . Special cases:

- ▶  $a = 0$ : horizontal line
- ▶  $b = 0$ : vertical line

Ex: Set point  $(-2, 3)$

Horizontal line through the point has the eq.:  $y = 3$  ( $b = 1$ )

Vertical line through this point has the eq.:

## Lines (II)

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$\begin{aligned}(y - y_1) \underbrace{\Delta x}_{=-1} &= (x - x_1) \underbrace{\Delta y}_{=2} \implies (y - 1) \cdot -1 = (x + 1) \cdot 2 \\ &\implies (y - 1) = -2(x + 1) \\ &\implies y = -2x - 1\end{aligned}$$

because  $\Delta x = x_2 - x_1 = -2 - (-1) = -1$ ,  $\Delta y = y_2 - y_1 = 3 - 1 = 2$ .  
Straight lines thus have the form  $ax + by = c$ . Special cases:

- ▶  $a = 0$ : horizontal line
- ▶  $b = 0$ : vertical line

Ex: Set point  $(-2, 3)$

Horizontal line through the point has the eq.:  $y = 3$  ( $b = 1$ )

Vertical line through this point has the eq.:  $x = -2$  ( $a = 1$ )

## Lines (III)

Intersections with the  $x$ - and  $y$ -axes are called “intercepts”

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$y = -2x - 1$$

Where does this line intersect the  $x$ -axis?

## Lines (III)

Intersections with the  $x$ - and  $y$ -axes are called “intercepts”

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$y = -2x - 1$$

Where does this line intersect the  $x$ -axis? Then  $y = 0$  thus

$$0 = -2x - 1 \implies -2x = 1 \implies x = -\frac{1}{2} \text{ (thus “x-intercept”} = -\frac{1}{2})$$

Where does this line intersect the  $y$ -axis?

## Lines (III)

Intersections with the  $x$ - and  $y$ -axes are called “intercepts”

Ex: The line through  $(x_1, y_1) = (-1, 1)$  and  $(x_2, y_2) = (-2, 3)$  is

$$y = -2x - 1$$

Where does this line intersect the  $x$ -axis? Then  $y = 0$  thus

$$0 = -2x - 1 \implies -2x = 1 \implies x = -\frac{1}{2} \text{ (thus “x-intercept”} = -\frac{1}{2})$$

Where does this line intersect the  $y$ -axis?

Then  $x = 0$  thus  $y = -2 \cdot 0 - 1 = -1$  (thus “y-intercept” = -1)

Do [A P.2 T38]

## Vertical and general lines (I)

A line is vertical if  $x_1 = x_2$ , then it follows

$$(y - y_1)\Delta x = (x - x_1)\Delta y \implies 0 = (x - x_1)\Delta y \implies 0 = x - x_1$$

A vertical line in  $\mathbb{R}^2$  is given by  $x = x_1$ .

Other lines:  $x_1 \neq x_2 \implies \Delta x \neq 0$  thus


$$\begin{aligned}(y - y_1)\Delta x &= (x - x_1)\Delta y \implies (y - y_1) = \frac{\Delta y}{\Delta x}(x - x_1) \implies \\(y - y_1) &= m(x - x_1)\end{aligned}$$

A general line in  $\mathbb{R}^2$  is described by the equation  $y = mx + b$

with  $b = m \cdot x_1 + y_1$  the  $y$ -intercept and  $m = \Delta y / \Delta x$  the slope.

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:



Ex: Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

The given line has the slope  $-2$

Thus the sought line has the slope  $\frac{1}{2}$

because  $-2 \cdot \frac{1}{2} = -1$


Thus  $y = \frac{1}{2}x + b$ , now fill in  $(1, 1)$ :

$$1 = \frac{1}{2} + b \implies b = \frac{1}{2} \text{ so the line } y = \frac{1}{2}x + \frac{1}{2}. \text{ [Do A P.2 T49]}$$

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$




Ex: Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$




Ex: Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

The given line has the slope

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$



**Ex:** Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?


The given line has the slope  $-2$

Thus the sought line has the slope

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$



**Ex:** Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

The given line has the slope  $-2$


Thus the sought line has the slope  $\frac{1}{2}$

because  $-2 \cdot \frac{1}{2} = -1$

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$



**Ex:** Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

The given line has the slope  $-2$

Thus the sought line has the slope  $\frac{1}{2}$


because  $-2 \cdot \frac{1}{2} = -1$

Thus  $y = \frac{1}{2}x + b$ ,

# Perpendicular lines

For lines with slopes  $m_1$  and  $m_2$  that are perpendicular to each other:

$$m_1 \cdot m_2 = -1$$



Ex: Which line is perpendicular to  $y = -2x - 1$  and goes through  $(1, 1)$ ?

The given line has the slope  $-2$

Thus the sought line has the slope  $\frac{1}{2}$

because  $-2 \cdot \frac{1}{2} = -1$

Thus  $y = \frac{1}{2}x + b$ , now fill in  $(1, 1)$ :

$$1 = \frac{1}{2} + b \implies b = \frac{1}{2} \text{ so the line } y = \frac{1}{2}x + \frac{1}{2}. \text{ [Do A P.2 T49]}$$

Circle with center  $(0, 0)$  and radius  $r$

Circle with radius  $r$ , center  $(0,0)$  has equation

## Circle with center $(0, 0)$ and radius $r$

Circle with radius  $r$ , center  $(0,0)$  has equation

$$x^2 + y^2 = r^2 \quad \text{or}$$

## Circle with center $(0, 0)$ and radius $r$

Circle with radius  $r$ , center  $(0, 0)$  has equation

$$x^2 + y^2 = r^2 \quad \text{or} \quad \left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1 \quad \text{or} \quad \frac{x^2}{r^2} + \frac{y^2}{r^2} = 1$$

Distance between  $(x, y)$  and  $(1, 2)$  is

## Circle with center $(0, 0)$ and radius $r$

Circle with radius  $r$ , center  $(0, 0)$  has equation

$$x^2 + y^2 = r^2 \quad \text{or} \quad \left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1 \quad \text{or} \quad \frac{x^2}{r^2} + \frac{y^2}{r^2} = 1$$

Distance between  $(x, y)$  and  $(1, 2)$  is  $\sqrt{(x - 1)^2 + (y - 2)^2}$

If the distance is equal to  $r$  is then holds

$$\sqrt{(x - 1)^2 + (y - 2)^2} = r \iff (x - 1)^2 + (y - 2)^2 = r^2.$$

General equation of a circle in  $\mathbb{R}^2$  with center  $(a, b)$  and radius  $r$ :

$$(x - a)^2 + (y - b)^2 = r^2$$

Circle with center  $(1, 2)$  and radius 3:

## Circle with center $(0, 0)$ and radius $r$

Circle with radius  $r$ , center  $(0, 0)$  has equation

$$x^2 + y^2 = r^2 \quad \text{or} \quad \left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1 \quad \text{or} \quad \frac{x^2}{r^2} + \frac{y^2}{r^2} = 1$$

Distance between  $(x, y)$  and  $(1, 2)$  is  $\sqrt{(x - 1)^2 + (y - 2)^2}$

If the distance is equal to  $r$  is then holds

$$\sqrt{(x - 1)^2 + (y - 2)^2} = r \iff (x - 1)^2 + (y - 2)^2 = r^2.$$

General equation of a circle in  $\mathbb{R}^2$  with center  $(a, b)$  and radius  $r$ :

$$(x - a)^2 + (y - b)^2 = r^2$$

Circle with center  $(1, 2)$  and radius 3:

$$(x - 1)^2 + (y - 2)^2 = 3^2$$

## Disc, circle and “hole”

The **circle** with radius  $r$  and center  $(a, b)$  consists of all points  $(x, y)$  for which it holds

$$(x - a)^2 + (y - b)^2 = r^2 \quad \text{circle line}$$

The **disk** with radius  $r$  and center  $(a, b)$  consists of all points  $(x, y)$  **inside** the circle for which it holds

$$(x - a)^2 + (y - b)^2 < r^2 \quad \text{circle interior} \rightarrow \text{open disk}$$

$$(x - a)^2 + (y - b)^2 \leq r^2 \quad \text{circle interior + line} \rightarrow \text{closed disk}$$

The “**hole**” with radius  $r$  and center  $(a, b)$  consists of alle points  $(x, y)$  **outside** the circle for which it holds

$$(x - a)^2 + (y - b)^2 > r^2 \quad \text{circle exterior}$$

# Ellipse with center $(0, 0)$ and “radii” $r_x$ and $r_y$

We saw:


Circle with radius  $r$  and center  $(0, 0)$ :

$$\left(\frac{x}{r}\right)^2 + \left(\frac{y}{r}\right)^2 = 1$$

ellipse

$$\left(\frac{x}{r_x}\right)^2 + \left(\frac{y}{r_y}\right)^2 = 1$$

is thus of the form  $cx^2 + dy^2 = e$




General ellipse with center  $(a, b)$  and semi-axes  $r_x$  and  $r_y$

$$\left(\frac{x-a}{r_x}\right)^2 + \left(\frac{y-b}{r_y}\right)^2 = 1$$

# Parabolas


Parabolas with the origin as vertex and the y-axis as axis of symmetry:  
 $y = ax^2$  with  $a \neq 0$ .

Parabolas with  $(x_V, y_V)$  as vertex and the y-axis as axis of symmetry:




# Hyperbola

For example:  $\left(\frac{x}{r_x}\right)^2 - \left(\frac{y}{r_y}\right)^2 = 1$  and  $\left(\frac{x}{r_x}\right)^2 - \left(\frac{y}{r_y}\right)^2 = -1$



Also:  $xy = c$  thus  $y = \frac{c}{x}$



# Overview of quadratic equations

Standard form has “= 1” on the right hand side

Circle ex:  $\left(\frac{x-a}{r}\right)^2 + \left(\frac{y-b}{r}\right)^2 = 1 \quad \text{or} \quad x^2 + y^2 = r^2$

Ellipse ex:  $\left(\frac{x-a}{r_x}\right)^2 + \left(\frac{y-b}{r_y}\right)^2 = 1 \quad \text{or} \quad cx^2 + dy^2 = e$

Parabola ex:  $\left(\frac{x-a}{r_x}\right)^2 + \frac{y-b}{r_y} = 1 \quad \text{or} \quad y = cx^2 + d$

Hyperbola ex:  $\left(\frac{x-a}{r_x}\right)^2 - \left(\frac{y-b}{r_y}\right)^2 = \pm 1 \quad \text{or} \quad xy = c$

# Definition of Curve

A curve is an equation of the form  $f(x, y) = 0$ .

Ex:

- ▶  $f(x, y) = x^2 - (2x - y^2) = 0$   
 $\rightarrow x^2 = 2x - y^2$

- ▶  $f(x, y) = 2y - 3x + 2 = 0$   
 $\rightarrow 2y = 3x - 2$  (straight line)

- ▶  $f(x, y) = y - (3x + 2)$   
 $\rightarrow y = 3x + 2$  (straight line)

What kind of curve is  $x^2 = 2x - y^2$ ?

ellipse, parabola, ...?

## Determining the type of curve

Now what does  $x^2 = 2x - y^2$  represent?

## Determining the type of curve

Now what does  $x^2 = 2x - y^2$  represent?

Complete the square

$$x^2 - 2x + y^2 = 0$$

## Determining the type of curve

Now what does  $x^2 = 2x - y^2$  represent?

Complete the square

$$x^2 - 2x + y^2 = 0$$

$$\implies x^2 - 2x + 1 - 1 + y^2 = 0$$

## Determining the type of curve

Now what does  $x^2 = 2x - y^2$  represent?

Complete the square

$$x^2 - 2x + y^2 = 0$$

$$\implies x^2 - 2x + 1 - 1 + y^2 = 0$$

$$\implies (x - 1)^2 + y^2 = 1$$


# Determining the type of curve

Now what does  $x^2 = 2x - y^2$  represent?

Complete the square

$$\begin{aligned}x^2 - 2x + y^2 &= 0 \\ \implies x^2 - 2x + 1 - 1 + y^2 &= 0 \\ \implies (x - 1)^2 + y^2 &= 1\end{aligned}$$

Circle with radius 1 and center (1,0)



See [A page 18], do [A, P.2 T9,11] and [A P.3 T7,15]

# Determining the type of curve

What does  $x^2 = 3x - y^2$  represent?

# Determining the type of curve

What does  $x^2 = 3x - y^2$  represent?

Complete the square

$$x^2 - 3x + y^2 = 0$$

# Determining the type of curve

What does  $x^2 = 3x - y^2$  represent?

Complete the square

$$x^2 - 3x + y^2 = 0$$

$$\implies x^2 - \left(\frac{3}{2} + \frac{3}{2}\right)x + \left(\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2 + y^2 = 0$$

# Determining the type of curve

What does  $x^2 = 3x - y^2$  represent?

Complete the square

$$x^2 - 3x + y^2 = 0$$

$$\implies x^2 - \left(\frac{3}{2} + \frac{3}{2}\right)x + \left(\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2 + y^2 = 0$$

$$\implies \left(x - \frac{3}{2}\right)^2 + y^2 = \left(\frac{3}{2}\right)^2$$

# Determining the type of curve

What does  $x^2 = 3x - y^2$  represent?

Complete the square

$$x^2 - 3x + y^2 = 0$$

$$\implies x^2 - \left(\frac{3}{2} + \frac{3}{2}\right)x + \left(\frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2 + y^2 = 0$$

$$\implies \left(x - \frac{3}{2}\right)^2 + y^2 = \left(\frac{3}{2}\right)^2$$

Circle with radius  $3/2$  and center  $(3/2, 0)$

# Functions and graphs


Function  $f$  is a rule that:

prescribes for each  $x \in D$  (domain) exactly one  $f(x)$

Notation:  $y = f(x)$

$x$  is the independent,  $y$  is the dependent variable


Function or not?



$$y = f(x) = x^2$$

function


Do [A P.4 T7]



$$y = \pm\sqrt{x}$$

no function

but curve



$$y = \sqrt{x}$$

function on  $[0, \infty)$

no function on  $\mathbb{R}$

# Functions and curves

Function  $y = g(x)$  is described by the curve

$$y - g(x) = 0$$

that means by  $f(x, y) = 0$  with  $f(x, y) = y - g(x)$ . Each function “is” therefore a curve.

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined  
("all possible inputs  $x$ "). Pay attention:

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined  
("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$       (also for  ${}^{10}\log$ )

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined  
("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$       (also for  ${}^{10}\log$ )

**Range** of  $f$ : set of all possible outputs  $y = f(x)$

Ex:  $f(x) = \frac{1}{x}$ : domain =

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined  
("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$       (also for  ${}^{10}\log$ )

**Range** of  $f$ : set of all possible outputs  $y = f(x)$

Ex:  $f(x) = \frac{1}{x}$ : domain =  $(-\infty, 0) \cup (0, \infty)$ , (or  $\mathbb{R} \setminus \{0\}$ )

range =

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined ("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$  (also for  ${}^{10}\log$ )

**Range** of  $f$ : set of all possible outputs  $y = f(x)$

Ex:  $f(x) = \frac{1}{x}$ : domain =  $(-\infty, 0) \cup (0, \infty)$ , (or  $\mathbb{R} \setminus \{0\}$ )  
range =  $(-\infty, 0) \cup (0, \infty)$

Ex:  $f(x) = \sqrt{x - 2}$ : domain =

# Domain and range

**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined ("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$  (also for  ${}^{10}\log$ )

**Range** of  $f$ : set of all possible outputs  $y = f(x)$

Ex:  $f(x) = \frac{1}{x}$ : domain =  $(-\infty, 0) \cup (0, \infty)$ , (or  $\mathbb{R} \setminus \{0\}$ )

$$\text{range} = (-\infty, 0) \cup (0, \infty)$$

Ex:  $f(x) = \sqrt{x - 2}$ : domain =  $[2, \infty)$ , range =

# Domain and range



**Domain** of  $f$ : the set of all  $x$  for which  $f(x)$  is defined ("all possible inputs  $x$ "). Pay attention:

- ▶ number under root must be  $\geq 0$
- ▶ do not divide by 0
- ▶  $\ln(x)$ :  $x$  must be  $> 0$  (also for  ${}^{10}\log$ )

**Range** of  $f$ : set of all possible outputs  $y = f(x)$

Ex:  $f(x) = \frac{1}{x}$ : domain =  $(-\infty, 0) \cup (0, \infty)$ , (or  $\mathbb{R} \setminus \{0\}$ )  
range =  $(-\infty, 0) \cup (0, \infty)$

Ex:  $f(x) = \sqrt{x-2}$ : domain =  $[2, \infty)$ , range =  $[0, \infty)$



Do [A P.4 T6] and [A P.6 T16]

# Even/odd function I

## Even function

- ▶  $f(x) = f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as if it is mirrored at the  $y$ -axis
- ▶ ex:  $f(x) = x^2$ ,  $f(x) = x^6 + x^2$ ,  
 $f(x) = 1$ ,  $f(x) = \cos(x)$
- ▶ sums of even powers in  $x$  are even

## Odd function

- ▶  $f(x) = -f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as first being mirrored at the  $y$ -axis, then at  $x$ -axis
- ▶ ex:  $f_1(x) = x$ ,  $f_2(x) = x^3$ ,  
 $f_3(x) = x^7 + x^3$ ,  $f(x) = \sin(x)$
- ▶ sums of odd powers in  $x$  are odd

Ex:  $f(x) = x^2 + x \Rightarrow f(-x) = (-x)^2 + (-x) = x^2 - x$

Is function even? Is  $x^2 - x = x^2 + x$  for all  $x$ ?

# Even/odd function I

## Even function

- ▶  $f(x) = f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as if it is mirrored at the  $y$ -axis
- ▶ ex:  $f(x) = x^2$ ,  $f(x) = x^6 + x^2$ ,  
 $f(x) = 1$ ,  $f(x) = \cos(x)$
- ▶ sums of even powers in  $x$  are even

## Odd function

- ▶  $f(x) = -f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as first being mirrored at the  $y$ -axis, then at  $x$ -axis
- ▶ ex:  $f_1(x) = x$ ,  $f_2(x) = x^3$ ,  
 $f_3(x) = x^7 + x^3$ ,  $f(x) = \sin(x)$
- ▶ sums of odd powers in  $x$  are odd

Ex:  $f(x) = x^2 + x \Rightarrow f(-x) = (-x)^2 + (-x) = x^2 - x$

Is function even? Is  $x^2 - x = x^2 + x$  for all  $x$ ? No

Is function odd? Is  $x^2 - x = -(x^2 + x)$  for all  $x$ ?

# Even/odd function I

## Even function

- ▶  $f(x) = f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as if it is mirrored at the  $y$ -axis
- ▶ ex:  $f(x) = x^2$ ,  $f(x) = x^6 + x^2$ ,  
 $f(x) = 1$ ,  $f(x) = \cos(x)$
- ▶ sums of even powers in  $x$  are even

## Odd function

- ▶  $f(x) = -f(-x)$  for all  $x$
- ▶ means: graph of  $f$  is the same as first being mirrored at the  $y$ -axis, then at  $x$ -axis
- ▶ ex:  $f_1(x) = x$ ,  $f_2(x) = x^3$ ,  
 $f_3(x) = x^7 + x^3$ ,  $f(x) = \sin(x)$
- ▶ sums of odd powers in  $x$  are odd

Ex:  $f(x) = x^2 + x \Rightarrow f(-x) = (-x)^2 + (-x) = x^2 - x$

Is function even? Is  $x^2 - x = x^2 + x$  for all  $x$ ? No

Is function odd? Is  $x^2 - x = -(x^2 + x)$  for all  $x$ ? No

A function does NOT have to be even or odd!

# Symmetries


Even and odd are special cases of (anti-)symmetry:  
namely around  $x = 0$

One can also have symmetries around  $x = 1$ :

# Symmetries

Even and odd are special cases of (anti-)symmetry:  
namely around  $x = 0$

One can also have symmetries around  $x = 1$ :  $f(x) = (x - 1)^2$




This can easily be seen by substituting  $w = x - 1$ :  $y = w^2$ , even function:  
symmetric around  $w = 0$ ,  
this coincides with  $x = 1$

**Ex:** function which is anti-symmetric around  $x = 2$ ?

# Symmetries

Even and odd are special cases of (anti-)symmetry:  
namely around  $x = 0$

One can also have symmetries around  $x = 1$ :  $f(x) = (x - 1)^2$



This can easily be seen by substituting  $w = x - 1$ :  $y = w^2$ , even function:  
symmetric around  $w = 0$ ,  
this coincides with  $x = 1$

**Ex:** function which is anti-symmetric around  $x = 2$ ?

$f(x) = x - 2$  or  $f(x) = (x - 2)^3$  or ...

# Functions and their graphs

- ▶ Give domain and range of  $f(x) = -x^2 + 2x - 7$
- ▶ Give domain and range of for  $f(x) = \sqrt{9 - x^2}$
- ▶ Draw the graph of  $y = -x^2$ . How does the equation change for shifting by 7 left, right, up, down?
- ▶ Give function and domain for shifting  $f(x) = \sqrt{x}$  1 to the left
- ▶ Even, odd or nothing?

$$\sqrt{1 - x^2}, \quad x + 1, \quad x^3 + x$$

# Addition and Multiplication

If  $f$  and  $g$  functions are, then also sums and products of them are functions:

operation		example
$(fg)(x)$	$= f(x) \cdot g(x)$	$x^2 \sin(x)$
$(f + g)(x)$	$= f(x) + g(x)$	$x^2 + \sin(x)$
$(f - g)(x)$	$= f(x) - g(x)$	$x^2 - \sin(x)$
$\left(\frac{f}{g}\right)(x)$	$= \frac{f(x)}{g(x)}$	$x^2 / \sin(x)$

Do [A P.5 T2]

# Composite functions – one after the other

Composite functions:  $(f \circ g)(x) = f(g(x))$  “ $f$  after  $g$ ”

- ▶  $f(x) = 4x - 1$ ,  $g(x) = x^2 - 1$   
 $(f \circ g)(x) =$

# Composite functions – one after the other

Composite functions:  $(f \circ g)(x) = f(g(x))$  “ $f$  after  $g$ ”

- ▶  $f(x) = 4x - 1, \quad g(x) = x^2 - 1$   
 $(f \circ g)(x) = 4(x^2 - 1) - 1$   
 $(g \circ f)(x) =$

# Composite functions – one after the other

Composite functions:  $(f \circ g)(x) = f(g(x))$  “ $f$  after  $g$ ”

- ▶  $f(x) = 4x - 1, \quad g(x) = x^2 - 1$   
 $(f \circ g)(x) = 4(x^2 - 1) - 1$   
 $(g \circ f)(x) = (4x - 1)^2 - 1$
- ▶  $\sqrt{\sin(x^2)} = f(g(h(x)))$  with:

# Composite functions – one after the other

Composite functions:  $(f \circ g)(x) = f(g(x))$  “ $f$  after  $g$ ”

- ▶  $f(x) = 4x - 1, \quad g(x) = x^2 - 1$   
 $(f \circ g)(x) = 4(x^2 - 1) - 1$   
 $(g \circ f)(x) = (4x - 1)^2 - 1$
- ▶  $\sqrt{\sin(x^2)} = f(g(h(x)))$  with:  
 $f(x) = \sqrt{x}, \quad g(x) = \sin(x), \quad h(x) = x^2$


$D_{f \circ g}$  consists of all  $x \in D_g$  for which  $g(x) \in D_f$

Ex:  $f(x) = 1/x, \quad g(x) = 1/x$ . Then is  $(f \circ g)(x) = x$  and  $D_{f \circ g} = \mathbb{R} - \{0\}$ .  
Do [A P.5 T6,7,9,13,15]

## Week 2: Trigonometrics, powers, and vectors

- ▶ Trigonometric functions ( $\sin, \dots$ )
- ▶ Powers
- ▶ Vectors in 2 and 3 dimensions
- ▶ Lines and planes
- ▶ Distances, angles and the inner product
- ▶ The Cross Product

# Sound



Sound, water, electro-magnetic and other waves are a sum of elementary waves:  $A \sin(f \cdot 2\pi \cdot t)$

►  $\sum_{k=1}^n A_k \sin(2\pi \cdot f_k \cdot t)$

where the frequency  $f$  determines the pitch and  
the amplitude  $A$  the intensity of the sound.  $\sin$  is the [sine](#) function

# Trigonometric functions: sin, cos, tan


$$\sin = \frac{\text{opp}}{\text{hyp}}, \cos = \frac{\text{adj}}{\text{hyp}}, \tan = \frac{\text{opp}}{\text{adj}}$$

Relation degrees – radians:

$$30^\circ = \frac{\pi}{6}, 45^\circ = \frac{\pi}{4}, 60^\circ = \frac{\pi}{3}, 90^\circ = \frac{\pi}{2}$$

Radians = the length of a piece of the circle


$$\text{Complete circle} = 2\pi = 360^\circ$$



Point on a circle with radius 1 has the coordinates  $(\cos(t), \sin(t))$

This agrees with  $\sin^2(t) + \cos^2(t) = (\sin(t))^2 + (\cos(t))^2 = 1$

Similar: point on the circle with radius  $a$  has the coordinates  $(a\cos(t), a\sin(t)) = a(\cos(t), \sin(t))$




# The 4 quadrants


- 1  $t \in [0, \pi/2] \Rightarrow (\cos(t), \sin(t))$  is in the 1st quadrant
- 2  $t \in [\pi/2, \pi] \Rightarrow (\cos(t), \sin(t))$  is in the 2nd quadrant
- 3  $t \in [\pi, 3/2 \cdot \pi] \Rightarrow (\cos(t), \sin(t))$  is in the 3rd quadrant
- 4  $t \in [3/2 \cdot \pi, 2\pi] \Rightarrow (\cos(t), \sin(t))$  is in the 4th quadrant

so that we define

- 1 the 1st quadrant:  $t \in [0, \pi/2]$
- 2 the 2nd quadrant:  $t \in [\pi/2, \pi]$
- 3 the 3rd quadrant:  $t \in [\pi, 3/2 \cdot \pi]$
- 4 the 4th quadrant:  $t \in [3/2 \cdot \pi, 2\pi]$



# Sine, Cosine, and Tangens functions



$$\sin(t + 2\pi) = \sin(t)$$

odd function:

$$\sin(-t) = -\sin(t)$$

$$\sin(t) = 0$$


$$\implies t = k\pi \quad (k \in \mathbb{Z})$$

$$\sin(t) = 1$$

$$\implies t = \frac{\pi}{2} + 2k\pi$$

$$\sin(t) = -1$$

$$\implies t = \frac{3\pi}{2} + 2k\pi$$



$$\cos(t + 2\pi) = \cos(t)$$

even function:


$$\cos(-t) = \cos(t)$$

$$\cos(t) = 0$$

$$\implies t = \frac{\pi}{2} + k\pi$$

$$\sin(t) = \cos(\frac{\pi}{2} - t)$$

$$\cos(t) = \sin(\frac{\pi}{2} - t)$$



$$\tan(t) = \frac{\sin(t)}{\cos t}$$

odd function:

$$\tan(-t) = -\tan(t)$$


$$\tan(t) = 0$$

$$\implies t = k\pi \quad (k \in \mathbb{Z})$$

vertical asymptote in

$$\frac{\pi}{2} + k\pi, \quad k \in \mathbb{Z}$$

# Sine, Cosine, and Tangens functions



$$\sin(x + 2\pi) = \sin(x)$$

odd function:

$$\sin(-x) = -\sin(x)$$

$$\sin(x) = 0$$


$$\implies x = k\pi \quad (k \in \mathbb{Z})$$

$$\sin(x) = 1$$

$$\implies x = \frac{\pi}{2} + 2k\pi$$

$$\sin(x) = -1$$

$$\implies x = \frac{3\pi}{2} + 2k\pi$$



$$\cos(x + 2\pi) = \cos(x)$$

even function:


$$\cos(-x) = \cos(x)$$

$$\cos(x) = 0$$

$$\implies x = \frac{\pi}{2} + k\pi$$

$$\sin(x) = \cos(\frac{\pi}{2} - x)$$

$$\cos(x) = \sin(\frac{\pi}{2} - x)$$



$$\tan(x) = \frac{\sin(x)}{\cos x}$$

odd function:

$$\tan(-x) = -\tan(x)$$

$$\tan(x) = 0$$

$$\implies x = k\pi \quad (k \in \mathbb{Z})$$

vertical asymptote in

$$\frac{\pi}{2} + k\pi, \quad k \in \mathbb{Z}$$

# Trigonometric identities

Know at least the red ones by heart!

$$\sin^2(x) + \cos^2(x) = 1$$

$$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$$

$$\sin(2x) = 2\sin(x)\cos(x)$$

$$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$$

$$\cos(2x) = \cos^2(x) - \sin^2(x)$$


$$= 2\cos^2(x) - 1$$

$$= 1 - 2\sin^2(x)$$

Furthermore, we can use the last two lines to rewrite  $\sin^2(x)$  and  $\cos^2(x)$

$$\sin^2(x) = \frac{1}{2} - \frac{1}{2}\cos(2x) \quad \cos^2(x) = \frac{1}{2} + \frac{1}{2}\cos(2x)$$

# Angles on the unit circle



Do we need to remember all of those?!?  
No, just those in the 1st quadrant:

$x$	0	$\frac{1}{6}\pi$	$\frac{1}{4}\pi$	$\frac{1}{3}\pi$	$\frac{1}{2}\pi$
$\sin(x)$	$\frac{1}{2}\sqrt{0}$	$\frac{1}{2}\sqrt{1}$	$\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}\sqrt{4}$
$\cos(x)$	$\frac{1}{2}\sqrt{4}$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{1}$	$\frac{1}{2}\sqrt{0}$
$\tan(x)$	0	$\frac{1}{3}\sqrt{3}$	1	$\sqrt{3}$	$ng$

$$\begin{aligned}\sin(\pi - x) &= \sin(\pi + (-x)) \\ &= \sin(\pi)\cos(-x) + \cos(\pi)\sin(-x) \\ &= 0 \cdot \cos(-x) + -1 \cdot \sin(-x) \\ &= \sin(x)\end{aligned}$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [\frac{\pi}{2}, 2\pi]$ (I)

Use the formulas (that you know by heart)

- ▶  $\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$
- ▶  $\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$

one can relate angles in Q2-4 to one in Q1!

Very practical: use  $\pi$  and  $2\pi$ !

$\sin(x) = \sin(\pi - x)$	$\cos(x) = -\cos(\pi - x)$	$\tan(x) = -\tan(\pi - x)$
$\sin(x) = -\sin(x - \pi)$	$\cos(x) = -\cos(x - \pi)$	$\tan(x) = \tan(x - \pi)$
$\sin(x) = -\sin(-x)$	$\cos(x) = \cos(-x)$	$\tan(x) = -\tan(-x)$

Ex: Calculate  $\sin(x)$  for  $x = \frac{5}{6}\pi$ .

$\frac{5}{6}\pi$  is in the 2nd quadrant and

$\pi - \frac{5}{6}\pi$  in the 1st:  $\sin(\frac{5}{6}\pi) = \sin(\pi - \frac{5}{6}\pi) = \sin(\frac{1}{6}\pi) = \frac{1}{2}$ .


# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [\frac{\pi}{2}, 2\pi]$ (II)

Who can (and wants to) remember all 9 relations?

Instead, remember to change the argument in the trigonometric function according to:

- ▶ if  $x$  is in 2nd quadrant  $[\frac{1}{2}\pi, \pi]$ :  $x \rightarrow \pi - x$
- ▶ if  $x$  is in 3rd quadrant  $[\pi, \frac{3}{2}\pi]$ :  $x \rightarrow x - \pi$
- ▶ if  $x$  is in 4th quadrant  $[\frac{3}{2}\pi, 2\pi]$ :  $x \rightarrow 2\pi - x$

and the **CAST** rule to check for a **changing sign**.



Ex: Calculate  $\sin(x)$  for  $x = \frac{7}{6}\pi$ .

$\frac{7}{6}\pi$  is in the 3rd quadrant.

$$\frac{7}{6}\pi \rightarrow \frac{7}{6}\pi - \pi = \frac{1}{6}\pi$$

$\sin(x)$  changes sign from Q3 to Q1

$$\text{so that } \sin\left(\frac{7}{6}\pi\right) = -\sin\left(\frac{7}{6}\pi - \pi\right) = -\sin\left(\frac{1}{6}\pi\right) = -\frac{1}{2}.$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \sin(x - \pi)$$

$$\cos\left(\frac{7}{6}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \sin(x - \pi)$$

$$\cos\left(\frac{7}{6}\pi\right) = -\cos\left(\frac{1}{6}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \sin(x - \pi)$$

$$\cos\left(\frac{7}{6}\pi\right) = -\cos\left(\frac{1}{6}\pi\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \cos(x - \pi)$$

$$\tan\left(\frac{11}{6}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$

$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \sin(x - \pi)$$

$$\cos\left(\frac{7}{6}\pi\right) = -\cos\left(\frac{1}{6}\pi\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \cos(x - \pi)$$

$$\tan\left(\frac{11}{6}\pi\right) = -\tan\left(\frac{1}{6}\pi\right) =$$

# Determining $\sin(x)$ , $\cos(x)$ , $\tan(x)$ , $x \in [0, 2\pi]$ (III)

$$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$\cos\left(\frac{2}{3}\pi\right) = -\cos\left(\frac{1}{3}\pi\right) = -\frac{1}{2} \quad \text{qu 2, } \cos(\pi - x)$$

$$\tan\left(-\frac{\pi}{4}\right) = -\tan\left(\frac{\pi}{4}\right) = -1 \quad \text{qu 1, tan is odd}$$


$$\sin\left(\frac{4\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \sin(x - \pi)$$

$$\cos\left(\frac{7}{6}\pi\right) = -\cos\left(\frac{1}{6}\pi\right) = -\frac{1}{2}\sqrt{3} \quad \text{qu 3, } \cos(x - \pi)$$

$$\tan\left(\frac{11}{6}\pi\right) = -\tan\left(\frac{1}{6}\pi\right) = -\frac{1}{3}\sqrt{3} \quad \text{qu 4, } \tan(x - 2\pi)$$

Do [A P.7 T5]


# Sine and Cosine Rules



Sine rule

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

# Sine and Cosine Rules



Sine rule

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

Cosine rule

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = c^2 + a^2 - 2ca \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

Do [A P.7 T31,37].

# Trigonometric Equations and Inequalities

$$\sin(x) = \sin(a)$$

# Trigonometric Equations and Inequalities

$$\begin{aligned}\sin(x) = \sin(a) &\implies x = a + 2k\pi \text{ or } x = \pi - a + 2k\pi \\ \cos(x) = \cos(a)\end{aligned}$$

# Trigonometric Equations and Inequalities

$$\begin{aligned}\sin(x) = \sin(a) &\implies x = a + 2k\pi \text{ or } x = \pi - a + 2k\pi \\ \cos(x) = \cos(a) &\implies x = \pm a + 2k\pi \\ \tan(x) = \tan(a) &\end{aligned}$$

# Trigonometric Equations and Inequalities

$$\sin(x) = \sin(a) \implies x = a + 2k\pi \text{ or } x = \pi - a + 2k\pi$$

$$\cos(x) = \cos(a) \implies x = \pm a + 2k\pi$$

$$\tan(x) = \tan(a) \implies x = a + k\pi$$

$$\sin(x) = \cos(a)$$

# Trigonometric Equations and Inequalities

$$\sin(x) = \sin(a) \implies x = a + 2k\pi \text{ or } x = \pi - a + 2k\pi$$

$$\cos(x) = \cos(a) \implies x = \pm a + 2k\pi$$

$$\tan(x) = \tan(a) \implies x = a + k\pi$$

$\sin(x) = \cos(a)$   $\implies$  rewrite cos to sin :

$$\cos(a) \underset{\sin(x+y)=...}{=} \sin\left(\frac{\pi}{2} - a\right)$$

# Recipe: Trigonometric Equations/Inequalities

- 1 Use **substitution** to rewrite to an equation with (powers of) only 1 unknown, eg  $\sin(x)$  and with only 1 argument (so no mix of  $\sin(x)$  and  $\sin(2x)$ ) or products of different functions (avoid roots as much as possible).
- 2 Maybe use substitution, eg  $y = \sin(x)$ , and solve the equation / inequality
- 3 Rewrite  $y$  back to eg  $\sin(x)$ , and solve, eg  
 $\sin(x) = \frac{1}{2} \implies x = \frac{\pi}{6}$  or  $x = \frac{5\pi}{6}$
- 4 Pay attention to “ $+k 2\pi$ ” ( $\sin$  and  $\cos$ ) or “ $+k \pi$ ” for  $\tan$

# Exercises

- 1 For  $\sin(x) = -\frac{5}{13}$  and  $x \in [3/2\pi, 2\pi]$ , determine  $\cos(x)$  and  $\tan(x)$ .
- 2 For  $\tan(x) = -\frac{7}{24}$ , and  $x$  in the 4th quadrant, determine  $\sin(x)$  and  $\cos(x)$ .
- 3 Calculate  $\sin(\frac{5\pi}{12})$ . Hint: make use of the  $\sin(x + y)$  identity.
- 4 Solve  $3\cos^2(x) - \sin^2(x) = 0$  for  $x \in (0, 2\pi)$ .
- 5 Solve  $\cos^2(x) + \cos(x) = \sin^2(x)$ .

# Exercises

- 1 For  $\sin(x) = -\frac{5}{13}$  and  $x \in [3/2\pi, 2\pi]$ , determine  $\cos(x)$  and  $\tan(x)$ .  
 $\cos(x) = \frac{12}{13}$  and  $\tan(x) = -\frac{5}{12}$
- 2 For  $\tan(x) = -\frac{7}{24}$ , and  $x$  in the 4th quadrant, determine  $\sin(x)$  and  $\cos(x)$ .  
 $\cos(x) = \frac{24}{25}$  and  $\sin(x) = -\frac{7}{25}$
- 3 Calculate  $\sin(\frac{5\pi}{12})$ . Hint: make use of the  $\sin(x + y)$  identity.  
 $\frac{1}{4}(\sqrt{6} + \sqrt{2})$
- 4 Solve  $3\cos^2(x) - \sin^2(x) = 0$  for  $x \in (0, 2\pi)$ .  
 $x = \frac{\pi}{3} \vee x = \frac{4\pi}{3} \vee x = \frac{2\pi}{3} \vee x = \frac{5\pi}{3}$
- 5 Solve  $\cos^2(x) + \cos(x) = \sin^2(x)$ .  
 $x = \pi + 2n\pi \vee x = \frac{\pi}{3} + 2n\pi \vee x = \frac{5\pi}{3} + 2n\pi$  with  $n \in \mathbb{Z}$

# Powers and roots

Reminder: Rules for calculating powers and roots

$$a^p a^q = a^{p+q} \quad (ab)^p = a^p b^p \quad a^{-p} = \frac{1}{a^p}$$

$$a^{1/2} = \sqrt{a} \quad a^{1/n} = \sqrt[n]{a} \quad a^{\frac{p}{q}} = \sqrt[q]{a^p}$$

$$(a^p)^q = a^{pq} \quad a^{-\frac{p}{q}} = \frac{1}{a^{\frac{p}{q}}} = \frac{1}{\sqrt[q]{a^p}} \quad \left(\frac{a}{b}\right)^p = \frac{a^p}{b^p}$$

Useful to rewrite expressions into more simplified forms:

Ex: 
$$\frac{(3a^{-\frac{2}{3}}b)^2}{2a^2b^{-\frac{1}{3}}} = \frac{3^2(a^{-\frac{2}{3}})^2b^2}{2a^2b^{-\frac{1}{3}}} = \frac{9a^{-\frac{4}{3}}b^2}{2a^2b^{-\frac{1}{3}}} = \frac{9}{2}a^{-\frac{10}{3}}b^{\frac{7}{3}}$$

# Equations with powers

Rewrite to

$$a^{\text{expression}_1} = a^{\text{expression}_2}$$

conclude that the exponents must be equal  
because:

# Equations with powers

Rewrite to


$$a^{\text{expression}_1} = a^{\text{expression}_2}$$

conclude that the exponents must be equal because:

- ▶  $f(x) = a^x$  everywhere increasing function if  $a > 1$
- ▶  $f(x) = a^x$  everywhere decreasing function if  $0 < a < 1$

Ex:

$$2^{x+1} + 2^{x+3} = 320$$



# Equations with powers

Rewrite to

$$a^{\text{expression}_1} = a^{\text{expression}_2}$$


conclude that the exponents must be equal because:

- ▶  $f(x) = a^x$  everywhere increasing function if  $a > 1$
- ▶  $f(x) = a^x$  everywhere decreasing function if  $0 < a < 1$

Ex:

$$2^{x+1} + 2^{x+3} = 320 \implies 2^x(2 + 2^3) = 320 \implies 2^x = 32 = 2^5 \implies x = 5$$

Ex:  $16^{3x+3} = 8^{x^2+4}$



# Equations with powers

Rewrite to

$$a^{\text{expression}_1} = a^{\text{expression}_2}$$


conclude that the exponents must be equal because:

- ▶  $f(x) = a^x$  everywhere increasing function if  $a > 1$
- ▶  $f(x) = a^x$  everywhere decreasing function if  $0 < a < 1$

Ex:

$$2^{x+1} + 2^{x+3} = 320 \implies 2^x(2 + 2^3) = 320 \implies 2^x = 32 = 2^5 \implies x = 5$$

Ex:  $16^{3x+3} = 8^{x^2+4} \implies (2^4)^{3x+3} = (2^3)^{x^2+4} \implies 2^{12x+12} = 2^{3x^2+12}$   
 $\implies 12x + 12 = 3x^2 + 12 \implies 3x(x - 4) = 0 \implies x = 0 \text{ or } x = 4$



# Inequalities with powers

Again: get the same base number, then compare the powers

Remember:

- ▶  $f(x) = a^x$  is increasing everywhere if  $a > 1$
- ▶  $f(x) = a^x$  is decreasing everywhere if  $0 < a < 1$

Ex: the base number: is it  $< 1$  or  $> 1$ ?

- ▶  $f(x) = 2^x$  is increasing everywhere because  $2 > 1$
- ▶  $f(x) = 2^{-x}$

# Inequalities with powers

Again: get the same base number, then compare the powers

Remember:

- ▶  $f(x) = a^x$  is increasing everywhere if  $a > 1$
- ▶  $f(x) = a^x$  is decreasing everywhere if  $0 < a < 1$

Ex: the base number: is it  $< 1$  or  $> 1$ ?

- ▶  $f(x) = 2^x$  is increasing everywhere because  $2 > 1$
- ▶  $f(x) = 2^{-x} = (2^{-1})^x = \left(\frac{1}{2}\right)^x$  is decreasing everywhere because  $0 < 1/2 < 1$

Ex: Inequalities

- ▶  $2^x > 2^4 \implies x > 4$  because  $2^x$  is increasing
- ▶  $\left(\frac{1}{2}\right)^x > \left(\frac{1}{2}\right)^4 \implies x < 4$  because  $\left(\frac{1}{2}\right)^x$  is decreasing

# Quiz: Powers

1 Solve  $e^{x^2} = 5e^{2x-1}$ .

2 Solve  $\frac{2^{2x}-8}{2^x-4} \leq 0$

3  $f(x) = 5^x$  is

- (a) increasing everywhere
- (b) decreasing everywhere
- (c) increasing for  $x > 0$ , decreasing for  $x < 0$
- (d) increasing for  $x < 0$ , decreasing for  $x > 0$

4 If  $x < 5$ , then

- (a)  $5^5 > 5^x$
- (b)  $5^x \leq 5^5$
- (c)  $5^5 < 5^x$
- (d)  $5^x \geq 5^5$

# Quiz: Powers

- 1 Solve  $e^{x^2} = 5e^{2x-1}$ .

$$x = 1 + \pm\sqrt{\ln(5)}$$

- 2 Solve  $\frac{2^{2x}-8}{2^x-4} \leq 0$

$$\frac{3}{2} \leq x < 2$$

- 3  $f(x) = 5^x$  is

- (a) increasing everywhere
- (b) decreasing everywhere
- (c) increasing for  $x > 0$ , decreasing for  $x < 0$
- (d) increasing for  $x < 0$ , decreasing for  $x > 0$

- 4 If  $x < 5$ , then

- (a)  $5^5 > 5^x$
- (b)  $5^x \leq 5^5$
- (c)  $5^5 < 5^x$
- (d)  $5^x \geq 5^5$

# Google Search


The screenshot shows a Microsoft Edge browser window with the title bar "Data Science Bachelor -". The address bar contains the URL "https://www.google.nl/search?q=Data+Science+Bachelor&ie=utf-8&oe=utf-8&qaws\_rd=cr\_ssl&ei=iEnnWM39GYm2aZKomL". The main content area displays two search results:

- Data Science Bachelor - ie.edu**  
[Ad] [www.ie.edu/data-science](http://www.ie.edu/data-science) ▾  
Join the Digital Revolution with a Degree in Data Science. Learn More! · Multiple Career Options · Innovative Methodology · Personalized Study Path · Courses: Software Development, Digital & Mobile Business, Databases & Data Modeling, IT Outsourc... · Programs Offered by IEU · Contact Us · 10 Reasons to Choose IE · Insight Sessions & Events
- Hackathon - Join the Data Science Game - datasciencegame.com**  
[Ad] [www.datasciencegame.com/](http://www.datasciencegame.com/) ▾  
Take part in the world's biggest competition in Data Science for students! · Worldwide challenge · Finals In Paris · More than 400 students · Real-world problem · Highlights: Improve Your Skills, Represent Your University And Push Your Limits... · Registration · Sponsors · Press · Previous edition

## Data Science - Technische Universiteit Eindhoven

<https://www.tue.nl> › Home › Education › TU/e Bachelor ... › Undergraduate ... ▾  
Data Science is analyzing and interpreting large amounts of data in order to retrieve ... Data Science is a joint bachelor of Tilburg University and Eindhoven ...

# Mathematical Model for Weblinks




Model the WWW as a **graph**:

- ▶ Squares (nodes, vertices) denote web pages
- ▶ Arrows (edges) denote links from one page to another

Relevance is translated into number of **incoming** links weighted by the importance of referring pages.


(What comes first?)

# System of Linear Equations



$$\begin{aligned}x_1 &= 1 \cdot x_3 + \frac{1}{2} \cdot x_4 \\x_2 &= \frac{1}{3} \cdot x_1 \\x_3 &= \frac{1}{3} \cdot x_1 + \frac{1}{2} \cdot x_2 + \frac{1}{2} \cdot x_4 \\x_4 &= \frac{1}{3} \cdot x_1 + \frac{1}{2} \cdot x_2\end{aligned}$$

# System of Linear Equations - Matrix and Vectors



$$\begin{bmatrix} 0 & 0 & 1 & \frac{1}{2} \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix} = A$$

$$A \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \Rightarrow \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \frac{1}{31} \cdot \begin{bmatrix} 12 \\ 4 \\ 9 \\ 6 \end{bmatrix}$$

Information about page ranks is determined as a **vector** of page relevance (solutions of linear systems).

# More Matrix-Vector problems

Alice	4			4
Bob		5	4	
Joe		5	4	
Sam	5			5



- ▶ User preferences/histories are encoded as vector data.
- ▶ How similar is your vector to those of others?
- ▶ completely similar = “parallel”
- ▶ completely different = “perpendicular”
- ▶ Trigonometric functions (just done) used with vector calculus (coming up)

# Introduction: Vectors in Analytic Geometry


We will introduce vectors and their calculus in a specific application:  
Analytic Geometry

- ▶ describe directions in two or three space dimensions
- ▶ addition and scaling of vectors
- ▶ linear combinations
- ▶ lines and line segments in  $\mathbb{R}^2$  and  $\mathbb{R}^3$
- ▶ Planes in  $\mathbb{R}^3$
- ▶ Orthogonality, angle between vectors
- ▶ Intersections of lines, lines and planes, planes

# Vectors in 2 and 3 dimensions

A vector is represented in plane or space by an arrow with **direction** and **length** (magnitude).

Vector  $\begin{bmatrix} -2 \\ 5 \end{bmatrix}$  (2 to the left, 5 up)



These vectors are equivalent, even though they have different starting points.

# Vectors in 2 and 3 dimensions: Notation

In a plane (2D): 2 to the left and 5 up is a vector:

- ▶ Notation: Write  $\begin{pmatrix} -2 \\ 5 \end{pmatrix}$  or  $\begin{bmatrix} -2 \\ 5 \end{bmatrix}$  (lectures: latter notation)
- ▶ Sometimes “also”:  $\langle -2, 5 \rangle$ : eg in books to save space...
- ▶  $\mathbf{v}$ ,  $\underline{\mathbf{v}}$ ,  $\vec{\mathbf{v}}$ , or simply  $v$  (on the slides  $\mathbf{v}$ , on paper, I write  $v$  of  $\underline{\mathbf{v}}$ )
- ▶ Sketch: Arrow (a combination of a line segment and directions)



A vector has a direction and length (here  $\sqrt{(-2)^2 + 5^2}$ ).

# Vectors: Addition and subtraction

- ▶ add or subtract component-wise:

$$\begin{bmatrix} 8 \\ 13 \end{bmatrix} + \begin{bmatrix} 26 \\ 7 \end{bmatrix} = \begin{bmatrix} 34 \\ 20 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix}$$

- ▶ graphically: parallelogram



## Vectors from point $A$ to point $B$

Because  $A = (x_1, y_1)$  and  $B = (x_2, y_2)$  then the vector from  $A$  to  $B$  is:



$$B - A = \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} - \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}$$

Notation:  $\overrightarrow{AB}$  or  $\vec{AB}$ .

Do [V 5 T1]


# Scaling of vectors

Scaling (scalar multiplication) changes the length and evt the direction:



- ▶ scalar means using (real) number
- ▶ multiply component-wise:  $2 \cdot \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$

# Scaling of vectors



- ▶ multiplication by 2:  
vector becomes 2 times as long, keeps the same direction
- ▶ multiplication by  $-2$ :  
vector becomes 2 times as long, in opposite direction

(Later for more about length)

# Subtraction of vectors

Convention: Subtraction is addition of the vector with opposite direction:

$$\mathbf{b} - \mathbf{a} = \mathbf{b} + ((-1) \cdot \mathbf{a})$$

Ex:  $\begin{bmatrix} 1 \\ -2 \end{bmatrix} - \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \end{bmatrix} + (-1) \cdot \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \end{bmatrix} + \begin{bmatrix} 3 \\ -1 \end{bmatrix}$

# Length of a vector

Notation:

- ▶  $\|\mathbf{x}\|$  (chosen)
- ▶ or sometimes also  $|\mathbf{x}|$

Calculate: if  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  then

$$\|\mathbf{x}\| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} \quad \text{thus} \quad \|\mathbf{x}\|^2 = x_1^2 + x_2^2 + \cdots + x_n^2$$

Distance between two points  $\mathbf{x}$  and  $\mathbf{y}$ :  $\|\mathbf{x} - \mathbf{y}\|$

Ex: Distance between  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 2 \\ 3 \end{bmatrix} = \sqrt{5}$

is also the length of the difference vector  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  (and  $\begin{bmatrix} -1 \\ -2 \end{bmatrix}$ )

# Length of a vector

Ex:  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  has length

## Length of a vector

Ex:  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  has length  $\sqrt{2}$

Sometimes we seek a vector in the same direction, but with length 1

$$\text{Ex: } \begin{bmatrix} 1 \\ 1 \end{bmatrix} / \sqrt{2} = \begin{bmatrix} \frac{1}{2}\sqrt{2} \\ \frac{1}{2}\sqrt{2} \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} = \sqrt{2} \cdot \underbrace{\begin{bmatrix} \frac{1}{2}\sqrt{2} \\ \frac{1}{2}\sqrt{2} \end{bmatrix}}_{\text{length 1}}$$

# Angle between vectors

Let  $\mathbf{a} = (a_1, a_2)$  and  $\mathbf{b} = (b_1, b_2)$  be vectors in  $\mathbb{R}^2$

Inner product in 2D:  $\mathbf{a} \cdot \mathbf{b} = (a_1, a_2) \cdot (b_1, b_2) = a_1 b_1 + a_2 b_2$

Inner product  $n$ D (so also for  $n = 3$ ):

$$\mathbf{a} \cdot \mathbf{b} = (a_1, \dots, a_n) \cdot (b_1, \dots, b_n) = a_1 b_1 + \cdots + a_n b_n$$

Useful for:

► length:  $\|\mathbf{a}\| = \sqrt{\mathbf{a} \cdot \mathbf{a}} = \sqrt{a_1^2 + \cdots + a_n^2}$

► angle:  $\cos(\mathbf{a}, \mathbf{b}) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|}$   $\Leftarrow$   $\underbrace{\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| \cdot |\mathbf{b}| \cdot \cos(\mathbf{a}, \mathbf{b})}_{\text{inner product rule}}$

Thus also  $\mathbf{a} \cdot \mathbf{b} = \cos(\alpha) \|\mathbf{a}\| \|\mathbf{b}\|$ ,  $\alpha = \angle(\mathbf{a}, \mathbf{b})$

Different notation for inner product:  $\langle \mathbf{a}, \mathbf{b} \rangle$  and  $\mathbf{a}^T \mathbf{b}$

# Angle between vectors

Ex: Calculate the angle  $\theta$  between  $\underbrace{\begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix}}_{\mathbf{v}}$  and  $\underbrace{\begin{bmatrix} -1 \\ \sqrt{3} \end{bmatrix}}_{\mathbf{w}}$ : Inner product rule:

$$\begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix} \cdot \begin{bmatrix} -1 \\ \sqrt{3} \end{bmatrix} = \left| \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix} \right| \cdot \left| \begin{bmatrix} -1 \\ \sqrt{3} \end{bmatrix} \right| \cdot \cos \theta$$

or (length of both vectors is  $\sqrt{(\pm 1)^2 + (\sqrt{3})^2} = \sqrt{1+3} = 2$ ):

$$\underbrace{1 \cdot -1 + \sqrt{3} \cdot \sqrt{3}}_{\mathbf{v} \cdot \mathbf{w}} = \underbrace{2}_{|\mathbf{v}|} \cdot \underbrace{2}_{|\mathbf{w}|} \cdot \cos \theta$$

and  $2 = 4 \cdot \cos \theta \implies \cos \theta = \frac{1}{2} \implies \theta = \pm \pi/3$  (solution with minus sign goes away)

(the angle between two vectors is maximal  $\pi$ !)

# Distances, angles and the inner product

- $\mathbf{x} \cdot \mathbf{x} \geq 0$

logical, is same as  $\|\mathbf{x}\|^2$ , thus  $\geq 0$

- $\mathbf{x} \cdot \mathbf{x} = 0$  only if  $\mathbf{x} = \mathbf{0}$

only the **0**-vector has length 0

- $\mathbf{x} \cdot \mathbf{y} = \mathbf{y} \cdot \mathbf{x}$

just as for the multiplication of numbers, the order does not matter

- $(\lambda \mathbf{x}) \cdot \mathbf{y} = \lambda (\mathbf{x} \cdot \mathbf{y})$

if  $\mathbf{x}$  eg 2 times as long, then the inner product is 2 times as big

logical, since  $\mathbf{x} \cdot \mathbf{y} = \|\mathbf{x}\| \|\mathbf{y}\| \cos(\alpha)$

# Orthogonal vectors

Vectors  $\mathbf{a}$  and  $\mathbf{b}$  are called **orthogonal** (or perpendicular) if  $\mathbf{a} \cdot \mathbf{b} = 0$

Notation:  $\mathbf{a} \perp \mathbf{b}$

Works with:  $\cos(\mathbf{a}, \mathbf{b}) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|}$

because then the numerator is 0, so  $\cos(\mathbf{a}, \mathbf{b}) = 0$ , thus  $\angle(\mathbf{a}, \mathbf{b}) = \frac{\pi}{2}$

Ex:  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$  are indeed orthogonal

but also  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} -2 \\ 1 \end{bmatrix}$

# Orthogonal vectors

In  $\mathbb{R}^2$  the vectors that are orthogonal to a given vector can be calculated

Ex: Which vectors are  $\perp (1, 2)$  ?

We need to have:  $(a_1, a_2) \cdot (1, 2) = a_1 + 2a_2 = 0$

# Orthogonal vectors

In  $\mathbb{R}^2$  the vectors that are orthogonal to a given vector can be calculated

**Ex:** Which vectors are  $\perp (1, 2)$  ?

We need to have:  $(a_1, a_2) \cdot (1, 2) = a_1 + 2a_2 = 0$

Useful trick:

switch the components, and give one an extra minus sign, thus:

- ▶  $(2, -1)$ , because  $(2, -1) \cdot (1, 2) = 0$

# Orthogonal vectors

In  $\mathbb{R}^2$  the vectors that are orthogonal to a given vector can be calculated

**Ex:** Which vectors are  $\perp (1, 2)$ ?

We need to have:  $(a_1, a_2) \cdot (1, 2) = a_1 + 2a_2 = 0$

Useful trick:

switch the components, and give one an extra minus sign, thus:

- ▶  $(2, -1)$ , because  $(2, -1) \cdot (1, 2) = 0$
- ▶  $(-2, 1)$ , because  $(-2, 1) \cdot (1, 2) = 0$

# Orthogonal vectors

In  $\mathbb{R}^2$  the vectors that are orthogonal to a given vector can be calculated

**Ex:** Which vectors are  $\perp (1, 2)$ ?

We need to have:  $(a_1, a_2) \cdot (1, 2) = a_1 + 2a_2 = 0$

Useful trick:

switch the components, and give one an extra minus sign, thus:

- ▶  $(2, -1)$ , because  $(2, -1) \cdot (1, 2) = 0$
- ▶  $(-2, 1)$ , because  $(-2, 1) \cdot (1, 2) = 0$
- ▶ and also all multiples of  $(-2, 1)$  and  $(2, -1)$ :  
 $(-2\lambda, \lambda) = \lambda(-2, 1)$  for  $\lambda \in \mathbb{R}$

(In  $\mathbb{R}^2$  a line of vectors is perpendicular on a given vector!)

## Quiz: Vector calculus

- 1 Given vectors  $\mathbf{v} = \langle -2, 3 \rangle$  and  $\mathbf{u} = \langle 4, 6 \rangle$ , find  
 $\mathbf{v} + 2\mathbf{u}$  and  $\mathbf{u} - 4\mathbf{u}$
- 2 Given vectors  $\mathbf{v} = \langle 1, -2 \rangle$  and  $\mathbf{u} = \langle u_1, u_2 \rangle$ , find components of vector  $\mathbf{u}$  so that  $\mathbf{v} + 3\mathbf{u} = \mathbf{0}$
- 3 Given vectors  $\mathbf{v} = \langle 4, 1 \rangle$  and  $\mathbf{u} = \langle u_1, u_2 \rangle$ , find components of vector  $\mathbf{u}$  so that  $2\mathbf{v} - 3\mathbf{u} = \mathbf{0}$
- 4 Calculate the distance between the points  $A(3, 1, 7)$  and  $B(5, -1, 3)$
- 5 Calculate the angle between the vectors  $\mathbf{v} = \langle 1, 1, 1 \rangle$  and  $\mathbf{w} = \langle 4, 0, -3 \rangle$

# Quiz: Vector calculus

- 1 Given vectors  $\mathbf{v} = \langle -2, 3 \rangle$  and  $\mathbf{u} = \langle 4, 6 \rangle$ , find  
 $\mathbf{v} + 2\mathbf{u}$  and  $\mathbf{u} - 4\mathbf{u}$   
 $\mathbf{v} + 2\mathbf{u} = \langle 6, 15 \rangle$  and  $\mathbf{u} - 4\mathbf{u} = \langle -12, -18 \rangle$
- 2 Given vectors  $\mathbf{v} = \langle 1, -2 \rangle$  and  $\mathbf{u} = \langle u_1, u_2 \rangle$ , find components of vector  $\mathbf{u}$  so that  $\mathbf{v} + 3\mathbf{u} = \mathbf{0}$   
 $\mathbf{u} = \langle -\frac{1}{3}, \frac{2}{3} \rangle$
- 3 Given vectors  $\mathbf{v} = \langle 4, 1 \rangle$  and  $\mathbf{u} = \langle u_1, u_2 \rangle$ , find components of vector  $\mathbf{u}$  so that  $2\mathbf{v} - 3\mathbf{u} = \mathbf{0}$   
 $\mathbf{u} = \langle \frac{8}{3}, \frac{2}{3} \rangle$
- 4 Calculate the distance between the points  $A(3, 1, 7)$  and  $B(5, -1, 3)$   
 $d = 2\sqrt{5}$
- 5 Calculate the angle between the vectors  $\mathbf{v} = \langle 1, 1, 1 \rangle$  and  $\mathbf{w} = \langle 4, 0, -3 \rangle$   
 $\cos(\phi) = \frac{1}{5\sqrt{2}}$

# Linear combination of vectors

In linear algebra, we say for vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  and real numbers  $a_1, \dots, a_n$  that the sum (-vector)  $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n$  is a **linear combination**.

Let's get to work with that.

# Linear functions and combinations

A function is called linear if:

- ▶ in 1 variable:  $f(x) = ax$  (or sometimes also with “ $+b$ ”)
- ▶ in 2 variables:  $f(x, y) = ax + by$
- ▶ in  $n$  variables:  $f(x_1, \dots, x_n) = a_1x_1 + a_2x_2 + \dots + a_nx_n$
- ▶ in  $n$  vectors:  $f(\mathbf{v}_1, \dots, \mathbf{v}_n) = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n$

Thus linear is always:

# Linear functions and combinations

A function is called linear if:

- ▶ in 1 variable:  $f(x) = ax$  (or sometimes also with “ $+b$ ”)
- ▶ in 2 variables:  $f(x, y) = ax + by$
- ▶ in  $n$  variables:  $f(x_1, \dots, x_n) = a_1x_1 + a_2x_2 + \dots + a_nx_n$
- ▶ in  $n$  vectors:  $f(\mathbf{v}_1, \dots, \mathbf{v}_n) = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n$

Thus linear is always: sum of constants times variables

What is not-linear?

# Linear functions and combinations

A function is called linear if:

- ▶ in 1 variable:  $f(x) = ax$  (or sometimes also with “ $+b$ ”)
- ▶ in 2 variables:  $f(x, y) = ax + by$
- ▶ in  $n$  variables:  $f(x_1, \dots, x_n) = a_1x_1 + a_2x_2 + \dots + a_nx_n$
- ▶ in  $n$  vectors:  $f(\mathbf{v}_1, \dots, \mathbf{v}_n) = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n$

Thus linear is always: sum of constants times variables

What is not-linear? Everything else, eq:

- ▶ everything containing multiplications of variables, eg  $x^2$ ,  $xy$ , etc
- ▶  $\sin$ ,  $\cos$ ,  $\tan$ ,  $e^x$ ,  $\ln(x)$ , etc etc

Read examples [V 1.11]

# Lines: An alternative to $ax + by = c$ in $\mathbb{R}^2$ and also $\mathbb{R}^3$

We have seen: Every line in  $\mathbb{R}^2$  is described by:

All  $(x, y)$  for which:  $ax + by = c$

Different  $a, b, c$  represent **possibly** different lines.

But it must not:  $2x - 3y = 1$  is the same line as  $3y - 2x = -1$   
(multiply left/right with  $-1$ ).

In  $\mathbb{R}^3$

$$ax + by + cz = d$$

does not represent a line.

# Lines: An alternative to $ax + by = c$ in $\mathbb{R}^2$ and also $\mathbb{R}^3$

Ex: What does  $ax + by + cz = d$  describe in  $\mathbb{R}^3$ ? Let's take an example:

$$0 \cdot x + 0 \cdot y + 1 \cdot z = 0$$

( $a = b = d = 0$  and  $c = 1$ ) – or  $z = 0$ . This is the  $xy$ -plane!: Table:

x	y	z	$0 \cdot x + 0 \cdot y + z$
0	0	0	$0 + 0 + 0 = 0$
1	0	0	$0 \cdot 1 + 0 + 0 = 0$
0	1	0	$0 + 0 \cdot 1 + 0 = 0$
1	1	0	$0 \cdot 1 + 0 \cdot 1 + 0 = 0$
:	:	0	$0 = 0$

The 4 points  $(0, 0, 0)$ ,  $(0, 1, 0)$ ,  $(1, 0, 0)$  and  $(1, 1, 0)$  are the corners of a square in the  $xy$ -plane. It cannot be a straight line!

We need: **An expression for a line that works in  $\mathbb{R}^2$ ,  $\mathbb{R}^3$ ,  $\mathbb{R}^4$  ...**

## Parameter representation of a Line

Let  $\mathbf{a} = \begin{bmatrix} x_A \\ y_A \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} x_B \\ y_B \end{bmatrix}$  be two coordinate vectors of the points  $A$  and  $B$ . (the same for  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ ). It holds that

$$\mathbf{x} = \mathbf{x}(\lambda) = \mathbf{a} + \lambda (\mathbf{b} - \mathbf{a}) \text{ for all } \lambda \in \mathbb{R}$$

describes all points on the line through  $A$  and  $B$ . Specifically

- ▶  $\lambda = 0$ :  $\mathbf{x} = \mathbf{a} + 0 \cdot (\mathbf{b} - \mathbf{a}) = \mathbf{a} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \mathbf{a}$
- ▶  $\lambda = 1$ :  $\mathbf{x} = \mathbf{a} + 1 \cdot (\mathbf{b} - \mathbf{a}) = \mathbf{a} + \mathbf{b} - \mathbf{a} = \mathbf{b}$

The representation  $\mathbf{x} = \mathbf{x}(\lambda) = \mathbf{a} + \lambda (\mathbf{b} - \mathbf{a})$  with the parameter  $\lambda$  is called **parameter representation** or **vector representation**. Definitions:

- ▶  $\mathbf{a}$  is the **support vector** or the **start point**
- ▶  $\mathbf{b} - \mathbf{a}$  is the **direction vector**


## Parameter representation of a Line, now in $\mathbb{R}^3$

Ex: Parameter representation of a line through points  $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$ :

$$\begin{aligned}\begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 2-1 \\ 3-1 \\ 1-1 \end{bmatrix} \\ &= \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \lambda \underbrace{\begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}}_{\text{direction vector}} \\ &= \begin{bmatrix} 1+\lambda \\ 1+2\lambda \\ 1+0 \cdot \lambda \end{bmatrix} = \begin{bmatrix} 1+\lambda \\ 1+2\lambda \\ 1 \end{bmatrix}\end{aligned}$$

# Lines, line segments, straight lines

For all  $\lambda \in \mathbb{R}$  is  $\mathbf{x} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$  on the line through  $A$  en  $B$ :



- ▶  $\lambda = 0: \mathbf{x} = \mathbf{a} + 0 \cdot (\mathbf{b} - \mathbf{a}) = \mathbf{a} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \mathbf{a}$  (point A)
- ▶  $\lambda = 1: \mathbf{x} = \mathbf{a} + 1 \cdot (\mathbf{b} - \mathbf{a}) = \mathbf{a} + \mathbf{b} - \mathbf{a} = \mathbf{b}$  (point B)
- ▶  $\lambda = \frac{1}{2}: \mathbf{x} = \mathbf{a} + \frac{1}{2} \cdot (\mathbf{b} - \mathbf{a}) = \frac{1}{2}(\mathbf{a} + \mathbf{b})$  middle between  $\mathbf{a}$  en  $\mathbf{b}$
- ▶  $\lambda = \frac{1}{4}: \mathbf{x} = \frac{3}{4}\mathbf{a} + \frac{1}{4}\mathbf{b}$  middle between M and A.

# Complete lines, half lines and line segments

Parameter representation

$$\mathbf{x} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$$

yields for each  $\lambda \in \mathbb{R}$

one point  $\mathbf{x}$  on the line through points  $B$  and  $A$

Depending on what  $\lambda$  can be, this is a line segment, line etc:

- ▶  $\lambda \in [0, 1]$ : line segment between  $A$  and  $B$  including these points
- ▶  $\lambda \geq 0$ : half line from  $A$  through  $B$  and further
- ▶  $\lambda \in \mathbb{R}$ : (complete) line through points  $A$  and  $B$

Note that  $\mathbf{a} + \lambda(\mathbf{b} - \mathbf{a})$  is a linear combination because

$\mathbf{a} + \lambda(\mathbf{b} - \mathbf{a}) = \mathbf{a} + \lambda\mathbf{b} - \lambda\mathbf{a} = \lambda\mathbf{b} + (1 - \lambda)\mathbf{a}$  is of the form: scalar times vector plus scalar times vector.

# Lines, line segments, straight lines

Ex: A line has several parameter representations:

Parameter representations are not unique!

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} -2 \\ -4 \\ 0 \end{bmatrix} \quad \text{just take } \mu = -\frac{1}{2}\lambda$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} -2 \\ -4 \\ 0 \end{bmatrix} \quad \text{just take } \mu = -\frac{1}{2}\lambda - \frac{1}{2}$$

If your neighbor gets a different answer, it does not mean anything! . . .

But: All direction vectors are “the same” except for a minus sign.

# Lines in $\mathbb{R}^2$ : Function vs. parameter representation

Change from function to parameter representation:

**Ex:**  $y = 2x + 3$  goes through  $(0, 3)$  and  $(1, 5)$ , thus

parameter representation  $\begin{bmatrix} 0 \\ 3 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} \lambda \\ 3 + 2\lambda \end{bmatrix}$

**Ex:** Let a parameter representation be  $\begin{bmatrix} 1 + 2\lambda \\ 2 + 3\lambda \end{bmatrix}$

then the line goes through  $(1, 2)$  (for  $\lambda = 0$ ) and  $(3, 5)$  (for  $\lambda = 1$ )

thus it's the equation  $y = \frac{3}{2}x + \frac{1}{2}$

# Lines in $\mathbb{R}^3$

Can be given as in  $\mathbb{R}^2$  by parametrisation:  
starting point and direction vector


$$\begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} + \lambda \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix}$$

A line can also be given by the intersection of two planes


# Lines

- ▶ Determine a parametric representation of the lines through:
  - ▶  $\begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}$  and  $\begin{bmatrix} 5 \\ -1 \\ 4 \end{bmatrix}$
  - ▶  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$
- ▶ Give an equation for the line:
$$\mathbf{x} = \begin{bmatrix} 1 \\ 3 \end{bmatrix} + \lambda \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$
- ▶ Find a parametric representation for the following lines in  $\mathbb{R}^2$   
$$3x - 4y + 7 = 0$$


# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$



# Vectors in $\mathbb{R}^2$




# Vectors in $\mathbb{R}^2$



start point  $p = (0, 0)$ , to reach  $(3, 2)$

direction  $v_1 = (1, 0) - (0, 0) = (1, 0)$

# Vectors in $\mathbb{R}^2$




start point  $\mathbf{p} = (0, 0)$ , to reach  $(3, 2)$

$$\text{direction } \mathbf{v}_1 = (1, 0) - (0, 0) = (1, 0)$$

$$\text{direction } \mathbf{v}_2 = (0, 1) - (0, 0) = (0, 1)$$

# Vectors in $\mathbb{R}^2$




start point  $p = (0, 0)$ , to reach  $(3, 2)$

$$\text{direction } v_1 = (1, 0) - (0, 0) = (1, 0)$$

$$\text{direction } v_2 = (0, 1) - (0, 0) = (0, 1)$$

$$\text{way to } \binom{3}{2} = \binom{0}{0} + 3 \binom{1}{0} + \dots$$

# Vectors in $\mathbb{R}^2$



start point  $\mathbf{p} = (0, 0)$ , to reach  $(3, 2)$


$$\text{direction } \mathbf{v}_1 = (1, 0) - (0, 0) = (1, 0)$$

$$\text{direction } \mathbf{v}_2 = (0, 1) - (0, 0) = (0, 1)$$

$$\text{way to } \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \dots$$

$$\text{way to } \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

# Vectors in $\mathbb{R}^2$



start point  $\mathbf{p} = (0, 0)$ , to reach  $(3, 2)$

$$\text{direction } \mathbf{v}_1 = (1, 0) - (0, 0) = (1, 0)$$


$$\text{direction } \mathbf{v}_2 = (0, 1) - (0, 0) = (0, 1)$$

$$\text{way to } \binom{3}{2} = \binom{0}{0} + 3 \binom{1}{0} + \dots$$

$$\text{way to } \binom{3}{2} = \binom{0}{0} + 3 \binom{1}{0} + 2 \binom{0}{1}$$

$$\text{way to } \binom{3}{2} = \mathbf{p} + 3\mathbf{v}_1 + 2\mathbf{v}_2$$

# Vectors in $\mathbb{R}^2$



start point  $\mathbf{p} = (0, 0)$ , to reach  $(3, 2)$

$$\text{direction } \mathbf{v}_1 = (1, 0) - (0, 0) = (1, 0)$$

$$\text{direction } \mathbf{v}_2 = (0, 1) - (0, 0) = (0, 1)$$

$$\text{way to } \binom{3}{2} = \binom{0}{0} + 3 \binom{1}{0} + \dots$$


$$\text{way to } \binom{3}{2} = \binom{0}{0} + 3 \binom{1}{0} + 2 \binom{0}{1}$$

$$\text{way to } \binom{3}{2} = \mathbf{p} + 3\mathbf{v}_1 + 2\mathbf{v}_2$$

xy-plane: Each point  $(x, y)$  determined by

$$\text{unique } \lambda, \mu: \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

# Vectors in $\mathbb{R}^2$



start point  $\mathbf{p} = (0, 0)$ , to reach  $(3, 2)$

$$\text{direction } \mathbf{v}_1 = (1, 0) - (0, 0) = (1, 0)$$

$$\text{direction } \mathbf{v}_2 = (0, 1) - (0, 0) = (0, 1)$$

$$\text{way to } \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \dots$$

$$\text{way to } \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 0 \end{pmatrix} + 2 \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\text{way to } \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \mathbf{p} + 3\mathbf{v}_1 + 2\mathbf{v}_2$$

xy-plane: Each point  $(x, y)$  determined by

$$\text{unique } \lambda, \mu: \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\text{xy-plane: } \begin{pmatrix} x \\ y \end{pmatrix} = \mathbf{p} + \lambda\mathbf{v}_1 + \mu\mathbf{v}_2$$

# Why the parameter representation?

The 2D notation  $y = ax + b$  does not work for equations in the 3D space!

**Ex:** What object is  $x - y + z = 1$  in  $\mathbb{R}^3$ ?

Answer: That depends on the set of points  $(x, y, z)$  that satisfies the equation.

Let's try to find a description for that:

- ▶ 1 equation with 3 unknowns. Choose 2 arbitrarily:
- ▶  $z = \lambda$  and  $y = \mu$  and calculate  $x$ :
- ▶  $x = 1 - \lambda + \mu$

So for all  $\lambda, \mu \in \mathbb{R}$  the resulting  $\mathbf{x}$  satisfies  $x - y + z = 1$ :

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 - \lambda + \mu \\ \mu \\ \lambda \end{bmatrix}$$

# Why the parameter representation?

Different notation:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 - \lambda + \mu \\ \mu \\ \lambda \end{bmatrix}$$
$$= \begin{bmatrix} 1 + (-1) \cdot \lambda + 1 \cdot \mu \\ 0 + 0 \cdot \lambda + 1 \cdot \mu \\ 0 + 1 \cdot \lambda + 0 \cdot \mu \end{bmatrix}$$
$$= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

represents all linear combinations of 2 directions from 1 support point.

Such an object is always a plane!

Summarized:

- ▶ in 2D:  $ax + by = c$  is the equation of a line
- ▶ in 3D:  $ax + by = c \iff ax + by + 0 \cdot z = c$  is the equation of a plane!

# Planes in $\mathbb{R}^3$

Planes can be given by a parametrisation:

start point and 2 direction vectors:

$$\mathbf{a} + \lambda(\mathbf{b} - \mathbf{a}) + \mu(\mathbf{c} - \mathbf{a}) \text{ for all } \lambda, \mu \in \mathbb{R}$$

$\lambda$  and  $\mu$  are the parameters

$$\begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} + \lambda \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix} + \mu \begin{bmatrix} x_3 - x_1 \\ y_3 - y_1 \\ z_3 - z_1 \end{bmatrix}$$

Ex: the xy-plane:  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

## The parameter representation of the $xy$ -plane

Addition yields  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu \\ 0 \end{bmatrix}$  that the points  $(\lambda, \mu, 0)$  are described for  $\lambda, \mu \in \mathbb{R}$ .

**Ex:** The equation  $z = 0$  describes the same  $xy$ -plane:

- ▶ 1 equation with 3 unknowns:  $0 \cdot x + 0 \cdot y + z = 0$ . Choose 2 arbitrarily:
- ▶  $x = \lambda$  and  $y = \mu$  and calculate  $z$ :
- ▶  $z = 0$ .

So that for all  $\lambda, \mu \in \mathbb{R}$  the resulting  $\mathbf{x}$  satisfy  $z = 0$ :

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

# The parameter representation is not unique

Take  $V$  the xy-plane:

$$V: \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu \\ 0 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{v}} + \lambda \underbrace{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{v}} + \mu \underbrace{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{w}}$$

All points  $P = (a, b, 0)$  are in  $V$ :

$$V: \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{v}} + a \underbrace{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{v}} + b \underbrace{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{w}}$$

We obtain the point  $P$  by choosing  $\lambda = a$  and  $\mu = b$

# The parameter representation is not unique

Take  $V$  the  $xy$ -plane:

$$V: \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu \\ 0 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{r}} + \lambda \underbrace{\begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}}_{\mathbf{r}} + \mu \underbrace{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{s}}$$

All points  $P = (a, b, 0)$  are in  $V$ :

$$V: \begin{bmatrix} a \\ b \\ 0 \end{bmatrix} = \underbrace{\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}}_{\mathbf{r}} + \frac{1}{2}a \underbrace{\begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}}_{\mathbf{r}} + \left(b + \frac{1}{2}a\right) \underbrace{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{s}}$$

We obtain the point  $P$  by choosing  $\lambda = \frac{1}{2}a$  and  $\mu = b + \frac{1}{2}a$

A plane can be described by many different directions

Here  $\mathbf{r} = 2\mathbf{v} - \mathbf{w}$  and  $\mathbf{s} = \mathbf{w}$

# Orthogonal vectors

In  $\mathbb{R}^3$

# Orthogonal vectors

In  $\mathbb{R}^3$  a plane of vectors is orthogonal to a given vector.

Ex: Which vectors  $v \in \mathbb{R}^3$  are perpendicular on  $\begin{bmatrix} -3 \\ 4 \\ 1 \end{bmatrix}$ ? For such a vector

$$v = \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} \text{ it must hold: } \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} \cdot \begin{bmatrix} -3 \\ 4 \\ 1 \end{bmatrix} = 0 \iff -3v_x + 4v_y + v_z = 0$$

- ▶ 1 equation with 3 unknown. Choose 2 arbitrarily:
- ▶  $v_x = \lambda$  and  $v_y = \mu$  and calculate  $v_z$ :
- ▶  $v_z = 3\lambda - 4\mu$

which is a plane through the origin:

$$\begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix} = \begin{bmatrix} \lambda \\ \mu \\ 3\lambda - 4\mu \end{bmatrix} = \lambda \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ -4 \end{bmatrix}$$

# Orthogonal vectors

Ex: For which numbers  $a \in \mathbb{R}$  is  $\begin{bmatrix} a \\ 4 \\ 1 \end{bmatrix}$  orthogonal to  $\begin{bmatrix} a \\ a \\ 3 \end{bmatrix}$ ?


We must have:

$$\begin{aligned}(a, 4, 1) \cdot (a, a, 3) &= 0 \\ \iff a^2 + 4a + 3 &= 0 \\ \iff (a+1)(a+3) &= 0 \\ \iff a = -1 \vee a = -3\end{aligned}$$

# Planes and normals (see also [A 10.4])

Let  $P_0 = (x_0, y_0, z_0)$  be an arbitrary point in the plane

and  $\mathbf{n} = (n_1, n_2, n_3)$  a normal(vector) on this plane (length does not matter):



normal vector = vector that is orthogonal to the plane

The plane consists of all points  $P = (x, y, z)$  for which it holds:

$(x - x_0, y - y_0, z - z_0)$  is orthogonal to  $\mathbf{n}$

Thus for  $(x, y, z)$  in the plane:

$$0 = \mathbf{n} \cdot (x - x_0, y - y_0, z - z_0) = n_1(x - x_0) + n_2(y - y_0) + n_3(z - z_0)$$

$$\text{So } n_1x + n_2y + n_3z = \underbrace{n_1x_0 + n_2y_0 + n_3z_0}_d$$

Normal vector form of a plane:  $\mathbf{n} \cdot \mathbf{x} = d$

# Planes and normals

Earlier, we saw that the plane  $V$  with equation  $x - y + z = 1$  has the vector representation:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \text{and normal } \mathbf{n} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

The normal  $\mathbf{n}$  must be orthogonal to both directions  $V$ !

Check:

$$\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} = 1 \cdot -1 + -1 \cdot 0 + 1 \cdot 1 = 0 \text{ works}$$

$$\begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = 1 \cdot 1 + -1 \cdot 1 + 1 \cdot 0 = 0 \text{ works too}$$

## Planes and normals

The plane  $V$  through point  $P = (1, 2, 3)$  with normal  $\mathbf{n} = \begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$ ?

has the equation  $n_1x + n_2y + n_3z = d$

Thus  $-x + 0 \cdot y + z = d \implies -x + z = d$

Point  $P$  lies in  $V$ : Filling in yields  $-1 + 3 = 2 = d$ .

## Planes and normals

The plane  $V$  through point  $P = (1, 2, 3)$  with normal  $\mathbf{n} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$ ?

has the equation  $-x + z = 2$  (previous slide)

But also the equation  $-2x + 2z = 4$  with normal  $\mathbf{n} = \begin{bmatrix} -2 \\ 0 \\ 2 \end{bmatrix}$

and equation  $x - z = -2$  with normal  $\mathbf{n} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$

The normals: All parallel!

## Equations for a plane and normal on plane

$$n_1x + n_2y + n_3z = \underbrace{n_1x_0 + n_2y_0 + n_3z_0}_d$$

Formula for  $d$  unimportant, but:

- ▶ From the equation of the plane  $n_1x + n_2y + n_3z = d$  you can see:  
normal =  $(n_1, n_2, n_3)$
- ▶ And vice versa: from the normal =  $(n_1, n_2, n_3)$  you know the equation of the plane:  $n_1x + n_2y + n_3z = d$  where you still need to determine  $d$  via

# Equations for a plane and normal on plane

$$n_1x + n_2y + n_3z = \underbrace{n_1x_0 + n_2y_0 + n_3z_0}_d$$

Formula for  $d$  unimportant, but:

- ▶ From the equation of the plane  $n_1x + n_2y + n_3z = d$  you can see:  
normal =  $(n_1, n_2, n_3)$
- ▶ And vice versa: from the normal =  $(n_1, n_2, n_3)$  you know the equation of the plane:  $n_1x + n_2y + n_3z = d$  where you still need to determine  $d$  via filling in 1 point  $P$  in the plane

# Lines in $\mathbb{R}^3$

Reminder: 2 options: given by

- ▶ parameter representation
- ▶ or 2 equations: intersection of 2 planes

Parameter representation:  $(x, y, z) = (x_1, y_1, z_1) + t(a, b, c)$ ,  $t \in \mathbb{R}$   
support point +  $t \cdot$  direction vector

$$x = 2+t$$


Ex:  $y = 3$  is rewritten as  $(x, y, z) = (2, 3, 0) + t(1, 0, -4)$   
 $z = -4t$

Ex: Line through  $(1, -2, 3)$  orthogonal to plane  $x - 2y + 4z = 5$  has parameter representation

$$(x, y, z) = (1, -2, 3) + t(1, -2, 4) = (1 + t, -2 - 2t, 3 + 4t)$$

# Lines in $\mathbb{R}^3$

Also possible by 2 equations: intersection of 2 planes



One can make a parameter representation out of this by finding 2 points,  
eg:

choose  $x = 0$  then  $y = 0$  and  $z = 1$

choose  $x = 1$  then  $y = 2$  and  $z = -2$

A parameter representation is then

$$(0, 0, 1) + t((1, 2, -2) - (0, 0, 1)) = (0, 0, 1) + t(1, 2, -3)$$

# Planes

- ▶ Give a parameter representation of the line through  $(5, -2, -5)$  parallel to a vector  $(7, 7, 10)$
- ▶ The same for the line through  $(-8, -7, 2)$  orthogonal to the plane  $3x + 3y + 3z = 8$
- ▶ Give the equation for the plane through  $(3, 5, 5)$  and normal to the vector  $(8, 6, 9)$
- ▶ Where does the line  $x = -8 + 3t, y = 4 + 7t, z = -9 - 8t$  intersect the plane  $9x - 6y + 3z = -11$

# Line in $\mathbb{R}^2$ and plane in $\mathbb{R}^3$

Note the similarities and differences:

Line in  $\mathbb{R}^2$ :  $ax + by = c$

( $a, b, c$  fixed,  $x, y$  variable)

Plane in  $\mathbb{R}^3$ :  $ax + by + cz = d$



( $a, b, c, d$  fixed,  $x, y, z$  variable)

# The cross product vector

Cross product vector: A vector orthogonal to two different vectors.

Three points  $A$ ,  $B$  and  $C$  define a start point (eg  $A$ ) and two directions from  $A$  ( $\vec{AB} = \mathbf{b} - \mathbf{a}$  and  $\vec{AC} = \mathbf{c} - \mathbf{a}$ ) and with that a plane with vector representation:  $\mathbf{x} = \mathbf{a} + \lambda(\mathbf{b} - \mathbf{a}) + \mu(\mathbf{c} - \mathbf{a})$ .

The normal  $N$  (vector  $\mathbf{n}$ ) is orthogonal to both vectors.



and can be calculated with the so called **cross product**.

## Definition of the cross product

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \times \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} = \begin{bmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{bmatrix}$$

Ex:  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

Ex:  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} \times \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 0 \\ 0 \end{bmatrix}$

Ex:  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$

Ex:  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$

What is noteworthy about these examples for  $\mathbf{v} \times \mathbf{w}$  ?

# Definition of the cross product

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} \times \begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix} = \begin{bmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{bmatrix}$$

Ex:  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

Ex:  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix} \times \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 0 \\ 0 \end{bmatrix}$

Ex:  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$

Ex:  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$

What is noteworthy about these examples for  $\mathbf{v} \times \mathbf{w}$  ?

- ▶ Cross product is orthogonal to the 2 vectors  $\mathbf{v}$  and  $\mathbf{w}$
- ▶ Length is the same as the surface of a parallelogram formed by  $\mathbf{v}$  and  $\mathbf{w}$

This always turns out to be true.

# Cross product and normal vector

The cross product is useful for calculating the normal vector on the plane

Ex: Plane with parametrisation  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

this is the

# Cross product and normal vector

The cross product is useful for calculating the normal vector on the plane

Ex: Plane with parametrisation  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

this is the  $(x, y)$ -plane

Normal is

# Cross product and normal vector

The cross product is useful for calculating the normal vector on the plane

Ex: Plane with parametrisation  $\begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

this is the  $(x, y)$ -plane

Normal is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \times \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

# Cross product: properties

►  $\mathbf{x} \times \mathbf{x} = \mathbf{0}$

cross product of a vector with itself is the  $\mathbf{0}$ -vector

logical, because then the parallelogram has the surface 0

►  $(\mathbf{x} \times \mathbf{y}) \cdot \mathbf{x} = \mathbf{0}$

cross product of one vector with another is perpendicular to the other

this is 1 of the 2 basic properties of the cross product

►  $\mathbf{y} \times \mathbf{x} = -(\mathbf{x} \times \mathbf{y})$

when you swap the order, you get the opposite direction

"corkscrew-rule"

►  $(\lambda \mathbf{x}) \times \mathbf{y} = \lambda (\mathbf{x} \times \mathbf{y})$

if  $\mathbf{x}$  is eg 2 times longer, then the cross product is 2 times as long

logical, because the length of the cross product is surface of a parallelogram

# Distance of a point to a plane

Distance of point  $(p, q, r)$  to a plane  $ax + by + cz = d$

- 1 Set up the eq. for the line through point orthogonal to plane:

$$(x, y, z) = (p, q, r) + \lambda (a, b, c)$$

- 2 Calculate the intersection point of the line with the plane:  
fill in  $x = p + \lambda a$ ,  $y = q + \lambda b$ ,  $z = r + \lambda c$  into the plane equation
- 3 Calculate distance between  $(p, q, r)$  and intersection point

# Distance of a point to a line

Distance of the point to a line  $\mathbb{R}^3$

- 1 Set up the eq. of a plane through point orthogonal to the line;  
 $ax + by + cz = d$ , with  $(a, b, c)$  the direction vector of the line
- 2 Calculate  $d$  by filling in the point
- 3 calculate intersection of line with plane
- 4 calculate the distance between point and intersection

# Surface of a triangle in plane and space

Surface triangle  $(a, b, c), (d, e, f), (g, h, i)$  in  $\mathbb{R}^3$

- 1 calculate cross product of  $(d, e, f) - (a, b, c)$  and  $(g, h, i) - (a, b, c)$
- 2 surface =  $\frac{1}{2} \cdot$  length of this vector

Surface triangle  $(a, b), (c, d), (e, f)$  in  $\mathbb{R}^2$

- 1 Calculate cross product of  $(c, d, 0) - (a, b, 0)$  and  $(e, f, 0) - (a, b, 0)$
- 2 surface =  $\frac{1}{2} \cdot$  length of this vector

Note: there are also other methods you can use.

# When inner product/when cross product

Let  $\mathbf{u}, \mathbf{v}$  be two vectors.

Inner product:

- ▶ To see if  $\mathbf{u} \perp \mathbf{v} \iff \mathbf{u} \cdot \mathbf{v} = 0$
- ▶ Calculation of the angle  $\theta$  between  $\mathbf{u}$  and  $\mathbf{v}$ :  $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}||\mathbf{v}| \cos \theta$

Cross product:

- ▶ Calculation of a vector  $\mathbf{n}$  orthogonal to  $\mathbf{u}$  and  $\mathbf{v}$ :  $\mathbf{n} = \mathbf{u} \times \mathbf{v}$
- ▶ To see if  $\mathbf{u} \parallel \mathbf{v} \iff \mathbf{u} \times \mathbf{v} = 0$
- ▶ Calculation of the surface  $O$  of the parallelogram spanned by  $\mathbf{u}$  and  $\mathbf{v}$ :  $O = |\mathbf{u} \times \mathbf{v}|$

## Week 2: We have seen

- ▶ Trigonometric functions  $\sin, \dots$
- ▶ Powers  $e^x \ 2^x, \dots$
- ▶ Vectors in 2 and 3 dimensions
- ▶ Lines and planes
- ▶ Distance, angle, inner product
- ▶ Cross product



# Week 3: Limits, Continuity and Differentiation I

- ▶ Limits of functions
- ▶ Limits at Infinity and Infinite limits
- ▶ Continuity, maxima and minima
- ▶ The Intermediate-Value Theorem
- ▶ The Derivative
- ▶ Derivatives of trigonometric functions
- ▶ Rules for Calculating Derivatives
- ▶ The Chain Rule
- ▶ Tangent lines and their direction

# Limits in Computer Science and Data Science

- ▶ Foundation to understand complexity analysis. Without this, you'll never understand algorithms, theory of computation.
- ▶ Concepts and definitions allow to go from *continuous* to **discrete**, **numerical** analysis, leading to numerical algorithms.
- ▶ Needed in statistics (law of large numbers) → data science
- ▶ software testing and deployment

# Ranking Different Algorithms

- ▶ How to formulate this more generally?  
→ Limits

- ▶
$$\lim_{n \rightarrow \infty} \frac{f}{g} = 0 \quad g \text{ grows faster}$$

$$\lim_{n \rightarrow \infty} \frac{f}{g} = c \quad g \text{ grows same as } f$$

$$\lim_{n \rightarrow \infty} \frac{f}{g} = \infty \quad g \text{ grows slower}$$

# Limits not always straightforward

- ▶  $f(n) = n \log(n)$

- ▶  $g(n) = n^{3/2}$

- ▶


$$\lim_{n \rightarrow \infty} \frac{n^{3/2}}{n \log(n)} = ?$$

- ▶ To evaluate we will need:

- ▶ derivatives
- ▶ Taylor series/polynomials
- ▶ Exponential functions and Logarithms
- ▶ rational functions
- ▶ ...



# Statistical estimate of $\pi$

$$\frac{\text{area of circle}}{\text{area of square}} = \frac{\pi \cdot r^2}{(2r)^2} = \frac{\pi}{4}$$



# Statistical estimate of $\pi$

$$\frac{\text{area of circle}}{\text{area of square}} = \frac{\pi \cdot r^2}{(2r)^2} = \frac{\pi}{4}$$



Relative Error  $\sim \frac{1}{\sqrt{N}}$   
 $N$  larger, error smaller

- ▶ randomly select a large number of points within the square
- ▶ ratio of number of points within the circle to the number points within the square provides an estimate for  $\pi/4$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, \quad x_2 = \frac{2}{3}, \quad \dots,$$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, x_2 = \frac{2}{3}, \dots, x_k = \frac{k}{k+1}$$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, x_2 = \frac{2}{3}, \dots, x_k = \frac{k}{k+1}$$

Which value does  $x_k$  take for  $k = 1, 2, 3, \dots$ ?

$$x_k = \frac{k}{k+1} =$$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, x_2 = \frac{2}{3}, \dots, x_k = \frac{k}{k+1}$$

Which value does  $x_k$  take for  $k = 1, 2, 3, \dots$ ?

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} =$$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, x_2 = \frac{2}{3}, \dots, x_k = \frac{k}{k+1}$$

Which value does  $x_k$  take for  $k = 1, 2, 3, \dots$ ?

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \frac{k+1}{k+1} - \frac{1}{k+1} =$$

# The limit of a series of numbers

Consider the sequence of numbers:

$$\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \dots$$

Can we describe the sequence with a simple expression?

Let  $x_k$  be the  $k$ -th entry:

$$x_1 = \frac{1}{2}, x_2 = \frac{2}{3}, \dots, x_k = \frac{k}{k+1}$$

Which value does  $x_k$  take for  $k = 1, 2, 3, \dots$ ?

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \frac{k+1}{k+1} - \frac{1}{k+1} = 1 - \frac{1}{k+1}$$

# The limit of a series of numbers

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \dots =$$

# The limit of a series of numbers

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \dots = 1 - \frac{1}{k+1}$$

Table

$k$	$x_k$
1	0.500000000000000
2	0.666666666666667
3	0.750000000000000
4	0.800000000000000
5	0.833333333333333
:	:
98	0.989898989898990
99	0.990000000000000
100	0.990099009900990

Clearly for  $k = 1, 2, 3, \dots \rightarrow \infty$ :  $x_k \rightarrow 1$

# The limit of a series of numbers

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \dots =$$

# The limit of a series of numbers

$$x_k = \frac{k}{k+1} = \frac{(k+1)-1}{k+1} = \dots = 1 - \frac{1}{k+1}$$

Clearly for  $k = 1, 2, 3, \dots \rightarrow \infty$ :  $x_k \rightarrow 1$

For  $k = 1, 2, \dots \rightarrow \infty$ ,  $x_k$  approaches the limit 1, notation:

$$\lim_{k \rightarrow \infty} x_k = 1$$

Not every series of numbers has a limit:  $x_k = k \rightarrow \infty$  for  $k = 1, 2, \dots, \infty$

# Limits: $x \rightarrow a$

There are **THREE** possible directions to approach  $a$ :

$x \rightarrow a^-$  for “ $x$  approaches  $a$ ” from left

$x \rightarrow a^+$  for “ $x$  approaches  $a$ ” from right

$x \rightarrow a$  for “ $x$  approaches  $a$ ”, from both sides

Some examples:

$x \rightarrow 0^-$ :  $x = -1, -0.1, -0.01, -0.001, \dots$  approaches 0 “quickly”

$x \rightarrow 0^-$ :  $x = -1, -1/2, -1/3, -1/4, \dots$  approaches 0 “slowly”

# Limits: $x \rightarrow a$

There are **THREE** possible directions to approach  $a$ :

$x \rightarrow a^-$  for “ $x$  approaches  $a$ ” from left

$x \rightarrow a^+$  for “ $x$  approaches  $a$ ” from right

$x \rightarrow a$  for “ $x$  approaches  $a$ ”, from both sides

Some examples:

$x \rightarrow 0^-$ :  $x = -1, -0.1, -0.01, -0.001, \dots$  approaches 0 “quickly”

$x \rightarrow 0^-$ :  $x = -1, -1/2, -1/3, -1/4, \dots$  approaches 0 “slowly”

$x \rightarrow 0^+$ :  $x = 1, 0.1, 0.01, 0.001, \dots$

# Limits: $x \rightarrow a$

There are **THREE** possible directions to approach  $a$ :

- $x \rightarrow a^-$  for “ $x$  approaches  $a$ ” from left
- $x \rightarrow a^+$  for “ $x$  approaches  $a$ ” from right
- $x \rightarrow a$  for “ $x$  approaches  $a$ ”, from both sides

Some examples:

$x \rightarrow 0^-$ :  $x = -1, -0.1, -0.01, -0.001, \dots$  approaches 0 “quickly”

$x \rightarrow 0^-$ :  $x = -1, -1/2, -1/3, -1/4, \dots$  approaches 0 “slowly”

$x \rightarrow 0^+$ :  $x = 1, 0.1, 0.01, 0.001, \dots$

$x \rightarrow 2^-$ :  $x = 1.9, 1.99, 1.999, 1.9999, \dots$

$x \rightarrow 2^+$ :  $x = 2.1, 2.01, 2.001, 2.0001, \dots$

# Limits: $x \rightarrow a$

There are **THREE** possible directions to approach  $a$ :

$x \rightarrow a^-$  for “ $x$  approaches  $a$ ” from left

$x \rightarrow a^+$  for “ $x$  approaches  $a$ ” from right

$x \rightarrow a$  for “ $x$  approaches  $a$ ”, from both sides

Some examples:

$x \rightarrow 0^-$ :  $x = -1, -0.1, -0.01, -0.001, \dots$  approaches 0 “quickly”

$x \rightarrow 0^-$ :  $x = -1, -1/2, -1/3, -1/4, \dots$  approaches 0 “slowly”

$x \rightarrow 0^+$ :  $x = 1, 0.1, 0.01, 0.001, \dots$

$x \rightarrow 2^-$ :  $x = 1.9, 1.99, 1.999, 1.9999, \dots$

$x \rightarrow 2^+$ :  $x = 2.1, 2.01, 2.001, 2.0001, \dots$

$x \rightarrow -2^-$ :  $x = -2.1, -2.01, -2.001, -2.0001, \dots$

$x \rightarrow -2^+$ :  $x = -1.9, -1.99, -1.999, -1.9999, \dots$

$x \rightarrow 0$ :  $x = 1, 0.1, -0.1, 0.01, -0.01, 0.001, -0.001, \dots$

$x \rightarrow 0$ :  $x = 1, 0.1, 0.05, 0.01, 0.005, 0.001, 0.0005, \dots$



If  $x \rightarrow a$  then  $f(x) \rightarrow ?$

Ex: If  $x \rightarrow 0^-$  then  $f(x) = x^2 \rightarrow^+ 0$  because:

$x \rightarrow 0^-$	$f(x) = x^2$
-1	1
-0.1	0.01
-0.01	0.001
-0.001	0.00001
-0.0001	0.0000001

Without filling 0 into  $f$ :  $x \rightarrow 0^-$  then  $f(x) \rightarrow f(0)$

If  $x \rightarrow a$  then  $f(x) \rightarrow ?$

Ex: If  $x \rightarrow 0^+$  then  $f(x) = x^2 \rightarrow 0$  because:

$x \rightarrow 0^+$	$f(x) = x^2$
1	1
0.1	0.01
0.01	0.001
0.001	0.00001
0.0001	0.0000001

Without filling 0 into  $f$ :

$$x \rightarrow 0^+$$

$$\implies$$

$$f(x) \rightarrow f(0)$$

$$x \rightarrow 0^-$$

$$\implies$$

previous slide

$$f(x) \rightarrow f(0)$$

$$\implies x \rightarrow 0 \implies f(x) \rightarrow 0$$

# Three types limits of functions

The left  $\lim_{x \rightarrow a^-} f(x)$ , right limit  $\lim_{x \rightarrow a^+} f(x)$  and “THE” limit  $\lim_{x \rightarrow a} f(x)$

Notation: The limit reached by  $f$  for  $x \rightarrow a$  is:

$$\begin{array}{lll} (x \rightarrow a^- \implies f(x) \rightarrow L) & \xrightleftharpoons[\text{notation}]{\iff} & \lim_{x \rightarrow a^-} f(x) = L \\ (x \rightarrow a^+ \implies f(x) \rightarrow R) & \xrightleftharpoons[\text{notation}]{\iff} & \lim_{x \rightarrow a^+} f(x) = R \\ (x \rightarrow a \implies f(x) \rightarrow M) & \xrightleftharpoons[\text{definition}]{\iff} & \lim_{x \rightarrow a} f(x) = M = \lim_{x \rightarrow a^+} f(x) \end{array}$$

Limits  $L$  and  $R$  can exist even though  $f(a)$  does not exist/is not defined!

If  $f(a)$  is a real number, then this is the limit!

Attention:

If  $L = R = \infty$  or  $L = R = -\infty$ , the limit exists and is  $\infty$  resp.  $-\infty$ .

If  $x \rightarrow a$  then  $f(x) \rightarrow ?$

Ex: If  $x \rightarrow 1^-$  then  $f(x) = x + 1 \rightarrow 2$  because:

$x \rightarrow 1^-$	$f(x) = x + 1$
0.9	1.9
0.99	1.99
0.999	1.999
0.9999	1.9999
0.99999	1.99999

So without filling 1 into  $f$ :  $\lim_{x \rightarrow 1^-} x + 1 = 2 = f(1)$ ,

If  $x \rightarrow a$  then  $f(x) \rightarrow ?$

Ex: If  $x \rightarrow 1^-$  then  $f(x) = x + 1 \rightarrow 2$  because:

$x \rightarrow 1^-$	$f(x) = x + 1$
0.9	1.9
0.99	1.99
0.999	1.999
0.9999	1.9999
0.99999	1.99999

So without filling 1 into  $f$ :  $\lim_{x \rightarrow 1^-} x + 1 = 2 = f(1)$ ,  
because  $1.99999\dots = 1.\bar{9} = 2$ .

# Limit: Functions with jumps

Ex: The sign-function (signum) jumps at  $x = 0$  from  $-1$  to  $1$ :


$$\text{sign}(x) = \begin{cases} 1 & \text{for } x > 0 \\ 0 & \text{for } x = 0 \\ -1 & \text{for } x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^-} \text{sign}(x) = -1$$


$$\lim_{x \rightarrow 0^+} \text{sign}(x) = 1$$

$$\lim_{x \rightarrow 0} \text{sign}(x) \text{ does not exist}$$

Now:  $\text{sign}(0)$  exists but is different to the left and right limits!



# Limits: Functions with holes/jumps



$$\lim_{x \rightarrow 1^-} g(x) = 1$$

$$\lim_{x \rightarrow 1^+} g(x) = 0 \quad \lim_{x \rightarrow 1} g(x) \text{ does not exist; } g(1) = 0$$

$$\lim_{x \rightarrow 2^-} g(x) = 1$$

$$\lim_{x \rightarrow 2^+} g(x) = 1 \quad \lim_{x \rightarrow 2} g(x) = 1; \text{ is equal to } g(2) = 1$$

$$\lim_{x \rightarrow 3^-} g(x) = 0$$

$$\lim_{x \rightarrow 3^+} g(x) = 0 \quad \lim_{x \rightarrow 3} g(x) = 0; \text{ but } g(3) = 1$$

The property that for  $a$  a function  $\lim_{x \rightarrow a} f(x) = f(a)$  is special. Such functions are called **continuous** at/in  $a$  (see [A 1.4]).

# Calculation rules limits

Let  $\lim_{x \rightarrow a} f(x) = L$  and  $\lim_{x \rightarrow a} g(x) = M$  exist, then

- $\lim_{x \rightarrow a} [f(x) \pm g(x)] = L \pm M$
- $\lim_{x \rightarrow a} f(x) \cdot g(x) = LM$
- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{L}{M}$  als  $M \neq 0$
- If  $f(x) \leq g(x)$  in an interval around  $a$  then  $L \leq M$

Ex:  $\lim_{x \rightarrow a} x + x^2 = \lim_{x \rightarrow a} x + \lim_{x \rightarrow a} x^2 = a + a^2$

It is necessary that both  $\lim_{x \rightarrow a} f(x) = L$  and  $\lim_{x \rightarrow a} g(x) = M$  exist:

$$1 = \lim_{x \rightarrow 0} 1 \neq \frac{\lim_{x \rightarrow 0} x^2}{\lim_{x \rightarrow 0} x^2} = \frac{0}{0} \quad \dots$$

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

Horizontal, vertical asymptote

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

Horizontal, vertical asymptote

$0 \cdot \infty$

(Week 5, ex  $\lim_{x \rightarrow 0^+} x \ln(x)$ )

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

Horizontal, vertical asymptote

$0 \cdot \infty$

(Week 5, ex  $\lim_{x \rightarrow 0^+} x \ln(x)$ )

$1^\infty$

(ex  $\lim_{x \rightarrow \infty} (1 + \frac{1}{x})^x$ )

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

## Horizontal, vertical asymptote

$0 \cdot \infty$

(Week 5, ex  $\lim_{x \rightarrow 0^+} x \ln(x)$ )

$1^\infty$

(ex  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x$ )

$\infty^0$

(ex  $\lim_{x \rightarrow \infty} \sqrt[x]{x}$ )

# Limits: The difficult cases

$\frac{\neq 0}{0}$

(coming soon!)

$\frac{0}{0}$

(divide factor out, l'Hôpital, Taylor)

$\frac{\infty}{\infty}$

(Week 5, ex  $\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1}$ )

$\infty - \infty$

(Week 5)

## Horizontal, vertical asymptote

$0 \cdot \infty$

(Week 5, ex  $\lim_{x \rightarrow 0^+} x \ln(x)$ )

$1^\infty$

(ex  $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x$ )

$\infty^0$




(ex  $\lim_{x \rightarrow \infty} \sqrt[x]{x}$ )

$0^0$

(ex  $\lim_{x \rightarrow 0^+} x^x$ )

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Typical (simple) cases:



$$\lim_{x \rightarrow 0^+} \frac{1}{x} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x} = -\infty$$

$$\lim_{x \rightarrow 0} \frac{1}{x} \text{ does not exist.}$$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^2} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^2} = \infty$$

$$\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^3} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^3} = -\infty$$

$$\lim_{x \rightarrow 0} \frac{1}{x^3} \text{ does not exist}$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+}$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} =$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} = \frac{-2^+}{0^+ \cdot 4^+} =$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} = \frac{-2^+}{0^+ \cdot 4^+} = \frac{-2}{0^+ \cdot 4} =$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} = \frac{-2^+}{0^+ \cdot 4^+} = \frac{-2}{0^+ \cdot 4} = \frac{-2}{0^+} =$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} = \frac{-2^+}{0^+ \cdot 4^+} = \frac{-2}{0^+ \cdot 4} = \frac{-2}{0^+} = -\infty$$

$$(-2^+ + 2 = (-2 + 2)^+ = 0^+)$$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{\neq 0}{0}$

Ex:  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$ . Filling  $x = -2$  in yields  $\frac{-2}{0}$

1 Factorize the denominator!

2 Analyze both  $\lim_{x \rightarrow -2^+}$  and  $\lim_{x \rightarrow -2^-}$

Because  $x^2 + 8x + 12 = (x + 2)(x + 6)$ :

$$\lim_{x \rightarrow -2^+} \frac{x}{(x + 2)(x + 6)} = \frac{-2^+}{0^+ \cdot 4^+} = \frac{-2}{0^+ \cdot 4} = \frac{-2}{0^+} = -\infty$$

$$(-2^+ + 2 = (-2 + 2)^+ = 0^+)$$

Same:  $\lim_{x \rightarrow -2^-} \frac{x}{(x + 2)(x + 6)} = \infty$

thus  $\lim_{x \rightarrow -2} \frac{x}{x^2 + 8x + 12}$  does not exist

# Why is $3 - 3^+ = 0^-$ ?

$x \rightarrow 3^+$	$3 - 3^+$	$0^-$
3.1	-0.1	-0.1
3.01	-0.01	-0.01
3.001	-0.001	-0.001
3.0001	-0.0001	-0.0001
3.00001	-0.00001	-0.00001

Or: Sequence of numbers  $x \rightarrow 3^+$  of the form  $x_k = 3 + h_k$  with  $h_k > 0$ .  
Then

$$3 - x_k = 3 - (3 + h_k) = -h_k \rightarrow 0^-.$$

## Limits: $f(x) = t(x)/n(x)$ , generals

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = ? \quad \text{if} \quad \lim_{x \rightarrow a} g(x) = g(a) = 0$$

- If numerator does **not** approach 0, then

$$\lim_{x \rightarrow a^-} \frac{f(x)}{g(x)} = \pm\infty \quad \text{and} \quad \lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = \pm\infty$$

Check sign for both cases. Are left and right limits the same?

3 possibilities for  $\lim_{x \rightarrow a}$ :  $\infty$ ,  $-\infty$  or **does not exist**

Useful to factorize the denominator

- If numerator **also** approaches 0:

(a) try to divide a factor out

$$\text{Ex: } \lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3} = \lim_{x \rightarrow 3} \frac{(x - 3)(x + 3)}{x - 3} = \lim_{x \rightarrow 3} x + 3 = 6$$

(b) or with l'Hôpital (later in Calculus)

(c) or with Taylor series (later in Calculus)

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5}$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2}{2-x} =$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2}{2-x} = \frac{4}{0^+} = \infty \quad \text{thus}$$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2}{2-x} = \frac{4}{0^+} = \infty \quad \text{thus} \quad \lim_{x \rightarrow 2} \frac{x^2}{2-x} \text{ does not exist}$$

►  $\lim_{x \rightarrow 2} \frac{x}{x^2 - 4x + 4} =$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2}{2-x} = \frac{4}{0^+} = \infty \quad \text{thus} \quad \lim_{x \rightarrow 2} \frac{x^2}{2-x} \text{ does not exist}$$

►  $\lim_{x \rightarrow 2} \frac{x}{x^2 - 4x + 4} = \lim_{x \rightarrow 2} \frac{x}{(x-2)^2} \quad \text{thus}$

# Limits of functions

►  $\lim_{x \rightarrow 0} \frac{1}{x^5} = ?$

$$\lim_{x \rightarrow 0^+} \frac{1}{x^5} = \frac{1}{0^+} = \infty$$

$$\lim_{x \rightarrow 0^-} \frac{1}{x^5} = \frac{1}{0^-} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{1}{x^5} \text{ does not exist}$$

►  $\lim_{x \rightarrow 0} \frac{x-2}{x^4} = ?$

$$\lim_{x \rightarrow 0^+} \frac{x-2}{x^4} = \frac{-2^+}{0^+} = -\infty$$

$$\lim_{x \rightarrow 0^-} \frac{x-2}{x^4} = \frac{-2^-}{0^+} = -\infty \quad \text{thus} \quad \lim_{x \rightarrow 0} \frac{x-2}{x^4} = -\infty$$

►  $\lim_{x \rightarrow 2} \frac{x^2}{2-x} = ?$

$$\lim_{x \rightarrow 2^+} \frac{x^2}{2-x} = \frac{4}{0^-} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2}{2-x} = \frac{4}{0^+} = \infty \quad \text{thus} \quad \lim_{x \rightarrow 2} \frac{x^2}{2-x} \text{ does not exist}$$

►  $\lim_{x \rightarrow 2} \frac{x}{x^2 - 4x + 4} = \lim_{x \rightarrow 2} \frac{x}{(x-2)^2} \quad \text{thus} = \infty !$

Because the limit from both sides is  $\frac{2}{0^+} = \infty$

Limits:  $f(x) = t(x)/n(x)$ , type  $\frac{0}{0}$

What results for  $\frac{0}{0}$  depends on whether numerator goes faster towards 0 than the denominator:

Ex:  $\lim_{x \rightarrow 0} \frac{x^2}{x} = \lim_{x \rightarrow 0} x = 0$  (numerator faster)

Ex:  $\lim_{x \rightarrow 0} \frac{x}{x^2} = \lim_{x \rightarrow 0} \frac{1}{x}$  does not exist (denominator faster)

Ex:  $\lim_{x \rightarrow 0} \frac{x}{x} = \lim_{x \rightarrow 0} 1 = 1$  (equally fast)

Ex:  $\lim_{x \rightarrow 0} \frac{3x}{x} = \lim_{x \rightarrow 0} 3 = 3$  (both same powers, but different constants)

# Limits: Type $\frac{0}{0}$ with variable substitution and $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$


Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} = \lim_{y \rightarrow 0^+} \frac{\sin(y)}{y} = 1$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{\sin(3x)} = \lim_{x \rightarrow 0} \frac{1}{3} \frac{\sin(x)}{x} \frac{3x}{\sin(3x)}$$

$$= \frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(x)}{x} \cdot \lim_{x \rightarrow 0} \frac{3x}{\sin(3x)} = \frac{1}{3}$$

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\tan(11x)}{x} &= \lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} = \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\ &= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11 \end{aligned}$$



Accepting the fact above, you can calculate other limits, with substitution ↗ ↘

Limits: Type  $\frac{0}{0}$  with variable substitution and  $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$


Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} =$$

$$\lim_{x \rightarrow 0} \frac{1}{3} \frac{\sin(x) 3x}{x \sin(3x)}$$

$$= \frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(x)}{x} \cdot \lim_{x \rightarrow 0} \frac{3x}{\sin(3x)} = \frac{1}{3}$$

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\tan(11x)}{x} &= \lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} = \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\ &= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11\end{aligned}$$



Accepting the fact above, you can calculate other limits, with substitution ↗ ↘


Limits: Type  $\frac{0}{0}$  with variable substitution and  $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} =$$



$$\frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(x)}{x} \cdot \lim_{x \rightarrow 0} \frac{3x}{\sin(3x)} = \frac{1}{3}$$

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\tan(11x)}{x} &= \lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} = \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\ &= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11\end{aligned}$$

Accepting the fact above, you can calculate other limits, with substitution ↗ ↘


Limits: Type  $\frac{0}{0}$  with variable substitution and  $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} =$$



$$\frac{1}{3}$$

$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\tan(11x)}{x} &= \lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} = \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\ &= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11\end{aligned}$$

Accepting the fact above, you can calculate other limits, with substitution ↗ ↘


Limits: Type  $\frac{0}{0}$  with variable substitution and  $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} =$$



$$\begin{aligned}\lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} &= \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\&= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11\end{aligned}$$

Accepting the fact above, you can calculate other limits, with substitution ↗ ↘


Limits: Type  $\frac{0}{0}$  with variable substitution and  $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$

Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} =$$



$$\begin{aligned}& \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\&= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11\end{aligned}$$

Accepting the fact above, you can calculate other limits, with substitution ↗ ↘

# Limits: Type $\frac{0}{0}$ with variable substitution and $\cdot \frac{g(x)}{g(x)}$

Accept now:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$$


Determine now:

$$\lim_{x \rightarrow 0^+} \frac{\sin(\sqrt{4x})}{\sqrt{4x}} = \lim_{y \rightarrow 0^+} \frac{\sin(y)}{y} = 1$$

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{\sin(3x)} = \lim_{x \rightarrow 0} \frac{1}{3} \frac{\sin(x)}{x} \frac{3x}{\sin(3x)}$$

$$= \frac{1}{3} \lim_{x \rightarrow 0} \frac{\sin(x)}{x} \cdot \lim_{x \rightarrow 0} \frac{3x}{\sin(3x)} = \frac{1}{3}$$

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\tan(11x)}{x} &= \lim_{x \rightarrow 0} \frac{\sin(11x)}{\cos(11x)x} = \lim_{x \rightarrow 0} \frac{11 \sin(11x)}{\cos(11x) 11x} \\ &= 11 \cdot \lim_{x \rightarrow 0} \frac{\sin(11x)}{11x} \cdot \lim_{x \rightarrow 0} \frac{1}{\cos(11x)} = 11 \end{aligned}$$



Accepting the fact above, you can calculate other limits, with substitution ↗ ↘

# Limits: Type $\lim_{x \rightarrow \pm\infty}$

- ▶ Already seen: limits themselves can be  $\pm\infty$  ( $\notin \mathbb{R}$ )
- ▶ New (now!): Calculate limits for  $x \rightarrow \pm\infty$

For  $\lim_{x \rightarrow \pm\infty}$  of polynomials: largest power is important

$$\begin{array}{lll} \lim_{x \rightarrow \infty} x & = & \infty \\ \lim_{x \rightarrow -\infty} x & = & -\infty \end{array}$$
$$\begin{array}{lll} \lim_{x \rightarrow \infty} x^2 & = & \infty \\ \lim_{x \rightarrow -\infty} x^2 & = & \infty \end{array}$$
$$\begin{array}{lll} \lim_{x \rightarrow \infty} x^2 - x & = & \infty \\ \lim_{x \rightarrow -\infty} x^2 - x & = & \infty \end{array}$$

The limit can be a number:

Ex:  $\lim_{x \rightarrow -\infty} e^x = 0$

Ex:  $\lim_{x \rightarrow \pm\infty} \frac{1}{x} = 0$

Limits:  $\lim_{x \rightarrow \pm\infty} t(x)/n(x)$ , type  $\frac{\infty}{\infty}$  or  $\frac{0}{0}$




Rational function:  $\frac{p(x)}{q(x)}$ , with  $p(x)$  and  $q(x)$  polynomials

Divide numerator and denominator by largest power of  $x$  in the denominator

$$\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x + 1} = \lim_{x \rightarrow \infty} \frac{2x + 1}{1 + \frac{1}{x}} = \frac{\infty}{1} = \infty$$

$$\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^2 + 1} = \lim_{x \rightarrow \infty} \frac{2 + \frac{1}{x}}{1 + \frac{1}{x^2}} = \frac{2}{1} = 2$$

$$\lim_{x \rightarrow \infty} \frac{2x^2 + x}{x^3 + 1} = \lim_{x \rightarrow \infty} \frac{\frac{2}{x} + \frac{1}{x^2}}{1 + \frac{1}{x^3}} = \frac{0}{1} = 0$$



$\frac{\infty}{\infty}$  can have any answer

It depends on whether the numerator or denominator approaches  $\infty$  faster,  
equally fast fast etc

Ex:  $\lim_{x \rightarrow \infty} \frac{x^2}{x} = \lim_{x \rightarrow \infty} x = \infty$  (numerator faster)


Ex:  $\lim_{x \rightarrow \infty} \frac{x}{x^2} = \lim_{x \rightarrow \infty} \frac{1}{x} = 0$  (denominator faster)

Ex:  $\lim_{x \rightarrow \infty} \frac{x}{x} = \lim_{x \rightarrow \infty} 1 = 1$  (equally fast)

Ex:  $\lim_{x \rightarrow \infty} \frac{3x}{x} = \lim_{x \rightarrow \infty} 3 = 3$  (same power, different constant)

(See also examples on previous slide)

# Squeeze theorem



If  $f(x) \leq g(x) \leq h(x)$  around  $x = a$ , then

$$\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x) \leq \lim_{x \rightarrow a} h(x)$$

If also  $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x)$  then

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x)$$

Ex: Let  $f(x) = \sin(x)$ , assume  $\frac{x}{2} \leq f(x) \leq x$  for  $x > 0$ . Then

$$0 = \lim_{x \rightarrow 0^+} \frac{x}{2} \leq \lim_{x \rightarrow 0^+} \sin(x) \leq \lim_{x \rightarrow 0^+} x = 0 \text{ thus } \lim_{x \rightarrow 0^+} \sin(x) = 0$$

$$\lim_{x \rightarrow \pm\infty} t(x)/n(x)$$


Squeeze Theorem is the last resort:

Ex:  $\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}$  ? Use  $-1 \leq \sin(x) \leq 1$  for all  $x$ :

$$\lim_{x \rightarrow \pm\infty} t(x)/n(x)$$

Squeeze Theorem is the last resort:



Ex:  $\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}$  ? Use  $-1 \leq \sin(x) \leq 1$  for all  $x$ :



$$\lim_{x \rightarrow \pm\infty} t(x)/n(x)$$

Squeeze Theorem is the last resort:

Ex:  $\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}$  ? Use  $-1 \leq \sin(x) \leq 1$  for all  $x$ :





Solve these with

$$\lim_{x \rightarrow \pm\infty} t(x)/n(x)$$

Squeeze Theorem is the last resort:

Ex:  $\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}$  ? Use  $-1 \leq \sin(x) \leq 1$  for all  $x$ :





Solve these with squeeze theorem:

$$\lim_{x \rightarrow \pm\infty} t(x)/n(x)$$

Squeeze Theorem is the last resort:

Ex:  $\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}$  ? Use  $-1 \leq \sin(x) \leq 1$  for all  $x$ :



Solve these with squeeze theorem:  $-\frac{1}{x} \leq \frac{\sin(x)}{x} \leq \frac{1}{x}$  thus


$$\underbrace{\lim_{x \rightarrow \infty} -\frac{1}{x}}_{=0} \leq \underbrace{\lim_{x \rightarrow \infty} \frac{\sin(x)}{x}}_{\text{must therefore also be } 0!} \leq \underbrace{\lim_{x \rightarrow \infty} \frac{1}{x}}_{=0}$$

Do [A 1.2 T75]

# Horizontal asymptote

If  $\lim_{x \rightarrow \pm\infty} f(x) = a$  ( $a \in \mathbb{R}$ , number,  $\neq \pm\infty$ ),

then  $f$  has a horizontal asymptote  $y = a$  for  $x \rightarrow \pm\infty$



Ex:  $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2} \sqrt{1 + \frac{1}{x^2}}} = \lim_{x \rightarrow \infty} \frac{x}{|x| \sqrt{1 + \frac{1}{x^2}}} =$


$$\lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{x^2}}} = 1$$

$$\lim_{x \rightarrow -\infty} \frac{x}{\sqrt{x^2 + 1}} =$$

# Horizontal asymptote

If  $\lim_{x \rightarrow \pm\infty} f(x) = a$  ( $a \in \mathbb{R}$ , number,  $\neq \pm\infty$ ),

then  $f$  has a horizontal asymptote  $y = a$  for  $x \rightarrow \pm\infty$



Ex:  $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} = \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2} \sqrt{1 + \frac{1}{x^2}}} = \lim_{x \rightarrow \infty} \frac{x}{|x| \sqrt{1 + \frac{1}{x^2}}} =$

$$\lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{x^2}}} = 1$$

$$\lim_{x \rightarrow -\infty} \frac{x}{\sqrt{x^2 + 1}} = -1 \text{ because then } |x| = -x$$

Thus horizontal asymptote  $y = 1$  to the right and  $y = -1$  to the left

# Vertical asymptote

$f$  has a **vertical asymptote**  $x = a$

if  $\lim_{x \rightarrow a^+} f(x) = \pm\infty$  or  $\lim_{x \rightarrow a^-} f(x) = \pm\infty$

Here it is allowed that the limit does not exist, thus eg  $\lim_{x \rightarrow a^+} f(x) = \infty$

and  $\lim_{x \rightarrow a^-} f(x) = -\infty$

**Ex:**  $\frac{-6}{x+4}$  has a vertical asymptote in  $x = -4$  because

$$\lim_{x \rightarrow -4^+} \frac{-6}{x+4} = -\infty \quad \text{and} \quad \lim_{x \rightarrow -4^-} \frac{-6}{x+4} = \infty$$

**Ex:**  $\ln(x)$  has a vertical asymptote in  $x = 0$  because

$$\lim_{x \rightarrow 0^+} \ln(x) = -\infty \quad \left( \lim_{x \rightarrow 0^-} \ln(x) \text{ does not exist, domain} \right)$$

# Some standard limits

If  $a > 0$ , then:

$$\lim_{x \rightarrow \infty} \frac{x^a}{e^x} = 0$$

$$\lim_{x \rightarrow \infty} \frac{\ln(x)}{x^a} = 0$$

$$\lim_{x \rightarrow -\infty} |x| e^x = 0$$

$$\lim_{x \rightarrow 0^+} x^a \ln(x) = 0$$

This also holds for  $b^x$  ( $b > 1$ ) in particular for  $e^x$

For  $x \rightarrow \infty$ ,  $b^x$  ( $b > 1$ ) approaches  $\infty$  faster than any polynomial  
(or any positive power), and any polynomial faster than  $\ln(x)$

For  $x \rightarrow -\infty$ ,  $b^x$  ( $b > 1$ ) approaches 0 faster than any polynomial approaches  $\infty$

For  $x \rightarrow 0^+$  approaches any positive power 0 faster than  $\ln(x)$  approaches  $-\infty$

# Limits of functions

- ▶  $\lim_{x \rightarrow \pm\infty} \frac{-8 + \frac{5}{x}}{7 - \frac{4}{x^2}}$
- ▶  $\lim_{x \rightarrow \infty} \frac{3x^7 + 4x^6 + 6}{9x^8}$
- ▶  $\lim_{x \rightarrow \infty} \frac{7\sqrt{x} + x^{-4}}{4x - 5}$
- ▶  $\lim_{x \rightarrow -\infty} \frac{8 - 8x + \sin(8x)}{8x - \cos(8x)}$
- ▶  $\lim_{x \rightarrow \infty} \frac{2^{3x} + 3^{2x}}{7^{x+9} + 9^x}$
- ▶  $\lim_{x \rightarrow \infty} \frac{7x^3 + 8x^5 + 2^{2x}}{1 + 4^{x-1}}$


# Continuity of function $f$ in point $a$

$f$  is right continuous in  $a$  if  $\lim_{x \rightarrow a^+} f(x) = f(a)$  [A 1.4 D6]


$f$  is left continuous in  $a$  if  $\lim_{x \rightarrow a^-} f(x) = f(a)$  [A 1.4 D6]

$f$  is continuous in  $a$  if  $\lim_{x \rightarrow a} f(x) = f(a)$  [A 1.4 D4]


Thus  $f$  is continuous in  $a \iff$  left and right continuous in  $a$




left continuous in  $x = 1$   
right continuous in  $x = 1$   
continuous in  $x = 1$



not left continuous in  $x = 0$   
right continuous in  $x = 0$   
no continuous in  $x = 0$



not left continuous in  $x = 0$   
not right continuous in  $x = 0$   
not continuous in  $x = 0$



not continuous in  $x = 1$   
but make continuous by defining  
 $f(1) = 2$   
"removable discontinuity"

# Continuity of function $f$ in point $a$

Repeat: In mathematical terms  $f$  is continuous in  $a$

- (I)  $f(a)$  exists
- (II)  $\lim_{x \rightarrow a^+} f(x) = f(a)$  exists
- (III)  $\lim_{x \rightarrow a^-} f(x) = f(a)$  exists;

It means (graphically): There is no hole/jump in the graph of  $f$  at  $a$

# Continuity of function $f$ in $x \in [a, b]$

The definition:

$f$  is continuous on an interval  $[a, b]$  (or on  $\mathbb{R}$ )

if  $f$  is continuous in every  $x \in (a, b)$  (or in every  $x \in \mathbb{R}$ )

If an endpoint as  $a$  is involved:

- ▶  $f(a)$  must exist and
- ▶ the right limit must be  $R = f(a)$

Similar for right endpoint.

Ex:  $f(x) = 1/x$  is continuous on  $(0, \infty)$

If  $f$  is continuous on a CLOSED interval  $[a, b]$  then there are no holes/jumps in the graph  $f$  over  $[a, b]$ , so then  $f$  obviously has a minimal and an maximal value!

# Intermediate-Value Theorem

If  $f$  is continuous on  $[a, b]$  then  $f$  takes every value between  $f(a)$  and  $f(b)$  because  $f$  does not have any holes or jumps in  $[a, b]$ :

If  $f$  is continuous on  $[a, b]$  then there is for every  $s \in [f(a), f(b)]$  a  $c$  with  $f(c) = s$ :





Figure 1.32 The continuous function  $f$  takes on the value  $s$  at some point  $c$  between  $a$  and  $b$

# Intermediate-Value Theorem

Ex:  $f(x) = x^3 - 15x + 1$  has three zeros on  $[-4, 4]$ :



x	f(x)	sign
-4	-3	-
-3	19	+
-2	23	+
-1	15	+
-0	1	+
1	-13	-
2	-21	-
3	-17	-
4	5	+

At  $x = -4$  is  $f(-4) < 0$  and  $x = -3$   $f(-3) > 0$ .  $f$  is continuous thus it has at least one zero on  $[-3, -4]$ . Same for zeros in  $(-1, 0)$  and in  $(3, 4)$

# Continuity: Examples

Ex: Continuous on its complete domain  $D$  are:

- ▶ polynomials  $D = \mathbb{R}$
- ▶ powers  $a^x$  on  $D = \mathbb{R}$  for  $a \geq 0$ ,  $\sqrt{x}$  for  $D = [0, \infty)$
- ▶ sin, cos on  $D = \mathbb{R}$
- ▶ tan on  $\mathbb{R} - \{\frac{\pi}{2} + k\pi\}$  for all integers  $k$
- ▶ sums, differences, products of continuous functions
- ▶ quotient of functions:  $\frac{f(x)}{g(x)}$  on

# Continuity: Examples

Ex: Continuous on its complete domain  $D$  are:

- ▶ polynomials  $D = \mathbb{R}$
- ▶ powers  $a^x$  on  $D = \mathbb{R}$  for  $a \geq 0$ ,  $\sqrt{x}$  for  $D = [0, \infty)$
- ▶ sin, cos on  $D = \mathbb{R}$
- ▶ tan on  $\mathbb{R} - \{\frac{\pi}{2} + k\pi\}$  for all integers  $k$
- ▶ sums, differences, products of continuous functions
- ▶ quotient of functions:  $\frac{f(x)}{g(x)}$  on  $D_f \cap D_g \cap \{x \in \mathbb{R}: g(x) \neq 0\}$
- ▶ compositions of continuous functions

# Continuity: Examples

Ex:  $\sin(\sqrt{x^2 - 1})$  is continuous on its domain  $(-\infty, -1] \cup [1, \infty)$   
because:

# Continuity: Examples

Ex:  $\sin(\sqrt{x^2 - 1})$  is continuous on its domain  $(-\infty, -1] \cup [1, \infty)$   
because:

- ▶  $x^2$  continuous, 1 continuous,  $x^2 - 1$  continuous on  $\mathbb{R}$

# Continuity: Examples

Ex:  $\sin(\sqrt{x^2 - 1})$  is continuous on its domain  $(-\infty, -1] \cup [1, \infty)$   
because:

- ▶  $x^2$  continuous, 1 continuous,  $x^2 - 1$  continuous on  $\mathbb{R}$
- ▶  $\sqrt{y}$  continuous for  $y \geq 0$ ,  $\sqrt{x^2 - 1}$  continuous on  $(-\infty, -1] \cup [1, \infty)$

# Continuity: Examples

Ex:  $\sin(\sqrt{x^2 - 1})$  is continuous on its domain  $(-\infty, -1] \cup [1, \infty)$   
because:

- ▶  $x^2$  continuous, 1 continuous,  $x^2 - 1$  continuous on  $\mathbb{R}$
- ▶  $\sqrt{y}$  continuous for  $y \geq 0$ ,  $\sqrt{x^2 - 1}$  continuous on  $(-\infty, -1] \cup [1, \infty)$
- ▶  $\sin(y)$  continuous,  $\sin(\sqrt{x^2 - 1})$  continuous on  $(-\infty, -1] \cup [1, \infty)$

# Continuity: Limits

If  $\lim_{x \rightarrow a} g(x)$  exists and  $f$  is continuous at  $g(a)$  then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(g(a))$$

Holds too for left and right limits. Ex: Calculate:

$$\lim_{x \rightarrow 0} \cos\left(\frac{\pi - \pi \cos^2 x}{x^2}\right).$$

The cosine is a continuous function (for all inputs), thus first:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\pi - \pi \cos^2 x}{x^2} &= \lim_{x \rightarrow 0} \frac{\pi \sin^2 x}{x^2} = \pi \lim_{x \rightarrow 0} \frac{\sin^2 x}{x^2} \\ &= \pi \lim_{x \rightarrow 0} \left(\frac{\sin x}{x}\right)^2 = \pi \cdot 1^2 = \pi \end{aligned}$$

so that

$$\lim_{x \rightarrow 0} \cos\left(\frac{\pi - \pi \cos^2 x}{x^2}\right) = \cos\left(\lim_{x \rightarrow 0} \frac{\pi - \pi \cos^2 x}{x^2}\right) = \cos(\pi) = -1$$

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$   
Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$

Do [A 1.4 T9,15]

# Continuity

- For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?  $x \in \mathbb{R}$   
because composition of continuous functions
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$   
Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?  $x \in \mathbb{R}$   
because composition of continuous functions
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$   
Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$   
 $f(5)$  is not defined because 0/0

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?  $x \in \mathbb{R}$   
because composition of continuous functions
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$

Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$

$f(5)$  is not defined because 0/0

$$\lim_{x \rightarrow 5} \frac{6x^2 - 150}{6x - 30}$$

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?  $x \in \mathbb{R}$   
because composition of continuous functions
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$

Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$

$f(5)$  is not defined because 0/0

$$\lim_{x \rightarrow 5} \frac{6x^2 - 150}{6x - 30} = \lim_{x \rightarrow 5} \frac{x^2 - 25}{x - 5}$$

# Continuity

- ▶ For which  $x$  is  $f(x) = \frac{8}{x-4} - 3x$  continuous?  
 $x \in \mathbb{R} \setminus \{4\}$  (or: only not for  $x = 4$ )
- ▶ For which  $x$  is  $f(x) = |9x - 4| - 4 \sin(5x)$  continuous?  $x \in \mathbb{R}$   
because composition of continuous functions
- ▶ Let  $f(x) = \frac{6x^2 - 150}{6x - 30}$

Determine  $f(5)$  so that  $f$  is continuous for  $x = 5$

$f(5)$  is not defined because 0/0

$$\lim_{x \rightarrow 5} \frac{6x^2 - 150}{6x - 30} = \lim_{x \rightarrow 5} \frac{x^2 - 25}{x - 5} = \lim_{x \rightarrow 5} \frac{(x - 5)(x + 5)}{x - 5} = 10$$

So we define  $f(5) = 10$ , then  $f$  is continuous everywhere

Do [A 1.4 T9,15]

# Secure Key Exchange: Elliptic Diffie Hellman

- ▶ based on addition of points on an *elliptic curve*
- ▶ mathematically, *groups* consist of elements (points on elliptic curve) and operations (addition of points) that fulfil the four fundamental properties of closure, associativity, the identity property, and the inverse property.
- ▶ in other words: the operation "addition" (and so multiplication) can be a very strange looking operation compared to the common addition of real numbers
- ▶  $P \oplus Q$  is the addition of two points on an elliptic curve
- ▶  $nP = P \oplus P \oplus \dots \oplus P$  is the multiplication of a point

# Secure Key Exchange: Elliptic Diffie Hellman

Elliptic curve:  $y^2 = x^3 + ax + b$

- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.

# Secure Key Exchange: Elliptic Diffie Hellman

Elliptic curve:  $y^2 = x^3 + ax + b$

- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.
- 2 User1 and User2 both generate a private key number,  $n_1$  and  $n_2$ .

# Secure Key Exchange: Elliptic Diffie Hellman

Elliptic curve:  $y^2 = x^3 + ax + b$

- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.
- 2 User1 and User2 both generate a private key number,  $n_1$  and  $n_2$ .
- 3 User1 calculates  $n_1 P$  and User2  $n_2 P$ . Both send the result over.

# Secure Key Exchange: Elliptic Diffie Hellman

Elliptic curve:  $y^2 = x^3 + ax + b$

- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.
- 2 User1 and User2 both generate a private key number,  $n_1$  and  $n_2$ .
- 3 User1 calculates  $n_1P$  and User2  $n_2P$ . Both send the result over.
- 4 User1 calculates  $n_1(n_2P)$  and User2  $n_2(n_1P)$ .

## Secure Key Exchange: Elliptic Diffie Hellman

Elliptic curve:  $y^2 = x^3 + ax + b$


- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.
- 2 User1 and User2 both generate a private key number,  $n_1$  and  $n_2$ .
- 3 User1 calculates  $n_1P$  and User2  $n_2P$ . Both send the result over.
- 4 User1 calculates  $n_1(n_2P)$  and User2  $n_2(n_1P)$ .
- 5 Both users should now know the same point, without knowing each other's private key number.

# Secure Key Exchange: Elliptic Diffie Hellman


Elliptic curve:  $y^2 = x^3 + ax + b$

- 1 User1 initiates a connection with User2, and selects a generator  $P$  (a point on the curve) and the parameters  $a$  and  $b$  of the elliptic curve equation, and sends them across the wire as plain text.
- 2 User1 and User2 both generate a private key number,  $n_1$  and  $n_2$ .
- 3 User1 calculates  $n_1P$  and User2  $n_2P$ . Both send the result over.
- 4 User1 calculates  $n_1(n_2P)$  and User2  $n_2(n_1P)$ .
- 5 Both users should now know the same point, without knowing each other's private key number.
- 6 The point is used as the symmetrical encryption key for all data transfers.


# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman




# Secure Key Exchange: Elliptic Diffie Hellman



# Secure Key Exchange: Elliptic Diffie Hellman



# Secure Key Exchange: Elliptic Diffie Hellman



# The Derivative

You might have learned at school :

- ▶ The derivative of  $x^n = n \cdot x^{n-1}$
- ▶ The derivative of  $\cos(x) = -\sin(x)$

How is that determined/calculated?

# The Derivative

You might have learned at school :

- ▶ The derivative of  $x^n = n \cdot x^{n-1}$
- ▶ The derivative of  $\cos(x) = -\sin(x)$

How is that determined/calculated? With limits from calculus ...

# Derivative of a function $f$ in point $x$

If both

Left derivative  $\lim_{h \rightarrow 0^-} \frac{f(x+h) - f(x)}{h} = L$

Right derivative  $\lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h} = R$

exist and  $L = R \neq \pm\infty$  then **the** derivative of  $f$  in  $x$  exists:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

This is the **slope of the tangent line** in  $(x, f(x))$ .

If this limit is a number, then the function  $f$  is called **differentiable in  $x$**

## Derivative of function $f$ in point $x$

Derivative  $f'$  indicates:  $f$  increasing/rising, decreasing/falling, or horizontal.

If  $f'(x) > 0$  then the function rises at  $x$

If  $f'(x) < 0$  then the function falls at  $x$

If  $f'(x) = 0$  then the function  $f$  has a maximum, minimum or inflection point at  $x$

If  $f''(x) = 0$  and  $f''$  changes sign at  $x$  then  $f$  has an inflection point at  $x$

Convention: If the derivative is  $\pm\infty$ , it does **not** exist.

For limits, we learned that they CAN exist if they are  $\pm\infty$  ....  
A convention must not be logical. So, just learn it.

# Derivative: Notation

Different notation for the derivative  $f'$  is:

$$f' = \frac{df}{dx} = \frac{d}{dx} f$$

If we call the function  $y$ , it's the same (eg  $y(x) = x^2$ ):  
Different notation for  $y'$  is:

$$y' = \frac{dy}{dx} = \frac{d}{dx} y$$

# Derivative: Examples

Ex: Function  $f(x) = x^2$ :

- ▶ at  $x = 0$ :  $f'(0) = \lim_{h \rightarrow 0} \frac{f(0 + h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h^2 - 0}{h} = \lim_{h \rightarrow 0} h = 0$
- ▶ at  $x = 1$ :  
$$f'(1) = \lim_{h \rightarrow 0} \frac{f(1 + h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{(1 + h)^2 - 1}{h} = \lim_{h \rightarrow 0} \frac{2h + h^2}{h} = 2$$
- ▶ For general  $x$ :

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x + h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{(x^2 + 2xh + h^2) - x^2}{h} = \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = 2x\end{aligned}$$

# Derivative: Example $f(x) = |x|$


If  $x > 0$ :  $f(x) = x$ , thus  $f'(x) = 1$

If  $x < 0$ :  $f(x) = -x$ , thus  $f'(x) = -1$

If  $x = 0$ : Left and right derivative:

$$\begin{aligned} \blacktriangleright \lim_{h \rightarrow 0^+} \frac{|0+h| - |0|}{h} &= \lim_{h \rightarrow 0^+} \frac{|0+h|}{h} = \\ \lim_{h \rightarrow 0^+} \frac{h}{h} &= 1 \quad (h > 0!) \end{aligned}$$

$$\begin{aligned} \blacktriangleright \lim_{h \rightarrow 0^-} \frac{|0+h| - |0|}{h} &= \lim_{h \rightarrow 0^-} \frac{|0+h|}{h} = \\ \lim_{h \rightarrow 0^-} \frac{-h}{h} &= -1 \quad (h < 0!) \end{aligned}$$




are different thus  $\lim_{h \rightarrow 0} \frac{|0+h| - |0|}{h}$  does not exist

Conclusion:  $f(x) = |x|$  is not differentiable in  $x = 0$

Equivalent: There is no clear tangent to the graph at  $(0,0)$

$$f(x) = |x^2 - 1|/x \text{ for } x = 1$$

Right derivative:  $h \rightarrow 0^+ \implies h > 0 \implies |(1+h)^2 - 1| = h^2 + 2h.$




$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h} &\stackrel{x=1}{=} \lim_{h \rightarrow 0^+} \left( \frac{|(1+h)^2 - 1|}{1} - \underbrace{\frac{|1^2 - 1|}{1}}_{=0} \right) / h \\ &= \lim_{h \rightarrow 0^+} \frac{h^2 + 2h}{h} \\ &= \lim_{h \rightarrow 0^+} h + 2 \\ &= 2 =: R \end{aligned}$$

$$f(x) = |x^2 - 1|/x \text{ for } x = 1$$

$f(x)$  for  $x \rightarrow 1^-$ :  $(1 + h)^2 - 1 = h^2 + 2h$  for  $h < 0$ :

$h \rightarrow 0^-$	$h^2 + 2h$
-3	$9 - 6 = 3$
-2	$4 - 4 = 0$
-1	$1 - 2 = -1$
-0.1	$0.01 - 0.2 = -0.19$
-0.01	$0.0001 - 0.02 = -0.0199$
-0.001	$0.000001 - 0.002 = -0.001999$
-0.0001	$0.00000001 - 0.0001 = -0.00019999$




thus  $(1 + h)^2 - 1 < 0$  for  $h \rightarrow -0$  and with that  
 $|(1 + h)^2 - 1| = -(h^2 + 2h)$  for  $h \rightarrow 0^-$

$$f(x) = |x^2 - 1|/x \text{ for } x \neq 1$$

Left derivative:  $|1 + h|^2 - 1| = -(h^2 + 2h)$  for  $h \rightarrow 0^-$ :

$$\begin{aligned} \lim_{h \rightarrow 0^-} \frac{f(x+h) - f(x)}{h} &\stackrel{x=1}{=} \lim_{h \rightarrow 0^-} \left( \frac{|(1+h)^2 - 1|}{1} - \underbrace{\frac{|1^2 - 1|}{1}}_{=0} \right) / h \\ &= \lim_{h \rightarrow 0^-} \frac{-h^2 - 2h}{h} \\ &= \lim_{h \rightarrow 0^-} -(h+2) \\ &= -2 =: L \end{aligned}$$



Left and right derivatives are different:  $-2 = L \neq R = 2$  thus  
**the** derivative does not exist.

# Derivatives: Differentiable on $[a, b]$

For endpoints  $a$  and  $b$  only a right or left derivative can exist:

Left derivative  $f'(b) = \lim_{h \rightarrow 0^-} \frac{f(b+h) - f(b)}{h}$

Right derivative  $f'(a) = \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h}$

For all other points in  $[a, b]$  it must hold that for the left derivative  $L$  and right derivative  $R$  exist and that  $L = R \neq \pm\infty$ .


# Derivatives: Differentiable on $[a, b]$

Ex:  $\sqrt{x}$ : domain  $[0, \infty)$ .

Thus for  $x = 0$  only  $\lim_{h \rightarrow 0^+}$  needs to exist:  
 $\sqrt{x}$  is not right differentiable at  $x = 0$  since

$$\lim_{h \rightarrow 0^+} \frac{\sqrt{h} - \sqrt{0}}{h} = \lim_{h \rightarrow 0^+} \frac{1}{\sqrt{h}} = \infty$$

This is a vertical tangent.




# Tangents to graph of a function

The tangent to the graph of  $f$  at point  $x = a$

- 1 includes the point of the graph, thus  $(a, f(a))$
- 2 has direction coefficient  $f'(a)$

and has the equation:  $y - f(a) = f'(a)(x - a)$ .

Thus  $y = f(a) + f'(a)(x - a)$  – see Taylor series later!



Ex: Function  $f(x) = x^2$ .

Tangent to the graph at  $a = 0$  is x-axis:

$$f(x) = x^2, a = 0 \implies f(a) = a^2 = 0$$

$$f'(x) = 2x, a = 0 \implies f'(a) = 0.$$


$$\text{Thus tangent: } y - 0 = 0 \cdot (x - 0)$$

# Normal to the graph of a function

Normal in a point

= line orthogonal to the graph in a point

= line orthogonal to the tangent in that point



Tangent in  $x = \frac{1}{2}$  has slope  $m = 1$ , goes through  $(\frac{1}{2}, \frac{1}{4})$ ,  
thus it has the equation  $y = x - \frac{1}{4}$

Normal has slope  $m_2 = -1$ , also goes through  $(\frac{1}{2}, \frac{1}{4})$ ,  
thus it has the equation  $y = -x + \frac{3}{4}$

## Quiz: Derivatives with Definition

Determine:

- ▶  $f'(0)$  and  $f'(7)$  for  $f(x) = 5 + x^2$
- ▶  $\frac{dy}{dx}$  if  $y = -8x^3 + 6x$
- ▶  $\frac{dy}{dx}$  if  $y = \frac{5}{\sqrt{x+7}}$
- ▶ Derivative of  $f(x) = 4x + \frac{5}{x}$  and slope of the tangent at  $x = 1$
- ▶ Derivative of  $f(x) = 4 + \sqrt{3-x}$  and equation for the tangent at  $(x, y) = (-6, 7)$

# The derivative with help of the definitions

Just seen: Lots of work. Often, we can do that easier: Recipe:

- 1 Learn a number of derivatives: eg  $\sin'(x) = \cos(x)$ ,  $\ln'(x) = 1/x$ , etc
- 2 Use calculation rules:  $(f + g)' = f' + g'$ ,  $(f \cdot g)' = f' \cdot g + f \cdot g'$ , etc

Attention: You must be able to calculate the derivative of a (simple?) function using the definition! That could happen in an exam!

# Learn these derivatives!

$f(x)$	$f'(x)$
$c$	0
$x^r$	$r x^{r-1}$
$a^x$	$a^x \ln(a)$
$e^x$	$e^x$
$\ln(x)$	$\frac{1}{x}$
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$
$\tan(x)$	$1 + \tan^2(x) = \frac{1}{\cos^2(x)}$
$f(x)g(x)$	via $e^{g(x)\ln(f(x))}$

# Differentiation rules

$f'$  via definition is a lot of work  $\Rightarrow$  therefore rules, eg  $(x^r)' = rx^{r-1}$

$$\begin{aligned}(f+g)'(x) &= f'(x) + g'(x) \\(f-g)'(x) &= f'(x) - g'(x) \\(fg)'(x) &= f'(x)g(x) + f(x)g'(x) \\\left(\frac{f}{g}\right)'(x) &= \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2} \\(cg)'(x) &= 0 \cdot g(x) + cg'(x) = cg'(x) \\\left(\frac{1}{g}\right)'(x) &= \frac{0 \cdot g(x) - 1 \cdot g'(x)}{(g(x))^2} = -\frac{g'(x)}{(g(x))^2}\end{aligned}$$

Proof: with the definition ..., eg:

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{f(x+h)(g(x+h) - g(x))}{h} + \frac{(f(x+h) - f(x))g(x)}{h} \\&= f'(x)g(x) + f(x)g'(x)\end{aligned}$$



# Derivatives of sin, cos, tan

We have already calculated the derivative of sine.

The one of the cosine works just the same way.

For the one of the tan function:

$$\begin{aligned}\tan'(x) &= \left( \frac{\sin(x)}{\cos(x)} \right)' = \frac{\cos(x)\cos(x) - \sin(x)(-\sin(x))}{\cos^2(x)} \\ &= \frac{1}{\cos^2(x)} = 1 + \tan^2(x)\end{aligned}$$

# Chain Rule: Examples

Chain rule for  $(f \circ g)(x) = f(g(x))$ :

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

Ex:  $y = (7x - 3)^{10} = f(g(x))$ , with

# Chain Rule: Examples

Chain rule for  $(f \circ g)(x) = f(g(x))$ :

$$(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$$

Ex:  $y = (7x - 3)^{10} = f(g(x))$ , with  $f(x) = x^{10}$  and  $g(x) = 7x - 3$

$$f'(x) = 10x^9, \quad g'(x) = 7$$

$$y' = f'(g(x)) \cdot g'(x) = 10(7x - 3)^9 \cdot 7 = 70(7x - 3)^9$$

$$\text{Ex: } \frac{d}{dx}\left(\frac{1}{f}\right)(x) = \frac{d}{dx}(f(x))^{-1} = (-1) \cdot (f(x))^{-2} \cdot f'(x) = -\frac{f'(x)}{(f(x))^2}$$

Also seen already with quotient rule!

# Chain Rule: Examples

Ex: Determine  $f'(x)$  if  $f(x) = (x^2 + 1)^2$

Here we can first rewrite  $f$ :  $f(x) = x^4 + 2x^2 + 1$

and then calculate the derivative:  $f(x) = 4x^3 + 4x$

or with chain rule:  $f'(x) = 2(x^2 + 1) \cdot 2x = 4x(x^2 + 1)$  same!

(because  $f(x) = g(h(x))$  with  $g(x) = x^2$  and  $h(x) = x^2 + 1$ )

# Chain Rule: Examples

Ex: Determine  $f'(x)$  if  $f(x) = (x^2 + 1)^2$

Here we can first rewrite  $f$ :  $f(x) = x^4 + 2x^2 + 1$

and then calculate the derivative:  $f(x) = 4x^3 + 4x$

or with chain rule:  $f'(x) = 2(x^2 + 1) \cdot 2x = 4x(x^2 + 1)$  same!

(because  $f(x) = g(h(x))$  with  $g(x) = x^2$  and  $h(x) = x^2 + 1$ )

Ex: But now: determine  $f'(x)$  if  $f(x) = (x^2 + 1)^{40}$

# Chain Rule: Examples

Ex: Determine  $f'(x)$  if  $f(x) = (x^2 + 1)^2$

Here we can first rewrite  $f$ :  $f(x) = x^4 + 2x^2 + 1$

and then calculate the derivative:  $f(x) = 4x^3 + 4x$

or with chain rule:  $f'(x) = 2(x^2 + 1) \cdot 2x = 4x(x^2 + 1)$  same!

(because  $f(x) = g(h(x))$  with  $g(x) = x^2$  and  $h(x) = x^2 + 1$ )

Ex: But now: determine  $f'(x)$  if  $f(x) = (x^2 + 1)^{40}$

Rewriting  $f$  first is not very nice!

But with chain rule:

$$f'(x) = 40(x^2 + 1)^{39} \cdot 2x = 80x(x^2 + 1)^{39} \quad \text{very quick!}$$

(because  $f(x) = g(h(x))$  with  $g(x) = x^{40}$  and  $h(x) = x^2 + 1$ )

## Quiz: Sum, product, quotient and chain rule

Calculate  $y'(x)$  for:

- ▶  $y = \left(-4 - \frac{5x}{7}\right)^{-7}$
- ▶  $y = \sqrt{-4 + 3x}$
- ▶  $y = \sqrt{-9 + \sqrt{3x}}$
- ▶  $y = \frac{\sin(x)}{1+\sin(x)}$
- ▶  $y = \cos(\sin(x))$
- ▶  $y = x(7x^2 + 1)^{1/2}$
- ▶  $y = \sqrt{\frac{1}{3x^5}}$
- ▶  $y = (5x + 2)^4(2x + 5)^{-3}$
- ▶  $y = \sin^2(3x - \pi)$
- ▶  $y = \sin((8x - 3)^{-9/2})$
- ▶ Let  $f(x) = x^2 - 1$  and  $g(x) = x^2 + 1$   
Calculate  $(f \circ g)'$  in 2 ways:
  - "usual" way: first  $(f \circ g)(x)$  and then differentiate
  - with the chain rule  $(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$
- ▶ Tangent and normal to  $y = \sqrt{x}$  in  $(4, 2)$
- ▶ Slope and equation for the tangent to  $y = 2 - 9x^2$  in  $(3, -79)$
- ▶ Slope and equation for the tangent to  $y = 6\sqrt{x}$  in  $(1, 6)$

## Week 3: We have seen

- ▶ Limits  $\lim_{x \rightarrow a} f(x)$
- ▶ Limits  $\lim_{x \rightarrow \pm\infty} f(x)$  and  $\lim_{x \rightarrow a} f(x) = \pm\infty$   
(horizontal/vertical asymptotes)
- ▶ Definition continuous (without holes and jumps)
- ▶ Intermediate-Value Theorem
- ▶ Tangent to  $(a, f(a))$ :  $y - f(a) = f'(a)(x - a)$
- ▶ Definition derivative  $\lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}$
- ▶ Calculation rules: derivative/chain rule  $(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$
- ▶ Derivatives of trigonometric functions

$$\tan'(x) = \frac{1}{\cos^2(x)} = 1 + \tan^2(x)$$

