

# Additional Hypothesis Test Applications Solutions

Lt Col Ken Horton

Lt Col Kris Pruitt

Professor Bradley Warner

26 October, 2020

## Exercises

### 1. Golf balls

Repeat the analysis of the golf ball problem from earlier this semester.

- a. Load the data and tally the data into a table. The data is in `golf_balls.csv`.

```
golf_balls <- read_csv("data/golf_balls.csv")
```

```
head(golf_balls)
```

```
## # A tibble: 6 x 1
##   number
##   <dbl>
## 1      3
## 2      2
## 3      1
## 4      4
## 5      4
## 6      3
```

```
tally(~number, data=golf_balls)
```

```
## number
##   1    2    3    4
## 137 138 107 104
```

- b. Using the function `chisq.test` conduct a hypothesis test of equally likely distribution of balls. You may have to read the help menu.

```
chisq.test(tally(~number, data=golf_balls), p=c(.25, .25, .25, .25))
```

```
##
## Chi-squared test for given probabilities
##
## data:  tally(~number, data = golf_balls)
## X-squared = 8.4691, df = 3, p-value = 0.03725
```

- c. Repeat part b. but assume balls with the numbers 1 and 2 occur 30% of the time and balls with 3 and 4 occur 20%.

```
chisq.test(tally(~number,data=golf_balls),p=c(.3,.3,.2,.2))
```

```
##  
## Chi-squared test for given probabilities  
##  
## data:  tally(~number, data = golf_balls)  
## X-squared = 2.4122, df = 3, p-value = 0.4914
```

## 2. Bootstrap hypothesis testing

Repeat the analysis of the MLB data from the lesson but this time generate a bootstrap distribution of the  $F$  statistic.

First, read in the data.

```
mlb_obp <- read_csv("data/mlb_obp.csv")
```

Convert position to a factor.

```
mlb_obp <- mlb_obp %>%  
  mutate(position=as.factor(position))
```

Summarize the data.

```
favstats(obp~position,data=mlb_obp)
```

	position	min	Q1	median	Q3	max	mean	sd	n	missing
## 1	C	0.219	0.30000	0.3180	0.35700	0.405	0.3226154	0.04513175	39	0
## 2	DH	0.287	0.31625	0.3525	0.36950	0.412	0.3477857	0.03603669	14	0
## 3	IF	0.174	0.30800	0.3270	0.35275	0.437	0.3315260	0.03709504	154	0
## 4	OF	0.265	0.31475	0.3345	0.35300	0.411	0.3342500	0.02944394	120	0

We need a function to resample the data, we will use the `resample()` from the `mosaic` package.


```
library(broom)
```

```
f_boot <- function(x){  
  aov(obp~position,data=resample(x)) %>%  
  tidy() %>%  
  summarize(stat=meansq[1]/meansq[2]) %>%  
  pull()  
}
```

```
set.seed(541)  
results<-do(1000)*f_boot(mlb_obp)
```

Let's plot our sampling distribution.

```
results %>%
  gf_histogram(~f_boot) %>%
  gf_theme(theme_classic()) %>%
  gf_labs(title="Bootstrap sampling distribution of F test statistic",
          x="Test statistic")
```



Now the confidence interval for the F-statistic is:

```
cdata(~f_boot, data=results)
```

```
##           lower    upper central.p
## 2.5% 0.3546682 8.724895      0.95
```

We are 95% confident that the  $F$  statistic is in the interval (0.35, 8.72) which includes 1 so we fail to reject.

### 3. Test of variance

We have not performed a test of variance so we will create our own.

- a. Using the MLB from the lesson, subset on IF and OF.

```
mlb_prob3 <- mlb_obp %>%
  filter(position=="IF" | position=="OF") %>%
  droplevels()
```

```
summary(mlb_prob3)
```

```
## position      obp
## IF:154   Min.   :0.1740
## OF:120   1st Qu.:0.3100
##           Median :0.3310
##           Mean   :0.3327
##           3rd Qu.:0.3530
##           Max.   :0.4370
```

b. Create a side-by-side boxplot.

Boxplot showing the distribution of obp (y-axis, ranging from 0.20 to 0.45) across two positions (x-axis: IF and OF). The IF position shows a median obp of approximately 0.33, with a box spanning from 0.31 to 0.35 and whiskers extending from 0.27 to 0.35. The OF position shows a median obp of approximately 0.34, with a box spanning from 0.32 to 0.35 and whiskers extending from 0.27 to 0.35. Outliers are present for both positions.

position	obp (approximate values)
IF	0.18, 0.24, 0.29, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.40, 0.41, 0.42, 0.43, 0.44
OF	0.27, 0.28, 0.29, 0.30, 0.31, 0.32, 0.33, 0.34, 0.35, 0.36, 0.37, 0.38, 0.39, 0.40, 0.41, 0.42

$H_0: \sigma_{IF}^2 = \sigma^2 OF$ . There is no difference in the variance of on base percentage for infielders and outfielders.  
 $H_A: \sigma_{IF}^2 \neq \sigma^2 OF$ . There is a difference in variances.

```
mlb_prob3 %>%
  group_by(position) %>%
  summarize(stat=sd(obp))
```

```
## # A tibble: 2 x 2
##   position    stat
##   <fct>      <dbl>
## 1 IF         0.0371
## 2 OF         0.0294
```


```
## [1] 0.007651101
```

Let's write a function to shuffle the position.

```
perm_stat <- function(x){  
  x %>%  
  mutate(position=shuffle(position)) %>%  
  summarize(stat=sd(obp[position=="IF"])-sd(obp[position=="OF"])) %>%  
  pull()  
}
```

```
set.seed(443)  
results<-do(1000)*perm_stat(mlb_prob3)
```

```
results %>%  
  gf_histogram(~perm_stat) %>%  
  gf_vline(xintercept=obs,color="red") %>%  
  gf_theme(theme_classic()) %>%  
  gf_labs(title="Sampling distribution of difference in variances",  
          subtitle="Randomization permutation test",  
          x="Test statistic")
```



The p-value is

```
2*prop1(~(perm_stat>=obs),data=results)
```

```
## prop_TRUE  
## 0.04395604
```

This is a two sided test since we did not know in advance which variance would be larger. We reject the hypothesis of equal variance but the p-value is too close to the significance level. The conclusion is suspect. We need more data.


- d. Create a bootstrap distribution of the differences in sample standard deviations, and report a 95% confidence interval. Compare with part c.

Let's write a function.

```
var_stat <- function(x){  
  resample(x) %>%  
  summarize(stat=sd(obp[position=="IF"])-sd(obp[position=="OF"])) %>%  
  pull()  
}
```

```
set.seed(827)  
results<-do(1000)*var_stat(mlb_prob3)
```

```
results %>%  
  gf_histogram(~var_stat) %>%  
  gf_vline(xintercept=obs,color="red")%>%  
  gf_theme(theme_classic()) %>%  
  gf_labs(title="Bootstrap sampling of difference in variances",  
          x="Difference in variances")
```



## File Creation Information

- File creation date: 2020-10-26
- Windows version: Windows 10 x64 (build 18362)
- R version 3.6.3 (2020-02-29)
- mosaic package version: 1.7.0
- tidyverse package version: 1.3.0
- openintro package version: 2.0.0
- broom package version: 0.7.0