

Logistic Regression Applications Solutions

Lt Col Ken Horton

Professor Bradley Warner

11 August, 2020

Exercises

1. Possum classification

Let's investigate the `possum` data set again. This time we want to model a binary outcome variable. As a reminder, the common brushtail possum of the Australia region is a bit cuter than its distant cousin, the American opossum. We consider 104 brushtail possums from two regions in Australia, where the possums may be considered a random sample from the population. The first region is Victoria, which is in the eastern half of Australia and traverses the southern coast. The second region consists of New South Wales and Queensland, which make up eastern and northeastern Australia.

We use logistic regression to differentiate between possums in these two regions. The outcome variable, called `pop`, takes value `Vic` when a possum is from Victoria and `other` when it is from New South Wales or Queensland. We consider five predictors: `sex`, `head_l`, `skull_w`, `total_l`, and `tail_l`.

- a. Explore the data by making histograms of the quantitative variables, and bar charts of the discrete variables.
Are there any outliers that are likely to have a very large influence on the logistic regression model?
- b. Build a logistic regression model with all the variable.
- c. Using the p-values decide if you want to remove a variable(S) and if so build that model.
- d. For any variable you decide to remove, build a bootstrap 95% confidence interval.
- e. Explain why the remaining parameter estimates change between the two models.
- f. Write out the form of the model. Also identify which of the following variables are positively associated (when controlling for other variables) with a possum being from Victoria: `head_l`, `skull_w`, `total_l`, and `tail_l`.
- g. Suppose we see a brushtail possum at a zoo in the US, and a sign says the possum had been captured in the wild in Australia, but it doesn't say which part of Australia. However, the sign does indicate that the possum is male, its skull is about 63 mm wide, its tail is 37 cm long, and its total length is 83 cm. What is the reduced model's computed probability that this possum is from Victoria? How confident are you in the model's accuracy of this probability calculation?

File Creation Information

- File creation date: 2020-08-11
- Windows version: Windows 10 x64 (build 18362)
- R version 3.6.3 (2020-02-29)
- `mosaic` package version: 1.7.0
- `tidyverse` package version: 1.3.0
- `openintro` package version: 2.0.0