

Linear Regression Basics Applications Solutions

Lt Col Ken Horton

Professor Bradley Warner

29 July, 2020

Exercises

1. Nutrition at Starbucks

In the `openintro` package there is a data set called `starbucks`. Use it to answer the questions below.

- Create a scatterplot of between number of calories and amount of carbohydrates.

```
library(openintro)
```

```
## Loading required package: airports
```

```
## Loading required package: cherryblossom
```

```
## Loading required package: usdata
```

```
##
```

```
## Attaching package: 'openintro'
```

```
## The following object is masked from 'package:mosaic':
```

```
##
```

```
## dotPlot
```

```
## The following objects are masked from 'package:lattice':
```

```
##
```

```
## ethanol, lsegments
```

```
glimpse(starbucks)
```

```
## Rows: 77
```

```
## Columns: 7
```

```
## $ item      <chr> "8-Grain Roll", "Apple Bran Muffin", "Apple Fritter", "Ban...
```

```
## $ calories  <int> 350, 350, 420, 490, 130, 370, 460, 370, 310, 420, 380, 320...
```

```
## $ fat       <dbl> 8, 9, 20, 19, 6, 14, 22, 14, 18, 25, 17, 12, 17, 21, 5, 18...
```


```
## $ carb      <int> 67, 64, 59, 75, 17, 47, 61, 55, 32, 39, 51, 53, 34, 57, 52...
```

```
## $ fiber     <int> 5, 7, 0, 4, 0, 5, 2, 0, 0, 0, 2, 3, 2, 2, 3, 3, 2, 3, 0, 2...
```

```
## $ protein   <int> 10, 6, 5, 7, 0, 6, 7, 6, 5, 7, 4, 6, 5, 5, 12, 7, 8, 6, 0,...
```

```
## $ type      <fct> bakery, bakery, bakery, bakery, bakery, bakery, bakery, ba...
```

```
starbucks %>%
  gf_point(calories~carb) %>%
  gf_labs(x="Carbohydrate Content (g)",y="Calories") %>%
  gf_theme(theme_classic())
```



We put `calories` as the response.

- b. Describe the relationship in the graph.

There is a positive, moderate, linear association between number of calories and amount of carbohydrates. In addition, the amount of carbohydrates is more variable for menu items with higher calories, indicating non-constant variance. There also appear to be two clusters of data: a patch of about a dozen observations in the lower left and a larger patch on the right side. There might be some natural groupings of these points. For example, the points in the lower left might come from a *light* menu.

- c. In this scenario, what are the explanatory and response variables?


Response: number of calories. Explanatory: amount of carbohydrates (in grams).

- d. Why might we want to fit a regression line to these data?

With a regression line, we can predict the amount of calories for a given number of carbohydrates. This may be useful if you are concerned about your carb intake and its impact on calorie consumption. Typically you can get both on the menu so this model might not be that valuable.

- e. Create a scatterplot of between number of calories and amount of carbohydrates with the regression line included.

```
starbucks %>%
  gf_point(calories~carb) %>%
  gf_labs(x="Carbohydrate Content (g)",y="Calories") %>%
  gf_lm() %>%
  gf_theme(theme_classic())
```



f. Using 'lm()' fit a least squares line to the data.

```
star_mod <- lm(calories~carb,data=starbucks)
```

```
summary(star_mod)
```

```
##
## Call:
## lm(formula = calories ~ carb, data = starbucks)
##
## Residuals:
```

	Min	1Q	Median	3Q	Max
	-151.962	-70.556	-0.636	54.908	179.444

```
##
## Coefficients:
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	146.0204	25.9186	5.634	2.93e-07 ***
carb	4.2971	0.5424	7.923	1.67e-11 ***

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 78.26 on 75 degrees of freedom
## Multiple R-squared:  0.4556, Adjusted R-squared:  0.4484
## F-statistic: 62.77 on 1 and 75 DF, p-value: 1.673e-11
```

g. Report and interpret the slope coefficient.

The estimated slope is 4.297 so one additional gram of carbohydrates results in an average increase in calories of 4.297.

h. For a menu item with 51 g of carbs, what is the estimated calorie count?

```
146.0204+4.2971*51
```

```
## [1] 365.1725
```

- i. Could we use the model for a menu item with 100 g of carbs?

```
summary(starbucks)
```

```
##      item      calories      fat      carb
## Length:77      Min.   : 80.0    Min.   : 0.00    Min.   :16.00
## Class :character 1st Qu.:300.0    1st Qu.: 9.00    1st Qu.:31.00
## Mode  :character Median :350.0    Median :13.00    Median :45.00
##              Mean  :338.8    Mean  :13.77    Mean  :44.87
##              3rd Qu.:420.0    3rd Qu.:18.00    3rd Qu.:59.00
##              Max.   :500.0    Max.   :28.00    Max.   :80.00
##
##      fiber      protein      type
## Min.   :0.000    Min.   : 0.000    bakery      :41
## 1st Qu.:0.000    1st Qu.: 5.000    bistro box   : 8
## Median :2.000    Median : 7.000    hot breakfast: 8
## Mean   :2.221    Mean   : 9.481    parfait      : 3
## 3rd Qu.:4.000    3rd Qu.:15.000    petite       : 9
## Max.   :7.000    Max.   :34.000    salad        : 1
##              sandwich   : 7
```

The maximum carb value is 80 so 100 is outside of the observed data. It would be suspect to extrapolate to that value.

- j. Does the assumption of constant variance seem reasonable for this problem?

It seems that the variance in the second group is larger than the first, so it may not be a reasonable assumption.

- k. Verify that the line passes through the mean carb and mean calories, do this mathematically.

```
146.0204+4.2971*44.87
```

```
## [1] 338.8313
```

It checks.

- l. What is the estimate of the standard deviation of the residuals? How could you use this information?

The estimate is 78.26. If the normal assumption is accurate, we would expect a majority of observations to be within ± 78.26 calories of the line.

File Creation Information

- File creation date: 2020-07-29
- Windows version: Windows 10 x64 (build 18362)
- R version 3.6.3 (2020-02-29)
- `mosaic` package version: 1.7.0
- `tidyverse` package version: 1.3.0
- `openintro` package version: 2.0.0
- `broom` package version: 0.7.0