

Multivariate Optimization

$$z = f(x_1, x_2, \dots, x_n)$$


Gradient

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \\ \dots \\ \frac{\partial f}{\partial x_n} \end{bmatrix}$$

Hessian

$$\nabla^2 f = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \dots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \dots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \dots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

$$z = \sqrt{x_1^2 + x_2^2}$$





- Gradient of a function at a point is orthogonal to the contours
- Gradient points in the direction of greatest increase of the function
- Negative gradient points in the direction of the greatest decrease of the function
- Hessian is a symmetric matrix

Unconstrained multivariate optimization - Descent direction and movement

- Iterative

$$x^{k+1} = x^k + \alpha^k s^k$$


Starting point Step length Search direction




- In ML techniques, this is called as the learning rule
- In neural networks
 - Back-propagation algorithm
 - Same gradient descent with application of chain rule
- In clustering
 - Minimization of an Euclidean distance norm

Steepest descent and optimum step size

- Minimize $f(x_1, x_2, \dots, x_n) = f(x)$
- **Steepest descent**
 - At iteration k starting point is x^k
 - Search direction $s^k =$ Negative of gradient of $f(x) = -\nabla f(x^k)$
 - New point is $x^{k+1} = x^k + \alpha^k s^k$ where α^k is the value of α for which $f(x^{k+1}) = f(\alpha)$ is a minimum (univariate minimization)



$$f(X) = 4x_1^2 + 3x_1x_2 + 2.5x_2^2 - 5.5x_1 - 4x_2$$



$$f'(X) = \begin{bmatrix} 8x_1 + 3x_2 - 5.5 \\ 3x_1 + 5x_2 - 4 \end{bmatrix}$$

Learning parameter (α) = 0.135

Initial guess (X_0) = $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ $f(X_0) = 19$

Step 1: $X_1 = X_0 - \alpha f'(X_0)$ ← Gradient Descent (or)
Learning Rule in ML

$$X_1 = \begin{bmatrix} 2 \\ 2 \end{bmatrix} - 0.135 \begin{bmatrix} 8x_{0,1} + 3x_{0,2} - 5.5 \\ 3x_{0,1} + 5x_{0,2} - 4 \end{bmatrix}$$


$$X_1 = \begin{bmatrix} 2 \\ 2 \end{bmatrix} - 0.135 \begin{bmatrix} 8(2) + 3(2) - 5.5 \\ 3(2) + 5(2) - 4 \end{bmatrix}$$

$$X_1 = \begin{bmatrix} -0.2275 \\ 0.3800 \end{bmatrix} \quad f(X_1) = 0.0399$$

Constant objective function contour plots

$$f(X) = 4x_1^2 + 3x_1x_2 + 2.5x_2^2 - 5.5x_1 - 4x_2 = K$$

Quadratic in this case - ellipse




First iteration $(X_1) = \begin{bmatrix} -0.2275 \\ 0.3800 \end{bmatrix}$

Step 2: $X_2 = X_1 - \alpha f'(X_1)$

$$X_2 = \begin{bmatrix} -0.2275 \\ 0.3800 \end{bmatrix} - 0.135 \begin{bmatrix} 8x_{1,1} + 3x_{1,2} - 5.5 \\ 3x_{1,1} + 5x_{1,2} - 4 \end{bmatrix}$$

$$X_2 = \begin{bmatrix} -0.2275 \\ 0.3800 \end{bmatrix} - 0.135 \begin{bmatrix} 8(-0.2275) + 3(0.3800) - 5.5 \\ 3(-0.2275) + 5(0.3800) - 4 \end{bmatrix}$$

$$X_2 = \begin{bmatrix} 0.6068 \\ 0.7556 \end{bmatrix} \quad f(X_2) = -2.0841$$




Second iteration $(X_2) = \begin{bmatrix} 0.6068 \\ 0.7556 \end{bmatrix}$

Step 3: $X_3 = X_2 - \alpha f'(X_2)$

$$X_3 = \begin{bmatrix} 0.6068 \\ 0.7556 \end{bmatrix} - 0.135 \begin{bmatrix} 8x_{2,1} + 3x_{2,2} - 5.5 \\ 3x_{2,1} + 5x_{2,2} - 4 \end{bmatrix}$$

$$X_3 = \begin{bmatrix} 0.6068 \\ 0.7556 \end{bmatrix} - 0.135 \begin{bmatrix} 8(0.6068) + 3(0.7556) - 5.5 \\ 3(0.6068) + 5(0.7556) - 4 \end{bmatrix}$$

$$X_3 = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix} \quad f(X_3) = -2.3342$$



Third iteration $(X_3) = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix}$

Step 4: $X_4 = X_3 - \alpha f'(X_3)$


$$X_4 = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix} - 0.135 \begin{bmatrix} 8x_{3,1} + 3x_{3,2} - 5.5 \\ 3x_{3,1} + 5x_{3,2} - 4 \end{bmatrix}$$

$$X_4 = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix} - 0.135 \begin{bmatrix} 8(0.3879) + 3(0.5398) - 5.5 \\ 3(0.3879) + 5(0.5398) - 4 \end{bmatrix}$$

$$X_4 = \begin{bmatrix} 0.4928 \\ 0.5583 \end{bmatrix} \quad f(X_4) = -2.3675$$

Optimal solution $(X_{opti}) = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} \quad f(X_{opti}) = -2.3750$

Gradient is zero at the optimum point



Third iteration $(X_3) = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix}$ ✓

Handwritten notes in red:

$$\begin{aligned} X^{k+1} &\leftarrow X^k \\ f(X^{k+1}) &\leftarrow f(X^k) \\ \nabla f(X^{k+1}) &\leftarrow \nabla f(X^k) \end{aligned}$$

Step 4: $X_4 = X_3 - \alpha \nabla f(X_3)$

$$X_4 = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix} - 0.135 \begin{bmatrix} 8x_{3,1} + 3x_{3,2} - 5.5 \\ 3x_{3,1} + 5x_{3,2} - 4 \end{bmatrix}$$

$$X_4 = \begin{bmatrix} 0.3879 \\ 0.5398 \end{bmatrix} - 0.135 \begin{bmatrix} 8(0.3879) + 3(0.5398) - 5.5 \\ 3(0.3879) + 5(0.5398) - 4 \end{bmatrix}$$

$$X_4 = \begin{bmatrix} 0.4928 \\ 0.5583 \end{bmatrix} \quad f(X_4) = -2.3675$$

Optimal solution $(X_{opti}) = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix} \quad f(X_{opti}) = -2.3750$

Gradient is zero at the optimum point

