		L,I	L,D = 1,1		L,D =	= 2,2	L,D = 2,2 L,D = 3,3 L,D = 4,4 L,D = 5,5 L,D = 6,6	3,3	L,D =	4,4	L,D	= 5,5	L,	0 = 0	3,6
	F	0.003	0.003	0.002	0.4 0.3	0.3	0.003 0.003 0.002 0.4 0.3 0.3 1.9 1.6 1.1 3.8 3.4 2.1 5.7 5.2 2.9 7.6 7.1	1.1	3.8 3.4	2.1	5.7 5.	2 2.9	9.7	7.1	3.4
$(x_i) \mapsto (\exp(x_i))$	Me	0.002	0.002	0.002	0.002 0.2 0.2	0.2	1.1 1.0	9.0	2.2 2.1	1.0	3.1 3.0 1	1.1	1.1 4.1 3.9	3.9	1.2
K = 100	Mo	0.002	0.002	0.002	$0.002 \mid 0.2 \mid 0.2$	0.1	0.1 1.0 0.9 0.5 2.2 2.1 0.9 48.1	0.5	2.2 2.1	0.0	48.1 48	1 48.7 23.0	23.0 48.1 46.5	46.5	16.8
N = 50000	Yme	0.003	0.003	0.002	0.002 0.4 0.3	0.2	0.2 1.7 1.5	6.0	0.9 3.1 2.9 1.4 4.7	1.4	4.7 4.4	4 1.9	1.9 6.4 6.3	6.3	2.2
Local = 100	Ymo	0.01	0.005	0.009	0.005 0.009 $ 1.1 0.7 $	1.1	1.1 4.7 3.3	4.4	4.4 7.7 6.1 6.0 77.4 74.8	6.0	77.4 74	8 39.9	39.9 77.5 72.4	72.4	30.1
	$^{\mathrm{Ap}}$	0.003	0.003	0.002	0.003 0.002 0.3 0.3		0.2 1.5 1.4	0.7	0.7 2.9 2.8 1.1 76.2 72.7 42.9 76.0 70.1 32.2	1.1	76.272	7 42.9	0.92	70.1	32.2
	>	100.	0 - 100	0:	100.0 -	100.0	100.0 - 100.0 100.0 - 100.0 100.0 - 100.0 100.0 - 99.7 99.6 - 99.2	100.0	100.0 -	99.7	9.66	99.2		98.2 - 97.1	7.1

Table 1: **Influence de la dimension**. La même fonction générique est apprise et inversée pour différentes dimensions.