

## Research Medical Engineering—Review

# Nanotechnology and Nanomedicine: A Promising Avenue for Lung Cancer Diagnosis and Therapy



Wei Yin <sup>a,b</sup>, Feng Pan <sup>c</sup>, Junjie Zhu <sup>c</sup>, Junwu Xu <sup>d</sup>, Diego Gonzalez-Rivas <sup>c,e</sup>, Meinoshin Okumura <sup>f</sup>, Zhiyong Tang <sup>g</sup>, Yang Yang <sup>c,\*</sup>

<sup>a</sup> Key Laboratory of Oral Biomedical Engineering (Wuhan University), Ministry of Education, Hospital of Stomatology, School of Stomatology, Wuhan University, Wuhan 430079, China

<sup>b</sup> Department of Biomedical Data Science, Geisel School of Medicine, Dartmouth College, Hanover, NH 03756, USA

<sup>c</sup> Department of Thoracic Surgery, Shanghai Pulmonary Hospital, School of Medicine, Tongji University, Shanghai 200433, China

<sup>d</sup> Tus-Medical Health Technology Investment (Jiaxing) Co., Ltd., Jiaxing 314033, China

<sup>e</sup> Minimally Invasive Thoracic Surgery Unit (UCTMI), Hospital San Rafael, Coruña 15006, Spain

<sup>f</sup> Osaka Toneyama Medical Center, Osaka 560-8552, Japan

<sup>g</sup> CAS Key Laboratory of Nanosystem and Hierarchical Fabrication, National Center for Nanoscience and Technology, Beijing 100190, China

## ARTICLE INFO

### Article history:

Received 11 December 2019

Revised 1 April 2020

Accepted 24 April 2020

Available online 7 October 2021

### Keywords:

Nanomedicine

Lung cancer

Diagnosis

Therapy

## ABSTRACT

Lung cancer is a leading cause of cancer-related death worldwide, with a very poor overall five-year survival rate. The intrinsic limitations associated with the conventional diagnosis and therapeutic strategies used for lung cancer have motivated the development of nanotechnology and nanomedicine approaches, in order to improve early diagnosis rate and develop more effective and safer therapeutic options for lung cancer. Cancer nanomedicines aim to individualize drug delivery, diagnosis, and therapy by tailoring them to each patient's unique physiology and pathological features—on both the genomic and proteomic levels—and have attracted widespread attention in this field. Despite the successful application of nanomedicine techniques in lung cancer research, the clinical translation of nanomedicine approaches remains challenging due to the limited understanding of the interactions that occur between nanotechnology and biology, and the challenges posed by the toxicology, pharmacology, immunology, and large-scale manufacturing of nanoparticles. In this review, we highlight the progress and opportunities associated with nanomedicine use for lung cancer treatment and discuss the prospects of this field, together with the challenges associated with clinical translation.

© 2021 THE AUTHORS. Published by Elsevier LTD on behalf of Chinese Academy of Engineering and Higher Education Press Limited Company. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Despite recent advances in lung cancer detection and treatment, lung cancer remains the most lethal cancer worldwide, due to the failure to detect cancer occurrence early and the lack of effective treatments for advanced-stage patients. Lung cancer is the most common newly diagnosed cancer type in North America and Asia [1,2]. The primary difficulty associated with lung cancer detection is that existing detection methods, including bronchial biopsy and computed tomography (CT), depend heavily on tumor size and require specific medical equipment, which is often associated with high costs. Nanotechnology offers the promise of new detection approaches, as nanoparticle (NP) surfaces

can be modified to bind to overexpressed receptors in tumor cells, which can act as cancer imaging contrast agents, to increase the sensitivity and specificity of cancer detection methods [3,4]. In addition, microfluidic arrays and array-based sensing methods that use NPs are promising and novel cancer diagnostic approaches, with ultralow detection thresholds, short assay times, high-throughput capabilities, and low sample consumption [5,6].

NPs can also be used to improve lung cancer treatment. Precision nanomedicines, which have unique properties, including nanoscale sizes, high surface-to-volume ratios, and favorable physicochemical characteristics, could potentially be used to modulate the pharmacokinetic and pharmacodynamic profiles of cancer drugs, enhancing their therapeutic indexes. These characteristics are necessary properties of current precision medicines (PMs) [7–10] because many PMs, such as nucleic acid-based therapies and antibodies, suffer from poor targeting abilities and

\* Corresponding author.

E-mail address: [timyangsh@tongji.edu.cn](mailto:timyangsh@tongji.edu.cn) (Y. Yang).

plasma stability, suboptimal pharmacokinetic properties, and immunological toxicities, which have prohibited their clinical translation [11,12].

An increasing number of studies have indicated the potential benefits of precision nanomedicine techniques for the early diagnosis and targeted therapy of lung cancer. Therefore, a critical review is necessary to provide a more complete understanding of these new strategies.

## 2. Personalized diagnosis of lung cancer

### 2.1. Nanotechnology for the *in vivo* diagnosis of lung cancer

Successful early diagnosis of lung cancer can improve survival rates. Conventional medical imaging technologies, such as magnetic resonance-guided focused ultrasound surgery (MRgFUS), are currently limited by the insensitivity of magnetic resonance imaging (MRI) for the visualization of small tumors [13]. NPs can be used as imaging contrast agents to increase the resolution and improve the anatomic definition of lesions. Recently, Wang et al. [14] constructed an active-targeting nanosized, theranostic, superparamagnetic iron oxide (SPIO) platform to increase the imaging sensitivity and energy-deposition efficiency of a clinical MRgFUS system. The surfaces of these polyethylene glycol (PEG)-ylated SPIO NPs were decorated with anti-epidermal growth factor receptor (EGFR) monoclonal antibodies, for the targeted delivery of NPs to EGFR-overexpressing lung cancer cells. The researchers demonstrated that using these NPs significantly improved MRI sensitivity for the visualization of EGFR-overexpressing lung cancer cells in a rat model.

Quantum dots (QDs) are semiconductor nanocrystals (2–100 nm in size), with unique optical and electrical properties. Compared with organic dye molecules, the bright fluorescence and high photochemical endurance of QDs make them promising for use in fluorescence imaging approaches [15]. Near-infrared (NIR)-emitting QDs, which exhibit high molar excitation coefficients, are particularly well-suited for *in vivo* whole-body imaging techniques because NIR light penetrates the body more deeply than light in the visible spectrum. The Papagiannaros et al. [16] prepared a tumor-targeted NIR-imaging agent composed of a cancer-specific monoclonal anti-nucleosome antibody, 2C5, coupled with QD-containing polymeric micelles. They demonstrated that this fluorescent imaging molecule exhibited excellent imaging properties, with a tumor fluorescence intensity 1 h after injection that was two-fold that of nontargeted, QD-loaded PEG–polyethylene micelles.

In addition to exploring the development of a single, powerful imaging modality, the development of multimodal approaches has been attempted, through the integration of several imaging technologies, in order to overcome the individual shortcomings of each technique [17]. For example, Xiao et al. [18] incorporated gadolinium-doped mesoporous silica nanoparticles (MSNs) and gold nanoparticles (AuNPs) into a single nanosystem ( $\text{Gd}_2\text{O}_3/\text{MCM-41@Au}$ ) and found that  $\text{Gd}_2\text{O}_3/\text{MCM-41@Au}$  served as an efficient MRI contrast agent during cancer imaging and successfully targeted EGFR molecules, for surface-enhanced Raman scattering (SERS) detection.

### 2.2. Nanotechnology for the *in vitro* diagnosis of lung cancer

Accurate diagnoses require the determination of how and where to collect biopsy tissue samples. Critical advances in molecular biology have allowed the capture and analysis of tumor-derived substances from body fluids [19,20]. The identification of clinically relevant alterations is now possible at the DNA, RNA,

and protein levels [21]. Therefore, physicians have multiple biopsy options when evaluating lung cancer patients, including lesions in the lungs, peripheral blood, and pleural effusion (PE). For population screening, noninvasive analyses such as peripheral blood analyses are preferable because tissues may not be available. In recent years, nanotechnology has made great contributions toward achieving the high sensitivity and high specificity that are necessary for accurate diagnoses. NPs possess large surface areas that can be linked with multiple diagnostic agents, improving the efficiency and sensitivity of diagnoses. Compared with traditional polymerase chain reaction (PCR)-based sequencing techniques, microfluidic (lab-on-a-chip) technologies and array-based sensing that use NPs are attractive alternatives for cancer diagnoses.


#### 2.2.1. Tumor-derived DNA in peripheral blood

Extracellular DNA detected in the peripheral blood that shares genetic information with the host is referred to as cell-free DNA (cfDNA) [22], and cfDNA that originates from tumor cells is termed circulating tumor DNA (ctDNA). Currently, the exact mechanism through which ctDNA is released into the blood remains unknown. ctDNA is thought to be released from apoptotic and necrotic tumor cells or secreted from live tumor cells [23]. Cancer patients present increased amounts of both cfDNA and ctDNA compared with healthy controls [24]. Because ctDNA carries the genomic variations and heterogeneity that have emerged during tumor evolution, ctDNA monitoring presents an obvious possible avenue for monitoring tumor status, progression, and the occurrence of treatment resistance in cancer patients.

To achieve the goal of using ctDNA to monitor tumor progression, ctDNA must first be enriched from peripheral blood samples. Both cfDNA and ctDNA in peripheral blood have low molecular weights, which require improved enrichment techniques because conventional methods for DNA isolation and extraction from peripheral blood are better suited for DNA molecules with medium and high molecular weights. Furthermore, the proportion of ctDNA in cfDNA varies widely, depending on individual variations. Therefore, isolation approaches are also necessary to distinguish ctDNA from cfDNA. Fortunately, technological advances have already facilitated the enrichment of rare ctDNA [25,26]. Nanomaterials have already been developed to enrich ctDNA using specific markers, such as epithelial cell adhesion molecule (EpCAM) and cytokeratin (CK).

Biocompatible AuNPs [27], in combination with polymer beads [28] and immunomagnetic beads [29], have been investigated for this scheme. In addition, nanostructure substrates that offer increased surface areas, such as nanotubes, nanopillars, nanowires, and nanotextured surfaces, could facilitate increased interactions with biomolecules, resulting in biosensing platforms that are capable of capturing or isolating ctDNA. For example, after being functionalized with an aptamer, nanotopographic substrates were able to selectively capture more than 90% of cancer cells [30]. Nanostructured electric materials [31], such as the electroactive conducting polymer polypyrrole [32] and gold (Au) nanowires coated with polypyrrole [33], have also exhibited high efficiency levels for the isolation of ctDNA, with high yields and purities.

Early lung cancer detection is just one potential application of ctDNA analysis. ctDNA can reveal genetic variations associated with lung cancer and increase the reliability of diagnoses. Thus far, several researchers have successfully enriched ctDNA for the performance of molecular genotyping analyses in lung cancer patients [34,35]. A silicon nanowire substrate (SiNS) embedded in a microfluidic chip (Fig. 1), combined with the rationally designed, cell-based, systematic evolution of ligands by exponential enrichment (SELEX)-derived aptamers, has successfully enhanced the differential capture of circulating tumor cells from non-small cell lung cancer (NSCLC) patients [36].



**Fig. 1.** Schematic depiction of a microfluidic, circulating tumor cell chip, composed of an aptamer-grafted silicon nanowire substrate (SiNS) and an overlaid polydimethylsiloxane (PDMS) chaotic mixer. Reproduced from Ref. [36] with permission of Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany ©2016.

### 2.2.2. Tumor-derived DNA in PE

The emergence of PE is a common clinical manifestation of lung cancer. This excess fluid, which accumulates in the pleural cavity, contains tumor-derived DNA. Sampling PE is a practical approach to analyze the molecular profiles of lung cancer lesions [37]. Furthermore, in some patients, PE is the initial symptom of disease, and thus provides an early opportunity to detect tumor-derived DNA.

The enrichment of tumor-derived DNA from PE can be difficult because PE often contains various inflammatory and mesothelial cells. Existing strategies for isolating tumor-derived DNA rely on the macro-dissection, manual micro-dissection, or laser-capture micro-dissection of smears or cell blocks. Several successful cases using these strategies have been previously described [38,39].

### 2.2.3. Exosomes

Exosomes range from 30 to 150 nm in size and are released during the fusion between multivesicular endosomes (MVEs) and the plasma membrane [40]. Exosomes can alter the functions of various cells by regulating cell communications. A previous study indicated that exosomes may be involved in the oncogenic process, through the regulation of tumor immune responses, pre-metastatic niches, and chemotherapeutic resistance [41]. Thus, exosomes are potential specific targets for diagnosis and therapeutic interventions. In addition, exosomes have been identified and enriched from peripheral blood [42] and may, therefore, be clinically applicable specimens for cancer analysis.

Current exosome isolation methods, such as sucrose gradient ultracentrifugation and ultrafiltration, rely heavily on multiple-step ultracentrifugation processes, which are tedious and time-consuming, as well as having low efficiency for discriminating exosomes from other biomaterials. Similar to existing techniques for the isolation and enrichment of ctDNA, microfluidic technologies are being developed to capture exosomes. These technologies have the advantages of small sample volume requirements, low costs, short operation times, and high sensitivity [43]. The capture of targeted exosomes depends on the use of microfluidic channels that have been functionalized with antibodies, in order to enhance their affinities for targeted exosomes. For example, He et al. [44] developed an integrated microfluidic exosome analysis platform by combining a magnetic bead-based strategy with a multi-step

analysis (Fig. 2). This device was able to successfully assess the total expression and phosphorylation levels of insulin-like growth factor 1 receptor (IGF-1R) in NSCLC patients by directly probing plasma exosomes. In addition to specific antibodies, aptamers [45], PEG-ylated lipids [46], and graphene oxide/polydopamine (GO/PDA) nano-interfaces [47] have demonstrated satisfactory performances as ligands. Aptamers are particularly stable in various solutions, including a wide range of salt concentrations, and under ionic and denaturing conditions. Furthermore, cancer cells can be sorted according to physical plasticity and diameter.

## 3. Precision nanomedicine for lung cancer

Compared with chemotherapy, which inhibits the proliferation of all cells, targeted cancer therapy specifically and accurately enacts tumor suppression by targeting the specific molecules involved in oncogenesis. The efficient and precise delivery of drugs to target lesions is a critical factor for the success of targeted treatments.


### 3.1. Drug delivery

The unique physicochemical properties of nanomaterials have made them premier options as both drugs and drug delivery systems (DDSs) for the targeted treatment of cancer [48]. Cancers demonstrate irregular cell growth, aided by the development of new blood vessel networks that are highly porous, with large spaces between endothelial cells. Taking advantage of the anatomical and pathophysiological differences between normal and tumor tissues, nanodrugs are designed to circulate in the blood for long periods and to accumulate at tumor sites through the enhanced permeability and retention (EPR) effect [49]. In addition, nano-based DDSs have received attention for their potential to overcome problems associated with the solubility, stability, diffusivity, blood circulation half-time, and immunogenicity of chemotherapy drugs and to improve the specificity of drug release during cancer treatment [50]. At present, nano-based DDSs include conventional liposomes, polymer NPs, dendritic polymers, and micelles, in addition to inorganic nanomaterials such as AuNPs, MSNs, and metal–organic frameworks. Furthermore, the favorable physicochemical, biochemical, and electrical properties of inorganic NP-based imaging contrast agents have resulted in improved sensitivity for positron emission tomography (PET), MRI, and single-photon emission computed tomography (SPECT) platforms, facilitating real-time observations of cancer progression during the treatment process.

#### 3.1.1. Gold nanoparticles

The confirmed driver oncogene associated with NSCLC is a top potential treatment target. During recent decades, inorganic NPs have been widely investigated for use as drugs or DDSs in targeted cancer treatment [51], and AuNPs have attracted particular attention due to their unique optical properties, low toxicity, and the ease with which they can be prepared and functionalized. To achieve the maximum accumulation of NPs in tumor tissues and to increase efficiency, AuNPs have been modified with various active ligands. AuNPs bound to the anti-cancer drug methotrexate (MTX) have demonstrated high levels of tumor retention and enhanced therapeutic efficacy in a Lewis lung carcinoma mouse model, compared with an equal dose of free MTX, which may be attributed to the “concentrated effect” of MTX–AuNPs [52]. EGFR is a cell-surface receptor that is overexpressed in several tumor types, including NSCLC. In recent years, EGFR-targeted antibodies have become a popular targeting strategy for NSCLC treatment. Yokoyama et al. [53] have reported that Clone 225 antibody-






**Fig. 2.** Integrated microfluidic exosome analysis for NSCLC patients. (a) A PDMS chip, containing a microchannel network for exosome analysis; (b) integration of the streamlined, lab-on-a-chip, immunomagnetic isolation of exosomes, exosome lysis, protein capture, and intravesicular protein analysis. Reproduced from Ref. [44] with permission of the Royal Society of Chemistry, ©2014.

conjugated hybrid plasmonic magnetic NPs (C225-AuFe NPs) exhibited enhanced antitumor activity through the induction of apoptosis and autophagy. Furthermore, a novel radioimmunotherapeutic agent,  $^{131}\text{I}$ -C225-AuNPs-PEG, was successfully synthesized (Fig. 3) and exhibited enhanced endocytosis and cytotoxicity against high-EGFR-expressing human A549 lung carcinoma cells; it also actively targeted an A549 tumor xenograft in a mouse model [54]. AuNPs have exhibited enhanced internalization via antibody-mediated endocytosis. Due to the strong and selective

interaction that occurs between gold and sulfur groups, the use of sulfur-containing targeted ligands, which are chemisorbed onto the NP surface via sulfur bonds, can improve efficiency [55]. AuNPs are also ideal carriers of micro-RNAs (miRNAs), which can further function as specific inhibitors [56].

### 3.1.2. Mesoporous silica nanoparticles

Due to their large surface areas, large pore volumes, and high levels of biochemical and physicochemical stability, MSNs have



**Fig. 3.** Scheme depicting the radioiodination of C225-AuNPs-PEG using the iodogen method. Reproduced from Ref. [54] with permission of Elsevier Ltd., ©2013.

attracted substantial interest as potential DDSs [57,58]. MSN-coated gold nanorods (AuNRs) loaded with doxorubicin (DOX) have been designed to act as light-mediated, multifunctional, theranostic carriers for lung cancer treatments, and have been shown to result in enhanced cancer cell death, due to the synergistic effects of chemotherapy and hyperthermia-based treatments [59]. A tumor microenvironment-cascade pH-responsive DDS was constructed by functionalizing hollow MSNs (HMSNs) using  $\beta$ -cyclodextrin ( $\beta$ -CD) and adenosine deaminase (Ada)-PEG, and further loading them with DOX (HMSNs- $\beta$ -CD/Ada-PEG@DOX). This approach effectively resolved the “PEG dilemma” and facilitated the specific release of loaded drugs in cancer cells, inducing cell apoptosis and inhibiting tumor growth with minimal toxic side effects (Fig. 4) [60]. EGFR monoclonal antibody-capped MSNs have been demonstrated to specifically target EGFR-mutant lung cancer cells and to efficiently release loaded drugs in the cancer cells. The release speed of therapeutic reagents from MSNs can also be adjusted by developing a smart nanomedicine system.

### 3.1.3. Nanoscale coordination polymers and nanoscale metal–organic frameworks

Nanoscale coordination polymers (NCPs) and nanoscale metal–organic frameworks (NMOFs), which are constructed from the self-assembly of metal-connecting points and organic bridging ligands, have also been developed as DDSs for cancer therapy, due to their tunable compositions, sizes, and shapes; ease of surface modification; high drug-loading capacities; and intrinsic biodegradability properties. Several reports have focused on the application of NCPs and NMOFs as DDSs. Lipid-coated and anisamide-targeted NCPs displayed enhanced cytotoxicity against human lung cancer cells, compared with as-synthesized particles or free bisphosphonates [61]. The Liu et al. [62] further constructed zinc (Zn)-bisphosphonate NCPs carrying either  $48\% \pm 3\%$  (weight percent) cisplatin prodrug or  $45\% \pm 5\%$  (weight percent) oxaliplatin prodrug. The NCPs were PEG-ylated to further stabilize the particles and to inhibit the burst release of drugs. Both NCPs demonstrated enhanced antitumor activities when compared with free drugs in three different tumor models. Folate (Fol)-targeted calcium


zoledronate (CaZol) NMOFs have been fabricated by incorporating Fol-targeted ligands into CaZol NMOFs, which possess excellent chemical and colloidal stability under physiological conditions. The encapsulated zoledronate was released from the NMOFs in mid-endosomes during endocytosis and exhibited increased efficiency for the inhibition of NSCLC proliferation and the induction of apoptosis, compared with small-molecule zoledronate [63].

### 3.2. Nanotheranostics

Recently, the early diagnosis and targeted therapeutic results associated with the use of nanomedicines during lung cancer have encouraged scientists to explore “nanotheranostics,” which are sub-micrometer-sized carrier materials that contain both drugs and imaging agents within a single formulation [64,65]. Nanotheranostics hold the potential to contribute to the development of personalized approaches to cancer management. Common diagnostic agents include SPIOs, QDs, radionuclides, and heavy elements, such as iodine. An ideal theranostic nanomedicine would recognize a specific target, bind to specific receptors on the targeted cell membrane, diagnose the cancer morphology, and provide effective therapy, while simultaneously possessing biocompatibility and biodegradability. Nanotheranostics are expected to provide practical solutions for cancer treatments and cures during the early stages of lung cancer.

The first example of theranostic nanomedicine, which combined chemotherapeutic and photothermal therapy (PTT) *in vivo*, was constructed by conjugating Taxol-loaded poly(lactide-co-glycolic acid) (PLGA) NPs with iron oxide NPs and QDs [66]. QD/Fe<sub>3</sub>O<sub>4</sub>/Taxol-loaded PLGA NPs can potentially serve as contrast agents for MRIs, and AuNRs can convert NIR light into heat, in order to simultaneously achieve the photothermal ablation of tumor tissue and destroy spherical PLGAs, efficiently releasing encapsulated Taxol. *In vivo*, AuNR/QD/Fe<sub>3</sub>O<sub>4</sub>/Taxol-loaded PLGA NPs were intratumorally injected into transplanted tumors in mice, resulting in the progressive reduction of tumor volumes.

The Jing et al. [67] successfully fabricated a robust theranostic cerasome named ICG@DPDCs-<sup>177</sup>Lu for NIR fluorescence imaging



**Fig. 4.** Schematic illustration showing the drug delivery process of the HMSNs- $\beta$ -CD/Ada-PEG system in a tumor microenvironment, *in vivo*. CTAB: *N*-cetyltrimethylammonium bromide; HepG2: human hepatocellular carcinoma. Reproduced from Ref. [60] with permission of Elsevier Ltd., ©2016.

and the photothermal ablation of cancer cells by encapsulating indocyanine green (ICG) in 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine-*N*-[carboxy(polyethylene glycol)2000]-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid monoamide (DSPE-PEG<sub>2000</sub>-DOTA) and then chelating the radioisotope <sup>177</sup>Lu. ICG@DPDCs-<sup>177</sup>Lu can be used as an attractive radionuclide cancer therapy; *in vivo*, the ICG@DPDCs effectively ablated cancer cells through photothermal effects (Fig. 5).

#### 4. Improvements in traditional lung cancer diagnosis methods using nanomedicine

Traditionally, histological examinations of resected tumors have been necessary in order to make reliable diagnoses. For pre-clinical diagnoses, CT and MRI are the most commonly used approaches. However, the use of nontargeted contrast agents during diagnosis has intrinsic limitations, such as low sensitivity and specificity, which can affect accurate tumor localization. NPs have been synthesized to overcome these problems. By conjugating NPs with other moieties, which act as markers, the morphologies of tumors can be more clearly delineated [68].

Inorganic nanomaterials have long been used for DDSs, imaging, tumor treatment, diagnosis, and prognosis. The most commonly employed materials are gold, silver (Ag), silica, and iron oxides. Several studies have focused on AuNPs, due to their biocompatible, multifunctional, and theranostic properties. For example, silica-gold nanoshells, modified with PEG, were synthesized for use during PTT, which can be applied to solid tumors using NIR light [69]. Knights and McLaughlan [70] demonstrated the size-dependent effects of AuNRs on both the photoacoustic (PA) imaging response and pulsed-wave photothermal therapeutic efficacy. A study demonstrated the *in vivo* lung cancer antitumor activity of silver nanoparticles (AgNPs) against lung cancer cells and a xenograft mouse model [71]. However, the toxicity of inorganic nanomaterials can result in harmful impacts on normal cells. Because the mononuclear phagocyte system (MPS) is responsible for the clearance of drugs, non-biodegradable inorganic NPs and toxic macromolecules can be difficult to clear from the human body. Cytotoxicity tests are usually performed using trypan blue and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays.

To overcome the difficulties associated with inorganic NPs, biocompatible organic nanomaterials and ultrasmall biodegradable nanomedicines have been developed [72,73]. Ultrasmall SPIOs (USPIOs), for example, are biodegradable NPs that are smaller than 50 nm, which allows them to pass through even the smallest blood vessels and remain in circulation [74,75]. In addition, these NPs are nontoxic to major MPS organs, such as the bone marrow, spleen,


and kidney, and are easily cleared from the body; thus, these novel nanomaterials have a potentially bright future.

#### 5. Improvements in traditional lung cancer treatment methods, using nanomedicine

Several lung cancer treatments are currently available, including surgical resections, chemotherapy, and radiation; however, none of these options are optimal for lung cancer therapy [76]. Although the surgical resection of lung cancer is a relatively effective treatment method, by the time most lung cancers are diagnosed, the patients have often reached an advanced stage of lung cancer or metastasis has occurred, which can make operations difficult and sometimes unfeasible to perform. The differentiation between tumors and surrounding normal tissues can be vague, which poses a great challenge during surgery and increases the likelihood of either overtreatment or relapse. Multiple-drug resistance (MDR) is a major problem associated with chemotherapy, and most chemotherapeutic drugs are toxic to both tumor cells and normal tissues. Radiation can cause serious side effects in patients and reduce their quality of life.

The surfaces of NPs can be modified with fluorescent dyes, hydrophilic ligands, and specific molecules that have affinities for unique proteins that are only expressed on the tumor surface; these fluorescently tagged nanomaterials can then be used to guide surgery [77,78]. NPs can increase the concentration of chemotherapeutic drugs near tumors, either through the EPR effect or through active-target delivery [79,80]. Angiogenic blood vessels found in tumors differ from their normal counterparts due to the presence of gaps between adjacent endothelial cells, which are large enough to induce the EPR effect. Active targeting involves the conjugation to the NP surface of either a targeting ligand or antibodies that specifically recognize tumor cells. Many factors, including the size, charge, surface modifications, and angiogenesis of the tumor, the tumor microenvironment, and the half-lives of NPs, can affect the final accumulation of NPs in tumors. Several liposomal NPs have been used during clinical treatments, including the most popular liposome formulation, DOX; some are available on the market, including liposomal daunorubicin and stealth liposomal DOX [81–83]. However, the majority of DOX-encapsulated liposomes remain in the clinical trial phase. By modifying the liposome surface, NPs can extend the circulation time of liposomes [84]. Moreover, chemotherapy drugs can be loaded into the nanoliposome capsule, thereby combining chemotherapy with NPs [85]. Liposomal paclitaxel is another example of a liposomal NP that has been used for the clinical treatment of solid tumors [86–88].

NPs can enhance the response to radiation and improve immunotherapeutic efficacy by regulating the tumor microenvironment [89,90] or integrating chemotherapies with other neoadjuvant therapies or adjuvant treatments [91–93]. Some nanomaterials possess unique properties that can be used to attack cancer cells. Photodynamic therapy (PDT) [94,95] and PTT [96,97] are based on nanomaterials with high efficacy during light-to-heat conversions. PDT and PTT can be used as alternative treatments for patients who are unable to undergo surgical resections. The administration of both therapies requires photosensitizers (PSs), which have low efficacies [98,99]. Coupling PSs with targeting molecules or encapsulating them within the cores of nanomaterials can improve the efficiencies of both PDT and PTT. Because of their light-to-heat conversion properties, these NPs can also be utilized to perform photoacoustic imaging (PAI) [100,101]. Combining therapy with imaging is consistent with the theory of developing multifunctional materials. Other novel therapies, such as ion interference and chemo-dynamic treatments, are also NP-dependent.



**Fig. 5.** Structural illustration of <sup>177</sup>Lu-labeled cerasomes encapsulating indocyanine green (ICG@DPDCs-<sup>177</sup>Lu). CSS: cholesteryl succinyl silane. Reproduced from Ref. [67] with permission of American Chemical Society, ©2015.



## 6. Perspective

Tumorigenesis is a complicated process that is associated with multiple changes in molecular biology. To date, functionalized nanomaterials and nanotechnologies, such as microfluidic devices, have achieved great success in improving the efficiency and specificity of ctDNA, tumor-derived DNA, and exosome isolation and detection during the diagnosis of lung cancer. However, single biomarkers may not be adequate for every cancer patient. As an alternative, PM, using a panel of molecular biomarkers identified by genomic and proteomic studies, may be more effective for the early screening of lung cancer. In the future, identifying additional biomarkers and developing a new generation of biosensors, with the help of microfluidics, will facilitate decreases in the cancer mortality rate.

In addition, precision nanomedicine is a promising tool for cancer treatment. Several precision nanomedicine platforms have been developed and used during clinical cancer care. However, concerns regarding the safety of nanomedicines persist. Thus, the comprehensive characterization of nanomedicine products and standards, using both *in vitro* and *in vivo* models, remains necessary to predict the performance of nanomedicines when translated to clinical applications.

## Acknowledgments

This work was supported by the National Program on Key Basic Research Project (2020YFA0211100), National Natural Science Foundation of China (51872205, 51922077, and 81602412), Fundamental Research Funds for the Central Universities, Training Plan of Outstanding Young Medical Talents, Shanghai Municipal Commission of Health and Family Planning (2017YQ050), Scientific Research Project of Shanghai Municipal Commission of Health and Family Planning (2016Y0121), Natural Scientific Foundation of Shanghai (134119b1002), and Outstanding Young Scientific Researcher of Shanghai Pulmonary Hospital.

## Compliance with ethics guidelines

Wei Yin, Feng Pan, Junjie Zhu, Junwu Xu, Diego Gonzalez-Rivas, Meinoshin Okumura, Zhiyong Tang, and Yang Yang declare that they have no conflict of interest or financial conflicts to disclose.

## References

- [1] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019;69(1):7–34.
- [2] Chen W, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, et al. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016;66(2):115–32.
- [3] Sun IC, Ahn CH, Kim K, Emelianov S. Photoacoustic imaging of cancer cells with glycol-chitosan-coated gold nanoparticles as contrast agents. *J Biomed Opt* 2019;24(12):1–5.
- [4] Garcia VB, de Carvalho TG, da Silva Gasparotto LH, da Silva HFO, de Araújo AA, Guerra GCB, et al. Environmentally compatible bioconjugated gold nanoparticles as efficient contrast agents for inflammation-induced cancer imaging. *Nanoscale Res Lett* 2019;14(1):166.
- [5] Tang CK, Vaze A, Shen M, Rusling JF. High-throughput electrochemical microfluidic immunoarray for multiplexed detection of cancer biomarker proteins. *ACS Sens* 2016;1(8):1036–43.
- [6] Munge BS, Stracensky T, Gamez K, DiBiase D, Rusling JF. Multiplex immunosensor arrays for electrochemical detection of cancer biomarker proteins. *Electroanalysis* 2016;28(11):2644–58.
- [7] Cheng Z, Yan X, Sun X, Shen B, Gambhir SS. Tumor molecular imaging with nanoparticles. *Engineering* 2016;2(1):132–40.
- [8] Hood RL, Andriani RT, Ecker TE, Robertson JL, Rylander CG. Characterizing thermal augmentation of convection-enhanced drug delivery with the fiberoptic microneedle device. *Engineering* 2015;1(3):344–50.
- [9] Tang T, Azuma T, Iwahashi T, Takeuchi H, Kobayashi E, Sakuma I. A high-precision US-guided robot-assisted HIFU treatment system for breast cancer. *Engineering* 2018;4(5):702–13.
- [10] Hu G, Guan K, Lu L, Zhang J, Lu N, Guan Y. Engineered functional surfaces by laser microprocessing for biomedical applications. *Engineering* 2018;4(6):822–30.
- [11] He W. Cell therapy: pharmacological intervention enters a third era. *Engineering* 2019;5(1):5–9.
- [12] Tonelli MR, Shirts BH. Knowledge for precision medicine: mechanistic reasoning and methodological pluralism. *JAMA* 2017;318(17):1649–50.
- [13] Badrzadeh F, Rahmati-Yamchi M, Badrzadeh K, Valizadeh A, Zarghami N, Farkhani SM, et al. Drug delivery and nanodetection in lung cancer. *Artif Cells Nanomed Biotechnol* 2016;44(2):618–34.
- [14] Wang Z, Qiao R, Tang N, Lu Z, Wang H, Zhang Z, et al. Active targeting theranostic iron oxide nanoparticles for MRI and magnetic resonance-guided focused ultrasound ablation of lung cancer. *Biomaterials* 2017;127:25–35.
- [15] Ghasemi Y, Peymani P, Afifi S. Quantum dot: magic nanoparticle for imaging, detection and targeting. *Acta Biomed* 2009;80(2):156–65.
- [16] Papagiannaros A, Upponi J, Hartner W, Mongayt D, Levchenko T, Torchilin V. Quantum dot loaded immunomicelles for tumor imaging. *BMC Med Imaging* 2010;10(1):22.
- [17] Jin Y, Jia C, Huang SW, O'Donnell M, Gao X. Multifunctional nanoparticles as coupled contrast agents. *Nat Commun* 2010;1(1):41.
- [18] Xiao L, Tian X, Harihar S, Li Q, Li L, Welch DR, et al. Gd<sub>2</sub>O<sub>3</sub>-doped silica @ Au nanoparticles for *in vitro* imaging cancer biomarkers using surface-enhanced Raman scattering. *Spectrochim Acta A* 2017;181:218–25.
- [19] Neoh KH, Hassan AA, Chen A, Sun Y, Liu P, Xu KF, et al. Rethinking liquid biopsy: microfluidic assays for mobile tumor cells in human body fluids. *Biomaterials* 2018;150:112–24.
- [20] Ooki A, Maleki Z, Tsay JC, Goparaju C, Brait M, Turaga N, et al. A panel of novel detection and prognostic methylated DNA markers in primary non-small cell lung cancer and serum DNA. *Clin Cancer Res* 2017;23(22):7141–52.
- [21] Campbell JD, Alexandrov A, Kim J, Wala J, Berger AH, Pedamallu CS, et al.; Cancer Genome Atlas Research Network. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat Genet* 2016;48(6):607–16.
- [22] Mandel P, Métais P. Les acides nucléiques du plasma sanguin chez l'Homme. *CR Seances Soc Biol Fil* 1948;142(3–4):241–3. French.
- [23] Jahr S, Hentze H, Engelsch S, Hardt D, Fackelmayer FO, Hesch RD, et al. DNA fragments in the blood plasma of cancer patients: quantitations and evidence for their origin from apoptotic and necrotic cells. *Cancer Res* 2001;61(4):1659–65.
- [24] Leon SA, Shapiro B, Sklaroff DM, Yaros MJ. Free DNA in the serum of cancer patients and the effect of therapy. *Cancer Res* 1977;37(3):646–50.
- [25] Chaudhuri AA, Chabon JJ, Lovejoy AF, Newman AM, Stehr H, Azad TD, et al. Early detection of molecular residual disease in localized lung cancer by circulating tumor DNA profiling. *Cancer Discov* 2017;7(12):1394–403.
- [26] Liang Z, Cheng Y, Chen Y, Hu Y, Liu WP, Lu Y, et al. EGFR T790M ctDNA testing platforms and their role as companion diagnostics: correlation with clinical outcomes to EGFR-TKIs. *Cancer Lett* 2017;403:186–94.
- [27] Nguyen AH, Sim SJ. Nanoplasmonic biosensor: detection and amplification of dual bio-signatures of circulating tumor DNA. *Biosens Bioelectron* 2015;67:443–9.
- [28] Dianat S, Bordbar AK, Tangestaninejad S, Yadollahi B, Amiri R, Zarkesh-Esfahani SH, et al. *In vitro* antitumor activity of free and nano-encapsulated Na<sub>5</sub>[PMo<sub>10</sub>V<sub>2</sub>O<sub>40</sub>]·nH<sub>2</sub>O and its binding properties with ctDNA by using combined spectroscopic methods. *J Inorg Biochem* 2015;152:74–81.
- [29] Makarovskiy AN, Ackerley W, Wojcik L, Halpert GK, Stein BS, Carreiro MP, et al. Application of immunomagnetic beads in combination with RT-PCR for the detection of circulating prostate cancer cells. *J Clin Lab Anal* 1997;11(6):346–50.
- [30] Simitzi C, Efstathiopoulos P, Kourgiantaki A, Ranella A, Charalampopoulos I, Fotakis C, et al. Laser fabricated discontinuous anisotropic microconical substrates as a new model scaffold to control the directionality of neuronal network outgrowth. *Biomaterials* 2015;67:115–28.
- [31] Sonnenberg A, Marciniak JY, McCanna J, Krishnan R, Rassenti L, Kipps TJ, et al. Dielectrophoretic isolation and detection of cfc-DNA nanoparticle biomarkers and virus from blood. *Electrophoresis* 2013;34(7):1076–84.
- [32] Jeon SH, Hong WY, Lee ES, Cho Y. High-purity isolation and recovery of circulating tumor cells using conducting polymer-deposited microfluidic device. *Theranostics* 2014;4(11):1123–32.
- [33] Jeon SH, Lee HJ, Bae K, Yoon KA, Lee ES, Cho Y. Efficient capture and isolation of tumor-related circulating cell-free DNA from cancer patients using electroactive conducting polymer nanowire platforms. *Theranostics* 2016;6(6):828–36.
- [34] Bernabé R, Hickson N, Wallace A, Blackhall FH. What do we need to make circulating tumour DNA (ctDNA) a routine diagnostic test in lung cancer? *Eur J Cancer* 2017;81:66–73.
- [35] Xu T, Kang X, You X, Dai L, Tian D, Yan W, et al. Cross-platform comparison of four leading technologies for detecting EGFR mutations in circulating tumor DNA from non-small cell lung carcinoma patient plasma. *Theranostics* 2017;7(6):1437–46.
- [36] Zhao L, Tang C, Xu L, Zhang Z, Li X, Hu H, et al. Enhanced and differential capture of circulating tumor cells from lung cancer patients by microfluidic assays using aptamer cocktail. *Small* 2016;12(8):1072–81.
- [37] Zhou B, Nie J, Yang W, Huang C, Huang Y, Zhao H. Effect of hydrothorax EGFR gene mutation and EGFR-TKI targeted therapy on advanced non-small cell lung cancer patients. *Oncol Lett* 2016;11(2):1413–7.

- [38] Roscilli G, De Vitis C, Ferrara FF, Noto A, Cherubini E, Ricci A, et al. Human lung adenocarcinoma cell cultures derived from malignant pleural effusions as model system to predict patients chemosensitivity. *J Transl Med* 2016;14(1):61.
- [39] Vallée A, Marcq M, Bizieux A, Kouri CE, Lacroix H, Bennouna J, et al. Plasma is a better source of tumor-derived circulating cell-free DNA than serum for the detection of *EGFR* alterations in lung tumor patients. *Lung Cancer* 2013;82(2):373–4.
- [40] Latifi Z, Fattahi A, Ranjbaran A, Nejabati HR, Imakawa K. Potential roles of metalloproteinases of endometrium-derived exosomes in embryo-maternal crosstalk during implantation. *J Cell Physiol* 2018;233(6):4530–45.
- [41] Yang MQ, Du Q, Varley PR, Goswami J, Liang Z, Wang R, et al. Interferon regulatory factor 1 priming of tumour-derived exosomes enhances the antitumour immune response. *Br J Cancer* 2018;118(1):62–71.
- [42] Sansone P, Savini C, Kurelac I, Chang Q, Amato LB, Strillacci A, et al. Packaging and transfer of mitochondrial DNA via exosomes regulate escape from dormancy in hormonal therapy-resistant breast cancer. *Proc Natl Acad Sci USA* 2017;114(43):E9066–75. Corrected in: *Proc Natl Acad Sci USA* 2017;114(47):E10255.
- [43] Son KJ, Rahimian A, Shin DS, Siltanen C, Patel T, Revzin A. Microfluidic compartments with sensing microbeads for dynamic monitoring of cytokine and exosome release from single cells. *Analyst* 2016;141(2):679–88.
- [44] He M, Crow J, Roth M, Zeng Y, Godwin AK. Integrated immunoisolation and protein analysis of circulating exosomes using microfluidic technology. *Lab Chip* 2014;14(19):3773–80.
- [45] Mansur N, Raziul Hasan M, Kim YT, Iqbal SM. Functionalization of nanotextured substrates for enhanced identification of metastatic breast cancer cells. *Nanotechnology* 2017;28(38):385101.
- [46] Wan Y, Cheng G, Liu X, Hao SJ, Nisic M, Zhu CD, et al. Rapid magnetic isolation of extracellular vesicles via lipid-based nanoprobe. *Nat Biomed Eng* 2017;1(4):0058.
- [47] Zhang P, He M, Zeng Y. Ultrasensitive microfluidic analysis of circulating exosomes using a nanostructured graphene oxide/polydopamine coating. *Lab Chip* 2016;16(16):3033–42.
- [48] In GK, Nieva J. Emerging chemotherapy agents in lung cancer: nanoparticle therapeutics for non-small cell lung cancer. *Transl Cancer Res* 2015;4(4):340–55.
- [49] Nazir S, Hussain T, Ayub A, Rashid U, MacRobert AJ. Nanomaterials in combating cancer: therapeutic applications and developments. *Nanomed-Nanotechnol* 2014;10(1):19–34.
- [50] Wicki A, Witzigmann D, Balasubramanian V, Huwyler J. Nanomedicine in cancer therapy: challenges, opportunities, and clinical applications. *J Control Release* 2015;200:138–57.
- [51] Kumar D, Mutreja I, Chitcholtan K, Sykes P. Cytotoxicity and cellular uptake of different sized gold nanoparticles in ovarian cancer cells. *Nanotechnology* 2017;28(47):475101.
- [52] Chen YH, Tsai CY, Huang PY, Chang MY, Cheng PC, Chou CH, et al. Methotrexate conjugated to gold nanoparticles inhibits tumor growth in a syngeneic lung tumor model. *Mol Pharmacol* 2007;4(5):713–22.
- [53] Yokoyama T, Tam J, Kuroda S, Scott AW, Aaron J, Larson T, et al. EGFR-targeted hybrid plasmonic magnetic nanoparticles synergistically induce autophagy and apoptosis in non-small cell lung cancer cells. *PLoS ONE* 2011;6(11):e25507.
- [54] Kao HW, Lin YY, Chen CC, Chi KH, Tien DC, Hsia CC, et al. Evaluation of EGFR-targeted radioimmuno-gold-nanoparticles as a theranostic agent in a tumor animal model. *Bioorg Med Chem Lett* 2013;23(11):3180–5.
- [55] Bernardes VHF, Qu Y, Du Z, Beaton J, Vargas MD, Farrell NP. Interaction of the HIV NCP7 protein with platinum(II) and gold(III) complexes containing tridentate ligands. *Inorg Chem* 2016;55(21):11396–407.
- [56] Li Q, Wang Q, Yang X, Wang K, Zhang H, Nie W. High sensitivity surface plasmon resonance biosensor for detection of microRNA and small molecule based on graphene oxide-gold nanoparticles composites. *Talanta* 2017;174:521–6.
- [57] Knežević NŽ, Durand JO. Targeted treatment of cancer with nanotherapeutics based on mesoporous silica nanoparticles. *ChemPlusChem* 2015;80(1):26–36.
- [58] Bardhan M, Majumdar A, Jana S, Ghosh T, Pal U, Swarnakar S, et al. Mesoporous silica for drug delivery: interactions with model fluorescent lipid vesicles and live cells. *J Photochem Photobiol B* 2018;178:19–26.
- [59] Zhang Z, Wang L, Wang J, Jiang X, Li X, Hu Z, et al. Mesoporous silica-coated gold nanorods as a light-mediated multifunctional theranostic platform for cancer treatment. *Adv Mater* 2012;24(11):1418–23.
- [60] Liu J, Luo Z, Zhang J, Luo T, Zhou J, Zhao X, et al. Hollow mesoporous silica nanoparticles facilitated drug delivery via cascade pH stimuli in tumor microenvironment for tumor therapy. *Biomaterials* 2016;83:51–65.
- [61] Liu D, Kramer SA, Huxford-Phillips RC, Wang S, Della Rocca J, Lin W. Coercing bisphosphonates to kill cancer cells with nanoscale coordination polymers. *Chem Commun* 2012;48(21):2668–70.
- [62] Liu D, Poon C, Lu K, He C, Lin W. Self-assembled nanoscale coordination polymers with trigger release properties for effective anticancer therapy. *Nat Commun* 2014;5(1):4182.
- [63] Au KM, Satterlee A, Min Y, Tian X, Kim YS, Caster JM, et al. Folate-targeted pH-responsive calcium zoledronate nanoscale metal-organic frameworks: turning a bone antiresorptive agent into an anticancer therapeutic. *Biomaterials* 2016;82:178–93.
- [64] Muthu MS, Mei L, Feng SS. Nanotheranostics: advanced nanomedicine for the integration of diagnosis and therapy. *Nanomedicine* 2014;9(9):1277–80.
- [65] Mura S, Couvreur P. Nanotheranostics for personalized medicine. *Adv Drug Deliv Rev* 2012;64(13):1394–416.
- [66] Cheng FY, Su CH, Wu PC, Yeh CS. Multifunctional polymeric nanoparticles for combined chemotherapeutic and near-infrared photothermal cancer therapy *in vitro* and *in vivo*. *Chem Commun* 2010;46(18):3167–9.
- [67] Jing L, Shi J, Fan D, Li Y, Liu R, Dai Z, et al. <sup>177</sup>Lu-labeled cerasomes encapsulating indocyanine green for cancer theranostics. *ACS Appl Mater Interfaces* 2015;7(39):22095–105.
- [68] Zhang L, Li J, Liu K. Recent advances in gadolinium-based MRI metal responsive agent. *Sci China Technol Sci* 2018;61(9):1329–33.
- [69] Anselmo AC, Mitragotri S. Nanoparticles in the clinic. *Bioeng Transl Med* 2016;1(1):10–29.
- [70] Knights OB, McLaughlan JR. Gold nanorods for light-based lung cancer theranostics. *Int J Mol Sci* 2018;19(11):3318.
- [71] He Y, Du Z, Ma S, Liu Y, Li D, Huang H, et al. Effects of green-synthesized silver nanoparticles on lung cancer cells *in vitro* and grown as xenograft tumors *in vivo*. *Int J Nanomed* 2016;11:1879–87.
- [72] Svehkarev D, Mohs AM. Organic fluorescent dye-based nanomaterials: advances in the rational design for imaging and sensing applications. *Curr Med Chem* 2019;26(21):4042–64.
- [73] Le Duc G, Roux S, Paruta-Tuarez A, Dufort S, Brauer E, Marais A, et al. Advantages of gadolinium based ultrasmall nanoparticles vs molecular gadolinium chelates for radiotherapy guided by MRI for glioma treatment. *Cancer Nanotechnol* 2014;5(1):4.
- [74] Ivashchenko O, Peplińska B, Gapiński J, Flak D, Jarek M, Załęski K, et al. Silver and ultrasmall iron oxides nanoparticles in hydrocolloids: effect of magnetic field and temperature on self-organization. *Sci Rep* 2018;8(1):4041.
- [75] Nguyen H, Tinet E, Chauveau T, Geinguenaud F, Lalatonne Y, Michel A, et al. Bimodal fucoidan-coated zinc oxide/iron oxide-based nanoparticles for the imaging of atherothrombosis. *Molecules* 2019;24(5):962.
- [76] Gonzalez-Rivas D, Yang Y, Ng C. Advances in unipolar video-assisted thoracoscopic surgery: pushing the envelope. *Thorac Surg Clin* 2016;26(2):187–201.
- [77] Ban I, Markuš S, Gyergyek S, Drofenik M, Korenák J, Helix-Nielsen C, et al. Synthesis of poly-sodium-acrylate (PSA)-coated magnetic nanoparticles for use in forward osmosis draw solutions. *Nanomaterials* 2019;9(9):1238.
- [78] Edelman R, Assaraf YG, Slavkin A, Dolev T, Shahar T, Livney YD. Developing body-components-based theranostic nanoparticles for targeting ovarian cancer. *Pharmaceutics* 2019;11(5):216.
- [79] Ahlawat J, Henriquez G, Narayan M. Enhancing the delivery of chemotherapeutics: role of biodegradable polymeric nanoparticles. *Molecules* 2018;23(9):2157.
- [80] Muhamad N, Plengsuriyakarn T, Na-Bangchang K. Application of active targeting nanoparticle delivery system for chemotherapeutic drugs and traditional/herbal medicines in cancer therapy: a systematic review. *Int J Nanomed* 2018;13:3921–35.
- [81] Zylberberg C, Gaskill K, Pasley S, Matosevic S. Engineering liposomal nanoparticles for targeted gene therapy. *Gene Ther* 2017;24(8):441–52.
- [82] Mayer LD, Tardi P, Louie AC. CPX-351: a nanoscale liposomal co-formulation of daunorubicin and cytarabine with unique biodistribution and tumor cell uptake properties. *Int J Nanomed* 2019;14:3819–30.
- [83] Caracciolo G, Palchetti S, Digiacomo L, Chiozzi RZ, Capriotti AL, Amenitsch H, et al. Human biomolecular corona of liposomal doxorubicin: the overlooked factor in anticancer drug delivery. *ACS Appl Mater Interfaces* 2018;10(27):22951–62.
- [84] Fathi S, Oyeler AK. Liposomal drug delivery systems for targeted cancer therapy: is active targeting the best choice? *Future Med Chem* 2016;8(17):2091–112.
- [85] Hwang H, Jeong HS, Oh PS, Kim M, Lee TK, Kwon J, et al. PEGylated nanoliposomes encapsulating angiogenic peptides improve perfusion defects: radionuclide imaging-based study. *Nucl Med Biol* 2016;43(9):552–8.
- [86] Wagner U, Marth C, Largillier R, Kaern J, Brown C, Heywood M, et al. Final overall survival results of phase III GCIG CALYPSO trial of pegylated liposomal doxorubicin and carboplatin vs paclitaxel and carboplatin in platinum-sensitive ovarian cancer patients. *Br J Cancer* 2012;107(4):588–91.
- [87] Nehate C, Jain S, Saneja A, Khare V, Alam N, Dubey RD, et al. Paclitaxel formulations: challenges and novel delivery options. *Curr Drug Deliv* 2014;11(6):666–86.
- [88] Pujade-Lauraine E, Hilpert F, Weber B, Reuss A, Poveda A, Kristensen G, et al. Bevacizumab combined with chemotherapy for platinum-resistant recurrent ovarian cancer: the AURELIA open-label randomized phase III trial. *J Clin Oncol* 2014;32(13):1302–8.
- [89] Rajendrakumar SK, Uthaman S, Cho CS, Park IK. Nanoparticle-based phototriggered cancer immunotherapy and its domino effect in the tumor microenvironment. *Biomacromolecules* 2018;19(6):1869–87.
- [90] Dougan M, Dougan SK. Targeting immunotherapy to the tumor microenvironment. *J Cell Biochem* 2017;118(10):3049–54.
- [91] Riley RS, June CH, Langer R, Mitchell MJ. Delivery technologies for cancer immunotherapy. *Nat Rev Drug Discov* 2019;18(3):175–96.
- [92] Cisterna BA, Kamaly N, Choi WI, Tavakkoli A, Farokhzad OC, Vilos C. Targeted nanoparticles for colorectal cancer. *Nanomedicine* 2016;11(18):2443–56.
- [93] Geiger BC, Wang S, Padera RF, Grodzinsky AJ, Hammond PT. Cartilage-penetrating nanocarriers improve delivery and efficacy of growth factor treatment of osteoarthritis. *Sci Transl Med* 2018;10(469):eaat8800.



- [94] Battogtokh G, Ko YT. Mitochondrial-targeted photosensitizer-loaded folate-albumin nanoparticle for photodynamic therapy of cancer. *Nanomedicine* 2017;13(2):733–43.
- [95] Sivasubramanian M, Chuang YC, Lo LW. Evolution of nanoparticle-mediated photodynamic therapy: from superficial to deep-seated cancers. *Molecules* 2019;24(3):520.
- [96] Ghalandari B, Asadollahi K, Shakerizadeh A, Komeili A, Riazi G, Kamrava SK, et al. Microtubule network as a potential candidate for targeting by gold nanoparticle-assisted photothermal therapy. *J Photochem Photobiol B* 2019;192:131–40.
- [97] Sweeney EE, Cano-Mejia J, Fernandes R. Photothermal therapy generates a thermal window of immunogenic cell death in neuroblastoma. *Small* 2018;14(20):1800678.
- [98] Daraee H, Eatemadi A, Abbasi E, Fekri Aval S, Kouhi M, Akbarzadeh A. Application of gold nanoparticles in biomedical and drug delivery. *Artif Cells Nanomed Biotechnol* 2016;44(1):410–22.
- [99] Kruger CA, Abrahamse H. Utilisation of targeted nanoparticle photosensitiser drug delivery systems for the enhancement of photodynamic therapy. *Molecules* 2018;23(10):2628.
- [100] Kim T, Zhang Q, Li J, Zhang L, Jokerst JV. A gold/silver hybrid nanoparticle for treatment and photoacoustic imaging of bacterial infection. *ACS Nano* 2018;12(6):5615–25.
- [101] Donnelly EM, Kubelick KP, Duman DS, Emelianov SY. Photoacoustic image-guided delivery of plasmonic-nanoparticle-labeled mesenchymal stem cells to the spinal cord. *Nano Lett* 2018;18(10):6625–32.