

Segmentectomy or lobectomy for early stage lung cancer: a meta-analysis

Feichao Bao, Peng Ye, Yunhai Yang, Luming Wang, Chong Zhang, Xiayi Lv and Jian Hu*

Department of Thoracic Surgery, First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, China

* Corresponding author. Department of Thoracic Surgery, First Hospital, College of Medicine, Zhejiang University, No. 79, Qingchun Road, Hangzhou, Zhejiang 310003, China. Tel: +86-571-87236770; fax: +86-571-87236770; e-mail: hujian_med@163.com (J. Hu).

Received 15 August 2013; received in revised form 11 October 2013; accepted 21 October 2013

Summary

Early stage lung cancer is routinely treated by lobectomy whenever clinically feasible, whereas the role of segmentectomy is controversial. The purpose of this study was to investigate the benefits of segmentectomy vs lobectomy for early stage lung cancer through a meta-analysis of published data. Eligible studies were identified from MEDLINE through February 2013. The manual selection of relevant studies was based on the summary analysis. We used published hazard ratios (HRs) if available or estimates from the published survival data. Lobectomy was chosen as the reference in all HR calculations. We compared the effect of segmentectomy and lobectomy for Stage I, Stage IA, Stage IA with tumours larger than 2 cm but smaller than 3 cm in size and Stage IA with tumours of 2 cm or smaller in 22 observational studies. The HRs of overall and cancer-specific survival indicated significant benefits of lobectomy for Stage I, Stage IA and Stage IA with tumours larger than 2 cm but smaller than 3 cm at 1.20 (95% confidence interval [CI] 1.04–1.38; $P = 0.011$), 1.24 (95% CI 1.08–1.42; $P = 0.002$) and 1.41 (95% CI 1.14–1.71; $P = 0.001$), respectively. For tumours 2 cm or smaller, segmentectomy provided an effect equivalent to that of lobectomy (HR 1.05; 95% CI 0.89–1.24; $P = 0.550$). No significant publication bias was detected in any part of the analysis. These findings should be interpreted in the context of the inherent limitations of meta-analyses of retrospective studies, including the heterogeneity of patient characteristics.

Keywords: Lung cancer • Segmentectomy • Lobectomy • Survival • Meta-analysis

INTRODUCTION

Lung cancer is the leading cause of cancer-related mortality worldwide, with ~1.4 million deaths yearly [1]. Lobectomy with lymph node dissection is the standard surgical treatment for Stage I non-small-cell lung cancer (NSCLC), whereas segmentectomy remains controversial because it lacks supporting data. In 1995, a prospective, multi-institutional, randomized trial reported by the Lung Cancer Study Group (LCSG) compared limited pulmonary resection (wedge resection and segmentectomy) with lobectomy for T1N0M0 NSCLC patients. This study found that limited resection was associated with inferior overall survival (OS) and a higher regional recurrence rate compared with lobectomy [2]. Other studies have also confirmed the results of the LCSG [3–5]. Since then, pulmonary segmentectomy has been applied only to those who are unable to tolerate lobectomy because of limited cardio-pulmonary function. However, the unfavourable results for limited resection in the LCSG study may have been related to 40 of the 122 patients who received limited resection undergoing wedge resection, which could have resulted in recurrence and poor outcomes compared with segmental resection [4]. Pulmonary wedge resection, a non-anatomical resection in which the hilar lymph nodes cannot be dissected efficiently, is associated with poor rates of overall and cancer-specific survival (CSS) when compared with segmentectomy, which allows the dissection of the hilar

lymph nodes and provides precise staging data that facilitate post-operative treatment [4]. Although lung cancers are increasingly diagnosed at early stages through the widespread use of computed tomography [6], it is unclear whether these patients with very early stage lung cancer should undergo lobectomy, a more aggressive procedure than segmentectomy. Several studies have demonstrated that limited resection, and especially segmentectomy, for Stage I NSCLC may have an effect equivalent to that of lobectomy [5, 7–12]. Additionally, segmentectomy can improve postoperative quality of life by preserving the pulmonary function significantly better than lobectomy [13].

Based on these considerations, we conducted a meta-analysis of published studies to quantitatively review survival data for the segmentectomy of Stage I lung cancer compared with those for standard lobectomy.

MATERIALS AND METHODS

Eligibility criteria

This meta-analysis was limited to studies comparing survival data of pulmonary segmentectomy with those of pulmonary lobectomy. The following eligibility criteria were established before collecting articles. (i) Operative approaches could include

thoracotomy, video-assisted thoracoscopic surgery or robotic surgery. (ii) Survival rates for a specific time interval after operation were stated in the article. (iii) Study subjects were limited to clinical Stage I patients. (iv) Median follow-up time was to exceed 1 year. (v) Articles were published or accepted in English in the periodical medical literature from January 1990 to March 2013. (vi) OS or CSS was assessed in detail. (vii) Twenty or more patients were included. (viii) When multiple articles by the same author or the study group analysed the same series of patients, the single most informative article was chosen for the meta-analysis. The exclusion criteria included studies published in letters, reviews and editorials, and papers not published in English.

Collection of published studies

The primary sources of the reviewed studies were MEDLINE abstracts published up to February 2013. The search included the following terms: ('lung neoplasms/surgery' [MESH terms]) and ('segmentectomy' or 'segmental resection') and 'lobectomy' and 'survival'. The computer search was supplemented with manual searches of the reference lists of all retrieved review articles, primary studies and abstracts from meetings. The manual selection of relevant studies was based on the summary analysis. Overlapping or unrelated articles were excluded, and items from manually searched bibliographies were added.

Statistical analysis

OS and CSS were estimated as the hazard ratio (HR) for each study. When possible, the HR and associated variance were

obtained directly from each publication or from individual patient data. HRs not reported were calculated by the Parmar and Tierney methods [11, 12]. Because HRs for OS/CSS could not be obtained directly from the data presented in some studies, the Kaplan–Meier survival curves of these studies were analysed to extract the data and to calculate the HR and standard of error for OS/CSS.

HRs and 95% confidence intervals (CIs) were obtained for each study, and the pooled estimate was calculated from an inverse variance-weighted average of the individual studies [13]. χ^2 tests were used to assess statistical heterogeneity. All statistical tests were two-sided, and the significance level was set at 5%. Publication bias was examined using the approach of Egger *et al.* [14], who suggested the presence of significant publication bias when the *P*-value was <0.1. Kaplan–Meier curves were read by Engauge Digitizer version 4.1 (free software downloaded from <http://sourceforge.net/projects/digitizer/files/Engauge%20Digitizer/digitizer-4.1/>). Data combining and tests of heterogeneity were performed with Stata version 12.0 (StatCorp, College Station, TX, USA). DerSimonian–Laird random-effects analysis [15] was used to combine the HR and 95% CI values to estimate the total effect.

RESULTS

Selected articles and description of the studies

The meta-analysis included 22 observational studies, the characteristics of which are summarized in Table 1. The search on segmentectomy vs lobectomy for lung cancer is illustrated in the flow diagram in Fig. 1. No randomized controlled trials were found. Most of the studies included were based on retrospective data or prospective, non-randomized, controlled studies except for that

Table 1: Characteristics of the included studies

Authors	Publication year	Study design	Stage	No. of seg.	No. of lob.	Survival difference	Reason for segmentectomy
Warren and Faber [3]	1994	RS	I	66	103	LB	Poor cardiopulmonary function and smaller lesions
Okada <i>et al.</i> [16]	2001	PS	IA ≤ 2 cm	68	104	NS	Intentional resection for small lesions of ≤ 2 cm
Campione <i>et al.</i> [17]	2004	RS	IA	21	100	NS	Poor cardiopulmonary function
Keenan <i>et al.</i> [8]	2004	RS	I	54	147	NS	Poor pulmonary function
Martin-Ucar <i>et al.</i> [10]	2005	MPS	I	17	17	NS	Poor pulmonary function
Watanabe <i>et al.</i> [18]	2005	PS	IA ≤ 2 cm	20	57	NS	Intentional resection for small lesions of ≤ 2 cm
Okada <i>et al.</i> [19]	2005	RS	I	211	496	NS	ND
Okumura <i>et al.</i> [20]	2007	RS	IA	84	418	NS	ND
Schuchert <i>et al.</i> [21]	2007	PS	I	182	246	NS	ND
Iwasaki <i>et al.</i> [22]	2007	RS	IA ≤ 2 cm	31	55	NS	Intentional resection for small lesions of ≤ 2 cm
Shapiro <i>et al.</i> [7]	2009	RS	I	31	113	NS	Small lesions of ≤ 2 cm, poor pulmonary function
Kilic <i>et al.</i> [5]	2009	PS	I	78	106	NS	Age >75 years
Sugi <i>et al.</i> [23]	2010	PS	IA	43	95	NS	Intentional resection for small lesions of ≤ 2 cm
Whitson <i>et al.</i> [24]	2011	RS	I	581	13 892	LB	ND
Nakamura <i>et al.</i> [25]	2011	RS	I	38	289	NS	ND
Zhong <i>et al.</i> [26]	2012	PS	IA ≤ 2 cm	39	81	NS	Intentional resection for small lesions of ≤ 2 cm
Yamashita <i>et al.</i> [27]	2012	PS	IA	90	124	NS	ND
Cheng <i>et al.</i> [9]	2012	PS	I	32	32	NS	Age ≥ 70 years, poor pulmonary function
Soukiasian <i>et al.</i> [28]	2012	RS	IA	56	178	NS	ND
Yendamuri <i>et al.</i> [29]	2012	RS	IA ≤ 2 cm	311	5142	NS	Intentional resection for small lesions of ≤ 2 cm
Hamatake <i>et al.</i> [30]	2012	RS	IA ≤ 1 cm	32	77	NS	Intentional resection for small lesions of ≤ 1 cm
Carr <i>et al.</i> [31]	2012	RS	IA	178	251	NS	ND

Seg: segmentectomy; Lob: lobectomy; RS: retrospective study; PS: prospective non-randomized study; NS: not significant; LB: lobectomy better; ND: not described; MPS: matched-pair study.




Figure 1: Flow diagram of the search results for segmentectomy vs lobectomy.

by Martin-Ucar *et al.* [10], which was a case-matched study. Cohort sizes ranged from 34 to 14 473 patients. Lobectomy was chosen as the reference in all HR calculations.

Effect of operative approach on the overall survival/cancer-specific survival of Stage I NSCLC patients

Sixteen studies analysed OS and three analysed CSS. Studies from three institutions used stratified analysis and did not provide the results of the whole cohort, and these data were analysed separately [19, 20, 28]. The combined HR of OS was 1.07 (95% CI 0.90–1.28; $P = 0.441$), that of CSS was 1.38 (95% CI 1.20–1.59; $P < 0.001$) and that of OS/CSS was 1.28. In combining the data from univariate analysis, we found that the OS/CSS of patients treated with segmentectomy was inferior to that of patients treated with lobectomy (HR 1.28; 95% CI 1.15–1.42; $P < 0.001$; Fig. 2 and Table 2). The tests for heterogeneity in the subanalysis of OS and CSS showed results of $\chi^2 = 17.25$ ($P = 0.438$, $df = 17$) and $\chi^2 = 2.34$ ($P = 0.673$, $df = 4$), respectively, which suggested no significant inconsistency. A P -value of 0.770 suggested no evidence of publication bias (Table 2).

Effect of operative approach on the overall survival/cancer-specific survival of Stage IA NSCLC patients

Twelve studies focused on the effect of segmentectomy or lobectomy on the OS of Stage IA patients by univariate analysis, and

two focused on the CSS of Stage IA patients. Five studies used stratified analysis and did not provide the results of the whole cohort [3, 20, 24, 31, 32], and these data were analysed separately. The combined HR value in the subanalysis of OS was 1.23 (95% CI 1.07–1.41; $P = 0.003$), that of CSS was 1.35 (95% CI 0.74–2.47; $P = 0.324$) and that of OS/CSS was 1.24. Segmentectomy resulted in inferior OS/CSS compared with that for lobectomy (HR 1.24; 95% CI 1.09–1.41; $P = 0.002$; Fig. 3 and Table 2). The tests for heterogeneity in the subanalysis of OS and CSS gave results of $\chi^2 = 10.25$ ($P = 0.744$, $df = 14$) and $\chi^2 = 1.45$ ($P = 0.693$, $df = 3$), respectively, which suggested no inconsistency among the results of the included studies (Fig. 3 and Table 2). A P -value of 0.265 suggested a lack of publication bias (Table 2).

Effect of operative approach on the overall survival/cancer-specific survival of Stage IA NSCLC patients with tumours larger than 2 cm but smaller than 3 cm

Six studies focused on the effect of segmentectomy or lobectomy on the OS/CSS of Stage IA NSCLC patients with tumours larger than 2 cm but smaller than 3 cm. These data were analysed separately. The combined HR value in the subanalysis of OS was 1.81 (95% CI 1.21–2.71; $P = 0.004$), that of CSS was 1.27 (95% CI 1.00–1.64; $P = 0.500$) and that of OS/CSS was 1.39. Segmentectomy resulted in inferior OS/CSS compared with that for lobectomy (HR 1.41; 95% CI 1.14–1.71; $P = 0.001$; Fig. 4 and Table 2). The tests for heterogeneity in the subanalysis of OS and CSS showed the results of $\chi^2 = 1.10$ ($P = 0.578$, $df = 2$) and $\chi^2 = 1.39$ ($P = 0.497$, $df = 2$), respectively, suggesting no inconsistency among the results of the included studies (Fig. 4 and Table 2). On the basis of a P -value of 0.678, no publication bias existed (Table 2).

Effect of operative approach on the overall survival/cancer-specific survival of Stage IA NSCLC patients with tumours of 2 cm or smaller

Ten studies focused on the effect of segmentectomy or lobectomy on the OS/CSS of Stage IA NSCLC patients with tumours of 2 cm or smaller. One study used stratified analysis and did not provide the results of the whole cohort [29], and these data were also analysed separately. The combined HR value in the subanalysis of OS was 1.05 (95% CI 0.89–1.24; $P = 0.599$), that of CSS was 1.14 (95% CI 0.56–2.32; $P = 0.709$) and that of OS/CSS was 1.05. The rates of OS/CSS with segmentectomy were equivalent to those with lobectomy (HR 1.05; 95% CI 0.89–1.24; $P = 0.550$; Fig. 5 and Table 2). The tests for heterogeneity in the subanalysis of OS and CSS showed the results of $\chi^2 = 4.15$ ($P = 0.843$, $df = 8$) and $\chi^2 = 0.07$ ($P = 0.876$, $df = 1$), respectively, which suggested no inconsistency among the results of the included studies (Fig. 5 and Table 2). No publication bias was detected, as indicated by a P -value of 0.802 (Table 2).

DISCUSSION

In 1995, a prospective, multi-institutional, randomized trial compared sublobar resection with lobectomy for T1N0M0 NSCLC patients and showed a discouraging outcome for sublobar




Figure 2: Overall survival/cancer-specific survival estimates for segmentectomy compared with lobectomy for Stage I NSCLC. Seg: segmentectomy; Lob: lobectomy.

resection due to an inferior OS rate and a higher recurrence rate than for lobectomy [2]. Since then, studies comparing the effect of sublobar resection and lobectomy have been limited to retrospective reviews with a small sample size [7, 19, 20, 22]. Advances in minimally invasive surgical techniques, critical care and imaging have caused the issue of sublobar resection, especially segmentectomy, to remerge [24]. The poor prognosis associated with sublobar resection may be due to the use of non-anatomical wedge resection, which does not allow efficient dissection of the pulmonary hilar lymph nodes, whereas anatomical segmentectomy does allow the dissection of these nodes and may be suitable for candidates with small-sized NSCLC. Furthermore, the accurate lymph node staging data available with segmentectomy but not with wedge resection are important for postoperative treatment.

Recently, thoracic surgeons have focused on the evaluation of segmentectomy and lobectomy for patients with small-sized NSCLC, with both encouraging and discouraging outcomes reported (Table 1). Meta-analyses reported by Nakamura *et al.* [33] and Fan *et al.* [34] compared sublobar resection and lobectomy. Nakamura combined differences in survival (survival rate with lobectomy minus that with limited resection) at 1, 3 and 5 years after resection and found no significant differences in survival, suggesting that survival after sublobar resection for Stage I lung cancer is comparable with that after lobectomy [33]. However, their report was based on data published about 10 years ago and much additional data have been published in

Table 2: Summary of comparison results

Series	Survival	Comparison of survival			Egger's test	
		HR	95% CI	P-value	Bias	P-value
I	OS/CSS	1.20	1.04–1.38	0.011	0.12	0.770
IA	OS/CSS	1.24	1.08–1.42	0.002	–0.41	0.265
IA (2–3 cm)	OS/CSS	1.41	1.14–1.71	0.001	0.35	0.678
IA (≤ 2 cm)	OS/CSS	1.05	0.89–1.24	0.550	–0.15	0.802

recent years. Fan *et al.* conducted a meta-analysis to compare the outcomes of segmentectomy with lobectomy for stage I NSCLC patients based on eight published studies, the results showed segmentectomy produced an outcome similar to that with lobectomy [34]; nonetheless, their findings are limited, with many more data reported in recent years. Moreover, these two meta-analyses do not compare segmentectomy with lobectomy using stratified data based on different tumour sizes [33, 34], and it is necessary to conduct a more stratified meta-analysis comparing segmentectomy and lobectomy that is based on updated data.

Meta-analysis is often used in randomized clinical trials (RCTs), but the application of meta-analysis to observational studies is controversial [35]. However, when no RCTs are available, as is the




Figure 3: Overall survival/cancer-specific survival estimates for segmentectomy compared with lobectomy for Stage IA NSCLC. Seg: segmentectomy; Lob: lobectomy.




Figure 4: Overall survival/cancer-specific survival estimates for segmentectomy compared with lobectomy for Stage IA NSCLC patients with tumours larger than 2 cm but smaller than 3 cm. Seg: segmentectomy; Lob: lobectomy.




Figure 5: Overall survival/cancer-specific survival estimates for segmentectomy compared with lobectomy for Stage IA NSCLC patients with tumour 2 cm or smaller. Seg: segmentectomy; Lob: lobectomy.

case for segmentectomy in Stage I NSCLC, a meta-analysis of observational studies may be an available method for evaluating efficacy and effectiveness [36]. Performance of this type of meta-analysis may be challenging because of the differences in the baseline characteristics of each study [37]. We addressed this issue by evaluating the statistical heterogeneity of the component studies and by performing subgroup analyses. Admittedly, our meta-analysis covers a mix of small retrospective studies with various indications for segmentectomy, including poor cardiopulmonary function, elderly patients and small lesions, and thus further comparison is needed to clarify which indication provides the largest benefit.

We combined data from 22 studies published from 1990 to 2012 and performed a meta-analysis by combining HRs. Our results indicate that compared with lobectomy, segmentectomy is statistically significantly associated with OS/CSS, with an increased HR for Stage I NSCLC patients. These results are inconsistent with those of Fan *et al.*, who found an equivalent effect of segmentectomy and lobectomy on Stage I NSCLC patients in eight studies from 1990 to 2010 [34]. The different outcomes in these two meta-analyses are probably due to the more recent articles included in the present analysis.

To further explore NSCLC candidates who can benefit from segmentectomy, we analysed the effect of segmentectomy or lobectomy on the OS/CSS of patients in three subgroups: Stage IA, Stage IA with tumours larger than 2 cm but smaller than 3 cm and Stage IA with tumours 2 cm or smaller. We found that segmentectomy resulted in inferior survival compared with lobectomy in the subgroups Stage IA and Stage IA with tumours larger than 2 cm but smaller than 3 cm. However, segmentectomy did

provide survival similar to that of lobectomy for subgroup Stage IA with tumours 2 cm or smaller. In general, if there are no anatomical contraindications to segmentectomy, a lesion of 2 cm or smaller should be amenable to anatomical segmentectomy. Examination of this issue is currently being undertaken by the Cancer and Leukaemia Group B (CALGB) in a large, randomized, controlled trial (CALGB 140503), which is also using the 2-cm-or-smaller criterion. The CALGB 140503 clinical trial will provide results in several years.

In conclusion, our meta-analysis suggests that segmentectomy provides survival inferior to that following lobectomy for Stage I NSCLC patients, Stage IA patients and Stage IA patients with tumours larger than 2 cm but smaller than 3 cm. However, for Stage IA patients with tumours 2 cm or smaller, segmentectomy had an effect equivalent to that of lobectomy. Based on these results, we recommend segmentectomy for lung cancer patients with tumours 2 cm or smaller but not with tumours larger than 2 cm. However, because of possible bias in the primary studies, heterogeneity between the studies and inherent limitations of the meta-analysis of observational studies, this conclusion should be interpreted cautiously. More evidence is needed to establish the role of segmentectomy in early NSCLC. We expect to explore the answer in future RCTs such as that of CALGB 140503.

Funding

The study was supported by Health Bureau of Zhejiang Province, with the identification number 2008A067. The funders had no

role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Conflict of interest: none declared.

REFERENCES

- [1] Jemal A, Center MM, DeSantis C, Ward EM. Global patterns of cancer incidence and mortality rates and trends. *Cancer Epidemiol Biomarkers Prev* 2010;19:1893–907.
- [2] Ginsberg RJ, Rubinstein LV. Randomized trial of lobectomy versus limited resection for T1 N0 non-small cell lung cancer. Lung Cancer Study Group. *Ann Thorac Surg* 1995;60:615–22; discussion 22–3.
- [3] Warren WH, Faber LP. Segmentectomy versus lobectomy in patients with stage I pulmonary carcinoma. Five-year survival and patterns of intrathoracic recurrence. *J Thorac Cardiovasc Surg* 1994;107:1087–93; discussion 93–4.
- [4] Smith CB, Swanson SJ, Mhango G, Wisnivesky JP. Survival after segmentectomy and wedge resection in stage I non-small-cell lung cancer. *J Thorac Oncol* 2013;8:73–8.
- [5] Kilic A, Schuchert MJ, Pettiford BL, Pennathur A, Landreneau JR, Landreneau JP et al. Anatomic segmentectomy for stage I non-small cell lung cancer in the elderly. *Ann Thorac Surg* 2009;87:1662–6; discussion 67–8.
- [6] Aberle DR, DeMello S, Berg CD, Black WC, Brewer B, Church TR et al. Results of the two incidence screenings in the National Lung Screening Trial. *N Engl J Med* 2013;369:920–31.
- [7] Shapiro M, Weiser TS, Wisnivesky JP, Chin C, Arustamyan M, Swanson SJ. Thorascopic segmentectomy compares favorably with thorascopic lobectomy for patients with small stage I lung cancer. *J Thorac Cardiovasc Surg* 2009;137:1388–93.
- [8] Keenan RJ, Landreneau RJ, Maley RH Jr, Singh D, Macherey R, Bartley S et al. Segmental resection spares pulmonary function in patients with stage I lung cancer. *Ann Thorac Surg* 2004;78:228–33; discussion 28–33.
- [9] Cheng YD, Duan CJ, Dong S, Zhang H, Zhang SK, Wang SQ et al. Clinical controlled comparison between lobectomy and segmental resection for patients over 70 years of age with clinical stage I non-small cell lung cancer. *Eur J Surg Oncol* 2012;38:1149–55.
- [10] Martin-Ucar AE, Nakas A, Pilling JE, West KJ, Waller DA. A case-matched study of anatomical segmentectomy versus lobectomy for stage I lung cancer in high-risk patients. *Eur J Cardiothorac Surg* 2005;27:675–9.
- [11] Tierney JF, Stewart LA, Ghersi D, Burdett S, Sydes MR. Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials* 2007;8:16.
- [12] Parmar MK, Torri V, Stewart L. Extracting summary statistics to perform meta-analyses of the published literature for survival endpoints. *Stat Med* 1998;17:2815–34.
- [13] Yusuf S, Peto R, Lewis J, Collins R, Sleight P. Beta blockade during and after myocardial infarction: an overview of the randomized trials. *Prog Cardiovasc Dis* 1985;27:335–71.
- [14] Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- [15] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- [16] Okada M, Yoshikawa K, Hatta T, Tsubota N. Is segmentectomy with lymph node assessment an alternative to lobectomy for non-small cell lung cancer of 2 cm or smaller? *Ann Thorac Surg* 2001;71:956–60; discussion 61.
- [17] Campione A, Ligabue T, Luzzi L, Ghiribelli C, Voltolini L, Paladini P et al. Comparison between segmentectomy and larger resection of stage IA non-small cell lung carcinoma. *J Cardiovasc Surg (Torino)* 2004;45:67–70.
- [18] Watanabe T, Okada A, Imakiire T, Koike T, Hirono T. Intentional limited resection for small peripheral lung cancer based on intraoperative pathologic exploration. *Jpn J Thorac Cardiovasc Surg* 2005;53:29–35.
- [19] Okada M, Nishio W, Sakamoto T, Uchino K, Yuki T, Nakagawa A et al. Effect of tumor size on prognosis in patients with non-small cell lung cancer: the role of segmentectomy as a type of lesser resection. *J Thorac Cardiovasc Surg* 2005;129:87–93.
- [20] Okumura M, Goto M, Ideguchi K, Tamura M, Sasaki H, Tanaka H et al. Factors associated with outcome of segmentectomy for non-small cell lung cancer: long-term follow-up study at a single institution in Japan. *Lung Cancer* 2007;58:231–7.
- [21] Schuchert MJ, Pettiford BL, Keeley S, D'Amato TA, Kilic A, Close J et al. Anatomic segmentectomy in the treatment of stage I non-small cell lung cancer. *Ann Thorac Surg* 2007;84:926–32; discussion 32–3.
- [22] Iwasaki A, Hamanaka W, Hamada T, Hiratsuka M, Yamamoto S, Shiraishi T et al. Comparison between a case-matched analysis of left upper lobe trisegmentectomy and left upper lobectomy for small size lung cancer located in the upper division. *Thorac Cardiovasc Surg* 2007;55:454–7.
- [23] Sugi K, Kobayashi S, Sudou M, Sakano H, Matsuda E, Okabe K. Long-term prognosis of video-assisted limited surgery for early lung cancer. *Eur J Cardiothorac Surg* 2010;37:456–60.
- [24] Whitson BA, Groth SS, Andrade RS, Maddaus MA, Habermann EB, D'Cunha J. Survival after lobectomy versus segmentectomy for stage I non-small cell lung cancer: a population-based analysis. *Ann Thorac Surg* 2011;92:1943–50.
- [25] Nakamura H, Taniguchi Y, Miwa K, Adachi Y, Fujioka S, Haruki T et al. Comparison of the surgical outcomes of thorascopic lobectomy, segmentectomy, and wedge resection for clinical stage I non-small cell lung cancer. *Thorac Cardiovasc Surg* 2011;59:137–41.
- [26] Zhong C, Fang W, Mao T, Yao F, Chen W, Hu D. Comparison of thorascopic segmentectomy and thorascopic lobectomy for small-sized stage IA lung cancer. *Ann Thorac Surg* 2012;94:362–7.
- [27] Yamashita S, Tokuiishi K, Anami K, Moroga T, Miyawaki M, Chujo M et al. Thorascopic segmentectomy for T1 classification of non-small cell lung cancer: a single center experience. *Eur J Cardiothorac Surg* 2012;42:83–8.
- [28] Soukiasian HJ, Hong E, McKenna RJ Jr. Video-assisted thorascopic trisegmentectomy and left upper lobectomy provide equivalent survivals for stage IA and IB lung cancer. *J Thorac Cardiovasc Surg* 2012;144:S23–6.
- [29] Yendamuri S, Sharma R, Demmy M, Groman A, Hennon M, Dexter E et al. Temporal trends in outcomes following sublobar and lobar resections for small (<2 cm) non-small cell lung cancers—a Surveillance Epidemiology End Results database analysis. *J Surg Res* 2013;183:27–32.
- [30] Hamatake D, Yoshida Y, Miyahara S, Yamashita S, Shiraishi T, Iwasaki A. Surgical outcomes of lung cancer measuring less than 1 cm in diameter. *Interact Cardiovasc Thorac Surg* 2012;15:854–8.
- [31] Carr SR, Schuchert MJ, Pennathur A, Wilson DO, Siegfried JM, Luketich JD et al. Impact of tumor size on outcomes after anatomic lung resection for stage 1A non-small cell lung cancer based on the current staging system. *J Thorac Cardiovasc Surg* 2012;143:390–7.
- [32] Okada M, Koike T, Higashiyama M, Yamato Y, Kodama K, Tsubota N. Radical sublobar resection for small-sized non-small cell lung cancer: a multicenter study. *J Thorac Cardiovasc Surg* 2006;132:769–75.
- [33] Nakamura H, Kawasaki N, Taguchi M, Kabasawa K. Survival following lobectomy vs limited resection for stage I lung cancer: a meta-analysis. *Br J Cancer* 2005;92:1033–7.
- [34] Fan J, Wang L, Jiang GN, Gao W. Sublobectomy versus lobectomy for stage I non-small-cell lung cancer, a meta-analysis of published studies. *Ann Surg Oncol* 2012;19:661–8.
- [35] Shrier I, Boivin JF, Steele RJ, Platt RW, Furlan A, Kakuma R et al. Should meta-analyses of interventions include observational studies in addition to randomized controlled trials? A critical examination of underlying principles. *Am J Epidemiol* 2007;166:1203–9.
- [36] Egger M, Schneider M, Davey Smith G. Spurious precision? Meta-analysis of observational studies. *BMJ* 1998;316:140–4.
- [37] Anello C, Fleiss JL. Exploratory or analytic meta-analysis: should we distinguish between them? *J Clin Epidemiol* 1995;48:109–16; discussion 17–8.