

Realising food webs: downsampling assumptions propagate into dynamic extinction models

Tanya Strydom ¹; Andrew P. Beckerman ¹

Abstract: Dynamic food-web models require realised interaction networks, yet empirical data frequently describe regional metawebs containing many potential interactions that cannot occur simultaneously. Converting metawebs into realised networks is therefore an essential but often overlooked modelling step. Existing approaches implicitly assume different ecological mechanisms governing which interactions are realised, but the consequences of these assumptions remain largely unexplored. Here we compare four ecologically motivated downsampling approaches representing alternative hypotheses of interaction realisation. Using empirical metawebs, we generated ensembles of realised networks and examined how these assumptions influenced network topology, burn-in dynamics and extinction simulations under both topological and dynamical frameworks. Rather than identifying a single optimal downsampling procedure, our objective was to evaluate how alternative ecological assumptions propagate through subsequent analyses. Different realisation processes consistently produced distinct network structures and altered the behaviour of extinction models despite beginning from identical regional interaction pools. These results suggest that realised-network generation should be regarded as an explicit component of ecological modelling rather than a purely technical preprocessing step. As dynamic food-web models become increasingly common, uncertainty arising from assumptions about interaction realisation may be as important to ecological interpretation as the choice of dynamical model itself.

Keywords: food webs

1 Introduction

Ecological networks provide one of the most powerful frameworks for understanding how biodiversity is organised and how ecological communities respond to environmental change. Historically, food-web research focused largely on describing structural properties such as connectance, trophic organisation and degree distributions, revealing remarkably consistent patterns across ecosystems (Williams & Martinez, 2000). Increasingly, however, the focus has shifted from describing network architecture to predicting ecological processes. Mechanistic dynamic food-web models now allow researchers to investigate community stability, biomass dynamics, extinction cascades and ecosystem responses to perturbation by explicitly simulating the transfer of energy through interacting species (Brose et al., 2006; Curtsdotter et al., 2011; Delmas et al., 2017; Yodzis & Innes, 1992). These approaches represent an important advance over purely topological analyses because secondary extinctions emerge from ecological dynamics—including resource limitation, interaction strengths and feedbacks—rather than simply from the structural loss of trophic links. As a result, dynamic food-web models are increasingly used to address questions spanning conservation biology, ecosystem functioning and macroecological responses to environmental change.

Despite these advances, the application of dynamic food web models remains constrained by an often overlooked problem - the nature of the network used as model input. Dynamic simulations require a realised interaction network representing trophic interactions that can occur simultaneously within a community, yet the networks used to construct these models vary considerably in both their provenance and ecological representation. Synthetic networks, such as those generated by the Cascade (Cohen et al., 1985) and Niche (Williams & Martinez, 2000) models, provide useful theoretical abstractions of food web structure and can be readily coupled to dynamic models, but are not intended to represent particular communities. More detailed networks can instead be constructed from empirical observations or inferred using biological information such as body size, energetics and trophic constraints. Mechanistic approaches such as the Allometric Diet Breadth Model (ADBM; Petchey et al. (2008)) and Allometric Trophic Networks (ATNs; Brose et al. (2006)), for example, can generate empirically parameterised networks. However, these models require detailed data (abundances and body sizes) and it remains to be seen if they are able to accurately recover specific pairwise interactions as they do not account for trait-based compatibility between specific species pairs (Strydom, Dunhill, et al., 2026; Strydom, Karapınar, et al., 2026). Conversely, a literature- or inference-derived metaweb only represents feasible pairwise interactions for species in a community (Dunne, 2006).

This distinction is important because empirical provenance does not necessarily imply realised ecological structure. Metawebs can incorporate substantial information about real ecological interactions while aggregating interactions observed or inferred across locations, seasons, sampling periods or environmental contexts (Dunne,

2006). They therefore provide a potentially powerful empirical basis for dynamic modelling, but typically contain more interactions than are realised simultaneously within any particular community. For regional metawebs, exploiting this information in a dynamic framework consequently requires an additional network realisation step - selecting from the pool of potential interactions to generate plausible community-level networks. Such network realisation is more than a computational reduction in connectance; it is an ecological modelling decision that determines which interactions are assumed to occur together.

The need for ‘metaweb realisation’ is particularly apparent when considering ecological systems for which dynamic predictions could be most informative but empirical data are inherently incomplete. Palaeoecological records, for example, can provide unusually valuable opportunities to study the ecological consequences of major extinction events because fossil assemblages can document communities before and after large-scale biodiversity loss (Dunhill et al., 2024; Fricke et al., 2022; Karapınar et al., 2026; Roopnarine, 2006; **roopnarineNetworksExtinctionPaleocommunity2017?**). Yet even where species composition and trophic interactions can be reconstructed, palaeoecological data rarely provide a complete observation of a single realised food web. Instead, they are more naturally used to reconstruct empirically constrained interaction pools from which plausible community networks can be inferred. This creates an opportunity for dynamic food-web modelling - provided metawebs can be converted into dynamically viable realised networks. Empirically-informed ecological change has the potential be used to test mechanistic explanations through comparisons before and after extinction event or more complex hindcasting exercises (Strydom et al., 2021).

However, the process of metaweb realisation itself has the potential to introduce assumptions that may propagate into subsequent ecological inference. In this sense, realised-network generation constitutes an ecological model in its own right, introducing structural assumptions before any population dynamics are simulated. These assumptions may then be compounded by the dynamics themselves. Once a realised network has been generated, dynamic models typically undergo a burn-in period during which biomass redistributes and species unable to persist under the chosen dynamical formulation are lost. Burn-in therefore represents a further filtering step, transforming a structurally feasible network into a dynamically viable community. Only after these successive stages of network realisation and dynamical filtering are perturbations, such as species removals or environmental change, imposed. Ecological inference from dynamic food web models is consequently conditioned not only by the dynamical formulation of the model, but also by assumptions introduced during network construction and by the subsequent process of dynamical realisation.

Here we investigate whether alternative approaches to network realisation influence the behaviour of subsequent ecological models. Using empirical metawebs, we generate ensembles of realised food webs using four ecologically motivated downsampling algorithms representing contrasting hypotheses of interaction realisation, alongside

realised networks generated using Niche and ATN models as additional benchmarks. We then evaluate how these assumptions influence network structure, dynamical burn-in and extinction simulations. Our objective is not to identify an optimal downsampling algorithm, nor to suggest that any single approach provides the correct realised community. Rather, we present network realisation as a proof of concept for integrating empirical interaction pools into dynamic food-web models, while testing whether the ecological signatures introduced during this step remain detectable throughout the subsequent modelling process. More broadly, we ask whether ‘realised metawebs’ can provide a viable bridge between observed ecological interaction structure and mechanistic dynamic modelling, and whether the assumptions required to construct that bridge ultimately become as influential as the ecological dynamics themselves.

2 Methods

2.1 Data

Seven locations (Karapınar et al., 2026). Here we use the guild-level community data to keep the size of the networks manageable and well within the bounds used by Curtsdotter et al. (2011). Additionally we added plankton and zooplankton data from EcoGenie models [REF] as a way to enlarge the ‘base’ of the network (again to better represent a realised network).

2.2 Network construction

For the empirical networks we used the PFWIM model (Shaw et al., 2024). *Insert brief description about trait assignments and mechanistic assumptions, mention we wanted to go as broad as possible so we used fully categorical rules.* This was then used as the base metaweb for each community. Each metaweb was then downsampled to a target connectance (see below for the different downsampling approaches). Additionally the target connectance and species richness was used to parameterise a niche model network (Williams & Martinez, 2000). We also constructed ATN networks (Brose et al., 2006).

2.3 Downsampling approaches

We treat downsampling as a process of **ecological realisation** rather than observation. Starting from a metaweb containing all potential trophic interactions, each downsampling approach represents an alternative ecological hypothesis describing how local ecological processes filter these potential interactions into the subset that is realised within a particular community. The different approaches therefore differ not only in their implementation, but also in the ecological assumptions they embed about how realised food webs emerge

from a common regional interaction pool.

2.3.1 Niche

The original niche model from Williams & Martinez (2000) assumes that realised interactions represent the core of a consumer's trophic niche, and that these diets can be arranged along a single niche axis. In order to create a downsampling approach that mimics this we infer a latent trophic axis directly from the metaweb using Singular Value Decomposition (SVD). It has been shown that the latent spaces described by the left and right singular vectors are closely aligned with the generality and vulnerability of species (Strydom et al., 2022) and so the SVD composition should retain the diet of the species.

Let A denote the $S \times S$ adjacency matrix of the metaweb, where $A_{ij} = 1$ if consumer i feeds on resource j . We compute its singular value decomposition,

$$A = U\Sigma V^\top,$$

where U and V are the left and right singular vectors, respectively, and Σ is the diagonal matrix of singular values. Using the leading singular value, σ_1 , and corresponding singular vectors, $\mathbf{u}_1$ and $\mathbf{v}_1$, each species is assigned latent consumer and resource coordinates,

$$c_i = \sqrt{\sigma_1} u_{i1}, \quad r_i = \sqrt{\sigma_1} v_{i1}.$$

Because species can act both as consumers and as resources, these coordinates are combined to obtain a single niche position,

$$x_i = \frac{c_i + r_i}{2},$$

which is subsequently rescaled to lie between 0 and 1. Species with similar values of x_i therefore occupy similar positions along the dominant trophic gradient of the metaweb.

For each consumer i , the centroid of its realised niche is

$$\mu_i = \frac{1}{k_i} \sum_{j \in R_i} x_j,$$

111 where R_i is the set of resources consumed by species i , and k_i is its trophic breadth (generality). Niche
112 breadth is estimated as the standard deviation of prey positions,

$$\sigma_i = \max(\text{sd}(x_j), 0.1),$$

113 where the lower bound prevents unrealistically narrow niches for specialists. Niche breadth can optionally be
114 scaled to further constrain realised diets,

$$\sigma'_i = \alpha \sigma_i,$$

115 where α is acts as a scaling parameter.

116 The probability of retaining an interaction between consumer i and resource j is then

$$P_{ij} \propto \exp \left[- \left(\frac{x_j - \mu_i}{\sigma'_i} \right)^2 \right].$$

117 Interactions closest to the consumer's niche centre are therefore most likely to be realised (retained), whereas
118 interactions towards the margins of the niche are preferentially removed. Ecologically, this assumes that
119 local communities realise the central portion of a consumer's potential trophic niche, with peripheral or
120 opportunistic interactions occurring less consistently.

121 2.3.2 Power-law

122 The power-law approach assumes that interaction realisation depends primarily on **consumer trophic**
123 **breadth**. Following Roopnarine (2006), the retention probability for consumer i is

$$P_i \propto \exp \left(\frac{k_i}{E} \right),$$

124 where k_i is consumer generality and

$$E = S^{(y-1)/y},$$

125 with S denoting network size and y the scaling exponent. Probabilities are subsequently normalised to lie
126 between 0 and 1. Every interaction belonging to consumer i inherits the same retention probability,

$$P_{ij} = P_i.$$

Ecologically, this represents the hypothesis that generalist consumers realise a larger fraction of their potential trophic niche than specialist consumers, irrespective of the identity of individual resources.

2.3.3 Degree-product

The degree-product approach assumes that interaction realisation depends on the centrality of both interacting species. Each interaction is assigned a weight

$$w_{ij} = k_i^{\text{out}} k_j^{\text{in}},$$

where k_i^{out} is the consumer's out-degree (generality) and k_j^{in} is the resource's in-degree (vulnerability).

Retention probabilities are then calculated as

$$P_{ij} = 1 - \frac{w_{ij}}{\max(w)},$$

before being constrained to the interval $[0, 1]$. Under iterative downsampling, degrees are recalculated following each interaction removal, allowing probabilities to evolve as the realised network changes. Ecologically, this assumes that highly connected species possess many potential interactions that are not simultaneously realised within any local community, whereas interactions involving more specialised or weakly connected species are more likely to represent essential realised links.

2.3.4 Random

Random downsampling provides the ecological null model. Every interaction has an equal probability of being retained,

$$P_{ij} = c,$$

where c is constant for all existing interactions. Interactions are removed uniformly at random until the target connectance is reached, subject to the constraint that species cannot become isolated. This assumes that no ecological process preferentially favours the realisation of one potential interaction over another, such that

realised food webs constitute an unbiased subset of the regional metaweb that is ultimately only constrained by the number of links.

2.4 Simulations

For each empirical community, we generated 30 replicate network ensembles. Species body masses were sampled independently from truncated log-normal distributions constrained by their categorical body-size classifications. Initial biomasses were drawn from a uniform distribution. These trait values were used to construct a series of alternative networks, representing different two realised networks (ATN and niche model), one metaweb, and four downsampled (realised) metawebs (niche, power-law, degree-product, and random).

For the allometric trophic network (ATN), interaction strengths were generated by varying the feeding threshold between 0.01 and 0.12 and selecting the network whose connectance most closely matched a target connectance. Target connectance was independently sampled for each replicate from a truncated normal distribution bounded between 0.05 and 0.15. The target connectance and richness of the metaweb were used to build a niche model and the metaweb was downsampled until the target connectance was reached. These represent the ‘creation’ networks.

Each creation network was then parameterised using the END package (Delmas et al., 2017) and simulated for up to 5000 time steps. Simulations terminated early if a dynamical equilibrium was reached. Species whose biomass declined below the survival threshold (10^{-12}) during this burn-in period were considered extinct and removed from the network. The surviving species and their interactions constituted the corresponding ‘realised’ networks.

Extinction experiments were subsequently performed on both the creation and realised networks. Topological extinction simulations were conducted on both network stages, whereas dynamic extinction simulations were performed only on realised networks. In all simulations, primary extinctions followed one of the prescribed extinction scenarios. During topological simulations, secondary extinctions occurred when species lost all prey, these were allowed to cascade. During dynamic simulations, primary extinctions were imposed by setting the biomass of the focal species to the survival threshold (10^{-12}), after which the community was simulated for a further 5000 time steps to allow biomass dynamics to relax. Species whose biomass subsequently declined below the survival threshold were recorded as secondary extinctions. This procedure was repeated until the entire community collapsed.

For each extinction scenario, robustness was quantified from the resulting extinction curves as the proportion of primary extinctions required to produce a 50% reduction in species richness (R_{50} (Jonsson et al., 2015)),

estimated by linear interpolation between extinction events.

2.5 Software

Network construction and extinction simulations were run in Julia v1.12.6 (Bezanson et al., 2017), we used `EcologicalNetworksDynamics.jl` (Lajaaity et al., 2025) for dynamic simulations, `PFIM.jl` [REF] for construction of metawebs, `FoodWebTools.jl` [REF] for network downsampling, and `Extinctions.jl` [REF] for extinctions and robustness calculations. Statistical analyses were run in R v4.5.2 [REF]. All code can be found at *TODO*

Keep the narrative extinction centric for now but remember we also have the topology of each network before the various extinction simulations begin

3 Downsampling and burn-in alter network topology in method-dependent ways

Redundancy analysis (RDA), using permutation-based significance testing while accounting for community-level variation, revealed that downsampling approach significantly affected network topology, but that this effect depended on the magnitude of connectance reduction required to reach the target connectance (999 permutations; $F_{7,707} = 112.8$, $p = 0.001$; Figure 1, panel A). The model explained 31.8% of total topology variation (adjusted $R^2 = 0.318$), with community identity accounting for an additional 39.7% of variation. The significant interaction between downsampling approach and connectance change (marginal test; $F_{3,707} = 18.90$, $p = 0.001$) demonstrated that topology trajectories differed among downsampling methods as connectance declined. Thus, the extent to which network structure diverged from the original topology depended on both the severity of downsampling and the mechanism by which interactions were removed.

[Figure 1 about here.]

To identify the source of this interaction, the response of the primary RDA axis was examined across the connectance gradient. RDA1 was strongly associated with connectance change ($F_{1,712} = 2076.86$, $p < 0.001$), with additional effects of downsampling approach ($F_{3,712} = 33.28$, $p < 0.001$) and a significant downsampling approach $\times$ connectance change interaction ($F_{3,712} = 4.81$, $p = 0.003$). Pairwise comparisons of estimated trajectory slopes showed that link-based downsampling followed a distinct trajectory, differing from niche-, power-, and random-based downsampling, whereas the latter three approaches exhibited statistically indistinguishable responses (Figure 1, panel B). These results indicate that all downsampling methods

alter network topology as connectance declines, but that removing interactions according to their realised connectivity produces a fundamentally different pathway of structural change from approaches that incorporate species-level ecological information or random link loss.

We next asked whether these structurally distinct networks responded differently during dynamic burn-in. Structural reorganisation was quantified as the Euclidean displacement between network positions before and after burn-in in the first three principal component dimensions, with community included as a random effect to account for the substantial variation among communities. Burn-in displacement differed significantly among network types ($F = 62.0$, $p < 0.001$). Metawebs underwent the greatest structural reorganisation, whereas ATNs exhibited the smallest displacement. Among the downsampled networks, niche-, power-, and link-based approaches underwent comparable structural change during burn-in, while random downsampling showed a modestly greater displacement (Figure 1, panels C–D).

Together, these analyses demonstrate that downsampling approach influences network topology both during network construction and subsequent dynamic relaxation. Differences among downsampling methods become increasingly pronounced as greater proportions of interactions are removed, consistent with the expectation that the ecological assumptions underlying each removal strategy exert a stronger influence under more aggressive downsampling. Importantly, despite generating distinct initial topologies, the different downsampling approaches underwent remarkably similar structural reorganisation during burn-in, changing in ways more similar to the niche model than the original metaweb. In contrast, random downsampling remained distinct throughout burn-in, consistent with its lack of an underlying ecological mechanism governing interaction removal. This suggests that the ecologically informed downsampling on metawebs allows us to produce networks whose subsequent structural dynamics more closely resemble those of the niche model. Notably the two realised models (Niche and ATN) do not behave similarly and still end up at different ‘end points’ post burn-in to the empirical webs, highlighting that network reconstruction approach will still have a strong effect on the inferred dynamics and behaviour of the system (Strydom, Karapınar, et al., 2026)

4 Differences in inferred robustness of topological and dynamic are unaffected by network type

To determine whether the qualitative differences between topological and dynamic extinction simulations previously demonstrated using niche-model food webs generalise across alternative network reconstructions, we compared the difference in robustness (ΔR_50) among network types for each extinction scenario while accounting for community-level variation. Network type significantly affected ΔR_50 in all four extinction

scenarios (low-to-high biomass: $F_{6,1248} = 22.3$, $p < 0.001$; high-to-low biomass: $F_{6,1248} = 46.7$, $p < 0.001$; random basal: $F_{6,1248} = 109.0$, $p < 0.001$; random consumer: $F_{6,1248} = 22.9$, $p < 0.001$), indicating that the magnitude of the difference between topological and dynamic robustness depended on network reconstruction.

Despite these quantitative differences, the qualitative pattern identified by Curtsdotter et al. (2011) was largely maintained. Under all but random basal removal, topological extinction simulations inferred greater robustness than dynamic simulations for every network type, although the magnitude of this discrepancy varied among reconstruction methods Figure 2. Random basal extinction represented the only notable departure from this general pattern. Here, the discrepancy between topological and dynamic robustness was substantially reduced, with link- and power-downsampled networks exhibiting slightly positive values of ΔR_{50} and random downsampled networks showing little difference between the two approaches, whereas niche-derived networks retained lower dynamic than topological robustness. Thus, the conclusion that topological analyses provide more conservative estimates of robustness was not universal, but remained remarkably robust across reconstruction methods and extinction scenarios.

[Figure 2 about here.]

Together, these analyses demonstrate that the principal conclusion of Curtsdotter et al. (2011) extends well beyond the niche model. While the precise magnitude of the discrepancy between topological and dynamic robustness depends on both network reconstruction and extinction mechanism, the qualitative tendency for topological extinction simulations to infer greater robustness than dynamic simulations was recovered across the vast majority of reconstructed food webs examined here.

Dynamic extinctions re-enforce ecological assumptions

To evaluate how interaction filtering influences dynamic behavior, we calculated pairwise contrasts of inferred robustness (R_{50}) across all combinations of community, extinction mechanism, and network type. In aggregate, topological extinctions (evaluated both at network creation and after dynamic burn-in (realisation)) yielded consistent robustness estimates across all empirically derived networks (Figure 3, left column). This structural cohesion was further reflected in hierarchical clustering, where empirically derived networks consistently formed a distinct cluster separate from the two synthetic benchmark networks (niche model and ATN; Figure 3, right column).

[Figure 3 about here.]

However, under dynamic extinction simulations, this grouping broke down. Introducing dynamic biomass interactions caused a distinct reshuffling in the relative relationships among network types (Figure 3, bottom row). Most notably, the downsampled niche networks yielded robustness estimates that converged with those

of the synthetic niche model, whereas the remaining empirically derived networks maintained their mutual similarity. Additionally, the ATN model shifted to become more similar in robustness to the empirical cluster. When examining topological extinctions, it is clear why empirically derived networks behave similarly to one another - they share a common baseline architecture inherited from the regional metaweb (Figure 1, panel B), and topological cascades depend strictly on network topology. For dynamic extinctions, however, network behaviour is not determined solely by initial structure. Instead, higher-order differences emerge from how functional parameterisation interacts with network topology. In the bio-energetic food web (BEFW) framework, species body mass dictates key functional parameters, including metabolic and consumption rates (Delmas et al., 2017). Because species body masses were assigned based on trophic position, the dynamic convergence of the synthetic niche model and the SVD-downsampled niche network likely reflects their shared trophic organisation. Consequently, dynamic simulations do not merely evaluate network structure; they actively re-enforce the ecological assumptions embedded during network construction.

5 Conclusions

Reconstructing local food webs from regional metawebs requires explicit assumptions about how ecological processes filter potential interactions into realised local networks. By comparing four downsampling approaches across topological and dynamic extinction scenarios, we demonstrate that the mechanism of interaction removal influences both the structural topology of the resulting networks and their inferred robustness to extinction.

5.1 Structural vs. dynamic fragility

Our results confirm that the qualitative discrepancy between topological and dynamic robustness identified by Curtsdotter et al. (2011) extends across alternative network reconstructions. Under most extinction scenarios, topological simulations inferred greater robustness than dynamic simulations regardless of downsampling approach. Topological extinctions treat networks as static topological dependency graphs, whereas dynamic extinctions incorporate biomass drawdown, competitive release, and rate-dependent secondary collapses. Consequently, static metrics consistently provide a conservative estimate of community vulnerability.

Importantly, while downsampling methods generated distinct initial topologies, their effects on topological robustness were relatively modest—likely because initial metawebs were already relatively sparse (mas Co = 0.20). However, once bio-energetic dynamics were introduced, subtle structural differences translated into substantial variations in dynamic persistence. Dynamic extinctions are strongly driven by node-level properties and metabolic rates rather than link topology alone. Thus, inferred dynamic robustness reflects an

interaction between network structure and the physiological constraints imposed upon it.

5.2 Mapping empirical networks or model philosophy?

A central question arising from our findings is whether dynamic simulations evaluate empirical reality or simply mirror the assumptions of the underlying models (Strydom, Karapınar, et al., 2026). The dynamic convergence between SVD-downsampled metawebs and synthetic niche models highlights a fundamental circularity - because frameworks like the BEFW rely heavily on body-mass ratios and trophic positioning to derive interaction rates, networks constructed around niche-ordering assumptions naturally align with these dynamic engines. Conversely, downsampling methods that do not enforce trophic hierarchy (such as degree-product or random link removal) introduce structural configurations that interact differently with allometric scaling rules, leading to divergent dynamic behaviors.

This distinction is particularly evident when comparing the ATN and empirical downsampled networks. Although the ATN model is widely used as a realistic benchmark, its dynamic behaviour diverged from both the niche model. This underscores that no single synthetic network model can be treated as an absolute benchmark for realised food-web dynamics.

5.3 Implications for network reconstruction

While there is no single correct downsampling approach, our results offer practical considerations for empirically-derived metaweb downsampling:

- Alignment with dynamic frameworks: If the objective is to parameterise dynamic biomass simulations using allometric or bio-energetic models, SVD-based niche downsampling provides an internally consistent approach that preserves trophic hierarchy without imposing arbitrary degree distributions.
- Caution with unconstrained null models: Random downsampling introduces topological distortions that increase structural displacement during burn-in (Figure 1) and generate unpredictable dynamic responses, making it unsuitable as an ecologically informative filter.
- Evaluating static and dynamic stability together: Because topological extinctions represent an upper bound of structural robustness and dynamic extinctions reflect rate-dependent persistence, reporting both provides a clearer picture of network robustness.

Ultimately, downsampling a metaweb is an act of modelling ecological realisation rather than simple network reduction. Acknowledging the assumptions embedded within these downsampling filters is essential for accurately inferring community robustness in both ancient and modern ecosystems.

References

- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. (2017). Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>
- Brose, U., Jonsson, T., Berlow, E. L., Warren, P., Banasek-Richter, C., Bersier, L.-F., Blanchard, J. L., Brey, T., Carpenter, S. R., Blandenier, M.-F. C., Cushing, L., Dawah, H. A., Dell, T., Edwards, F., Harper-Smith, S., Jacob, U., Ledger, M. E., Martinez, N. D., Memmott, J., ... Cohen, J. E. (2006). Consumer–Resource Body-Size Relationships in Natural Food Webs. *Ecology*, 87(10), 2411–2417. [https://doi.org/10.1890/0012-9658\(2006\)87%5B2411:CBRINF%5D2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87%5B2411:CBRINF%5D2.0.CO;2)
- Cohen, J. E., Newman, C. M., & Steele, J. H. (1985). A stochastic theory of community food webs I. Models and aggregated data. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 224(1237), 421–448. <https://doi.org/10.1098/rspb.1985.0042>
- Curtsdotter, A., Binzer, A., Brose, U., De Castro, F., Ebenman, B., Eklöf, A., Riede, J. O., Thierry, A., & Rall, B. C. (2011). Robustness to secondary extinctions: Comparing trait-based sequential deletions in static and dynamic food webs. *Basic and Applied Ecology*, 12(7), 571–580. <https://doi.org/10.1016/j.baae.2011.09.008>
- Delmas, E., Brose, U., Gravel, D., Stouffer, D. B., & Poisot, T. (2017). Simulations of biomass dynamics in community food webs. *Methods in Ecology and Evolution*, 8(7), 881–886. <https://doi.org/10.1111/2041-210X.12713>
- Dunhill, A. M., Zarzychny, K., Shaw, J. O., Atkinson, J. W., Little, C. T. S., & Beckerman, A. P. (2024). Extinction cascades, community collapse, and recovery across a Mesozoic hyperthermal event. *Nature Communications*, 15(1), 8599. <https://doi.org/10.1038/s41467-024-53000-2>
- Dunne, J. (2006). The network structure of food webs. In *Ecological Networks: Linking Structure to Dynamics in Food Webs* (pp. 27–86).
- Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning, J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene. *Science*, 377(6609), 1008–1011. <https://doi.org/10.1126/science.abn4012>
- Jonsson, T., Berg, S., Pimenov, A., Palmer, C., & Emmerson, M. (2015). The reliability of R50 as a measure of vulnerability of food webs to sequential species deletions. *Oikos*, 124(4), 446–457. <https://doi.org/10.1111/oik.01588>
- Karapınar, B., Strydom, T., Beckerman, A. P., Ridgwell, A., Wignall, P. B., Dunne, J. A., Little, C. T. S., Hull, P., Pimiento, C., & Dunhill, A. (2026, February 25). *No global collapse of food webs across the Permian-Triassic Mass Extinction*. <https://doi.org/10.64898/2026.02.24.707709> (Pre-published)
- Lajaaity, I., Bonnici, I., Kéfi, S., Mayall, H., Danet, A., Beckerman, A. P., Malpas, T., & Delmas, E. (2025).

EcologicalNetworksDynamics.jl: A Julia package to simulate the temporal dynamics of complex ecological networks. *Methods in Ecology and Evolution*, 16(3), 520–529. <https://doi.org/10.1111/2041-210X.14497>

Petchey, O. L., Eklöf, A., Borrvall, C., & Ebenman, B. (2008). Trophically Unique Species Are Vulnerable to Cascading Extinction. *The American Naturalist*, 171(5), 568–579. <https://doi.org/10.1086/587068>

Roopnarine, P. D. (2006). Extinction Cascades and Catastrophe in Ancient Food Webs. *Paleobiology*, 32(1), 1–19. <https://www.jstor.org/stable/4096814>

Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024, January 30). *A framework for reconstructing ancient food webs using functional trait data*. <https://doi.org/10.1101/2024.01.30.578036> (Pre-published)

Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*, 13(12). <https://doi.org/10.1111/2041-210X.13835>

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz, N. R., Higinio, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap towards predicting species interaction networks (across space and time). *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>

Strydom, T., Dunhill, A. M., Dunne, J. A., Poisot, T., & Beckerman, A. P. (2026). *From Metawebs to Realised Webs: A Framework for Ecological Network Representation under Global Change*. <https://ecoevorxiv.org/repository/view/11514/>

Strydom, T., Karapınar, B., Dunne, J. A., Hull, P., Little, C. T. S., Pimiento, C., Ridgwell, A., Beckerman, A. P., & Dunhill, A. M. (2026). *Are We Mapping Ecosystems or Models? Framework Choices Dominate Food Web Topology and Extinction Inferences*. <https://ecoevorxiv.org/repository/view/13495/>

Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, 404(6774), 180–183. <https://doi.org/10.1038/35004572>

Yodzis, P., & Innes, S. (1992). Body Size and Consumer-Resource Dynamics. *The American Naturalist*, 139(6), 1151–1175. <https://doi.org/10.1086/285380>




Figure 1: **Downsampling approach influences both network topology and subsequent structural reorganisation during burn-in.** (A) Redundancy analysis (RDA) ordination showing differences in network topology among downsampling approaches across the connectance gradient after accounting for community-level variation. (B) Estimated slopes ($\pm 95\%$ confidence intervals) describing the relationship between the primary RDA axis and connectance change for each downsampling approach; letters denote post-hoc groupings




Figure 2: **Differences between topological and dynamic estimates of network robustness.** Boxplot showing aggregated realised topological and dynamic robustness for the different network types. Each extinction scenario is presented in the supp matt but note that the with the exception of the random basal extinctions this pattern remains consistent




Figure 3: TODO