From metawebs to dynamics: How food webs are built shapes how they behave

Tanya Strydom [1*], **Baran Karapunar** [2], **Andrew P. Beckerman** [1], **Alexander M. Dunhill** [2]

[1] School of Biosciences, University of Sheffield, Sheffield, UK
[2] School of Earth, Environment and Sustainability, University of Leeds, Leeds, UK, LS2 9JT

[*]Correspondence: t.strydom@sheffield.ac.uk

Abstract: Dynamic food web models provide a mechanistic framework for understanding ecological stability, extinction cascades, and responses to environmental change, but their predictions depend on the network supplied as model input. Empirical metawebs provide rich information on potential species interactions yet contain interactions that may not occur simultaneously within a community. Converting these interaction pools into realised networks therefore requires assumptions about which interactions are likely to co-occur, potentially introducing uncertainty before population dynamics are simulated. Here, we investigate how alternative approaches to network realisation influence network structure and subsequent ecological dynamics. Using empirical metawebs, we generated realised food webs using four ecologically motivated downsampling approaches, alongside networks generated using Niche and allometric trophic network models as reference approaches. We then compared network topology, structural change during dynamical burn-in, and inferred robustness under topological and dynamic extinction simulations. Downsampling approach altered network topology, with differences becoming increasingly pronounced as greater proportions of interactions were removed. Dynamical burn-in further reorganised network structure but did not erase the signatures introduced during network construction; instead, downsampled networks underwent structural changes that were often more like the niche model than the original metaweb. Despite these differences, the broad tendency for topological extinction simulations to infer greater robustness than dynamic simulations were consistent across network reconstruction approaches. However, dynamic simulations revealed differences in network behaviour that were not apparent from topology alone, including convergence of niche-based downsampling towards the behaviour of the synthetic niche model. These results demonstrate that network realisation is not simply a preprocessing step, but an additional source of ecological model uncertainty. Dynamic food web predictions are therefore shaped by successive assumptions about which interactions are realised and which of those realised networks can persist dynamically, highlighting the importance of explicitly considering network reconstruction when using empirical interaction data to make mechanistic predictions about ecological change.

Keywords: food webs, ecological networks, network reconstruction, network realisation, metawebs, dynamical modelling, extinction cascades, ecological model uncertainty

1 Introduction

Understanding how the loss of individual species propagates through ecological communities is a central challenge in ecology. The loss of a species triggers secondary extinctions that propagate throughout the food web. Being able to predict extinction cascades resulting from the removal of a specific species requires more than knowing how many interactions occur within a community or how species are connected. Rather it requires understanding which specific species interact, and how those particular interactions underpin the persistence of other species. Food web approaches to studying extinction have traditionally addressed this problem through topological simulations, in which species are sequentially removed and secondary extinctions are inferred from the structure of the network. These approaches provide a tractable, but simplified means of assessing community robustness as they are only driven by bottom-up processes and do not explicitly account for the population processes governing species persistence (Curtsdotter et al., 2011). Dynamic food web models offer a complementary approach by simulating biomass changes and energy transfer among interacting species, allowing secondary extinctions to emerge from changes in resource availability and population dynamics (Brose et al., 2006; Delmas et al., 2017; Yodzis & Innes, 1992). However, the ecological realism of these predictions depends fundamentally on the network supplied to the model. In particular, whether the species interactions represented in the network adequately reflect those occurring within the community.

A limitation of existing approaches to dynamic food web modelling is that networks are commonly generated using theoretical rules rather than empirical knowledge of species interactions. The niche model, for example, generates plausible food web structures by arranging species along a continuous niche axis, with consumers feeding within a defined region of that axis (Williams & Martinez, 2000). This approach reproduces several characteristic properties of empirical food webs and provides a convenient framework for coupling network structure to population dynamics. However, it is fundamentally species-agnostic where interactions are generated from generalised structural rules rather than the known ecological relationships between particular species. Although the resulting networks may reproduce realistic networks structure they do not necessarily represent the interactions that actually occur within a given community. Allometric trophic network models provide a step towards greater biological realism by incorporating consumer–resource body-size relationships and associated assumptions about feeding interactions (Brose et al., 2006;

Petchey et al., 2008). However body size is only one component of trophic feasibility. Species may differ in their feeding strategies, habitat requirements, functional morphology and other ecological characteristics that constrain their interactions [REF]. Networks generated primarily from body-size relationships may therefore include interactions that are theoretically plausible but biologically impossible *i.e.* predict forbidden links. While these approaches provide useful representations of food web structure and trophic processes, they do not necessarily preserve the species-specific interaction identity required to reconstruct realistic, empirically grounded, extinction pathways.

Empirical metawebs offer an opportunity to address this limitation. By compiling potential trophic interactions among particular species from observations, published literature and trait-based inference, metawebs retain information about the identity of consumer–resource relationships that is absent from species-agnostic network models (Dunne, 2006). This distinction is particularly important for extinction modelling. The loss of a species does not simply remove a node with a given number of connections, it removes a particular set of ecological relationships, potentially affecting consumers in ways that depend on the identity and availability of alternative resources. Empirically constrained interactions therefore provide a more biologically meaningful basis for predicting how the loss of a specific species propagates through a community. However, metawebs represent pools of potential interactions rather than necessarily realised community networks. Because interactions may have been recorded or inferred across different locations, environmental conditions and periods, they need not all occur simultaneously within a single ecological community (Strydom, Dunhill, et al., 2026). Thus the distribution of links within a metaweb are not constrained by community context. Directly using metawebs as inputs to population dynamic models is a philosophical misalignment as the resulting structure of the network does not necessarily represent a system that is compatible with the energetic assumptions of dynamic models.

Translating empirical metawebs into networks suitable for dynamic modelling therefore requires a means of distinguishing potential interactions from those likely to be realised within a particular community. Downsampling provides a promising approach to this problem by reducing the number of interactions while retaining the identity of the species involved [EG ROOPNARINE]. Rather than generating interactions from generalised structural rules, downsampling starts with an empirically constrained pool of potential interactions and selects a subset to represent a plausible realised food web. This offers a

potential bridge between empirical interaction data and mechanistic dynamic modelling, retaining species-specific trophic relationships (and importantly no forbidden links) while producing networks with more realistically constrained structures. However, the selection of interactions to remove or retain is not necessarily neutral. Different downsampling approaches will embody different assumptions about how potential interactions are filtered into realised communities. These different assumptions may preserve different subsets of species interactions, with potentially important consequences for the structure of the resulting food web, the secondary extinction pathways that it supports, and importantly the exact identity of species lost. Thus, although downsampling provides a means of incorporating empirical interaction identity into dynamic models, the ecological assumptions used to select interactions may themselves influence subsequent predictions.

Importantly, the construction of a network is not the only process determining which interactions ultimately contribute to community persistence. Once a network has been parameterised and subjected to population dynamics, an initial burn-in period allows biomass to change and species unable to persist under the specified model conditions to become extinct. This process effectively imposes a second filter on the network, transforming the initially constructed topology into a dynamically viable community. Interactions that are retained during downsampling may therefore not necessarily persist as part of the resulting community, because their associated species may fail to maintain viable populations. Conversely, the removal of particular species during burn-in can alter the structure of the surviving network. The resulting community reflects both the assumptions used to select interactions from the original metaweb and the constraints imposed by the population dynamics themselves. Understanding how these successive filters influence network structure is therefore important for establishing whether the empirical information retained during downsampling persists through dynamic modelling, and whether alternative approaches to network construction ultimately produce different ecological outcomes.

The importance of these considerations becomes particularly apparent when comparing topological and dynamic approaches to extinction robustness. Topological simulations infer secondary extinctions from the loss of trophic connections, whereas dynamic simulations additionally account for changes in biomass and the rate-dependent processes governing species persistence. Previous work has demonstrated that these approaches can produce different estimates of food web robustness, with topological simulations generally indi-

cating greater robustness than dynamic simulations (Curtsdotter et al., 2011). However, this comparison has largely been developed using theoretically constructed food webs, leaving it unclear whether the same conclusions extend to networks retaining empirical, species-specific interaction identity. Moreover, the extent to which network reconstruction influences the relationship between topological and dynamic extinction responses remains poorly understood. Networks with similar structural properties may differ in the identity of their interactions, while networks derived from the same empirical metaweb may retain different subsets of potential interactions following downsampling. These differences may be inconsequential for some topological measures of robustness but become important when population dynamics determine how species respond to the loss of particular resources. Comparing extinction responses across alternative network representations therefore provides an opportunity to establish whether incorporating empirical interaction identity changes inferred robustness, and whether dynamic modelling reveals differences that are not apparent from network topology alone.

probably need a version of the paragraph below..

These questions are particularly relevant when modelling ecological communities for which complete observations of realised food webs are unavailable, including palaeoecological systems. Fossil assemblages provide opportunities to investigate ecological change across major extinction events, but the interactions reconstructed from the fossil record are more appropriately understood as empirically constrained pools of potential trophic relationships than complete representations of a single realised community (Dunhill et al., 2024; Fricke et al., 2022; Karapunar et al., 2026; Roopnarine, 2017; Roopnarine, 2006). Developing approaches that translate these interaction pools into plausible, dynamically viable food webs could therefore provide a means of moving beyond structural reconstructions towards mechanistic explanations of ecological change, including the processes underlying extinction cascades and community collapse. More broadly, integrating empirical interaction identity with population dynamics represents an important step towards making predictions from ecological network models more representative of the communities they are intended to describe (Strydom, Catchen, et al., 2021).

Here, we investigate how empirically informed network reconstruction influences food web structure and inferred extinction robustness under topological and dynamic extinction simulations. Using empirical metawebs, we generate community-level networks through four alternative downsampling approaches and compare these with networks constructed

using the niche and allometric trophic network models. By retaining species-specific interactions from empirical metawebs, our approach provides a means of incorporating interaction identity into dynamic extinction modelling while investigating the consequences of alternative assumptions about how potential interactions are realised. Specifically, we address three questions. First, how does downsampling approach influence network topology, and to what extent do these structural differences persist following dynamical burn-in? Second, does the difference between topological and dynamic estimates of extinction robustness generalise across alternative approaches to network construction and extinction scenarios? Third, do alternative network types retain similar relationships in inferred robustness under topological and dynamic extinction simulations, or does incorporating population dynamics reveal differences that are not apparent from topology alone? Rather than identifying a single optimal reconstruction method, we use these comparisons to investigate the extent to which empirical interaction identity can be retained through dynamic modelling, and how assumptions made during network construction propagate into subsequent ecological predictions. We show that downsampling approaches produce increasingly distinct network topologies as greater proportions of interactions are removed, that dynamical burn-in acts as an additional structural filter, and that the broad difference between topological and dynamic estimates of robustness is maintained across alternative network representations. However, dynamic simulations also reveal differences in network behaviour that are not apparent from topology alone. Together, these findings demonstrate the potential of empirical metawebs and downsampling to bridge the gap between species-specific interaction data and mechanistic extinction modelling, while highlighting network reconstruction as an important source of uncertainty in ecological predictions.

2 Methods

2.1 Fossil data

We used community data from seven locations spanning the Permian-Triassic mass extinction (PTME) event compiled by Karapunar et al. (2026). This dataset comprises community-level trophic data representing potentially interacting taxa across seven locations across the Permian-Triassic mass extinction (PTME; ~250 Ma) event. Here we used the guild-level assemblages from the pre-extinction (Permian) time point along with the plankton and zooplankton data from the outputs of the EcoGEnIE models (Ward et

al., 2018) also used in Karapunar et al. (2026). These represent distinct communities of potentially interacting guilds. We excluded the community from Russia because it contained no intermediate species and was therefore unsuitable for the subsequent extinction analyses as it would produce distinct results. The seven locations act as replicates to better determine if observed extinction patterns are driven by specific network structures (location) or if results are more generalisable. Detailed descriptions of the underlying community data, including trait assignment, their composition and spatial structure, are provided in Karapunar et al. (2026).

2.2 Network construction

Metawebs were constructed using the Paleo Food Web Inference Model (PFWIM; Shaw et al. (2024)). The PFWIM applies a set of mechanistic feasibility rules to consumer–resource pairs based on different trait classes. An interaction is deemed as being feasible only when all trait combinations satisfy the specified rules. These metawebs were subsequently downsampled to the specified target connectance (see Section 2.3 for details on downsampling). Using the community data we also constructed networks using the allometric trophic network (ATN) model (Brose et al., 2006), which represents a realised but empirically grounded network. For the ATN species body masses were sampled independently from a truncated log-normal distribution constrained by species categorical body-size classifications. Interaction probabilities were generated by varying the feeding threshold between 0.01 and 0.12 and selecting the network whose connectance most closely matched a target connectance. Finally for each community we also constructed a niche model network (Williams & Martinez, 2000) using the guild richness and target connectance as input parameters (see Section 2.4 for details on parametrisation)

2.3 Downsampling approaches

We treat downsampling as an assumption on the process of ecological realisation rather than observation. Starting from a metaweb containing all potential trophic interactions, each downsampling approach represents an alternative ecological hypothesis describing how local ecological processes may filter these potential interactions into the subset that is realised within a particular community. The different approaches therefore differ not only in their implementation, but also in the ecological assumptions they embed about how realised food webs emerge from a common regional interaction pool.

2.3.1 Niche

The original niche model from Williams & Martinez (2000) assumes that realised interactions represent the core of a consumer's trophic niche, and that these diets can be arranged along a single niche axis. In order to create a downsampling approach that mimics this we infer a latent trophic axis directly from the metaweb using Singular Value Decomposition (SVD). It has been shown that the latent spaces described by the left and right singular vectors are closely aligned with the generality and vulnerability of species (Strydom et al., 2022) and so the SVD composition should retain the dietary signal of the species. This is done as follows:

Let A denote the $S \times S$ adjacency matrix of the metaweb, where $A_{ij} = 1$ if consumer i feeds on resource j. We compute its singular value decomposition,

$$A = U\Sigma V^{\top}$$

where U and V are the left and right singular vectors, respectively, and Σ is the diagonal matrix of singular values. Using the leading singular value, σ_1, and corresponding singular vectors, $\mathbf{u}_1$ and $\mathbf{v}_1$, each species is assigned latent consumer and resource coordinates,

$$c_i = \sqrt{\sigma_1}\, u_{i1}, \qquad r_i = \sqrt{\sigma_1}\, v_{i1}$$

Because species can act both as consumers and as resources, these coordinates are combined to obtain a single niche position,

$$x_i = \frac{c_i + r_i}{2}$$

which is subsequently rescaled to lie between 0 and 1. Species with similar values of x_i therefore occupy similar positions along the dominant trophic gradient of the metaweb.

For each consumer i, the centroid of its realised niche is

$$\mu_i = \frac{1}{k_i} \sum_{j \in R_i} x_j$$

where R_i is the set of resources consumed by species i, and k_i is its trophic breadth (generality). Niche breadth is estimated as the standard deviation of prey positions,

$$\sigma_i = \max\left(sd(x_j),\, 0.1\right)$$

where the lower bound prevents unrealistically narrow niches for specialists. Niche breadth can optionally be scaled to further constrain realised diets,

$$\sigma_i' = \alpha\sigma_i$$

where α is acts as a scaling parameter.

The probability of retaining an interaction between consumer i and resource j is then

$$P_{ij} \propto \exp\left[-\left(\frac{x_j - \mu_i}{\sigma_i'}\right)^2\right]$$

Interactions closest to the consumer's niche centre are therefore most likely to be realised (retained), whereas interactions towards the margins of the niche are preferentially removed. Ecologically, this assumes that local communities realise the central portion of a consumer's potential trophic niche, with peripheral or opportunistic interactions occurring less consistently.

2.3.2 Power-law

The power-law approach assumes that interaction realisation depends primarily on consumer trophic breadth. Following Roopnarine (2006), the retention probability for consumer i is

$$P_i \propto \exp\left(\frac{k_i}{E}\right)$$

where k_i is consumer generality and

$$E = S^{(y-1)/y}$$

with S denoting network size and y the scaling exponent. Probabilities are subsequently normalised to lie between 0 and 1. Every interaction belonging to consumer i inherits the same retention probability,

$$P_{ij} = P_i$$

Ecologically, this represents the hypothesis that generalist consumers realise a larger fraction of their potential trophic niche than specialist consumers, irrespective of the identity of individual resources.

2.3.3 Degree-product

The degree-product approach assumes that interaction realisation depends on the centrality of both interacting species. Here each interaction is assigned a weight

$$w_{ij} = k_i^{out} k_j^{in}$$

where k_i^{out} is the consumer's out-degree (generality) and k_j^{in} is the resource's in-degree (vulnerability). Retention probabilities are then calculated as

$$P_{ij} = 1 - \frac{w_{ij}}{\max(w)}$$

before being constrained to the interval $[0, 1]$. Under iterative downsampling, degrees are recalculated following each interaction removal, allowing probabilities to evolve as the realised network changes. Ecologically, this assumes that highly connected species possess many potential interactions that are not simultaneously realised within any local community, whereas interactions involving more specialised species are more likely to represent essential realised links.

2.3.4 Random

Random downsampling provides the ecological null model. Every interaction has an equal probability of being retained,

$$P_{ij} = c$$

where c is constant for all existing interactions. Interactions are removed uniformly at random until the target connectance is reached, subject to the constraint that species cannot become isolated. This assumes that no ecological process preferentially favours

the realisation of one potential interaction over another, such that realised food webs constitute an unbiased subset of the regional metaweb that is ultimately only constrained by the number of links.

2.4 Simulations

For each empirical community, we generated 30 network replicates ($n = 30$ replicates $\times$ 7 network types $\times$ 6 locations). For each replicate a target connectance was independently sampled from a truncated normal distribution bounded between 0.05 and 0.15. This target connectance was used to determine the 'degree of downsampling' of the metaweb for each downsampling approach as well as constraining the ATN and Niche networks. These represent the 'creation' networks

For the dynamic simulations body masses were assigned using the predator-prey body mass distribution where body mass (M_i) is assigned deterministically from the trophic level (TL) as follows $M_i = Z^{TL_i - 1}$, where Z represents the assumed constant predator–prey body-mass ratio, here set to 10. Note that even for the ATN networks we used this theoretical body size specification despite the network itself being constructed with empirical bodysizes. This is because we found that for networks which had empirically specified body sizes the community would almost completely collapse (see Supplementary Materials), which suggests the dynamic model itself is sensitive to this specification. Initial biomasses were drawn from a uniform distribution. These trait values were used to construct a series of alternative networks.

Each creation network was then parameterised using the default parameters (Delmas et al., 2017) and ran the model until we had stable biomass fluctuation or up to 5000 time steps. Species whose biomass declined below the survival threshold (10^{-12}) during this burn-in period were considered extinct and removed from the network. The surviving species and their interactions constituted the corresponding 'realised' networks.

For each 'creation' and 'realised' network we measured 32 network structural metrics (described in Table S1). These span a different components of network structure from macro-scale to node-level properties.

2.4.1 Extinction simulations

For extinctions we followed the framework of Curtsdotter et al. (2011) and simulated both topological and dynamic extinctions. Topological extinctions were done on both the

'creation' as well as 'realised' networks. For each type of extinction simulation we used the same ten extinction scenarios from Curtsdotter et al. (2011) to determine the order of species to be removed for primary extinctions. For topological extinctions secondary extinctions occurred when species lost all prey, these secondary extinctions were allowed to cascade before the next primary extinction occurred. For dynamic extinctions primary extinctions were imposed by setting the biomass of the focal species to the survival threshold (10^{-12}), after which the community was simulated for a further 5000 time steps to allow for the downstream effects of the primary extinction to propagate through the system. Species whose biomass subsequently declined below the survival threshold were recorded as secondary extinctions and removed at the end of the 5000 time steps. This procedure was repeated until the entire community collapsed.

For each extinction scenario, robustness was quantified from the resulting extinction curves as the proportion of primary extinctions required to produce a 50% reduction in total species richness (R_{50} (Jonsson et al., 2015)). In instances when a primary extinction would 'overshoot' the 50% robustness was estimated by linear interpolation between extinction events.

2.5 Software

Network construction and extinction simulations were run in Julia v1.12.6 (Bezanson et al., 2017), we used `EcologicalNetworksDynamics.jl` (Lajaaiti et al., 2025) for dynamic simulations, `PFIM.jl` [REF] for construction of metawebs, `FoodWebTools.jl` [REF] for network downsampling, and `Extinctions.jl` [REF] for extinctions and robustness calculations. Statistical analyses were run in R v4.5.2 (R Core Team, 2024). All code can be found at *TODO*

3 Downsampling and burn-in alter network topology in method-dependent ways

Redundancy analysis (RDA), using permutation-based significance testing while accounting for community-level variation, revealed that downsampling approach significantly affected network topology, but that this effect depended on the magnitude of connectance reduction required to reach the target connectance (999 permutations; $F_{7,707} = 112.8$, $p = 0.001$; Figure 1, panel A). The type of network (*i.e.,* downsampling approach) explained 31.8%

of total topology variation (adjusted $R^2 = 0.318$), with community identity accounting for an additional 39.7% of variation. The significant interaction between downsampling approach and connectance change (marginal test; $F_{3,707} = 18.90$, $p = 0.001$) demonstrated that topology trajectories differed among downsampling methods depending on the degree of downsampling required. Thus, the extent to which network structure diverged from the original topology depended on both the severity of downsampling and the mechanism by which interactions were removed.

[Figure 1 about here.]

We next asked whether these structurally distinct networks responded differently during the burn-in phase (Figure 1, panel B). Structural reorganisation was quantified as the Euclidean displacement between network positions before and after burn-in in the first three principal component dimensions, with location included as a random effect to account for the substantial variation among location. Burn-in displacement differed significantly among network types ($F = 62.0$, $p < 0.001$). Metawebs underwent the greatest structural reorganisation, whereas ATN networks exhibited the smallest displacement. Among the downsampled networks, all but the randomly downsampled networks had a displacement similar to that of the Niche model (Figure S2), while all downsampled networks showed displacement distinct from the original metaweb. When looking at the loadings of the first two PC axes (Figure 1, panel C), the primary changes in structure were related to node-level measures and, unsurprisingly, the number of links. This is to be expected since there is no rewiring of links permitted in this system, and as such any changes should be driven by the loss of nodes rather than a major reorganisation of network structure.

Together, these analyses demonstrate that downsampling approach influences network topology both during the initial downsampling step and during the subsequent burn-in phase. Differences among downsampling methods become increasingly pronounced as greater proportions of interactions are removed, consistent with the expectation that the ecological assumptions underlying each removal strategy would become more embedded as more links are removed. Importantly, despite generating distinct initial topologies, the different downsampling approaches underwent remarkably similar structural reorganisation during burn-in, changing in ways more similar to the niche model than the original metaweb. This suggests that ecologically informed downsampling of metawebs allows us to produce networks whose subsequent structure more closely resembles that of realised networks, while also suggesting that the burn-in phase is, in and of itself, an additional network

realisation step. Notably, the two realised models (Niche and ATN) do not behave similarly and still end up at different 'end points' following burn-in compared with the empirical webs, highlighting that the choice of network reconstruction approach will still have a strong effect on the structure and behaviour of the system (Strydom, Karapunar, et al., 2026).

4 Differences in inferred robustness of topological and dynamic extinctions are unaffected by network type

To determine whether the qualitative differences between topological and dynamic extinction simulations previously demonstrated by Curtsdotter et al. (2011) generalise across alternative network reconstructions, we compared the difference in robustness (ΔR_{50}) among network types for each extinction scenario while accounting for community-level variation. Network type significantly affected ΔR_{50} across extinction scenarios (model summaries in Table S2), indicating that the magnitude of the difference between topological and dynamic robustness depended on network reconstruction.

Despite these quantitative differences, the qualitative pattern identified by Curtsdotter et al. (2011) was largely maintained. Under all but random basal removal, topological extinction inferred greater robustness than dynamic extinction for every network type, although the magnitude of this discrepancy varied among reconstruction methods (Figure 2). Random basal extinction represented the only notable departure from this general pattern (Figure S3). Here, the discrepancy between topological and dynamic robustness was substantially reduced, with power-law and degree-product downsampled networks exhibiting a slightly higher ΔR_{50} and random downsampled networks showing little difference between the two approaches. Thus, the conclusion that topological analyses provide more conservative estimates of robustness, although not universal, remains remarkably robust across reconstruction methods and extinction scenarios.

[Figure 2 about here.]

Together, these analyses demonstrate that the principal conclusion of Curtsdotter et al. (2011) extends well beyond the niche model. While the precise magnitude of the discrepancy between topological and dynamic robustness depends on both network reconstruction and extinction mechanism, the qualitative tendency for topological extinction simulations to infer greater robustness than dynamic simulations were recovered across most reconstructed

food webs examined here.

5 Dynamic extinctions re-enforce ecological assumptions

For each community × extinction scenario × time point, R_{50} was modelled as a function of network type, and estimated marginal means were compared using Tukey-adjusted pairwise contrasts (Figure S4-6). Network-type pairs with adjusted $p \geq 0.05$ were considered indistinguishable, and the frequency of indistinguishable comparisons was summarised across scenarios (Figure S7). To identify broader groupings of network types, estimated marginal means were standardised, Euclidean distances among network types were calculated, and hierarchical clustering was performed using Ward's D^2 method (Figure 3).

[Figure 3 about here.]

In terms of topological robustness, metaweb derived networks consistently formed a distinct cluster separate from the two realised networks (niche model and ATN; Figure 3, top two rows). However, under dynamic extinction simulations, this grouping broke down, with a reshuffling of the relative similarity among network types (Figure 3, bottom row). Most notably, the niche downsampled networks converged with those of the Niche model, whereas the remaining empirically derived networks maintained their mutual similarity.

Topological extinction simulations primarily interrogate network architecture and therefore retain the common structural signature inherited from the metaweb. Dynamic simulations introduce biomass- and rate-dependent interactions, revealing differences among networks that are not apparent from topology alone. When examining topological extinctions, it is clear why empirically derived networks behave similarly to one another: they share a common baseline architecture inherited from the regional metaweb (Figure 1, panel B), and topological cascades depend strictly on network topology. For dynamic extinctions, however, network behaviour is not determined solely by initial structure. Instead, higher-order differences emerge from how model parameterisation interacts with network topology. Thus, network structure appears sufficient to explain much of the similarity in responses to topological extinctions, but is insufficient to predict responses to dynamic extinctions, where functional parameterisation introduces additional sources of variation. Although the clustering pattern was not universal across communities (Figures S8-10), suggesting that initial network structure still exerts an influence.

This raises an important question about the role of network connectance – to what extent does specifying connectance constrain the structural and dynamical properties of generated networks? Work from Poisot & Gravel (2014) and Strydom, Dalla Riva, et al. (2021) shows that less-connected networks can exhibit greater structural complexity, because there are many ways to arrange a small number of links but progressively fewer ways to arrange links as a network becomes fully connected. At low connectance, we therefore expect a much larger number of possible network configurations, meaning that in specifying connectance we place constraints on the precise structure of the resulting network. This provides one possible explanation for why increasingly aggressive downsampling can amplify differences among reconstruction approaches: as fewer interactions are retained, the ecological assumptions used to determine which interactions remain have greater scope to shape the resulting network.

6 Conclusions

Reconstructing realised food webs from regional interaction pools requires assumptions about which potential interactions are likely to occur together. Here, we demonstrate that alternative downsampling approaches can produce dynamically viable food webs from the same empirical metaweb, but that the resulting networks retain detectable signatures of the assumptions underlying their construction. These differences become increasingly pronounced as greater proportions of interactions are removed, indicating that downsampling is not a neutral step between empirical data collection and dynamical modelling.

Our results further suggest that dynamical burn-in represents a second form of network realisation. Although burn-in altered the structure of all networks, it did not erase the differences introduced during initial network construction. Instead, the dynamical model acted as an additional filter, determining which structurally feasible networks could persist under its assumptions about biomass, interaction strengths, and species persistence. The ecological behaviour of a simulated community is therefore shaped by successive filters. First, how potential interactions are realised within a network, and second, how that network is filtered into a dynamically viable community.

Despite these differences in network structure and dynamical behaviour, the broad relationship between topological and dynamic robustness was consistent across network reconstruction approaches. Topological extinction simulations generally indicated greater

robustness than dynamic simulations, consistent with previous work (Curtsdotter et al., 2011). This suggests that some conclusions drawn from network topology may be relatively robust to uncertainty in network reconstruction (Strydom, Karapunar, et al., 2026). However, dynamic simulations also revealed differences among networks that were not apparent from topology alone, demonstrating that networks generated from the same empirical interaction pool can exhibit different dynamical behaviour depending on the assumptions used to realise that pool.

These findings highlight an important challenge for the growing use of dynamic food web models to move beyond describing network structure towards predicting ecological change. Increasingly sophisticated approaches can reconstruct networks from empirical observations or biological traits, while regional metawebs provide a means of retaining broad empirical information about potential interactions. Yet neither empirical provenance nor mechanistic parameterisation necessarily produces a realised community as both require assumptions about which interactions occur together and (in dynamic models) which of those interactions can persist. Network realisation therefore represents a necessary but currently underappreciated bridge between empirical interaction data and mechanistic dynamic modelling.

Developing a clear framework for explicitly incorporating empirical metawebs into dynamic models is an important step towards bridging theoretical work on network stability with real-world communities and ecological scenarios. Our case study reinforces the findings of Curtsdotter et al. (2011), suggesting that topological extinction simulations alone may underestimate the importance of top-down effects in determining secondary extinctions and community robustness. Intentional downsampling provides a means of approximating the structure of realised communities (Roopnarine, 2006), helping to integrate ecological realism with theoretical predictions. If palaeoecological communities can be reconstructed as plausible, dynamically viable food webs, dynamic modelling could provide a route to testing mechanistic explanations of observed ecological change rather than relying solely on topological reconstructions. Our results represent a first step towards this goal, while demonstrating that the assumptions underlying network construction can remain detectable throughout the modelling process.

Rather than identifying a single correct method for constructing realised networks, we argue that network realisation should be recognised as a methodological choice that can influence model outcomes. We therefore recommend that researchers who use downsampling explic-

itly test the sensitivity of their results to alternative downsampling approaches. As food web modelling increasingly combines empirical interaction data with dynamic approaches, predictions will depend not only on the equations governing population dynamics, but also on how empirical knowledge is translated into the networks on which those equations operate. Understanding and quantifying this additional source of uncertainty will be essential if dynamic food web models are to provide increasingly empirically grounded predictions of how ecological communities respond to environmental change.

References

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. (2017). Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, *59*(1), 65–98. https://doi.org/10.1137/141000671

Brose, U., Jonsson, T., Berlow, E. L., Warren, P., Banasek-Richter, C., Bersier, L.-F., Blanchard, J. L., Brey, T., Carpenter, S. R., Blandenier, M.-F. C., Cushing, L., Dawah, H. A., Dell, T., Edwards, F., Harper-Smith, S., Jacob, U., Ledger, M. E., Martinez, N. D., Memmott, J., … Cohen, J. E. (2006). Consumer–Resource Body-Size Relationships in Natural Food Webs. *Ecology*, *87*(10), 2411–2417. https://doi.org/10.1890/0012-9658(2006)87%5B2411:CBRINF%5D2.0.CO;2

Curtsdotter, A., Binzer, A., Brose, U., De Castro, F., Ebenman, B., Eklöf, A., Riede, J. O., Thierry, A., & Rall, B. C. (2011). Robustness to secondary extinctions: Comparing trait-based sequential deletions in static and dynamic food webs. *Basic and Applied Ecology*, *12*(7), 571–580. https://doi.org/10.1016/j.baae.2011.09.008

Delmas, E., Brose, U., Gravel, D., Stouffer, D. B., & Poisot, T. (2017). Simulations of biomass dynamics in community food webs. *Methods in Ecology and Evolution*, *8*(7), 881–886. https://doi.org/10.1111/2041-210X.12713

Dunhill, A. M., Zarzyczny, K., Shaw, J. O., Atkinson, J. W., Little, C. T. S., & Beckerman, A. P. (2024). Extinction cascades, community collapse, and recovery across a Mesozoic hyperthermal event. *Nature Communications*, *15*(1), 8599. https://doi.org/10.1038/s41467-024-53000-2

Dunne, J. (2006). The network structure of food webs. In *Ecological Networks: Linking Structure to Dynamics in Food Webs* (pp. 27–86).

Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning, J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal

food webs since the Late Pleistocene. *Science*, *377*(6609), 1008–1011. https://doi.org/10.1126/science.abn4012

Jonsson, T., Berg, S., Pimenov, A., Palmer, C., & Emmerson, M. (2015). The reliability of R50 as a measure of vulnerability of food webs to sequential species deletions. *Oikos*, *124*(4), 446–457. https://doi.org/10.1111/oik.01588

Karapunar, B., Strydom, T., Beckerman, A. P., Ridgwell, A., Wignall, P. B., Dunne, J. A., Little, C. T. S., Hull, P., Pimiento, C., & Dunhill, A. (2026, February 25). *No global collapse of food webs across the Permian-Triassic Mass Extinction*. https://doi.org/10.64898/2026.02.24.707709 (Pre-published)

Lajaaiti, I., Bonnici, I., Kéfi, S., Mayall, H., Danet, A., Beckerman, A. P., Malpas, T., & Delmas, E. (2025). EcologicalNetworksDynamics.jl: A Julia package to simulate the temporal dynamics of complex ecological networks. *Methods in Ecology and Evolution*, *16*(3), 520–529. https://doi.org/10.1111/2041-210X.14497

Petchey, O. L., Eklöf, A., Borrvall, C., & Ebenman, B. (2008). Trophically Unique Species Are Vulnerable to Cascading Extinction. *The American Naturalist*, *171*(5), 568–579. https://doi.org/10.1086/587068

Poisot, T., & Gravel, D. (2014). When is an ecological network complex? Connectance drives degree distribution and emerging network properties. *PeerJ*, *2*, e251. https://doi.org/10.7717/peerj.251

R Core Team. (2024). *R: A Language and Environment for Statistical Computing* [Computer software]. R Foundation for Statistical Computing. https://www.r-project.org/

Roopnarine, P. D. (2017). Ecological Modelling of Paleocommunity Food Webs. In *Conservation Paleobiology: Using the Past to Manage for the Future* (pp. 201–226). University of Chicago Press.

Roopnarine, P. D. (2006). Extinction Cascades and Catastrophe in Ancient Food Webs. *Paleobiology*, *32*(1), 1–19. https://www.jstor.org/stable/4096814

Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024, January 30). *A framework for reconstructing ancient food webs using functional trait data*. https://doi.org/10.1101/2024.01.30.578036 (Pre-published)

Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*, *13*(12). https:

//doi.org/10.1111/2041-210X.13835

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz, N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap towards predicting species interaction networks (across space and time). *Philosophical Transactions of the Royal Society B: Biological Sciences*, *376*(1837), 20210063. https://doi.org/10.1098/rstb.2021.0063

Strydom, T., Dalla Riva, G. V., & Poisot, T. (2021). SVD Entropy Reveals the High Complexity of Ecological Networks. *Frontiers in Ecology and Evolution*, *9*. https://doi.org/10.3389/fevo.2021.623141

Strydom, T., Dunhill, A. M., Dunne, J. A., Poisot, T., & Beckerman, A. P. (2026). *From Metawebs to Realised Webs: A Framework for Ecological Network Representation under Global Change*. https://ecoevorxiv.org/repository/view/11514/

Strydom, T., Karapunar, B., Dunne, J. A., Hull, P., Little, C. T. S., Pimiento, C., Ridgwell, A., Beckerman, A. P., & Dunhill, A. M. (2026). *Are We Mapping Ecosystems or Models? Framework Choices Dominate Food Web Topology and Extinction Inferences*. https://ecoevorxiv.org/repository/view/13495/

Ward, B. A., Wilson, J. D., Death, R. M., Monteiro, F. M., Yool, A., & Ridgwell, A. (2018). EcoGEnIE 1.0: Plankton ecology in the cGEnIE Earth system model. *Geoscientific Model Development*, *11*(10), 4241–4267. https://doi.org/10.5194/gmd-11-4241-2018

Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, *404*(6774), 180–183. https://doi.org/10.1038/35004572

Yodzis, P., & Innes, S. (1992). Body Size and Consumer-Resource Dynamics. *The American Naturalist*, *139*(6), 1151–1175. https://doi.org/10.1086/285380




Figure 1: **Downsampling approach influences both network topology and subsequent structural reorganisation during burn-in.** (A) Redundancy analysis (RDA) ordination showing differences in network topology among downsampling approaches across the downsamping (ΔCo) gradient after accounting for community-level variation. (B) Principal component trajectories illustrating structural change between network creation and burn-in for each downsampling approach. (C) Loadings of the top ten contributing structure metrics.




Figure 2: **Differences between topological and dynamic estimates of network robustness.** Boxplot showing aggregated realised topological and dynamic robustness across the different communities and extinction scenarios for the different network types.




Figure 3: **Clustering of network types based on similarity in inferred robustness (R_{50}).** Dendrograms show hierarchical clustering of network types based on their estimated marginal mean (R_{50}) values across community ($\times$) extinction scenario combinations. Clustering was performed separately for each extinction time point after standardising estimated marginal means across network types, using Euclidean distances and Ward's (D^2) method. Branch colours indicate three clusters of network types