

# Navigating food web prediction; assumptions, rationale, and methods

Tanya Strydom <sup>1</sup>; Jennifer A. Dunne <sup>2</sup>; Timothée Poisot <sup>3,4</sup>; Andrew P. Beckerman <sup>1</sup>

**Abstract:** TODO

**Keywords:** food web, network construction, scientific ignorance

1 At the heart of modern biodiversity science are a set of concepts about biodiversity, community structure,  
2 productivity, and asynchrony, and how they define the stability, resilience, and dynamics of complex com-  
3 munities. The use of species interaction networks provides a powerful abstraction that one can use to help  
4 quantify, conceptualise, and understand these concepts. However, network ecology has its own nuance and  
5 idiosyncrasies that not only provide a barrier to entry but causes dissonance even within the field (Dormann,  
6 2023). This is perhaps particularly pervasive within the space of network prediction...

7 One of the fundamental challenges that we are faced when working with and studying interaction networks  
8 (and, within the context of this manuscript, specifically food webs) is that there is a scarcity of ‘real world’  
9 interaction data (Hortal et al., 2015; Poisot et al., 2021). The difficulty of recording interactions in the field  
10 (Jordano, 2016a, 2016b) has necessitated that researchers find and develop alternative means to construct  
11 and build food webs using **models** (Morales-Castilla et al., 2015; Strydom et al., 2021). Over the past  
12 decade, there has been a proliferation of tools and processes for characterising food webs, these models  
13 span a wide range of philosophies that rely on different approaches, data, and definitions, which ultimately  
14 determine how the food web is constructed and coded. Although the development of these different models  
15 have carved out the path for constructing either synthetic, ecologically plausible networks (Poisot, Gravel,  
16 et al., 2016), or providing ‘first draft’ networks that can be utilised in real world settings (Strydom et al.,  
17 2022) we are still lacking in discussions that are explicitly comparing and contrasting how the way one  
18 chooses to approach the task of constructing a food web is introducing (and ultimately embedding) specific  
19 assumptions and hypotheses (Petchey et al., 2008). Most attempts that focus on comparing and contrasting  
20 models are focused on the same group of **model families** (Pichler et al., 2020; Williams & Martinez, 2008)  
21 and only benchmark the different models as opposed to contextualising them within the bigger framework  
22 of understanding the data needs of the different models, as well as how the resulting network is defined and  
23 structured. As food webs become a more integrated part of some of the broader fields of ecology (Bhatia et  
24 al., 2023; Thuiller et al., 2024) it is critical that we review these different model families as a whole (not only  
25 in isolation), and move away from simply benchmarking the performance of these different model families.  
26 This is important because different models impose different constraints upon themselves and will not only  
27 delimit and dictate the potential questions one will be able to ask (Petchey et al., 2011) but also determine  
28 the appropriate research setting for which the model (and resulting network) can be used. For example  
29 the use of ‘structural food webs’ are useful for developing additional theory such as re-wiring of networks  
30 (Staniczenko et al., 2010) but would be meaningless if one’s intention is to produce a location-specific network  
31 [do we need an *e.g.*, ref??]. This will allow us to ensure the right models are being used to answer the right  
32 questions, particularly within the context of trying to accelerate cross-cutting research in the face of global

change.

When navigating the seas of using and constructing food webs the researcher needs to be able to clearly articulate and define the parameters that are used to define their food web(s) of interest. This will aid them in being able to select the correct model to help them to reach their goal. In order to be able to make informed decisions it is important that one has a strong grasp of exactly what it means to ‘code’/define a food web (Section 1), a clear understanding of why one wants to predict a food web (Section 2), and ultimately one needs to be able to assess and evaluate which model family is going to best match up with the goal of network prediction (Section 3). Here we specifically aim to look at not only the performance of the different models but also initiate a (thus far lacking) discussion around how the interplay between the language used to define networks and the underlying theory/philosophy should also be a part of the broader discussion when it comes to the task of ‘model selection’.

[Figure 1 about here.]

## 1 The anatomy of a food web

Defining a food web seems simple, it is the representation of the interactions (edges) between species (nodes), however the definition of ‘edges’ and ‘nodes’, as well as the scale at which they are aggregated can take many forms. As highlighted in Poisot, Stouffer, et al. (2016) networks can be constructed at the population (the links among individuals), community (the links between species), or metacommunity (fluxes between locations) level. Even if one were to limit their scope to thinking of interaction networks only in terms of food webs at the community-level there are still many ways to define the various components of the network. Panel A of 1, one needs to understand the different intentions/assumptions that are made when a food web is constructed. Although the main intention of a food web is to capture and represent the feeding links between species there are many ways to define the nodes (*e.g.*, species or taxonomic group), edges (*e.g.*, **potential** or **realised feeding links**), the magnitude of the edges (*e.g.*, binary vs probabilistic), and even how the network itself is delimited (does it represent an aggregation of interactions over time?).

### 1.1 How do we define a node?

Although this may seem an elementary question in the context of food webs — a node *should* represent a (taxonomic) species, the reality is that nodes can often represent an aggregation of different species - so called ‘trophic species’ or segregation of species by life stages. Representing nodes as non-taxonomic species can be useful in certain contexts (Williams & Martinez, 2000) and in cases where the adult and larval stages of a

species have different diets it may make ecological sense (Clegg et al., 2018) meaning that it is not uncommon that networks often have nodes that have different definitions of a ‘species’ *e.g.* consisting of both taxonomic and trophic species. Practical implications of how we are aggregating the nodes is that the resolution may not always be ‘pixel perfect’ *i.e.*, we may be unable to assess the co-extinction risk of a species pair, however there is value in having nodes that represent an aggregation of species, as these convey a much more general overview of how the links are distributed within the community.

## 1.2 What is meant by an edge?

As discussed earlier there are many ways to define the links between species — even feeding links. At its core links within food webs can be thought of as a representation of either the flow of a resource [ref], realised (Pringle, 2020) or potential (Dunne, 2006) feeding links, or energy transfer and material flow (Lindeman, 1942). How we specify links will influence the resulting structure of the network - and the inferences we will make thereof. For example taking a food web that consists of links representing *potential* feeding links between species will be meaningless if you are interested in understanding *e.g.*, the flow of energy through the system as the links within the network are over overrepresented. In addition to the various ways of defining the links between species pairs there are also a myriad of ways in which the links themselves can be quantified. Links between species are often treated as being present or absent (*i.e.*, binary) but it is also possible to use probabilities (which quantifies how likely an interaction is to occur, Poisot, Cirtwill, et al., 2016) or continuous measurements (which quantifies the strength of of an interaction, Berlow et al., 2004). Moving away from a purely binary way of representing allows us to quantify a level of (un)certainty of our knowledge of interactions (*i.e.*, moving from being able to ask if are they occurring to quantifying how likely they are to occur) does add an additional level of ‘complexity’ to the construction and interpretation of networks, but ultimately it allows us to capture more information at different scales (Banville, in prep).

## 1.3 Putting the parts together; what does it mean?

The ingredients one uses to construct networks from nodes and edges generates a unique representation of the mechanisms (see Box 1 - Mechanisms that determine feeding links) that allow inference and reasoning about the structure, aspects of dynamics (*e.g.*, stability), and potentially the function of communities (*e.g.*, flux). It is thus beneficial to keep in mind that in the process of ‘codifying’ a network one is already embedding some sort of hypothesis as to the nature of the feeding links between species (Brimacombe et al., 2023; Proulx et al., 2005). Here it may be meaningful to contextualise the different ‘types’ of food webs within the larger research programmes (or even practical needs) that have been driving the construction of them.

## **i** Box 1 - Moving between the feasibility and reality of interactions

Before thinking about the ways in which we can predict networks it is perhaps meaningful to take a step back and think about the different criteria that must be met in order for an interaction to be able to occur between two species, specifically thinking of this in terms of distinguishing between the feasibility versus realisation of an interaction and how these are determined (and defined by) different ‘rules’/mechanisms. If we look at this feasibility-reality continuum (Figure 2) it is clear how the different predictive approaches (methods) tend to fall within one of the broader categories identified (distinguished) in the triangle. This is not to say that this shortcoming should be viewed as a ‘bug’ but rather a ‘feature’ of the field as it allows one to engage with, as well as construct networks at different scales, which is particularly valuable if one takes into consideration the considerable ‘data cost’ of predicting well resolved, realised networks in comparison to constructing high-level metawebs. However, it is important that there is an awareness and acknowledgement of where within this feasibility-reality one is working at and how this will impact and limit the contexts in which the resulting network can be used and applied within.

[Figure 2 about here.]

Aside: this is maybe not the IRL ‘assembly mechanisms’ per se (e.g. species must co-occur, then they must meet, then we can determine if the interaction will happen) but it makes modelling sense? IDK... anyway we can shuffle things around

## **‘Mechanisms’**

### **1. Feasibility interactions**

This is based on the idea of forbidden links introduced by Jordano (2016b), specifically that there must be some degree of *trait complementarity* that allows a predator to chase, capture, kill, and consume, its prey. This is probably the level that the idea of a metaweb (Dunne, 2006) is most applicable to. Within the network prediction ‘field’ this is perhaps the most developed space. Predictive models run the gamut including mechanistic models (Morales-Castilla et al., 2015), binary classifiers (Pichler et al., 2020), and graph embedding (Strydom et al., 2023) and use either traits (or phylogeny as a proxy for the conservation thereof) as a means to ‘evaluate’ if an interaction is *possible* between two species (again not the likelihood of it happening but the likelihood of its feasibility). It is probably worth having a brief interlude here to be really clear that just because an interaction is probabilistic it does not make it weighted (at least not in the traditional sense of weighted networks, *e.g.*, J. T. Wootton &

Emmerson (2005)) - it is still ‘binary’, it just happens to be defined by a binomial distribution (*sensu* Banville, in prep).

## **2. (Co)occurrence interactions**

Although the assumption that because two species are co-occurring it must mean that they are interacting is presumptive (Blanchet et al., 2020) it is of course impossible for two species to interact (at least in terms of feeding links) if they are not co-occurring in time and space. An example of this would be Dansereau et al. (2023) where the metaweb is downsampled into smaller realisations based on better data/knowledge as to which species are occurring at a specific location - however arguably these are still firmly in the space of feasible interactions for the specific location but are approaching a better approximation of ‘reality’...

## **3. Mass effect interactions**

Not sure if there are models that ‘only’ consider abundance (barring the neutral model) and that it is rather more of a building block in some of the models that are more relevant to the next steps. Maybe there is an argument that this ‘rule’ is ‘irrelevant’ in the context of network prediction and more so a data parameter one needs...

This is probably the point where we start to shift from a *potential* (presence/absence) way of defining interactions and start moving into the ‘qualitative’/weighted interaction space - we are not ‘determining’ if the interaction is feasible but rather making an assumption on prey selection based on the species likelihood of ‘meeting’, although Banville (in prep) presents a compelling case that this could still be considered something that falls under the ‘feasibility’ and not ‘reality’ side of the spectrum... (maybe, past Tanya seemed to think so)

## **4. Consumer–resource interactions**

This is where we begin to move into the foraging ecology space - specifically consumption rate and how that pertains to energy acquisition. In the loosest sense I think this is the ‘prey choice’ space - but specifically in the context of how prey choice is informed by energetic cost (not just purely based on *e.g.*, the most abundant species). If we think about ways that people have approached this there are the diet models of (Beckerman et al., 2006) and (Petchey et al., 2008) that take a high-level (imo) approach to accounting for energy acquisition, then you get the ‘trait’ framework developed by K. L. Wootton et al. (2023) that moves the ‘energy’ into different ‘modules’ related to the process of the consumer acquiring energy from the resource (however there is a disregard for the ‘Rule 1’ requirement of forbidden links, again not bad just pointing it out). The idea of the consumer search space developed by Pawar et al. (2012) is also an interesting consideration.

## 5. Consumer–environment interactions

I think this should be it's own rule since its really more about the idea of how the environment is imposing energy costs on the predator as opposed the energetic costs (and gain) of consuming the prey. Basically the ideas presented in Cherif et al. (2024), which is essentially a take on movement ecology? What it boils down to is being able to quantify the cost of movement *i.e.*, the physical constraints that the environment imposes on a species... Maybe we can also think of it more in terms of metabolic rate?

The role of the environment: Although we only explicitly take the environment into consideration in the final 'rule' the role of the environment (at least from a modelling perspective) is perhaps also meaningful at the first step (we expect environmental filtering to be the main determinant of species distributions in the absence of species interactions) I think the way in which we perceive the environment at the extreme of these two ends of the spectrum comes down to a question of scale?? Or at least the degree of information that we need at need at these two points is very different. RE occurrence it is a case of survival and for movement it is a case of performance??

### Notes on the 'ensemble' approach

IDK I just really want to articulate that we can't feasibly be functioning at a 'Rule 5' level at all times because of other real world constraints such as the fact that the amount of data one needs is simply not going to be feasible in most cases... Also that you (probably) need to go through the whole step-wise process because (at least for now) models at the 'higher' levels are overlooking some of the 'lower' levels *e.g.*, the the ADBM model (in its current form) has no way of accounting for forbidden links and so it is probably going to end up predicting some links that are strictly speaking 'forbidden' at the trait level.

94

## 2 Why do we want to predict food webs?

95

96 As discussed in Section 1 there are many ways to define a food web, meaning that there are equally as many  
97 reasons one might be interested in predicting a food web. However we may think of two primary drivers  
98 for wanting to predict networks (Panel B Figure 1), namely an interest in generating a set of ecologically  
99 plausible networks (*i.e.*, being able to describe networks using a model) or being able to recover (predict)  
100 location specific, 'realised', interactions for a specific species community (*i.e.*, being able to predict/infer the  
101 interactions between species). Of course these two categories are not distinct, mutually exclusive, groups but

can rather be viewed as operating on a continuum ranging from a need for generality (*i.e.*, creating a network that, when taken in aggregate, the distribution of links (interactions) between nodes (species) are ecologically plausible) to a need for specificity (*i.e.*, local-level predictions between specific species pairs). Although the ability to predict ‘real-world’ interactions (and the resulting food webs) can have more intuitive ‘real world’ applications *e.g.*, being able to ‘recover’ food webs that have since gone extinct (Dunne et al., 2008; Yeakel et al., 2014), using pairwise interactions to understand species distributions (Pollock et al., 2014) or even co-extinction risk (Dunn et al., 2009), a more structural approach to network construction affords one an opportunity to interrogate some of the more high-level mechanisms that are structuring networks (Box 1). It is perhaps more important that when one is talking about ‘why’ they want to predict networks to articulate exactly what anatomical part of the food web we are interested in scrutinising.

### 3 How do we predict food webs?

Selecting a model for the task of network prediction should come down to two things; what *aspect* of a food web one is interested in predicting, and what data are available, necessary, and sufficient. As shown in panel B of Figure 1 the interest in a network is (usually) at either the ‘structural’ or ‘interaction’ level and the development of models for the task of network prediction often focus on high fidelity (performance) at one of these scales. With this in mind it is beneficial to think of the different model families relative to these two different goals; here we refer to models that are used to predict the structure of a network as **topology generators** and models developed to infer the interactions for a given species pool as **interaction predictors**. It is meaningful to make this distinction because although it is possible to construct a food web given using an *interaction predictor* the models themselves lack any sort of parametrisation of the network structure and so the resulting network is a poor reflection of the actual network structure (Caron et al., 2024). This is primarily because *interaction predictors* are models that evaluate the feasibility of an interaction between species pairs and not in the context of feasibility at the community level. Models themselves are a reflection of the different goals and intentions of the research program from which they are developed and are often ‘described’ by a specific mechanism that will determine the resulting structure or interactions (Box 1). Models such as the niche (Williams & Martinez, 2000) or cascade (Cohen et al., 1990) were developed with the intent of being used to understand the *structural* aspects of food webs, specifically how links are distributed amongst species in the community, whereas bayesian (Cirtwill et al., 2019) or trait hierarchy (Shaw et al., 2024) models have been developed on the basis that the traits of a species are the underlying mechanism in determining the feasibility of interactions (*i.e.*, species *a* has the capacity to eat species *b*). Along with predicting different anatomical parts of a food web the different models have varying



degrees of data that are needed to ‘parametrise’ the network. Once these two limitations are assessed and addressed it is then possible to select the model (or model family) that will best be able to capture food web feature that the researcher is most interested in (see Box 2 - Assessing model outputs). It is thus clear that (realistically) there will probably never be a ‘best fit’ tool that is able to construct a food web that will span the entire range of needs, and rather the responsibility lies with the researcher to be aware of not only the underlying philosophy of the specific toolset (as this could have knock-on effects when using those networks for downstream analyses/simulations; pers. comms. Beckerman, 2024), but also how well the tool is able to retrieve the specific network or interaction properties that is of interest.

In order for a model to formalise a ‘complete’ food web it is necessary to formalise two aspects of the network, ‘who eats whom’ (to determine the links between nodes) as well as the structure of the network (to limit the distribution of links), however most models are inclined to focus on one of the two aspects panel B of 1.

Crucially most topology generators lack some key data on the interaction between species (this can be because of how the model itself defines species or the way in which links are assigned in the network) and interaction predictors lack some sort of parametrisation of network structure (just because two species can interact it does not mean that they will, Poisot et al., 2015).

What is the purpose of generating a network? Is it an element of a bigger question we are asking, *e.g.*, I want to generate a series of networks to do some extinction simulations/bioenergetic stuff OR are we looking for a ‘final product’ network that is relevant to a specific location? (this can still be broad in geographic scope).

### 3.1 Model families

As there are many food web models to choose from it is perhaps useful to think about the models in terms of model families, a summary of these families is presented in Table 1 and along with Figure 3 highlights the differences and similarities of the philosophies and assumptions that determine a network. A more extensive overview of the different models that fall within the different model families can be found in SuppMat 1 and for a more detailed breakdown of the different ‘traits’ of the model families refer to SuppMat 2.

Table 1: A summary of the different families of tools that can be used to generate food webs, this includes a brief description of the underlying philosophy of the family as well as how the different elements (nodes and edges) of the generated network represents.

Model		Network	Nodes	Links		Key
family	Theory	predicted	represent	represent	Interaction	reference
null	Links are randomly distributed within a network	structural	agnostic	feeding links	binary	
neutral	Network structure is random, but species abundance determines links between nodes	structural	species	feeding links	binary	
resource	Networks are interval, species can be ordered on a ‘niche axis’	structural	trophic species	subdivision of resource	binary	Williams & Martinez (2008)
generative	Networks are determined by their structural features	structural	agnostic	links	binary	

Model family	Theory	Network predicted	Nodes represent	Links represent	Interaction	Key reference
energetic	Interactions are determined by foraging theory (feeding links)	interaction	species	feeding links	quantitative	
graph embedding	Interactions can be predicted from the latent traits of networks	interaction	species	potential feeding links	probabilistic	Strydom et al. (2023)
trait matching	Interactions can be inferred by a mechanistic framework/relationships	interaction	species	feeding links	binary	Morales-Castilla et al. (2015)
binary classifiers	Interactions can be predicted by learning the relationship between interactions and ecologically relevant predictors	interaction	species	feeding links	binary	Pichler et al. (2020)

Model family	Theory	Network predicted	Nodes represent	Links represent	Interaction	Key reference
expert knowledge	‘Boots on the ground’ ecological knowledge and observations	interaction	species	feeding links	binary	
data scavenging	Webscraping to create networks from online databases	interaction	species	feeding links	binary	Poisot, Gravel, et al. (2016) (f you squint?)
co-occurrence	co-occurrence patterns arise from interactions so we can use these patterns to reverse engineer the interactions	co-occurrence patterns	species	association links	binary	Kusch et al. (2023) (although more plant-plant <i>i.e.</i> non-trophic...)

[Figure 3 about here.]

### **i** Box 2 - Assessing model outputs

Although understanding the underlying philosophy of the different model families is beneficial it is also important to understand in what situations the different families are likely to perform well or poorly. When we are assessing the performance of the different model families it is beneficial to think of benchmarking these assessments based on a broader basis than just its ability to correctly recover

network structure or pairwise interactions. When thinking about how to benchmark models it is perhaps beneficial to take a step back and once again assess what are the needs of the researcher (Section 2) and linking this back to what aspects of the network (Section 1) are of importance and assess the performance of a model within those parameters.

### **Benchmarking**

Benchmarking how well a model is doing to capture the desired elements of a network is also a task that required some thought and contemplation. Even if we think about the predicting the structure of a network it is possible that two networks may have the same number of nodes and links but that those links may be distributed in very different ways. Thus it is important to think critically about the suite of summary statistics that are used to assess a model, since there is no one ‘silver bullet’ summary statistic that will be able to assess if a model is able to fully replicate an empirical network (Allesina et al., 2008). One of the main challenges when assessing the ability to retrieve pairwise interactions is that food webs are sparse (that means that there are few links given the number of species) and it is important that we are able to discern between a model that is able to correctly predict interactions that do (true positives) and not (true negatives) occur and one that is simply predicting a lack of interactions (Poisot, 2023). For more detailed methods as to how benchmarking was done refer to [SuppMat 3](#)

[Figure 4 about here.]

### **Data cost**

This includes thinking about the need for additional data sources (such as trait or phylogenetic data), the computational cost, as well as the time it might take to generate a network, *e.g.*, binary classifiers require an (often times) extensive list of additional trait data for the model training process, which limits predictions to communities for which you do have the relevant auxiliary data available.

### **Philosophical constraints**

Probably mentioned elsewhere but basically are we constructing networks because we want to make real-world, case-specific predictions *e.g.*, for a conservation area or do we want to just have a set of ecologically plausible networks we can use for theoretical stuffs. Need to discuss the key differences and implications between predicting a **metaweb** (*sensu* Dunne (2006)) and a network realisation. (In a way the idea of predicting a metaweb vs realisation is what makes me hesitant to use the Mangal networks to test the structural models because do we even know what the Mangal networks represent and what the structural models are predicting...) Maybe also Poisot et al. (2015) that discuss how the local factors are going to play a role.

Also need to take into consideration inherent constraints that the model imposes on itself and how it will

affect our ability to test hypotheses/ask questions using the *e.g.*, from Petchey et al. (2011) - models that are constrained by connectance means that we are unable to explain connectance itself and you would need a different approach if understanding connectance is your goal. Another way of phrasing this is thinking about what is needed (input data/parameters), produced (final network characteristics), and desired (end-use).

An interesting thing to also think about is data dependant and data independent ‘parametrisation’ of the models...

## 4 Concluding remarks

- As discussion about the different model families and in what areas they do/do not do well. This will depend probably a fair bit on how Figure 4 end up looking... But it will also be important to tie in some of the other considerations/constraints that are listed in what is currently Box 2
  - In certain situations structure is ‘enough’ but there may be use cases where we are really interested in the node-level interactions *i.e.*, species identity is a thing we care about and need to be able to retrieve specific interactions at specific nodes correctly.
- Why do interaction models do so badly at predicting structure? Nuance of metaweb vs realisation but also time? At the core of it interaction models are trained on existing interaction data; this is data that are most likely closer to a metaweb than a local realisation even if they are being inventoried at a small scale...
  - We can briefly shoehorn downsampling here maybe??
- It will be interesting to bring up the idea that if a model is missing a specific pairwise link but doing well overall then when does it matter?
  - The fact that *some* people are concerned about the taxonomic resolution and cascading effects those might have on our understanding of network structure (Pringle, 2020; Pringle & Hutchinson, 2020), but that puts us in a place where we are at risk of losing our ability to distinguish the wood from the tree - are we not (at least at times) concerned more with understanding ecosystem level processes than with needing to understand things *perfectly* at the species level.
  - I don’t think these ‘rare’/nuanced links (e.g. carnivorous hippos) are going to rock the boat when we think about networks at the structural level.

“The resolution of food-web data is demonic because it can radically change network topology and associated biological inferences in ways that are unknowable in the absence of better data.” - Pringle & Hutchinson (2020) The counter to this is that structural models are often not working at the species level and thus the structure remains ‘unchanged’ when you increase the resolution - I don’t think that people are that concerned with the structure of real world networks barring connectance and since that scales with species richness anyway your final proportion will probably still remain the same...

- I think a big take home will (hopefully) be how different approaches do better in different situations and so you as an end user need to take this into consideration and pick accordingly. I think Petchey et al. (2011) might have (and share) some thoughts on this. I feel like I need to look at Berlow et al. (2008) but maybe not exactly in this context but vaguely adjacent.

- I think this is sort of the crux of the argument presented in Brimacombe et al. (2024) as well.

*“we highlight an interesting paradox: the models with the best performance measures are not necessarily the models with the closest reconstructed network structure.”* - Poisot (2023)

- Do we need network models to predict interactions and interaction models to predict structure?

- “Another argument for the joint prediction of networks and interactions is to reduce circularity and biases in the predictions. As an example, models like linear filtering generate probabilities of non-observed interactions existing, but do so based on measured network properties.” - Strydom et al. (2021)

- Aligning (dove-tailing) with this the idea of ensemble modelling as presented by Becker et al. (2022)

- Close out with a call to action that we have models that predict networks very well and models that predict interactions very well but nothing that is doing well at predicting both - this is where we should be focusing our attention when it comes to furthering model development...

- Do we expect there to be differences when thinking about unipartite vs bipartite networks? Is there underlying ecology/theory that would assume that different mechanisms (and thus models) are relevant in these two ‘systems’.

- The Terry & Lewis (2020) paper looks at some methods but is specifically looking at a bipartite world...

## 4.1 Downsampling

do we bring this up? this could be a box... if we have the ‘finances’ for it... otherwise it should go to the outstanding questions for sure

- Dansereau et al. (2023)
- “That being said, there is a compelling argument for the need to ‘combine’ these smaller functional units with larger spatial networks (Fortin et al., 2021) and that we should also start thinking about the interplay of time and space (Estay et al., 2023). Although deciding exactly what measure might actually be driving differences between local networks and the regional metaweb might not be that simple (Saravia et al., 2022).”

## 4.2 Time

We lack a clear agenda (and conceptualisation) as to what the appropriate level of aggregation is for a ‘network’. Realistically most empirical networks are more aligned with ‘feasibility networks’ as opposed to ‘realised networks’ as they are often the result of some sort of aggregation of observations across time. This ‘problem’ is two-fold. Firstly we need to think about how this affects any sort of development of theory that sits closer to the ‘realised network’ side of the spectrum - how often are we trying to ask and answer questions about realised networks using feasible networks? The second is that this lack of ‘direction’ as to how we should define a network is (actually) probably one of the biggest barriers that is affecting the use of networks in applied settings...

Another time perspective question is when do we determine a link to be ‘real’... In the context of feasible networks this is perhaps clearer - all things equal would the predator be able to consume the prey. However in the realised space there is also the question of the long term ‘energetic feasibility’ of an interaction - just because an interaction is possible in the now is it able to sustain a population in the long term. And what is the scale for that long term - are we thinking at the generational scale? Because ultimately when we are constructing a network we are aggregating not only across space but also across time.

## Glossary

Term	Definition
food web	a representation of feeding links between species



Term	Definition
topology generator	a model that predicts a network based on assumptions of structure, this network is species agnostic in the sense that it does not necessarily contain information at the node level
interaction predictor	a model that predicts species interactions, these interactions can be used to construct a network but there are no <i>a priori</i> assumptions as that will constrain the network structure
model	A tool that can be used to construct food webs, where the resulting network is a representation of a real world network. Models typically only capture specific elements of real world networks and are intended to be used in specific settings
model family	A family of models that share an underlying philosophy when it comes to the mapping, pragmatism, and reduction of a network. Families have the same underlying philosophies and assumptions that determine the links between nodes as well as how these may be encoded
metaweb	A network that represents <i>all</i> the potential links between species. Importantly these links will not necessarily all be realised in a specific location for a specific time
realised network	A network that represents the links between species that are occurring. These networks represent a very localised network...
potential feeding link	links that indicate that an interaction is ecologically feasible but not realised <i>per se</i> (a metaweb would contain potential feeding links)

Term	Definition
realised feeding link	links that indicate that the interaction is realised ‘in the field’. (a realised network contains realised feeding links)
confusion matrix	captures the number of true positives (interaction predicted as present when it is present), false negatives (interaction predicted as absent when it is present), false positives (interaction predicted as present when it is absent), and true negatives (interaction predicted as absent when it is absent)

## Outstanding questions

- non-consumptive effects
- can we develop a model that is both a topology generator as well as an interaction predictor?
- how do we define the spatial and temporal ‘boundaries’ of a network

## References

- Allesina, S., Alonso, D., & Pascual, M. (2008). A General Model for Food Web Structure. *Science*, 320(5876), 658–661. <https://doi.org/10.1126/science.1156269>
- Becker, D. J., Albery, G. F., Sjodin, A. R., Poisot, T., Bergner, L. M., Chen, B., Cohen, L. E., Dallas, T. A., Eskew, E. A., Fagre, A. C., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Stock, M., Teeling, E. C., & Carlson, C. J. (2022). Optimising predictive models to prioritise viral discovery in zoonotic reservoirs. *The Lancet Microbe*, 3(8), e625–e637. [https://doi.org/10.1016/S2666-5247\(21\)00245-7](https://doi.org/10.1016/S2666-5247(21)00245-7)
- Beckerman, A. P., Petchey, O. L., & Warren, P. H. (2006). Foraging biology predicts food web complexity. *Proceedings of the National Academy of Sciences*, 103(37), 13745–13749. <https://doi.org/10.1073/pnas.0603039103>
- Berlow, E. L., Brose, U., & Martinez, N. D. (2008). The “Goldilocks factor” in food webs. *Proceedings of the National Academy of Sciences*, 105(11), 4079–4080. <https://doi.org/10.1073/pnas.0800967105>
- Berlow, E. L., Neutel, A.-M., Cohen, J. E., de Ruiter, P. C., Ebenman, B., Emmerson, M., Fox, J. W., Jansen, V. A. A., Iwan Jones, J., Kokkoris, G. D., Logofet, D. O., McKane, A. J., Montoya, J. M., & Petchey, O.

- (2004). Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73(3), 585–598. <https://doi.org/10.1111/j.0021-8790.2004.00833.x>
- Bhatia, U., Dubey, S., Gouhier, T. C., & Ganguly, A. R. (2023). Network-based restoration strategies maximize ecosystem recovery. *Communications Biology*, 6(1), 1–10. <https://doi.org/10.1038/s42003-023-05622-3>
- Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23(7), 1050–1063. <https://doi.org/10.1111/ele.13525>
- Brimacombe, C., Bodner, K., & Fortin, M.-J. (2024). *Applying a method before its proof-of-concept: A cautionary tale using inferred food webs*. <https://doi.org/10.13140/RG.2.2.22076.65927>
- Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4), e3002068. <https://doi.org/10.1371/journal.pbio.3002068>
- Caron, D., Brose, U., Lurgi, M., Blanchet, F. G., Gravel, D., & Pollock, L. J. (2024). Trait-matching models predict pairwise interactions across regions, not food web properties. *Global Ecology and Biogeography*, 33(4), e13807. <https://doi.org/10.1111/geb.13807>
- Cherif, M., Brose, U., Hirt, M. R., Ryser, R., Silve, V., Albert, G., Arnott, R., Berti, E., Cirtwill, A., Dyer, A., Gauzens, B., Gupta, A., Ho, H.-C., Portalier, S. M. J., Wain, D., & Wootton, K. (2024). The environment to the rescue: Can physics help predict predator–prey interactions? *Biological Reviews*, n/a(n/a). <https://doi.org/10.1111/brv.13105>
- Cirtwill, A. R., Eklf, A., Roslin, T., Wootton, K., & Gravel, D. (2019). A quantitative framework for investigating the reliability of empirical network construction. *Methods in Ecology and Evolution*, 10(6), 902–911. <https://doi.org/10.1111/2041-210X.13180>
- Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on food web structure. *Ecology*, 99(12), 2712–2720. <https://doi.org/10.1002/ecy.2523>
- Cohen, J. E., Briand, F., & Newman, C. (1990). *Community Food Webs: Data and Theory*. Springer-Verlag.
- Dansereau, G., Barros, C., & Poisot, T. (2023). *Spatially explicit predictions of food web structure from regional level data*.
- Dormann, C. F. (2023). The rise, and possible fall, of network ecology. In *Defining Agroecology – A Festschrift for Teja Tscharnke* (pp. 143–159.). Tredition.
- Dunn, R. R., Harris, N. C., Colwell, R. K., Koh, L. P., & Sodhi, N. S. (2009). The sixth mass coextinction: Are most endangered species parasites and mutualists? *Proceedings. Biological Sciences*, 276(1670), 3037–3045. <https://doi.org/10.1098/rspb.2009.0413>
- Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological*

networks: *Linking structure and dynamics* (pp. 27–86). Oxford University Press.

Dunne, J. A., Williams, R. J., Martinez, N. D., Wood, R. A., & Erwin, D. H. (2008). Compilation and Network Analyses of Cambrian Food Webs. *PLOS Biology*, 6(4), e102. <https://doi.org/10.1371/journal.pbio.0060102>

Estay, S. A., Fortin, M.-J., & López, D. N. (2023). Editorial: Patterns and processes in ecological networks over space. *Frontiers in Ecology and Evolution*, 11.

Fortin, M.-J., Dale, M. R. T., & Brimacombe, C. (2021). Network ecology in dynamic landscapes. *Proceedings of the Royal Society B: Biological Sciences*, 288(1949), rspb.2020.1889, 20201889. <https://doi.org/10.1098/rspb.2020.1889>

Hortal, J., de Bello, F., Diniz-Filho, J. A. F., Lewinsohn, T. M., Lobo, J. M., & Ladle, R. J. (2015). Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 523–549. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>

Jordano, P. (2016a). Chasing Ecological Interactions. *PLOS Biology*, 14(9), e1002559. <https://doi.org/10.1371/journal.pbio.1002559>

Jordano, P. (2016b). Sampling networks of ecological interactions. *Functional Ecology*. <https://doi.org/10.1111/1365-2435.12763>

Kusch, E., Bimler, M., Lutz, J. A., & Ordonez, A. (2023). *Ecological network inference is not consistent across scales or approaches* (p. 2023.07.13.548816). bioRxiv. <https://doi.org/10.1101/2023.07.13.548816>

Lindeman, R. L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23(4), 399–417. <https://doi.org/10.2307/1930126>

Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from proxies. *Trends in Ecology & Evolution*, 30(6), 347–356. <https://doi.org/10.1016/j.tree.2015.03.014>

Pawar, S., Dell, A. I., & Savage, V. M. (2012). Dimensionality of consumer search space drives trophic interaction strengths. *Nature*, 486(7404), 485–489. <https://doi.org/10.1038/nature11131>

Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2008). Size, foraging, and food web structure. *Proceedings of the National Academy of Sciences*, 105(11), 4191–4196. <https://doi.org/10.1073/pnas.0710672105>

Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2011). Fit, efficiency, and biology: Some thoughts on judging food web models. *Journal of Theoretical Biology*, 279(1), 169–171. <https://doi.org/10.1016/j.jtbi.2011.03.019>

Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine learning algorithms to infer trait-matching and predict species interactions in ecological networks. *Methods in Ecology and Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>

- Poisot, T. (2023). Guidelines for the prediction of species interactions through binary classification. *Methods in Ecology and Evolution*, 14(5), 1333–1345. <https://doi.org/10.1111/2041-210X.14071>
- Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., Mercier, B., Violet, C., & Vissault, S. (2021). Global knowledge gaps in species interaction networks data. *Journal of Biogeography*, 48(7), 1552–1563. <https://doi.org/10.1111/jbi.14127>
- Poisot, T., Cirtwill, A., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. (2016). The structure of probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312. <https://doi.org/10.1111/2041-210X.12180>
- Poisot, T., Gravel, D., Leroux, S., Wood, S. A., Fortin, M.-J., Baiser, B., Cirtwill, A. R., Araújo, M. B., & Stouffer, D. B. (2016). Synthetic datasets and community tools for the rapid testing of ecological hypotheses. *Ecography*, 39(4), 402–408. <https://doi.org/10.1111/ecog.01941>
- Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>
- Poisot, T., Stouffer, D. B., & Kéfi, S. (2016). Describe, understand and predict: Why do we need networks in ecology? *Functional Ecology*, 30(12), 1878–1882. <https://www.jstor.org/stable/48582345>
- Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., Veski, P. A., & McCarthy, M. A. (2014). Understanding co-occurrence by modelling species simultaneously with a Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution*, 5(5), 397–406. <https://doi.org/10.1111/2041-210X.12180>
- Pringle, R. M. (2020). Untangling Food Webs. In *Unsolved Problems in Ecology* (pp. 225–238). Princeton University Press. <https://doi.org/10.1515/9780691195322-020>
- Pringle, R. M., & Hutchinson, M. C. (2020). Resolving Food-Web Structure. *Annual Review of Ecology, Evolution and Systematics*, 51(Volume 51, 2020), 55–80. <https://doi.org/10.1146/annurev-ecolsys-110218-024908>
- Proulx, S. R., Promislow, D. E. L., & Phillips, P. C. (2005). Network thinking in ecology and evolution. *Trends in Ecology & Evolution*, 20(6), 345–353. <https://doi.org/10.1016/j.tree.2005.04.004>
- Saravia, L. A., Marina, T. I., Kristensen, N. P., De Troch, M., & Momo, F. R. (2022). Ecological network assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, 91(3), 630–642. <https://doi.org/10.1111/1365-2656.13652>
- Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024). A framework for reconstructing ancient food webs using functional trait data (p. 2024.01.30.578036). bioRxiv. <https://doi.org/10.1101/2024.01.30.578036>
- Staniczenko, P. P. A., Lewis, O. T., Jones, N. S., & Reed-Tsochas, F. (2010). Structural dynamics and robustness of food webs. *Ecology Letters*, 13(7), 891–899. <https://doi.org/10.1111/j.1461-0248.2010.024908.x>

- Stouffer, D. B., Camacho, J., Jiang, W., & Nunes Amaral, L. A. (2007). Evidence for the existence of a robust pattern of prey selection in food webs. *Proceedings of the Royal Society B: Biological Sciences*, 274(1621), 1931–1940. <https://doi.org/10.1098/rspb.2007.0571>
- Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*, 13(12), 2838–2849. <https://doi.org/10.1111/2041-210X.13835>
- Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2023). Graph embedding and transfer learning can help predict potential species interaction networks despite data limitations. *Methods in Ecology and Evolution*, 14(12), 2917–2930. <https://doi.org/10.1111/2041-210X.14228>
- Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz, N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap towards predicting species interaction networks (across space and time). *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>
- Terry, J. C. D., & Lewis, O. T. (2020). Finding missing links in interaction networks. *Ecology*, 101(7), e03047. <https://doi.org/10.1002/ecy.3047>
- Thuiller, W., Calderón-Sanou, I., Chalmandrier, L., Gaüzère, P., O'Connor, L. M. J., Ohlmann, M., Poggiato, G., & Münkemüller, T. (2024). Navigating the integration of biotic interactions in biogeography. *Journal of Biogeography*, 51(4), 550–559. <https://doi.org/10.1111/jbi.14734>
- Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, 404(6774), 180–183. <https://doi.org/10.1038/35004572>
- Williams, R. J., & Martinez, N. D. (2008). Success and its limits among structural models of complex food webs. *Journal of Animal Ecology*, 77(3), 512–519. <https://doi.org/10.1111/j.1365-2656.2008.01362.x>
- Wootton, J. T., & Emmerson, M. (2005). Measurement of Interaction Strength in Nature. *Annual Review of Ecology, Evolution, and Systematics*, 36(1), 419–444. <https://doi.org/10.1146/annurev.ecolsys.36.091704.175535>
- Wootton, K. L., Curtsdotter, A., Roslin, T., Bommarco, R., & Jonsson, T. (2023). Towards a modular theory of trophic interactions. *Functional Ecology*, 37(1), 26–43. <https://doi.org/10.1111/1365-2435.13954>
- Yeakel, J. D., Pires, M. M., Rudolf, L., Dominy, N. J., Koch, P. L., Guimarães, P. R., & Gross, T. (2014). Collapse of an ecological network in Ancient Egypt. *PNAS*, 111(40), 14472–14477. <https://doi.org/10.1073/pnas.1408471111>




Figure 1: Panel **A** shows the many ways in which a food web can be defined and described at the node, edge, and even network level. Panel **B** (will) shows how the way in which we predict networks also limited and often focuses only on predicting the structure of a network (the final network is parametrised by the expected structure of the network) or the interactions between species (the final network is determined by the behaviour of the nodes). These different models also encode different philosophies/hypotheses not only as to what determines how a network will look like but also how the final network itself is encoded *i.e.*, its anatomy. (*aside*: there is the potential to either try and visually summarise how the different model families define a network (so repeating the motifs used in the ANATOMY panel) alternatively it would be cool to try and have a panel C that tries to quantify the different ‘data ingredients’ you would need to try and construct a network, this would probably be very visually overwhelming though...)








Figure 2: A schematic overview of the feasibility-reality axis that defines species interactions (and the resulting networks), specifically in the context of the underlying mechanisms that determine the 'rules of engagement'. As one moves from feasibility towards reality the way in which the edges between nodes are defined (but not necessarily quantified) begins to shift from representing the *possibility* of an interaction occurring between two species to the quantification of *strength* of the interaction. This is because as we move down this axis




Figure 3: Dendrogram of the trait table using a hierarchical clustering model, This is based off of the traits table in SuppMat 2)




Figure 4: Difference between real and model network property. S1 - S5 represent the different motif structures identified in Stouffer et al. (2007) which are S1: Number of linear chains, S2: Number of omnivory motifs, S4: Number of apparent competition motifs, and S5: Number of direct competition motifs