

# **Unveiling the Complexity of Food Webs: A Comprehensive Overview of Definitions, Scales, and Mechanisms**

Tanya Strydom <sup>1</sup>; Jennifer A. Dunne <sup>2</sup>; Timothée Poisot <sup>3,4</sup>; Andrew P. Beckerman <sup>1</sup>

**Abstract:** Food webs are a useful abstraction and representation of the feeding links between species in a community and are used to infer many ecosystem level processes. However, the different theories, mechanisms, and criteria that underpin how a food web is defined and, ultimately, constructed means that not all food webs are representing the same ecological process. Here we present a synthesis of the different assumptions, scales and mechanisms that are used to define different ecological networks ranging from metawebs (an inventory of all potential interactions) to fully realised networks (interactions that occur within a given community over a certain timescale). Illuminating the assumptions, scales, and mechanisms of network inference allows a formal categorisation of how to use networks to answer key ecological and conservation questions and defines guidelines to prevent unintentional misuse or misinterpretation.

**Keywords:** food web, network construction, scientific ignorance

1 At the heart of modern biodiversity science are a set of concepts and theories about biodiversity, stability  
2 and function. These relate to the abundance, distribution and services that biodiversity provides, and how  
3 biodiversity – as an interconnected set of species – responds to multiple stressors. The interaction between  
4 species (or individuals) is one of the fundamental building blocks of ecological communities provide a powerful  
5 abstraction that can help quantify, conceptualise, and understand biodiversity dynamics, and ultimately,  
6 one hopes, make prediction, mitigate change and manage services [ref]. Such network representations of  
7 biodiversity (including within species diversity) are increasingly argued to be an asset to predictive ecology,  
8 climate change mitigation and resource management. Here, it is argued that characterising biodiversity in  
9 a network will allow deeper capacity to understand and predict the abundance, distribution, dynamics and  
10 services provided by multiple species facing multiple stressors.

11 However, the way that a network is constructed (encoded) defines an epistemology of the network concept  
12 which, we argue, can influence the resulting observations and conclusions about pattern and mechanisms  
13 that are made (Brimacombe et al., 2023; Proulx et al., 2005). This process of constructing networks has two  
14 major pillars: the data and theory, the latter representing an expression of mechanism and process giving  
15 rise to patterns that emerge from collating interactions among species. Each of these pillars carries with it  
16 a set of practical, semantic and conceptual constraints that not only influence progress in making network  
17 ecology more valuable and potentially predictive, but help define the spatial, temporal and evolutionary scale  
18 of assumptions we make and predictions we might generate from the networks.

19 With respect to data, it is extremely challenging to actually record species interactions in the field (Jordano,  
20 2016a, 2016b). Despite notable herculean efforts (**Woodward? Benguela?** Maiorano et al. (2020)), actual  
21 coverage of ‘real world’ interaction data remains sparse (Poisot et al., 2021). Against this practical challenge,  
22 there is additionally high variance in the terminology we use to define networks. Finally, the mathematical  
23 and statistical tools we use to construct, conceptualise, analyse and predict with these networks are also  
24 highly variable.

25 1. what are the underlying assumptions about nodes, edges, scale and process that are made when we  
26 attempt to delimit and describe a food webs;

27 2. are there families of commonly used tools that map onto assumptions about scales and processes;

28 The provision of this detail ultimately leads to a set of insights and conclusions about whether, when and  
29 under what conditions network representations of biodiversity can contribute to the advancement of ecological  
30 theory and generate value in predictive ecology. Specifically, we finish this perspective with an overview of  
31 fundamental questions in ecology that we think can benefit from network thinking and a proposal that such

32 thinking can accelerate our capacity to predict the impact of multiple stressors on biodiverse communities.

## 33 1 Setting the Scene: The Not So Basics of Nodes and Edges

34 Defining a food web seems simple; it is the representation of the interactions (edges) between species (nodes),  
35 however the definition of ‘edges’ and ‘nodes’, as well as the scale at which they are aggregated can take many  
36 forms (Poisot, Stouffer, et al., 2016), which ultimately encodes a series of assumptions and criteria within  
37 a network. An awareness of variance in the way a food web can be defined is critical as a network (or its  
38 adjacency matrix) is both the ‘object’ from which inferences are made (*e.g.*, the interactions between species,  
39 or how the structure influences ecosystem level processes) as well as the ‘product’ of either the data collection  
40 (Brimacombe et al., 2023) or prediction process (Banville et al., 2024). One thus needs to be aware of both  
41 the criteria that is used to define nodes and edges, and what processes or mechanisms the aggregation of the  
42 two represents, as this will ultimately determine and delimit the way in which a network can and should be  
43 used.

### 44 1.0.1 How do we define a node?

45 Although this may seem an elementary question in the context of food webs — a node *should* represent a  
46 (taxonomic) species, the reality is that nodes can often represent an aggregation of different species - so called  
47 ‘trophic species’ (Williams & Martinez, 2000; Yodzis, 1982) or segregation of species by life stages (Clegg  
48 et al., 2018). Practical implications of how we are aggregating the nodes is that the resolution may not  
49 always be ‘pixel perfect’, which limits the ability to make (taxonomic) species specific inferences *e.g.*, does  
50 species *a* eat species *b*, however there is value in having nodes that represent an aggregation of species, as  
51 the distribution of the links between them are more meaningful in terms of understanding energy flow and  
52 distribution within the system.

### 53 1.0.2 What is meant by an edge?

54 At its core, links within food webs can be thought of as a representation of either feeding links between  
55 species - be that realised (Pringle, 2020) or potential (Dunne, 2006), or representative of fluxes within  
56 the community/system *e.g.*, energy transfer or material flow (Lindeman, 1942). How we specify links will  
57 influence the resulting structure of the network - and the inferences we will make thereof. For example taking  
58 a food web that consists of links representing all *potential* feeding links for a community (*i.e.*, a metaweb)  
59 will be meaningless if one is interested in understanding the flow of energy through the network as the links  
60 within a metaweb do not represent environmental/energetic constraints, making them poor representations

of which interactions are *realised* in a specific location (Caron et al., 2024). In addition to the various ways of defining the links between species pairs there are also a myriad of ways in which the links themselves can be quantified. Links between species are often treated as being present or absent (*i.e.*, binary) but it is also possible to use probabilities (Banville et al., 2024; which quantifies how likely an interaction is to occur, Poisot, Cirtwill, et al., 2016) or continuous measurements (which quantifies the strength of an interaction, Berlow et al., 2004).

### 1.0.3 Network representations

Broadly, networks can be thought of to fall into two different ‘types’; namely metawebs; traditionally defined as all of the *potential* interactions for a specific species pool (Dunne, 2006), and realised networks; which is the subset of interactions in a metaweb that are *realised* ‘on the ground’. The fundamental difference between these two different types of networks is that a metaweb provides insight as to the viability of an interaction between two species occurring and is a means to identify links that are not ecologically plausible, *i.e.*, forbidden links (Jordano, 2016b), or an idea of the *complete* diet of a species (Strydom et al., 2023). Although metawebs are typically ‘constrained’ to a collection of species that also co-occur, there is no reason that a metaweb cannot include species that do not co-occur (although this would require some degree of prediction/assumptions to identify those possible interactions). In contrast realised networks are highly localised and contingent on both the co-occurrence of species as well as the influence of the environment, and population and community dynamics on predator choice. In the context of definitions and semantics the links that are represented by a metaweb and a realised network are different; links that are absent in a metaweb can be treated as being truly absent, however links that are absent in a realised network cannot be considered to be truly absent but are rather as absent due to the broader environmental/community context. Importantly, a realised network is *not* simply the downscaling of a metaweb to a smaller scale (*e.g.*, moving from the country to the 1x1 km<sup>2</sup> scale based on fine-scale species co-occurrence) but represents a shift towards capturing the higher level processes that determine the *realisation* of an interaction. Thus, different network representations are determined and constrained by different sets of assumptions as to what the processes are that determine the presence/absence of an interaction between two species as well as the resulting network structure.

## 2 From Nodes and Edges to Scales, Context, and Processes

The interplay between network representation and network definition is primarily governed by the process(es) that determine the interaction between species, however these processes are also scale and context dependent.

91 Here we start by introducing the five core processes that determine either the feasibility or realisation of  
92 interactions, namely: evolutionary compatibility, co-occurrence, feasibility, abundance, predator choice, and  
93 non-trophic interactions; while simultaneously contextualising them within, and linking them to the different  
94 network representations Figure 1; specifically if the processes captures an all-or-nothing (possibility) vs  
95 context dependent (likelihood) determination of interactions between species. Of course these processes do  
96 not function in a vacuum and do interact with/influence one another, but it is still beneficial to present them  
97 in a categorical manner as these different processes are often the underpinning logic in the development of  
98 prediction/network models, the criteria for data collection in the field, and the scale of organisation for which  
99 they are relevant (species, population, or community).

100 [Figure 1 about here.]

## 101 **2.1 The processes that determine species interactions**

### 102 **Evolutionary compatibility**

103 There is compelling evidence that the possibility of an interaction occurring between two species is the result  
104 of their shared (co)evolutionary history (Dalla Riva & Stouffer, 2016; Gómez et al., 2010; Segar et al., 2020).  
105 In the more proximal sense this is manifested as the ‘trait complementarity’ between two species, whereby  
106 one species (the predator) has the ‘correct’ set of traits that allow it to chase, capture, kill, and consume  
107 the other species (the prey). For species pairs where this condition is not met the link is deemed to be  
108 forbidden (Jordano, 2016b); *i.e.*, not physically possible and will always be absent within the network. In the  
109 context of trying to determine the feasibility (*i.e.*, the *possibility*) of an interaction, phylogeny is an excellent  
110 predictor (Fricke et al., 2022; Strydom et al., 2022) and allows one to construct what can be considered to  
111 be a metaweb. In terms of thinking about the anatomy of an ‘feasibility network’ one should be aware that  
112 it is possible to represent interactions as either binary (feasible/forbidden; *i.e.*, the traditional definition of a  
113 metaweb Dunne (2006)) or as a probability (Banville et al., 2024), where the probability represents how likely  
114 that the interaction between two species is feasible (what is the possibility of this interaction occurring?).

### 115 **(Co)occurrence**

116 Although the outright assumption that because two species are co-occurring it must mean that they are  
117 interacting is inherently flawed (Blanchet et al., 2020), it is of course impossible for two species to interact (at  
118 least in terms of feeding links) if they are not co-occurring in time and space. Thus co-occurrence data alone is  
119 insufficient to build an accurate and ecologically meaningful representation of a food web having information  
120 on the co-occurrence of species can further aid us in refining metawebs by allowing us to downsample the

network based on the species found in a specific location, or even add additional uncertainty based in how likely species are to co-occur (Dansereau et al., 2024). Additionally the interplay between the interaction between a species pair and their co-occurrence is meaningful when one is operating in the space of trying to determine the distribution of a species (Higino et al., 2023), and forms a key component of some of the next generation species distribution models *e.g.*, joint SDMs (Pollock et al., 2014).

## 126 Abundance

The abundance of the different species within the community can influence the likelihood of an interaction occurring in a myriad of ways. There is the argument that networks (and the interactions that make them up) are driven by only the abundance of the different species and not the characteristics (traits), *sensu* neutral processes and have been formalised with the neutral model (Canard et al., 2012), as well as statistical tools (Momal et al., 2020). Alternatively the abundance of species in a community can influence which interactions are ultimately realised (Banville et al., 2024; Poisot et al., 2015).

## 133 Predator choice (energetic cost)

Ultimately, predator choice is underpinned by the energetic cost-benefit of trying to catch, kill, and consume prey, and is well described within both optimal foraging theory [ref] and metabolic theory [ref], which rests on the idea that the prey a predator chooses to target is one that will have the greatest return on energy with the lowest energetic cost. With a body of evidence that suggests that body size might be the underlying driver, and thus suitable proxy for understanding these processes (Yodzis & Innes, 1992) There are additional bodies of work that attempt to include the cost of movement that the environment imposes on an individual (Cherif et al., 2024) as well as 2D/3D search space (Pawar et al., 2012).

## 141 Indirect interactions

The realisation (presence/absence) or strength of trophic interactions themselves can also be modified by other, indirect (non-trophic), interactions (Golubski & Abrams, 2011; Pilosof et al., 2017), this can be either ‘directly’ through *e.g.*, competition or ‘indirectly’ *e.g.*, mutualistic/facilitative interactions will alter the fine-scale distribution and abundance of some species (Kéfi et al., 2012, 2015).

## 146 2.2 Contextualising the processes that determine species interactions

It should be self evident that the different processes discussed above are all ultimately going to influence the realisation of interactions as well as the structure of a network, however they are acting at different scales of organisation. Both the **co-occurrence** and the **evolutionary compatibility** are valid at the scale of the species pair of interest, that is the *possibility* of an interaction being present/absent is assessed at the

pairwise level and one is left with a ‘list’ of interactions that are present/absent. Although it is possible to build a network (*i.e.*, metaweb) from this information it is important to be aware that the structure of this network is not constrained by real-world dynamics or conditions (*i.e.*, community context), and so just because species are able to interact does not mean that they will (Poisot et al., 2015). In order to construct a network whose structure is a closer approximation of reality (localised interactions) one needs to take into consideration properties of the community as a whole and not just the two species of interest, which requires more data at the community scale, such as the abundance of species.

*something about ‘physical’/landscape scale as well as time scale??*

### 3 Network construction is nuanced

The act of constructing a ‘real world’ network will ultimately be delimited by its intended use, however the reality is that the empirical collection of interaction data is both costly [ref] and challenging to execute in a way that captures the different processes (owing to the different time and spatial scales they may be operating at). Thus we often turn to models to either predict networks, be that the interaction between two species, or its structure (Strydom, Catchen, et al., 2021), or as a means to identify missing interactions (gap fill) within existing empirical dataset (Biton et al., 2024; Stock, 2021), and so for the purpose of this discussion network construction will be synonymous with using a model as a means to represent or predict a network — it can be argued that even the collection of empirical data is in and of itself a ‘model’ as it is still only a *representation* of the system. Different models have different underlying philosophies that often only capture one or a few of the processes discussed in Section 2, has implications for how the resulting network is defined Section 1, which will ultimately delimit and define what inferences can be made from the resulting network. Here we will introduce the three different types of network representations, how they link back to the different processes determining interactions Figure 1, and broadly discuss some of the modelling approaches that are used to construct these different network types. This is paralleled by a hypothetical case study (Box 1) where we showcase the utility/applicability of the different network representation in the context of trying to understand the feeding dynamics of a seasonal community.

#### **i** Box 1 - Why we need to aggregate networks at different scales: A hypothetical case study

Although it might seem most prudent to be predicting, constructing, and defining networks that are the closest representation of reality there are pros and cons of constructing both realised networks as well as metawebs. Let us take for example a community across time/through seasons. In this community we

expect species to be either present or absent depending on the season (*i.e.*, changes in co-occurrence) as well as some species exhibiting seasonal diet shifts, these details would be lost at the scale of the metaweb and it would be valuable to construct either smaller metawebs for the different seasonal communities (thereby capturing the changes in community diversity), or realised networks for each season (to capture diet or ecosystem process shifts). However, these small-scale networks lack the context of the bigger picture that is available at the metaweb - that is it gives us a more holistic idea of the entire diet range of a specific species, which is important when one needs to make conservation-based/applied decisions (*e.g.*, conserving the entire diet of a species and not just seasonal prey items) as well as providing information on interactions that may be possible regardless of the environmental/community context (species may have the capacity to consume certain prey items but do not do so due to local conditions).

With this in mind let us see how the different network aggregations can be used

#### **1: A global metaweb**

Knowledge of the entire diet breadth of a species is valuable especially in terms of understanding how a species will respond to changes in the community - *e.g.*, invasions/rewilding exercises (where does the new species ‘fit’ within the network?) as well as potential capacity to shift its diet. Although this might make sense across space and not time but certain species act as links across the landscape [Rooney]

#### **2: A seasonal metaweb**

Knowledge at the finer scale is also valuable to understand/identify that there are in fact differences between the seasons

#### **3: A seasonal realised network**

Dynamics are useful because they are a representation of the different configurations/energy flows/ecosystem processes. Also to detect more nuanced shifts in diet - *e.g.*, seasonal diet shifts.

#### **4: A structural network**

##### **Data trade off**

Above we highlight the practical uses of the different network configurations but we also need to take into consideration the barriers to construction/associated data needs/cost and acknowledge them. Basically in the ideal world we would have all this information at hand but in reality we might be sitting with seasonal metawebs...

178 **3.1 How do we predict food webs?**

179 There as many ways to predict networks as what there is to define them and along with taking into consid-  
180 eration the points raised in the previous section it is also beneficial to think about the context in which the  
181 different models were developed - and how this will influence the networks that they produce...

182 There is a bit of a ‘point of conflict’ between those calling for ‘pixel perfect’, regional scale data (Pringle,  
183 2020; Pringle & Hutchinson, 2020) and for the means to generate networks that are ecologically plausible  
184 *representations* (*sensu* structural networks). This represents two challenges; one is that models that repre-  
185 sent generalisations of networks often lack the ability to retrieve any species/community specificity which  
186 limits their utility for real world, species-driven scenarios *e.g.*, species driven conservation efforts (Dunn et  
187 al., 2009), however networks that are constructed through either (most) empirical observations or through  
188 predictive means are fundamentally going to represent metawebs, *i.e.*, lack constrained links, a representation  
189 of structure, or energy flow...

190 **3.1.1 Models that predict metawebs (feasible interactions)**

191 This is perhaps the most developed group of models; with a variety of approaches having been developed that  
192 typically determine the feasibility of an interaction based on the trait compatibility between predator and  
193 prey (*i.e.* their evolutionary compatibility) to determine ‘feeding rules’ (Morales-Castilla et al., 2015). These  
194 feeding rules are broadly elucidated in two different ways; mechanistic feeding rules can be explicitly defined  
195 and applied to a community (Dunne et al., 2008; *e.g.*, Shaw et al., 2024) or they are inferred from a community  
196 for which there is interaction data and the ‘rules’ are then applied to a different community (Caron et al.,  
197 2022; Cirtwill et al., 2019; Desjardins-Proulx et al., 2017; Eklöf et al., 2013; Llewelyn et al., 2023; Pichler et  
198 al., 2020; Strydom et al., 2022; *e.g.*, Strydom et al., 2023). The fundamental difference between these two  
199 model groups is that ‘mechanistic models’ rely on expert knowledge and make assumptions on trait-feeding  
200 relationships, whereas the ‘pattern finding’ models are dependent on existing datasets from which to elucidate  
201 feeding rules. These models are useful for determining all feasible interactions for a specific community, and  
202 owing to the availability of datasets (Gray et al., 2015; *e.g.*, Poelen et al., 2014; Poisot, Baiser, et al., 2016),  
203 as well as the development of model testing/benchmarking tools (Poisot, 2023), means that these models  
204 can be validated and (with relative confidence) be used to construct first draft networks for communities for  
205 which we have no data (Strydom et al., 2022), and are valuable for constructing networks where we lack any  
206 interaction data *e.g.*, prehistoric networks (Fricke et al., 2022; Yeakel et al., 2014).

207 **3.1.2 Models that predict realised networks (realised interactions)**

208 In order to construct realised networks models need to incorporate *both* the feasibility of interactions (*i.e.*,  
209 determine the entire diet breadth of a species) as well as then determine which interactions are realised (*i.e.*,  
210 incorporate the ‘cost’ of interactions). As far as we are aware there is no model that explicitly accounts for  
211 both of these ‘rules’ and rather *only* account for processes that determine the realisation of an interaction (*i.e.*,  
212 abundance, predator choice, or non-trophic interactions). Although the use of allometric scaling *i.e.*, body  
213 size (Beckerman et al., 2006; *e.g.*, Valdovinos et al., 2023) may represent a first step in capturing evolutionary  
214 compatibility one still needs to account for other feeding traits. In terms of models that do formalise these  
215 processes, diet models (Beckerman et al., 2006; Petchey et al., 2008) have been used construct networks  
216 based on both predator choice (as determined by the handling time, energy content, and predator attack  
217 rate) as well as abundance (prey density). Wootton et al. (2023) developed a model that moves the energy  
218 of the system into different modules related to the process of the predator acquiring energy from the prey  
219 *i.e.*, compartmentation in food webs (Krause et al., 2003).

220 **3.1.3 Models that predict structure (interaction agnostic)**

221 Although we identify mechanisms that determine species interactions in Section 2 not all models that are used  
222 to predict networks explicitly operate at the ‘process’ level, but rather represent the *structure* of a network  
223 based on a series of *a priori* assumptions as to the distribution of links between species (typically trophic not  
224 taxonomic species) by parametrising an aspect of the network structure, (*e.g.*, the niche model (Williams &  
225 Martinez, 2000) makes an assumption as to the expected connectance of the network, although see Allesina  
226 & Pascual (2009) for a parameter-free model) or alternatively uses structural features of an exiting *realised*  
227 network (*e.g.*, stochastic block model, Xie et al. (2017)). Importantly these structural models do not make  
228 species specific predictions (they are usually species agnostic and treat nodes as trophic species) and so cannot  
229 be used to determine if an interaction is either possible *or* realised between two species (*i.e.*, one cannot use  
230 these models to determine if species *a* eats species *b*). Although this means this suite of models are unsuitable  
231 as tools for predicting species-specific interactions, they have been shown to be sufficient tools to predict the  
232 structure of networks (Williams & Martinez, 2008), and provide a data-light (the models often only require  
233 species richness) but assumption heavy (the resulting network structure is determined by an assumption of  
234 network structure) way to construct a network.

235 **4 Making Progress with Networks**

236 **4.1 Further development of models and tools**

237 There has been a suite of models that have been developed to predict trophic links, however we are lacking in  
238 tools that are explicitly taking into consideration estimating both the feasibility as well as realisation of links,  
239 *i.e.*, both interactions and structure simultaneously (Strydom, Catchen, et al., 2021). This could be addressed  
240 either through the development of tools that do both (predict both interactions and structure), or to develop  
241 an ensemble modelling approach (Becker et al., 2022). Alternatively the development of tools that will allow  
242 for the downsampling of metawebs into realised networks (*e.g.*, Roopnarine, 2006), although deciding exactly  
243 what is driving differences between local networks and the regional metaweb might not be that simple (Saravia  
244 et al., 2022). Probably also something that aligns with trying to predict interaction strength - because that  
245 would be the gold standard. Probably also worth just plainly stating that feasibility of developing a model  
246 that is both broadly generalisable, but also has local specificity is probably not attainable (Stouffer, 2019),  
247 and more specifically the potential use in models untangling/identifying the different processes that shape  
248 interaction networks (Song & Levine, 2024), *e.g.*, Curtsdotter et al. (2019) showcasing the use of models to  
249 disentangle the drivers of community function and Strydom, Dalla Riva, et al. (2021) who identified that  
250 networks are less complex than they could be, suggesting that there are constraints on network assembly.  
251 In addition to the more intentional development of models we also need to consider the validation of these  
252 models, there have been developments and discussions for assessing how well a model recovers pairwise  
253 interactions (Poisot, 2023; Strydom, Catchen, et al., 2021), although the rate of false-negatives that may be  
254 present in the testing data still present a challenge (Catchen et al., 2023), and we still lack clear strategies  
255 for benchmarking the ability of models to recover structure (Allesina et al., 2008).

256 **4.1.1 At what scale should we be predicting and using networks?**

257 Look at Hutchinson et al. (2019)

258 We lack a clear agenda (and conceptualisation) as to what the appropriate level of aggregation is for a  
259 ‘network’. Realistically most empirical networks are more aligned with metawebs as opposed to realised  
260 networks as they are often the result of some sort of aggregation of observations across time, this creates a  
261 two-fold problem. Firstly, we need to think about how this affects any sort of development of theory that sits  
262 closer to the ‘realised network’ side of the spectrum - how often are we trying to ask and answer questions  
263 about realised networks using feasible networks? The second is that this lack of ‘direction’ as to how we  
264 should define a network is (actually) probably one of the biggest barriers that is affecting the use of networks

in applied settings... By define I mean both delimiting the time and geographic scale at which a network is aggregated at (Estay et al., 2023). This is important because it can influence the inferences made, *e.g.*, the large body of work (landscape theory for food web architecture) that showcases how different species use the landscape will influence network dynamics (Rooney et al., 2008). There is also a bit of an interplay with time and data and the different scales that they may be integrated at - co-occurrence may span decades and just because two species have been recorded in the same space does not mean it was at the same timescale (Brimacombe et al., 2024)

#### 4.1.2 Feasible, realised, or sustainable?

When do we determine a link to be ‘real’... In the context of metawebs this is perhaps clearer - if all things were equal (*i.e.*, community context is irrelevant) would the predator be able to consume the prey. However in the realised space there is also the question of the long term ‘energetic feasibility’ of an interaction - just because an interaction is possible in the now is it able to sustain a population in the long term. And what is the scale for that long term - are we thinking at the generational scale? Because ultimately when we are constructing a network we are aggregating not only across space but also across time... This is probably again a Lokta-Volterra space question and something that the dynamic foodweb model (Curtsdotter et al., 2019; Delmas et al., 2017; Lajaaiti et al., 2024) is addressing, but again it is integrating this with the feasible/realised axis.

## 4.2 How should we use different networks?

What for and how we can use networks is perhaps one of the biggest ‘gaps’ we have in network ecology (Tim’s EBV ms), and there is a serious need to start drawing clear, ecological links between network form and function (although see Delmas et al., 2019). That being said one of the most important things we can do is to be aware of the parameter space that is possible given a specific definition of a network and operate within those parameters. Here we can maybe tie it back to scale - specifically the idea that the fact that metawebs operate at evolutionary scales they are not suitable for ‘dynamic’ processes, although they do have the potential to think about novel species entering the network/community...

## 5 Concluding remarks

I think a big take home will (hopefully) be how different approaches do better in different situations and so you as an end user need to take this into consideration and pick accordingly. I think Petchey et al. (2011) might have (and share) some thoughts on this. I feel like I need to look at Berlow et al. (2008) but maybe not exactly in this context but vaguely adjacent. This is sort of the crux of the argument presented in

- 294 Brimacombe et al. (2024) as well.
- 295 Do we expect there to be differences when thinking about unipartite vs bipartite networks? Is there underlying  
296 ecology/theory that would assume that different mechanisms (and thus models) are relevant in these two  
297 ‘systems’.
- 298 • The Terry & Lewis (2020) paper looks at some methods but is specifically looking at a bipartite world...

## 299 References

- 300 Allesina, S., Alonso, D., & Pascual, M. (2008). A General Model for Food Web Structure. *Science*, 320(5876),  
301 658–661. <https://doi.org/10.1126/science.1156269>
- 302 Allesina, S., & Pascual, M. (2009). Food web models: A plea for groups. *Ecology Letters*, 12(7), 652–662.  
303 <https://doi.org/10.1111/j.1461-0248.2009.01321.x>
- 304 Banville, F., Strydom, T., Blyth, P., Brimacombe, C., Catchen, M. D., Dansereau, G., Higino, G., Malpas,  
305 T., Mayall, H., Norman, K., Gravel, D., & Poisot, T. (2024). *Deciphering probabilistic species interaction*  
306 *networks*. EcoEvoRxiv. <https://doi.org/10.32942/X28G8Z>
- 307 Becker, D. J., Albery, G. F., Sjodin, A. R., Poisot, T., Bergner, L. M., Chen, B., Cohen, L. E., Dallas, T.  
308 A., Eskew, E. A., Fagre, A. C., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Stock, M., Teeling,  
309 E. C., & Carlson, C. J. (2022). Optimising predictive models to prioritise viral discovery in zoonotic  
310 reservoirs. *The Lancet Microbe*, 3(8), e625–e637. [https://doi.org/10.1016/S2666-5247\(21\)00245-7](https://doi.org/10.1016/S2666-5247(21)00245-7)
- 311 Beckerman, A. P., Petchey, O. L., & Warren, P. H. (2006). Foraging biology predicts food web complexity.  
312 *Proceedings of the National Academy of Sciences*, 103(37), 13745–13749. <https://doi.org/10.1073/pnas.0603039103>
- 313 Berlow, E. L., Brose, U., & Martinez, N. D. (2008). The “Goldilocks factor” in food webs. *Proceedings of  
314 the National Academy of Sciences*, 105(11), 4079–4080. <https://doi.org/10.1073/pnas.0800967105>
- 315 Berlow, E. L., Neutel, A.-M., Cohen, J. E., de Ruiter, P. C., Ebenman, B., Emmerson, M., Fox, J. W., Jansen,  
316 V. A. A., Iwan Jones, J., Kokkoris, G. D., Logofet, D. O., McKane, A. J., Montoya, J. M., & Petchey, O.  
317 (2004). Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73(3),  
318 585–598. <https://doi.org/10.1111/j.0021-8790.2004.00833.x>
- 319 Biton, B., Puzis, R., & Pilosof, S. (2024). *Inductive link prediction boosts data availability and enables  
320 cross-community link prediction in ecological networks*.
- 321 Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions.  
322 *Ecology Letters*, 23(7), 1050–1063. <https://doi.org/10.1111/ele.13525>

- 324 Brimacombe, C., Bodner, K., & Fortin, M.-J. (2024). *Applying a method before its proof-of-concept: A*  
325 *cautionary tale using inferred food webs.* <https://doi.org/10.13140/RG.2.2.22076.65927>
- 326 Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings  
327 of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4),  
328 e3002068. <https://doi.org/10.1371/journal.pbio.3002068>
- 329 Canard, E., Mouquet, N., Marescot, L., Gaston, K. J., Gravel, D., & Mouillot, D. (2012). Emergence of  
330 Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7(8), e38295. <https://doi.org/10.1371/journal.pone.0038295>
- 332 Caron, D., Brose, U., Lurgi, M., Blanchet, F. G., Gravel, D., & Pollock, L. J. (2024). Trait-matching models  
333 predict pairwise interactions across regions, not food web properties. *Global Ecology and Biogeography*,  
334 33(4), e13807. <https://doi.org/10.1111/geb.13807>
- 335 Caron, D., Maiorano, L., Thuiller, W., & Pollock, L. J. (2022). Addressing the Eltonian shortfall with  
336 trait-based interaction models. *Ecology Letters*, 25(4), 889–899. <https://doi.org/10.1111/ele.13966>
- 337 Catchen, M. D., Poisot, T., Pollock, L. J., & Gonzalez, A. (2023). *The missing link: Discerning true from*  
338 *false negatives when sampling species interaction networks.*
- 339 Cherif, M., Brose, U., Hirt, M. R., Ryser, R., Silve, V., Albert, G., Arnott, R., Berti, E., Cirtwill, A.,  
340 Dyer, A., Gauzens, B., Gupta, A., Ho, H.-C., Portalier, S. M. J., Wain, D., & Wootton, K. (2024). The  
341 environment to the rescue: Can physics help predict predator-prey interactions? *Biological Reviews*,  
342 n/a(n/a). <https://doi.org/10.1111/brv.13105>
- 343 Cirtwill, A. R., Ekklf, A., Roslin, T., Wootton, K., & Gravel, D. (2019). A quantitative framework for  
344 investigating the reliability of empirical network construction. *Methods in Ecology and Evolution*, 10(6),  
345 902–911. <https://doi.org/10.1111/2041-210X.13180>
- 346 Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on food web structure.  
347 *Ecology*, 99(12), 2712–2720. <https://doi.org/10.1002/ecy.2523>
- 348 Curtsdotter, A., Banks, H. T., Banks, J. E., Jonsson, M., Jonsson, T., Laubmeier, A. N., Traugott, M., &  
349 Bommarco, R. (2019). Ecosystem function in predator-prey food webs—confronting dynamic models with  
350 empirical data. *Journal of Animal Ecology*, 88(2), 196–210. <https://doi.org/10.1111/1365-2656.12892>
- 351 Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food webs' backbones  
352 using functional traits. *Oikos*, 125(4), 446–456. <https://doi.org/10.1111/oik.02305>
- 353 Dansereau, G., Barros, C., & Poisot, T. (2024). Spatially explicit predictions of food web structure from  
354 regional-level data. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 379(1909).  
355 <https://doi.org/10.1098/rstb.2023.0166>
- 356 Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Riva, G. V. D., Fortin, M.-J., Gravel, D., Guimarães,

- 357 P. R., Hembry, D. H., Newman, E. A., Olesen, J. M., Pires, M. M., Yeakel, J. D., & Poisot, T. (2019).  
358 Analysing ecological networks of species interactions. *Biological Reviews*, 94(1), 16–36. <https://doi.org/10.1111/brv.12433>
- 360 Delmas, E., Brose, U., Gravel, D., Stouffer, D. B., & Poisot, T. (2017). Simulations of biomass dynamics in  
361 community food webs. *Methods in Ecology and Evolution*, 8(7), 881–886. <https://doi.org/10.1111/2041-210X.12713>
- 363 Desjardins-Proulx, P., Laigle, I., Poisot, T., & Gravel, D. (2017). Ecological interactions and the Netflix  
364 problem. *PeerJ*, 5, e3644. <https://doi.org/10.7717/peerj.3644>
- 365 Dunn, R. R., Harris, N. C., Colwell, R. K., Koh, L. P., & Sodhi, N. S. (2009). The sixth mass coextinction:  
366 Are most endangered species parasites and mutualists? *Proceedings. Biological Sciences*, 276(1670),  
367 3037–3045. <https://doi.org/10.1098/rspb.2009.0413>
- 368 Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological  
369 networks: Linking structure and dynamics* (pp. 27–86). Oxford University Press.
- 370 Dunne, J. A., Williams, R. J., Martinez, N. D., Wood, R. A., & Erwin, D. H. (2008). Compilation and  
371 Network Analyses of Cambrian Food Webs. *PLOS Biology*, 6(4), e102. <https://doi.org/10.1371/journal.pbio.0060102>
- 373 Eklöf, A., Tang, S., & Allesina, S. (2013). Secondary extinctions in food webs: A Bayesian network approach.  
374 *Methods in Ecology and Evolution*, 4(8), 760–770. <https://doi.org/10.1111/2041-210X.12062>
- 375 Estay, S. A., Fortin, M.-J., & López, D. N. (2023). Editorial: Patterns and processes in ecological networks  
376 over space. *Frontiers in Ecology and Evolution*, 11.
- 377 Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning,  
378 J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene.  
379 *Science*, 377(6609), 1008–1011. <https://doi.org/10.1126/science.abn4012>
- 380 Golubski, A. J., & Abrams, P. A. (2011). Modifying modifiers: What happens when interspecific interactions  
381 interact? *Journal of Animal Ecology*, 80(5), 1097–1108. <https://doi.org/10.1111/j.1365-2656.2011.01852.x>
- 383 Gómez, J. M., Verdú, M., & Perfectti, F. (2010). Ecological interactions are evolutionarily conserved across  
384 the entire tree of life. *Nature*, 465(7300), 918–921. <https://doi.org/10.1038/nature09113>
- 385 Gray, C., Figueroa, D. H., Hudson, L. N., Ma, A., Perkins, D., & Woodward, G. (2015). Joining the dots:  
386 An automated method for constructing food webs from compendia of published interactions. *Food Webs*,  
387 5, 11–20. <https://doi.org/10.1016/j.fooweb.2015.09.001>
- 388 Higino, G. T., Banville, F., Dansereau, G., Muñoz, N. R. F., Windsor, F., & Poisot, T. (2023). Mismatch  
389 between IUCN range maps and species interactions data illustrated using the Serengeti food web. *PeerJ*,

- 390 11, e14620. <https://doi.org/10.7717/peerj.14620>
- 391 Hutchinson, M. C., Bramon Mora, B., Pilosof, S., Barner, A. K., Kéfi, S., Thébault, E., Jordano, P., &  
392 Stouffer, D. B. (2019). Seeing the forest for the trees: Putting multilayer networks to work for community  
393 ecology. *Functional Ecology*, 33(2), 206–217. <https://doi.org/10.1111/1365-2435.13237>
- 394 Jordano, P. (2016a). Chasing Ecological Interactions. *PLOS Biology*, 14(9), e1002559. <https://doi.org/10.1371/journal.pbio.1002559>
- 395 Jordano, P. (2016b). Sampling networks of ecological interactions. *Functional Ecology*. <https://doi.org/10.1111/1365-2435.12763>
- 396 Jordano, P. (2016c). Network structure beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky  
397 shores. *Ecology*, 96(1), 291–303. <https://doi.org/10.1890/13-1424.1>
- 398 Kéfi, S., Berlow, E. L., Wieters, E. A., Joppa, L. N., Wood, S. A., Brose, U., & Navarrete, S. A. (2015).  
399 Network structure beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky  
400 shores. *Ecology*, 96(1), 291–303. <https://doi.org/10.1890/13-1424.1>
- 401 Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A., Boit, A., Joppa, L. N.,  
402 Lafferty, K. D., Williams, R. J., Martinez, N. D., Menge, B. A., Blanchette, C. A., Iles, A. C., & Brose,  
403 U. (2012). More than a meal... integrating non-feeding interactions into food webs: More than a meal ....  
404 *Ecology Letters*, 15(4), 291–300. <https://doi.org/10.1111/j.1461-0248.2011.01732.x>
- 405 Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E., & Taylor, W. W. (2003). Compartments  
406 revealed in food-web structure. *Nature*, 426(6964), 282–285. <https://doi.org/10.1038/nature02115>
- 407 Lajaaiti, I., Bonnici, I., Kéfi, S., Mayall, H., Danet, A., Beckerman, A. P., Malpas, T., & Delmas, E. (2024).  
408 *EcologicalNetworksDynamics.jl A Julia package to simulate the temporal dynamics of complex ecological*  
409 *networks* (p. 2024.03.20.585899). bioRxiv. <https://doi.org/10.1101/2024.03.20.585899>
- 410 Lindeman, R. L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23(4), 399–417. <https://doi.org/10.2307/1930126>
- 411 Llewelyn, J., Strona, G., Dickman, C. R., Greenville, A. C., Wardle, G. M., Lee, M. S. Y., Doherty, S.,  
412 Shabani, F., Saltré, F., & Bradshaw, C. J. A. (2023). Predicting predator–prey interactions in terrestrial  
413 endotherms using random forest. *Ecography*, 2023(9), e06619. <https://doi.org/10.1111/ecog.06619>
- 414 Maiorano, L., Montemaggiori, A., Ficetola, G. F., O'Connor, L., & Thuiller, W. (2020). TETRA-EU 1.0: A  
415 species-level trophic metaweb of European tetrapods. *Global Ecology and Biogeography*, 29(9), 1452–1457.  
416 <https://doi.org/10.1111/geb.13138>
- 417 Momal, R., Robin, S., & Ambroise, C. (2020). Tree-based inference of species interaction networks from  
418 abundance data. *Methods in Ecology and Evolution*, 11(5), 621–632. <https://doi.org/10.1111/2041-210X.13380>
- 419 Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from  
420 proxies. *Trends in Ecology & Evolution*, 30(6), 347–356. <https://doi.org/10.1016/j.tree.2015.03.014>

- 423 Pawar, S., Dell, A. I., & Savage, V. M. (2012). Dimensionality of consumer search space drives trophic  
424 interaction strengths. *Nature*, 486(7404), 485–489. <https://doi.org/10.1038/nature11131>
- 425 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2008). Size, foraging, and food web  
426 structure. *Proceedings of the National Academy of Sciences*, 105(11), 4191–4196. <https://doi.org/10.1073/pnas.0710672105>
- 427 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2011). Fit, efficiency, and biology: Some  
428 thoughts on judging food web models. *Journal of Theoretical Biology*, 279(1), 169–171. <https://doi.org/10.1016/j.jtbi.2011.03.019>
- 429 Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine learning algorithms  
430 to infer trait-matching and predict species interactions in ecological networks. *Methods in Ecology and  
431 Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>
- 432 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological networks.  
433 *Nature Ecology & Evolution*, 1(4), 101. <https://doi.org/10.1038/s41559-017-0101>
- 434 Poelen, J. H., Simons, J. D., & Mungall, C. J. (2014). Global biotic interactions: An open infrastructure to  
435 share and analyze species-interaction datasets. *Ecological Informatics*, 24, 148–159. <https://doi.org/10.1016/j.ecoinf.2014.08.005>
- 436 Poisot, T. (2023). Guidelines for the prediction of species interactions through binary classification. *Methods  
437 in Ecology and Evolution*, 14(5), 1333–1345. <https://doi.org/10.1111/2041-210X.14071>
- 438 Poisot, T., Baiser, B., Dunne, J., Kéfi, S., Massol, F., Mouquet, N., Romanuk, T. N., Stouffer, D. B., Wood, S.  
439 A., & Gravel, D. (2016). Mangal – making ecological network analysis simple. *Ecography*, 39(4), 384–390.  
440 <https://doi.org/10.1111/ecog.00976>
- 441 Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., Mercier, B., Violet, C., &  
442 Vissault, S. (2021). Global knowledge gaps in species interaction networks data. *Journal of Biogeography*,  
443 48(7), 1552–1563. <https://doi.org/10.1111/jbi.14127>
- 444 Poisot, T., Cirtwill, A., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. (2016). The structure of  
445 probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312. <https://doi.org/10.1111/2041-210X.12502>
- 446 Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary  
447 through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>
- 448 Poisot, T., Stouffer, D. B., & Kéfi, S. (2016). Describe, understand and predict: Why do we need networks  
449 in ecology? *Functional Ecology*, 30(12), 1878–1882. <https://www.jstor.org/stable/48582345>
- 450 Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O’Hara, R. B., Parris, K. M., Vesk, P. A., &  
451 McCarthy, M. A. (2014). Understanding co-occurrence by modelling species simultaneously with a Joint  
452 Species Distribution Model (JSDM). *Methods in Ecology and Evolution*, 5(5), 397–406. <https://doi.org/10.1111/2041-210X.12502>
- 453

- 456 10.1111/2041-210X.12180
- 457 Pringle, R. M. (2020). Untangling Food Webs. In *Unsolved Problems in Ecology* (pp. 225–238). Princeton  
458 University Press. <https://doi.org/10.1515/9780691195322-020>
- 459 Pringle, R. M., & Hutchinson, M. C. (2020). Resolving Food-Web Structure. *Annual Review of Ecology,*  
460 *Evolution and Systematics*, 51(Volume 51, 2020), 55–80. <https://doi.org/10.1146/annurev-ecolsys-110218-024908>
- 461
- 462 Proulx, S. R., Promislow, D. E. L., & Phillips, P. C. (2005). Network thinking in ecology and evolution.  
463 *Trends in Ecology & Evolution*, 20(6), 345–353. <https://doi.org/10.1016/j.tree.2005.04.004>
- 464 Rooney, N., McCann, K. S., & Moore, J. C. (2008). A landscape theory for food web architecture. *Ecology*  
465 *Letters*, 11(8), 867–881. <https://doi.org/10.1111/j.1461-0248.2008.01193.x>
- 466 Roopnarine, P. D. (2006). Extinction Cascades and Catastrophe in Ancient Food Webs. *Paleobiology*, 32(1),  
467 1–19. <https://www.jstor.org/stable/4096814>
- 468 Saravia, L. A., Marina, T. I., Kristensen, N. P., De Troch, M., & Momo, F. R. (2022). Ecological network  
469 assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, 91(3),  
470 630–642. <https://doi.org/10.1111/1365-2656.13652>
- 471 Segar, S. T., Fayle, T. M., Srivastava, D. S., Lewinsohn, T. M., Lewis, O. T., Novotny, V., Kitching, R. L.,  
472 & Maunsell, S. C. (2020). The Role of Evolution in Shaping Ecological Networks. *Trends in Ecology &*  
473 *Evolution*, 35(5), 454–466. <https://doi.org/10.1016/j.tree.2020.01.004>
- 474 Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024). *A framework for*  
475 *reconstructing ancient food webs using functional trait data* (p. 2024.01.30.578036). bioRxiv. <https://doi.org/10.1101/2024.01.30.578036>
- 476
- 477 Song, C., & Levine, J. M. (2024). *Rigorous (in)validation of ecological models* (p. 2024.09.19.613075). bioRxiv.  
478 <https://doi.org/10.1101/2024.09.19.613075>
- 479 Stock, M. (2021). Pairwise learning for predicting pollination interactions based on traits and phylogeny.  
480 *Ecological Modelling*, 14.
- 481 Stouffer, D. B. (2019). All ecological models are wrong, but some are useful. *Journal of Animal Ecology*,  
482 88(2), 192–195. <https://doi.org/10.1111/1365-2656.12949>
- 483 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V.,  
484 Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction  
485 through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*,  
486 13(12), 2838–2849. <https://doi.org/10.1111/2041-210X.13835>
- 487 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Mercier, B.,  
488 Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2023). Graph embedding and transfer

- 489 learning can help predict potential species interaction networks despite data limitations. *Methods in*  
490 *Ecology and Evolution*, 14(12), 2917–2930. <https://doi.org/10.1111/2041-210X.14228>
- 491 Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz,  
492 N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap  
493 towards predicting species interaction networks (across space and time). *Philosophical Transactions of*  
494 *the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>
- 495 Strydom, T., Dalla Riva, G. V., & Poisot, T. (2021). SVD Entropy Reveals the High Complexity of Ecological  
496 Networks. *Frontiers in Ecology and Evolution*, 9. <https://doi.org/10.3389/fevo.2021.623141>
- 497 Terry, J. C. D., & Lewis, O. T. (2020). Finding missing links in interaction networks. *Ecology*, 101(7), e03047.  
498 <https://doi.org/10.1002/ecy.3047>
- 499 Valdovinos, F. S., Hale, K. R. S., Dritz, S., Glaum, P. R., McCann, K. S., Simon, S. M., Thébault, E., Wetzel,  
500 W. C., Wootton, K. L., & Yeakel, J. D. (2023). A bioenergetic framework for aboveground terrestrial  
501 food webs. *Trends in Ecology & Evolution*, 38(3), 301–312. <https://doi.org/10.1016/j.tree.2022.11.004>
- 502 Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, 404(6774), 180–183.  
503 <https://doi.org/10.1038/35004572>
- 504 Williams, R. J., & Martinez, N. D. (2008). Success and its limits among structural models of complex food  
505 webs. *Journal of Animal Ecology*, 77(3), 512–519. <https://doi.org/10.1111/j.1365-2656.2008.01362.x>
- 506 Wootton, K. L., Curtsdotter, A., Roslin, T., Bommarco, R., & Jonsson, T. (2023). Towards a modular theory  
507 of trophic interactions. *Functional Ecology*, 37(1), 26–43. <https://doi.org/10.1111/1365-2435.13954>
- 508 Xie, J.-R., Zhang, P., Zhang, H.-F., & Wang, B.-H. (2017). Completeness of Community Structure in  
509 Networks. *Scientific Reports*, 7(1), 5269. <https://doi.org/10.1038/s41598-017-05585-6>
- 510 Yeakel, J. D., Pires, M. M., Rudolf, L., Dominy, N. J., Koch, P. L., Guimarães, P. R., & Gross, T. (2014).  
511 Collapse of an ecological network in Ancient Egypt. *PNAS*, 111(40), 14472–14477. <https://doi.org/10.1073/pnas.1408471111>
- 512 Yodzis, P. (1982). The Compartmentation of Real and Assembled Ecosystems. *The American Naturalist*,  
513 120(5), 551–570. <https://doi.org/10.1086/284013>
- 514 Yodzis, P., & Innes, S. (1992). Body Size and Consumer-Resource Dynamics. *The American Naturalist*,  
515 139(6), 1151–1175. <https://doi.org/10.1086/285380>

the "driver" of the interaction  
≠ the underlying theory (1:1) & the  
'way' we get to the answer is diff  
e.g. co-occurrence is about sharing  
space/time but we use niche theory/  
env. filtering as the means to determine  
co-occurrence.

JUN	27	THU	木	179
			(mechanism)	
			what determines	
			interaction	
6		Body of theory		
15	Species	fund. niches env. filter	co- occurrence	
21		buck bones (mora)	"capacity"	
0	pop.	truit-match		
3	indiv.	neutral. funct. resp.	pop size / dynamics	
?? allometric scaling ??				
				: THEORY OF WHAT DETERMINES FEEDING LINKS (INTERACTIONS) BETWEEN SPECIES

Figure 1: TODO.