

# **Unveiling the Complexity of Food Webs: A Comprehensive Overview of Definitions, Scales, and Mechanisms**

Tanya Strydom <sup>1</sup>; Jennifer A. Dunne <sup>2</sup>; Timothée Poisot <sup>3,4</sup>; Andrew P. Beckerman <sup>1</sup>

**Abstract:** Food webs are a useful abstraction and representation of the feeding links between species in a community and are used to infer many ecosystem level processes. However, the different theories, mechanisms, and criteria that underpin how a food web is defined and, ultimately, constructed means that not all food webs are representing the same ecological process. Here we present a synthesis of the different assumptions, scales and mechanisms that are used to define different ecological networks ranging from metawebs (an inventory of all potential interactions) to fully realised networks (interactions that occur within a given community over a certain timescale). Illuminating the assumptions, scales, and mechanisms of network inference allows a formal categorisation of how to use networks to answer key ecological and conservation questions and defines guidelines to prevent unintentional misuse or misinterpretation.

**Keywords:** food web, network construction, scientific ignorance

1 At the heart of modern biodiversity science are a set of concepts and theories about biodiversity, stability  
2 and function. These relate to the abundance, distribution and services that biodiversity provides, and how  
3 biodiversity – as an interconnected set of species – responds to multiple stressors. The interaction between  
4 species is one of the fundamental building blocks of ecological communities, providing a powerful abstraction  
5 that can help quantify, conceptualise, and understand biodiversity dynamics, and ultimately, make predic-  
6 tions, mitigate change, and manage services. Such network representations of biodiversity (including within  
7 species diversity) are increasingly argued to be an asset to predictive ecology, climate change mitigation  
8 and resource management. Here, it is argued that characterising biodiversity in a network will allow deeper  
9 capacity to understand and predict the abundance, distribution, dynamics and services provided by multiple  
10 species facing multiple stressors.

11 A ‘network’ can be defined and conceptualised in a myriad of ways, which means that different networks  
12 will be embedding different processes (or determinants) of interactions, ultimately influencing the patterns  
13 and mechanisms that are inferred (Brimacombe et al., 2023; Proulx et al., 2005). The different ways in  
14 which a network can be represented is the result of *how* the network is constructed, which itself rests on two  
15 pillars: the data used to construct the network and the underlying theory as to what drives the interactions  
16 between species. The latter represents an expression of mechanism and process that gives rise to the patterns  
17 that emerge from collating interactions among species, and will ultimately inform which data are deemed  
18 important in the determination of interactions occurring. Each of these pillars carries with it a set of practical,  
19 semantic and conceptual constraints that not only influence progress in making network ecology more valuable  
20 and potentially predictive, but help define the spatial, temporal, and evolutionary scale of assumptions we  
21 make and the predictions we might generate from different network representations.

22 In this perspective we aim to provide an overview of the different **food web** representations, particularly how  
23 these relate to the terminology used to define a network, and how this influenced by both the processes that  
24 determine networks as well as how this relates to the way in which we construct networks. The provision of  
25 this detail ultimately leads to a set of insights and conclusions about whether, when and under what conditions  
26 network representations of biodiversity can contribute to the advancement of ecological theory and generate  
27 value in predictive ecology. Specifically, we finish this perspective with an overview of fundamental questions  
28 in ecology that we think can benefit from network thinking and a proposal that such thinking can accelerate  
29 our capacity to predict the impact of multiple stressors on biodiverse communities.

# **30 1 Setting the Scene: The Not So Basics of Nodes and Edges**

**31** Defining a food web seems simple; it is the representation of the interactions (edges) between species (nodes)  
**32** in the form of a network, however the definition of ‘edges’ and ‘nodes’, as well as the scale at which they  
**33** are aggregated can take many forms (Poisot, Stouffer, et al., 2016), which ultimately encodes a series of  
**34** assumptions and criteria within a network. An awareness of variance in the way a food web can be defined  
**35** is critical as a network (or its adjacency matrix) is both the ‘object’ from which inferences are made (*e.g.*,  
**36** the interactions between species, or how the structure influences ecosystem level processes) as well as the  
**37** ‘product’ of either the data collection (Brimacombe et al., 2023) or prediction process (Banville et al., 2024).  
**38** One thus needs to be aware of both the criteria that is used to define nodes and edges, and what processes  
**39** or mechanisms the aggregation of the two represents, as this will determine what the network can be used  
**40** for.

## **41 1.1 How do we define a node?**

**42** Although this may seem an elementary question in the context of food webs — a node *should* represent a  
**43** (taxonomic) species, the reality is that nodes can often represent an aggregation of different species - so called  
**44** ‘trophic species’ (Williams & Martinez, 2000; Yodzis, 1982) or segregation of species by life stages (Clegg  
**45** et al., 2018). Practical implications of how we are aggregating the nodes is that the resolution may not  
**46** always be ‘pixel perfect’, which limits the ability to make (taxonomic) species specific inferences *e.g.*, does  
**47** species *a* eat species *b*, however there is value in having nodes that represent an aggregation of species, as  
**48** the distribution of the links between them are more meaningful in terms of understanding energy flow and  
**49** distribution within the system.

## **50 1.2 What is captured by an edge?**

**51** At its core, links within food webs can be thought of as a representation of either feeding links between species  
**52** - be that realised (Pringle, 2020) or potential (Dunne, 2006), alternative links can represent fluxes within the  
**53** system *e.g.*, energy transfer or material flow as the result of the feeding links between species (Lindeman,  
**54** 1942). Fundamentally this means that the links within a network represent different ‘currencies’ (either the  
**55** feasibility of a link existing between two species or the energy that is moving through the system) and how the  
**56** links within a network are specified will influence the resulting structure of the network. For example taking  
**57** a food web that consists of links representing all *potential* feeding links for a community (*i.e.*, a metaweb)  
**58** will be meaningless if one is interested in understanding the flow of energy through the network as the links  
**59** within a metaweb do not represent environmental/energetic constraints, making them poor representations

of which interactions are *realised* in a specific location (Caron et al., 2024). In addition to the various ways of defining the links between species pairs there are also a myriad of ways in which the links themselves can be quantified. Links between species are often treated as being present or absent (*i.e.*, binary) but it is also possible to use probabilities (Banville et al., 2024; which quantifies how likely an interaction is to occur, Poisot, Cirtwill, et al., 2016) or continuous measurements (which quantifies the strength of an interaction, Berlow et al., 2004).

### 1.3 Network representations

Broadly, networks can be thought of to fall into two different ‘types’; namely metawebs; traditionally defined as all of the *potential* interactions for a specific species pool (Dunne, 2006), and realised networks; which is the subset of interactions in a metaweb that are *realised* for a specific community at a given time and place. The fundamental difference between these two different network representations is that a metaweb provides insight as to the viability of an interaction between two species occurring and is a means to identify links that are not ecologically plausible, *i.e.*, forbidden links (Jordano, 2016b), or provide an idea of the *complete* diet of a species (Strydom et al., 2023). In contrast realised networks are highly localised and links between species are contingent on both the co-occurrence of species as well as the influence of the environment, and population and community dynamics on predator choice. In the context of definitions and semantics the links that are represented by a metaweb and a realised network are different; links that are absent in a metaweb can be treated as being truly absent, however links that are absent in a realised network cannot be considered to be truly absent but rather as absent due to the broader environmental/community context. Importantly, a realised network is *not* simply the downscaling of a metaweb to a smaller scale (*e.g.*, moving from the country to the 1x1 km<sup>2</sup> scale based on fine-scale species co-occurrence) but represents a shift towards capturing the higher level processes that determine the *realisation* of an interaction, *i.e.*, the definition of an edges shifts from being determined by interaction feasibility to that of energetic choices/consequences. Thus, different network representations are determined and constrained by different sets of assumptions as to what the processes are that determine the presence/absence of an interaction between two species as well as the resulting network structure.

## 2 From Nodes and Edges to Scale, Context, and Process

The interplay between network representation and network (node and edge) definition is primarily governed by the process(es) that determine the interaction between species, however these processes are also scale and context dependent. Here we start by introducing the five core processes that determine either the feasibility

90 or the realisation of interactions, namely: evolutionary compatibility, co-occurrence, abundance, predator  
91 choice, and non-trophic interactions; while simultaneously contextualising them within, and linking them  
92 to the different network representations Figure 1. We can think of the different network representations  
93 to be conceptually analogous to the fundamental and realised niche, whereby the metaweb represents the  
94 ‘fundamental diet niche’ of a species and a realised network represents the ‘realised diet’ of a species. Of  
95 course these processes do not function in a vacuum and do interact with/influence one another, but it is still  
96 beneficial to present them in a categorical manner as these different processes are often the underpinning  
97 logic in the development of prediction/network models, the criteria for data collection in the field, and the  
98 scale of organisation for which they are relevant (species, population, or community).

99 [Figure 1 about here.]

## 100 2.1 The processes that determine species interactions

### 101 Evolutionary compatibility

102 There is compelling evidence that an interaction occurring between two species is the result of their shared  
103 (co)evolutionary history (Dalla Riva & Stouffer, 2016; Gómez et al., 2010; Segar et al., 2020) which, in the  
104 more proximal sense, is manifested as the ‘trait complementarity’ between two species (Benadi et al., 2022),  
105 whereby one species (the predator) has the ‘correct’ set of traits that allow it to chase, capture, kill, and  
106 consume the other species (the prey). For species pairs where this condition is not met the link is deemed  
107 to be forbidden (Jordano, 2016b); *i.e.*, not physically possible and will always be absent within a network.  
108 A network constructed on the basis of evolutionary compatible links is most closely aligned with a metaweb,  
109 although it would not be required that the species co-occur (as shown in Figure 1), and arguably makes for  
110 a good approximation of the ‘Eltonian niche’ of species (Soberón, 2007). Finally, one should be aware that  
111 it is possible to represent evolutionary compatible interactions as either binary (possible vs forbidden) or as  
112 a probability (Banville et al., 2024), where the probability represents how likely the interaction between two  
113 species is to be possible.

### 114 (Co)occurrence

115 Although the outright assumption that because two species are co-occurring it must mean that they are  
116 interacting is flawed (Blanchet et al., 2020), it is of course impossible for two species to interact (at least in  
117 terms of feeding links) if they are not co-occurring in time and space. Thus, although co-occurrence data  
118 alone is insufficient to build an accurate and ecologically meaningful representation of *feeding links* it is still  
119 a critical process that determines the realisation of feeding links and allows us to constrain a global metaweb

120 to only consider ‘realised’ communities (Dansereau et al., 2024) and an understanding of the intersection of  
121 species interactions and their co-occurrence is meaningful when one is operating in the space of trying to  
122 determine the distribution of a species (Higino et al., 2023; Pollock et al., 2014), representing something of  
123 a fusion of the the Grinnellian and Eltonian niches (Gravel et al., 2019).

124 **Abundance**

125 The abundance of different the species within the community is thought to influence the realisation of feeding  
126 links primarily in two ways. Firstly there is the argument that that structure of networks (and the interactions  
127 that they are composed of) are driven *only* by the abundance of the different species and that interactions  
128 are not contingent on there being any compatibility (trait matching) between them, *sensu* neutral processes  
129 (Canard et al., 2012; Momal et al., 2020). However, a more ecologically sound assumption would be that  
130 the abundance of different prey species will influence the distribution of links in a network (Vázquez et al.,  
131 2009), be influencing which prey are targeted or preferred by the predator as abundance influences factors  
132 such as the likelihood of two species (individuals) meeting (Banville et al., 2024; Poisot et al., 2015), or in  
133 the dynamic sense will influence the persistence of viable populations.

134 **Profitability (predator choice)**

135 Ultimately, predator choice is underpinned by the energetic cost-benefit (profitability) of trying to catch, kill,  
136 and consume prey (where a predator will optimise energy while minimising handling and search time), and  
137 is well described within both optimal foraging (Pyke, 1984) and metabolic theory (Brown et al., 2004). The  
138 energetic cost of feeding is itself can be deconstructed as the energy content as well as the density (abundance)  
139 of prey (as this influences search time) and how these will influence which links are realised Figure 1, with an  
140 argument that body size represents a key trait that may capture and influence these processes (White et al.,  
141 2007; Yodzis & Innes, 1992). Additional work on on understanding the energetic cost that the environment  
142 imposes on an individual (Cherif et al., 2024) as well as the way a predator uses the landscape to search for  
143 prey (Pawar et al., 2012) is bringing us closer to accounting for the energetic cost of realising feeding links.

144 **Non-trophic interactions**

145 Perhaps not as intuitive when thinking about the processes that determine feeding links (trophic interac-  
146 tions) is thinking about the role of the ability of non-trophic interactions to modify either the realisation or  
147 strength of trophic interactions (Golubski & Abrams, 2011; Pilosof et al., 2017). Non-trophic interactions  
148 can modify interactions either ‘directly’ e.g., predator *a* outcompetes predator *b* or ‘indirectly’ e.g., mutu-  
149 alistic/facilitative interactions will alter the fine-scale distribution and abundance of species as well as their  
150 persistence (Buche et al., 2024; Kéfi et al., 2012, 2015). The ‘unobservable’ nature of non-trophic interactions

151 makes them a challenge to quantify, however their importance in network dynamics (Staniczenko et al., 2010)  
152 as well as cascading effects [e.g., Kamaru et al. (2024)] should not be overlooked.

## 153 2.2 Contextualising the processes that determine species interactions

154 It should be self evident that the different processes discussed above are all ultimately going to influence the  
155 realisation of interactions as well as the structure of a network, however they are acting at different scales  
156 of organisation. Both the **co-occurrence** and the **evolutionary compatibility** are valid at the scale of  
157 the species pair of interest, that is the *possibility* of an interaction being present/absent is assessed at the  
158 pairwise level and one is left with a ‘list’ of interactions that are present/absent. Although it is possible  
159 to build a network (*i.e.*, metaweb) from this information it is important to be aware that the structure of  
160 this network is not constrained by real-world dynamics or conditions, and so just because species are able to  
161 interact does not mean that they will (Poisot et al., 2015). In order to construct a network who’s structure is  
162 a closer approximation of reality (localised interactions) one needs to take into consideration the properties  
163 of the community as a whole and information about the individuals it is comprised of (Quintero et al., 2024),  
164 which requires more data at the community scale, such as the abundance of species.

## 165 3 Network construction is nuanced

166 The act of constructing a ‘real world’ network will ultimately be delimited by its intended use, however the  
167 reality is that the empirical collection of interaction data is both costly and challenging to execute (Jordano,  
168 2016a, 2016b), especially if one wants to capture *all* aspects of the processes discussed in Section 2 (owing  
169 to the different time and spatial scales they may be operating at). Thus we often turn to models to either  
170 predict networks (be that the interaction between two species, or network structure (Strydom et al., 2021)),  
171 or as a means to identify missing interactions (gap fill) within an existing empirical dataset (Biton et al.,  
172 2024; Dallas et al., 2017; Stock, 2021), and so for the purpose of this discussion network construction will be  
173 synonymous with using a model as a means to represent or predict a network. That is not to say that there is  
174 no need for empirical data collection but rather that using a model for food web prediction (or reconstruction)  
175 is a more feasible approach as it allows us to make inferences about interactions that are not happening in  
176 the ‘observable now’ (Strydom et al., 2021), with the added benefit that one is able to build some uncertainty  
177 into the resulting network (Banville et al., 2024). Additionally different models have different underlying  
178 philosophies that allow us to capture one or a few of the processes discussed in Section 2, and although the  
179 delimits and defines what inferences can be made from the resulting network it also allows us to isolate and  
180 understand how different processes determine interactions (Song & Levine, 2024; Stouffer, 2019). Here we will

introduce the three different types of network representations (metawebs, realised networks, and structural networks), how they link back to (and encode) the different processes determining interactions Figure 1, and broadly discuss some of the modelling approaches that are used to construct these different network types. This is paralleled by a hypothetical case study (Box 1) where we showcase the utility/applicability of the different network representations in the context of trying to understand the feeding dynamics of a seasonal community.

**i** Box 1 - Why we need to aggregate networks at different scales: A hypothetical case study

note I am using a figure for layout experimentation purposes

Although it might seem most prudent to be predicting, constructing, and defining networks that are the closest representation of reality there are pros and cons of constructing both realised networks as well as metawebs. Let us take for example a community that experiences a degree of species turnover between seasons. In this community we expect species to be either present or absent depending on the season (*i.e.*, changes in co-occurrence) as well as some species exhibiting seasonal shifts in their diets (be that due to changes in species occurrence or predator choice). If one were to construct a metaweb that disregards these season shifts ('global metaweb') these details would be lost and it would be valuable to construct either smaller metawebs for the different seasonal communities (thereby capturing the changes in community diversity), or realised networks for each season (to capture diet or ecosystem process shifts). However, these small-scale networks lack the context of the bigger picture that is available at the metaweb - that is it gives us a more holistic idea of the entire diet range of a specific species, which is important when one needs to make conservation-based/applied decisions (*e.g.*, conserving the entire diet of a species and not just seasonal prey items) as well as providing information on interactions that may be possible regardless of the environmental/community context (species may have the capacity to consume certain prey items but do not do so due to local conditions). With this in mind let us see how the different network aggregations can be used

[Figure 2 about here.]

187

### 188 3.1 Models that predict metawebs (feasible interactions)

189 This is perhaps the most developed group of models; with a variety of approaches having been developed  
190 that typically determine the feasibility of an interaction using the trait compatibility between predator and  
191 prey (*i.e.* their evolutionary compatibility) to determine 'feeding rules' (Morales-Castilla et al., 2015). These  
192 feeding rules are broadly elucidated in two different ways; mechanistic feeding rules can be explicitly defined

and applied to a community (Dunne et al., 2008; Roopnarine, 2017; *e.g.*, Shaw et al., 2024) or they are inferred from a community for which there are interaction data and the ‘rules’ are then applied to a different community (Caron et al., 2022; Cirtwill et al., 2019; Desjardins-Proulx et al., 2017; Eklöf et al., 2013; Llewelyn et al., 2023; Pichler et al., 2020; Strydom et al., 2022; *e.g.*, Strydom et al., 2023). The fundamental difference between these two model groups is that ‘mechanistic models’ rely on expert knowledge and make explicit assumptions on trait-feeding relationships, whereas the ‘pattern finding’ models are dependent on existing datasets from which to elucidate feeding rules. These models are useful for determining all feasible interactions for a specific community, and owing to the availability of empirical interaction datasets (Gray et al., 2015; *e.g.*, Poelen et al., 2014; Poisot, Baiser, et al., 2016), as well as the development of model testing/benchmarking tools (Poisot, 2023), means that these models can be validated and (with relative confidence) be used to construct first draft networks for communities for which we have no interaction data (Strydom et al., 2022), and are valuable not only in data poor regions but also for predicting interactions for ‘unobservable’ communities *e.g.*, prehistoric networks (Fricke et al., 2022; Yeakel et al., 2014) or future, novel community assemblages. Importantly metawebs are inherently ‘static’ in the sense that they are *not* able to capture dynamic processes (since the notion of feasibility is all or nothing), however they provide a bigger picture context (*e.g.*, understanding the *entire* diet breadth of a species) and often require little data to construct.

### 3.2 Models that predict realised networks (realised interactions)

In order to construct realised networks models need to incorporate *both* the feasibility of interactions (*i.e.*, determine the entire diet breadth of a species) as well as then determine which interactions are realised (*i.e.*, incorporate the ‘cost’ of interactions). As far as we are aware there is no model that explicitly accounts for both of these ‘rules’ (although see Olivier et al. (2019)) and rather *only* account for processes that determine the realisation of an interaction (*i.e.*, abundance, predator choice, or non-trophic interactions). Although the use of allometry *i.e.*, body size (Beckerman et al., 2006; *e.g.*, Valdovinos et al., 2023) may represent a first step in capturing ‘evolutionary compatibility’ alongside more energy (predator choice) driven processes we still need to account for other traits that determine feeding compatibility (*e.g.*, Van De Walle et al., 2023 show how incorporating prey defensive properties alongside body size improves predictions). In terms of constructing realised networks, diet models (Beckerman et al., 2006; Petchey et al., 2008) have been used to construct networks based on both predator choice (as determined by the handling time, energy content, and predator attack rate) as well as abundance (prey density) and progress has also been made in understanding the compartmentation of energy in networks and how this influences energy acquisition (Krause et al., 2003;

224 Wootton et al., 2023). As realised networks are build on the concept of dynamic processes (the abundance  
225 of species will always be in flux) these networks are valuable for understanding the behaviour of networks  
226 over time or their response to change (Curtsdotter et al., 2019; Delmas et al., 2017; Lajaaiti et al., 2024).  
227 However, they are ‘costly’ to construct (requiring data about the entire community as it is the behaviour of  
228 the system that determines the behaviour of the part) and also lack the larger diet niche context afforded by  
229 metawebs.

### 230 3.3 Models that predict structure (interaction agnostic)

231 Although we identify mechanisms that determine species interactions in Section 2 not all models that are used  
232 to predict networks explicitly operate at the ‘process’ level, but rather represent the *structure* of a network  
233 based on a series of *a priori* assumptions as to the distribution of links between species (typically trophic  
234 not taxonomic species). These models operate by parametrising an aspect of the network structure, (*e.g.*,  
235 the niche model (Williams & Martinez, 2000) makes an assumption as to the expected connectance of the  
236 network, although see Allesina & Pascual (2009) for a parameter-free model) or alternatively uses structural  
237 features of an exiting *realised* network (*e.g.*, stochastic block model, Xie et al. (2017)). Importantly these  
238 structural models do not make species specific predictions (they are usually species agnostic and treat nodes  
239 as trophic species) and so cannot be used to determine if an interaction is either possible *or* realised between  
240 two species (*i.e.*, one cannot use these models to determine if species *a* eats species *b*). Although this means  
241 this suite of models are unsuitable as tools for predicting species-specific interactions, they have been shown  
242 to be sufficient tools to predict the structure of networks (Williams & Martinez, 2008), and provide a data-  
243 light (the models often only require species richness) but assumption heavy (the resulting network structure  
244 is determined by an assumption of network structure) way to construct a network.

## 245 4 Making Progress with Networks

### 246 4.1 Further development of models and tools

247 There has been a suite of models that have been developed to predict feeding links, however we are lacking  
248 in tools that are explicitly taking into consideration estimating both the feasibility as well as realisation of  
249 links, *i.e.*, both interactions and structure simultaneously (Strydom et al., 2021). This could be addressed  
250 either through the development of tools that do both (predict both interactions and structure), or to develop  
251 an ensemble modelling approach (Becker et al., 2022; Terry & Lewis, 2020) or tools that will allow for the  
252 downsampling of metawebs into realised networks (*e.g.*, Roopnarine, 2006). Additionally although realised

networks are more closely aligned with capturing interaction strength we lack models that allow us to quantify this (Strydom et al., 2021; Wells & O'Hara, 2013). In addition to the more intentional development of models we also need to consider the validation of these models, there have been developments and discussions for assessing how well a model recovers pairwise interactions (Poisot, 2023; Strydom et al., 2021), although the rate of false-negatives that may be present in the testing data still present a challenge (Catchen et al., 2023), and we still lack clear set of guidelines for benchmarking the ability of models to recover structure (Allesina et al., 2008).

## 4.2 At what scale should we be predicting and using networks?

We lack an understanding of which processes drive the differences between different scales (Saravia et al., 2022), as well as to what the appropriate level of aggregation is for a ‘network’ (Estay et al., 2023). Which presents a challenge both in deciding what the appropriate spatial and time scales are for constructing not only a network but also which type of network representation. Space influences both network properties (Galiana et al., 2018), as well as dynamics (Fortin et al., 2021; Rooney et al., 2008), and time has implications when it comes to accounting for seasonal turnover in communities (Brimacombe et al., 2021; Laender et al., 2010) as well as thinking co-occurrence, particularly the records used to determine it (Brimacombe et al., 2024). Although multilayer networks may allow us to encode the nuances of space and time (Hutchinson et al., 2019) we still need to understand the implications of *e.g.*, constructing networks that are not at ecologically but rather politically relevant scales (Strydom et al., 2022) and what we can learn or infer from networks at these scales.

## 5 The future value of networks

developing a dictionary of use... that helps navigate between the levels and assumptions

It should be clear that there is a high degree of interrelatedness and overlap between the way a network is constructed (modelled or predicted) and the process(es) it captures, these are encoded (embedded) within the network representation and ultimately influences how the network can and should be used (Berlow et al., 2008; Petchey et al., 2011), with different network representations yielding different interpretations of processes (Keyes et al., 2024). It is probably both this nuance as well as a lack of clear boundaries and guidelines as to the links between network form and function (although see Delmas et al., 2019) that has stifled the ‘productive use’ of networks beyond inventorying the interactions between species. Although, progress with using networks as a means to address questions within larger bodies of ecological theory *e.g.*,

<sup>282</sup> invasion biology (Hui & Richardson, 2019) and co-existence theory (García-Callejas et al., 2023), has been  
<sup>283</sup> made we still need to have a discussion on what the appropriate network representation for the task at hand  
<sup>284</sup> would be. This is highlighted in Box 1, and underscores that we need to evaluate exactly what process a  
<sup>285</sup> specific network representation captures as well as its suitability for the question of interest.

<sup>286</sup> **5.1 How will novel communities interact?**

<sup>287</sup> Here we can talk about the effects of range shifts and invasions and how this will result in new/novel  
<sup>288</sup> community assemblages. And then also the intentional changes of species compositions through rewilding.

<sup>289</sup> **5.2 How will changes in the community influence ecosystem processes?**

<sup>290</sup> Linking to dynamic networks and how this lets us build spatially/temporally explicit networks which can be  
<sup>291</sup> used to infer form and function. Also bring in the discussion on the suitable aggregation (and the fact that  
<sup>292</sup> we don't know)

<sup>293</sup> **5.3 How do species persist/co-exist?**

<sup>294</sup> Specific sub points to consider here is persistence, especially persistence to perturbation. Again, dynamic  
<sup>295</sup> networks and network/community assembly and finally extinctions (Dunhill et al., 2024).

<sup>296</sup> [Figure 3 about here.]

Table 1: An informative table

Question (broad)	Question (specific)	Network representation
Species invasions	What species will the invading species interact with?	Regional metaweb but need to derive information from a global metaweb since these are interactions that are 'novel'
Species invasions	How does the invading species alter network dynamics and function?	Realised network (after having moved through the global metaweb to understand which interactions are feasible)
Range shifts and novel communities	Under global change how will novel community assemblages interact?	Global metaweb, need context of broader community

Question (broad)	Question (specific)	Network representation
Extinctions	Cascading effect of the loss of a species from the network	Regional metaweb - need to account for entire diet, a realised network will exclude the entire diet but will allow to elucidate the final structure
Species/community persistence	Dynamics over time. Stability/resilience. How does a change in pop A affect pop B?	Realised networks - but dynamic!
Synthetic networks	Creating ecologically plausible communities for synthetic analyses	Structural networks - data light!
Practical use	What is both attainable (data constraints) but also of practical use to 'real world' decision making. So moving from theory to applied	??Regional metawebs??

## 297 References

- 298 Allesina, S., Alonso, D., & Pascual, M. (2008). A General Model for Food Web Structure. *Science*, 320(5876),  
 299 658–661. <https://doi.org/10.1126/science.1156269>
- 300 Allesina, S., & Pascual, M. (2009). Food web models: A plea for groups. *Ecology Letters*, 12(7), 652–662.  
 301 <https://doi.org/10.1111/j.1461-0248.2009.01321.x>
- 302 Banville, F., Strydom, T., Blyth, P., Brimacombe, C., Catchen, M. D., Dansereau, G., Higino, G., Malpas,  
 303 T., Mayall, H., Norman, K., Gravel, D., & Poisot, T. (2024). *Deciphering probabilistic species interaction*  
 304 *networks*. EcoEvoRxiv. <https://doi.org/10.32942/X28G8Z>
- 305 Becker, D. J., Albery, G. F., Sjodin, A. R., Poisot, T., Bergner, L. M., Chen, B., Cohen, L. E., Dallas, T.  
 306 A., Eskew, E. A., Fagre, A. C., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Stock, M., Teeling,  
 307 E. C., & Carlson, C. J. (2022). Optimising predictive models to prioritise viral discovery in zoonotic  
 308 reservoirs. *The Lancet Microbe*, 3(8), e625–e637. [https://doi.org/10.1016/S2666-5247\(21\)00245-7](https://doi.org/10.1016/S2666-5247(21)00245-7)
- 309 Beckerman, A. P., Petchey, O. L., & Warren, P. H. (2006). Foraging biology predicts food web complexity.  
 310 *Proceedings of the National Academy of Sciences*, 103(37), 13745–13749. <https://doi.org/10.1073/pnas.0603039103>
- 312 Benadi, G., Dormann, C. F., Fründ, J., Stephan, R., & Vázquez, D. P. (2022). Quantitative Prediction of

- 313      Interactions in Bipartite Networks Based on Traits, Abundances, and Phylogeny. *The American Naturalist*,  
314      199(6), 841–854. <https://doi.org/10.1086/714420>
- 315      Berlow, E. L., Brose, U., & Martinez, N. D. (2008). The “Goldilocks factor” in food webs. *Proceedings of  
316      the National Academy of Sciences*, 105(11), 4079–4080. <https://doi.org/10.1073/pnas.0800967105>
- 317      Berlow, E. L., Neutel, A.-M., Cohen, J. E., de Ruiter, P. C., Ebenman, B., Emmerson, M., Fox, J. W., Jansen,  
318      V. A. A., Iwan Jones, J., Kokkoris, G. D., Logofet, D. O., McKane, A. J., Montoya, J. M., & Petchey, O.  
319      (2004). Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73(3),  
320      585–598. <https://doi.org/10.1111/j.0021-8790.2004.00833.x>
- 321      Biton, B., Puzis, R., & Pilosof, S. (2024). *Inductive link prediction boosts data availability and enables  
322      cross-community link prediction in ecological networks*.
- 323      Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions.  
324      *Ecology Letters*, 23(7), 1050–1063. <https://doi.org/10.1111/ele.13525>
- 325      Brimacombe, C., Bodner, K., & Fortin, M.-J. (2021). Inferred seasonal interaction rewiring of a freshwater  
326      stream fish network. *Ecography*, 44(2), 219–230. <https://doi.org/10.1111/ecog.05452>
- 327      Brimacombe, C., Bodner, K., & Fortin, M.-J. (2024). *Applying a method before its proof-of-concept: A  
328      cautionary tale using inferred food webs*. <https://doi.org/10.13140/RG.2.2.22076.65927>
- 329      Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings  
330      of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4),  
331      e3002068. <https://doi.org/10.1371/journal.pbio.3002068>
- 332      Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a Metabolic Theory  
333      of Ecology. *Ecology*, 85(7), 1771–1789. <https://doi.org/10.1890/03-9000>
- 334      Buche, L., Bartomeus, I., & Godoy, O. (2024). Multitrophic Higher-Order Interactions Modulate Species  
335      Persistence. *The American Naturalist*, 203(4), 458–472. <https://doi.org/10.1086/729222>
- 336      Canard, E., Mouquet, N., Marescot, L., Gaston, K. J., Gravel, D., & Mouillot, D. (2012). Emergence of  
337      Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7(8), e38295. <https://doi.org/10.1371/journal.pone.0038295>
- 338      Caron, D., Brose, U., Lurgi, M., Blanchet, F. G., Gravel, D., & Pollock, L. J. (2024). Trait-matching models  
339      predict pairwise interactions across regions, not food web properties. *Global Ecology and Biogeography*,  
340      33(4), e13807. <https://doi.org/10.1111/geb.13807>
- 341      Caron, D., Maiorano, L., Thuiller, W., & Pollock, L. J. (2022). Addressing the Eltonian shortfall with  
342      trait-based interaction models. *Ecology Letters*, 25(4), 889–899. <https://doi.org/10.1111/ele.13966>
- 343      Catchen, M. D., Poisot, T., Pollock, L. J., & Gonzalez, A. (2023). *The missing link: Discerning true from  
344      false negatives when sampling species interaction networks*.

- 346 Cherif, M., Brose, U., Hirt, M. R., Ryser, R., Silve, V., Albert, G., Arnott, R., Berti, E., Cirtwill, A.,  
347 Dyer, A., Gauzens, B., Gupta, A., Ho, H.-C., Portalier, S. M. J., Wain, D., & Wootton, K. (2024). The  
348 environment to the rescue: Can physics help predict predator–prey interactions? *Biological Reviews*,  
349 n/a(n/a). <https://doi.org/10.1111/brv.13105>
- 350 Cirtwill, A. R., Ekklf, A., Roslin, T., Wootton, K., & Gravel, D. (2019). A quantitative framework for  
351 investigating the reliability of empirical network construction. *Methods in Ecology and Evolution*, 10(6),  
352 902–911. <https://doi.org/10.1111/2041-210X.13180>
- 353 Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on food web structure.  
354 *Ecology*, 99(12), 2712–2720. <https://doi.org/10.1002/ecy.2523>
- 355 Curtsdotter, A., Banks, H. T., Banks, J. E., Jonsson, M., Jonsson, T., Laubmeier, A. N., Traugott, M., &  
356 Bommarco, R. (2019). Ecosystem function in predator–prey food webs—confronting dynamic models with  
357 empirical data. *Journal of Animal Ecology*, 88(2), 196–210. <https://doi.org/10.1111/1365-2656.12892>
- 358 Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food webs' backbones  
359 using functional traits. *Oikos*, 125(4), 446–456. <https://doi.org/10.1111/oik.02305>
- 360 Dallas, T., Park, A. W., & Drake, J. M. (2017). Predicting cryptic links in host-parasite networks. *PLOS  
361 Computational Biology*, 13(5), e1005557. <https://doi.org/10.1371/journal.pcbi.1005557>
- 362 Dansereau, G., Barros, C., & Poisot, T. (2024). Spatially explicit predictions of food web structure from  
363 regional-level data. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 379(1909).  
364 <https://doi.org/10.1098/rstb.2023.0166>
- 365 Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Riva, G. V. D., Fortin, M.-J., Gravel, D., Guimarães,  
366 P. R., Hembry, D. H., Newman, E. A., Olesen, J. M., Pires, M. M., Yeakel, J. D., & Poisot, T. (2019).  
367 Analysing ecological networks of species interactions. *Biological Reviews*, 94(1), 16–36. <https://doi.org/10.1111/brv.12433>
- 368 Delmas, E., Brose, U., Gravel, D., Stouffer, D. B., & Poisot, T. (2017). Simulations of biomass dynamics in  
369 community food webs. *Methods in Ecology and Evolution*, 8(7), 881–886. [https://doi.org/10.1111/2041-210X.12713](https://doi.org/10.1111/2041-<br/>370 210X.12713)
- 371 Desjardins-Proulx, P., Laigle, I., Poisot, T., & Gravel, D. (2017). Ecological interactions and the Netflix  
372 problem. *PeerJ*, 5, e3644. <https://doi.org/10.7717/peerj.3644>
- 373 Dunhill, A. M., Zarzyczny, K., Shaw, J. O., Atkinson, J. W., Little, C. T. S., & Beckerman, A. P. (2024).  
374 Extinction cascades, community collapse, and recovery across a Mesozoic hyperthermal event. *Nature  
375 Communications*, 15(1), 8599. <https://doi.org/10.1038/s41467-024-53000-2>
- 376 Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological  
377 networks: Linking structure and dynamics* (pp. 27–86). Oxford University Press.
- 378

- 379 Dunne, J. A., Williams, R. J., Martinez, N. D., Wood, R. A., & Erwin, D. H. (2008). Compilation and  
380 Network Analyses of Cambrian Food Webs. *PLOS Biology*, 6(4), e102. <https://doi.org/10.1371/journal.pbio.0060102>
- 382 Eklöf, A., Tang, S., & Allesina, S. (2013). Secondary extinctions in food webs: A Bayesian network approach.  
383 *Methods in Ecology and Evolution*, 4(8), 760–770. <https://doi.org/10.1111/2041-210X.12062>
- 384 Estay, S. A., Fortin, M.-J., & López, D. N. (2023). Editorial: Patterns and processes in ecological networks  
385 over space. *Frontiers in Ecology and Evolution*, 11.
- 386 Fortin, M.-J., Dale, M. R. T., & Brimacombe, C. (2021). Network ecology in dynamic landscapes. *Proceedings  
387 of the Royal Society B: Biological Sciences*, 288(1949), rspb.2020.1889, 20201889. <https://doi.org/10.1098/rspb.2020.1889>
- 389 Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning,  
390 J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene.  
391 *Science*, 377(6609), 1008–1011. <https://doi.org/10.1126/science.abn4012>
- 392 Galiana, N., Lurgi, M., Claramunt-López, B., Fortin, M.-J., Leroux, S., Cazelles, K., Gravel, D., & Montoya,  
393 J. M. (2018). The spatial scaling of species interaction networks. *Nature Ecology & Evolution*, 2(5),  
394 782–790. <https://doi.org/10.1038/s41559-018-0517-3>
- 395 García-Callejas, D., Godoy, O., Buche, L., Hurtado, M., Lanuza, J. B., Allen-Perkins, A., & Bartomeus, I.  
396 (2023). Non-random interactions within and across guilds shape the potential to coexist in multi-trophic  
397 ecological communities. *Ecology Letters*, 26(6), 831–842. <https://doi.org/10.1111/ele.14206>
- 398 Golubski, A. J., & Abrams, P. A. (2011). Modifying modifiers: What happens when interspecific interactions  
399 interact? *Journal of Animal Ecology*, 80(5), 1097–1108. <https://doi.org/10.1111/j.1365-2656.2011.01852.x>
- 401 Gómez, J. M., Verdú, M., & Perfectti, F. (2010). Ecological interactions are evolutionarily conserved across  
402 the entire tree of life. *Nature*, 465(7300), 918–921. <https://doi.org/10.1038/nature09113>
- 403 Gravel, D., Baiser, B., Dunne, J. A., Kopelke, J.-P., Martinez, N. D., Nyman, T., Poisot, T., Stouffer, D. B.,  
404 Tylianakis, J. M., Wood, S. A., & Roslin, T. (2019). Bringing Elton and Grinnell together: A quantitative  
405 framework to represent the biogeography of ecological interaction networks. *Ecography*, 42(3), 401–415.  
406 <https://doi.org/10.1111/ecog.04006>
- 407 Gray, C., Figueroa, D. H., Hudson, L. N., Ma, A., Perkins, D., & Woodward, G. (2015). Joining the dots:  
408 An automated method for constructing food webs from compendia of published interactions. *Food Webs*,  
409 5, 11–20. <https://doi.org/10.1016/j.fooweb.2015.09.001>
- 410 Higino, G. T., Banville, F., Dansereau, G., Muñoz, N. R. F., Windsor, F., & Poisot, T. (2023). Mismatch  
411 between IUCN range maps and species interactions data illustrated using the Serengeti food web. *PeerJ*,

- 412 11, e14620. <https://doi.org/10.7717/peerj.14620>
- 413 Hui, C., & Richardson, D. M. (2019). How to Invade an Ecological Network. *Trends in Ecology & Evolution*,  
414 34(2), 121–131. <https://doi.org/10.1016/j.tree.2018.11.003>
- 415 Hutchinson, M. C., Bramon Mora, B., Pilosof, S., Barner, A. K., Kéfi, S., Thébault, E., Jordano, P., &  
416 Stouffer, D. B. (2019). Seeing the forest for the trees: Putting multilayer networks to work for community  
417 ecology. *Functional Ecology*, 33(2), 206–217. <https://doi.org/10.1111/1365-2435.13237>
- 418 Jordano, P. (2016a). Chasing Ecological Interactions. *PLOS Biology*, 14(9), e1002559. <https://doi.org/10.1371/journal.pbio.1002559>
- 420 Jordano, P. (2016b). Sampling networks of ecological interactions. *Functional Ecology*. <https://doi.org/10.1111/1365-2435.12763>
- 422 Kamaru, D. N., Palmer, T. M., Riginos, C., Ford, A. T., Belnap, J., Chira, R. M., Githaiga, J. M., Gituku,  
423 B. C., Hays, B. R., Kavwele, C. M., Kibungei, A. K., Lamb, C. T., Maiyo, N. J., Milligan, P. D., Mutisya,  
424 S., Ng’weno, C. C., Ongutu, M., Pietrek, A. G., Wildt, B. T., & Goheen, J. R. (2024). Disruption of  
425 an ant-plant mutualism shapes interactions between lions and their primary prey. *Science*, 383(6681),  
426 433–438. <https://doi.org/10.1126/science.adg1464>
- 427 Kéfi, S., Berlow, E. L., Wieters, E. A., Joppa, L. N., Wood, S. A., Brose, U., & Navarrete, S. A. (2015).  
428 Network structure beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky  
429 shores. *Ecology*, 96(1), 291–303. <https://doi.org/10.1890/13-1424.1>
- 430 Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A., Boit, A., Joppa, L. N.,  
431 Lafferty, K. D., Williams, R. J., Martinez, N. D., Menge, B. A., Blanchette, C. A., Iles, A. C., & Brose,  
432 U. (2012). More than a meal... integrating non-feeding interactions into food webs: More than a meal ....  
433 *Ecology Letters*, 15(4), 291–300. <https://doi.org/10.1111/j.1461-0248.2011.01732.x>
- 434 Keyes, A. A., Barner, A. K., & Dee, L. E. (2024). Synthesising the Relationships Between Food Web Structure  
435 and Robustness. *Ecology Letters*, 27(10), e14533. <https://doi.org/10.1111/ele.14533>
- 436 Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E., & Taylor, W. W. (2003). Compartments  
437 revealed in food-web structure. *Nature*, 426(6964), 282–285. <https://doi.org/10.1038/nature02115>
- 438 Laender, F. D., Oevelen, D. V., Soetaert, K., & Middelburg, J. J. (2010). Carbon transfer in a herbivore-  
439 and microbial loop-dominated pelagic food webs in the southern Barents Sea during spring and summer.  
440 *Marine Ecology Progress Series*, 398, 93–107. <https://doi.org/10.3354/meps08335>
- 441 Lajaaiti, I., Bonnici, I., Kéfi, S., Mayall, H., Danet, A., Beckerman, A. P., Malpas, T., & Delmas, E. (2024).  
442 *EcologicalNetworksDynamics.jl* A Julia package to simulate the temporal dynamics of complex ecological  
443 networks (p. 2024.03.20.585899). bioRxiv. <https://doi.org/10.1101/2024.03.20.585899>
- 444 Lindeman, R. L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23(4), 399–417. <https://doi.org/>




- 445 10.2307/1930126
- 446 Llewelyn, J., Strona, G., Dickman, C. R., Greenville, A. C., Wardle, G. M., Lee, M. S. Y., Doherty, S.,  
447 Shabani, F., Saltré, F., & Bradshaw, C. J. A. (2023). Predicting predator–prey interactions in terrestrial  
448 endotherms using random forest. *Ecography*, 2023(9), e06619. <https://doi.org/10.1111/ecog.06619>
- 449 Momal, R., Robin, S., & Ambroise, C. (2020). Tree-based inference of species interaction networks from  
450 abundance data. *Methods in Ecology and Evolution*, 11(5), 621–632. <https://doi.org/10.1111/2041-210X.13380>
- 451 13380
- 452 Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from  
453 proxies. *Trends in Ecology & Evolution*, 30(6), 347–356. <https://doi.org/10.1016/j.tree.2015.03.014>
- 454 Olivier, P., Frelat, R., Bonsdorff, E., Kortsch, S., Kröncke, I., Möllmann, C., Neumann, H., Sell, A. F., &  
455 Nordström, M. C. (2019). Exploring the temporal variability of a food web using long-term biomonitoring  
456 data. *Ecography*, 42(12), 2107–2121. <https://doi.org/10.1111/ecog.04461>
- 457 Pawar, S., Dell, A. I., & Savage, V. M. (2012). Dimensionality of consumer search space drives trophic  
458 interaction strengths. *Nature*, 486(7404), 485–489. <https://doi.org/10.1038/nature11131>
- 459 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2008). Size, foraging, and food web  
460 structure. *Proceedings of the National Academy of Sciences*, 105(11), 4191–4196. <https://doi.org/10.1073/pnas.0710672105>
- 461
- 462 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2011). Fit, efficiency, and biology: Some  
463 thoughts on judging food web models. *Journal of Theoretical Biology*, 279(1), 169–171. <https://doi.org/10.1016/j.jtbi.2011.03.019>
- 464
- 465 Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine learning algorithms  
466 to infer trait-matching and predict species interactions in ecological networks. *Methods in Ecology and  
467 Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>
- 468 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological networks.  
469 *Nature Ecology & Evolution*, 1(4), 101. <https://doi.org/10.1038/s41559-017-0101>
- 470 Poelen, J. H., Simons, J. D., & Mungall, C. J. (2014). Global biotic interactions: An open infrastructure to  
471 share and analyze species-interaction datasets. *Ecological Informatics*, 24, 148–159. <https://doi.org/10.1016/j.ecoinf.2014.08.005>
- 472
- 473 Poisot, T. (2023). Guidelines for the prediction of species interactions through binary classification. *Methods  
474 in Ecology and Evolution*, 14(5), 1333–1345. <https://doi.org/10.1111/2041-210X.14071>
- 475 Poisot, T., Baiser, B., Dunne, J., Kéfi, S., Massol, F., Mouquet, N., Romanuk, T. N., Stouffer, D. B., Wood, S.  
476 A., & Gravel, D. (2016). Mangal – making ecological network analysis simple. *Ecography*, 39(4), 384–390.  
477 <https://doi.org/10.1111/ecog.00976>

- 478 Poisot, T., Cirtwill, A., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. (2016). The structure of  
479 probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312. <https://doi.org/10>
- 480 Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary  
481 through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>
- 482 Poisot, T., Stouffer, D. B., & Kéfi, S. (2016). Describe, understand and predict: Why do we need networks  
483 in ecology? *Functional Ecology*, 30(12), 1878–1882. <https://www.jstor.org/stable/48582345>
- 484 Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O’Hara, R. B., Parris, K. M., Vesk, P. A., &  
485 McCarthy, M. A. (2014). Understanding co-occurrence by modelling species simultaneously with a Joint  
486 Species Distribution Model (JSDM). *Methods in Ecology and Evolution*, 5(5), 397–406. <https://doi.org/>  
487 [10.1111/2041-210X.12180](https://doi.org/10.1111/2041-210X.12180)
- 488 Pringle, R. M. (2020). Untangling Food Webs. In *Unsolved Problems in Ecology* (pp. 225–238). Princeton  
489 University Press. <https://doi.org/10.1515/9780691195322-020>
- 490 Proulx, S. R., Promislow, D. E. L., & Phillips, P. C. (2005). Network thinking in ecology and evolution.  
491 *Trends in Ecology & Evolution*, 20(6), 345–353. <https://doi.org/10.1016/j.tree.2005.04.004>
- 492 Pyke, G. (1984). Optimal Foraging Theory: A Critical Review. *Annual Review of Ecology, Evolution and  
493 Systematic*, 15, 523–575. <https://doi.org/10.1146/annurev.ecolsys.15.1.523>
- 494 Quintero, E., Arroyo-Correa, B., Isla, J., Rodríguez-Sánchez, F., & Jordano, P. (2024). *Downscaling mu-  
495 tualistic networks from species to individuals reveals consistent interaction niches and roles within plant  
496 populations* (p. 2024.02.02.578595). bioRxiv. <https://doi.org/10.1101/2024.02.02.578595>
- 497 Rooney, N., McCann, K. S., & Moore, J. C. (2008). A landscape theory for food web architecture. *Ecology  
498 Letters*, 11(8), 867–881. <https://doi.org/10.1111/j.1461-0248.2008.01193.x>
- 499 Roopnarine, P. D. (2006). Extinction Cascades and Catastrophe in Ancient Food Webs. *Paleobiology*, 32(1),  
500 1–19. <https://www.jstor.org/stable/4096814>
- 501 Roopnarine, P. D. (2017). Ecological Modelling of Paleocommunity Food Webs. In *Conservation Paleobiology:  
502 Using the Past to Manage for the Future* (pp. 201–226). University of Chicago Press.
- 503 Saravia, L. A., Marina, T. I., Kristensen, N. P., De Troch, M., & Momo, F. R. (2022). Ecological network  
504 assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, 91(3),  
505 630–642. <https://doi.org/10.1111/1365-2656.13652>
- 506 Segar, S. T., Fayle, T. M., Srivastava, D. S., Lewinson, T. M., Lewis, O. T., Novotny, V., Kitching, R. L.,  
507 & Maunsell, S. C. (2020). The Role of Evolution in Shaping Ecological Networks. *Trends in Ecology &  
508 Evolution*, 35(5), 454–466. <https://doi.org/10.1016/j.tree.2020.01.004>
- 509 Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024). *A framework for  
510 reconstructing ancient food webs using functional trait data* (p. 2024.01.30.578036). bioRxiv. <https://doi.org/10.1101/2024.01.30.578036>


- 511 //doi.org/10.1101/2024.01.30.578036
- 512 Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters*,  
513 10(12), 1115–1123. <https://doi.org/10.1111/j.1461-0248.2007.01107.x>
- 514 Song, C., & Levine, J. M. (2024). *Rigorous (in)validation of ecological models* (p. 2024.09.19.613075). bioRxiv.  
515 <https://doi.org/10.1101/2024.09.19.613075>
- 516 Staniczenko, P. P. A., Lewis, O. T., Jones, N. S., & Reed-Tsochas, F. (2010). Structural dynamics and  
517 robustness of food webs. *Ecology Letters*, 13(7), 891–899. <https://doi.org/10.1111/j.1461-0248.2010.01485.x>
- 518 Stock, M. (2021). Pairwise learning for predicting pollination interactions based on traits and phylogeny.  
519 *Ecological Modelling*, 14.
- 520 Stouffer, D. B. (2019). All ecological models are wrong, but some are useful. *Journal of Animal Ecology*,  
521 88(2), 192–195. <https://doi.org/10.1111/1365-2656.12949>
- 522 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V.,  
523 Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction  
524 through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*,  
525 13(12), 2838–2849. <https://doi.org/10.1111/2041-210X.13835>
- 526 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Mercier, B.,  
527 Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2023). Graph embedding and transfer  
528 learning can help predict potential species interaction networks despite data limitations. *Methods in  
529 Ecology and Evolution*, 14(12), 2917–2930. <https://doi.org/10.1111/2041-210X.14228>
- 530 Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz,  
531 N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap  
532 towards predicting species interaction networks (across space and time). *Philosophical Transactions of  
533 the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>
- 534 Terry, J. C. D., & Lewis, O. T. (2020). Finding missing links in interaction networks. *Ecology*, 101(7), e03047.  
535 <https://doi.org/10.1002/ecy.3047>
- 536 Valdovinos, F. S., Hale, K. R. S., Dritz, S., Glaum, P. R., McCann, K. S., Simon, S. M., Thébaud, E., Wetzel,  
537 W. C., Wootton, K. L., & Yeakel, J. D. (2023). A bioenergetic framework for aboveground terrestrial  
538 food webs. *Trends in Ecology & Evolution*, 38(3), 301–312. <https://doi.org/10.1016/j.tree.2022.11.004>
- 539 Van De Walle, R., Logghe, G., Haas, N., Massol, F., Vandegehuchte, M. L., & Bonte, D. (2023). Arthro-  
540 pod food webs predicted from body size ratios are improved by incorporating prey defensive properties.  
541 *Journal of Animal Ecology*, 92(4), 913–924. <https://doi.org/10.1111/1365-2656.13905>
- 542 Vázquez, D. P., Blüthgen, N., Cagnolo, L., & Chacoff, N. P. (2009). Uniting pattern and process in plant–  
543

- 544 animal mutualistic networks: A review. *Annals of Botany*, 103(9), 1445–1457. <https://doi.org/10.1093/aob/mcp057>
- 545
- 546 Wells, K., & O'Hara, R. B. (2013). Species interactions: Estimating per-individual interaction strength and  
547 covariates before simplifying data into per-species ecological networks. *Methods in Ecology and Evolution*,  
548 4(1), 1–8. <https://doi.org/10.1111/j.2041-210x.2012.00249.x>
- 549 White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between body size  
550 and abundance in ecology. *Trends in Ecology & Evolution*, 22(6), 323–330. <https://doi.org/10.1016/j.tree.2007.03.007>
- 551
- 552 Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, 404(6774), 180–183.  
553 <https://doi.org/10.1038/35004572>
- 554 Williams, R. J., & Martinez, N. D. (2008). Success and its limits among structural models of complex food  
555 webs. *Journal of Animal Ecology*, 77(3), 512–519. <https://doi.org/10.1111/j.1365-2656.2008.01362.x>
- 556 Wootton, K. L., Curtsdotter, A., Roslin, T., Bommarco, R., & Jonsson, T. (2023). Towards a modular theory  
557 of trophic interactions. *Functional Ecology*, 37(1), 26–43. <https://doi.org/10.1111/1365-2435.13954>
- 558 Xie, J.-R., Zhang, P., Zhang, H.-F., & Wang, B.-H. (2017). Completeness of Community Structure in  
559 Networks. *Scientific Reports*, 7(1), 5269. <https://doi.org/10.1038/s41598-017-05585-6>
- 560 Yeakel, J. D., Pires, M. M., Rudolf, L., Dominy, N. J., Koch, P. L., Guimarães, P. R., & Gross, T. (2014).  
561 Collapse of an ecological network in Ancient Egypt. *PNAS*, 111(40), 14472–14477. <https://doi.org/10.1073/pnas.1408471111>
- 562
- 563 Yodzis, P. (1982). The Compartmentation of Real and Assembled Ecosystems. *The American Naturalist*,  
564 120(5), 551–570. <https://doi.org/10.1086/284013>
- 565 Yodzis, P., & Innes, S. (1992). Body Size and Consumer-Resource Dynamics. *The American Naturalist*,  
566 139(6), 1151–1175. <https://doi.org/10.1086/285380>


WED  
水 276  
PROCESS.



## REPRESENTATION



"Global" metaweb.  
All possible interactions for collection of spp.




"Regional" metaweb.  
All possible interactions for co-occurring species.

Realised networks.  
The dynamic interactions that occur.  
Dynamic config.


Figure 1: Aligning the various processes that determine interactions with the different network representations. First we start with a 'global metaweb' this network which captures all possible interactions for an arbitrary collection of species, we can further refine this network by taking in to consideration the co-occurrence of these difference species - as shown here we have two regions with some species (blue) that are found in both regions and others endemic to either region one (pink) or region two (orange). These regional metawebs to capture all possible interactions, however it only considers species that co-occur. However even within a region we do not expect all interactions to be realised but rather that there are multiple configurations of the regional metaweb over both space and time. The 'state' of the different network realisations are ultimately influenced not just by the co-occurrence of a species pair but rather the larger community context such as the abundance of different species, maximising energy gain, or indirect/higher order interactions.






### 1. A 'global' metaweb

Knowledge of the entire diet breadth of a species is valuable especially in terms of understanding how a species will respond to changes in the community - \*e.g.,\* invasions/rewilding exercises (where does the new species 'fit' within the network?) as well as potential capacity to shift its diet. Although this might make sense across space and not time but certain species act as links across the landscape.



### 2. Seasonal metaweb

- Knowledge at the finer scale is also valuable to understand/identify that there are in fact differences between the seasons
- Information of seasonal diet of species



### 3. Seasonal (elliptical networks)

Dynamics are useful because they are a representation of the different configurations/energy flows/ecosystem processes. Also to detect more nuanced shifts in diet - \*e.g.,\* seasonal diet shifts.

- Structurally informative
- can be @ even finer scale & time / space

Figure 2




Figure 3: An attempt to try and visualise a way to map the different scales of network representations to the way in which we can interrogate/ask questions about them?