

Unveiling the Complexity of Food Webs: A Comprehensive Overview of Definitions, Scales, and Mechanisms

Tanya Strydom ¹; Jennifer A. Dunne ²; Timothée Poisot ^{3,4}; Andrew P. Beckerman ¹

Abstract: Food webs are a useful abstraction and representation of the feeding links between species in a community and are used to infer many ecosystem level processes. However, the different theories, mechanisms, and criteria that underpin how a food web is defined and, ultimately, constructed means that not all food webs are representing the same ecological process. Here we present a synthesis of the different assumptions, scales and mechanisms that are used to define different ecological networks ranging from metawebs (an inventory of all potential interactions) to fully realised networks (interactions that occur within a given community over a certain timescale). Illuminating the assumptions, scales, and mechanisms of network inference allows a formal categorisation of how to use networks to answer key ecological and conservation questions and defines guidelines to prevent unintentional misuse or misinterpretation.

Keywords: food web, network construction, scientific ignorance

1 At the heart of modern biodiversity science are a set of concepts and theories about biodiversity, stability
2 and function. These relate to the abundance, distribution and services that biodiversity provides, and how
3 biodiversity – as an interconnected set of species – responds to multiple stressors. The interaction between
4 species (or individuals) is one of the fundamental building blocks of ecological communities provide a powerful
5 abstraction that can help quantify, conceptualise, and understand biodiversity dynamics, and ultimately,
6 one hopes, make prediction, mitigate change and manage services [ref]. Such network representations of
7 biodiversity (including within species diversity) are increasingly argued to be an asset to predictive ecology,
8 climate change mitigation and resource management. Here, it is argued that characterising biodiversity in
9 a network will allow deeper capacity to understand and predict the abundance, distribution, dynamics and
10 services provided by multiple species facing multiple stressors.

11 However, the way that a network is constructed (encoded) defines an epistemology of the network concept
12 which, we argue, can influence the resulting observations and conclusions about pattern and mechanisms
13 that are made (Brimacombe et al., 2023; Proulx et al., 2005). This process of constructing networks has two
14 major pillars: the data and theory, the latter representing an expression of mechanism and process giving
15 rise to patterns that emerge from collating interactions among species. Each of these pillars carries with it
16 a set of practical, semantic and conceptual constraints that not only influence progress in making network
17 ecology more valuable and potentially predictive, but help define the spatial, temporal and evolutionary scale
18 of assumptions we make and predictions we might generate from the networks.

19 With respect to data, it is extremely challenging to actually record species interactions in the field (Jordano,
20 2016a, 2016b). Despite notable herculean efforts (**Woodward? Benguela?** Maiorano et al. (2020)), actual
21 coverage of ‘real world’ interaction data remains sparse (Poisot et al., 2021). Against this practical challenge,
22 there is additionally high variance in the terminology we use to define networks. Finally, the mathematical
23 and statistical tools we use to construct, conceptualise, analyse and predict with these networks are also
24 highly variable.

25 1. what are the underlying assumptions about nodes, edges, scale and process that are made when we
26 attempt to delimit and describe a food webs;

27 2. are there families of commonly used tools that map onto assumptions about scales and processes;

28 The provision of this detail ultimately leads to a set of insights and conclusions about whether, when and
29 under what conditions network representations of biodiversity can contribute to the advancement of ecological
30 theory and generate value in predictive ecology. Specifically, we finish this perspective with an overview of
31 fundamental questions in ecology that we think can benefit from network thinking and a proposal that such

32 thinking can accelerate our capacity to predict the impact of multiple stressors on biodiverse communities.

33 1 Setting the Scene: The Not So Basics of Nodes and Edges

34 Defining a food web seems simple; it is the representation of the interactions (edges) between species (nodes),
35 however the definition of ‘edges’ and ‘nodes’, as well as the scale at which they are aggregated can take
36 many forms (Poisot, Stouffer, et al., 2016). Networks can be constructed at the population (the links among
37 individuals), community (the links between species), or metacommunity (changes between locations) level.
38 Even if one were to limit their scope to thinking of interaction networks only in terms of food webs at the
39 community-level there are still many ways to define the various components of the network Panel A of 1,
40 one needs to understand the different intentions/assumptions that are made when a food web is constructed.
41 Although the main goal of constructing a food web is to capture and represent the feeding links between
42 species there are many ways to define the nodes (*e.g.*, species or taxonomic group), edges (*e.g.*, *potential* or
43 *realised* feeding links), the magnitude of the edges (*e.g.*, binary vs probabilistic), and even how the network
44 itself is delimited (does it represent an aggregation of interactions over time?).

45 [Figure 1 about here.]

46 1.0.1 How do we define a node?

47 Although this may seem an elementary question in the context of food webs — a node *should* represent a
48 (taxonomic) species, the reality is that nodes can often represent an aggregation of different species - so called
49 ‘trophic species’ or segregation of species by life stages. Representing nodes as non-taxonomic species can be
50 useful in certain contexts (Williams & Martinez, 2000) and in cases where the adult and larval stages of a
51 species have different diets it may make ecological sense (Clegg et al., 2018) meaning that it is not uncommon
52 that networks often have nodes that have different definitions of a ‘species’ *e.g.* consisting of both taxonomic
53 and trophic species. Practical implications of how we are aggregating the nodes is that the resolution may
54 not always be ‘pixel perfect’ *i.e.*, we may be unable to assess the co-extinction risk of a species pair, however
55 there is value in having nodes that represent an aggregation of species, as these convey a much more general
56 overview of how the links are distributed within the community.

57 1.0.2 What is meant by an edge?

58 At its core links within food webs can be thought of as a representation of either feeding links between
59 species - be that realised (Pringle, 2020) or potential (Dunne, 2006), or representative of fluxes within the
60 community/system *e.g.*, energy transfer or material flow (Lindeman, 1942). How we specify links will influence

the resulting structure of the network - and the inferences we will make thereof. For example taking a food web that consists of links representing all *potential* feeding links for a community (*i.e.*, a metaweb) will be meaningless if you are interested in understanding the flow of energy through the system as the links within a metaweb do not represent environmental/energetic constraints. In addition to the various ways of defining the links between species pairs there are also a myriad of ways in which the links themselves can be quantified. Links between species are often treated as being present or absent (*i.e.*, binary) but it is also possible to use probabilities (which quantifies how likely an interaction is to occur, Poisot, Cirtwill, et al., 2016) or continuous measurements (which quantifies the strength of of an interaction, Berlow et al., 2004). Moving away from a purely binary way of representing allows us to quantify a level of (un)certainty of our knowledge of interactions (*i.e.*, moving from being able to ask if they are occurring to quantifying how likely they are to occur) does add an additional level of ‘complexity’ to the construction and interpretation of networks, but ultimately it will allow us to capture more information at different scales (Banville et al., 2024).

1.0.3 Putting the parts together

The ingredients one uses to construct networks from nodes and edges generates a unique representation of the mechanisms (see Section 2) that allow inference and reasoning about the structure, aspects of dynamics (*e.g.*, stability), and potentially the function of communities (*e.g.*, flux). It is thus important to keep in mind that different networks are going to be representing different processes and that can only be used for inference of some but not all aspects of the community at large. Here it may be meaningful to contextualise the different ‘types’ of food webs within the larger research programmes (or even practical needs) that have been driving the construction of them.

Before thinking about the ways in which we can predict networks it is perhaps meaningful to take a step back and think about the different criteria that must be met in order for an interaction to be able to occur between two species, specifically thinking of this in terms of distinguishing between the feasibility versus realisation of an interaction and how these are determined (and defined by) different ‘rules’/mechanisms. If we look at this feasibility-reality continuum (Figure 2) it is clear how the different predictive approaches (methods) tend to fall within one of the broader categories identified (distinguished) in the triangle. This is not to say that this shortcoming should be viewed as a ‘bug’ but rather a ‘feature’ of the field as it allows one to engage with, as well as construct networks at different scales, which is particularly valuable if one takes into consideration the considerable ‘data cost’ of predicting well resolved, realised networks in comparison to constructing high-level metawebs. However, it is important that there is an awareness and acknowledgement of where within this feasibility-reality one is working at and how this will impact and limit the contexts in

92 which the resulting network can be used and applied within.

93 **2 From Nodes and Edges to Scales, Context, and Processes**

94 Armed with these basics, it is now possible to review the scales and assumptions that are made by a wide
95 range of tools to assist in constructing networks against poor data with the hope of capturing important
96 processes that underpin accurate prediction. Our thesis centres on a four-tier conceptualization of networks:
97 co-occurrence, feasibility, mass effects and energetics. In the following sections we review each of these and
98 then provide a synthesis among them.

99 **2.1 Understanding the processes that determine species interactions**

100 Processes that are all-or-nothing (possibility) vs processes that are context dependent (likelihood). Processes
101 form the underlying logic of models (and arguably even empirical data capture). Processes influence how we
102 define the network (its anatomy). Ultimately when we put this all together it will influence how we can and
103 should use the resulting network.

104 Important goal here is to introduce the idea that there are multiple facets as to what determines the interaction
105 between species and that there is some sort of ‘scale of organisation’ Figure 2. We can then introduce these
106 different scales/theories and I think some key points to highlights are the features, limits, and descriptions
107 of these different scales (by that I mean what rules them, what finds them, and what binds them - sorry not
108 sorry). I think it is also worth either in this section or in the one where we talk about model families to
109 discuss the idea of ‘moving between’ different levels - e.g. downsampling but also feasibility - can we actually
110 do that? Another interesting discussion here (or maybe actually something that can make its way into the
111 concluding remarks) is thinking about what determines interactions vs what determines structure...

- 112 • These different theories are shown in Figure 2 and we can see there is some element of scaling (species
113 - population - individual)

114 [Figure 2 about here.]

115 **Evolutionary compatibility**

116 There is compelling evidence that the possibility of an interaction occurring between two species is the result
117 of their shared (co)evolutionary history (Dalla Riva & Stouffer, 2016; Gómez et al., 2010). In the more
118 proximal sense this is manifested as the ‘trait complementarity’ between two species, whereby one species
119 (the predator) has the ‘correct’ set of traits that allow it to chase, capture, kill, and consume the other species

120 (the prey). For species pairs where this condition is not met the link is deemed to be forbidden (Jordano,
121 2016b); *i.e.*, not physically possible and will always be absent within the network. In the context of trying
122 to determine the feasibility (*i.e.*, the *possibility*) of an interaction, phylogeny is an excellent predictor (Fricke
123 et al., 2022; Strydom et al., 2022) and allows one to construct what can be considered to be a metaweb.
124 In terms of thinking about the anatomy of an ‘feasibility network’ one should be aware that it is possible
125 to represent interactions as either binary (feasible/forbidden; *i.e.*, the traditional definition of a metaweb
126 Dunne (2006)) or as a probability (Banville et al., 2024), where the probability represents how likely that
127 the interaction between two species is feasible (what is the possibility of this interaction occurring?).

128 **(Co)occurrence**

129 Although the outright assumption that because two species are co-occurring it must mean that they are
130 interacting is inherently flawed (Blanchet et al., 2020), it is of course impossible for two species to interact (at
131 least in terms of feeding links) if they are not co-occurring in time and space. Thus co-occurrence data alone is
132 insufficient to build an accurate and ecologically meaningful representation of a food web having information
133 on the co-occurrence of species can further aid us in refining metawebs by allowing us to downsample the
134 network based on the species found in a specific location, or even add additional uncertainty based in how
135 likely species are to co-occur (Dansereau et al., 2023). Additionally the interplay between the interaction
136 between a species pair and their co-occurrence is meaningful when one is operating in the space of trying to
137 determine the distribution of a species (Higino et al., 2023), and forms a key component of some of the next
138 generation species distribution models *e.g.*, joint SDMs (Pollock et al., 2014).

139 **Abundance**

140 The abundance of the different species within the community can influence the likelihood of an interaction
141 occurring in a myriad of ways. There is the argument that networks (and the interactions that make them up)
142 are driven by only the abundance of the different species and not the characteristics (traits), *sensu* neutral
143 processes (Canard et al., 2012). Alternatively the abundance of species in a community can influence which
144 interactions are ultimately realised (Banville et al., 2024; Poisot et al., 2015).

145 **Predator choice (energetic cost)**

146 Ultimately, predator choice is underpinned by the energetic cost-benefit of trying to catch, kill, and consume
147 prey, and is well described within optimal foraging theory [ref] and rests on the idea that the prey a predator
148 chooses to target is one that will have the greatest return on energy with the lowest energetic cost. There are
149 additional bodies of work that attempt to include the cost of movement that the environment imposes on an
150 individual (Cherif et al., 2024) as well as 2D/3D search space (Pawar et al., 2012). In terms of formalising

151 these processes in the context of predicting networks using diet models (Beckerman et al., 2006; Petchey
152 et al., 2008) that have predator choice determined by the handling time, energy content, prey density, and
153 predator attack rate. Wootton et al. (2023) developed a model that moves the energy of the system into
154 different modules related to the process of the predator acquiring energy from the prey.

155 **Indirect interactions**

156 The realisation (presence/absence) or strength of trophic interactions themselves can also be modified by
157 other, indirect (non-trophic), interactions (Golubski & Abrams, 2011; Pilosof et al., 2017), this can be either
158 ‘directly’ through *e.g.*, competition or ‘indirectly’ *e.g.*, mutualistic/facilitative interactions will alter the fine-
159 scale distribution and abundance of some species (Kéfi et al., 2012, 2015).

160 It should be self evident that the different processes discussed above are all ultimately going to influence the
161 realisation of interactions as well as the structure of a network, however they are acting at different scales
162 of organisation. Both the **co-occurrence** and the **evolutionary compatibility** are valid at the scale of
163 the species pair of interest, that is the *possibility* of an interaction being present/absent is assessed at the
164 pairwise level and one is left with a ‘list’ of interactions that are present/absent. Although it is possible to
165 build a network (*i.e.*, metaweb) from this information it is important to be aware that the structure of this
166 network is not constrained by real-world dynamics or conditions, just because species are able to interact
167 does not mean that they will (Poisot et al., 2015). In order to construct a network who’s structure is a
168 closer approximation of reality (localised interactions) one needs to take into consideration properties of the
169 community as a whole and not just the two species of interest.

170 **downsampling paragraph??**

171 **3 Network prediction is nuanced**

172 The different models that are used to either predict or construct networks have an underlying philosophy
173 that often only captures one or a few of the processes discussed in Section 2.1, has implications for how
174 the resulting network is defined Section 1, which will ultimately delimit and define what inferences can
175 be made from the resulting network. Selecting a model for the task of network prediction should come
176 down to two things; what *aspect* of a food web one is interested in predicting, and what data are available,
177 necessary, and sufficient, and what is the purpose of wanting to predict a network? It is important that
178 a researcher is aware of this to ensure that the appropriate model is selected. Broadly researchers will be
179 interested in predicting/constructing two different types of networks; *metawebs*, which is essentially a list of
180 all interactions that are *possible* for a specific community (*i.e.*, at the scale of the species pairs), or being

able to predict location specific, *realised*, networks for the community (*i.e.*, at the scale of the community). The nature of metawebs means that they are unable to capture the structural metrics of realised/‘real-world’ networks (Caron et al., 2024). The researcher is also constrained by the data needs of both the model as well as the network type; for example in order to predict a realised network one needs additional data (*e.g.*, abundance), making metawebs a more feasible choice in data-poor contexts (*e.g.*, Strydom et al. (2023) construct a metaweb using a species list and a phylogenetic tree). The final question is assessing the purpose of predicting a network - is it to create a series of simulated, species agnostic but still ecologically plausible, networks [*e.g.*,] or to predict a network for a specific community at a specific location. It is these three points that will ultimately dictate which model is going to best allow one to predict the appropriate network.

Although the ability to predict ‘real-world’ interactions (and the resulting food webs) can have more intuitive ‘real world’ applications *e.g.*, being able to ‘recover’ food webs that have since gone extinct (Dunne et al., 2008; Yeakel et al., 2014), using pairwise interactions to understand species distributions (Pollock et al., 2014) or even co-extinction risk (Dunn et al., 2009), a more structural approach to network construction affords one an opportunity to interrogate some of the more high-level mechanisms that are structuring networks.

3.1 Models that predict structure

Although we identify mechanisms that determine species interactions in Section 2.1 not all models that are used to predict networks operate at this ‘mechanistic’ level, but rather represent the *structure* of a network based on a series of *a priori* assumptions of network connectance (*e.g.*, the niche model Williams & Martinez (2000)) or other structural features (*e.g.*, stochastic block model, Xie et al. (2017)) of a *realised* network. Importantly these structural models do not make species specific predictions (they are species agnostic and usually treat nodes as trophic species) and so cannot be used to determine if an interaction is either possible or realised between two species (*i.e.*, one cannot use these models to determine if species *a* eats species *b*). Although this means this suite of models are unsuitable as tools for predicting interactions, they have been shown to be sufficient tools to predict the structure of networks (Williams & Martinez, 2008).

3.2 How do we predict food webs?

There as many ways to predict networks as what there is to define them and along with taking into consideration the points raised in the previous section it is also beneficial to think about the context in which the different models were developed - and how this will influence the networks that they produce... Also it is not feasibly possible to list every single approach that has been developed to predict networks and so we will

211 present what we believe to be the broad families that represent the different approaches to predicting net-
212 works, particularly how these relate to the processes identified in Section 2.1, as well as models that predict
213 network structure (see Section 3.1).

214 In order for a model to formalise a ‘complete’ food web it is necessary to formalise two aspects of the network,
215 ‘who eats whom’ (to determine the links between nodes) as well as the structure of the network (to limit
216 the distribution of links), however most models are inclined to focus on one of the two aspects. As there
217 are many food web models to choose from it is perhaps useful to think about the models in terms of model
218 families, a summary of these families is presented in Table 1 highlights the differences and similarities of the
219 philosophies and assumptions that determine a network. A more extensive overview of the different models
220 that fall with in the different model families can be found in SuppMat 1 and for a more detailed breakdown
221 of the different ‘traits’ of the model families refer to SuppMat 2.

Table 1: A summary of the different families of tools that can be used to generate food webs. Here ‘type’ leans on the ideas from Momal et al. (2020) in terms of reconstructing networks from other known (observed) interactions and inferring interactions without relying on observed interactions.

Model family	Assumption (process)	Theory	Type	Key reference
null	Links are randomly distributed within a network		structural network	
neutral	Network structure is random, but species abundance determines links between nodes	abundance	structural network	Canard et al. (2012)
resource	Networks are interval, species can be ordered on a ‘niche axis’		structural network	Williams & Martinez (2008)
generative	Networks are determined by their structural features		structural network	
energetic	Interactions are determined by energetic costs	abundance + energy	‘energy’ network	

Model family	Assumption (process)	Theory	Type	Key reference
graph embedding	Interactions can be predicted from the latent traits of networks	evolutionary compatibility	metaweb	Strydom et al. (2023)
trait matching	Interactions can be inferred by a mechanistic framework/relationships	evolutionary compatibility	metaweb	Morales-Castilla et al. (2015)
binary classifiers	Interactions can be predicted by learning the relationship between interactions and ecologically relevant predictors	evolutionary compatibility	metaweb	Pichler et al. (2020)
expert knowledge	‘Boots on the ground’ ecological knowledge and observations	evolutionary compatibility	metaweb	
data scavenging	Webscraping to create networks from online databases		metaweb	Poisot, Gravel, et al. (2016) (if you squint?)
co-occurrence	co-occurrence patterns arise from interactions so we can use these patterns to reverse engineer the interactions	co-occurrence	co-occurrence network	

²²² 4 Making Progress with Networks

²²³ In this section I want to highlight that we don’t actually have any clear guidelines as to how we can ‘use’

²²⁴ networks - which probably stems from both the fact that when I am talking about a network and when

²²⁵ someone else is talking about a network we may actually be talking about two very different conceptualisations
²²⁶ of ‘a network’ (this should actually be a selling point in the intro - may have just found my *raison d’être*) as
²²⁷ well as that a lot of the ideas that we have about networks are not really tied to any sort of tangible function
²²⁸ (i.e. Tim’s GeoBon ms thing-y). However we can maybe at least try to present some guidelines - but I think
²²⁹ specifically within the sort of Petchy dilemma space and clearly tied to the ideas we discuss in the ms. This
²³⁰ includes: understanding the limits of how a network is defined and how the underlying theory impacts the use
²³¹ as well as data?? IDK we need to shoehorn data in here somehow... We can also use this as a gap identifying
²³² space and I think the framing can still rest under the limits concept particularly time, space, and boundaries
²³³ - which will all probably fall under some aspect of biological scale... We can also raise the idea of trust - as in
²³⁴ which methods have more support/trust than others. Also what even a ‘real’ network entails (and this links
²³⁵ again back to Tim’s stuff) as well as a subtle jab at Pringles notion that the most critical issue in the world
²³⁶ of food webs is being able to identify every. single. link. even though there is no real discussion as to what
²³⁷ is an ‘opportunistic’ link vs a link that represents a sustainable energy source for a population (or would it
²³⁸ be an individual)...

²³⁹ We need to be aware of the parameter space that is possible given a specific definition of a network and
²⁴⁰ operate within those parameters.

²⁴¹ 5 Concluding remarks

²⁴² I think the idea of time and how we are aggregating networks across that should be a prominent feature
²⁴³ here...

- ²⁴⁴ • In certain situations structure is ‘enough’ but there may be use cases where we are really interested in
²⁴⁵ the node-level interactions *i.e.*, species identity is a thing we care about and need to be able to retrieve
²⁴⁶ specific interactions at specific nodes correctly.
- ²⁴⁷ • Why do interaction models do so badly at predicting structure? Nuance of metaweb vs realisation but
²⁴⁸ also time? At the core of it interaction models are trained on existing interaction data; this is data
²⁴⁹ that are most likely closer to a metaweb than a local realisation even if they are being inventoried at a
²⁵⁰ small scale...
 - ²⁵¹ – We can briefly shoehorn downsampling here maybe??
 - ²⁵² • It will be interesting to bring up the idea that if a model is missing a specific pairwise link but doing
²⁵³ well overall then when does it matter?

- 254 – The fact that *some* people are concerned about the taxonomic resolution and cascading effects
255 those might have on our understanding of network structure (Pringle, 2020; Pringle & Hutchinson,
256 2020), but that puts us in a place where we are at risk of losing our ability to distinguish the wood
257 from the tree - are we not (at least at times) concerned more with understanding ecosystem level
258 processes than with needing to understand things *perfectly* at the species level.
- 259 – I don't think these 'rare'/nuanced links (e.g. carnivorous hippos) are going to rock the boat when
260 we think about networks at the structural level.

261 "The resolution of food-web data is demonic because it can radically change network topology
262 and associated biological inferences in ways that are unknowable in the absence of better data." -
263 Pringle & Hutchinson (2020) The counter to this is that structural models are often not working
264 at the species level and thus the structure remains 'unchanged' when you increase the resolution
265 - I don't think that people are that concerned with the structure of real world networks barring
266 connectance and since that scales with species richness anyway your final proportion will probably
267 still remain the same...

- 268 • I think a big take home will (hopefully) be how different approaches do better in different situations
269 and so you as an end user need to take this into consideration and pick accordingly. I think Petchey
270 et al. (2011) might have (and share) some thoughts on this. I feel like I need to look at Berlow et al.
271 (2008) but maybe not exactly in this context but vaguely adjacent.

272 – I think this is sort of the crux of the argument presented in Brimacombe et al. (2024) as well.

273 "we highlight an interesting paradox: the models with the best performance measures are not
274 necessarily the models with the closest reconstructed network structure." - Poisot (2023)

- 275 • Do we need network models to predict interactions and interaction models to predict structure?

276 – "Another argument for the joint prediction of networks and interactions is to reduce circularity
277 and biases in the predictions. As an example, models like linear filtering generate probabilities of
278 non-observed interactions existing, but do so based on measured network properties." - Strydom
279 et al. (2021)

280 – Aligning (dove-tailing) with this the idea of ensemble modelling as presented by Becker et al.
281 (2022)

- 282 • Close out with a call to action that we have models that predict networks very well and models that
283 predict interactions very well but nothing that is doing well at predicting both - this is where we should

284 be focusing our attention when it comes to furthering model development...

- 285 • Do we expect there to be differences when thinking about unipartite vs bipartite networks? Is there
286 underlying ecology/theory that would assume that different mechanisms (and thus models) are relevant
287 in these two ‘systems’.

288 – The Terry & Lewis (2020) paper looks at some methods but is specifically looking at a bipartite
289 world...

290 do we bring this up? this could be a box... if we have the ‘finances’ for it... otherwise it should go to the
291 outstanding questions fur sure

292 “That being said, there is a compelling argument for the need to ‘combine’ these smaller functional units
293 with larger spatial networks (Fortin et al., 2021) and that we should also start thinking about the interplay
294 of time and space (Estay et al., 2023). Although deciding exactly what measure might actually be driving
295 differences between local networks and the regional metaweb might not be that simple (Saravia et al., 2022).”

296 **5.1 Time**

297 Look at Hutchinson et al. (2019)

298 We lack a clear agenda (and conceptualisation) as to what the appropriate level of aggregation is for a
299 ‘network’. Realistically most empirical networks are more aligned with ‘feasibility networks’ as opposed to
300 ‘realised networks’ as they are often the result of some sort of aggregation of observations across time. This
301 ‘problem’ is two-fold. Firstly we need to think about how this affects any sort of development of theory
302 that sits closer to the ‘realised network’ side of the spectrum - how often are we trying to ask and answer
303 questions about realised networks using feasible networks? The second is that this lack of ‘direction’ as to
304 how we should define a network is (actually) probably one of the biggest barriers that is affecting the use of
305 networks in applied settings...

306 Another time perspective question is when do we determine a link to be ‘real’... In the context of feasible
307 networks this is perhaps clearer - all things equal would the predator be bale to consume the prey. However
308 in the realised space there is also the question of the long term ‘energetic feasibility’ of an interaction - just
309 because an interaction is possible in the now is it able to sustain a population in the long term. And what
310 is the scale for that long term - are we thinking at the generational scale? Because ultimately when we are
311 constructing a network we are aggregating not only across space but also across time.

₃₁₂ **Glossary**

Term	Definition
food web	a representation of feeding links between species
topology generator	a model that predicts a network based on assumptions of structure, this network is species agnostic in the sense that it does not necessarily contain information at the node level
interaction predictor	a model that predicts species interactions, these interactions can be used to construct a network but there are no <i>a priori</i> assumptions as that will constrain the network structure
model	A tool that can be used to construct food webs, where the resulting network is a representation of a real world network. Models typically only capture specific elements of real world networks and are intended to be used in specific settings
model family	A family of models that share an underlying philosophy when it comes to the mapping, pragmatism, and reduction of a network. Families have the same underlying philosophies and assumptions that determine the links between nodes as well as how these may be encoded
metaweb	A network that represents <i>all</i> the potential links between species. Importantly these links will not necessarily all be realised in a specific location for a specific time
realised network	A network that represents the links between species that are occurring. These networks represent a very localised network...
potential feeding link	links that indicate that an interaction is ecologically feasible but not realised <i>per se</i> (a metaweb would contain potential feeding links)

Term	Definition
realised feeding link	links that indicate that the interaction is realised ‘in the field’. (a realised network contains realised feeding links)
confusion matrix	captures the number of true positives (interaction predicted as present when it is present), false negatives (interaction predicted as absent when it is present), false positives (interaction predicted as present when it is absent), and true negatives (interaction predicted as absent when it is absent)

313 Outstanding questions

- 314 • non-consumptive effects
- 315 • how do we define the spatial and temporal ‘boundaries’ of a network?
- 316 • how do we define a ‘real’ network?

317 References

- 318 Banville, F., Strydom, T., Blyth, P., Brimacombe, C., Catchen, M. D., Dansereau, G., Higino, G., Malpas,
 319 T., Mayall, H., Norman, K., Gravel, D., & Poisot, T. (2024). *Deciphering probabilistic species interaction*
 320 *networks*. EcoEvoRxiv. <https://doi.org/10.32942/X28G8Z>
- 321 Becker, D. J., Albery, G. F., Sjodin, A. R., Poisot, T., Bergner, L. M., Chen, B., Cohen, L. E., Dallas, T.
 322 A., Eskew, E. A., Fagre, A. C., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Stock, M., Teeling,
 323 E. C., & Carlson, C. J. (2022). Optimising predictive models to prioritise viral discovery in zoonotic
 324 reservoirs. *The Lancet Microbe*, 3(8), e625–e637. [https://doi.org/10.1016/S2666-5247\(21\)00245-7](https://doi.org/10.1016/S2666-5247(21)00245-7)
- 325 Beckerman, A. P., Petchey, O. L., & Warren, P. H. (2006). Foraging biology predicts food web complexity.
 326 *Proceedings of the National Academy of Sciences*, 103(37), 13745–13749. <https://doi.org/10.1073/pnas.0603039103>
- 328 Berlow, E. L., Brose, U., & Martinez, N. D. (2008). The “Goldilocks factor” in food webs. *Proceedings of*
 329 *the National Academy of Sciences*, 105(11), 4079–4080. <https://doi.org/10.1073/pnas.0800967105>

- 330 Berlow, E. L., Neutel, A.-M., Cohen, J. E., de Ruiter, P. C., Ebenman, B., Emmerson, M., Fox, J. W., Jansen,
331 V. A. A., Iwan Jones, J., Kokkoris, G. D., Logofet, D. O., McKane, A. J., Montoya, J. M., & Petchey, O.
332 (2004). Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73(3),
333 585–598. <https://doi.org/10.1111/j.0021-8790.2004.00833.x>
- 334 Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions.
335 *Ecology Letters*, 23(7), 1050–1063. <https://doi.org/10.1111/ele.13525>
- 336 Brimacombe, C., Bodner, K., & Fortin, M.-J. (2024). *Applying a method before its proof-of-concept: A
337 cautionary tale using inferred food webs*. <https://doi.org/10.13140/RG.2.2.22076.65927>
- 338 Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings
339 of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4),
340 e3002068. <https://doi.org/10.1371/journal.pbio.3002068>
- 341 Canard, E., Mouquet, N., Marescot, L., Gaston, K. J., Gravel, D., & Mouillot, D. (2012). Emergence of
342 Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7(8), e38295. <https://doi.org/10.1371/journal.pone.0038295>
- 343 Caron, D., Brose, U., Lurgi, M., Blanchet, F. G., Gravel, D., & Pollock, L. J. (2024). Trait-matching models
344 predict pairwise interactions across regions, not food web properties. *Global Ecology and Biogeography*,
345 33(4), e13807. <https://doi.org/10.1111/geb.13807>
- 346 Cherif, M., Brose, U., Hirt, M. R., Ryser, R., Silve, V., Albert, G., Arnott, R., Berti, E., Cirtwill, A.,
347 Dyer, A., Gauzens, B., Gupta, A., Ho, H.-C., Portalier, S. M. J., Wain, D., & Wootton, K. (2024). The
348 environment to the rescue: Can physics help predict predator-prey interactions? *Biological Reviews*,
349 n/a(n/a). <https://doi.org/10.1111/brv.13105>
- 350 Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on food web structure.
351 *Ecology*, 99(12), 2712–2720. <https://doi.org/10.1002/ecy.2523>
- 352 Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food webs' backbones
353 using functional traits. *Oikos*, 125(4), 446–456. <https://doi.org/10.1111/oik.02305>
- 354 Dansereau, G., Barros, C., & Poisot, T. (2023). *Spatially explicit predictions of food web structure from
355 regional level data*.
- 356 Dunn, R. R., Harris, N. C., Colwell, R. K., Koh, L. P., & Sodhi, N. S. (2009). The sixth mass coextinction:
357 Are most endangered species parasites and mutualists? *Proceedings. Biological Sciences*, 276(1670),
358 3037–3045. <https://doi.org/10.1098/rspb.2009.0413>
- 359 Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological
360 networks: Linking structure and dynamics* (pp. 27–86). Oxford University Press.
- 361 Dunne, J. A., Williams, R. J., Martinez, N. D., Wood, R. A., & Erwin, D. H. (2008). Compilation and

- 363 Network Analyses of Cambrian Food Webs. *PLOS Biology*, 6(4), e102. <https://doi.org/10.1371/journal.pbio.0060102>
- 364
- 365 Estay, S. A., Fortin, M.-J., & López, D. N. (2023). Editorial: Patterns and processes in ecological networks
366 over space. *Frontiers in Ecology and Evolution*, 11.
- 367 Fortin, M.-J., Dale, M. R. T., & Brimacombe, C. (2021). Network ecology in dynamic landscapes. *Proceedings
368 of the Royal Society B: Biological Sciences*, 288(1949), rspb.2020.1889, 20201889. <https://doi.org/10.1098/rspb.2020.1889>
- 369
- 370 Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning,
371 J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene.
372 *Science*, 377(6609), 1008–1011. <https://doi.org/10.1126/science.abn4012>
- 373 Golubski, A. J., & Abrams, P. A. (2011). Modifying modifiers: What happens when interspecific interactions
374 interact? *Journal of Animal Ecology*, 80(5), 1097–1108. <https://doi.org/10.1111/j.1365-2656.2011.01852.x>
- 375
- 376 Gómez, J. M., Verdú, M., & Perfectti, F. (2010). Ecological interactions are evolutionarily conserved across
377 the entire tree of life. *Nature*, 465(7300), 918–921. <https://doi.org/10.1038/nature09113>
- 378 Higino, G. T., Banville, F., Dansereau, G., Muñoz, N. R. F., Windsor, F., & Poisot, T. (2023). Mismatch
379 between IUCN range maps and species interactions data illustrated using the Serengeti food web. *PeerJ*,
380 11, e14620. <https://doi.org/10.7717/peerj.14620>
- 381 Hutchinson, M. C., Bramon Mora, B., Pilosof, S., Barner, A. K., Kéfi, S., Thébault, E., Jordano, P., &
382 Stouffer, D. B. (2019). Seeing the forest for the trees: Putting multilayer networks to work for community
383 ecology. *Functional Ecology*, 33(2), 206–217. <https://doi.org/10.1111/1365-2435.13237>
- 384 Jordano, P. (2016a). Chasing Ecological Interactions. *PLOS Biology*, 14(9), e1002559. <https://doi.org/10.1371/journal.pbio.1002559>
- 385
- 386 Jordano, P. (2016b). Sampling networks of ecological interactions. *Functional Ecology*. <https://doi.org/10.1111/1365-2435.12763>
- 387
- 388 Kéfi, S., Berlow, E. L., Wieters, E. A., Joppa, L. N., Wood, S. A., Brose, U., & Navarrete, S. A. (2015).
389 Network structure beyond food webs: Mapping non-trophic and trophic interactions on Chilean rocky
390 shores. *Ecology*, 96(1), 291–303. <https://doi.org/10.1890/13-1424.1>
- 391 Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A., Boit, A., Joppa, L. N.,
392 Lafferty, K. D., Williams, R. J., Martinez, N. D., Menge, B. A., Blanchette, C. A., Iles, A. C., & Brose,
393 U. (2012). More than a meal... integrating non-feeding interactions into food webs: More than a meal
394 *Ecology Letters*, 15(4), 291–300. <https://doi.org/10.1111/j.1461-0248.2011.01732.x>
- 395 Lindeman, R. L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23(4), 399–417. <https://doi.org/>

- 396 10.2307/1930126
- 397 Maiorano, L., Montemaggiori, A., Ficetola, G. F., O'Connor, L., & Thuiller, W. (2020). TETRA-EU 1.0: A
398 species-level trophic metaweb of European tetrapods. *Global Ecology and Biogeography*, 29(9), 1452–1457.
- 399 <https://doi.org/10.1111/geb.13138>
- 400 Momal, R., Robin, S., & Ambroise, C. (2020). Tree-based inference of species interaction networks from
401 abundance data. *Methods in Ecology and Evolution*, 11(5), 621–632. <https://doi.org/10.1111/2041-210X.13380>
- 402 13380
- 403 Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from
404 proxies. *Trends in Ecology & Evolution*, 30(6), 347–356. <https://doi.org/10.1016/j.tree.2015.03.014>
- 405 Pawar, S., Dell, A. I., & Savage, V. M. (2012). Dimensionality of consumer search space drives trophic
406 interaction strengths. *Nature*, 486(7404), 485–489. <https://doi.org/10.1038/nature11131>
- 407 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2008). Size, foraging, and food web
408 structure. *Proceedings of the National Academy of Sciences*, 105(11), 4191–4196. <https://doi.org/10.1073/pnas.0710672105>
- 409
- 410 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2011). Fit, efficiency, and biology: Some
411 thoughts on judging food web models. *Journal of Theoretical Biology*, 279(1), 169–171. <https://doi.org/10.1016/j.jtbi.2011.03.019>
- 412
- 413 Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine learning algorithms
414 to infer trait-matching and predict species interactions in ecological networks. *Methods in Ecology and
415 Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>
- 416 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological networks.
417 *Nature Ecology & Evolution*, 1(4), 101. <https://doi.org/10.1038/s41559-017-0101>
- 418 Poisot, T. (2023). Guidelines for the prediction of species interactions through binary classification. *Methods
419 in Ecology and Evolution*, 14(5), 1333–1345. <https://doi.org/10.1111/2041-210X.14071>
- 420 Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., Mercier, B., Violet, C., &
421 Vissault, S. (2021). Global knowledge gaps in species interaction networks data. *Journal of Biogeography*,
422 48(7), 1552–1563. <https://doi.org/10.1111/jbi.14127>
- 423 Poisot, T., Cirtwill, A., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. (2016). The structure of
424 probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312. <https://doi.org/10.1111/2041-210X.12508>
- 425 Poisot, T., Gravel, D., Leroux, S., Wood, S. A., Fortin, M.-J., Baiser, B., Cirtwill, A. R., Araújo, M. B.,
426 & Stouffer, D. B. (2016). Synthetic datasets and community tools for the rapid testing of ecological
427 hypotheses. *Ecography*, 39(4), 402–408. <https://doi.org/10.1111/ecog.01941>
- 428 Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction networks vary

- 429 through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>
- 430 Poisot, T., Stouffer, D. B., & Kéfi, S. (2016). Describe, understand and predict: Why do we need networks
431 in ecology? *Functional Ecology*, 30(12), 1878–1882. <https://www.jstor.org/stable/48582345>
- 432 Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O’Hara, R. B., Parris, K. M., Vesk, P. A., &
433 McCarthy, M. A. (2014). Understanding co-occurrence by modelling species simultaneously with a Joint
434 Species Distribution Model (JSDM). *Methods in Ecology and Evolution*, 5(5), 397–406. <https://doi.org/10.1111/2041-210X.12180>
- 435 Pringle, R. M. (2020). Untangling Food Webs. In *Unsolved Problems in Ecology* (pp. 225–238). Princeton
436 University Press. <https://doi.org/10.1515/9780691195322-020>
- 437 Pringle, R. M., & Hutchinson, M. C. (2020). Resolving Food-Web Structure. *Annual Review of Ecology,
438 Evolution and Systematics*, 51(Volume 51, 2020), 55–80. <https://doi.org/10.1146/annurev-ecolsys-110218-024908>
- 439 Proulx, S. R., Promislow, D. E. L., & Phillips, P. C. (2005). Network thinking in ecology and evolution.
440 *Trends in Ecology & Evolution*, 20(6), 345–353. <https://doi.org/10.1016/j.tree.2005.04.004>
- 441 Saravia, L. A., Marina, T. I., Kristensen, N. P., De Troch, M., & Momo, F. R. (2022). Ecological network
442 assembly: How the regional metaweb influences local food webs. *Journal of Animal Ecology*, 91(3),
443 630–642. <https://doi.org/10.1111/1365-2656.13652>
- 444 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Hemming, V.,
445 Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2022). Food web reconstruction
446 through phylogenetic transfer of low-rank network representation. *Methods in Ecology and Evolution*,
447 13(12), 2838–2849. <https://doi.org/10.1111/2041-210X.13835>
- 448 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin, M.-J., Mercier, B.,
449 Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T. (2023). Graph embedding and transfer
450 learning can help predict potential species interaction networks despite data limitations. *Methods in
451 Ecology and Evolution*, 14(12), 2917–2930. <https://doi.org/10.1111/2041-210X.14228>
- 452 Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz,
453 N. R., Higino, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap
454 towards predicting species interaction networks (across space and time). *Philosophical Transactions of
455 the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>
- 456 Terry, J. C. D., & Lewis, O. T. (2020). Finding missing links in interaction networks. *Ecology*, 101(7), e03047.
457 <https://doi.org/10.1002/ecy.3047>
- 458 Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*, 404(6774), 180–183.
459 <https://doi.org/10.1038/35004572>

- 462 Williams, R. J., & Martinez, N. D. (2008). Success and its limits among structural models of complex food
463 webs. *Journal of Animal Ecology*, 77(3), 512–519. <https://doi.org/10.1111/j.1365-2656.2008.01362.x>
- 464 Wootton, K. L., Curtsdotter, A., Roslin, T., Bommarco, R., & Jonsson, T. (2023). Towards a modular theory
465 of trophic interactions. *Functional Ecology*, 37(1), 26–43. <https://doi.org/10.1111/1365-2435.13954>
- 466 Xie, J.-R., Zhang, P., Zhang, H.-F., & Wang, B.-H. (2017). Completeness of Community Structure in
467 Networks. *Scientific Reports*, 7(1), 5269. <https://doi.org/10.1038/s41598-017-05585-6>
- 468 Yeakel, J. D., Pires, M. M., Rudolf, L., Dominy, N. J., Koch, P. L., Guimarães, P. R., & Gross, T. (2014).
469 Collapse of an ecological network in Ancient Egypt. *PNAS*, 111(40), 14472–14477. <https://doi.org/10.1073/pnas.1408471111>
- 470



Figure 1: The many ways in which a food web can be defined and described at the node, edge, and even network level.

the "driver" of the interaction
≠ the underlying theory (1:1) & the
'way' we get to the answer is diff
e.g. co-occurrence is about sharing
space/time but we use niche theory/
env. filtering as the means to determine
co-occurrence.

JUN	27	THU	木	179
			(mechanism)	
			what determines	
			interaction	
6		Body of theory		
15	Species	fund. niches env. filter	co- occurrence	
21		buck bones (mora)	"capacity"	
0	pop.	truit-match		
3	indiv.	neutral. funct. resp.	pop size / dynamics	
?? allometric scaling ??				
				: THEORY OF WHAT DETERMINES FEEDING LINKS (INTERACTIONS) BETWEEN SPECIES

Figure 2: TODO.