

From Metawebs to Realised Webs: A Framework for Ecological Network Representation under Global Change

Tanya Strydom ^{1*}, Alexander M. Dunhill ², Jennifer A. Dunne ³, Timothée Poisot ^{4,5}, Andrew P. Beckerman ¹

¹ School of Biosciences, University of Sheffield, Sheffield, UK, S10 2TN

² School of Earth, Environment and Sustainability, University of Leeds, Leeds, UK, LS2 9JT

³ Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

⁴ Université de Montréal, Département de Sciences Biologiques, 1375 Avenue Thérèse-Lavoie-Roux, Montréal, QC H2V 0B3, Canada

⁵ Québec Centre for Biodiversity Sciences, 1205 Docteur Penfield Avenue, Montreal, Quebec, H3A 1B1, Canada

*Correspondence: t.strydom@sheffield.ac.uk

Abstract: Ecological networks are increasingly used to understand biodiversity and predict ecosystem responses to environmental change, yet their application is often limited by uncertainty about the assumptions embedded in different network representations. Here, we present a theory-driven framework that views network construction as a hierarchical transition from metawebs, which represent the feasibility of interactions, to realised webs, which represent interactions expressed in specific spatiotemporal contexts. We identify five key processes underlying this transition: evolutionary compatibility and co-occurrence, which define interaction feasibility, and abundance, diet choice, and non-trophic interactions, which determine interaction realisation. We further position these processes along a continuum of network construction methods, from inductive approaches that infer structure from observations to deductive approaches that generate structure from mechanistic principles. By explicitly linking network representations to their underlying assumptions and scale-dependent processes, our framework clarifies the questions that different networks can address, highlights key challenges in moving between potential and realised interactions, and provides a roadmap for developing predictive network ecology. We argue that greater conceptual clarity is essential for advancing biodiversity science and for the effective application of networks in conservation, environmental management, and forecasting ecological change in the Anthropocene.

Keywords: food web, network construction, biodiversity, scale and process, interaction modelling, trophic network

1 Introduction

At the heart of modern biodiversity science are a set of concepts and theories about species richness, stability, and function, which have been discussed since the foundational work of Elton (2001), Lindeman (1942), and MacArthur (1955). These relate to the abundance, distribution, functions, and services that biodiversity provides. Network representations of interactions among organisms are increasingly argued to be an asset to understanding and predicting the impacts of multiple, simultaneous stress on biodiversity (Cohen et al., 1990; Dunne et al., 2002; Martinez & Dunne, 1998; Simmons et al., 2021). Documenting interactions is thus one of the fundamental building blocks of community ecology and provides a powerful abstraction for mathematical and statistical modelling of biodiversity to make predictions, and to mitigate and manage threats (Galiana & Araújo, 2026; Windsor et al., 2023).

However, there is a growing discourse around limitations to the interpretation and applied use of networks, which have been recognised since early discussions of sampling effort and aggregation in food webs (Blüthgen, 2010; Cohen et al., 1990; Dormann, 2023; Martinez, 1991; Polis, 1991). Against this, it is important to evaluate the value and the limitations of the various network conceptualisations and how these relate to biodiversity concepts, such as community structure, community stability and ecosystem function (Blüthgen & Staab, 2024). In this perspective we aim to provide an overview of different food web representations, particularly how each representation embeds assumptions about the processes that determine interactions (Section 3) about the levels of organisation at which this occurs (*i.e.* the biological, ecological, spatial/temporal scales) and the way in which we construct the resulting networks (Section 4).

Network construction reflects both the data used and the theories governing species interactions. We still lack a clear explanation of the different assumptions and scale dependent processes that underpin network construction alongside extensive discussions about the challenges relating to data collection and observation (Blüthgen & Staab, 2024; Brimacombe et al., 2023; Brimacombe et al., 2024; Martinez & Dunne, 1998; Moulatlet et al., 2024; Pimm, 1984; Polis, 1991; Pringle & Hutchinson, 2020; Saberski et al., 2024). Such an understanding should deliver an acceleration in capacity to more effectively predict the impact of multiple stressors on biodiverse communities and their functions.

In their recent work, Gauzens et al. (2025) synthesised ecological network representations according to node resolution and link type, highlighting the advantages, limitations,

and methodological considerations associated with different approaches. Here, we take a complementary theory-driven perspective by framing network construction as a hierarchical transition from the feasibility to the realisation of interactions. Rather than classifying networks according to their structural characteristics alone, we focus on how assumptions about ecological entities, interactions, and processes give rise to different representations and the inferences they support. In the following sections, we review how nodes, edges, and interaction processes shape ecological networks, highlighting how these assumptions determine ecological scale, node and link resolution, the relative importance of evolutionary versus ecological processes, and the approaches used to construct networks. We conclude by linking alternative network representations to key questions in biodiversity science in the Anthropocene.

2 Setting the Scene: The Not So Basics of Nodes and Edges

Ecological networks represent an ‘object’ from which inferences can be made and thus serve multiple purposes. While many aspects of community structure can be analysed without networks (*e.g.*, trait distributions or abundance patterns) networks provide a formal framework for capturing the organisation of interactions among species. The study of network structure and topology has a long history in ecology, rooted in early theory on energy flow (Lindeman, 1942; Odum, 1968), and later extended to questions of robustness, stability, and complexity (*e.g.*, (Brose et al., 2006; Dunne et al., 2002; May, 1972; Montoya et al., 2006; Pimm, 1984)). More recent work has built on this foundation to link network structure to ecosystem functioning, persistence, and dynamical behaviour (*e.g.*, (Danet et al., 2024; Pilosof et al., 2017; Schneider et al., 2016)). Networks are therefore commonly treated as response variables in tests of ecological theory and statistical models of the generative processes that give rise to observed structure and are widely used to compare communities across environmental gradients or through time (*e.g.*, Hao et al., 2025; Pecuchet et al., 2020). They also provide a platform for evaluating downstream responses to perturbations, including secondary extinctions and robustness to species loss (*e.g.*, (Dunne et al., 2002; Keyes et al., 2021; Keyes et al., 2024; Staniczenko et al., 2010)), as well as inference about stability, ecosystem function, invasions, climate change, contaminants, and extinction cascades (*e.g.*, (Curtsdotter et al., 2019; Delmas et al., 2017; Terry et al., 2025)). Against this backdrop of multiple research agendas, the

66 definition of ‘edges’ and ‘nodes’, and the levels of organisation at which they are defined,
67 take many forms (Moulatlet et al., 2024; Poisot, Stouffer, et al., 2016), each of which
68 encode a series of assumptions within a network. Here we introduce an alignment between
69 these questions and baseline assumptions about nodes and edges.

70 2.1 How do we define a node?

71 While nodes are conventionally described as representing species, in practice they may
72 correspond to a range of taxonomic and non-taxonomic units, including sub-species,
73 genera, or families, as well as trophic species (*e.g.*, (Williams & Martinez, 2000; Yodzis,
74 1982)), feeding guilds (*e.g.*, (García-Callejas et al., 2023)), or life-stage-specific subsets of
75 species (*e.g.*, (Clegg et al., 2018)). These choices reflect differences in the level, type, and
76 consistency of resolution at which networks are constructed, rather than aggregation *per se*.
77 Representing nodes at coarser or mixed resolutions can limit taxon-specific inference (*e.g.*,
78 whether species *a* consumes species *b*), bias estimates of degree distributions (particularly
79 generality and vulnerability) and complicate downstream analyses such as extinction or
80 invasion dynamics, where species identity and the consequences of loss may be obscured
81 (Beckerman et al., 2006; Clegg et al., 2018). At the same time, there are clear justifications
82 for using aggregated representations when the distribution of interactions among functional
83 or trophic units is more informative than species-level detail, for example when analysing
84 extinction patterns across feeding guilds or other functional units (Dunhill et al., 2024).
85 Issues of resolution, scale, and sampling have long been recognised as central to the
86 construction and interpretation of food webs (*e.g.*, (Dunne, 2006; Martinez & Dunne,
87 1998)). Consequently, node definition is not merely a methodological choice but an
88 assumption about the ecological entities represented in a network, with direct implications
89 for the processes that can be represented and the questions that can be addressed (Gauzens
90 et al., 2025).

91 2.2 What is captured by an edge?

92 Understanding how edge assumptions impact ecological process representation requires
93 distinguishing between *potential* and *realised* links. Potential links reflect interaction
94 feasibility, whereas realised links reflect ecological fluxes such as energy transfer. Links
95 within food webs may therefore represent either potential interactions between species
96 (Dunne, 2006; Pringle, 2020) or realised fluxes within a system, such as the transfer of
97 energy or materials resulting from feeding interactions (Lindeman, 1942; Proulx et al.,

2005). Edges can thus correspond to different ecological ‘currencies’ (Gauzens et al., 2025). Moreover, links can be specified in multiple ways: they may be treated as present or absent (*i.e.*, binary), defined probabilistically (Banville et al., 2025; Poisot, Cirtwill, et al., 2016), or represented by continuous functions that quantify interaction strength (Berlow et al., 2004). Link definition therefore depends on both the ecological currency and how interactions are represented. Because different edge definitions encode different ecological processes, they also support different forms of inference. For example, networks composed of feasible interactions can characterise potential trophic structure but are generally insufficient for inferring the magnitude or distribution of energy flow because their links are not environmentally or energetically constrained.

2.3 Network representations

These definitions of nodes and edges and associated assumptions give rise to two well established types of ecological networks: **metawebs**, which represent all *potential* interactions within a species pool (Dunne, 2006), and **realised networks**, which represent the subset of interactions expressed within a particular community at a given place and time. Because of the assumptions made about nodes and edges in these networks, they differ in the scales they represent and the processes assumed to generate their structure.

A metaweb is fundamentally a list of *feasible* interactions between species. Metawebs identify evolutionarily and regionally plausible interactions, ecologically impossible (*i.e.*, forbidden) links (Jordano, 2016b), and the potential diet breadth of species (Strydom et al., 2023). Feasibility is thus determined by trait complementarity, typically related to feeding, and can be further constrained by species co-occurrence, producing a transition from *global* to *regional* metawebs.

In contrast, realised networks are localised in space and time, with links shaped by co-occurrence, environmental conditions, and diet choice. Even when represented as binary matrices, their links implicitly reflect interaction strength and energy transfer. Realised networks are therefore not simply spatial or temporal subsets of metawebs; they emerge from ecological processes that govern the occurrence and strength of interactions among species. Consequently, a metaweb and realised network containing the same species may differ substantially in structure because link presence is governed by different constraints. Critically, links absent from a metaweb represent infeasible interactions, whereas links absent from a realised network reflect context-dependent ecological constraints. At its most

130 general, a realised network must represent more than the potential for species interactions;
131 it must reflect the processes governing the distribution or strength of interactions within a
132 particular community.

133 **3 From Nodes and Edges to Process and Constraints**

134 In the previous section we discussed how the definition of nodes and edges, representing
135 different scales and processes, lead to the concept of a metaweb and a realised web. The
136 fundamental take-homes are that nodes vary in their resolution, edges vary in what kind
137 of process they represent and the intersection of these, defined by meta- vs. realised webs,
138 underpins distinct lines of inquiry and constraints on the type of inference we can make
139 with networks.

140 Following this, we here reveal five core constraints across evolutionary and ecological scales
141 that further delineate the transition from meta- to realised webs, exposing processes that
142 determine the nature of links among nodes: evolutionary compatibility, co-occurrence,
143 abundance, diet choice, and non-trophic interactions Figure 1.

144 [Figure 1 about here.]

145 **3.1 Processes that determine the feasibility of an interaction**

146 Evolutionary compatibility and co-occurrence are the two principal processes that define
147 the feasibility of an interaction between two species. The scale of inference and set
148 of processes embodied in these two constraints typically combine to define a ‘list’ of
149 interactions that are viable/feasible and defined strictly as present/absent. Reflecting on
150 the previous section, nodes are typically species and rules defining edges are defined by
151 trait complementarity (phylogenetic) and/or co-occurrence. Here we provide more insight
152 into each process.

153 **Evolutionary compatibility**

154 This constraint is defined by shared (co)evolutionary history between consumers and
155 resources (Dalla Riva & Stouffer, 2016; Gómez et al., 2010; Rossberg et al., 2006; Segar et
156 al., 2020) which is manifested as ‘trait complementarity’ between two species (Benadi et
157 al., 2022). In this body of theory, the consumer has the ‘correct’ set of traits that allow it
158 to acquire and consume the resource. Interactions that are not compatible are defined as
159 forbidden links (Jordano, 2016b); *i.e.*, they are not physically possible and will *always* be

absent within a network.

Networks do not explicitly arise from models based on this constraint. Instead, interacting species pairs are defined and these are represented as binary (possible vs forbidden) or probabilistic (Banville et al., 2025). For example, in the metaweb constructed by Strydom et al. (2022) probabilities are quantified as the confidence of a specific interaction being *possible* between two species. A network constructed based on evolutionary compatibility is conceptually aligned with a ‘global metaweb’, and gives us information as to the global feasibility of links between species pairs even though they do not co-occur (see Figure 1).

(Co)occurrence

The co-occurrence of species in both time and space is a fundamental requirement for an interaction between two species to occur (at least in terms of feeding links). Although co-occurrence data alone is insufficient for building an accurate and ecologically meaningful representation of *feeding links* (Blanchet et al., 2020), it is still a critical process that determines the possible realisation of a feeding. Knowledge on the co-occurrence of species allows us to spatially constrain a global metaweb to reflect regional metawebs (Dansereau, Barros, et al., 2024). In the context of Figure 1 this would be the metawebs for regions one and two.

We reinforce that these two constraints don’t deliver a network *per se*, but a list of feasible species pairs. Although it is possible to visualise a network from the list of interactions generated by these constraints, it is important to be aware that the structure of this network is not constrained by any community context - just because species are able to interact does not mean that they will (Caron et al., 2024; Poisot et al., 2015).

3.2 Processes that realise networks

In contrast to the above, here we highlight three processes that influence the *realisation* of an interaction between species and thus form the conceptual basis for realised networks. As we show in Figure 1, a ‘truly realised’ network is the product of properties of the community (**abundance** and **non-trophic interactions**) and the individual (**diet choice**). This represents a conceptual shift from considering the feasibility for species pairwise interactions to considering the edge as a representation of energy flow. Such a transition requires information about how the community, the environment and the individual *constrains* network topology as defined by consumer choice (Quintero et al. (2024), Section 2.3)

Abundance

Abundance as a realising process emerges from a null model for energy acquisition: organisms feeding randomly will consume resources in proportion to their abundance (Stephens & Krebs, 1986). Here, abundance of different prey species influences the distribution of links in a network (Vázquez et al., 2009) by defining a preference linked to individuals among species meeting (Banville et al., 2025; Poisot et al., 2015). Abundance data (linked to a derived metaweb) delivers a foundation ruleset that can define the distribution and strength of links. Of note, however, is that such abundance constrained interactions are not necessarily contingent on there being any compatibility between species (E. Canard et al., 2012; Momal et al., 2020; Pomeranz et al., 2019).

Diet choice

It is well established that consumers make more active decisions than eating items in proportion to their abundance (Stephens & Krebs, 1986). Ultimately, consumer choice is underpinned by an energetic cost-benefit framework centred around profitability and defined by traits associated with acquisition and consumption of a resource (Smith et al., 2021; Wootton et al., 2023). Energetic constraints are invoked to construct networks in a myriad of ways (*e.g.*, Beckerman et al., 2006; Cherif et al., 2024; Pawar et al., 2012; Portalier et al., 2019).

Unlike metaweb approaches, these models generate realised webs as emergent outcomes of consumer behaviour. We also here make a distinction, developed below, with models like the Niche Model (Williams & Martinez, 2000), where diet choice is implicit in its probabilistic network generating function, but it is working to replicate the *expected* structure of the network, and this structure does not emerge from node-based rules. Note that we select diet choice as a term to capture rules linked to optimal foraging (Pyke, 1984) and metabolic theory (Brown et al., 2004) for capturing the energetic constraints on the distribution and strength of interactions.

Non-trophic interactions

We include non-trophic interactions (see Miele et al., 2019) here not as a determinant of links, but a modifier of them - they are the community context above and beyond co-occurrence and abundance. Non-trophic interactions include competition for space, predator interference, refuge provisioning, recruitment facilitation as well as non-trophic effects that increase or decrease mortality. These interactions specifically modify either

the realisation or strength of trophic interactions (Golubski & Abrams, 2011; Ings et al., 2009; Kamaru et al., 2024; Pilosof et al., 2017; Staniczenko et al., 2010) and represent direct (*e.g.*, predator *a* outcompetes predator *b*) and indirect (*e.g.*, mutualistic/facilitative interactions) mechanisms.

Some interactions, such as pollination, occupy an intermediate position in this framework, as they combine trophic components (*e.g.*, resource consumption) with non-trophic effects that influence reproduction, recruitment, and population persistence (Bascompte & Jordano, 2007; Holland et al., 2002; Sauve et al., 2016). They operate on the realisation of a network by altering the fine-scale distribution and abundance of species and relative contributions of direct and indirect effects to biomass, persistence, stability, and the functioning of the communities (Buche et al., 2024; Kéfi et al., 2012; Kéfi et al., 2015; Miele et al., 2019).

4 Network construction

The above five processes are central to understanding the assumptions inherent in building different types of networks. Each of the processes, or combinations thereof, deliver a unique set of boundary conditions on what a network represents and can be used for. Here we build on the introduction of these five processes to further categorise the approaches to constructing networks ultimately moving to showcase how different construction approaches, encoding different assumptions about how interactions arise, can support different ecological questions.

4.1 Why construct networks?

Networks are a representation of biodiversity. In a perfect world, we might know about all interactions. However, the empirical collection of interaction data is both costly and challenging to execute (Jordano, 2016a, 2016b; Poisot et al., 2021). In the absence of robust empirical data, we use ‘models’ that facilitate interpolation and gap-filling of existing empirical datasets (*e.g.*, Biton et al., 2024; Dallas et al., 2017; Poisot et al., 2023; Stock et al., 2017), predict the feasibility of interaction among pairs of species, or directly predict network structure (see Strydom et al., 2021 for a broader discussion).

Networks are unique in delivering more than just estimates of species richness. As noted in the introduction, a network embodies the organising structure of biodiversity and allows numerous opportunities for ‘downstream’ analysis, including the comparison of

structures, estimation of energy flux or extinction dynamics and ultimately form the structural inputs to dynamical systems models that facilitate ecological and conservation relevant inference about productivity-diversity-stability-function relationships (Danet et al., 2024) in space and time. But making such inferences requires careful attention to one or more of the processes discussed in Section 3. While these network representations may be simplifications of the truth (Stouffer, 2019), they remain critically useful tools as they allow us to move beyond simple descriptions of species richness to test hypotheses about community architecture and ecosystem functioning that would be otherwise impossible to assess.

4.2 Construction through induction

Constructing feasible or realised networks can be framed as an ‘inductive reasoning’ process where insight and generalisation arise from a set of observations and relationships around feeding. Inductive reasoning as a foundation for network construction is implemented at node and network levels.

4.2.1 Species specific induction

Species-specific approaches construct networks from the expected feeding interactions between species pairs and are therefore fundamentally creating metawebs. All methods in this inference space rest on a set of three assumptions: there are a set of ‘feeding rules’ that underpin interaction feasibility (Morales-Castilla et al., 2015); these rules are phylogenetically conserved (Bramon Mora et al., 2018; Dalla Riva & Stouffer, 2016); and they can be specified by matching the traits between consumer and resource.

Evolutionary compatibility and co-occurrence constraints have been critical to the construction of ‘first draft’ networks for communities for which we have no interaction data (Strydom et al., 2022). They are also central to interpolation in data poor regions and predicting interactions for ‘unobservable’ communities *e.g.*, prehistoric networks (Dunhill et al., 2024; Fricke et al., 2022; Yeakel et al., 2014) or future, novel community assemblages (Van der Putten et al., 2010). Furthermore, they have the capacity to evaluate a role of interactions among species relative to their distribution by accounting for the role of the environment and the role of species interactions (Albouy et al., 2014; Gravel et al., 2019; Higinio et al., 2023; Pollock et al., 2014).

Feeding rules can be specified in several ways. Expert-based approaches define feasible

interactions using trait matching (Roopnarine, 2017; Shaw et al., 2024), while mechanistic approaches often rely on trait-based relationships between consumers and resources (Gravel et al., 2013; Rohr et al., 2010). Alternatively, statistical and machine-learning methods infer interaction probabilities from observed networks using ecological predictors such as traits and phylogeny (Pichler et al., 2020).

Rules are also defined by correlating real world interaction data with suitable ecological proxies for which data is more widely available (*e.g.*, traits) using some sort of binary classifier (see Pichler et al. (2020) for an overview). These include generalised linear models (*e.g.*, Caron et al., 2022), random forest (*e.g.*, Llewelyn et al., 2023), trait-based k-NN (*e.g.*, Desjardins-Proulx et al., 2017), and Bayesian models (*e.g.*, Eklöf et al., 2013; Cirtwill et al., 2019).

Finally, graph embedding uses the structural features of a known network to infer the position of species in an unknown network through the decomposition of the interaction onto the embedding space (see Strydom et al. (2023)). This decomposition relies on a combination of ecological proxies (*e.g.*, phylogenetic relatedness Strydom et al. (2022)) in conjunction with known interactions to infer the latent values of species, which can then be mapped onto decomposition of a known network.

4.2.2 Structure-based induction

These models generate ecologically realistic network structures using simple probabilistic rules. They are commonly used as null expectations, for comparative analyses, and as inputs to dynamical models. The determination of links between species is not directly linked to properties of the nodes. This means these networks are usually not species specific. Although they require little empirical information, they encode explicit assumptions about expected network structure.

Stochastic network models use a probabilistic ruleset about diet choice and niche breadth to reflect fundamental ideas of foraging biology. These models that are based on the compartmentalisation and acquisition of energy for species at different trophic levels (Allesina & Pascual, 2009; Krause et al., 2003) and that network structure can be determined by distributing interactions along single dimension (the ‘niche axis’; Allesina et al. (2008)). Typically, these models parametrise some aspect of the network structure (although see Allesina & Pascual (2009) for a parameter-free model). These models include the most used network generator, the Niche model (Williams & Martinez, 2000), as well as

the original Cascade model (Cohen et al., 1990) and the derived Nested hierarchy model (Cattin et al., 2004). Even though these networks are derived without any real-world data they are still able to recover the structure of empirical networks (Stouffer et al., 2005). These models often form the basis for dynamic models *e.g.*, the allometric trophic network (Brose et al., 2006; Schneider et al., 2016) and bioenergetic food web models (Delmas et al., 2017).

4.3 Construction through deduction

In contrast to metaweb construction, realised networks require assumptions about the ecological processes that determine whether feasible interactions are expressed. These approaches are deductive because interactions emerge from explicit assumptions about encounters, foraging behaviour, energetic constraints, or interaction modification. These approaches operationalise the abundance, diet-choice, and non-trophic processes introduced in Section 3.1. The resulting networks are widely used to study energy flux, extinction dynamics, and ecosystem functioning. They also provide the structural backbone for dynamical systems modelling to address questions about stability-structure-productivity-function relationships, secondary extinction dynamics, species invasion and climate change. There are two broad groups of models in this deductive category.

These models capture the *behaviour* of the nodes by explicitly considering the properties of the different species in the community. Which means that there is a degree of variance in which links are predicted between species unlike the more ‘static’ predictions made by inductive models. However, these networks are costly to construct in real world settings (requiring data about the entire community, as it is the behaviour of the system that determines the behaviour of the part) and lack the larger diet niche context afforded by metawebs.

4.3.1 Abundance-based models

Neutral networks are built on the assumption that foraging decisions are tied *only* to the abundance of species within the community (E. F. Canard et al., 2014; Krishna et al., 2008). Here links are solely determined by the relative abundance of the different species in the community. Although unrealistic as a complete explanation, neutral models can be combined with inductive approaches to generate more localised predictions (Pomeranz et al., 2019).

349 4.3.2 Energetics-constrained models

350 There is a broader group of models that focus on determining interactions in terms of
351 energetic constraints on diet breadth, often using the ratio of consumer-resource body
352 size as a proxy for capturing the energetic constraints of feeding. Models such as those
353 developed by Portalier et al. (2019) and Wootton et al. (2023) are similar to the mechanistic
354 approaches discussed in Section 4.2, however instead of determining interactions based on
355 mechanistic feasibility it is rather constrained by the energetic cost of predation. Note
356 that although these models do not place any explicit constraints on the expected structure
357 of the network, the links should still be considered as ‘realised’ owing to the energetic
358 constraint placed on links. A different subset of diet models (*e.g.*, Beckerman et al., 2006;
359 Petchey et al., 2008) use a diet choice approach, however like the stochastic network models
360 they also embed assumptions on network structure. Thus, these models predict both
361 interactions and network structure simultaneously, although they may benefit in being
362 refined by more explicitly accounting for trait-based (*i.e.*, feasibility) parameterisation
363 (Curtsdotter et al., 2019).

364 5 Making Progress with Networks

365 The motivation to leverage network ecology in conservation ecology, environmental risk
366 assessment and natural resource management stems from a shift away from species/popu-
367 lation specific measures of the effects of stress and disturbance to community level metrics
368 of these impacts. These metrics, such as resilience and more generally stability, ecosystem
369 function and biodiversity, are natural properties of networks. This suggests that modern
370 conservation, risk assessment and resource management would benefit from robust network
371 tools to support decision making.

372 This is also true in the disciplines of ecology and environmental science and their focus
373 on abundance, distribution, functions and services that biodiversity provides (Loreau &
374 de Mazancourt, 2013). Major questions remain unresolved, for example, about stability-
375 diversity-productivity relationships, the drivers and consequences of extinctions and
376 invasions, and the impacts of multiple stressors operating at multiple ecological scales
377 (Simmons et al., 2021). A network approach to answering these types of questions
378 specifically allows us to evaluate how environmental gradients and anthropogenic stress
379 map through direct and indirect effects among species in a complex community and reveal
380 fundamental patterns and understanding of processes in the natural world.

381 To effectively use networks to aid us in answering questions about conservation/risk
382 assessment/management and core ecological theory, we need to be mindful that we are
383 mapping the *correct* network representation to the question of interest (Gauzens et al.,
384 2025). Notably, there are certain questions that cannot be answered using specific network
385 representations as the scale of the question of interest is fundamentally misaligned with
386 the process captured by a specific network representation (Section 3.1), the underlying
387 data that is used to construct it (Section 4), or both.

388 Here we discuss and map the different network representations shown in Figure 1 to
389 ‘appropriate’ research questions and agendas (see also Table 1). We also highlight some of
390 the key methodological challenges that currently limit our conceptualisation of a ‘network’
391 and thus impact their effective practical application in real world settings.

Table 1: Showcasing some of the broader avenues of inquiry, specifically how they map to the different network representations. Additionally, we highlight some studies that address or present opening discussions around each research question. Superscripts at the research question indicate the strength of the current literature in addressing that research question: $\checkmark$ indicates area with strong foundational research, Δ partial/emerging areas of research, and $\times$ areas where research is weak/largely absent

Network		
Representation	Example Research Question	Representative Studies
Global Metaweb	How will novel communities respond to <i>e.g.</i> , extinction, turnover, invasion and rewilding $\checkmark$	Gravel, Albouy, et al. (2016); Dunne et al. (2002)
	Diet-based conservation focusing not only on the target species but the species it might depend on for food resources Δ	Rooney & McCann (2012); Curtsdotter et al. (2011); McDonald-Madden et al. (2016)
	Rewiring capacity/potential of species by looking at the <i>entire</i> diets of species Δ	Gilljam et al. (2015); Staniczenko et al. (2010); Su et al. (2024); Marjakangas et al. (2025)

Network		
Representation	Example Research Question	Representative Studies
Regional Metawebs	Eco-Evolutionary dynamics and how they relate to the conservation and origination of feeding strategies [×]	Poisot et al. (2015); Baskerville et al. (2011)
	Applied use potential of questions highlighted for global metawebs at the management scale <i>e.g.</i> , a protected area ^Δ	Albouy et al. (2014); Pellissier et al. (2018); Estrada & Bodin (2008)
	Refinement/extension of species distribution models by incorporating co-occurrence and species associations <i>e.g.</i> , predator and prey [✓]	Araújo & Luoto (2007); Kissling et al. (2012)
Realised Webs	The allocation of multiple stressors across networks ^Δ	Crain et al. (2008); Beauchesne et al. (2021)
	Temperature threshold to community collapse ^Δ	O’Gorman et al. (2019); Petchey et al. (2010)
	Extinction and persistence after harvesting/invasion/extinction [✓]	Allesina & Tang (2012); Yodzis (2001)
	Stability-diversity-productivity-function [✓]	Thébault & Fontaine (2010); Rooney & McCann (2012)
	Explicitly tying ecosystem level processes and nutrient flows to networks [×]	Moore et al. (2004)
	Meta communities and the idea of meta-network-communities ^Δ	Gravel, Massol, et al. (2016); Gilarranz et al. (2017)

5.1 Key Eco-Evo-Conservation Questions

5.1.1 Global Metawebs

Global metawebs are most appropriate for questions centred on interaction feasibility and potential diet breadth. They provide a platform for exploring hypothetical or novel communities under climate change (Hui & Richardson, 2019), species invasions, reintroductions, and rewilding, and the potential rewiring capacity (Marjakangas et al., 2025). Because they focus on feasible rather than realised interactions, global metawebs are also well suited to studying eco-evolutionary dynamics and how evolutionary history, natural selection, and phenotypic plasticity shape interaction niches.

5.1.2 Regional Metawebs

Regional metawebs extend these questions by explicitly incorporating species co-occurrence and therefore provide a more management-relevant perspective. They can be used to refine species distribution models and projections of future community composition (García-Callejas et al., 2023; Hao et al., 2025). However, caution is required when comparing regional metaweb structure across space or environmental gradients, as observed differences may reflect species turnover (*e.g.*, β -diversity) rather than changes in interaction processes.

5.1.3 Realised networks

Realised networks are best suited to questions concerning how community and environmental context shape interaction structure and ecosystem functioning (Thuiller et al., 2024). They provide the appropriate framework for studying stability, resilience, biodiversity dynamics, ecosystem functioning, extinction cascades, invasions, climate change impacts, and network rewiring through time. By capturing interactions that are actually occurring, realised networks allow investigation of how perturbations propagate through communities and influence persistence.

The increasing availability of long-term interaction datasets is expanding opportunities to address these questions (Danet et al., 2024; Wooster et al., 2024). However, empirical datasets often accumulate interactions across extended periods, potentially obscuring temporal variation in realised interactions (Polis, 1991). Developing approaches that better reconcile empirical networks with realised community dynamics therefore remains an important challenge.

5.2 Key methodological challenges

As noted above, the three network types highlight longstanding methodological challenges that limit both the precision and accuracy of ecological inference. Here we briefly review these challenges and emerging opportunities to address them.

Understanding what empirical data represents: Robust inference requires understanding what constitutes an observed interaction, whether recorded directly (predation events) or indirectly (*e.g.*, gut contents or stable isotopes). Because empirical networks often accumulate observations across space and time, they may be conceptually closer to metawebs than realised networks.

The validation of network structure: Considerable progress has been made in assessing how well models recover pairwise interactions (Poisot, 2023; Strydom et al., 2021), yet there remains no clear framework for evaluating their ability to recover network structure (Allesina et al., 2008; Tylianakis et al., 2010). This raises two related questions: what constitutes an appropriate benchmark, and which aspects of network recovery matter most? For metawebs, accurately identifying both present and forbidden links may be essential, whereas for realised webs it remains unclear whether recovering pairwise interactions, aggregate properties (*e.g.*, connectance), or both should be the primary objective.

Transitioning between metawebs and realised webs: Most approaches for modelling realised networks do not explicitly incorporate evolutionary constraints (although see Van De Walle et al. (2023); Wootton et al. (2023)). Progress will likely require either ensemble approaches (Becker et al., 2022; Terry & Lewis, 2020) or methods for downscaling metawebs into realised networks (*e.g.*, Roopnarine, 2006). However, any such transition must retain clarity about the meaning of links - structurally realistic networks do not necessarily represent realised prey choice or energy flow.

Developing frameworks that allow transitioning between metawebs and realised representations will also facilitate integration with metacommunity, metaecosystem, and ecosystem-level theory (Liu et al., 2025; Massol et al., 2011). Doing so requires that we develop an understanding of what the appropriate spatial and temporal scales of networks are (Fortin et al., 2021), and specifically how these relate to network type and the question of interest.

Making networks more tractable in applied spaces: The application of networks to conservation and management remains limited by difficulties in defining network

boundaries, aligning them with management units, and interpreting network metrics in policy-relevant ways (Dansereau, Braga, et al., 2024). Addressing these challenges will require stronger links between network structure and ecosystem function, alongside careful matching of analytical tools and network representations to management objectives (O'Connor et al., 2025; Pellissier et al., 2018).

Taken together, these challenges highlight three overarching messages. (i) Network representations are inseparable from the data and assumptions used to construct them; (ii) validation and benchmarking must be aligned with the intended network type and research question; and (iii) greater conceptual clarity is needed when transitioning between metawebs and realised networks, particularly in applied contexts. Explicitly articulating these distinctions is essential if networks are to be used rigorously and transparently across scales.

6 Concluding Remarks

Ecological network representations provide alternative abstractions of biodiversity, each defined by specific assumptions regarding ecological entities, interactions, and processes. Consequently, the suitability of a given representation depends on the ecological questions being addressed and the processes that the available data can meaningfully capture. Recognising these differences is essential for interpreting network structure, evaluating the scope of inference, and avoiding implicit assumptions associated with conventional approaches. By linking network representations to their underlying ecological processes and data requirements, the framework developed here provides a basis for systematically comparing alternative network constructions and identifying the contexts in which they are most informative. Establishing this standard is essential for preventing the misinterpretation of network data and for addressing foundational questions in biodiversity science.

References

- Albouy, C., Velez, L., Coll, M., Colloca, F., Le Loc'h, F., Mouillot, D., & Gravel, D. (2014). From projected species distribution to food-web structure under climate change. *Global Change Biology*, 20(3), 730–741. <https://doi.org/10.1111/gcb.12467>
- Allesina, S., Alonso, D., & Pascual, M. (2008). A General Model for Food Web Structure. *Science*, 320(5876), 658–661. <https://doi.org/10.1126/science.1156269>

Allesina, S., & Pascual, M. (2009). Food web models: A plea for groups. *Ecology Letters*, 12(7), 652–662. <https://doi.org/10.1111/j.1461-0248.2009.01321.x>

Allesina, S., & Tang, S. (2012). Stability criteria for complex ecosystems. *Nature*, 483(7388, 7388), 205–208. <https://doi.org/10.1038/nature10832>

Araújo, M. B., & Luoto, M. (2007). The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography*, 16(6), 743–753. <https://doi.org/10.1111/j.1466-8238.2007.00359.x>

Banville, F., Strydom, T., Blyth, P. S. A., Brimacombe, C., Catchen, M. D., Dansereau, G., Higino, G., Malpas, T., Mayall, H., Norman, K., Gravel, D., & Poisot, T. (2025). Deciphering Probabilistic Species Interaction Networks. *Ecology Letters*, 28(6), e70161. <https://doi.org/10.1111/ele.70161>

Bascompte, J., & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The Architecture of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 38(1), 567–593. <https://doi.org/10.1146/annurev.ecolsys.38.091206.095818>

Baskerville, E. B., Dobson, A. P., Bedford, T., Allesina, S., Anderson, T. M., & Pascual, M. (2011). Spatial Guilds in the Serengeti Food Web Revealed by a Bayesian Group Model. *PLOS Computational Biology*, 7(12), e1002321. <https://doi.org/10.1371/journal.pcbi.1002321>

Beauchesne, D., Cazelles, K., Archambault, P., Dee, L. E., & Gravel, D. (2021). On the sensitivity of food webs to multiple stressors. *Ecology Letters*, 24(10), 2219–2237. <https://doi.org/10.1111/ele.13841>

Becker, D. J., Albery, G. F., Sjödin, A. R., Poisot, T., Bergner, L. M., Chen, B., Cohen, L. E., Dallas, T. A., Eskew, E. A., Fagre, A. C., Farrell, M. J., Guth, S., Han, B. A., Simmons, N. B., Stock, M., Teeling, E. C., & Carlson, C. J. (2022). Optimising predictive models to prioritise viral discovery in zoonotic reservoirs. *The Lancet Microbe*, 3(8), e625–e637. [https://doi.org/10.1016/S2666-5247\(21\)00245-7](https://doi.org/10.1016/S2666-5247(21)00245-7)

Beckerman, A. P., Petchey, O. L., & Warren, P. H. (2006). Foraging biology predicts food web complexity. *Proceedings of the National Academy of Sciences*, 103(37), 13745–13749. <https://doi.org/10.1073/pnas.0603039103>

Benadi, G., Dormann, C. F., Fründ, J., Stephan, R., & Vázquez, D. P. (2022). Quantitative Prediction of Interactions in Bipartite Networks Based on Traits, Abundances, and Phylogeny. *The American Naturalist*, 199(6), 841–854. <https://doi.org/10.1086/714420>

Berlow, E. L., Neutel, A.-M., Cohen, J. E., de Ruiter, P. C., Ebenman, B., Emmerson, M., Fox, J. W., Jansen, V. A. A., Iwan Jones, J., Kokkoris, G. D., Logofet, D. O.,

- McKane, A. J., Montoya, J. M., & Petchey, O. (2004). Interaction strengths in food webs: Issues and opportunities. *Journal of Animal Ecology*, 73(3), 585–598. <https://doi.org/10.1111/j.0021-8790.2004.00833.x>
- Biton, B., Puzis, R., & Pilosof, S. (2024, August 2). *Inductive link prediction boosts data availability and enables cross-community link prediction in ecological networks*. <https://ecoevorxiv.org/repository/view/7492/>
- Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23(7), 1050–1063. <https://doi.org/10.1111/ele.13525>
- Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: A critique and an ecologist’s guide. *Basic and Applied Ecology*, 11(3), 185–195. <https://doi.org/10.1016/j.baae.2010.01.001>
- Blüthgen, N., & Staab, M. (2024). A Critical Evaluation of Network Approaches for Studying Species Interactions. *Annual Review of Ecology, Evolution, and Systematics*, 55(1), 65–88. <https://doi.org/10.1146/annurev-ecolsys-102722-021904>
- Bramon Mora, B., Gravel, D., Gilarranz, L. J., Poisot, T., & Stouffer, D. B. (2018). Identifying a common backbone of interactions underlying food webs from different ecosystems. *Nature Communications*, 9(1), 2603. <https://doi.org/10.1038/s41467-018-05056-0>
- Brimacombe, C., Bodner, K., Gravel, D., Leroux, S. J., Poisot, T., & Fortin, M.-J. (2024). Publication-driven consistency in food web structures: Implications for comparative ecology. *Ecology*, n/a(n/a), e4467. <https://doi.org/10.1002/ecy.4467>
- Brimacombe, C., Bodner, K., Michalska-Smith, M., Poisot, T., & Fortin, M.-J. (2023). Shortcomings of reusing species interaction networks created by different sets of researchers. *PLOS Biology*, 21(4), e3002068. <https://doi.org/10.1371/journal.pbio.3002068>
- Brose, U., Williams, R. J., & Martinez, N. D. (2006). Allometric scaling enhances stability in complex food webs. *Ecology Letters*, 9(11), 1228–1236. <https://doi.org/10.1111/j.1461-0248.2006.00978.x>
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a Metabolic Theory of Ecology. *Ecology*, 85(7), 1771–1789. <https://doi.org/10.1890/03-9000>
- Buche, L., Bartomeus, I., & Godoy, O. (2024). Multitrophic Higher-Order Interactions Modulate Species Persistence. *The American Naturalist*, 203(4), 458–472. <https://doi.org/10.1093/aes/203.4.458>

552 [//doi.org/10.1086/729222](https://doi.org/10.1086/729222)

553 Canard, E. F., Mouquet, N., Mouillot, D., Stanko, M., Miklisova, D., & Gravel, D. (2014).
 554 Empirical Evaluation of Neutral Interactions in Host-Parasite Networks. *The American*
 555 *Naturalist*, 183(4), 468–479. <https://doi.org/10.1086/675363>

556 Canard, E., Mouquet, N., Marescot, L., Gaston, K. J., Gravel, D., & Mouillot, D. (2012).
 557 Emergence of Structural Patterns in Neutral Trophic Networks. *PLOS ONE*, 7(8),
 558 e38295. <https://doi.org/10.1371/journal.pone.0038295>

559 Caron, D., Brose, U., Lurgi, M., Blanchet, F. G., Gravel, D., & Pollock, L. J. (2024). Trait-
 560 matching models predict pairwise interactions across regions, not food web properties.
 561 *Global Ecology and Biogeography*, 33(4), e13807. <https://doi.org/10.1111/geb.13807>

562 Caron, D., Maiorano, L., Thuiller, W., & Pollock, L. J. (2022). Addressing the Eltonian
 563 shortfall with trait-based interaction models. *Ecology Letters*, 25(4), 889–899. <https://doi.org/10.1111/ele.13966>

564 [//doi.org/10.1111/ele.13966](https://doi.org/10.1111/ele.13966)

565 Cattin, M.-F., Bersier, L.-F., Banašek-Richter, C., Baltensperger, R., & Gabriel, J.-P.
 566 (2004). Phylogenetic constraints and adaptation explain food-web structure. *Nature*,
 567 427(6977, 6977), 835–839. <https://doi.org/10.1038/nature02327>

568 Cherif, M., Brose, U., Hirt, M. R., Ryser, R., Silve, V., Albert, G., Arnott, R., Berti, E.,
 569 Cirtwill, A., Dyer, A., Gauzens, B., Gupta, A., Ho, H.-C., Portalier, S. M. J., Wain,
 570 D., & Wootton, K. (2024). The environment to the rescue: Can physics help predict
 571 predator–prey interactions? *Biological Reviews*, 138(1). [https://doi.org/10.1111/brv.](https://doi.org/10.1111/brv.13105)
 572 [13105](https://doi.org/10.1111/brv.13105)

573 Cirtwill, A. R., Ekl f, A., Roslin, T., Wootton, K., & Gravel, D. (2019). A quantitative
 574 framework for investigating the reliability of empirical network construction. *Methods*
 575 *in Ecology and Evolution*, 10(6), 902–911. <https://doi.org/10.1111/2041-210X.13180>

576 Clegg, T., Ali, M., & Beckerman, A. P. (2018). The impact of intraspecific variation on
 577 food web structure. *Ecology*, 99(12), 2712–2720. <https://doi.org/10.1002/ecy.2523>

578 Cohen, J. E., Briand, F., & Newman, C. (1990). *Community Food Webs: Data and Theory*.
 579 Springer-Verlag. <https://www.springer.com/gp/book/9783642837869>

580 Crain, C. M., Kroeker, K., & Halpern, B. S. (2008). Interactive and cumulative effects
 581 of multiple human stressors in marine systems. *Ecology Letters*, 11(12), 1304–1315.
 582 <https://doi.org/10.1111/j.1461-0248.2008.01253.x>

583 Curtsdotter, A., Banks, H. T., Banks, J. E., Jonsson, M., Jonsson, T., Laubmeier, A. N.,
 584 Traugott, M., & Bommarco, R. (2019). Ecosystem function in predator–prey food
 585 webs—confronting dynamic models with empirical data. *Journal of Animal Ecology*,

88(2), 196–210. <https://doi.org/10.1111/1365-2656.12892>

Curtsdotter, A., Binzer, A., Brose, U., De Castro, F., Ebenman, B., Eklöf, A., Riede, J. O., Thierry, A., & Rall, B. C. (2011). Robustness to secondary extinctions: Comparing trait-based sequential deletions in static and dynamic food webs. *Basic and Applied Ecology*, 12(7), 571–580. <https://doi.org/10.1016/j.baae.2011.09.008>

Dalla Riva, G. V., & Stouffer, D. B. (2016). Exploring the evolutionary signature of food webs’ backbones using functional traits. *Oikos*, 125(4), 446–456. <https://doi.org/10.1111/oik.02305>

Dallas, T., Park, A. W., & Drake, J. M. (2017). Predicting cryptic links in host-parasite networks. *PLOS Computational Biology*, 13(5), e1005557. <https://doi.org/10.1371/journal.pcbi.1005557>

Danet, A., Kéfi, S., Johnson, T. F., & Beckerman, A. P. (2024, August 30). *Response diversity is a major driver of temporal stability in complex food webs*. <https://doi.org/10.1101/2024.08.29.610288> (Pre-published)

Dansereau, G., Barros, C., & Poisot, T. (2024). Spatially explicit predictions of food web structure from regional-level data. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 379(1909). <https://doi.org/10.1098/rstb.2023.0166>

Dansereau, G., Braga, J., Ficetola, G. F., Galiana, N., Gravel, D., Maiorano, L., Montoya, J. M., O’Connor, L., Pollock, L. J., Thuiller, W., Poisot, T., & Barros, C. (2024). *Overcoming the disconnect between interaction networks and biodiversity conservation and management*. <https://ecoevorxiv.org/repository/view/8134/>

Delmas, E., Brose, U., Gravel, D., Stouffer, D. B., & Poisot, T. (2017). Simulations of biomass dynamics in community food webs. *Methods in Ecology and Evolution*, 8(7), 881–886. <https://doi.org/10.1111/2041-210X.12713>

Desjardins-Proulx, P., Laigle, I., Poisot, T., & Gravel, D. (2017). Ecological interactions and the Netflix problem. *PeerJ*, 5, e3644. <https://doi.org/10.7717/peerj.3644>

Dormann, C. F. (2023). The rise, and possible fall, of network ecology. In *Defining Agroecology – A Festschrift for Teja Tscharnkte* (pp. 143–159). Tredition.

Dunhill, A. M., Zarzychny, K., Shaw, J. O., Atkinson, J. W., Little, C. T. S., & Beckerman, A. P. (2024). Extinction cascades, community collapse, and recovery across a Mesozoic hyperthermal event. *Nature Communications*, 15(1), 8599. <https://doi.org/10.1038/s41467-024-53000-2>

Dunne, J. A. (2006). The Network Structure of Food Webs. In J. A. Dunne & M. Pascual (Eds.), *Ecological networks: Linking structure and dynamics* (pp. 27–86). Oxford

University Press.

Dunne, J. A., Williams, R. J., & Martinez, N. D. (2002). Network structure and biodiversity loss in food webs: Robustness increases with connectance. *Ecology Letters*, 5(4), 558–567. <https://doi.org/10.1046/j.1461-0248.2002.00354.x>

Eklöf, A., Tang, S., & Allesina, S. (2013). Secondary extinctions in food webs: A Bayesian network approach. *Methods in Ecology and Evolution*, 4(8), 760–770. <https://doi.org/10.1111/2041-210X.12062>

Elton, C. S. (2001). *Animal Ecology* (M. A. Leibold & T. J. Wootton, Eds.). University of Chicago Press. <https://press.uchicago.edu/ucp/books/book/chicago/A/bo25281897.html>

Estrada, E., & Bodin, Ö. (2008). Using Network Centrality Measures to Manage Landscape Connectivity. *Ecological Applications*, 18(7), 1810–1825. <https://doi.org/10.1890/07-1419.1>

Fortin, M.-J., Dale, M. R. T., & Brimacombe, C. (2021). Network ecology in dynamic landscapes. *Proceedings of the Royal Society B: Biological Sciences*, 288(1949), rspb.2020.1889, 20201889. <https://doi.org/10.1098/rspb.2020.1889>

Fricke, E. C., Hsieh, C., Middleton, O., Gorczynski, D., Cappello, C. D., Sanisidro, O., Rowan, J., Svenning, J.-C., & Beaudrot, L. (2022). Collapse of terrestrial mammal food webs since the Late Pleistocene. *Science*, 377(6609), 1008–1011. <https://doi.org/10.1126/science.abn4012>

Galiana, N., & Araújo, M. B. (2026). Biotic interactions biogeography: A framework for understanding how species interactions shape biodiversity patterns across scales. *PLOS Biology*, 24(6), e3003813. <https://doi.org/10.1371/journal.pbio.3003813>

García-Callejas, D., Godoy, O., Buche, L., Hurtado, M., Lanuza, J. B., Allen-Perkins, A., & Bartomeus, I. (2023). Non-random interactions within and across guilds shape the potential to coexist in multi-trophic ecological communities. *Ecology Letters*, 26(6), 831–842. <https://doi.org/10.1111/ele.14206>

Gauzens, B., Thouvenot, L., Srivastava, D. S., Kratina, P., Romero, G. Q., Berti, E., O’Gorman, E. J., González, A. L., Dézerald, O., Eisenhauer, N., Pires, M., Ryser, R., Farjalla, V. F., Rogy, P., Brose, U., Petermann, J. S., Geslin, B., & Hines, J. (2025). Tailoring interaction network types to answer different ecological questions. *Nature Reviews Biodiversity*, 1–10. <https://doi.org/10.1038/s44358-025-00056-7>

Gilarranz, L. J., Rayfield, B., Liñán-Cembrano, G., Bascompte, J., & Gonzalez, A. (2017). Effects of network modularity on the spread of perturbation impact in experimental

metapopulations. *Science (New York, N.Y.)*, 357(6347), 199–201. <https://doi.org/10.1126/science.aal4122>

Gilljam, D., Curtsdotter, A., & Ebenman, B. (2015). Adaptive rewiring aggravates the effects of species loss in ecosystems. *Nature Communications*, 6(1), 8412. <https://doi.org/10.1038/ncomms9412>

Golubski, A. J., & Abrams, P. A. (2011). Modifying modifiers: What happens when interspecific interactions interact? *Journal of Animal Ecology*, 80(5), 1097–1108. <https://doi.org/10.1111/j.1365-2656.2011.01852.x>

Gómez, J. M., Verdú, M., & Perfectti, F. (2010). Ecological interactions are evolutionarily conserved across the entire tree of life. *Nature*, 465(7300, 7300), 918–921. <https://doi.org/10.1038/nature09113>

Gravel, D., Albouy, C., & Thuiller, W. (2016). The meaning of functional trait composition of food webs for ecosystem functioning. *Philosophical Transactions: Biological Sciences*, 371(1694), 1–14. <http://www.jstor.org/stable/24768743>

Gravel, D., Baiser, B., Dunne, J. A., Kopelke, J.-P., Martinez, N. D., Nyman, T., Poisot, T., Stouffer, D. B., Tylianakis, J. M., Wood, S. A., & Roslin, T. (2019). Bringing Elton and Grinnell together: A quantitative framework to represent the biogeography of ecological interaction networks. *Ecography*, 42(3), 401–415. <https://doi.org/10.1111/ecog.04006>

Gravel, D., Massol, F., & Leibold, M. A. (2016). Stability and complexity in model metaecosystems. *Nature Communications*, 7, 12457. <https://doi.org/10.1038/ncomms12457>

Gravel, D., Poisot, T., Albouy, C., Velez, L., & Mouillot, D. (2013). Inferring food web structure from predator–prey body size relationships. *Methods in Ecology and Evolution*, 4(11), 1083–1090. <https://doi.org/10.1111/2041-210X.12103>

Hao, X., Holyoak, M., Zhang, Z., & Yan, C. (2025). Global Projection of Terrestrial Vertebrate Food Webs Under Future Climate and Land-Use Changes. *Global Change Biology*, 31(2), e70061. <https://doi.org/10.1111/gcb.70061>

Higino, G. T., Banville, F., Dansereau, G., Muñoz, N. R. F., Windsor, F., & Poisot, T. (2023). Mismatch between IUCN range maps and species interactions data illustrated using the Serengeti food web. *PeerJ*, 11, e14620. <https://doi.org/10.7717/peerj.14620>

Holland, J. N., DeAngelis, D. L., & Bronstein, J. L. (2002). Population Dynamics and Mutualism: Functional Responses of Benefits and Costs. *The American Naturalist*, 159(3), 231–244. <https://doi.org/10.1086/338510>

Hui, C., & Richardson, D. M. (2019). How to Invade an Ecological Network. *Trends in Ecology & Evolution*, 34(2), 121–131. <https://doi.org/10.1016/j.tree.2018.11.003>

Ings, T. C., Montoya, J. M., Bascompte, J., Blüthgen, N., Brown, L., Dormann, C. F.,
 Edwards, F., Figueroa, D., Jacob, U., Jones, J. I., Lauridsen, R. B., Ledger, M. E.,
 Lewis, H. M., Olesen, J. M., van Veen, F. J. F., Warren, P. H., & Woodward, G.
 (2009). Ecological networks—beyond food webs. *The Journal of Animal Ecology*, 78(1),
 253–269. <https://doi.org/10.1111/j.1365-2656.2008.01460.x>

Jordano, P. (2016a). Chasing Ecological Interactions. *PLOS Biology*, 14(9), e1002559.
<https://doi.org/10.1371/journal.pbio.1002559>

Jordano, P. (2016b). Sampling networks of ecological interactions. *Functional Ecology*.
<https://doi.org/10.1111/1365-2435.12763>

Kamaru, D. N., Palmer, T. M., Riginos, C., Ford, A. T., Belnap, J., Chira, R. M., Githaiga,
 J. M., Gituku, B. C., Hays, B. R., Kavwele, C. M., Kibungei, A. K., Lamb, C. T.,
 Maiyo, N. J., Milligan, P. D., Mutisya, S., Ng’weno, C. C., Ogutu, M., Pietrek, A.
 G., Wildt, B. T., & Goheen, J. R. (2024). Disruption of an ant-plant mutualism
 shapes interactions between lions and their primary prey. *Science*, 383(6681), 433–438.
<https://doi.org/10.1126/science.adg1464>

Kéfi, S., Berlow, E. L., Wieters, E. A., Joppa, L. N., Wood, S. A., Brose, U., & Navarrete,
 S. A. (2015). Network structure beyond food webs: Mapping non-trophic and trophic
 interactions on Chilean rocky shores. *Ecology*, 96(1), 291–303. <https://doi.org/10.1890/13-1424.1>

Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A.,
 Boit, A., Joppa, L. N., Lafferty, K. D., Williams, R. J., Martinez, N. D., Menge,
 B. A., Blanchette, C. A., Iles, A. C., & Brose, U. (2012). More than a meal...
 integrating non-feeding interactions into food webs. *Ecology Letters*, 15(4), 291–300.
<https://doi.org/10.1111/j.1461-0248.2011.01732.x>

Keyes, A. A., Barner, A. K., & Dee, L. E. (2024). Synthesising the Relationships
 Between Food Web Structure and Robustness. *Ecology Letters*, 27(10), e14533.
<https://doi.org/10.1111/ele.14533>

Keyes, A. A., McLaughlin, J. P., Barner, A. K., & Dee, L. E. (2021). An ecological
 network approach to predict ecosystem service vulnerability to species losses. *Nature*
Communications, 12(1), 1586. <https://doi.org/10.1038/s41467-021-21824-x>

Kissling, W. D., Dormann, C. F., Groeneveld, J., Hickler, T., Kühn, I., McInerny, G. J.,
 Montoya, J. M., Römermann, C., Schippers, K., Schurr, F. M., Singer, A., Svenning,
 J.-C., Zimmermann, N. E., & O’Hara, R. B. (2012). Towards novel approaches
 to modelling biotic interactions in multispecies assemblages at large spatial extents.

722 *Journal of Biogeography*, 39(12), 2163–2178. [https://doi.org/10.1111/j.1365-2699.2011.](https://doi.org/10.1111/j.1365-2699.2011.02663.x)
723 [02663.x](https://doi.org/10.1111/j.1365-2699.2011.02663.x)

724 Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E., & Taylor, W. W. (2003).
725 Compartments revealed in food-web structure. *Nature*, 426(6964), 282–285. <https://doi.org/10.1038/nature02115>
726 [//doi.org/10.1038/nature02115](https://doi.org/10.1038/nature02115)

727 Krishna, A., Guimarães Jr, P. R., Jordano, P., & Bascompte, J. (2008). A neutral-niche
728 theory of nestedness in mutualistic networks. *Oikos*, 117(11), 1609–1618. <https://doi.org/10.1111/j.1600-0706.2008.16540.x>
729 [//doi.org/10.1111/j.1600-0706.2008.16540.x](https://doi.org/10.1111/j.1600-0706.2008.16540.x)

730 Lindeman, R. L. (1942). The Trophic-Dynamic Aspect of Ecology. *Ecology*, 23(4), 399–417.
731 <https://doi.org/10.2307/1930126>

732 Liu, J., Liu, M., Yin, J., Zhang, Y., Guo, Y., Yu, T., Yu, H., & Ming, X. (2025).
733 Harnessing food webs as a framework for enhancing ecosystem multifunctionality.
734 *Biological Conservation*, 310, 111383. <https://doi.org/10.1016/j.biocon.2025.111383>

735 Llewelyn, J., Strona, G., Dickman, C. R., Greenville, A. C., Wardle, G. M., Lee, M. S.
736 Y., Doherty, S., Shabani, F., Saltr  , F., & Bradshaw, C. J. A. (2023). Predicting
737 predator–prey interactions in terrestrial endotherms using random forest. *Ecography*,
738 2023(9), e06619. <https://doi.org/10.1111/ecog.06619>

739 Loreau, M., & de Mazancourt, C. (2013). Biodiversity and ecosystem stability: A synthesis
740 of underlying mechanisms. *Ecology Letters*, 16(s1), 106–115. [https://doi.org/10.1111/](https://doi.org/10.1111/ele.12073)
741 [ele.12073](https://doi.org/10.1111/ele.12073)

742 MacArthur, R. (1955). Fluctuations of Animal Populations and a Measure of Community
743 Stability. *Ecology*, 36(3), 533–536. <https://doi.org/10.2307/1929601>

744 Marjakangas, E.-L., Dalsgaard, B., & Ordonez, A. (2025). Fundamental Interaction Niches:
745 Towards a Functional Understanding of Ecological Networks’ Resilience. *Ecology Letters*,
746 28(6), e70146. <https://doi.org/10.1111/ele.70146>

747 Martinez, N. D. (1991). Artifacts or Attributes? Effects of Resolution on the Little Rock
748 Lake Food Web. *Ecological Monographs*, 61(4), 367–392. [https://doi.org/10.2307/](https://doi.org/10.2307/2937047)
749 [2937047](https://doi.org/10.2307/2937047)

750 Martinez, N. D., & Dunne, J. A. (1998). Time, space, and beyond: Scale issues in food-web
751 research. In *Ecological Scale: Theory and Applications* (pp. 207–226). Columbia
752 University Press.

753 Massol, F., Gravel, D., Mouquet, N., Cadotte, M. W., Fukami, T., & Leibold, M. A.
754 (2011). Linking community and ecosystem dynamics through spatial ecology. *Ecology*
755 *Letters*, 14(3), 313–323. <https://doi.org/10.1111/j.1461-0248.2011.01588.x>

- May, R. M. (1972). Will a large complex system be stable? *Nature*, 238(5364), 413–414.
<https://doi.org/10.1038/238413a0>
- McDonald-Madden, E., Sabbadin, R., Game, E. T., Baxter, P. W. J., Chadès, I., & Possingham, H. P. (2016). Using food-web theory to conserve ecosystems. *Nature Communications*, 7(1), 10245. <https://doi.org/10.1038/ncomms10245>
- Miele, V., Guill, C., Ramos-Jiliberto, R., & Kéfi, S. (2019). Non-trophic interactions strengthen the diversity—functioning relationship in an ecological bioenergetic network model. *PLOS Computational Biology*, 15(8), e1007269. <https://doi.org/10.1371/journal.pcbi.1007269>
- Momal, R., Robin, S., & Ambroise, C. (2020). Tree-based inference of species interaction networks from abundance data. *Methods in Ecology and Evolution*, 11(5), 621–632. <https://doi.org/10.1111/2041-210X.13380>
- Montoya, J. M., Pimm, S. L., & Solé, R. V. (2006). Ecological networks and their fragility. *Nature*, 442(7100, 7100), 259–264. <https://doi.org/10.1038/nature04927>
- Moore, J. C., Berlow, E. L., Coleman, D. C., de Ruiter, P. C., Dong, Q., Hastings, A., Johnson, N. C., McCann, K. S., Melville, K., Morin, P. J., Nadelhoffer, K., Rosemond, A. D., Post, D. M., Sabo, J. L., Scow, K. M., Vanni, M. J., & Wall, D. H. (2004). Detritus, trophic dynamics and biodiversity. *Ecology Letters*, 7(7), 584–600. <https://doi.org/10.1111/j.1461-0248.2004.00606.x>
- Morales-Castilla, I., Matias, M. G., Gravel, D., & Araújo, M. B. (2015). Inferring biotic interactions from proxies. *Trends in Ecology & Evolution*, 30(6), 347–356. <https://doi.org/10.1016/j.tree.2015.03.014>
- Moulatlet, G., Luna, P., Dattilo, W., & Villalobos, F. (2024). *The scaling of trophic specialization in interaction networks across levels of organization*. <https://doi.org/10.22541/au.172977303.33335171/v1>
- O’Connor, L. M. J., Thuiller, W., Brose, U., Chenevois, É., Freund, C., Gauzens, B., Güzere, P., Graham, C., Harfoot, M., Hirt, M. R., Lavergne, S., Maiorano, L., Moilanen, A., Verburg, P. H., Visconti, P., & Pollock, L. J. (2025). *The untapped potential of food webs in systematic conservation planning*. <https://hal.science/hal-05331872>
- O’Gorman, E. J., Petchey, O. L., Faulkner, K. J., Gallo, B., Gordon, T. A. C., Neto-Cerejeira, J., Ólafsson, J. S., Pichler, D. E., Thompson, M. S. A., & Woodward, G. (2019). A simple model predicts how warming simplifies wild food webs. *Nature Climate Change*, 9(8, 8), 611–616. <https://doi.org/10.1038/s41558-019-0513-x>

790 Odum, E. P. (1968). Energy Flow in Ecosystems: A Historical Review. *American Zoologist*,
791 8(1), 11–18. <https://doi.org/10.1093/icb/8.1.11>

792 Pawar, S., Dell, A. I., & Savage, V. M. (2012). Dimensionality of consumer search space
793 drives trophic interaction strengths. *Nature*, 486(7404), 485–489. <https://doi.org/10.1038/nature11131>
794

795 Pecuchet, L., Blanchet, M.-A., Frainer, A., Husson, B., Jørgensen, L. L., Kortsch, S.,
796 & Primicerio, R. (2020). Novel feeding interactions amplify the impact of species
797 redistribution on an Arctic food web. *Global Change Biology*, 26(9), 4894–4906.
798 <https://doi.org/10.1111/gcb.15196>

799 Pellissier, L., Albouy, C., Bascompte, J., Farwig, N., Graham, C., Loreau, M., Maglianesi,
800 M. A., Melián, C. J., Pitteloud, C., Roslin, T., Rohr, R., Saavedra, S., Thuiller,
801 W., Woodward, G., Zimmermann, N. E., & Gravel, D. (2018). Comparing species
802 interaction networks along environmental gradients. *Biological Reviews*, 93(2), 785–800.
803 <https://doi.org/10.1111/brev.12366>

804 Petchey, O. L., Beckerman, A. P., Riede, J. O., & Warren, P. H. (2008). Size, foraging,
805 and food web structure. *Proceedings of the National Academy of Sciences*, 105(11),
806 4191–4196. <https://doi.org/10.1073/pnas.0710672105>

807 Petchey, O. L., Brose, U., & Rall, B. C. (2010). Predicting the effects of temperature on
808 food web connectance. *Philosophical Transactions of the Royal Society B: Biological
809 Sciences*, 365(1549), 2081–2091. <https://doi.org/10.1098/rstb.2010.0011>

810 Pichler, M., Boreux, V., Klein, A.-M., Schleuning, M., & Hartig, F. (2020). Machine
811 learning algorithms to infer trait-matching and predict species interactions in ecological
812 networks. *Methods in Ecology and Evolution*, 11(2), 281–293. <https://doi.org/10.1111/2041-210X.13329>
813

814 Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological
815 networks. *Nature Ecology & Evolution*, 1(4), 101. <https://doi.org/10.1038/s41559-017-0101>
816

817 Pimm, S. L. (1984). The complexity and stability of ecosystems. *Nature*, 307(5949),
818 321–326. <https://doi.org/10.1038/307321a0>

819 Poisot, T. (2023). Guidelines for the prediction of species interactions through binary
820 classification. *Methods in Ecology and Evolution*, 14(5), 1333–1345. <https://doi.org/10.1111/2041-210X.14071>
821

822 Poisot, T., Bergeron, G., Cazelles, K., Dallas, T., Gravel, D., MacDonald, A., Mercier, B.,
823 Violet, C., & Vissault, S. (2021). Global knowledge gaps in species interaction networks

824 data. *Journal of Biogeography*, 48(7), 1552–1563. <https://doi.org/10.1111/jbi.14127>

825 Poisot, T., Cirtwill, A., Cazelles, K., Gravel, D., Fortin, M.-J., & Stouffer, D. (2016). The
826 structure of probabilistic networks. *Methods in Ecology and Evolution*, 7(3), 303–312.
827 <https://doi.org/10>

828 Poisot, T., Ouellet, M.-A., Mollentze, N., Farrell, M. J., Becker, D. J., Brierley, L., Albery,
829 G. F., Gibb, R. J., Seifert, S. N., & Carlson, C. J. (2023). Network embedding
830 unveils the hidden interactions in the mammalian virome. *Patterns*, 4(6), 100738.
831 <https://doi.org/10.1016/j.patter.2023.100738>

832 Poisot, T., Stouffer, D. B., & Gravel, D. (2015). Beyond species: Why ecological interaction
833 networks vary through space and time. *Oikos*, 124(3), 243–251. <https://doi.org/10.1111/oik.01719>

834
835 Poisot, T., Stouffer, D. B., & Kéfi, S. (2016). Describe, understand and predict: Why
836 do we need networks in ecology? *Functional Ecology*, 30(12), 1878–1882. <https://www.jstor.org/stable/48582345>

837
838 Polis, G. A. (1991). Complex Trophic Interactions in Deserts: An Empirical Critique of
839 Food-Web Theory. *The American Naturalist*, 138(1), 123–155. <https://www.jstor.org/stable/2462536>

840
841 Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., Vesk,
842 P. A., & McCarthy, M. A. (2014). Understanding co-occurrence by modelling species
843 simultaneously with a Joint Species Distribution Model (JSDM). *Methods in Ecology*
844 *and Evolution*, 5(5), 397–406. <https://doi.org/10.1111/2041-210X.12180>

845 Pomeranz, J. P. F., Thompson, R. M., Poisot, T., & Harding, J. S. (2019). Inferring
846 predator–prey interactions in food webs. *Methods in Ecology and Evolution*, 10(3),
847 356–367. <https://doi.org/10.1111/2041-210X.13125>

848 Portalier, S. M. J., Fussmann, G. F., Loreau, M., & Cherif, M. (2019). The mechanics
849 of predator–prey interactions: First principles of physics predict predator–prey size
850 ratios. *Functional Ecology*, 33(2), 323–334. <https://doi.org/10.1111/1365-2435.13254>

851 Pringle, R. M. (2020). Untangling Food Webs. In *Unsolved Problems in Ecology* (pp.
852 225–238). Princeton University Press. <https://doi.org/10.1515/9780691195322-020>

853 Pringle, R. M., & Hutchinson, M. C. (2020). Resolving Food-Web Structure. *Annual*
854 *Review of Ecology, Evolution and Systematics*, 51, 55–80. <https://doi.org/10.1146/annurev-ecolsys-110218-024908>

855
856 Proulx, S. R., Promislow, D. E. L., & Phillips, P. C. (2005). Network thinking in ecology
857 and evolution. *Trends in Ecology & Evolution*, 20(6), 345–353. <https://doi.org/10>

1016/j.tree.2005.04.004

Pyke, G. (1984). Optimal Foraging Theory: A Critical Review. *Annual Review of Ecology, Evolution and Systematic*, 15, 523–575. <https://doi.org/10.1146/annurev.ecolsys.15.1.523>

Quintero, E., Arroyo-Correa, B., Isla, J., Rodríguez-Sánchez, F., & Jordano, P. (2024, July 16). *Downscaling mutualistic networks from species to individuals reveals consistent interaction niches and roles within plant populations*. <https://doi.org/10.1101/2024.02.02.578595> (Pre-published)

Rohr, R. P., Scherer, H., Kehrli, P., Mazza, C., & Bersier, L. (2010). Modeling Food Webs: Exploring Unexplained Structure Using Latent Traits. *The American Naturalist*, 176(2), 170–177. <https://doi.org/10.1086/653667>

Rooney, N., & McCann, K. S. (2012). Integrating food web diversity, structure and stability. *Trends in Ecology & Evolution*, 27(1), 40–46. <https://doi.org/10.1016/j.tree.2011.09.001>

Roopnarine, P. D. (2017). Ecological Modelling of Paleocommunity Food Webs. In *Conservation Paleobiology: Using the Past to Manage for the Future* (pp. 201–226). University of Chicago Press.

Roopnarine, P. D. (2006). Extinction Cascades and Catastrophe in Ancient Food Webs. *Paleobiology*, 32(1), 1–19. <https://www.jstor.org/stable/4096814>

Rossberg, A. G., Matsuda, H., Amemiya, T., & Itoh, K. (2006). Food webs: Experts consuming families of experts. *Journal of Theoretical Biology*, 241(3), 552–563. <https://doi.org/10.1016/j.jtbi.2005.12.021>

Saberski, E., Lorimer, T., Carpenter, D., Deyle, E., Merz, E., Park, J., Pao, G. M., & Sugihara, G. (2024). The impact of data resolution on dynamic causal inference in multiscale ecological networks. *Communications Biology*, 7(1), 1–10. <https://doi.org/10.1038/s42003-024-07054-z>

Sauve, A. M. C., Thébault, E., Pocock, M. J. O., & Fontaine, C. (2016). How plants connect pollination and herbivory networks and their contribution to community stability. *Ecology*, 97(4), 908–917. <https://doi.org/10.1890/15-0132.1>

Schneider, F. D., Brose, U., Rall, B. C., & Guill, C. (2016). Animal diversity and ecosystem functioning in dynamic food webs. *Nature Communications*, 7(1), 12718. <https://doi.org/10.1038/ncomms12718>

Segar, S. T., Fayle, T. M., Srivastava, D. S., Lewinsohn, T. M., Lewis, O. T., Novotny, V., Kitching, R. L., & Maunsell, S. C. (2020). The Role of Evolution in Shaping Ecological

892 Networks. *Trends in Ecology & Evolution*, 35(5), 454–466. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.tree.2020.01.004)
893 [tree.2020.01.004](https://doi.org/10.1016/j.tree.2020.01.004)

894 Shaw, J. O., Dunhill, A. M., Beckerman, A. P., Dunne, J. A., & Hull, P. M. (2024, January
895 30). *A framework for reconstructing ancient food webs using functional trait data.*
896 <https://doi.org/10.1101/2024.01.30.578036> (Pre-published)

897 Simmons, B. I., Blyth, P. S. A., Blanchard, J. L., Clegg, T., Delmas, E., Garnier, A.,
898 Griffiths, C. A., Jacob, U., Pennekamp, F., Petchey, O. L., Poisot, T., Webb, T.
899 J., & Beckerman, A. P. (2021). Refocusing multiple stressor research around the
900 targets and scales of ecological impacts. *Nature Ecology & Evolution*, 5(11), 1478–1489.
901 <https://doi.org/10.1038/s41559-021-01547-4>

902 Smith, J. G., Tomoleoni, J., Staedler, M., Lyon, S., Fujii, J., & Tinker, M. T. (2021).
903 Behavioral responses across a mosaic of ecosystem states restructure a sea otter–
904 urchin trophic cascade. *Proceedings of the National Academy of Sciences*, 118(11),
905 e2012493118. <https://doi.org/10.1073/pnas.2012493118>

906 Staniczenko, P. P. A., Lewis, O. T., Jones, N. S., & Reed-Tsochas, F. (2010). Structural
907 dynamics and robustness of food webs. *Ecology Letters*, 13(7), 891–899. <https://doi.org/10.1111/j.1461-0248.2010.01485.x>

908 Stephens, D. W., & Krebs, J. R. (1986). *Foraging Theory* (Vol. 1). Princeton University
909 Press. <https://doi.org/10.2307/j.ctvs32s6b>

910 Stock, M., Poisot, T., Waegeman, W., & Baets, B. D. (2017). Linear filtering reveals false
911 negatives in species interaction data. *Scientific Reports*, 7, 45908. [https://doi.org/10.](https://doi.org/10.1038/srep45908)
912 [1038/srep45908](https://doi.org/10.1038/srep45908)

913 Stouffer, D. B., Camacho, J., Guimerà, R., Ng, C. A., & Nunes Amaral, L. A. (2005).
914 Quantitative Patterns in the Structure of Model and Empirical Food Webs. *Ecology*,
915 86(5), 1301–1311. <https://doi.org/10.1890/04-0957>

916 Stouffer, D. B. (2019). All ecological models are wrong, but some are useful. *Journal of*
917 *Animal Ecology*, 88(2), 192–195. <https://doi.org/10.1111/1365-2656.12949>

918 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin,
919 M.-J., Hemming, V., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V.,
920 & Poisot, T. (2022). Food web reconstruction through phylogenetic transfer of low-
921 rank network representation. *Methods in Ecology and Evolution*, 13(12), 2838–2849.
922 <https://doi.org/10.1111/2041-210X.13835>

923 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M. J., Fortin,
924 M.-J., Mercier, B., Pollock, L. J., Runghen, R., Dalla Riva, G. V., & Poisot, T.
925

(2023). Graph embedding and transfer learning can help predict potential species interaction networks despite data limitations. *Methods in Ecology and Evolution*, 14(12), 2917–2930. <https://doi.org/10.1111/2041-210X.14228>

Strydom, T., Catchen, M. D., Banville, F., Caron, D., Dansereau, G., Desjardins-Proulx, P., Forero-Muñoz, N. R., Higinio, G., Mercier, B., Gonzalez, A., Gravel, D., Pollock, L., & Poisot, T. (2021). A roadmap towards predicting species interaction networks (across space and time). *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1837), 20210063. <https://doi.org/10.1098/rstb.2021.0063>

Su, M., Ma, Q., & Hui, C. (2024). Adaptive rewiring shapes structure and stability in a three-guild herbivore-plant-pollinator network. *Communications Biology*, 7(1), 103. <https://doi.org/10.1038/s42003-024-05784-8>

Terry, J. C. D., Bonsall, M. B., & Morris, R. J. (2025). The impact of structured higher-order interactions on ecological network stability. *Theoretical Ecology*, 18(1), 9. <https://doi.org/10.1007/s12080-025-00603-0>

Terry, J. C. D., & Lewis, O. T. (2020). Finding missing links in interaction networks. *Ecology*, 101(7), e03047. <https://doi.org/10.1002/ecy.3047>

Thébault, E., & Fontaine, C. (2010). Stability of ecological communities and the architecture of mutualistic and trophic networks. *Science (New York, N.Y.)*, 329(5993), 853–856. <https://doi.org/10.1126/science.1188321>

Thuiller, W., Calderón-Sanou, I., Chalmandrier, L., Gaüzère, P., O'Connor, L. M. J., Ohlmann, M., Poggiato, G., & Münkemüller, T. (2024). Navigating the integration of biotic interactions in biogeography. *Journal of Biogeography*, 51(4), 550–559. <https://doi.org/10.1111/jbi.14734>

Tylianakis, J. M., Laliberté, E., Nielsen, A., & Bascompte, J. (2010). Conservation of species interaction networks. *Biological Conservation, Conserving Complexity: Global Change and Community-Scale Interactions*, 143(10), 2270–2279. <https://doi.org/10.1016/j.biocon.2009.12.004>

Van De Walle, R., Logghe, G., Haas, N., Massol, F., Vandegheuchte, M. L., & Bonte, D. (2023). Arthropod food webs predicted from body size ratios are improved by incorporating prey defensive properties. *Journal of Animal Ecology*, 92(4), 913–924. <https://doi.org/10.1111/1365-2656.13905>

Van der Putten, W. H., Macel, M., & Visser, M. E. (2010). Predicting species distribution and abundance responses to climate change: Why it is essential to include biotic interactions across trophic levels. *Philosophical Transactions of the Royal Society B:*

960 *Biological Sciences*, 365(1549), 2025–2034. <https://doi.org/10.1098/rstb.2010.0037>

961 Vázquez, D. P., Blüthgen, N., Cagnolo, L., & Chacoff, N. P. (2009). Uniting pattern and
 962 process in plant–animal mutualistic networks: A review. *Annals of Botany*, 103(9),
 963 1445–1457. <https://doi.org/10.1093/aob/mcp057>

964 Williams, R. J., & Martinez, N. D. (2000). Simple rules yield complex food webs. *Nature*,
 965 404(6774, 6774), 180–183. <https://doi.org/10.1038/35004572>

966 Windsor, F. M., van den Hoogen, J., Crowther, T. W., & Evans, D. M. (2023). Using
 967 ecological networks to answer questions in global biogeography and ecology. *Journal*
 968 *of Biogeography*, 50(1), 57–69. <https://doi.org/10.1111/jbi.14447>

969 Wooster, E. I. F., Middleton, O. S., Wallach, A. D., Ramp, D., Sanisidro, O., Harris, V. K.,
 970 Rowan, J., Schowaneck, S. D., Gordon, C. E., Svenning, J.-C., Davis, M., Scharlemann,
 971 J. P. W., Nimmo, D. G., Lundgren, E. J., & Sandom, C. J. (2024). Australia’s recently
 972 established predators restore complexity to food webs simplified by extinction. *Current*
 973 *Biology*, 34(22), 5164–5172.e2. <https://doi.org/10.1016/j.cub.2024.09.049>

974 Wootton, K. L., Curtsdotter, A., Roslin, T., Bommarco, R., & Jonsson, T. (2023).
 975 Towards a modular theory of trophic interactions. *Functional Ecology*, 37(1), 26–43.
 976 <https://doi.org/10.1111/1365-2435.13954>

977 Yeakel, J. D., Pires, M. M., Rudolf, L., Dominy, N. J., Koch, P. L., Guimarães, P. R., &
 978 Gross, T. (2014). Collapse of an ecological network in Ancient Egypt. *PNAS*, 111(40),
 979 14472–14477. <https://doi.org/10.1073/pnas.1408471111>

980 Yodzis, P. (1982). The Compartmentation of Real and Assembled Ecosystems. *The*
 981 *American Naturalist*, 120(5), 551–570. <https://doi.org/10.1086/284013>

982 Yodzis, P. (2001). Must top predators be culled for the sake of fisheries? *Trends in Ecology*
 983 *& Evolution*, 16(2), 78–84. [https://doi.org/10.1016/S0169-5347\(00\)02062-0](https://doi.org/10.1016/S0169-5347(00)02062-0)




Figure 1: Aligning the processes that determine interactions (right) with different network representations (left). A **global metaweb** captures all feasible interactions among a species pool. Restricting this network by species co-occurrence yields **regional metawebs**, which differ in species composition. Species occurring in both regions are shown in red, whereas region-specific species are shown in blue and yellow. **Realised networks** represent subsets of regional metawebs expressed in specific spatial and temporal contexts. Their structure is shaped not only by co-occurrence, but also by community-level processes such as abundance, diet choice, and non-trophic interactions.