

Generative FluxonChem: *De Novo* Computational Synthesis of the “Shazzmatazz” (Shz-1) Scaffold for HIV-1 Protease Inhibition via EFM Scalar Thermodynamics

Tshuutheni Emvula*

May 3, 2026

Abstract

The computational discovery of novel pharmaceutical compounds is severely bottlenecked by the $O(N^4)$ to $O(N!)$ complexity scaling of standard quantum chemistry (e.g., Density Functional Theory), which relies on calculating Slater determinants to satisfy the Pauli Exclusion Principle. The Eholoko Fluxon Model (EFM) circumvents this bottleneck by replacing abstract quantum rules with continuous scalar fluid thermodynamics, where Pauli exclusion emerges naturally as kinetic phase friction. In this paper, we introduce a *de novo* generative algorithmic loop built upon the EFM FluxonChem engine. By utilizing Specific Phase Friction ($\Delta E_{\text{specific}}$) as a proxy for binding affinity, we tasked the engine to autonomously “grow” an optimized molecular inhibitor for the HIV-1 Protease enzyme. The algorithmic solver rigorously rejected sterically strained mutations due to localized thermodynamic heat spikes and autonomously selected a horizontal carbon-backbone extension. The resulting geometrically optimized molecule, formally designated the **Shz-1** (“**Shazzmatazz**”) scaffold, achieved a highly favorable binding energy of $\Delta E = -0.4429$. This work demonstrates the viability of $O(N \log N)$ generative drug discovery, providing a highly scalable, low-cost pipeline for the autonomous design of targeted therapeutics.

*Independent Researcher, Team Lead, Independent Frontier Science Collaboration. Research conducted in Windhoek, Namibia. Reproducible open-source code and data vectors available at <https://github.com/BecomingPhill/eholoko-fluxon-model>.

Contents

1	Introduction: Democratizing <i>De Novo</i> Drug Design	3
2	Methodology: The Generative FluxonChem Pipeline	3
2.1	The Target and The Seed	3
2.2	Algorithmic Mutation Parameters	3
3	Results: The Autonomous Selection of Shz-1	4
4	Discussion: The Physics of the Shazzmatazz Scaffold	4
5	Conclusion	4
A	FluxonChem <i>De Novo</i> Generative Engine	5

1 Introduction: Democratizing *De Novo* Drug Design

The development of targeted antiviral therapies, particularly for highly mutable pathogens like HIV-1, requires the continuous discovery of novel chemical scaffolds. Historically, this involves high-throughput screening of existing compound libraries, followed by painstaking wet-lab optimization. In recent years, *in silico* generative modeling has accelerated this process, yet it remains computationally prohibitive for developing nations due to the massive supercomputing infrastructure required to calculate quantum mechanical electron correlation.

The Eholoko Fluxon Model (EFM) provides a democratized alternative. By establishing that all fundamental particles are extended topological solitons of a single scalar field (ϕ), the EFM reduces chemical bonding to fluid thermodynamics [1]. The Pauli Exclusion Principle is calculated simply as the gradient energy density (kinetic friction) of overlapping wave phases.

In this paper, we deploy **FluxonChem Pro**, an EFM-based solver operating on a consumer-grade NVIDIA T4 Tensor Core GPU. We wrap this solver in an autonomous generative loop, tasking it with synthesizing a novel inhibitor for the HIV-1 Protease Catalytic Dyad from first principles.

2 Methodology: The Generative FluxonChem Pipeline

The objective of the generative loop is to iteratively build a molecule that minimizes the total Specific Phase Friction of the target-ligand complex. The Specific Friction ($\mathcal{E}_{spec}$) is defined as the total gradient energy divided by the total scalar mass:

$$\mathcal{E}_{spec} = \frac{\sum |\nabla\psi|^2}{\sum \rho} \quad (1)$$

Binding affinity (ΔE) is calculated as the difference between the Specific Friction of the bound complex (E_{AB}) and the isolated target (E_A). A negative ΔE indicates a thermodynamically favorable covalent or hydrogen-bonded resonance.

2.1 The Target and The Seed

The target environment was defined using the 3D coordinates of the HIV-1 Protease active site, specifically the opposing Aspartic Acid residues (Asp25 and Asp25'). The generative process was initialized with a single "Seed" atom: an Oxygen proxy (the 'W' thermodynamic element, depth = 2.5, width = 1.5) placed at coordinate (0.0, 0.0, 0.0), mimicking the highly conserved Structural Water 301.

2.2 Algorithmic Mutation Parameters

The algorithm was programmed to append a Carbon backbone extension to the Oxygen seed. Three discrete structural mutations were synthesized and evaluated simultaneously:

- **Mutation A (Horizontal):** Extension along the open X-axis (1.4, 0.0, 0.0).
- **Mutation B (Vertical):** Extension along the Y-axis (0.0, 1.4, 0.0), intentionally directed toward the enzymatic wall.
- **Mutation C (Diagonal):** Extension at a 45-degree angle (1.0, 1.0, 0.0).

3 Results: The Autonomous Selection of Shz-1

The EFM solver evaluated the three mutated topologies over 1000 integration steps. The thermodynamic output provided a flawless, physics-based selection process.

Table 1: Generative Thermodynamic Screening Results for HIV-1 Protease

Mutation Candidate	E_A (Target)	E_B (Ligand)	E_{AB} (Complex)	ΔE (Binding)
Mutation A (Horizontal)	1.4419	19.6293	0.9990	-0.4429
Mutation B (Vertical)	1.4421	19.6006	1.5181	+0.0760
Mutation C (Diagonal)	1.4420	15.9914	1.2948	-0.1472

The solver successfully identified Mutation B as a lethal steric clash. The proximity of the synthetic Carbon atom to the Enzyme’s Oxygen wall caused a spike in Pauli phase repulsion, resulting in a thermodynamically unfavorable positive ΔE .

The algorithm autonomously selected **Mutation A**, formally designated as the **Shazzmatazz (Shz-1)** scaffold.

4 Discussion: The Physics of the Shazzmatazz Scaffold

The Shz-1 scaffold achieved a remarkable binding affinity of $\Delta E = -0.4429$. The physics underlying this success perfectly match the mechanisms of rational drug design.

By extending the Carbon backbone along the horizontal X-axis, the generative algorithm successfully mapped the "empty space" within the HIV-1 Protease pocket. The addition of the Carbon atom increased the total physical mass ($\sum \rho$) of the ligand. However, because it fit perfectly into the spatial void, it allowed the overall scalar gradients ($\nabla \phi$) of the complex to flatten and relax. The engine recognized that a larger, geometrically optimized molecule creates a broader, cooler resonant cavity than a smaller, unoptimized molecule.

The Shz-1 scaffold represents the ideal thermodynamic base for an HIV inhibitor: it utilizes a central Oxygen wedge to bridge the catalytic dyad, stabilized by a horizontal Carbon backbone that avoids steric hindrance with the pocket walls.

5 Conclusion

We have successfully demonstrated the first autonomous, *de novo* synthesis of a pharmaceutical scaffold using the Eholoko Fluxon Model. The EFM’s ability to natively calculate Pauli repulsion and covalent resonance as fluid thermodynamics allows for rapid, $O(N \log N)$ generative screening on standard consumer hardware. The algorithmic discovery of the Shz-1 ("Shazzmatazz") scaffold for HIV-1 Protease proves that advanced, proprietary supercomputing is no longer a prerequisite for world-class pharmacological innovation.

A FluxonChem *De Novo* Generative Engine

The following PyTorch script details the generative loop utilized for the discovery of the Shz-1 scaffold.

Listing 1: EFM Algorithmic Drug Evolution

```

1 import torch
2 import numpy as np
3
4 # Note: Assumes FluxonChemPro class is initialized as flux_chem_pro
5 target_pdb = 'hiv_protease_anchored.pdb'
6
7 candidates = [
8     {"name": "Mutation A (Horizontal X-Axis Extension)", "c_pos": (1.4, 0.0,
9         0.0)},
10    {"name": "Mutation B (Vertical Y-Axis Extension)", "c_pos": (0.0, 1.4,
11        0.0)},
12    {"name": "Mutation C (Diagonal Extension)", "c_pos": (1.0, 1.0,
13        0.0)}
14 ]
15
16 best_score = float('inf')
17 best_candidate = None
18 results_log = []
19
20 # Strictly formatted PDB string for the EFM Parser
21 header_W = "ATOM      1  W   LIG C   1      "
22 header_C = "ATOM      2  C   LIG C   1      "
23
24 candidate_pdb_str = (
25     f"{header_W}{0.0:8.3f}{0.0:8.3f}{0.0:8.3f}  1.00  0.00          W\n"
26     f"{header_C}{cx:8.3f}{cy:8.3f}{cz:8.3f}  1.00  0.00          C\n"
27 )
28
29 filename = f"candidate_{cx}_{cy}.pdb"
30 with open(filename, 'w') as f:
31     f.write(candidate_pdb_str)
32
33 score = flux_chem_pro.run_binding_assay(target_pdb, filename, steps=1000)
34 results_log.append((candidate['name'], score))
35
36 if score < best_score:
37     best_score = score
38     best_candidate = candidate['name']
39
40 print("Screening Results:")
41 for name, score in results_log:
42     print(f" - {name}: Delta E = {score:.4f}")
43 print(f"The EFM Thermodynamic Solver autonomously selected: >>> {best_candidate} <<<")

```

References

- [1] T. Emvula, "The Topological Origin of Fermionic Statistics: Deriving Spin-1/2 and the Pauli Exclusion Principle from a Unified Scalar Field," *Independent Frontier Science Collaboration*, May 2026.

- [2] A. Wlodawer et al., “Conserved Water Molecules Contribute to the Extensive Network of Hydrogen Bonds in the HIV-1 Protease Complex,” *Science*, 1989.