

# Categorifying Algebra

Saxon Supple

February 2026

## 1 Introduction

This article will give categorical definitions of the standard objects from algebra, by way of defining universal properties which determine the objects up to unique isomorphism. For a given object  $A$  of interest, the general procedure will be to first define an object  $B$  as an object which satisfies a specific universal property, then show that  $A$  satisfies that universal property, and then show that all pairs of objects  $B$  which satisfy the universal property are related by a unique isomorphism. This will then demonstrate that the universal property determines  $A$  up to unique isomorphism. We will then apply these universal properties to prove additional useful categorical results, such as the functoriality of the abelianization of a group.

## 2 Sets

**Definition 2.1** (Universal property of the quotient set). *Let  $X$  be a set and let  $\sim$  be an equivalence relation on  $X$ . A pair  $(X', \pi : X \rightarrow X')$  is a quotient of  $X$  by  $\sim$  if given any  $f : X \rightarrow Y$  such that  $x_0 \sim x_1 \implies f(x_0) = f(x_1) \forall x_0, x_1 \in X$ , then there exists a unique  $\bar{f} : X' \rightarrow Y$  such that  $f = \bar{f} \circ \pi$ .*

**Theorem 2.2.** *The canonical projection  $\pi : X \rightarrow X/\sim : x \mapsto [x]$  satisfies the universal property of the quotient set.*

*Proof.* We define  $\bar{f} : X/\sim \rightarrow Y$  by  $\bar{f}(\pi(x)) = f(x)$ . This is well-defined, since if  $x_0 \sim x_1$ , then  $\bar{f}(\pi(x_0)) = f(x_0) = f(x_1) = \bar{f}(\pi(x_1))$ , and hence  $\bar{f}$  is independent of representatives. Uniqueness is also evident from the definition of  $\pi$ .  $\square$

**Theorem 2.3.** *Let  $X$  be a set with an equivalence relation  $\sim$ . Let  $(W, \pi_W)$  and  $(Z, \pi_Z)$  be two pairs of sets, along with maps  $\pi_W : X \rightarrow W$  and  $\pi_Z : X \rightarrow Z$  such that both pairs satisfy the universal property of the quotient set. Then there exists a unique bijection  $h : W \rightarrow Z$  such that  $\pi_Z = h \circ \pi_W$ .*

*Proof.* Let  $x_0 \sim x_1$ . Then  $\pi_Z(x_0) = \pi_Z(x_1)$ , by the definition of a quotient map. Hence, by the universal property of the quotient set, there exists a unique  $h : W \rightarrow Z$  such that  $\pi_Z = h \circ \pi_W$ . Similarly, there exists a  $g : Z \rightarrow W$  such that  $\pi_W = g \circ \pi_Z$ . Hence,  $\pi_Z = (h \circ g) \circ \pi_Z$ . However, we also have that  $\pi_Z = \text{id}_Z \circ \pi_Z$ , and hence, by uniqueness,  $h \circ g = \text{id}_Z$ . Similarly,  $g \circ h = \text{id}_W$ , implying that  $h$  is a bijection, as required.  $\square$

### 3 Groups

#### 3.1 Quotient Groups

**Definition 3.1** (Universal property of the quotient group). *Let  $G$  be a group, and let  $N \trianglelefteq G$  be a normal subgroup. A pair  $(Q, q : G \rightarrow Q)$ , consisting of a group  $Q$  and a group homomorphism  $q$ , is a quotient of  $G$  if  $N \subseteq \ker q$  and given any group homomorphism  $\phi : G \rightarrow H$  with  $N \subseteq \ker \phi$ , there exists a unique group homomorphism  $\bar{\phi} : Q \rightarrow H$  such that  $\phi = \bar{\phi} \circ q$ .*

**Theorem 3.2.** *Let  $G$  be a group and let  $N \trianglelefteq G$  be a normal subgroup. Then the canonical quotient homomorphism  $\pi : G \rightarrow G/N : g \mapsto gN$  is a quotient map.*

*Proof.* Let  $\sim$  be an equivalence relation on  $G$  which identifies the fibres of  $\pi$ ; that is,  $x \sim y \iff xN = yN$ . If  $\pi(x) = \pi(y)$ , then  $x$  and  $y$  are in the same coset, so there exists some  $n \in N$  such that  $x = yn$ . Hence, since  $N \subseteq \ker \phi$ , we have  $\phi(x) = \phi(yn) = \phi(y)\phi(n) = \phi(y)$ . Then by the universal property of the quotient set, there exists a unique map  $\bar{\phi} : G/\sim \rightarrow H$  such that  $\phi = \bar{\phi} \circ \pi$ . In particular,  $\bar{\phi}$  is defined as  $\bar{\phi}([x]) = \bar{\phi}(xN) = \phi(x)\forall x$ , and hence  $\bar{\phi}$  is simply a map on the cosets of  $N$ . Furthermore, given  $x, y \in G$ , we have  $\bar{\phi}(xyN) = \phi(xy) = \phi(x)\phi(y) = \bar{\phi}(xN)\bar{\phi}(yN)$ , and hence  $\bar{\phi}$  is a group homomorphism.  $\square$

**Corollary 3.3.** *Let  $\phi : G \rightarrow H$  be a group homomorphism, and let  $N \trianglelefteq G$  be a normal subgroup such that  $N \subseteq \ker \phi$ . Then  $N = \ker \phi$  if and only if  $\bar{\phi} : G/N \rightarrow H$  is injective.*

*Proof.* By the universal property of the quotient group, we know that  $\bar{\phi}$  is both unique and well-defined. First suppose that  $N = \ker \phi$ . Let  $gN \in \ker \bar{\phi}$ . Then  $\phi(g) = \bar{\phi}(gN) = e$ , and hence  $g \in \ker \phi = N$ , implying that  $gN = N$ . Hence,  $\ker \bar{\phi} = N$ , implying that  $\bar{\phi}$  is injective. Now suppose that  $\bar{\phi}$  is injective. That then means that  $\ker \bar{\phi} = N$ . Let  $g \in \ker \phi$ . Then  $\bar{\phi}(gN) = \phi(g) = e$ , implying that  $gN = N$ , or  $g \in N$ . Hence,  $N = \ker \phi$ .  $\square$

**Theorem 3.4.** *Let  $G$  be a group and let  $N \trianglelefteq G$  be a normal subgroup. Let  $q : G \rightarrow Q$  and  $q' : G \rightarrow Q'$  be two quotient homomorphisms. Then there exists a unique isomorphism  $h : Q \rightarrow Q'$  such that  $q' = h \circ q$ .*

*Proof.* By the universal property of the quotient group, there exists a unique group homomorphism  $h : Q \rightarrow Q'$  such that  $q' = h \circ q$ , and similarly, there exists a unique group homomorphism  $g : Q' \rightarrow Q$  such that  $q = g \circ q'$ . Hence,  $q' = (h \circ g) \circ q'$ . However, we also have  $q' = \text{id}_{Q'} \circ q'$ , and so the universal property implies that  $h \circ g = \text{id}_{Q'}$ . Similarly,  $g \circ h = \text{id}_Q$ , and hence  $h$  is an isomorphism.  $\square$

#### 3.2 Abelianization

**Definition 3.5** (Universal property of the abelianization of a group). *Let  $G$  be a group. An abelianization of  $G$  is a pair  $(G^{ab}, \pi)$ , where  $G^{ab}$  is an abelian group and  $\pi : G \rightarrow G^{ab}$  is a group homomorphism, such that given any abelian group  $H$  and group homomorphism  $\phi : G \rightarrow H$ , there exists a unique group homomorphism  $\phi^{ab} : G^{ab} \rightarrow H$  such that  $\phi^{ab} \circ \pi = \phi$ .*

**Theorem 3.6.** *Let  $G$  be a group, let  $[G, G]$  be the commutator subgroup, and let  $\pi : G \rightarrow G/[G, G]$  be the canonical quotient homomorphism. Then  $(G/[G, G], \pi)$  is an abelianization of  $G$ .*

*Proof.* We simply need to show that the commutator subgroup  $[G, G]$  is contained within the kernel of  $\phi$ , since that will allow us to apply the universal property of the quotient group to obtain the unique group homomorphism  $\phi^{ab} : G/[G, G] \rightarrow H$  such that  $\phi^{ab} \circ \pi = \phi$ . Indeed, let  $x, y \in G$ . Then  $\phi([x, y]) = \phi(x^{-1}y^{-1}xy) = \phi(x)^{-1}\phi(y)^{-1}\phi(x)\phi(y) = e$ , so  $[x, y] \in \ker \phi$ . Hence, since  $[G, G]$  is the subgroup generated by all commutators, it follows that  $[G, G] \subseteq \ker \phi$ , as required.  $\square$

**Theorem 3.7.** Let  $G$  be a group, and let  $(G^{ab}, \pi)$  and  $(G^{ab'}, \pi')$  be two abelianizations of  $G$ . Then there exists a unique isomorphism  $h : G^{ab} \rightarrow G^{ab'}$  such that  $\pi' = h \circ \pi$ .

*Proof.* By the universal property of the abelianization of a group, there exists a unique group isomorphism  $h : G^{ab} \rightarrow G^{ab'}$  such that  $\pi' = h \circ \pi$ . Similarly, there exists a unique group isomorphism  $g : G^{ab'} \rightarrow G^{ab}$  such that  $\pi = g \circ \pi'$ . Hence,  $\pi' = (h \circ g) \circ \pi'$ . However, we also have  $\pi' = \text{id}_{G^{ab'}} \circ \pi'$ , and hence, by uniqueness, we have  $h \circ g = \text{id}_{G^{ab'}}$ . Similarly,  $g \circ h = \text{id}_{G^{ab}}$ , implying that  $h$  is an isomorphism.  $\square$

**Lemma 3.8.** Let  $\phi : G \rightarrow H$  be a group homomorphism, and let  $\pi_G : G \rightarrow G/[G, G]$ ,  $\pi_H : H \rightarrow H/[H, H]$  be the quotient homomorphisms of  $G$  and  $H$  into their respective abelianizations. Then there exists a group homomorphism  $\phi^{ab} : G^{ab} \rightarrow H^{ab}$  given by  $\phi^{ab}(g[G, G]) = \phi(g)[H, H] \forall g[G, G] \in G^{ab}$ , which is the unique group homomorphism  $\psi : G^{ab} \rightarrow H^{ab}$  such that  $\psi \circ \pi_G = \pi_H \circ \phi$ .

*Proof.* Consider the map  $\pi_H \circ \phi : G \rightarrow H^{ab}$ . This is a map from a group to an abelian group, so by the universal property of the abelianization of a group, there exists a unique group homomorphism  $\phi^{ab} : G^{ab} \rightarrow H^{ab}$  such that  $\phi^{ab} \circ \pi_G = \pi_H \circ \phi$ . That is, given any  $g \in G$ , we have

$$\phi^{ab}(g[G, G]) = \phi^{ab} \circ \pi_G(g) = \pi_H \circ \phi(g) = \phi(g)[H, H],$$

as required.  $\square$

**Theorem 3.9.** There is a covariant functor, called the abelianization functor, from the category of groups to the category of abelian groups.

*Proof.* Let  $F$  be the functor which maps groups to their abelianizations, and group homomorphisms to group homomorphisms between the abelianizations of the groups. That is, given groups  $G$  and  $H$  and a group homomorphism  $\phi : G \rightarrow H$ , we have  $F(G) = G^{ab}$  and  $F(\phi) = \phi^{ab} : G^{ab} \rightarrow H^{ab} : g[G, G] \mapsto \phi(g)[H, H]$ .

We first show that  $F(\text{id}_G) = \text{id}_{F(G)}$  for any group  $G$ . Indeed,  $F(\text{id}_G)$  is given by  $\text{id}_{G^{ab}} : G^{ab} \rightarrow G^{ab}$ , where  $\text{id}_{G^{ab}}(g[G, G]) = \text{id}_G(g)[G, G] = g[G, G] \forall g \in G$ .  $F(\text{id}_G)$  is then the identity map on  $G^{ab} = F(G)$ , as required.

We now show that  $F$  preserves composition. Let  $G, H, K$  be groups and let  $\phi : G \rightarrow H$  and  $\psi : H \rightarrow K$  be group homomorphisms. Then

$$F(\phi) = \phi^{ab} : G^{ab} \rightarrow H^{ab} : g[G, G] \mapsto \phi(g)[H, H],$$

$$F(\psi) = \psi^{ab} : H^{ab} \rightarrow K^{ab} : h[H, H] \mapsto \psi(h)[K, K]$$

and

$$F(\psi \circ \phi) = (\psi \circ \phi)^{ab} : G^{ab} \rightarrow K^{ab} : g[G, G] \mapsto (\psi \circ \phi)(g)[K, K].$$

Hence,

$$\begin{aligned} (\psi^{ab} \circ \phi^{ab})(g[G, G]) &= \psi^{ab}(\phi(g)[H, H]) \\ &= \psi(\phi(g))[K, K] \\ &= (\psi \circ \phi)(g)[K, K] \\ &= (\psi \circ \phi)^{ab}(g[G, G]), \end{aligned}$$

as required. Hence,  $F$  is a covariant functor.  $\square$