

# Concerning Universal Properties and the Abelianization Functor

Saxon Supple

February 2026

**Theorem 1.1** (Universal property of the quotient set). *Let  $X$  and  $Y$  be sets, and let  $\sim$  be an equivalence relation on  $X$ . Let  $\pi : X \rightarrow X/\sim$  be the quotient map, and let  $f : X \rightarrow Y$  be such that  $x_0 \sim x_1 \implies f(x_0) = f(x_1) \forall x_0, x_1 \in X$ . Then there exists a unique  $\bar{f} : X/\sim \rightarrow Y$  such that  $f = \bar{f} \circ \pi$ .*

*Proof.* We define  $\bar{f} : X/\sim \rightarrow Y$  by  $\bar{f}([x]) = f(x)$ . This is well-defined, since if  $x_0 \sim x_1$ , then  $\bar{f}([x_0]) = f(x_0) = f(x_1) = \bar{f}([x_1])$ , and hence  $\bar{f}$  is independent of representatives. Uniqueness is also evident from the definition.  $\square$

**Theorem 1.2** (Universal property of the quotient group). *Let  $G, H$  be groups, and let  $N \trianglelefteq G$  be a normal subgroup. Let  $\pi : G \rightarrow G/N$  be the quotient map, and let  $\phi : G \rightarrow H$  be a group homomorphism, with  $N \subseteq \ker \phi$ . Then there exists a unique group homomorphism  $\bar{\phi} : G/N \rightarrow H$  such that  $\phi = \bar{\phi} \circ \pi$ .*

*Proof.* Let  $\sim$  be an equivalence relation on  $G$  which identifies the fibres of  $\pi$ ; that is,  $x \sim y \iff xN = yN$ . Furthermore, since  $N \subseteq \ker \phi$ , we have that if  $\pi(x) = \pi(y)$ , then  $x$  and  $y$  are in the same coset, so there exists some  $n \in N$  such that  $x = yn$ . Hence,  $\phi(x) = \phi(yn) = \phi(y)\phi(n) = \phi(y)$ . Then by the universal property of the quotient set, there exists a unique map  $\bar{\phi} : G/\sim \rightarrow H$  such that  $\phi = \bar{\phi} \circ \pi$ . In particular,  $\bar{\phi}$  is defined as  $\bar{\phi}([x]) = \bar{\phi}(xN) = \phi(x) \forall x$ , and hence  $\bar{\phi}$  is simply a map on the cosets of  $N$ . Furthermore, given  $x, y \in G$ , we have  $\bar{\phi}(xyN) = \phi(xy) = \phi(x)\phi(y) = \bar{\phi}(xN)\bar{\phi}(yN)$ , and hence  $\bar{\phi}$  is a group homomorphism.  $\square$

**Corollary 1.3.** *Let  $\phi : G \rightarrow H$  be a group homomorphism, and let  $N \trianglelefteq G$  be a normal subgroup such that  $N \subseteq \ker \phi$ . Then  $N = \ker \phi$  if and only if  $\bar{\phi} : G/N \rightarrow H$  is injective.*

*Proof.* By the universal property of the quotient group, we know that  $\bar{\phi}$  is both unique and well-defined. First suppose that  $N = \ker \phi$ . Let  $gN \in \ker \bar{\phi}$ . Then  $\phi(g) = \bar{\phi}(gN) = e$ , and hence  $g \in \ker \phi = N$ , implying that  $gN = N$ . Hence,  $\ker \bar{\phi} = N$ , implying that  $\bar{\phi}$  is injective. Now suppose that  $\bar{\phi}$  is injective. That then means that  $\ker \bar{\phi} = N$ . Let  $g \in \ker \phi$ . Then  $\bar{\phi}(gN) = \phi(g) = e$ , implying that  $gN = N$ , or  $g \in N$ . Hence,  $N = \ker \phi$ .  $\square$

**Theorem 1.4** (Universal property of the abelianization of a group). *Let  $G$  be a group, and let  $G^{ab} := G/[G, G]$  be its abelianization. Let  $\pi : G \rightarrow G^{ab}$  be the quotient group homomorphism. Let  $H$  be an abelian group. Let  $\phi : G \rightarrow H$  be a group homomorphism. Then there exists a unique group homomorphism  $\phi^{ab} : G^{ab} \rightarrow H$  such that  $\phi^{ab} \circ \pi = \phi$ .*

*Proof.* We simply need to show that the commutator subgroup  $[G, G]$  is contained within the kernel of  $\phi$ , since that will allow us to apply the universal property of the quotient group to obtain the unique group homomorphism  $\phi^{\text{ab}} : G^{\text{ab}} \rightarrow H$  such that  $\phi^{\text{ab}} \circ \pi = \phi$ . Indeed, let  $x, y \in G$ . Then  $\phi([x, y]) = \phi(x^{-1}y^{-1}xy) = \phi(x)^{-1}\phi(y)^{-1}\phi(x)\phi(y) = e$ , so  $[x, y] \in \ker \phi$ . Hence, since  $[G, G]$  is the subgroup generated by all commutators, it follows that  $[G, G] \subseteq \ker \phi$ , as required.  $\square$

**Lemma 1.5.** *Let  $\phi : G \rightarrow H$  be a group homomorphism, and let  $\pi_G : G \rightarrow G/[G, G]$ ,  $\pi_H : H \rightarrow H/[H, H]$  be the quotient homomorphisms of  $G$  and  $H$  into their respective abelianizations. Then there exists a group homomorphism  $\phi^{\text{ab}} : G^{\text{ab}} \rightarrow H^{\text{ab}}$  given by  $\phi^{\text{ab}}(g[G, G]) = \phi(g)[H, H] \forall g[G, G] \in G^{\text{ab}}$ , which is the unique group homomorphism  $\psi : G^{\text{ab}} \rightarrow H^{\text{ab}}$  such that  $\psi \circ \pi_G = \pi_H \circ \phi$ .*

*Proof.* Consider the map  $\pi_H \circ \phi : G \rightarrow H^{\text{ab}}$ . This is a map from a group to an abelian group, so by the universal property of the abelianization of a group, there exists a unique group homomorphism  $\phi^{\text{ab}} : G^{\text{ab}} \rightarrow H^{\text{ab}}$  such that  $\phi^{\text{ab}} \circ \pi_G = \pi_H \circ \phi$ . That is, given any  $g \in G$ , we have

$$\phi^{\text{ab}}(g[G, G]) = \phi^{\text{ab}} \circ \pi_G(g) = \pi_H \circ \phi(g) = \phi(g)[H, H],$$

as required.  $\square$

**Theorem 1.6.** *There is a covariant functor, called the abelianization functor, from the category of groups to the category of abelian groups.*

*Proof.* Let  $F$  be the functor which maps groups to their abelianizations, and group homomorphisms to group homomorphisms between the abelianizations of the groups. That is, given groups  $G$  and  $H$  and a group homomorphism  $\phi : G \rightarrow H$ , we have  $F(G) = G^{\text{ab}}$  and  $F(\phi) = \phi^{\text{ab}} : G^{\text{ab}} \rightarrow H^{\text{ab}} : g[G, G] \mapsto \phi(g)[H, H]$ .

We first show that  $F(\text{id}_G) = \text{id}_{F(G)}$  for any group  $G$ . Indeed,  $F(\text{id}_G)$  is given by  $\text{id}_{G^{\text{ab}}} : G^{\text{ab}} \rightarrow G^{\text{ab}}$ , where  $\text{id}_{G^{\text{ab}}}(g[G, G]) = \text{id}_G(g)[G, G] = g[G, G] \forall g \in G$ .  $F(\text{id}_G)$  is then the identity map on  $G^{\text{ab}} = F(G)$ , as required.

We now show that  $F$  preserves composition. Let  $G, H, K$  be groups and let  $\phi : G \rightarrow H$  and  $\psi : H \rightarrow K$  be group homomorphisms. Then

$$F(\phi) = \phi^{\text{ab}} : G^{\text{ab}} \rightarrow H^{\text{ab}} : g[G, G] \mapsto \phi(g)[H, H],$$

$$F(\psi) = \psi^{\text{ab}} : H^{\text{ab}} \rightarrow K^{\text{ab}} : h[H, H] \mapsto \psi(h)[K, K]$$

and

$$F(\psi \circ \phi) = (\psi \circ \phi)^{\text{ab}} : G^{\text{ab}} \rightarrow K^{\text{ab}} : g[G, G] \mapsto (\psi \circ \phi)(g)[K, K].$$

Hence,

$$\begin{aligned} (\psi^{\text{ab}} \circ \phi^{\text{ab}})(g[G, G]) &= \psi^{\text{ab}}(\phi(g)[H, H]) \\ &= \psi(\phi(g))[K, K] \\ &= (\psi \circ \phi)(g)[K, K] \\ &= (\psi \circ \phi)^{\text{ab}}(g[G, G]), \end{aligned}$$

as required. Hence,  $F$  is a covariant functor.  $\square$