



**POLITECNICO**  
**MILANO 1863**

M.Sc. Computer Science and Engineering  
Data Analysis for Smart Agriculture

## Data Analysis On An Eggs Farm

Davide Canali	-	10674880	-	davide1.canali@polimi.it
Matteo Cordioli	-	00000000	-	matteo.cordioli@polimi.it
Federico Camilletti	-	10619856	-	federico.camilletti@polimi.it
Shakiba Shahidiani	-	00000000	-	shakiba.shahidiani@polimi.it

December 13, 2022

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Data</b>	<b>2</b>
2.1	Data Collection . . . . .	2
2.2	Data Cleaning . . . . .	2
<b>3</b>	<b>Analysis of Each Cycle</b>	<b>3</b>
3.1	Clustering Techniques . . . . .	3
3.2	Cycle A . . . . .	3
3.3	Cycle B . . . . .	4
3.4	Cycle C . . . . .	5
3.5	Cycle X1 . . . . .	6
3.6	Cycle X2 . . . . .	7
3.7	Cycle Y . . . . .	8
3.8	Cycle Z . . . . .	9
<b>4</b>	<b>Common features</b>	<b>10</b>
<b>5</b>	<b>Organic vs non-organic cycles</b>	<b>12</b>
<b>6</b>	<b>Death-season correlation</b>	<b>14</b>
<b>7</b>	<b>Cycle C</b>	<b>15</b>
<b>8</b>	<b>Economic results</b>	<b>17</b>
<b>9</b>	<b>Data processing: shakiba</b>	<b>17</b>

# 1 Introduction

In this study, we are going to analyze data from an eggs farm near Mantova to see if it's possible to improve both animal welfare and farmer revenue.

The farm under analysis is [//TODO INSERT NAME] and has around 40'000 chickens that produce organic eggs. We have the data starting from 2014, the production of eggs is divided into cycles lasting about 13 - 15 months each. We have 5 complete cycles and the current 2022 cycle. The first 2 cycles (called X and Y) are non-organic which means the chickens are treated differently from the last 4 cycles (Z, A, B, C) which are organic.

The chickens are divided into 2 equal-sized barns, both of which receive the same treatment and operations.

The farmer is in an agistment that takes care of refurbishing the feed and the livestock.

Upon talking with the farmer we focus our attention on 3 main topics which involve:

- Understanding the mortality between different cycles and organic with non-organic.
- Improve the welfare of the chickens.
- Quantify the monetary loss when a chicken dies at the start of the cycle. //TODO
- Quantify the losses in cycle C

## 2 Data

### 2.1 Data Collection

The farmer collected the data thanks to sensors and by inspecting the barns each day. Before cycle A the data were only collected in paper spreadsheets and had to be imported in a digital format. The data on paper had 5 features: date, death, current chickens, eggs produced and sold when the data became saved digitally 2 extra features were collected which are feed and water.

### 2.2 Data Cleaning

Before starting to analyze the data we cleaned them. In all the cycles we performed these next steps:

- Removed tail and head (20 rows each more or less).
- Adjusted feed and water consumption to correct typos and switchboard errors.
- Adjusted eggs production when % of laid when above 100%, distributed in the days before assuming that those eggs were not collected.
- Adjusted death count to match the delta of death at the end of the cycle.
- Checked that the difference between the total amount of eggs produced and the number of eggs sold was less than 0.6%.

Specifically, in cycle A we also adjust a series of missing data regarding the water consumption adding them following the cycle distribution.

//TODO Teo add what u did in cycles Y and X

### 3 Analysis of Each Cycle

#### 3.1 Clustering Techniques

We used 3 different clustering techniques K-Means, Hierarchical and Density Based Clustering. To choose the parameters for these methods we used different solutions:

- K-Means: silhouette technique to find the optimal number of clusters.
- Hierarchical: plot the dendrogram to find the optimal number of clusters.
- Density: plot the k-NN to find the optimal  $\epsilon$ .


#### 3.2 Cycle A

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
19/7/2018	42.009	1/9/2018	19/5/2020	Yes	20.208.086

Significant features:

- Average Percentage of Laid Eggs each day: 83,8%
- Average Percentage of Death each day: 0,044%
- Average Temperature during the cycle: 12,99 °C
- Average Humidity during the cycle: 74,55%

Cluster:




### 3.3 Cycle B

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
09/08/2020	42.098	24/09/2020	02/05/2022	Yes	18.392.640

Significant features:

- Average Percentage of Laid Eggs each day: 81,39%
- Average Percentage of Death each day: 0,0494%
- Average Temperature during the cycle: 11,81 °C
- Average Humidity during the cycle: 73,24%

Cluster:




### 3.4 Cycle C

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
20/06/2022	42.098	08/08/2022	In progress	Yes	In progress

Significant features:

- Average Percentage of Laid Eggs each day: 55,85%
- Average Percentage of Death each day: 0,043%
- Average Temperature during the cycle: 18,61 °C
- Average Humidity during the cycle: 70,05%

Cluster:




### 3.5 Cycle X1

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
20/01/2014	33.743	18/03/2014	08/07/2015	No	12.375.840

Significant features:

- Average Percentage of Laid Eggs each day: 80,69%
- Average Percentage of Death each day: 0,032%
- Average Temperature during the cycle: 15,34 °C
- Average Humidity during the cycle: 72,83%

Cluster:



### 3.6 Cycle X2


Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
26/05/2014	23.898	15/07/2014	21/06/2015	No	7.558.799

Significant features:

- Average Percentage of Laid Eggs each day: 95,36%
- Average Percentage of Death each day: 0,021%
- Average Temperature during the cycle: 13,76 °C
- Average Humidity during the cycle: 76,04%

Cluster:






### 3.7 Cycle Y

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
11/08/2015	57.346	05/10/2015	27/09/2016	No	16.759.240

Significant features:

- Average Percentage of Laid Eggs each day: 83.70%
- Average Percentage of Death each day: 0,022%
- Average Temperature during the cycle: 14,20 °C
- Average Humidity during the cycle: 74,54%

Cluster:




### 3.8 Cycle Z

Arrival date	#Chickens	Frist Laid	End of cycle	Organic	#Eggs
17/11/2016	42.130	08/01/2017	27/05/2018	Yes	17.721.240

Significant features:


- Average Percentage of Laid Eggs each day: 87,78%
- Average Percentage of Death each day: 0,021%
- Average Temperature during the cycle: 13,07 °C
- Average Humidity during the cycle: 74,27%

Cluster:



## 4 Common features


In this section, we will discuss what we discovered by comparing all the cycles together. After seeing that each cycle can be clustered in two parts, those clusters also divide the cycle into two distinct sections in a plot based on date. We tried to apply the same idea in a dataset composed of all the cycles together.



Looking at the clusters we find that indeed the division of each cycle is respected. We find also that the only one that didn't respect this division is cycle C, which could be expected since it just started, but the interesting thing is that the whole cycle C is clustered as cluster 1 which is the one that identifies the end of every other cycle so we can see how bad is performing this new cycle both in death and laid rate. This can show how the ban on beak-cutting affected the welfare of the chickens, with the beak the violence between animals increased a lot and not only increase the death rate but increased a lot the stress of the chickens reducing their productivity.

To better compare the different cycles we made a spider plot with 4 features:

- Average Percentage of Laid Eggs each day
- Average Percentage of Death each day
- Average Temperature during the cycle
- Average Humidity during the cycle



## 5 Organic vs non-organic cycles

One of the main differences in the cycles under analysis is the breeding method. Cycles A, B, C and Z are grown with organic farming, a method regulated by the European Union with the regulation 589/2008 modified from text 2168/2017 while the cycles Y, X1 and X2 are non-organic. We can summarize the main differences:


Charachteristics	Organic	Non-organic
Chickens/ $m^2$	6	9
Outdoor access	✓	✓
Feed requirements	✓	×

Given the differences in the breeding methods, we analyze if there are implications in the cycles' behavior. Since the non-organic ones have no information on feed and water consumption, we can't go deep into the analysis because of the lack of inputs. For this reason, we will focus mainly on the outputs like the percentage of eggs laid and deaths.

One of the most important outputs is the productivity of a cycle expressed as EggsProduced#Chickens. In the following results, we exclude cycle C since it is in progress. Furthermore, we round the data to the second digit.

Cycle	Organic	Productivity	Breeding days	Productivity/Breeding days
A	✓	481.04	627	0.77
B	✓	436.90	586	0.75
Z	✓	420.63	505	0.83
X1	×	366.77	478	0.77
X2	×	316.29	342	0.92
Y	×	292.25	359	0.81


We can notice that non-organic cycles have a lower EggsProduced/#Chickens rate than the organic ones, but considering the days of breeding the non-organic cycles are better. The medium ProductivityBreeding days of an organic cycle is 0.78 while the one of non-organic is 0.84.



Beyond the productivity in terms of eggs produced there is another important aspect to consider which is the chickens' health. Since it is a difficult feature to value, we decided to consider the chickens' death percentage on the number of chickens as a health indicator.

Cycle	Organic	Deaths on chickens	Breeding days	Daily deaths average
A	✓	0.24	627	16.18
B	✓	0.25	586	18.07
Z	✓	0.10	505	8.53
X1	×	0.14	478	10.10
X2	×	0.07	342	4.93
Y	×	0.07	359	12.03


It is evident that in organic cycles the deaths on chickens' number is much higher than in the non-organic ones. So we decided to better understand the reason of this behavior. This question could have a different ways to be answered or a mix of them: - Different feeds quality - Different water consumption - ... So we decided to look for correlations:



## 6 Death-season correlation


In this section, we will analyze how the period when the chickens are introduced into the barn affects the mortality rate. This effect will be more visible in the organic cycles where the chickens are more fragile due to a lack of nutrients.


First, we plot the Death Rate of all the organic cycles (Z, A, B).



It's clear that the Z cycle outperforms the other two by a lot, so we will analyze what are the differences between those 3 cycles.

The number of chickens at the start of the cycle is identical ( $\sim 42.000$ ), and the feed in an organic farm is standard and checked so it doesn't change between those cycles. The difference in mortality cannot be explained with just "a lucky cycle" so we tried to analyze those graphs deeper.






We see as expected that cycle Z is clustered the same as the start of the other cycles. But the interesting analysis is over the season in which events happen.

- Both cycles A and B start to see an increase in mortality around mid-summer whereas cycle Z just saw a small peak.
- Cycles A and B started around early Autumn whereas cycle Z started in mid-winter

What also happens in the summer is that chickens start to eat less and so become weaker.



After discussing it with the farmer we conclude that the summer is the most challenging period for the chickens since:

- The barns don't have air-conditioning and the Temperature can be quite high.
- The chickens are weaker due to age and the quality of nutrients in the organic feed.


To overcome those problems it seems that starting the cycle in mid-winter improves the mortality rate since during the summer the chickens are younger and stronger.

To put it in perspective, cycle Z had 4308 deaths compared to cycles A and B which had respectively 7449 deaths and 9130 deaths across the same period. This help the production of cycle Z which, in the same period, produced 500.000 more eggs than cycle A and 1.200.000 more eggs than cycle B. Which is respectively an increment of 2% and a 7% in eggs produced.

## 7 Cycle C


Cycle C is a particular cycle, it is an organic cycle so it has the same characteristics as cycles Z, A and B but from this one, a new regulation has been introduced. This new law banned the cut of the beak from the chick which increased the violence between the animals and this led to a higher mortality rate.





A difference of 0,03% means on average that in this period every day die 12 chickens more than the normal rate. This has an effect also on production since the chickens have poorer welfare they produce less.

Using a gradient boost regression we were able to guess the death that this cycle will have in the first 400 days based on the data that we have with the other 2 cycles A and B.



Using this prediction we say that we guess that in the first 400 days the farmer will lose 9210 compared to the 4903 of cycle A and 6369 of cycle B.

- 8 Economic results
- 9 Data processing: shakiba