

期末考试试卷

课程名称: 力学基础 IV (广义相对论) 任课教师: 杨捷

学院: 专业: 年级:

姓名: 校园卡号:

题号	一	二	三	四	五	六	七	总分
分数								

注: 7 道任选 5 道, 每题 20 分, 共 100 分, 第 7 题研究生必做

- 1.计算牛顿近似中的所以非零联络 $\Gamma^{\lambda}_{\mu\nu}$
- 2.写出逆变、协变矢量与二阶张量的的坐标变换式,(已知 $A^{\mu}_{\nu}=\frac{\partial \bar{x}^{\mu}}{\partial x^{\nu}}$ ,  $\bar{A}^{\mu}_{\nu}=\frac{\partial x^{\mu}}{\partial \bar{x}^{\nu}}$ 且 $\det[A^{\mu}_{\nu}]\neq 0$ )并写出它们的协变微分表达式。
- 3.由毕安基恒等式证明 $\left(R^{\mu\nu}-\frac{1}{2}g^{\mu\nu}R\right)_{;\nu}=0$

4.写出爱因斯坦作用量, 并变分得到场方程。

已知 Palatini 公式

$$\sqrt{-g}g^{\mu\nu}\delta R_{\mu\nu}=\partial_{\mu}(\sqrt{-g}\varphi^{\mu})$$

且 $\delta g=gg^{\mu\nu}\delta g_{\mu\nu}=-gg_{\mu\nu}\delta g^{\mu\nu}$ 。

5.求弱场线性近似的里奇张量, 并导出线性化的场方程。

6.推导光线在行星附近的轨道方程, 并讨论其偏折, 这证明了广义相对论的什么效应?

7.推导史瓦西时空与闵可夫斯基时空的类光测地线, 简述并画出 krusal 沿拓图与彭若斯图。

## 参考提示

### 一、

我们在下列近似下化简运动方程(3.4.27)。

(1) 引力场是弱场,即

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (3.4.$$

其中  $\eta_{\mu\nu}$  是闵可夫斯基度规(2.8.10),而

$$|h_{\mu\nu}| \ll 1 \quad (3.4.$$

(2) 引力场是静态的,即

$$g_{\mu\nu,0} = h_{\mu\nu,0} = 0 \quad (3.4.$$

(3) 引力场是空间缓变的,即

$$|g_{\mu\nu,i}| = |h_{\mu\nu,i}| \ll 1, \quad i = 1, 2, 3 \quad (3.4.$$

(4) 粒子作低速运动

$$\left| \frac{dx^i}{dx^0} \right| \ll 1, \quad \left| \frac{dx^i}{d\tau} \right| \ll \left| \frac{dx^0}{d\tau} \right|, \quad \left| \frac{dx^i}{d\tau} \right| \ll c \quad (3.4.$$

注意,在自然单位制下  $c=1$ 。这些近似能把运动方程化成牛顿形式,所以称上述化简为顿近似。

利用近似条件(1)、(2)和(3),略去高于一阶的小量,联络可以化成

$$\Gamma_{\mu\nu}^{\lambda} = \frac{1}{2} \eta^{\lambda\rho} (h_{\rho\mu,\nu} + h_{\rho\nu,\mu} - h_{\mu\nu,\rho}) \quad (3.6.$$

具体算得其分量为

$$\begin{cases} \Gamma_{00}^0 = 0, & \Gamma_{0i}^0 = -\frac{1}{2} h_{00,i}, & \Gamma_{ij}^0 = -\frac{1}{2} (h_{0i,j} + h_{0j,i}), & \Gamma_{00}^i = -\frac{1}{2} h_{00,i} \\ \Gamma_{0j}^i = \frac{1}{2} (h_{0i,j} - h_{0j,i}), & \Gamma_{jk}^i = \frac{1}{2} (h_{ij,k} + h_{ik,j} - h_{jk,i}), & i, j, k = 1, 2, 3 \end{cases} \quad (3.6.$$

### 二、

$$A_{\mu;\nu} = A_{\mu,\nu} - \Gamma_{\mu\nu}^{\lambda} A_{\lambda}$$

$$A_{;\nu}^{\mu} = A_{,\nu}^{\mu} + \Gamma_{\lambda\nu}^{\mu} A^{\lambda}$$

$$T_{\mu\nu;\lambda} = T_{\mu\nu,\lambda} - \Gamma_{\mu\lambda}^{\rho} T_{\rho\nu} - \Gamma_{\nu\lambda}^{\rho} T_{\mu\rho}$$

$$T_{;\lambda}^{\mu\nu} = T_{,\lambda}^{\mu\nu} + \Gamma_{\rho\lambda}^{\mu} T^{\rho\nu} + \Gamma_{\rho\lambda}^{\nu} T^{\mu\rho}$$

$$T_{\nu;\lambda}^{\mu} = T_{\nu,\lambda}^{\mu} + \Gamma_{\rho\lambda}^{\mu} T_{\nu}^{\rho} - \Gamma_{\nu\lambda}^{\rho} T_{\rho}^{\mu}$$

### 三、

现在缩并毕安基恒等式(2.11.25)的指标  $\rho$  和  $\sigma$ ,得

$$R_{\lambda\mu\nu;\sigma}^{\sigma} - R_{\lambda\nu;\mu\sigma} + R_{\lambda\mu;\nu\sigma} = 0 \quad (2.11.30)$$

乘以  $g^{\nu\lambda}$ ,并注意到  $g_{;\alpha}^{\nu\lambda}=0$ ,上式可以化成

$$R_{\mu;\sigma}^{\sigma} - R_{;\mu} + R_{\mu;\nu}^{\nu} = 0 \quad (2.11.31)$$

即

$$R_{\mu;\nu}^{\nu} - \frac{1}{2} R_{;\mu} = 0 \quad (2.11.32)$$

也即

$$\left( R_{\mu}^{\nu} - \frac{1}{2} \delta_{\mu}^{\nu} R \right)_{;\nu} = 0 \quad (2.11.33)$$

(2.11.33)式还可以写成协变或逆变张量的形式

$$\left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right)^{;\nu} = 0 \quad (2.11.34)$$

$$\left( R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R \right)_{;\nu} = 0 \quad (2.11.35)$$

这三个式子表明爱因斯坦张量的协变散度为零

$$G_{\mu;\nu}^{\nu} = G_{\mu\nu}^{;\nu} = G_{\nu}^{\mu\nu} = 0 \quad (2.11.36)$$

爱因斯坦张量的这一性质,对于建立广义相对论的场方程极为重要。

四、

$$I_g = \int_M L_g \sqrt{-g} d^4x, \quad L_g = \frac{c^3}{16\pi G} R$$

$$I_n = \int_M L_n \sqrt{-g} d^4x$$

$$\delta I_m = \frac{1}{2c} \int_M T_{\mu\nu} \delta q_{\mu\nu} \sqrt{-g} d^4x$$

对前一式变分可利用公式得

$$\delta(R\sqrt{-g}) = (\delta g^{\mu\nu}) R_{\mu\nu} \sqrt{-g} + g^{\mu\nu} (\delta R_{\mu\nu}) \sqrt{-g} + g^{\mu\nu} R_{\mu\nu} (\delta \sqrt{-g})$$

$$\delta(R\sqrt{-g}) = \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) \sqrt{-g} \delta g^{\mu\nu} + \partial_\mu (\sqrt{-g} \phi^\mu)$$

故

$$\delta \int_M R \sqrt{-g} d^4x = \int_M \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) \sqrt{-g} \delta g^{\mu\nu} d^4x$$

由此得

$$\begin{aligned} \delta I_m &= \delta I_g + \delta I_n \\ &= \frac{c^3}{16\pi G} \int_M \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) \sqrt{-g} \delta g^{\mu\nu} d^4x \\ &\quad + \frac{1}{2c} \int_M T_{\mu\nu} \delta g_{\mu\nu} \sqrt{-g} d^4x \end{aligned}$$

由最小作用量原理

$$\frac{c^3}{16\pi G} \left( R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) + \frac{1}{2c} T_{\mu\nu} = 0 \quad (241)$$

由此可得 Einstein 引力场方程

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = -\frac{8\pi G}{c^4} T_{\mu\nu} \quad (242)$$

五、作业 3.2,3.3

计算曲率张量时，忽略二阶小量，可得

$$R_{\mu\nu} = \Gamma_{\mu\lambda,\nu}^\lambda - \Gamma_{\mu\nu,\lambda}^\lambda \equiv \frac{1}{2} (h_{\mu\nu,\alpha}^\alpha + h_{,\mu,\nu} - h_{\mu,\nu,\alpha}^\alpha - h_{\nu,\mu,\alpha}^\alpha)$$

场方程为

$$\bar{R}_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} R = -8\pi G T_{\mu\nu}$$

$$\Leftrightarrow 2R_{\mu\nu} - \eta_{\mu\nu}R = -16\pi GT_{\mu\nu}$$

$$\Leftrightarrow \left( h_{\mu\nu,\alpha}^{\alpha} + h_{,\mu,\nu}^{\alpha} - h_{\mu,\nu,\alpha}^{\alpha} - h_{\nu,\mu,\alpha}^{\alpha} \right) - \frac{1}{2} \eta_{\mu\nu} \left( h_{,\alpha}^{\alpha} + h_{,\alpha}^{\alpha} - h_{\beta,\alpha}^{\beta\alpha} - h_{\beta,\alpha}^{\beta\alpha} \right) = -16\pi GT_{\mu\nu}。$$

$$\because h_{\mu,\nu,\alpha}^{\alpha} = \eta^{\alpha\beta} h_{\alpha\mu,\nu,\beta} = h_{\mu\alpha,\nu}^{\alpha}, \quad h_{\nu,\mu,\alpha}^{\alpha} = \eta^{\alpha\beta} h_{\alpha\nu,\mu,\beta} = h_{\nu\alpha,\mu}^{\alpha},$$

$$h_{\beta,\alpha}^{\beta\alpha} = \eta^{\rho\alpha} \eta^{\varphi\beta} h_{\varphi\rho,\beta,\alpha} = h_{\varphi\rho}^{\varphi,\rho} = h_{\alpha\beta}^{\alpha,\beta},$$

$$\therefore -16\pi GT_{\mu\nu} = \left( h_{\mu\nu,\alpha}^{\alpha} + h_{,\mu,\nu}^{\alpha} - h_{\mu,\nu,\alpha}^{\alpha} - h_{\nu,\mu,\alpha}^{\alpha} \right) - \frac{1}{2} \eta_{\mu\nu} \left( h_{,\alpha}^{\alpha} + h_{,\alpha}^{\alpha} - h_{\beta,\alpha}^{\beta\alpha} - h_{\beta,\alpha}^{\beta\alpha} \right)$$

$$= \left( h_{\mu\nu,\alpha}^{\alpha} - \frac{1}{2} \eta_{\mu\nu} h_{,\alpha}^{\alpha} \right) + \frac{1}{2} h_{,\mu,\nu}^{\alpha} + \frac{1}{2} h_{,\mu,\nu}^{\alpha} - h_{\mu\alpha,\nu}^{\alpha} - h_{\nu\alpha,\mu}^{\alpha} - \frac{1}{2} \eta_{\mu\nu} \left( h_{,\alpha}^{\alpha} - 2h_{\alpha\beta}^{\alpha,\beta} \right)$$

$$= \left( h_{\mu\nu,\alpha}^{\alpha} - \frac{1}{2} \eta_{\mu\nu} h_{,\alpha}^{\alpha} \right) - \left( h_{\mu\alpha,\nu}^{\alpha} - \frac{1}{2} h_{,\mu,\nu}^{\alpha} \right) - \left( h_{\nu\alpha,\mu}^{\alpha} - \frac{1}{2} h_{,\nu,\mu}^{\alpha} \right) + \eta_{\mu\nu} \left( h_{\alpha\beta}^{\alpha,\beta} - \frac{1}{2} h_{,\alpha}^{\alpha} \right)$$

$$= \left( h_{\mu\nu,\alpha}^{\alpha} - \frac{1}{2} \eta_{\mu\nu} h_{,\alpha}^{\alpha} \right) + \eta_{\mu\nu} \left( h_{\alpha\beta}^{\alpha,\beta} - \frac{1}{2} \eta_{\alpha\beta} h^{\alpha,\beta} \right) - \left( h_{\mu\alpha,\nu}^{\alpha} - \frac{1}{2} \eta_{\mu\alpha} h_{,\nu}^{\alpha} \right) - \left( h_{\nu\alpha,\mu}^{\alpha} - \frac{1}{2} \eta_{\nu\alpha} h_{,\mu}^{\alpha} \right)$$

$$= \bar{h}_{\mu\nu,\alpha}^{\alpha} + \eta_{\mu\nu} \bar{h}_{\alpha\beta}^{\alpha,\beta} - \bar{h}_{\mu\alpha,\nu}^{\alpha} - \bar{h}_{\nu\alpha,\mu}^{\alpha},$$

$$\therefore \bar{h}_{\mu\nu,\alpha}^{\alpha} + \eta_{\mu\nu} \bar{h}_{\alpha\beta}^{\alpha,\beta} - \bar{h}_{\mu\alpha,\nu}^{\alpha} - \bar{h}_{\nu\alpha,\mu}^{\alpha} = -16\pi GT_{\mu\nu}。$$

故线性化的场方程为

$$\bar{h}_{\mu\nu,\alpha}^{\alpha} + \eta_{\mu\nu} \bar{h}_{\alpha\beta}^{\alpha,\beta} - \bar{h}_{\mu\alpha,\nu}^{\alpha} - \bar{h}_{\nu\alpha,\mu}^{\alpha} = -16\pi GT_{\mu\nu}.$$

六、见赵峥第四章 P96-99 或第三章 PPT3.3

守恒量方程：

$$\frac{dt}{d\lambda} = E \cdot \left( 1 - \frac{2M}{r} \right)^{-1}$$

$$\frac{d\varphi}{d\lambda} = \frac{\tilde{L}}{r^2}$$

$$\left( \frac{dr}{d\lambda} \right)^2 = E^2 - \left( 1 - \frac{2M}{r} \right) \frac{\tilde{L}^2}{r^2}$$

利用上式可得可得轨道

$$\frac{d^2 \tilde{u}}{d\varphi^2} + \tilde{u} = \frac{3GM}{c^2} \tilde{u}^2$$

作近似令修正项为  $A \cos^2 \varphi + B$  可得

$$u = \frac{1}{b} \sin \phi + \frac{GM}{c^2 \cdot b^2} (\cos^2 \phi + 1)$$

利用泰勒展开可得偏转角

$$\delta = \frac{4GM}{c^2 b}$$






图 6.2.5 闵氏时空彭若斯图




图 6.2.6 克鲁斯卡时空彭若斯图

- (1) 类时未来无穷远  $I^+$  :  $r$  有限,  $t \rightarrow +\infty$ ;
- (2) 类时过去无穷远  $I^-$  :  $r$  有限,  $t \rightarrow -\infty$ ;
- (3) 类空无穷远  $I^0$  :  $t$  有限,  $r \rightarrow \infty$ ;
- (4) 类光未来无穷远  $J^+$  :  $(t-r)$  有限,  $(t+r) \rightarrow +\infty$ ;
- (5) 类光过去无穷远  $J^-$  :  $(t+r)$  有限,  $(t-r) \rightarrow -\infty$ 。