
Fast Two-Time-Scale Noisy EM Algorithms

Anonymous Author(s)

Affiliation

Address

email

Abstract

Training latent data models using the EM algorithm is the most popular choice for current learning tasks. For today's modern and complex tasks, variants of the EM have been initially introduced by [Neal and Hinton, 1998], using incremental updates to scale to large dataset, and by [Wei and Tanner, 1990, Delyon et al., 1999], using Monte-Carlo (MC) approximations to bypass the impossible conditional expectation of the latent data for most nonconvex models. In this paper, we propose to combine both techniques in a single class of methods called Two-Time-Scale EM Methods. We motivate the choice of a double dynamics by invoking the variance reduction virtue of each stage of the method on both noises: the incremental update and the MC approximation. We establish finite-time and independent of the initialization convergence bounds for nonconvex objective function. Numerical applications are also presented in this article to illustrate our findings.

1 Introduction

Learning latent data models is critical for modern machine learning problems, see [McLachlan and Krishnan, 2007] for references. We formulate the training of such model as the following empirical risk minimization problem:

$$\min_{\theta \in \Theta} \bar{L}(\theta) := r(\theta) + L(\theta) \quad \text{with} \quad L(\theta) = \frac{1}{n} \sum_{i=1}^n L_i(\theta) := \frac{1}{n} \sum_{i=1}^n \{ -\log g(y_i; \theta) \}, \quad (1)$$

We denote the observations by $\{y_i\}_{i=1}^n$, $\Theta \subset \mathbb{R}^d$ is the convex parameters space. We consider a regularized model where $r : \Theta \rightarrow \mathbb{R}$ is a smooth convex regularization function and for $\theta \in \Theta$, $g(y; \theta)$ is the (incomplete) likelihood of each individual observation. The objective function $\bar{L}(\theta)$ is possibly *nonconvex* and is assumed to be lower bounded $\bar{L}(\theta) > -\infty$ for all $\theta \in \Theta$.

In the latent variable model, $g(y_i; \theta)$, is the marginal of the complete data likelihood defined as $f(z_i, y_i; \theta)$, i.e. $g(y_i; \theta) = \int_{\mathcal{Z}} f(z_i, y_i; \theta) \mu(dz_i)$, where $\{z_i\}_{i=1}^n$ are the (unobserved) latent variables. In this paper, we make the assumption of a complete model belonging to the curved exponential family, i.e.,

$$f(z_i, y_i; \theta) = h(z_i, y_i) \exp(\langle S(z_i, y_i) | \phi(\theta) \rangle - \psi(\theta)), \quad (2)$$

where $\psi(\theta)$, $h(z_i, y_i)$ are scalar functions, $\phi(\theta) \in \mathbb{R}^k$ is a vector function, and $S(z_i, y_i) \in \mathbb{R}^k$ is the complete data sufficient statistics.

Full batch EM [Dempster et al., 1977] is the method of reference for that kind of task and is a two steps procedure. The E-step amounts to computing the conditional expectation of the complete data sufficient statistics,

$$\bar{s}(\theta) = \frac{1}{n} \sum_{i=1}^n \bar{s}_i(\theta) \quad \text{where} \quad \bar{s}_i(\theta) = \int_{\mathcal{Z}} S(z_i, y_i) p(z_i | y_i; \theta) \mu(dz_i). \quad (3)$$

30 The M-step is given by

$$\text{M-step: } \hat{\theta} = \bar{\theta}(\bar{s}(\theta)) := \arg \min_{\vartheta \in \Theta} \{r(\vartheta) + \psi(\vartheta) - \langle \bar{s}(\theta) | \phi(\vartheta) \rangle\}, \quad (4)$$

31 Two caveats of this method are the following: (a) with the explosion of data, the first step of the EM
 32 is computationally inefficient as it requires a full pass over the dataset at each iteration and (b) the
 33 complexity of modern models makes the expectation intractable. So far, both challenges have been
 34 addressed separately, to the best of our knowledge, and we give an overview of current solutions in
 35 the sequel.

36 **Prior Work** Inspired by stochastic optimization procedures, [Neal and Hinton, 1998] and [Cappé
 37 and Moulines, 2009] developed respectively an incremental and an online variant of the E-step in
 38 models where the expectation is computable then extensively used and studied in [Nguyen et al.,
 39 2020, Liang and Klein, 2009, Cappé, 2011]. Some improvements of that methods have been pro-
 40 vided and analyzed, globally and in finite-time, in [Karimi et al., 2019] where variance reduction
 41 techniques taken from the optimization literature have been efficiently applied to scale the EM algo-
 42 rithm to large datasets.

43 Regarding the computation of the expectation under the posterior distribution, the first method was
 44 the Monte-Carlo EM (MCEM) introduced in the seminal paper [Wei and Tanner, 1990] where a MC
 45 approximation of this expectation is computed. A variant of that method is the Stochastic Approx-
 46 imation of the EM (SAEM) in [Delyon et al., 1999] leveraging the power of Robbins-Monro type of
 47 update [Robbins and Monro, 1951] to ensure pointwise convergence of the vector of estimated pa-
 48 rameters rather using a decreasing stepsize than increasing the number of MC samples. The MCEM
 49 and the SAEM have been successfully applied in mixed effects models [McCulloch, 1997, Hughes,
 50 1999, Baey et al., 2016] or to do inference for joint modelling of time to event data coming from
 51 clinical trials in [Chakraborty and Das, 2010], among other applications.

52 Recently, an incremental variant of the SAEM was proposed in [Kuhn et al., 2019] showing positive
 53 empirical results but its analysis is limited to asymptotic consideration. Gradient-based methods
 54 have been developed and analyzed in [Zhu et al., 2017] but they remain out of the scope of this
 55 paper as they tackle the high-dimensionality issue.

56 **Contributions** This paper *introduces* and *analyzes* a new class of methods which purpose is to
 57 combine both solutions proposed in the past years in a two-time-scale manner in order to optimize
 58 (1) for current modern examples and settings. The main contributions of the paper are:

- 59 • We propose a two-time-scale method based on Stochastic Approximation (SA), to alleviate
 60 the problem of MC computation, and on Incremental updates, to scale to large datasets.
 61 We describe in details the edges of each level of our method based on variance reduc-
 62 tion arguments. The derivation of such class of algorithms has two advantages. First, it
 63 combines two powerful ideas, commonly used separately, to tackle large scale and highly
 64 nonlinear learning tasks. Then, it gives a simple formulation as a *scaled-gradient method*,
 65 as introduced in [Karimi et al., 2019], which makes the global analysis accessible.
- 66 • We also establish global (independent of the initialization) and finite-time (true at each
 67 iteration) upper bounds on a classical suboptimality condition in the nonconvex literature,
 68 *i.e.*, the second order moment of the gradient of the objective function.

69 In Section 2 we give rigorous mathematical definitions of the various updates used for both incre-
 70 mental and Monte-Carlo EMs and we introduce the main class of new algorithms, based on two
 71 different dynamics, we are proposing to analyze and compare to baselines algorithms. Section 3
 72 presents the main theoretical guarantees of this newly introduced two-time-scale class of algorithms.
 73 Results are given both in finite-time and in the nonconvex setting. Finally, we illustrate the advan-
 74 tages of our method in Section 4 on two numerical experiments.

75 2 Two-Time-Scale Stochastic EM Algorithms

76 We recall and formalize in this section the different methods found in the literature that aim to solv-
 77 ing the large scale problem and the intractable expectation. We then provide the general framework
 78 of our method to efficiently tackle the optimization problem (1).

79 2.1 Monte Carlo Integration and Stochastic Approximation

80 As mentioned in the introduction, for complex and possibly nonlinear models, the expectation under
 81 the posterior distribution defined in (3) is not tractable. In that case, the first solution involves
 82 computing a Monte Carlo integration of that latter term. For all $i \in \llbracket 1, n \rrbracket$, draw for $m \in \llbracket 1, M \rrbracket$,
 83 samples $z_{i,m} \sim p(z_i|y_i; \theta)$ and compute the MC integration \tilde{s} of the deterministic quantity $\bar{s}(\theta)$:

$$\text{MC-step : } \tilde{s} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}, y_i) \quad (5)$$

84 and then update the parameter $\hat{\theta} = \bar{\theta}(\tilde{s})$. This algorithm bypasses the intractable expectation issue
 85 but is rather computationally expensive in order to reach point wise convergence (M needs to be
 86 large). An alternative to that stochastic algorithm is to use a Robbins-Monro (RM) type of update.
 87 We denote, at iteration k , the following quantity

$$\tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}^{(k)}, y_i) \quad \text{where } z_{i,m}^{(k)} \sim p(z_i|y_i; \theta^{(k)}) \quad (6)$$

88 Then, the RM updated of the sufficient statistics $\hat{s}^{(k+1)}$ reads:

$$\text{SA-step : } \hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad (7)$$

89 where $\{\gamma_k\}_{k \geq 1} \in (0, 1)$ is a sequence of decreasing step sizes to ensure asymptotic convergence.
 90 This is called the Stochastic Approximation of the EM (SAEM) and has been shown theoretically to
 91 converge to a maximum of the likelihood of the observations under very general conditions [Delyon
 92 et al., 1999]. In the simulation step (6), since the relation between the observed data y_i and the
 93 latent variable z_i can be non linear, sampling from the posterior distribution $p(z_i|y_i; \theta)$, under the
 94 current model θ , could require using an inference algorithm. [?] proved almost sure convergence
 95 of the sequence of parameters obtained by this algorithm coupled with an MCMC procedure during
 96 the simulation step. In simple scenarios, the samples $\{z_{i,m}\}_{m=0}^{M-1}$ are conditionally independent and
 97 identically distributed with distribution $p(z_i, \theta)$. Nevertheless, in most cases, sampling exactly from
 98 this distribution is not an option and the Monte Carlo batch is sampled by Monte Carlo Markov
 99 Chains (MCMC) algorithm. In the SA-step, the sequence of decreasing positive integers $\{\gamma_k\}_{k \geq 1}$
 100 controls the convergence of the algorithm. In practice, γ_k is set equal to 1 during the first few
 101 iterations to let the algorithm explore the parameter space without memory and converge quickly
 102 to a neighbourhood of the target estimate. The Stochastic Approximation is performed during the
 103 remaining iterations where $\gamma_k = 1/k^\alpha$, where $\alpha \in (0, 1)$, ensuring the almost sure convergence of
 104 the estimate. It is inappropriate to start with small values for step size γ_k and large values for the
 105 number of simulations M_k . Rather, it is recommended that one decrease γ_k and keep a constant
 106 and small number of MC samples M_k which shows a great advantage over the MC-step (5), which
 107 requires large M_k to converge.

108 This Robbins-Monro type of update represents the *first level* of our algorithm, needed to temper
 109 the variance and noise implied by MC integration. In the next section, we derive variants of this
 110 algorithm to adapt to the sheer size of data of today's applications and formalize the *second level* of
 111 our class of Two-Time-Scale EM methods.

112 2.2 Incremental and Bi-Level Inexact EM Methods

113 Strategies to scale to large datasets include classical incremental and variance reduced variants. We
 114 will explicit a general update that will cover those variants and that represents the *second level* of our
 115 algorithm, namely the incremental update of the noisy statistics $\hat{S}^{(k)}$ inside the RM type of update.

$$\text{Incremental-step : } \tilde{S}^{(k+1)} = \tilde{S}^{(k)} + \rho_{k+1}(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}), \quad (8)$$

116 Note $\{\rho_k\}_{k \geq 1} \in (0, 1)$ is a sequence of step sizes, $\mathcal{S}^{(k)}$ is a proxy for $\tilde{S}^{(k)}$, If the stepsize is equal
 117 to one and the proxy $\mathcal{S}^{(k)} = \hat{S}^{(k)}$, i.e., computed in a full batch manner as in (6), then we recover
 118 the SAEM algorithm. Also if $\rho_k = 1$, $\gamma_k = 1$ and $\mathcal{S}^{(k)} = \tilde{S}^{(k)}$, then we recover the Monte Carlo
 119 EM algorithm.

We now introduce three variants of the SAEM update depending on different definitions of the proxy $\mathcal{S}^{(k)}$ and the choice of the stepsize ρ_k . Let $i_k \in \llbracket 1, n \rrbracket$ be a random index drawn at iteration k and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ be the iteration index where $i \in \llbracket 1, n \rrbracket$ is last drawn prior to iteration k . For iteration $k \geq 0$, the fiTTSEM method draws *two* indices *independently* and uniformly as $i_k, j_k \in \llbracket 1, n \rrbracket$. In addition to τ_i^k which was defined w.r.t. i_k , we define $t_j^k = \{k' : j_{k'} = j, k' < k\}$ to be the iteration index where the sample $j \in \llbracket 1, n \rrbracket$ is last drawn as j_k prior to iteration k . With the initialization $\bar{\mathcal{S}}^{(0)} = \bar{s}^{(0)}$, we use a slightly different update rule from SAGA inspired by [Reddi et al., 2016]. Then, we obtain:

$$(iSAEM [Karimi, 2019, Kuhn et al., 2019]) \quad \mathcal{S}^{(k+1)} = \mathcal{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)}) \quad (9)$$

$$(vrTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}) \quad (10)$$

$$(fiTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \quad (11)$$

$$\bar{\mathcal{S}}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + n^{-1} (\tilde{S}_{j_k}^{(k)} - \tilde{S}_{j_k}^{(t_{j_k}^k)}). \quad (12)$$

The stepsize is set to $\rho_{k+1} = 1$ for the iSAEM method; $\rho_{k+1} = \gamma$ is constant for the vrTTSEM and fiTTSEM methods. Moreover, for iSAEM we initialize with $\mathcal{S}^{(0)} = \tilde{S}^{(0)}$; for vrTTSEM we set an epoch size of m and define $\ell(k) := m \lfloor k/m \rfloor$ as the first iteration number in the epoch that iteration k is in.

2.3 Two-Time-Scale Noisy EM methods

We now introduce the general method derived using the two variance reduction techniques described above. Algorithm 1 leverages both levels (7) and (8) in order to output a vector of fitted parameters $\hat{\theta}^{(K)}$ where K is some randomly chosen termination point.

The update in (14) is said to have two timescales as the step sizes satisfy $\lim_{k \rightarrow \infty} \gamma_k / \rho_k < 1$ such that $\tilde{S}^{(k+1)}$ is updated at a faster timescale than $\hat{s}^{(k+1)}$.

Algorithm 1 Two-Time-Scale Noisy EM methods.

- 1: **Input:** initializations $\hat{\theta}^{(0)} \leftarrow 0, \hat{s}^{(0)} \leftarrow \hat{S}^{(0)}, K_{\max} \leftarrow \text{max. iteration number}$.
- 2: Set the terminating iteration number, $K \in \{0, \dots, K_{\max} - 1\}$, as a discrete r.v. with:

$$P(K = k) = \frac{\gamma_k}{\sum_{\ell=0}^{K_{\max}-1} \gamma_{\ell}}. \quad (13)$$

- 3: **for** $k = 0, 1, 2, \dots, K$ **do**
- 4: Draw index $i_k \in \llbracket 1, n \rrbracket$ uniformly (and $j_k \in \llbracket 1, n \rrbracket$ for fiTTSEM).
- 5: Compute $\hat{S}_{i_k}^{(k)}$ using the MC-step (5), for the drawn indices.
- 6: Compute the surrogate sufficient statistics $\mathcal{S}^{(k+1)}$ using (9) or (10) or (11).
- 7: Compute $\hat{S}^{(k+1)}$ and $\hat{s}^{(k+1)}$ using respectively (8) and (7):

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho_{k+1} (\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{s}^{(k+1)} &= \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \end{aligned} \quad (14)$$

- 8: Compute $\hat{\theta}^{(k+1)}$ via the M-step (4).
 - 9: **end for**
 - 10: **Return:** $\hat{\theta}^{(K)}$.
-

3 Finite Time Analysis of the Two-Time-Scale Scheme

First, we consider the following minimization problem on the statistics space:

$$\min_{\mathbf{s} \in \mathcal{S}} V(\mathbf{s}) := \bar{\mathcal{L}}(\bar{\theta}(\mathbf{s})) = \mathbf{r}(\bar{\theta}(\mathbf{s})) + \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(\bar{\theta}(\mathbf{s})) \quad (15)$$

140 It has been shown that this minimization problem is equivalent to the optimization problem (1), see
 141 [Karimi et al., 2019, Lemma2]

142 **H1.** Θ is an open set of \mathbb{R}^d and the sets Z, S are measurable open sets such that:

$$S \supset \left\{ n^{-1} \sum_{i=1}^n u_i, u_i \in \text{conv}(\bar{s}_i(\theta)) \right\} \quad (16)$$

143 where $\bar{s}_i(\theta)$ is defined in (3).

144 **H2.** The conditional distribution is smooth on $\text{int}(\Theta)$. For any $i \in \llbracket 1, n \rrbracket$, $z \in Z$, $\theta, \theta' \in \text{int}(\Theta)^2$,
 145 we have $|p(z|y_i; \theta) - p(z|y_i; \theta')| \leq L_p \|\theta - \theta'\|$.

146 We also recall from the introduction that we consider curved exponential family models. besides:

147 **H3.** For any $s \in S$, the function $\theta \mapsto L(s, \theta) := r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$ admits a unique global
 148 minimum $\bar{\theta}(s) \in \text{int}(\Theta)$. In addition, $J_\phi^\theta(\bar{\theta}(s))$ is full rank and $\bar{\theta}(s)$ is L_θ -Lipschitz.

149 We denote by $H_L^\theta(s, \theta)$ the Hessian (w.r.t to θ for a given value of s) of the function $\theta \mapsto L(s, \theta) =$
 150 $r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$, and define

$$B(s) := J_\phi^\theta(\bar{\theta}(s)) \left(H_L^\theta(s, \bar{\theta}(s)) \right)^{-1} J_\phi^\theta(\bar{\theta}(s))^\top. \quad (17)$$

151 **H4.** It holds that $v_{\max} := \sup_{s \in S} \|B(s)\| < \infty$ and $0 < v_{\min} := \inf_{s \in S} \lambda_{\min}(B(s))$. There exists
 152 a constant L_B such that for all $s, s' \in S^2$, we have $\|B(s) - B(s')\| \leq L_B \|s - s'\|$.

153 We now formulate the main difference with the work done in [Karimi et al., 2019]. The class of
 154 algorithms we develop in this paper are two time-scale where the first stage corresponds to the
 155 variance reduction trick used in [Karimi et al., 2019] in order to accelerate incremental methods and
 156 reduce the variance induced by the index sampling. The second stage is the Robbins-Monro type of
 157 update that aims to reduce the variance induced by the MC approximations

158 Indeed the expectations (3) are never available and requires Monte Carlo approximation. Thus, at
 159 iteration $k + 1$, we introduce the errors when approximating the quantity $\bar{s}_i(\hat{\theta}(\hat{s}^{(k-1)}))$. For all
 160 $i \in \llbracket 1, n \rrbracket$, $r > 0$ and $\vartheta \in \Theta$, define:

$$\eta_i^{(r)} := \tilde{S}_i^{(r)} - \bar{s}_i(\vartheta^{(r)}) \quad (18)$$

161 For instance, we consider that the MC approximation is unbiased if for all $i \in \llbracket 1, n \rrbracket$ and $m \in$
 162 $\llbracket 1, M \rrbracket$, the samples $z_{i,m} \sim p(z_i|y_i; \theta)$ are i.i.d. under the posterior distribution, i.e., $\mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] = 0$
 163 where \mathcal{F}_r is the filtration up to iteration r . The following results are derived under the assumption
 164 of control of the fluctuations implied by the approximation stated as follows:

165 **H5.** There exist a positive sequence of MC batch size $\{M_r\}_{r>0}$ and constants (C, C_η) such that for
 166 all $k > 0$, $i \in \llbracket 1, n \rrbracket$ and $\vartheta \in \Theta$:

$$\mathbb{E} \left[\left\| \eta_i^{(r)} \right\|^2 \right] \leq \frac{C_\eta}{M_r} \quad \text{and} \quad \mathbb{E} \left[\left\| \mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] \right\|^2 \right] \leq \frac{C}{M_r} \quad (19)$$

167 In that setting, we can prove two important results on the Lyapunov function. The first one suggests
 168 smoothness:

169 **Lemma 1.** [Karimi et al., 2019] Assume H2, H3, H4. For all $s, s' \in S$ and $i \in \llbracket 1, n \rrbracket$, we have

$$\|\bar{s}_i(\bar{\theta}(s)) - \bar{s}_i(\bar{\theta}(s'))\| \leq L_s \|s - s'\|, \quad \|\nabla V(s) - \nabla V(s')\| \leq L_V \|s - s'\|, \quad (20)$$

170 where $L_s := C_Z L_p L_\theta$ and $L_V := v_{\max}(1 + L_s) + L_B C_S$.

171 and the second one suggests a growth condition on the gradient of V depending on the mean field
 172 of the algorithm:

173 **Lemma 2.** Assume H3, H4. For all $s \in S$,

$$v_{\min}^{-1} \langle \nabla V(s) | s - \bar{s}(\bar{\theta}(s)) \rangle \geq \|s - \bar{s}(\bar{\theta}(s))\|^2 \geq v_{\max}^{-2} \|\nabla V(s)\|^2, \quad (21)$$

174 See proofs of this Lemma in Appendix A.

3.1 Global Convergence of Incremental Noisy EM Algorithms

We present in this section a finite-time analysis of the incremental variant of the Stochastic Approximation of the EM algorithm. We want to draw the attention of the readers that the word "global" here does not mean for a global optimum of the nonconvex function, but of the independence of our analysis on the initialization and the iteration k (finite time).

The first intermediate result is the computation of the quantity $\hat{S}^{(k+1)} - \hat{s}^{(k)}$, which corresponds to the drift term of (7) and reads as follows:

Lemma 3. Assume H1. The update (9) is equivalent to the following update on the resulting statistics

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad \text{where} \quad \tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^{k+1})} \quad (22)$$

Also:

$$\mathbb{E} [\tilde{S}^{(k+1)} - \hat{s}^{(k)}] = \mathbb{E} [\bar{s}^{(k)} - \hat{s}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \quad (23)$$

where $\bar{s}^{(k)}$ is defined by (3) and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$.

See proofs of this Lemma in Appendix B.

The following main result for the iSAEM algorithm is derived under a control of the Monte Carlo fluctuations as described by assumption H 5. Typically, the controls exhibited below are of interest when the number of MC samples M_k increase with the iteration index f .

Theorem 1. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and consider the iSAEM sequence $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = 1$ for any $k > 0$. We also set $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$. Assume that $\hat{s}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$.

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} [\|\nabla V(\hat{s}^{(k)})\|^2] \leq \mathbb{E} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K)})] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} [\|\eta_{i_k}^{(k)}\|^2] \quad (24)$$

See proof in Appendix C.

3.2 Global Convergence of Two-Time-Scale Noisy EM Algorithms

We now proceed by giving our main result regarding the global convergence of the fiTTSEM algorithm. Two important auxiliary Lemmas, which proofs are given in Appendix D.1, are need in order to derive our finite-time bound. The first one derives an identity for the quantity $\mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2]$ using the vrTTSEM update:

Lemma 4. For any $k \geq 0$ and consider the vrTTSEM update in (10) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned} \mathbb{E} [\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E} [\|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E} [\|\hat{s}^{(k)} - \hat{s}^{(\ell(k))}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E} [\|\hat{s}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E} [\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (25)$$

where we recall that $\ell(k)$ is the first iteration number in the epoch that iteration k is in.

The second one derives an identity for the quantity $\mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2]$ using the fiTTSEM update:

Lemma 5. For any $k \geq 0$ and consider the fiTTSEM update in (11) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned} \mathbb{E} [\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E} [\|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E} [\|\hat{s}^{(k)} - \hat{s}^{(t_i^k)}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E} [\|\hat{s}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E} [\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (26)$$

206 We now state the main result regarding the vrTTSEM method.

207 **Theorem 2.** Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes
 208 and consider the vrTTSEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$.

209 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\bar{L} = \max\{L_S, L_V\}$, $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$
 210 and a constant $\mu \in (0, 1)$, we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (27)$$

211 See proof in Appendix E. We now state the main result regarding the fiTTSEM method.

212 **Theorem 3.** Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes
 213 and consider the fiTTSEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$.

214 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_S, L_V\}$,
 215 $\beta = \frac{c_1 \bar{L}}{n}$, $\rho = \frac{1}{n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$, we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (28)$$

216 See proof in Appendix F. Note that in those two bounds, the quantities $\tilde{\eta}^{(k+1)}$ and $\Xi^{(k+1)}$ are
 217 abstraction that depends only on the MC fluctuations $\mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]$ and some constants.

218 **Remarks:** The following remarks are worth noting on the quantity $\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right]$:

- 219 • This term is the price we pay for the two time scale dynamics and corresponds to the gap
 220 between the two asynchronous updates (one is on $\hat{\mathbf{s}}^{(k)}$ and the other on $\tilde{S}^{(k)}$).
- It is trivial to see that if $\rho = 1$, i.e., there is no variance reduction, then

$$\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] = \mathbb{E} \left[\left\| \mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)} \right\|^2 \right] = 0 \quad \text{with} \quad \hat{\mathbf{s}}^{(0)} = \tilde{S}^{(0)} = 0$$

221 which strengthen the fact that this quantity characterizes the impact of the variance reduc-
 222 tion technique introduced in our two stages class of methods.

223 The following lemma, which proof can be found in Appendix D.2, can be derived to characterize
 224 this gap:

225 **Lemma 6.** Consider a decreasing stepsize $\gamma_k \in (0, 1)$ and a constant $\rho \in (0, 1)$, then the following
 226 inequality holds:

$$\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_{\ell})^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}) \quad (29)$$

227 where $\mathcal{S}^{(k)}$ is defined either by (10) (vrTTSEM) or (11) (fiTTSEM).

4 Numerical Examples

4.1 Gaussian Mixture Models

Given n observations $\{y_i\}_{i=1}^n$, we want to fit a Gaussian Mixture Model (GMM) whose distribution is modeled as a Gaussian mixture of M components, each with a unit variance. Let $z_i \in \llbracket M \rrbracket$ be the latent labels of each component, the complete log-likelihood is defined as:

$$\log f(z_i, y_i; \boldsymbol{\theta}) = \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) [\log(\omega_m) - \mu_m^2/2] + \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) \mu_m y_i + \text{constant} . \quad (30)$$

where $\boldsymbol{\theta} := (\boldsymbol{\omega}, \boldsymbol{\mu})$ with $\boldsymbol{\omega} = \{\omega_m\}_{m=1}^{M-1}$ are the mixing weights with the convention $\omega_M = 1 - \sum_{m=1}^{M-1} \omega_m$ and $\boldsymbol{\mu} = \{\mu_m\}_{m=1}^M$ are the means. We use the penalization $r(\boldsymbol{\theta}) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \log \text{Dir}(\boldsymbol{\omega}; M, \epsilon)$ where $\delta > 0$ and $\text{Dir}(\cdot; M, \epsilon)$ is the M dimensional symmetric Dirichlet distribution with concentration parameter $\epsilon > 0$. The constraint set on $\boldsymbol{\theta}$ is given by

$$\Theta = \{\omega_m, m = 1, \dots, M-1 : \omega_m \geq 0, \sum_{m=1}^{M-1} \omega_m \leq 1\} \times \{\mu_m \in \mathbb{R}, m = 1, \dots, M\}. \quad (31)$$

Exact two time scale updates are given in Appendix G.1.

In the following experiments on synthetic data, we generate samples from a GMM model with $M = 2$ components with two mixtures with means $\mu_1 = -\mu_2 = 0.5$. We use $n = 10^5$ synthetic samples and run the bEM method until convergence (to double precision) to obtain the ML estimate μ^* averaged on 50 datasets. We compare the bEM, SAEM, iSAEM, vrTTSEM and fiTTSEM methods in terms of their precision measured by $|\mu - \mu^*|^2$. We set the step-size of the SA-step of all method as $\gamma_k = 1/k^\alpha$ with $\alpha = 0.5$, and the stepsizes of the Incremental-step for vrTTSEM and the fiTTSEM to a constant stepsize equal to $1/n^{2/3}$.

The number of MC samples is fixed to $M = 10$ chains. Figure 1 shows the convergence of the precision $|\mu - \mu^*|^2$ for the different methods against the epoch(s) elapsed (one epoch equals n iterations). We observe that the vrTTSEM and fiTTSEM methods outperform the other methods, supporting our analytical results.




Figure 1: TO COMPLETE

4.2 Deformable Template Model for Image Analysis

Let $(y_i, i \in \llbracket 1, n \rrbracket)$ be observed gray level images defined on a grid of pixels. Let $u \in \mathcal{U} \subset \mathbb{R}^2$ denotes the pixel index on the image and $x_u \in \mathcal{D} \subset \mathbb{R}^2$ its location. The model used in this experiment suggests that each image y_i is a deformation of a template, noted $I : \mathcal{D} \rightarrow \mathbb{R}$, common to all images of the dataset:

$$y_i(u) = I(x_u - \Phi_i(x_u, z_i)) + \varepsilon_i(u) \quad (32)$$

where $\phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a deformation function, z_i some latent variable parametrizing this deformation and $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$ is an observation error.

The template model, given $(p_k, k \in \llbracket 1, k_p \rrbracket)$ landmarks on the template, a fixed known kernel \mathbf{K}_p and a vector of parameters $\beta \in \mathbb{R}^{k_p}$ is defined as follows:

$$I_\xi = \mathbf{K}_p \beta, \quad \text{where} \quad (\mathbf{K}_p \beta)(x) = \sum_{k=1}^{k_p} \mathbf{K}_p(x, p_k) \beta_k \quad (33)$$

Besides, we parameterize the deformation model given some landmarks $(g_k, k \in \llbracket 1, k_g \rrbracket)$ and a fixed kernel \mathbf{K}_g as:

$$\Phi_i = \mathbf{K}_g z_i \quad \text{where} \quad (\mathbf{K}_g z_i)(x) = \sum_{k=1}^{k_g} \mathbf{K}_g(x, g_k) \left(z_i^{(1)}(k), z_i^{(2)}(k) \right) \quad (34)$$

265 where we put a Gaussian prior on the latent variables, $z_i \sim \mathcal{N}(0, \Gamma)$ and $z_i \in (\mathbb{R}^{k_g})^2$. The vector
 266 of parameters we ought to estimate is thus $\theta = (\beta, \Gamma, \sigma)$. The complete model belongs to the
 267 curved exponential family, see [Allasonnière et al., 2007], which vector of sufficient statistics $S =$
 268 $(S_1(z), S_2(z), S_3(z))$ read:

$$\begin{aligned} S_1(z) &= \sum_{i=1}^n S_1(y_i, z_i) = \sum_{i=1}^n (\mathbf{K}_p^{z_i})^t y_i \\ S_2(z) &= \sum_{i=1}^n S_2(y_i, z_i) = \sum_{i=1}^n (\mathbf{K}_p^{z_i})^t (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \sum_{i=1}^n S_3(y_i, z_i) = \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (35)$$

269 where for any pixel $u \in \mathbb{R}^2$ and $j \in \llbracket 1, k_g \rrbracket$ we noted:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j) \quad (36)$$

270 Finally, the Two-Time-Scale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z) \hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n} \hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z) \beta(\hat{s}) - 2\beta(\hat{s}) \hat{s}_1(z) \end{pmatrix} \quad (37)$$

271 where $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$ is the vector of statistics obtained via the **SA-step** (7) and using the
 272 MC approximation of the sufficient statistics $(S_1(z), S_2(z), S_3(z))$ defined in (35).

273 **Comparison using epochs credit**

274 **Comparison using number of training samples credit**

275 **5 Conclusion**

References

- S. Allasonnière, Y. Amit, and A. Trouvé. Towards a coherent statistical framework for dense deformable template estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):3–29, 2007.
- S. Allasonnière, E. Kuhn, A. Trouvé, et al. Construction of bayesian deformable models via a stochastic approximation algorithm: a convergence study. *Bernoulli*, 16(3):641–678, 2010.
- C. Baey, S. Trevezas, and P.-H. Cournède. A non linear mixed effects model of plant growth and estimation via stochastic variants of the em algorithm. *Communications in Statistics-Theory and Methods*, 45(6):1643–1669, 2016.
- O. Cappé. Online em algorithm for hidden markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- O. Cappé and E. Moulines. On-line expectation–maximization algorithm for latent data models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):593–613, 2009.
- A. Chakraborty and K. Das. Inferences for joint modelling of repeated ordinal scores and time to event data. *Computational and mathematical methods in medicine*, 11(3):281–295, 2010.
- B. Delyon, M. Lavielle, and E. Moulines. Convergence of a stochastic approximation version of the em algorithm. *Ann. Statist.*, 27(1):94–128, 03 1999. doi: 10.1214/aos/1018031103. URL <https://doi.org/10.1214/aos/1018031103>.
- A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the royal statistical society. Series B (methodological)*, pages 1–38, 1977.
- J. P. Hughes. Mixed effects models with censored data with application to hiv rna levels. *Biometrics*, 55(2):625–629, 1999.
- B. Karimi. *Non-Convex Optimization for Latent Data Models: Algorithms, Analysis and Applications*. PhD thesis, 2019.
- B. Karimi, H.-T. Wai, É. Moulines, and M. Lavielle. On the global convergence of (fast) incremental expectation maximization methods. In *Advances in Neural Information Processing Systems*, pages 2833–2843, 2019.
- E. Kuhn, C. Matias, and T. Rebafka. Properties of the stochastic approximation em algorithm with mini-batch sampling. *arXiv preprint arXiv:1907.09164*, 2019.
- P. Liang and D. Klein. Online em for unsupervised models. In *Proceedings of human language technologies: The 2009 annual conference of the North American chapter of the association for computational linguistics*, pages 611–619, 2009.
- C. E. McCulloch. Maximum likelihood algorithms for generalized linear mixed models. *Journal of the American statistical Association*, 92(437):162–170, 1997.
- G. McLachlan and T. Krishnan. *The EM algorithm and extensions*, volume 382. John Wiley & Sons, 2007.
- R. M. Neal and G. E. Hinton. A view of the EM algorithm that justifies incremental, sparse, and other variants. In *Learning in graphical models*, pages 355–368. Springer, 1998.
- H. D. Nguyen, F. Forbes, and G. J. McLachlan. Mini-batch learning of exponential family finite mixture models. *Statistics and Computing*, pages 1–18, 2020.

- 317 S. J. Reddi, S. Sra, B. Póczos, and A. Smola. Fast incremental method for nonconvex optimization.
318 *arXiv preprint arXiv:1603.06159*, 2016.
- 319 H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statis-*
320 *tics*, pages 400–407, 1951.
- 321 G. C. Wei and M. A. Tanner. A monte carlo implementation of the em algorithm and the poor man’s
322 data augmentation algorithms. *Journal of the American statistical Association*, 85(411):699–704,
323 1990.
- 324 R. Zhu, L. Wang, C. Zhai, and Q. Gu. High-dimensional variance-reduced stochastic gradient
325 expectation-maximization algorithm. In *Proceedings of the 34th International Conference on*
326 *Machine Learning-Volume 70*, pages 4180–4188. JMLR. org, 2017.

A Proof of Lemma 2

Lemma. Assume H3, H4. For all $\mathbf{s} \in \mathcal{S}$,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (38)$$

Proof Using H3 and the fact that we can exchange integration with differentiation and the Fisher's identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (39)$$

Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \mathbf{s}. \quad (40)$$

Taking the gradient of the above map w.r.t. \mathbf{s} and using assumption H3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left(\nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})}}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}). \quad (41)$$

The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))) \quad (42)$$

where we recall $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left(\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top}$. The proof of (38) follows directly from the assumption H4. \square

B Proof of Lemma 3

Lemma. Assume H1. The update (9) is equivalent to the following update on the resulting statistics

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}) \quad (43)$$

Also:

$$\mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \quad (44)$$

where $\bar{\mathbf{s}}^{(k)}$ is defined by (3) and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$.

Proof From update (9), we have:

$$\begin{aligned} \tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left(\tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned} \quad (45)$$

Since $\tilde{S}_{i_k}^{(k+1)} = \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) + \eta_{i_k}^{(k+1)}$ we have

$$\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)} \quad (46)$$

Taking the full expectation of both side of the equation leads to:

$$\begin{aligned} \mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] &= \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] \\ &\quad - \frac{1}{n} \mathbb{E} \left[\mathbb{E} [\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \end{aligned} \quad (47)$$

The following equalities:

$$\mathbb{E} [\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E} [\bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] = \bar{\mathbf{s}}^{(k)} \quad (48)$$

concludes the proof of the Lemma. \square

346 C Proof of Theorem 1

347 **Theorem.** Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and
 348 consider the iSAEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = 1$ for any $k > 0$. We also set
 349 $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$. Assume that
 350 $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$.

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (49)$$

351 **Proof** We begin our proof by giving this auxiliary Lemma setting an upper bound for the quantity
 352 $\mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$

353 **Lemma 7.** For any $k \geq 0$ and consider the iSAEM update in (9), it holds that

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &\leq 4\mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \frac{2L_{\mathbf{s}}^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right] \\ &\quad + 2\frac{C_{\eta}}{M_k} + 4\mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \end{aligned} \quad (50)$$

354 **Proof** Applying the iSAEM update yields:

$$\begin{aligned} \mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &= \mathbb{E} \left[\left\| \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} - \frac{1}{n} (\tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(k)}) \right\|^2 \right] \\ &\leq 4\mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + 4\mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\quad + \frac{2}{n^2} \mathbb{E} \left[\left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] + 2\frac{C_{\eta}}{M_k} \end{aligned} \quad (51)$$

355 The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E} \left[\left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\left\| \bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)} \right\|^2 \right] \stackrel{(a)}{\leq} \frac{2L_{\mathbf{s}}^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right], \quad (52)$$

356 where (a) is due to Lemma 1 and which concludes the proof of the Lemma.

357 □

358 Under the smoothness of the Lyapunov function V (cf. Lemma 1), we can write:

$$V(\hat{\mathbf{s}}^{(k+1)}) \leq V(\hat{\mathbf{s}}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \quad (53)$$

359 Taking the expectation on both sides yields:

$$\mathbb{E} \left[V(\hat{\mathbf{s}}^{(k+1)}) \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) \right] + \gamma_{k+1} \mathbb{E} \left[\langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (54)$$

360 Using Lemma 3, we obtain:

$$\begin{aligned}
& \mathbb{E} \left[\langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] = \\
& \mathbb{E} \left[\langle \bar{s}^{(k)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[\left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& \stackrel{(a)}{\leq} \left(v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{55}$$

361 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with $\beta \rightarrow 1$). Note

362 $a_k = \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$ and

$$\begin{aligned}
a_k \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{56}$$

363 We now give an upper bound of $\mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right]$ using Lemma 7 and plug it into (56):

$$\begin{aligned}
(a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] \\
& + \gamma_{k+1} \left(\frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right]
\end{aligned} \tag{57}$$

364 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^{k+1})} \right\|^2 \right] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{n-1}{n} \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \right) \tag{58}$$

365 where the equality holds as i_k and j_k are drawn independently. For any $\beta > 0$, it holds

$$\begin{aligned}
& \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 - 2\gamma_{k+1}\langle \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2 + \gamma_{k+1}\beta\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right]
\end{aligned} \tag{59}$$

366 where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2\right]
\end{aligned} \tag{60}$$

367 Observe that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)})$. Applying Lemma 7 yields

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\
&\leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\
&\quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\
&\quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right]
\end{aligned} \tag{61}$$

368 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] \tag{62}$$

369 From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} &\leq (1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}))\Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
&\quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right]
\end{aligned} \tag{63}$$

370 Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$,

371 $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$, $\alpha \geq 8$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1} \tag{64}$$

372 which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} =$
 373 $\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} \leq & 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\eta_{i_\ell}^{(\ell)}\|^2] \\ & + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)} \right\|^2 \right] \end{aligned} \quad (65)$$

374 Note $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ Summing on both sides over $k = 0$ to $k = K_{\max} - 1$ yields:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} \\ &= 4 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}[\|\eta_{i_\ell}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\leq \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}[\|\eta_{i_\ell}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \end{aligned} \quad (66)$$

375 We recall (57) where we have summed on both sides from $k = 0$ to $k = K_{\max} - 1$:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)})] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)} \end{aligned} \quad (67)$$

376 Plugging (66) into (67) results in:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \tilde{\alpha}_k \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \tilde{\beta}_k \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)})] \\ &+ \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2] \end{aligned} \quad (68)$$

377 where:

$$\begin{aligned}\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\beta}_k &= \gamma_{k+1} \left(\frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\Gamma}_k &= \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}}\end{aligned}$$

378 and

$$\begin{aligned}a_k &= \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1) + 1}{2n} \right) \\ \Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1} L_s^2}{n^2} (\gamma_{k+1} + \frac{1}{\beta}) \\ c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{L_s, L_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}\end{aligned}$$

379 When, for any $k > 0$, $\tilde{\alpha}_k \geq 0$, we have by Lemma 2 that:

$$\sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (69)$$

380 which yields an upper bound of the gradient of the Lyapunov function V along the path of the
381 iSAEM update and concludes the proof of the Theorem. \square

382 D Proofs of Auxiliary Lemmas

383 D.1 Proof of Lemma 4 and Lemma 5

384 **Lemma.** For any $k \geq 0$ and consider the vrTTSEM update in (10) with $\rho_k = \rho$, it holds for all
385 $k > 0$

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (70)$$

386 where we recall that $\ell(k)$ is the first iteration number in the epoch that iteration k is in.

387 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
388 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \end{aligned} \quad (71)$$

389 We observe, using the identity (71), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] \quad (72)$$

390 For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{\ell(k)}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{\ell(k)}) \right\|^2 \right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \stackrel{(b)}{\leq} L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (73)$$

391 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (72) proves the
392 lemma. \square

393 **Lemma.** For any $k \geq 0$ and consider the fiTTSEM update in (11) with $\rho_k = \rho$, it holds for all $k > 0$
394

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (74)$$

395 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
396 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) \\ &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right] \right) \end{aligned} \quad (75)$$

397 We observe, using the identity (75), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] \quad (76)$$

398 For the latter term, we obtain its upper bound as

$$\begin{aligned}\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{S}}_i^{(k)}) - (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_k^k)})\right\|^2\right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\end{aligned}\quad (77)$$

399 where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_k^k)}\|^2] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_k^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_k^k)}\|^2] \quad (78)$$

400 Substituting into (76) proves the lemma. \square

401 D.2 Proof of Lemma 6

402 **Lemma.** Consider a decreasing stepsize $\gamma_k \in (0, 1)$ and a constant ρ , then the following inequality
403 holds:

$$\mathbb{E}\left[\left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_\ell)^2 (\mathbf{S}^{(\ell)} - \tilde{\mathbf{S}}^{(\ell)}) \quad (79)$$

404 where $\mathbf{S}^{(k)}$ is defined either by (11) (fTTSEM) or (10) (vrTTSEM)

405 **Proof** We begin by writing the two-time-scale update:

$$\begin{aligned}\tilde{\mathbf{S}}^{(k+1)} &= \tilde{\mathbf{S}}^{(k)} + \rho(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k)}) \\ \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)})\end{aligned}\quad (80)$$

406 where $\mathbf{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(t_k^k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_k^k)})$ according to (11). Denote $\delta^{(k+1)} = \hat{\mathbf{s}}^{(k+1)} -$
407 $\tilde{\mathbf{S}}^{(k+1)}$. Then from (80), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1-\rho} (1 - \gamma_{k+1})(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k+1)}) \quad (81)$$

408 Using the telescoping sum and noting that $\delta^{(0)} = 0$, we have

$$\delta^{(k+1)} \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1 - \gamma_{\ell+1})^2 (\mathbf{S}^{(\ell+1)} - \tilde{\mathbf{S}}^{(\ell+1)}) \quad (82)$$

409 \square

410 D.3 Additional Intermediary Result

411 **Lemma 8.** Assume H1. At iteration $k+1$, the drift term of update (11), with $\rho_{k+1} = \rho$, is equivalent
412 to the following :

$$\begin{aligned}\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_k^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_k^k)}] \right] \\ &\quad + (1 - \rho) (\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)})\end{aligned}\quad (83)$$

413 where we recall that $\eta_{i_k}^{(k+1)}$, defined in (19), which is the gap between the MC approximation and
414 the expected statistics.

415 **Proof** Using the fTTSEM update $\tilde{S}^{(k+1)} = (1-\rho)\tilde{S}^{(k)} + \rho\mathcal{S}^{(k+1)}$ where $\mathcal{S}^{(k+1)} = \overline{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$ leads to the following decomposition:

$$\begin{aligned}
& \tilde{S}^{(k+1)} - \hat{s}^{(k)} \\
&= (1-\rho)\tilde{S}^{(k)} + \rho\left(\overline{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right) - \hat{s}^{(k)} + \rho\overline{\mathcal{S}}^{(k)} - \rho\overline{\mathcal{S}}^{(k)} \\
&= \rho(\overline{\mathcal{S}}^{(k)} - \hat{s}^{(k)}) + \rho(\tilde{S}_{i_k}^{(k)} - \overline{\mathcal{S}}_{i_k}^{(k)}) + (1-\rho)\left(\tilde{S}^{(k)} - \hat{s}^{(k)}\right) + \rho\left(\overline{\mathcal{S}}^{(k)} - \overline{\mathcal{S}}^{(k)} + (\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right) \\
&= \rho(\overline{\mathcal{S}}^{(k)} - \hat{s}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho\left[(\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}]\right] \\
&+ (1-\rho)\left(\tilde{S}^{(k)} - \hat{s}^{(k)}\right)
\end{aligned}$$

417 where we observe that $\mathbb{E}[\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] = \overline{\mathcal{S}}^{(k)} - \overline{\mathcal{S}}^{(k)}$ and which concludes the proof.

418 *Important Note:* Note that $\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}$ is not equal to $\eta_{i_k}^{(k+1)}$, defined in (19), which is the gap
419 between the MC approximation and the expected statistics. Indeed $\tilde{S}_{i_k}^{(t_{i_k}^k)}$ is not computed under the
420 same model as $\overline{\mathcal{S}}_{i_k}^{(k)}$. □

421 E Proof of Theorem 2

422 **Theorem.** Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and
 423 consider the vrTTSEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$.

424 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$, $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$
 425 and a constant $\mu \in (0, 1)$, we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (84)$$

426 **Proof** Using the smoothness of V and update (10), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (85)$$

427 Denote $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fiTTSEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.
 428 Taking expectations on both sides show that

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}\rho \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\langle \eta_{i_k}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \gamma_{k+1}(1-\rho) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \quad (86)$$

429 where we have used (71) in (a) and $\mathbb{E}[\mathbf{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$ in (b), the growth condition in
 430 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

431 Furthermore, for $k+1 \leq \ell(k) + m$ (i.e., $k+1$ is in the same epoch as k), we have

$$\begin{aligned} &\mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 \\ &\quad - 2\gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1-\rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle] \\ &\leq \mathbb{E}[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2], \end{aligned} \quad (87)$$

432 where we first used (71) and the last inequality is due to the Young's inequality.

433 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \quad (88)$$

434 where $b_k := \bar{b}_{k \bmod m}$ is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0. \quad (89)$$

435 Note that \bar{b}_i is decreasing with i and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2}, \quad i = 1, 2, \dots, m. \quad (90)$$

436 For $k+1 \leq \ell(k) + m$, we have the following inequality

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[\rho \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\frac{\gamma_{k+1}\rho}{\beta} \left\|\eta_{i_k}^{(k+1)}\right\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \end{aligned} \quad (91)$$

437 And using Lemma 4 we obtain:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2 - \gamma_{k+1}^2\rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \|\mathbf{h}_k\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[(\rho + \rho^2\gamma_{k+1} L_V) \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho - (1-\rho)^2\gamma_{k+1} L_V) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \left\|\eta_{i_k}^{(k+1)}\right\|^2 + \left(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2\right) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \end{aligned} \quad (92)$$

438 Rearranging the terms yields:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\quad + \underbrace{\left(b_{k+1}(1 + \gamma\beta + 2\gamma^2\rho^2 L_{\mathbf{s}}^2) + \gamma^2\rho^2 L_V L_{\mathbf{s}}^2\right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)} \end{aligned} \quad (93)$$

439 where

$$\begin{aligned} \tilde{\eta}^{(k+1)} &= \left(\gamma_{k+1}(\rho + \rho^2\gamma_{k+1} L_V) + b_{k+1}(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2)\right) \mathbb{E}\left[\left\|\eta_{i_k}^{(k+1)}\right\|^2\right] \\ \chi^{(k+1)} &= \left(b_{k+1}(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2) - \gamma_{k+1}(1-\rho - (1-\rho)^2\gamma_{k+1} L_V)\right) \\ \tilde{\chi}^{(k+1)} &= \chi^{(k+1)} \mathbb{E}\left[\left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \end{aligned} \quad (94)$$

440 This leads, using Lemma 2, that for any γ_{k+1} , ρ and β such that $\rho v_{\min} + v_{\max}^2 -$
 441 $\gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2) > 0$,

$$\begin{aligned} v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \\ &\quad + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \end{aligned} \quad (95)$$

442 We first remark that

$$\begin{aligned} & \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \\ & \geq \frac{\gamma_{k+1}\rho}{c_1}(1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1)) \end{aligned} \quad (96)$$

443 where $c_1 = v_{\min}^{-1}$. By setting $\bar{L} = \max\{L_s, L_V\}$, $\beta = \frac{c_1\bar{L}}{n^{1/3}}$, $\rho = \frac{\mu}{c_1\bar{L}n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$ and
 444 $\{\gamma_{k+1}\}$ any sequence of decreasing stepsizes in $(0, 1)$, it can be shown that there exists $\mu \in (0, 1)$,
 445 such that the following lower bound holds

$$\begin{aligned} & 1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1) \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V\mu^2}{c_1^2 n^{\frac{4}{3}}} \frac{(1 + \gamma\beta + 2\gamma^2 L_s^2)^m - 1}{\gamma\beta + 2\gamma^2 L_s^2} (\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2} (e - 1)(1 + \frac{2\mu}{n}) \geq 1 - \mu - \mu(1 + 2\mu) \frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2} \end{aligned} \quad (97)$$

446 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2 L_s^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2 n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \text{ and } (1 + \gamma\beta + 2\gamma^2 L_s^2)^m \leq e - 1. \quad (98)$$

447 and the required μ in (b) can be found by solving the quadratic equation.

448 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_{\max}})}{v_{\min}\rho} + 2 \sum_{k=0}^{K_{\max}-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min}\rho} \quad (99)$$

449 Note that $R_0 = \mathbb{E}[V(\hat{s}^{(0)})]$ and if K_{\max} is a multiple of m , then $R_{\max} = \mathbb{E}[V(\hat{s}^{(K_{\max})})]$. Under the
 450 latter condition, we have

$$\sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}] \quad (100)$$

451 This concludes our proof.

452 □

453 **F Proof of Theorem 3**

454 **Theorem.** Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and
 455 consider the fiTTSEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$.

456 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$,
 457 $\beta = \frac{c_1 \bar{L}}{n}$, $\rho = \frac{1}{n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$, we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (101)$$

458 **Proof** Using the smoothness of V and update (11), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (102)$$

459 Denote $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fiTTSEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.

460 Using Lemma 8 and the additional following identity:

$$\mathbb{E} \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] = 0 \quad (103)$$

461 we have:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \mathbb{E} \left[\langle \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1 - \rho) \mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(a)}{\leq} -v_{\min} \gamma_{k+1} \rho \mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1} \mathbb{E} \left[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(b)}{\leq} -(v_{\min} \gamma_{k+1} \rho + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \end{aligned} \quad (104)$$

462 where $\xi^{(k+1)} = \mathbb{E} \left[\left\| \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] \right\|^2 \right]$.

463 **Bounding** $\mathbb{E}[\|\mathbf{H}_{k+1}\|^2]$ Using Lemma 5, we obtain:

$$\begin{aligned} &\gamma_{k+1} (v_{\min} \rho + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)}) \right] + \tilde{\xi}^{(k+1)} + \left((1 - \rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V \rho^2 L_{\mathbf{s}}^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (105)$$

464 where $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 \mathbb{L}_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1}\rho^2}{2} \xi^{(k+1)}$. Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (106)$$

465 where the equality holds as i_k and j_k are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \end{aligned} \quad (107)$$

466 Note that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}\mathbf{H}_{k+1}$ and that in expectation we recall
467 that $\mathbb{E}[\mathbf{H}_{k+1}|\mathcal{F}_k] = \rho\mathbf{h}_k + \rho\mathbb{E}[\eta_{i_k}^{(k+1)}|\mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}]$ where $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$. Thus,
468 for any $\beta > 0$, it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (108)$$

469 where the last inequality is due to the Young's inequality. Plugging this into (106) yields:

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (109)$$

470 Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2] \\ &\quad + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (110)$$

471 We now use Lemma 5 on $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2 \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2$ and obtain:

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\
& \leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n} + \frac{(n-1)(1+\gamma_{k+1}\beta)}{n^2}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\
& \leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{111}$$

472 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \tag{112}$$

473 From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} & \leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{113}$$

474 Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$,
475 $\rho = \frac{1}{n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{4/3}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1} \tag{114}$$

476 which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} = \frac{1}{n} -$
477 $\gamma_{k+1}\beta - \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned}
\Delta^{(k+1)} & \leq \sum_{\ell=0}^k \omega_{k,\ell} \left(2\gamma_{\ell+1}^2 \rho^2 + \frac{\gamma_{\ell+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] \\
& + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left(2\gamma_{\ell+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\tilde{\mathbf{S}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)}
\end{aligned} \tag{115}$$

478 where $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ and $\tilde{\epsilon}^{(\ell+1)} = \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]$.

479 Summing on both sides over $k = 0$ to $k = K_{\max} - 1$ yields:

$$\begin{aligned}
\sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} & \leq \sum_{k=0}^{K_{\max}-1} \frac{2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
& + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}
\end{aligned} \tag{116}$$

480 We recall (105) where we have summed on both sides from $k = 0$ to $k = K_{\max} - 1$:

$$\begin{aligned}
& \mathbb{E}[V(\hat{\mathbf{s}}^{(K_{\max})}) - V(\hat{\mathbf{s}}^{(0)})] \\
& \leq \sum_{k=0}^{K_{\max}-1} \left\{ \gamma_{k+1}(-(\nu_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] + \gamma^2 L_V \rho^2 L_{\mathbf{s}}^2 \Delta^{(k)} \right\} \\
& + \sum_{k=0}^{K_{\max}-1} \left\{ \tilde{\xi}^{(k+1)} + \left((1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right\} \\
& \leq \sum_{k=0}^{K_{\max}-1} \left\{ -\gamma_{k+1}(\nu_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right\} \mathbb{E}[\|\mathbf{h}_k\|^2] \\
& + \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned} \tag{117}$$

where

$$\Xi^{(k+1)} = \tilde{\xi}^{(k+1)} + \frac{\gamma_{k+1}^3 L_V \rho^2 L_{\mathbf{s}}^2}{\Lambda_{(k+1)}} \epsilon^{(k+1)}$$

and

$$\Gamma_{k+1} = \left((1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_{\mathbf{s}}^2 (1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta} \right)}{\Lambda_{(k+1)}}$$

481 We now analyse the following quantity

$$\begin{aligned}
& -\gamma_{k+1}(\nu_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\
& = -\gamma_{k+1} \left[(\nu_{\min}\rho + v_{\max}^2) + \gamma_{k+1} \rho^2 L_V + \frac{\rho^2 \gamma_{k+1} L_V L_{\mathbf{s}}^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right]
\end{aligned} \tag{118}$$

482 Furthermore, we recall that $c_1 = \nu_{\min}^{-1}$, $\alpha = \max\{2, 1+2\nu_{\min}\}$, $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$,

483 $\beta = \frac{c_1 \bar{L}}{n}$, $\rho = \frac{1}{n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$. Then,

$$\begin{aligned}
& \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\
& \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} (k^2 \alpha^2 c_1^2 n^{4/3})^{-1} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1^2 \bar{L}^2 n^{1/3}} \right)}{\frac{1}{n} - \frac{1}{k \alpha n} - \frac{1}{k^2 \alpha^2 c_1^2 n^{4/3}}} \\
& = \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1^2 \bar{L}^2 n^{1/3}} \right)}{(k \alpha c_1 n^{1/3})(k \alpha - 1) c_1 - 1} \\
& \stackrel{(a)}{\leq} \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k \alpha c_1^2 \bar{L} n^{1/3}} \left(\frac{2}{k \alpha n} + 1 \right)}{2(\alpha c_1 n^{1/3}) - 1} \\
& \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{1}{k \alpha^2 c_1^3 \bar{L} n^{2/3}} \\
& \leq \frac{1}{\alpha c_1 \bar{L} n^{2/3}}
\end{aligned} \tag{119}$$

484 where (a) is due to $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ and $k \alpha c_1 n^{1/3} \geq 1$. Also, since $-\gamma_{k+1}(\nu_{\min}\rho +$
485 $v_{\max}^2) \leq -\gamma_{k+1}\rho \nu_{\min} = -$

486 Using the Lemma 2, we know that $v_{\max}^2 \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \leq \|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2$ and using (119) on (117)
 487 yields:

$$\begin{aligned} v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned} \quad (120)$$

488 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned} \quad (121)$$

489 **Bounding** $\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right]$ Remark that this term is the price we pay for the two time scale
 490 dynamics and corresponds to the gap between the two asynchronous updates (one is on $\hat{\mathbf{s}}^{(k)}$ and the
 491 other on $\tilde{S}^{(k)}$).

492 **FIND AN UPPER BOUND TO THAT GAP**

493

□

G Practical Implementations of Two-Time-Scale EM Methods

G.1 Gaussian mixture models

G.1.1 Model assumptions

We first recognize that the constraint set for θ is given by

$$\Theta = \Delta^M \times \mathbb{R}^M. \quad (122)$$

Using the partition of the sufficient statistics as $S(y_i, z_i) = (S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$, the partition $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ and the fact that $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$, the complete data log-likelihood can be expressed as in (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i) y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (123)$$

and $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$. We also define for each $m \in \llbracket 1, M \rrbracket$, $j \in \llbracket 1, 3 \rrbracket$, $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$. Consider the following latent sample used to compute an approximation of the conditional expected value $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}} | y = y_i]$:

$$z_{i,m} \sim \mathbb{P}(z_i = m | y_i; \theta) \quad (124)$$

where $m \in \llbracket 1, M \rrbracket$, $i \in \llbracket 1, n \rrbracket$ and $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$.

In particular, given iteration $k + 1$, the computation of the approximated quantity $\tilde{S}_{i_k}^{(k)}$ during Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left(\underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:=\tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1})y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})y_{i_k}}_{:=\tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:=\tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (125)$$

Recall that we have used the following regularizer:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (126)$$

It can be shown that the regularized M-step in (4) evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\mu}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (127)$$

where we have defined for all $m \in \llbracket 1, M \rrbracket$ and $j \in \llbracket 1, 3 \rrbracket$, $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$.

G.1.2 Algorithms updates

In the sequel, recall that, for all $i \in \llbracket n \rrbracket$ and iteration k , the computed statistic $\tilde{S}_{i_k}^{(k)}$ is defined by (125). At iteration k , the several E-steps defined by (9) or (10) and (11) leads to the definition of the quantity $\hat{\mathbf{s}}^{(k+1)}$. For the GMM example, after the initialization of the quantity $\hat{\mathbf{s}}^{(0)} = n^{-1} \sum_{i=1}^n \tilde{\mathbf{s}}_i^{(0)}$, those E-steps break down as follows:

Batch EM (EM): for all $i \in \llbracket 1, n \rrbracket$, compute $\tilde{\mathbf{s}}_i^{(k)}$ and set

$$\hat{\mathbf{s}}^{(k+1)} = n^{-1} \sum_{i=1}^n \tilde{\mathbf{s}}_i^{(k)}. \quad (128)$$

518 where $\bar{s}_i^{(k)}$ are computed using the exact conditional expected value $\mathbb{E}_{\boldsymbol{\theta}}[\mathbb{1}_{\{z_i=m\}}|y=y_i]$:

$$\tilde{\omega}_m(y_i; \boldsymbol{\theta}) := \mathbb{E}_{\boldsymbol{\theta}}[\mathbb{1}_{\{z_i=m\}}|y=y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)}, \quad (129)$$

519 **Incremental EM (iEM):** draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}. \quad (130)$$

520 **batch SAEM (SAEM):** draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}. \quad (131)$$

521 where $= \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$ with $\tilde{S}_i^{(k)}$ defined in (125).

522 **Incremental SAEM (iSAEM):** draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \left(\tilde{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_i^k)}) \right). \quad (132)$$

524 **Variance Reduced Two-Time-Scale EM (vrTTSEM):** draw an index i_k uniformly at random on
525 $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}))). \quad (133)$$

526 **Fast Incremental Two-Time-Scale EM (fiTTSEM):** draw an index i_k uniformly at random on $\llbracket n \rrbracket$,
527 compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}))). \quad (134)$$

528 Finally, the k -th update reads $\hat{\boldsymbol{\theta}}^{(k+1)} = \bar{\boldsymbol{\theta}}(\hat{s}^{(k+1)})$ where the function $s \rightarrow \bar{\boldsymbol{\theta}}(s)$ is defined by (127).