

434 A Proof of Auxiliary Lemmas

435 **Lemma 1.** *For the sequence defined in (9), we have*

$$Z_{t+1} - Z_t = \alpha \frac{\beta_1}{1 - \beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}}. \quad (6)$$

436 **Proof:** By update rule of Algorithm 2, we first have

$$\begin{aligned} \bar{X}_{t+1} &= \frac{1}{N} \sum_{i=1}^N x_{t+1,i} \\ &= \frac{1}{N} \sum_{i=1}^N \left(x_{t+0.5,i} - \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) \\ &= \frac{1}{N} \sum_{i=1}^N \left(\sum_{j=1}^N W_{ij} x_{t,j} - \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) \\ &\stackrel{(i)}{=} \left(\frac{1}{N} \sum_{j=1}^N x_{t,j} \right) - \frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \\ &= \bar{X}_t - \frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}}, \end{aligned}$$

437 where (i) is due to an interchange of summation and $\sum_{i=1}^N W_{ij} = 1$. Then, we have

$$\begin{aligned} Z_{t+1} - Z_t &= \bar{X}_{t+1} - \bar{X}_t + \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) - \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) \\ &= \frac{1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) - \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) \\ &= \frac{1}{1 - \beta_1} \left(-\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) - \frac{\beta_1}{1 - \beta_1} \left(-\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right) \\ &= \frac{1}{1 - \beta_1} \left(-\frac{1}{N} \sum_{i=1}^N \alpha \frac{\beta_1 m_{t-1,i} + (1 - \beta_1) g_{t,i}}{\sqrt{u_{t,i}}} \right) - \frac{\beta_1}{1 - \beta_1} \left(-\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right) \\ &= \alpha \frac{\beta_1}{1 - \beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}}, \end{aligned}$$

438 which is the desired result. \square

439 **Lemma 2.** *Given a set of numbers a_1, \dots, a_n and denote their mean to be $\bar{a} = \frac{1}{n} \sum_{i=1}^n a_i$. Define*
 440 $b_i(r) \triangleq \max(a_i, r)$ *and $\bar{b}(r) = \frac{1}{n} \sum_{i=1}^n b_i(r)$. For any r and r' with $r' \geq r$ we have*

$$\sum_{i=1}^n |b_i(r) - \bar{b}(r)| \geq \sum_{i=1}^n |b_i(r') - \bar{b}(r')| \quad (7)$$

441 and when $r \leq \min_{i \in [n]} a_i$, we have

$$\sum_{i=1}^n |b_i(r) - \bar{b}(r)| = \sum_{i=1}^n |a_i - \bar{a}|. \quad (8)$$

442 **Proof:** Without loss of generality, assume $a_i \leq a_j$ when $i < j$, i.e. a_i is a non-decreasing sequence.
 443 Define

$$h(r) = \sum_{i=1}^n |b_i(r) - \bar{b}(r)| = \sum_{i=1}^n \left| \max(a_i, r) - \frac{1}{n} \sum_{j=1}^n \max(a_j, r) \right|.$$

444 We need to prove that h is a non-increasing function of r . First, it is easy to see that h is a continuous
 445 function of r with non-differentiable points $r = a_i, i \in [n]$, thus h is a piece-wise linear function.

446 Next, we will prove that $h(r)$ is non-increasing in each piece. Define $l(r)$ to be the largest index
 447 with $a(l(r)) < r$, and $s(r)$ to be the largest index with $a_{s(r)} < \bar{b}(r)$. Note that we have for $i \leq l(r)$,
 448 $b_i(r) = r$ and for $i \leq s(r)$ $b_i(r) - \bar{b}(r) \leq 0$ since a_i is a non-decreasing sequence. Therefore, we
 449 have

$$h(r) = \sum_{i=1}^{l(r)} (\bar{b}(r) - r) + \sum_{i=l(r)+1}^{s(r)} (\bar{b}(r) - a_i) + \sum_{i=s(r)+1}^n (a_i - \bar{b}(r))$$

450 and

$$\bar{b}(r) = \frac{1}{n} \left(l(r)r + \sum_{i=l(r)+1}^n a_i \right).$$

451 Taking derivative of the above form, we know the derivative of $h(r)$ at differentiable points is

$$\begin{aligned} h'(r) &= l(r) \left(\frac{l(r)}{n} - 1 \right) + (s(r) - l(r)) \frac{l(r)}{n} - (n - s(r)) \frac{l(r)}{n} \\ &= \frac{l(r)}{n} ((l(r) - n) + (s(r) - l(r)) - (n - s(r))). \end{aligned}$$

452 Since we have $s(r) \leq n$ we know $(l(r) - n) + (s(r) - l(r)) - (n - s(r)) \leq 0$ and thus

$$h'(r) \leq 0,$$

453 which means $h(r)$ is non-increasing in each piece. Combining with the fact that $h(r)$ is continuous,
 454 (7) is proven. When $r \leq a(i)$, we have $b(i) = \max(a_i, r) = r$, for all $r \in [n]$ and $\bar{b}(r) =$
 455 $\frac{1}{n} \sum_{i=1}^n a_i = \bar{a}$ which proves (8). \square

456 B Proof of Theorem 2

457 To prove convergence of the algorithm, we first define an auxiliary sequence

$$Z_t = \bar{X}_t + \frac{\beta_1}{1 - \beta_1} (\bar{X}_t - \bar{X}_{t-1}), \quad (9)$$

458 with $\bar{X}_0 \triangleq \bar{X}_1$. Since $\mathbb{E}[g_{t,i}] = \nabla f(x_{t,i})$ and $u_{t,i}$ is a function of $G_{1:t-1}$ (which denotes
 459 G_1, G_2, \dots, G_{t-1}), we have

$$\mathbb{E}_{G_t | G_{1:t-1}} \left[\frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right] = \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}}.$$

460 Assuming smoothness (A1) we have

$$f(Z_{t+1}) \leq f(Z_t) + \langle \nabla f(Z_t), Z_{t+1} - Z_t \rangle + \frac{L}{2} \|Z_{t+1} - Z_t\|^2.$$

461 Using Lemma 1 into the above inequality and take expectation over G_t given $G_{1:t-1}$, we have

$$\begin{aligned} & \mathbb{E}_{G_t | G_{1:t-1}} [f(Z_{t+1})] \\ & \leq f(Z_t) - \alpha \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle + \frac{L}{2} \mathbb{E}_{G_t | G_{1:t-1}} [\|Z_{t+1} - Z_t\|^2] \\ & \quad + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E}_{G_t | G_{1:t-1}} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]. \end{aligned}$$

462 Then take expectation over $G_{1:t-1}$ and rearrange, we have

$$\alpha \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle \right] \quad (10)$$

$$\leq \mathbb{E}[f(Z_t)] - \mathbb{E}[f(Z_{t+1})] + \frac{L}{2} \mathbb{E}[\|Z_{t+1} - Z_t\|^2] \\ + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]. \quad (11)$$

463 In addition, we have

$$\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle \\ = \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\rangle + \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\rangle \quad (12)$$

464 and the first term on RHS of the equality can be lower bounded as

$$\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\rangle \\ = \frac{1}{2} \left\| \frac{\nabla f(Z_t)}{\bar{U}_t^{1/4}} \right\|^2 + \frac{1}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\nabla f(Z_t) - \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 \\ \geq \frac{1}{4} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \frac{1}{4} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\nabla f(Z_t) - \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 \\ - \frac{1}{2} \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \\ \geq \frac{1}{2} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{3}{2} \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{3}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2, \quad (13)$$

465 where the inequalities are all due to Cauchy-Schwartz. Substituting (13) and (12) into (10), we get

$$\frac{1}{2} \alpha \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \leq \mathbb{E}[f(Z_t)] - \mathbb{E}[f(Z_{t+1})] + \frac{L}{2} \mathbb{E}[\|Z_{t+1} - Z_t\|^2] \\ + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\ - \alpha \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\rangle \right] \\ + \frac{3}{2} \alpha \mathbb{E} \left[\left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right].$$

466 Then sum over the above inequality from $t = 1$ to T and divide both sides by $T\alpha/2$, we have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
& \leq \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + \frac{L}{T\alpha} \sum_{t=1}^T \mathbb{E} [\|Z_{t+1} - Z_t\|^2] \\
& \quad + \underbrace{\frac{2}{T} \frac{\beta_1}{1-\beta_1} \sum_{t=1}^T \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]}_{D_1} \\
& \quad + \underbrace{\frac{2}{T} \sum_{t=1}^T \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{\bar{U}_t}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]}_{D_2} \\
& \quad + \underbrace{\frac{3}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right]}_{D_3}. \tag{14}
\end{aligned}$$

467 Now we need to upper bound all the terms on RHS of the above inequality to get the convergence
 468 rate. For the terms composing D_3 in (14), we can upper bound them by

$$\begin{aligned}
\left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 & \leq \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \|\nabla f(Z_t) - \nabla f(\bar{X}_t)\|^2 \\
& \leq L \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \underbrace{\|Z_t - \bar{X}_t\|^2}_{D_4} \tag{15}
\end{aligned}$$

469 and

$$\begin{aligned}
\left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 & \leq \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \frac{1}{N} \sum_{i=1}^N \|\nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)\|^2 \\
& \leq L \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \frac{1}{N} \underbrace{\sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2}_{D_5}, \tag{16}
\end{aligned}$$

470 using Jensen's inequality, Lipschitz continuity of f_i , and the fact that $f = \frac{1}{N} \sum_{i=1}^N f_i$. Next we need
 471 to bound D_4 and D_5 . Recall the update rule of X_t , we have

$$X_t = X_{t-1}W - \alpha \frac{M_{t-1}}{\sqrt{U_{t-1}}} = X_1 W^{t-1} - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} W^k, \tag{17}$$

472 where we define $W^0 = \mathbf{I}$. Since W is a symmetric matrix, we can decompose it as $W = Q\Lambda Q^T$
 473 where Q is a orthonormal matrix and Λ is a diagonal matrix whose diagonal elements correspond
 474 to eigenvalues of W in an descending order, i.e. $\Lambda_{ii} = \lambda_i$ with λ_i being i th largest eigenvalue of
 475 W . In addition, because W is a doubly stochastic matrix, we know $\lambda_1 = 1$ and $q_1 = \frac{1}{\sqrt{N}}$. With
 476 eigen-decomposition of W , we can rewrite D_5 as

$$\sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 = \|X_t - \bar{X}_t \mathbf{1}_N^T\|_F^2 = \|X_t Q Q^T - X_t \frac{1}{N} \mathbf{1}_N \mathbf{1}_N^T\|_F^2 = \sum_{l=2}^N \|X_t q_l\|^2. \tag{18}$$

477 In addition, we can rewrite (17) as

$$X_t = X_1 W^{t-1} - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} W^k = X_1 - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} Q \Lambda^k Q^T, \quad (19)$$

478 where the last equality is because $x_{1,i} = x_{1,j}$, for all i, j and thus $X_1 W = X_1$. Then we have when
479 $l > 1$,

$$X_t q_l = (X_1 - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} Q \Lambda^k Q^T) q_l = -\alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} q_l \lambda_l^k, \quad (20)$$

480 since Q is orthonormal and $X_1 q_l = x_{1,1} \mathbf{1}_N^T q_l = x_{1,1} \sqrt{N} q_1^T q_l = 0$, for all $l \neq 1$.

481 Combining (18) and (20), we have

$$\begin{aligned} D_5 &= \sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 = \sum_{l=2}^N \|X_t q_l\|^2 \\ &= \sum_{l=2}^N \alpha^2 \left\| \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} \lambda_l^k q_l \right\|^2 \\ &\leq \alpha^2 \left(\frac{1}{1-\lambda} \right)^2 N d G_\infty^2 \frac{1}{\epsilon}, \end{aligned} \quad (21)$$

482 where the last inequality follows from the fact that $g_{t,i} \leq G_\infty$, $\|q_l\| = 1$, and $|\lambda_l| \leq \lambda < 1$. Now let
483 us turn to D_4 , it can be rewritten as

$$\begin{aligned} \|Z_t - \bar{X}_t\|^2 &= \left\| \frac{\beta_1}{1-\beta_1} (\bar{X}_t - \bar{X}_{t-1}) \right\|^2 = \left(\frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right\|^2 \\ &\leq \left(\frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 d \frac{G_\infty^2}{\epsilon}. \end{aligned} \quad (22)$$

484 Now we know both D_4 and D_5 are in the order of $\mathcal{O}(\alpha^2)$ and thus D_3 is in the order of
485 $\mathcal{O}(\alpha^2)$. Next we will bound D_2 and D_1 . Define $G_1 \triangleq \max_{t \in [T]} \max_{i \in [N]} \|\nabla f_i(x_{t,i})\|_\infty$,
486 $G_2 \triangleq \max_{t \in [T]} \|\nabla f(Z_t)\|_\infty$, $G_3 \triangleq \max_{t \in [T]} \max_{i \in [N]} \|g_{t,i}\|_\infty$ and $G_\infty = \max(G_1, G_2, G_3)$.
487 Then we have

$$\begin{aligned} D_2 &= \sum_{t=1}^T \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{\bar{U}_t}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\ &\leq \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[\bar{U}_t]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \right] \\ &= \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[\bar{U}_t]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \frac{\sqrt{[\bar{U}_t]_j} + \sqrt{[u_{t,i}]_j}}{\sqrt{[\bar{U}_t]_j} + \sqrt{[u_{t,i}]_j}} \right] \\ &= \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{[\bar{U}_t]_j - [u_{t,i}]_j}{[\bar{U}_t]_j \sqrt{[u_{t,i}]_j} + \sqrt{[\bar{U}_t]_j} [u_{t,i}]_j} \right| \right] \\ &\leq \underbrace{\mathbb{E} \left[\sum_{t=1}^T G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{[\bar{U}_t]_j - [u_{t,i}]_j}{2\epsilon^{1.5}} \right| \right]}_{D_6}, \end{aligned} \quad (23)$$

where the last inequality is due to $[u_{t,i}]_j \geq \epsilon$, for all t, i, j . To simplify notations, define $\|A\|_{abs} = \sum_{i,j} |A_{ij}|$ to be the entry-wise L_1 norm of a matrix A , then we obtain

$$\begin{aligned} D_6 &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\bar{U}_t \mathbf{1}^T - U_t\|_{abs} \leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\tilde{U}_t \mathbf{1}^T - \tilde{U}_t\|_{abs} \\ &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\tilde{U}_t \frac{1}{N} \mathbf{1}_N \mathbf{1}_N^T - \tilde{U}_t Q Q^T\|_{abs} \\ &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs}, \end{aligned}$$

where the second inequality is due to Lemma 2, introduced Section A, and the fact that $U_t = \max(\tilde{U}_t, \epsilon)$ (element-wise max operator). Recall from update rule of U_t , by defining $\hat{V}_{-1} \triangleq \hat{V}_0$ and $U_0 \triangleq U_{1/2}$, we have for all $t \geq 0$, $\tilde{U}_{t+1} = (\tilde{U}_t - \hat{V}_{t-1} + \hat{V}_t)W$. Thus, we obtain

$$\tilde{U}_t = \tilde{U}_0 W^t + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) W^k = \tilde{U}_0 + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) Q \Lambda^k Q^T.$$

Then we further obtain when $l \neq 1$,

$$\tilde{U}_t q_l = (\tilde{U}_0 + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) Q \Lambda^k Q^T) q_l = \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) q_l \lambda_l^k,$$

where the last equality is due to the definition $\tilde{U}_0 \triangleq U_{1/2} = \epsilon \mathbf{1}_d \mathbf{1}_N^T = \sqrt{N} \epsilon \mathbf{1}_d \mathbf{1}_N^T$ (recall that $q_1 = \frac{1}{\sqrt{N}} \mathbf{1}_N^T$) and $q_i^T q_j = 0$ when $i \neq j$. Note that by definition of $\|\cdot\|_{abs}$, we have for all A, B , $\|A + B\|_{abs} \leq \|A\|_{abs} + \|B\|_{abs}$, then

$$\begin{aligned} D_6 &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs} \\ &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_{abs} \\ &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \left\| \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_1 \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \\ &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \sqrt{N} \left\| \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_2 \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \\ &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \sqrt{N} \lambda^k \\ &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})\|_{abs} \sqrt{N} \lambda^k \\ &= \frac{G_\infty^2}{N} \frac{1}{2\epsilon^{1.5}} \sum_{o=0}^{T-1} \sum_{t=o+1}^T \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \sqrt{N} \lambda^{t-o} \\ &\leq \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs}, \end{aligned} \tag{24}$$

where $\lambda = \max(|\lambda_2|, |\lambda_N|)$. Combining (23) and (24), we have

$$D_2 \leq \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \frac{\lambda}{1-\lambda} \mathbb{E} \left[\sum_{o=0}^{T-1} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right].$$

498 Now we need to bound D_1 , we have

$$\begin{aligned}
D_1 &= \sum_{t=1}^T \mathbb{E} \left[\left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\
&\leq \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \right] \\
&= \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \left(\frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right) \frac{\sqrt{[u_{t,i}]_j} + \sqrt{[u_{t-1,i}]_j}}{\sqrt{[u_{t,i}]_j} + \sqrt{[u_{t-1,i}]_j}} \right| \right] \\
&\leq \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{2\epsilon^{1.5}} ([u_{t-1,i}]_j - [u_{t,i}]_j) \right| \right] \\
&\stackrel{(a)}{\leq} \sum_{t=1}^T \mathbb{E} \left[G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{2\epsilon^{1.5}} |([\tilde{u}_{t-1,i}]_j - [\tilde{u}_{t,i}]_j)| \right] \\
&= G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{N} \mathbb{E} \left[\sum_{t=1}^T \|\tilde{U}_{t-1} - \tilde{U}_t\|_{abs} \right],
\end{aligned} \tag{25}$$

499 where (a) is due to $[\tilde{u}_{t-1,i}]_j = \max([u_{t-1,i}]_j, \epsilon)$ and the function $\max(\cdot, \epsilon)$ is 1-Lipschitz. In
500 addition, by update rule of U_t , we have

$$\begin{aligned}
&\sum_{t=1}^T \|\tilde{U}_{t-1} - \tilde{U}_t\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1} - (\tilde{U}_{t-1} - \hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1}(QQ^T - Q\Lambda Q^T) + (-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1}(\sum_{l=2}^N q_l(1 - \lambda_l)q_l^T) + (-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&\leq \sum_{t=1}^T \left\| \sum_{k=1}^{t-1} (-\hat{V}_{t-2-k} + \hat{V}_{t-1-k}) \sum_{l=2}^N q_l \lambda_l^k (1 - \lambda_l) q_l^T \right\|_{abs} + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&\leq \sum_{t=1}^T \left(\sum_{k=1}^{t-1} \|-\hat{V}_{t-2-k} + \hat{V}_{t-1-k}\|_{abs} \sqrt{N} \lambda^k \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&= \sum_{t=1}^T \left(\sum_{o=1}^{t-1} \|-\hat{V}_{o-2} + \hat{V}_{o-1}\|_{abs} \sqrt{N} \lambda^{t-o} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&= \sum_{o=1}^{T-1} \sum_{t=o+1}^T \left(\|-\hat{V}_{o-2} + \hat{V}_{o-1}\|_{abs} \sqrt{N} \lambda^{t-o} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&\leq \sum_{o=1}^{T-1} \frac{\lambda}{1 - \lambda} \left(\|-\hat{V}_{o-2} + \hat{V}_{o-1}\|_{abs} \sqrt{N} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&\leq \frac{1}{1 - \lambda} \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \sqrt{N}.
\end{aligned} \tag{26}$$

501 Combining (25) and (26), we have

$$D_1 \leq G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{N} \mathbb{E} \left[\frac{1}{1-\lambda} \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \sqrt{N} \right]. \quad (27)$$

502 What remains is to bound $\sum_{t=1}^T \mathbb{E} [\|Z_{t+1} - Z_t\|^2]$. By update rule of Z_t , we have

$$\begin{aligned} & \|Z_{t+1} - Z_t\|^2 \\ &= \left\| \alpha \frac{\beta_1}{1-\beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\ &\leq 2\alpha^2 \left\| \frac{\beta_1}{1-\beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left(\frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\|^2 + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\ &\leq 2\alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{\sqrt{\epsilon}} \left| \frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\ &\leq 2\alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{\sqrt{\epsilon}} \left| \frac{[u_{t,i}]_j - [u_{t-1,i}]_j}{2\epsilon^{1.5}} \right| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\ &\leq 2\alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{2\epsilon^2} |\tilde{u}_{t,i}]_j - [\tilde{u}_{t-1,i}]_j| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\ &= 2\alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \frac{1}{2\epsilon^2} \|\tilde{U}_t - \tilde{U}_{t-1}\|_{abs} + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2, \end{aligned} \quad (28)$$

503 where the last inequality is again due to the definition that $[\tilde{u}_{t,i}]_j = \max([u_{t,i}]_j, \epsilon)$ and the fact that
504 $\max(\cdot, \epsilon)$ is 1-Lipschitz. Then, we have

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E} [\|Z_{t+1} - Z_t\|^2] \\ &\leq 2\alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \frac{1}{2\epsilon^2} \mathbb{E} \left[\sum_{t=1}^T \|\tilde{U}_t - \tilde{U}_{t-1}\|_{abs} \right] + 2\alpha^2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\ &\leq \alpha^2 \left(\frac{\beta_1}{1-\beta_1} \right)^2 \frac{G_\infty^2}{\sqrt{N}} \frac{1}{\epsilon^2} \frac{1}{1-\lambda} \mathbb{E} \left[\sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] + 2\alpha^2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right], \end{aligned}$$

505 where the last inequality is due to (26).

506 We now bound the last term on RHS of the above inequality. A trivial bound can be

$$\sum_{t=1}^T \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \leq \sum_{t=1}^T d G_\infty^2 \frac{1}{\epsilon},$$

507 due to $\|g_{t,i}\| \leq G_\infty$ and $[u_{t,i}]_j \geq \epsilon$, for all j (verified from update rule of $u_{t,i}$ and the assumption
508 that $[v_{t,i}]_j \geq \epsilon$, for all i). However, the above bound is independent of N , to get a better bound, we

509 need a more involved analysis to show its dependency on N . To do this, we first notice that

$$\begin{aligned}
& \mathbb{E}_{G_t|G_{1:t-1}} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\
&= \mathbb{E}_{G_t|G_{1:t-1}} \left[\frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left\langle \frac{\nabla f_i(x_{t,i}) + \xi_{t,i}}{\sqrt{u_{t,i}}}, \frac{\nabla f_j(x_{t,j}) + \xi_{t,j}}{\sqrt{u_{t,j}}} \right\rangle \right] \\
&\stackrel{(a)}{=} \mathbb{E}_{G_t|G_{1:t-1}} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] + \mathbb{E}_{G_t|G_{1:t-1}} \left[\frac{1}{N^2} \sum_{i=1}^N \left\| \frac{\xi_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\
&\stackrel{(b)}{=} \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{1}{N^2} \sum_{i=1}^N \sum_{l=1}^d \frac{\mathbb{E}_{G_t|G_{1:t-1}} [[\xi_{t,i}]_l^2]}{[u_{t,i}]_l} \\
&\stackrel{(c)}{\leq} \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{d}{N} \frac{\sigma^2}{\epsilon},
\end{aligned}$$

510 where (a) is due to $\mathbb{E}_{G_t|G_{1:t-1}} [\xi_{t,i}] = 0$ and $\xi_{t,i}$ is independent of $x_{t,j}$, $u_{t,j}$ for all j , and ξ_j , for all
511 $j \neq i$, (b) comes from the fact that $x_{t,i}$, $u_{t,i}$ are fixed given $G_{1:t}$, (c) is due to $\mathbb{E}_{G_t|G_{1:t-1}} [[\xi_{t,i}]_l^2] \leq \sigma^2$
512 and $[u_{t,i}]_l \geq \epsilon$ by definition. Then we have

$$\begin{aligned}
\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] &= \mathbb{E}_{G_{1:t-1}} \left[\mathbb{E}_{G_t|G_{1:t-1}} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \right] \\
&\leq \mathbb{E}_{G_{1:t-1}} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{d}{N} \frac{\sigma^2}{\epsilon} \right] \\
&= \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] + \frac{d}{N} \frac{\sigma^2}{\epsilon}. \tag{29}
\end{aligned}$$

513 In traditional analysis of SGD-like distributed algorithms, the term corresponding to
514 $\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right]$ will be merged with the first order descent when the stepsize is cho-
515 sen to be small enough. However, in our case, the term cannot be merged because it is different from
516 the first order descent in our algorithm. A brute-force upper bound is possible but this will lead to a
517 worse convergence rate in terms of N . Thus, we need a more detailed analysis for the term in the
518 following.

$$\begin{aligned}
& \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] \\
&= \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} + \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right] \\
&\leq 2\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right] \\
&\leq 2\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N \left\| \nabla f_i(x_{t,i}) \odot \left(\frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right]
\end{aligned}$$

$$\leq 2\mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right].$$

519 Summing over T , we have

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] \\ & \leq 2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right]. \end{aligned} \quad (30)$$

520 For the last term on RHS of (30), we can bound it similarly as what we did for D_2 from (23) to (24),
521 which yields

$$\begin{aligned} \sum_{t=1}^T \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right] & \leq \sum_{t=1}^T \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \frac{1}{2\epsilon^{1.5}} \|u_{t,i} - \bar{U}_t\|_1 \right] \\ & = \sum_{t=1}^T \mathbb{E} \left[\frac{1}{N} G_\infty^2 \frac{1}{2\epsilon^2} \|\bar{U}_t \mathbf{1}^T - U_t\|_{abs} \right] \\ & \leq \sum_{t=1}^T \mathbb{E} \left[\frac{1}{N} G_\infty^2 \frac{1}{2\epsilon^2} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs} \right] \\ & \leq \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[\sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right]. \end{aligned} \quad (31)$$

522 Further, we have

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] \\ & \leq 2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] \\ & = 2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] \end{aligned}$$

523 and the last term on RHS of the above inequality can be bounded following similar procedures from
524 (16) to (21), as what we did for D_3 . Completing the procedures yields

$$\begin{aligned} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] & \leq \sum_{t=1}^T \mathbb{E} \left[L \frac{1}{\epsilon} \frac{1}{N} \sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 \right] \\ & \leq \sum_{t=1}^T \mathbb{E} \left[L \frac{1}{\epsilon} \frac{1}{N} \alpha^2 \left(\frac{1}{1-\lambda} \right) N d G_\infty^2 \frac{1}{\epsilon} \right] \\ & = T L \frac{1}{\epsilon^2} \alpha^2 \left(\frac{1}{1-\lambda} \right) d G_\infty^2. \end{aligned} \quad (32)$$

525 Finally, combining (29) to (32), we get

$$\begin{aligned}
\sum_{t=1}^T \mathbb{E} \left[\left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] &\leq 4 \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 4TL \frac{1}{\epsilon^2} \alpha^2 \left(\frac{1}{1-\lambda} \right) dG_\infty^2 \\
&\quad + 2 \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[\sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + T \frac{d}{N} \frac{\sigma^2}{\epsilon} \\
&\leq 4 \frac{1}{\sqrt{\epsilon}} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] + 4TL \frac{1}{\epsilon^2} \alpha^2 \left(\frac{1}{1-\lambda} \right) dG_\infty^2 \\
&\quad + 2 \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[\sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + T \frac{d}{N} \frac{\sigma^2}{\epsilon}.
\end{aligned}$$

526 where the last inequality is due to each element of \bar{U}_t is lower bounded by ϵ by definition.

527 Combining all above, we obtain

$$\begin{aligned}
&\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
&\leq \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) \\
&\quad + \frac{L}{T} \alpha \left(\frac{\beta_1}{1-\beta_1} \right)^2 \frac{G_\infty^2}{\sqrt{N}} \frac{1}{\epsilon^2} \frac{1}{1-\lambda} \mathbb{E}[\mathcal{V}_T] \\
&\quad + \frac{8L}{T} \alpha \frac{1}{\sqrt{\epsilon}} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] + 8L^2 \alpha \frac{1}{\epsilon^2} \alpha^2 \left(\frac{1}{1-\lambda} \right) dG_\infty^2 \tag{33} \\
&\quad + \frac{4L}{T} \alpha \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[\sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + 2L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon} \\
&\quad + \frac{2}{T} \frac{\beta_1}{1-\beta_1} G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{\sqrt{N}} \mathbb{E} \left[\frac{1}{1-\lambda} \mathcal{V}_T \right] \\
&\quad + \frac{2}{T} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \frac{\lambda}{1-\lambda} \mathbb{E}[\mathcal{V}_T] \\
&\quad + \frac{3}{T} \left(\sum_{t=1}^T L \left(\frac{1}{1-\lambda} \right)^2 \alpha^2 dG_\infty^2 \frac{1}{\epsilon^{1.5}} + \sum_{t=1}^T L \left(\frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 d \frac{G_\infty^2}{\epsilon^{1.5}} \right) \\
&= \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + 2L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon} + 8L\alpha \frac{1}{\sqrt{\epsilon}} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
&\quad + 3\alpha^2 d \left(\left(\frac{\beta_1}{1-\beta_1} \right)^2 + \left(\frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 8\alpha^3 L^2 \left(\frac{1}{1-\lambda} \right) d \frac{G_\infty^2}{\epsilon^2} \\
&\quad + \frac{1}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left(L\alpha \left(\frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2L\alpha \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E}[\mathcal{V}_T].
\end{aligned}$$

528 where $\mathcal{V}_T := \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs}$. Set $\alpha = \frac{1}{\sqrt{dT}}$ and when $\alpha \leq \frac{\epsilon^{0.5}}{16L}$, we further have

$$\begin{aligned}
&\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
&\leq \frac{4}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + 4L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon}
\end{aligned}$$

$$\begin{aligned}
& + 6\alpha^2 d \left(\left(\frac{\beta_1}{1-\beta_1} \right)^2 + \left(\frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 16\alpha^3 L^2 \left(\frac{1}{1-\lambda} \right) d \frac{G_\infty^2}{\epsilon^2} \\
& + \frac{2}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left(L\alpha \left(\frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2L\alpha \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E}[\mathcal{V}_T] \\
& = \frac{4\sqrt{d}}{\sqrt{T}} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + 4L \frac{\sqrt{d}}{\sqrt{T}} \frac{1}{N} \frac{\sigma^2}{\epsilon} \\
& + 6 \frac{1}{T} \left(\left(\frac{\beta_1}{1-\beta_1} \right)^2 + \left(\frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 16 \frac{1}{T^{1.5} d^{0.5}} L^2 \left(\frac{1}{1-\lambda} \right) \frac{G_\infty^2}{\epsilon^2} \\
& + \frac{2}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left(\frac{L}{\sqrt{Td}} \left(\frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2 \frac{L}{\sqrt{Td}} \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E}[\mathcal{V}_T] \\
& \leq C_1 \frac{\sqrt{d}}{\sqrt{T}} \left(\mathbb{E}[f(Z_1)] - \min_z f(z) + \frac{\sigma^2}{N} \right) + \frac{1}{T} C_2 + \frac{1}{T^{1.5} d^{0.5}} C_3 \\
& + \left(\frac{1}{TN^{0.5}} C_4 + \frac{1}{T^{1.5} d^{0.5} N^{0.5}} C_5 \right) \mathbb{E}[\mathcal{V}_T],
\end{aligned}$$

529 where the first inequality is obtained by moving the term $8L\alpha \frac{1}{\sqrt{\epsilon}} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right]$ on the
530 RHS of (33) to the LHS to cancel it using the assumption $8L\alpha \frac{1}{\sqrt{\epsilon}} \leq \frac{1}{2}$ followed by multiplying both
531 sides by 2. The constants introduced in the last step are defined as following

$$\begin{aligned}
C_1 &= \max(4, 4L/\epsilon), \\
C_2 &= 6 \left(\left(\frac{\beta_1}{1-\beta_1} \right)^2 + \left(\frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}}, \\
C_3 &= 16L^2 \left(\frac{1}{1-\lambda} \right) \frac{G_\infty^2}{\epsilon^2}, \\
C_4 &= \frac{2}{\epsilon^{1.5}} \frac{1}{1-\lambda} \left(\lambda + \frac{\beta_1}{1-\beta_1} \right) G_\infty^2, \\
C_5 &= \frac{2}{\epsilon^2} \frac{1}{1-\lambda} L \left(\frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 + \frac{4}{\epsilon^2} \frac{\lambda}{1-\lambda} L G_\infty^2.
\end{aligned}$$

532 Substituting into $Z_1 = \bar{X}_1$ completes the proof. \square

533 C Proof of Theorem 3

534 Under some assumptions stated in Theorem 2, we have that

$$\begin{aligned}
\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] & \leq C_1 \frac{\sqrt{d}}{\sqrt{T}} \left(\mathbb{E}[f(\bar{X}_1)] - \min_z f(z) + \frac{\sigma^2}{N} \right) + \frac{1}{T} C_2 + \frac{1}{T^{1.5} d^{0.5}} C_3 \\
& + \left(\frac{1}{TN^{0.5}} C_4 + \frac{1}{T^{1.5} d^{0.5} N^{0.5}} C_5 \right) \mathbb{E} \left[\sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right],
\end{aligned} \tag{34}$$

535 where $\|\cdot\|_{abs}$ denotes the entry-wise L_1 norm of a matrix (i.e. $\|A\|_{abs} = \sum_{i,j} |A_{ij}|$) and
536 C_1, C_2, C_3, C_4, C_5 are defined in Theorem 2.

537 Since Algorithm 3 is a special case of 2, building on result of Theorem 2, we just need to characterize
538 the growth speed of $\mathbb{E} \left[\sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right]$ to prove convergence of Algorithm 3. By the

539 update rule of Algorithm 3, we know \hat{V}_t is non decreasing and thus

$$\begin{aligned}
\mathbb{E} \left[\sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] &= \mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^N \sum_{j=1}^d | -[\hat{v}_{t-2,i}]_j + [\hat{v}_{t-1,i}]_j | \right] \\
&= \mathbb{E} \left[\sum_{t=1}^T \sum_{i=1}^N \sum_{j=1}^d (-[\hat{v}_{t-2,i}]_j + [\hat{v}_{t-1,i}]_j) \right] \\
&= \mathbb{E} \left[\sum_{i=1}^N \sum_{j=1}^d (-[\hat{v}_{-1,i}]_j + [\hat{v}_{T-1,i}]_j) \right] \\
&= \mathbb{E} \left[\sum_{i=1}^N \sum_{j=1}^d (-[\hat{v}_{0,i}]_j + [\hat{v}_{T-1,i}]_j) \right],
\end{aligned}$$

540 where the last equality is because we defined $\hat{V}_{-1} \triangleq \hat{V}_0$ previously.

541 Further, because $\|g_{t,i}\|_\infty \leq G_\infty$ for all t, i and $v_{t,i}$ is a exponential moving average of $g_{k,i}^2, k =$
542 $1, 2, \dots, t$, we know $|\hat{v}_{t,i}]_j| \leq G_\infty^2$, for all t, i, j . In addition, by update rule of \hat{V}_t , we also know
543 each element of \hat{V}_t also cannot be greater than G_∞^2 , i.e. $|\hat{v}_{t,i}]_j| \leq G_\infty^2$, for all t, i, j . Given the fact
544 that $[\hat{v}_{0,i}]_j \geq 0$, we have

$$\mathbb{E} \left[\sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] = \mathbb{E} \left[\sum_{i=1}^N \sum_{j=1}^d (-[\hat{v}_{0,i}]_j + [\hat{v}_{T-1,i}]_j) \right] \leq \mathbb{E} \left[\sum_{i=1}^N \sum_{j=1}^d G_\infty^2 \right] = NdG_\infty^2.$$

545 Substituting the above into (34), we have

$$\begin{aligned}
\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] &\leq C_1 \frac{\sqrt{d}}{\sqrt{T}} \left(\mathbb{E}[f(\bar{X}_1)] - \min_z f(z) + \frac{\sigma^2}{N} \right) + \frac{1}{T} C_2 + \frac{1}{T^{1.5} d^{0.5}} C_3 \\
&\quad + \frac{d}{T} C_4 \sqrt{N} G_\infty^2 + \frac{\sqrt{d}}{T^{1.5}} C_5 \sqrt{N} G_\infty^2 \\
&= C'_1 \frac{\sqrt{d}}{\sqrt{T}} \left(\mathbb{E}[f(\bar{X}_1)] - \min_z f(z) + \frac{\sigma^2}{N} \right) + \frac{1}{T} C'_2 + \frac{1}{T^{1.5} d^{0.5}} C'_3 \\
&\quad + \frac{d}{T} \sqrt{N} C'_4 + \frac{\sqrt{d}}{T^{1.5}} \sqrt{N} C'_5,
\end{aligned} \tag{35}$$

546 where we have

$$C'_1 = C_1 \quad C'_2 = C_2 \quad C'_3 = C_3 \quad C'_4 = C_4 G_\infty^2 \quad C'_5 = C_5 G_\infty^2. \tag{36}$$

547 and we conclude the proof. \square

548 D Additional Experiments and Details

549 In this section, we compare the training loss and testing accuracy of different algorithms, namely
550 Decentralized Stochastic Gradient Descent (DGD), Decentralized Adam (DADAM) and our proposed
551 Decentralized AMSGrad, with different stepsizes on heterogeneous data distribution. We use 5 nodes
552 and the heterogeneous data distribution is created by assigning each node with data of only two labels.
553 Note that there are no overlapping labels between different nodes. For all algorithms, we compare
554 stepsizes in the grid $[10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}]$.

555 Figure 2 shows the training loss and test accuracy for DGD algorithm. We observe that the stepsize
556 10^{-3} works best for DGD in terms of test accuracy and 10^{-1} works best in terms of training loss.
557 This difference is caused by the inconsistency among the value of parameters on different nodes when

the stepsize is large. The training loss is calculated as the average of the loss value of different local models evaluated on their local training batch. Thus, while the training loss is small at a particular node, the test accuracy will be low when evaluating data with labels not seen by the node (recall that each node contains data with different labels since we are in the heterogeneous setting).




Figure 2: Performance comparison of different stepsizes for DGD




Figure 3: Performance comparison of different stepsizes for decentralized AMSGrad




Figure 4: Performance comparison of different stepsizes for DADAM

Figure 3 shows the performance of decentralized AMSGrad with different stepsizes. We see that its best performance is better than the one of DGD and the performance is more stable (the test performance is less sensitive to stepsize tuning).

565 Figure 4 displays the performance of Decentralized Adam algorithm. As expected, the performance of
566 DADAM is not as good as DGD or decentralized AMSGrad. Its divergence characteristic, highlighted
567 Section 2.3, coupled with the heterogeneity in the data amplify its non-convergence issue in our
568 experiments. From the experiments above, we can see the advantages of decentralized AMSGrad
569 in terms of both performance and ease of parameter tuning, and the importance of ensuring the
570 theoretical convergence of any newly proposed methods in the presented setting.