

---

# Fast Two-Timescale Stochastic EM Algorithms

---

Anonymous Author(s)

Affiliation

Address

email

## Abstract

The Expectation-Maximization (EM) algorithm is a popular choice for learning latent variable models. Variants of the EM have been initially introduced by [21], using incremental updates to scale to large datasets, and by [25, 9], using Monte Carlo (MC) approximations to bypass the intractable conditional expectation of the latent data for most nonconvex models. In this paper, we propose a general class of methods called Two-Timescale EM Methods based on a two-stage approach of stochastic updates to tackle an essential nonconvex optimization task for latent variable models. We motivate the choice of a double dynamic by invoking the variance reduction virtue of each stage of the method on both sources of noise: the index sampling for the incremental update and the MC approximation. We establish finite-time and global convergence bounds for nonconvex objective functions. Numerical applications are also presented to illustrate our findings.

## 1 Introduction

Learning latent variable models is critical for modern machine learning problems, see (e.g.,) [19] for references. We formulate the training of such model as an empirical risk minimization problem:

$$\min_{\theta \in \Theta} \bar{L}(\theta) := L(\theta) + r(\theta) \quad \text{with} \quad L(\theta) = \frac{1}{n} \sum_{i=1}^n L_i(\theta) := \frac{1}{n} \sum_{i=1}^n \{ -\log g(y_i; \theta) \}. \quad (1)$$

We denote the observations by  $\{y_i\}_{i=1}^n$ ,  $\Theta \subset \mathbb{R}^d$  is the convex parameters set. We consider a smooth convex regularization noted  $r : \Theta \rightarrow \mathbb{R}$  and  $g(y; \theta)$  is the (incomplete) likelihood of each observation. The objective function  $\bar{L}(\theta)$  is possibly *nonconvex* and is assumed to be lower bounded. In the latent variable model,  $g(y_i; \theta)$ , is the marginal of the complete data likelihood defined as  $f(z_i, y_i; \theta)$ , i.e.,  $g(y_i; \theta) = \int_{\mathcal{Z}} f(z_i, y_i; \theta) \mu(dz_i)$ , where  $\{z_i\}_{i=1}^n$  are the latent variables. In this paper, we make the assumption of a complete model belonging to the curved exponential family [11]:

$$f(z_i, y_i; \theta) = h(z_i, y_i) \exp \left( \langle S(z_i, y_i) | \phi(\theta) \rangle - \psi(\theta) \right), \quad (2)$$

where  $\psi(\theta)$ ,  $h(z_i, y_i)$  are scalar functions,  $\phi(\theta) \in \mathbb{R}^k$  is a vector function, and  $\{S(z_i, y_i) \in \mathbb{R}^k\}_{i=1}^n$  is the vector of sufficient statistics of the complete model. Full batch EM [10] is the method of reference for that type of task and is a two steps procedure. The **E-step** amounts to computing the conditional expectation of the complete data sufficient statistics,

$$\text{E-step: } \bar{s}(\theta) = \frac{1}{n} \sum_{i=1}^n \bar{s}_i(\theta) \quad \text{where} \quad \bar{s}_i(\theta) = \int_{\mathcal{Z}} S(z_i, y_i) p(z_i | y_i; \theta) \mu(dz_i), \quad (3)$$

and the **M-step** is given by

$$\text{M-step: } \hat{\theta} = \bar{\theta}(\bar{s}(\theta)) := \arg \min_{\vartheta \in \Theta} \{ r(\vartheta) + \psi(\vartheta) - \langle \bar{s}(\theta) | \phi(\vartheta) \rangle \}. \quad (4)$$

Two caveats of this method are the following: (a) with the explosion of data, the first step of the EM is computationally inefficient as it requires, at each iteration, a full pass over the dataset; and (b) the complexity of modern models makes the expectation in (3) intractable. So far, and to the best of our knowledge, both challenges have been addressed separately, as detailed in the sequel.

**Prior Work:** Inspired by stochastic optimization procedures, [21] and [5] develop respectively an incremental and an online variant of the E-step in models where the expectation is computable, and were then extensively used and studied in [22, 16, 4]. Some improvements of those methods have been provided and analyzed, globally and in finite-time, in [14] where variance reduction techniques taken from the optimization literature have been efficiently applied to scale the EM algorithm to large datasets. Regarding the computation of the expectation under the posterior distribution, the Monte Carlo EM (MCEM) has been introduced in the seminal paper [25] where an MC approximation for this expectation is computed. A variant of that algorithm is the Stochastic Approximation of the EM (SAEM) in [9] leveraging the power of Robbins-Monro update [24] to ensure pointwise convergence of the vector of estimated parameters using a decreasing stepsize rather than increasing the number of MC samples. The MCEM and the SAEM have been successfully applied in mixed effects models [18, 12, 3] or to do inference for joint modeling of time to event data coming from clinical trials in [7], among other applications. Recently, an incremental variant of the SAEM was proposed in [15] showing positive empirical results but its analysis is limited to asymptotic consideration. Gradient-based methods have been developed and analyzed in [26] but they remain out of the scope of this paper as they tackle the high-dimensionality issue.

**Contributions:** This paper *introduces* and *analyzes* a new class of methods which purpose is to update two proxies for the target expected quantities in a two-timescale manner. Those approximated quantities are then used to optimize the objective function (1) for modern examples and settings using the M-step of the EM algorithm. The main contributions of the paper are:

- We propose a two-timescale method based on (i) Stochastic Approximation (SA), to alleviate the problem of computing MC approximations, and on (ii) Incremental updates, to scale to large datasets. We describe in details the edges of each level of our method based on variance reduction arguments. Such class of algorithms has two advantages. First, it naturally leverages variance reduction and Robbins-Monro type of updates to tackle large-scale and highly nonlinear learning tasks. Then, it gives a simple formulation as a *scaled-gradient method* which makes the global analysis and the implementation accessible.
- We also establish global (independent of the initialization) and finite-time (true at each iteration) upper bounds on a classical sub-optimality condition in the nonconvex literature, *i.e.*, the second order moment of the gradient of the objective function.

In Section 2 we formalize both incremental and Monte Carlo variants of the EM. Then, we introduce our two-timescale class of EM algorithms for which we derive several global statistical guarantees in Section 3 for possibly *nonconvex* functions. Section 4 is devoted to numerical illustrations. The supplementary material of this paper includes proofs of our theoretical results.

## 2 Two-Timescale Stochastic EM Algorithms

We recall and formalize in this section the different methods found in the literature that aim at solving the intractable expectation and the large-scale problem. We then provide the general framework of our method that efficiently tackles the optimization problem (1).

### 2.1 Monte Carlo Integration and Stochastic Approximation

As mentioned in the Introduction, for complex and possibly nonconvex models, the expectation under the posterior distribution defined in (3) is not tractable. In that case, the first solution involves computing a Monte Carlo integration of that latter term. For all  $i \in [n]$ , draw  $\{z_{i,m} \sim p(z_i|y_i; \theta)\}_{m \in [1,M]}$  samples and compute the MC integration  $\tilde{s}$  of the quantity  $\bar{s}(\theta)$  (3):

$$\text{MC-step : } \tilde{s} := \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}, y_i). \quad (5)$$

Then update the parameter  $\hat{\theta} = \bar{\theta}(\tilde{s})$ . This algorithm bypasses the intractable expectation issue but is rather computationally expensive in order to reach point wise convergence ( $M$  needs to be large). An alternative to that stochastic algorithm is to use a Robbins-Monro (RM) type of update. We denote, at iteration  $k$ , the following quantity

$$\tilde{S}^{(k+1)} := \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}^{(k)}, y_i) \quad \text{where} \quad z_{i,m}^{(k)} \sim p(z_i|y_i; \theta^{(k)}). \quad (6)$$

80 Then, the RM update of the sufficient statistics  $\hat{\mathbf{s}}^{(k+1)}$  reads:

$$\text{SA-step : } \hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}), \quad (7)$$

81 where  $\{\gamma_k\}_{k>1} \in (0, 1)$  is a sequence of decreasing step sizes to ensure asymptotic convergence.  
 82 This is called the Stochastic Approximation of the EM (SAEM) and has been shown to converge to  
 83 a maximum likelihood of the observations under very general conditions [9]. In simple scenarios,  
 84 the samples  $\{z_{i,m}\}_{m=0}^{M-1}$  are conditionally independent and identically distributed with distribution  
 85  $p(z_i, \theta)$ . Nevertheless, in most cases, since the loss function between the observed data  $y_i$  and the  
 86 latent variable  $z_i$  can be nonconvex, sampling exactly from this distribution is not an option and the  
 87 MC batch is sampled by Markov Chain Monte Carlo (MCMC) algorithm.

88 **Role of the stepsize  $\gamma_k$ :** The sequence of decreasing positive integers  $\{\gamma_k\}_{k>1}$  controls the conver-  
 89 gence of the algorithm. It is inefficient to start with small values for step size  $\gamma_k$  and large values for  
 90 the number of simulations  $M_k$ . Rather, it is recommended that one decreases  $\gamma_k$ , as in  $\gamma_k = 1/k^\alpha$ ,  
 91 with  $\alpha \in (0, 1)$ , and keeps a constant and small number  $M_k$  bypassing the computationally involved  
 92 sampling step in (5). In practice,  $\gamma_k$  is set equal to 1 during the first few iterations to let the iterates  
 93 explore the parameter space without memory and converge quickly to a neighborhood of the target  
 94 estimate. The Stochastic Approximation is performed during the remaining iterations ensuring the  
 95 almost sure convergence of the vector of estimates.

96 This Robbins-Monro type of update constitutes the *first level* of our algorithm, needed to temper the  
 97 variance and noise introduced by the Monte Carlo integration. In the next section, we derive variants  
 98 of this algorithm to adapt to the sheer size of data of today's applications and formalize the *second*  
 99 *level* of our class of two-timescale EM methods.

## 100 2.2 Incremental and Two-Stage Stochastic EM Methods

101 Efficient strategies to scale to large datasets include incremental [21] and variance reduced [8] meth-  
 102 ods. We will explicit a general update that covers those latter variants and that represents the *second*  
 103 *level* of our algorithm, namely the incremental update of the noisy statistics  $\tilde{\mathbf{S}}^{(k+1)}$  in the SA-Step:

$$\text{Incremental-step : } \tilde{\mathbf{S}}^{(k+1)} = \tilde{\mathbf{S}}^{(k)} + \rho_{k+1}(\mathcal{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k)}). \quad (8)$$

105 Note that  $\{\rho_k\}_{k>1} \in (0, 1)$  is a sequence of step sizes,  $\mathcal{S}^{(k)}$  is a proxy for  $\tilde{\mathbf{S}}^{(k)}$ . If the stepsize  
 106 is equal to one and the proxy  $\mathcal{S}^{(k)} = \tilde{\mathbf{S}}^{(k)}$ , i.e., computed in a full batch manner as in (6), then  
 107 we recover the SAEM algorithm. Also if  $\rho_k = 1$ ,  $\gamma_k = 1$  and  $\mathcal{S}^{(k)} = \tilde{\mathbf{S}}^{(k)}$ , then we recover the  
 108 MCEM [25]. For all methods, we define a random index drawn at iteration  $k$ , noted  $i_k \in [n]$ , and  
 109  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$  as the iteration index where  $i \in [n]$  is last drawn prior to iteration  
 110  $k$ . The proposed fitTEM method draws *two* indices *independently* and uniformly as  $i_k, j_k \in [n]$ .  
 111 Thus, we define  $t_j^k = \{k' : j_{k'} = j, k' < k\}$  to be the iteration index where the sample  $j \in [n]$  is  
 112 last drawn as  $j_k$  prior to iteration  $k$  in addition to  $\tau_i^k$  which was defined *w.r.t.*  $i_k$ .

iSAEM	$\mathcal{S}^{(k+1)} = \mathcal{S}^{(k)} + \frac{1}{n}(\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(\tau_{i_k}^k)})$	(9)
vrTTEM	$\mathcal{S}^{(k+1)} = \tilde{\mathbf{S}}^{(\ell(k))} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(\ell(k))})$	(10)
fitTEM	$\mathcal{S}^{(k+1)} = \overline{\mathcal{S}}^{(k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}), \quad \overline{\mathcal{S}}^{(k+1)} = \overline{\mathcal{S}}^{(k)} + n^{-1}(\tilde{\mathbf{S}}_{j_k}^{(k)} - \tilde{\mathbf{S}}_{j_k}^{(t_{j_k}^k)})$	(11)

115 Recall  $\tilde{\mathbf{S}}_{i_k}^{(k)} = \frac{1}{M_k} \sum_{m=1}^{M_k} \mathcal{S}(z_{i_k,m}^{(k)}, y_{i_k})$  and  $z_{i_k,m}^{(k)} \sim p(z_{i_k} | y_{i_k}; \theta^{(k)})$ . The stepsize is set to  $\rho_{k+1} =$   
 116 1 for the iSAEM method and we initialize with  $\mathcal{S}^{(0)} = \tilde{\mathbf{S}}^{(0)}$ ;  $\rho_{k+1} = \rho$  is constant for the vrTTEM  
 117 and fitTEM methods. Note that we initialize as follows  $\overline{\mathcal{S}}^{(0)} = \tilde{\mathbf{S}}^{(0)}$  for the fitTEM which can be  
 118 seen as a slightly modified version of SAGA inspired by [23]. For vrTTEM we set an epoch size of  
 119  $m$  and define  $\ell(k) := m \lfloor k/m \rfloor$  as the first iteration number in the epoch that iteration  $k$  is in.

120 **Two-Timescale Stochastic EM methods:** We now introduce the general method derived using the  
 121 two variance reduction techniques described above. Algorithm 1 leverages both levels (7) and (8) in  
 122 order to output a vector of fitted parameters  $\hat{\theta}^{(K)}$  where  $K$  is a randomly chosen termination point.

---

**Algorithm 1** Two-Timescale Stochastic EM methods.

---

- 1: **Input:** Initializations  $\hat{\theta}^{(0)} \leftarrow 0, \hat{s}^{(0)} \leftarrow \tilde{S}^{(0)}, \gamma_k, \rho_k \leftarrow$  Stepsizes.
- 2: **for**  $k = 0, 1, 2, \dots, K_m - 1$  **do**
- 3:   Draw index  $i_k \in [n]$  uniformly (and  $j_k \in [n]$  for fitTEM).
- 4:   Compute  $\tilde{S}_{i_k}^{(k)}$  using the MC-step (5), for the drawn indices.
- 5:   Compute the surrogate sufficient statistics  $\mathcal{S}^{(k+1)}$  using (9) or (10) or (11).
- 6:   Compute  $\tilde{S}^{(k+1)}$  and  $\hat{s}^{(k+1)}$  using respectively (8) and (7):

$$\begin{aligned}\tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho_{k+1}(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{s}^{(k+1)} &= \hat{s}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{s}^{(k)})\end{aligned}\tag{12}$$

- 7:   Compute  $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$  via the M-step.
  - 8: **end for**
- 

The update in (12) is said to have two-timescale property as the step sizes satisfy  $\lim_{k \rightarrow \infty} \gamma_k / \rho_k < 1$  such that  $\tilde{S}^{(k+1)}$  is updated at a faster time-scale, determined by  $\rho_{k+1}$ , than  $\hat{s}^{(k+1)}$ , determined by  $\gamma_{k+1}$ . The next section introduces the main results of this paper and establishes global and finite-time bounds for the three different updates of our scheme.

### 3 Finite Time Analysis of the Two-Timescale Scheme

Following [5], it can be shown that stationary points of the objective function (1) corresponds to the stationary points of the following *nonconvex* Lyapunov function:

$$\min_{\mathbf{s} \in \mathcal{S}} V(\mathbf{s}) := \bar{L}(\bar{\theta}(\mathbf{s})) = \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(\bar{\theta}(\mathbf{s})) + r(\bar{\theta}(\mathbf{s})), \tag{13}$$

that we propose to study in this article.

#### 3.1 Assumptions and Intermediate Lemmas

Several important assumptions required to derive convergence guarantees read as follows:

**A1.** *The sets  $\mathcal{Z}, \mathcal{S}$  are compact. There exist constants  $C_S, C_Z$  such that:*

$$C_S := \max_{\mathbf{s}, \mathbf{s}' \in \mathcal{S}} \|\mathbf{s} - \mathbf{s}'\| < \infty, \quad C_Z := \max_{i \in [n]} \int_{\mathcal{Z}} |S(z, y_i)| \mu(dz) < \infty. \tag{14}$$

**A2.** *For any  $i \in [n]$ ,  $z \in \mathcal{Z}$ ,  $\theta, \theta' \in \text{int}(\Theta)^2$ , we have  $|p(z|y_i; \theta) - p(z|y_i; \theta')| \leq L_p \|\theta - \theta'\|$  where  $\text{int}(\Theta)$  denotes the interior of  $\Theta$ .*

We also recall from the introduction that we consider curved exponential family models with:

**A3.** *For any  $\mathbf{s} \in \mathcal{S}$ , the function  $\theta \mapsto L(\mathbf{s}, \theta) := r(\theta) + \psi(\theta) - \langle \mathbf{s} | \phi(\theta) \rangle$  admits a unique global minimum  $\bar{\theta}(\mathbf{s}) \in \text{int}(\Theta)$ . In addition,  $J_\phi^\theta(\bar{\theta}(\mathbf{s}))$  is full rank,  $L_\phi$ -Lipschitz and  $\bar{\theta}(\mathbf{s})$  is  $L_\theta$ -Lipschitz.*

We denote by  $H_L^\theta(\mathbf{s}, \theta)$  the Hessian (w.r.t to  $\theta$  for a given value of  $\mathbf{s}$ ) of the function  $\theta \mapsto L(\mathbf{s}, \theta) = r(\theta) + \psi(\theta) - \langle \mathbf{s} | \phi(\theta) \rangle$ , and define  $B(\mathbf{s}) := J_\phi^\theta(\bar{\theta}(\mathbf{s})) \left( H_L^\theta(\mathbf{s}, \bar{\theta}(\mathbf{s})) \right)^{-1} J_\phi^\theta(\bar{\theta}(\mathbf{s}))^\top$ .

**A4.** *It holds that  $v_{\max} := \sup_{\mathbf{s} \in \mathcal{S}} \|B(\mathbf{s})\| < \infty$  and  $0 < v_{\min} := \inf_{\mathbf{s} \in \mathcal{S}} \lambda_{\min}(B(\mathbf{s}))$ . There exists a constant  $L_B$  such that for all  $\mathbf{s}, \mathbf{s}' \in \mathcal{S}^2$ , we have  $\|B(\mathbf{s}) - B(\mathbf{s}')\| \leq L_B \|\mathbf{s} - \mathbf{s}'\|$ .*

The class of algorithms we develop in this paper is composed of two levels where the second stage corresponds to the variance reduction trick used in [14] in order to accelerate incremental methods and reduce the variance introduced by the index sampling. The first stage is the Robbins-Monro type of update that aims at reducing the Monte Carlo noise of the quantity  $\bar{s}_i(\hat{\theta}(\hat{s}^{(r)}))$  at iteration  $r$ . We denote those latter MC fluctuations terms as follows:

$$\eta_i^{(r)} := \tilde{S}_i^{(r)} - \bar{s}_i(\vartheta^{(r)}) \quad \text{for all } i \in [n], r > 0 \quad \text{and} \quad \vartheta \in \Theta. \tag{15}$$

For instance, we consider that the MC approximation is unbiased if for all  $i \in [n]$  and  $m \in [1, M]$ , the samples  $z_{i,m} \sim p(z_i|y_i; \theta)$  are i.i.d. under the posterior distribution, i.e.,  $\mathbb{E}[\eta_i^{(r)}|\mathcal{F}_r] = 0$  where  $\mathcal{F}_r$  is the filtration up to iteration  $r$ . The following results are derived under the assumption of control of the fluctuations implied by the approximation, and is stated as follows:

**A5.** *There exist a positive sequence of MC batch size  $\{M_r\}_{r>0}$  and constants  $(c, c_\eta)$  such that for all  $k > 0$ ,  $i \in [n]$  and  $\vartheta \in \Theta$ :*

$$\mathbb{E}[\|\eta_i^{(r)}\|^2] \leq \frac{c_\eta}{M_r} \quad \text{and} \quad \mathbb{E}[\|\mathbb{E}[\eta_i^{(r)}|\mathcal{F}_r]\|^2] \leq \frac{c}{M_r}. \quad (16)$$

We can prove two important results on the Lyapunov function. The first one suggests smoothness:

**Lemma 1.** *[14] Assume A1-A4. For all  $\mathbf{s}, \mathbf{s}' \in \mathcal{S}$  and  $i \in [n]$ , we have*

$$\|\bar{\mathbf{s}}_i(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \bar{\mathbf{s}}_i(\bar{\boldsymbol{\theta}}(\mathbf{s}'))\| \leq L_{\mathbf{s}} \|\mathbf{s} - \mathbf{s}'\|, \quad \|\nabla V(\mathbf{s}) - \nabla V(\mathbf{s}')\| \leq L_V \|\mathbf{s} - \mathbf{s}'\|, \quad (17)$$

where  $L_{\mathbf{s}} := C_Z L_p L_\theta$  and  $L_V := v_{\max}(1 + L_{\mathbf{s}}) + L_B C_S$ .

We also establish a growth condition on the gradient of  $V$  related to the mean field of the algorithm:

**Lemma 2.** *Assume A3,A4. For all  $\mathbf{s} \in \mathcal{S}$ ,*

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2. \quad (18)$$

### 3.2 Global Convergence of Incremental and Two-Timescale Stochastic EM Algorithms

We present in this section a finite-time and global (independent of the initialization) analysis of both the incremental and two-timescale variants of the Stochastic Approximation of the EM algorithm.

The following result for the iSAEM algorithm is derived under the control of the Monte Carlo fluctuations as described by Assumption A5 and is built upon an intermediary Lemma, detailed in the supplementary material, characterizing the quantity of interest  $(\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)})$ . Typically, the controls exhibited above are of interest when the number of MC samples  $M_k$  increase with  $k$ .

**Theorem 1.** *Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and consider the iSAEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ . We also set  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k^\alpha \alpha c_1 \bar{L}}$  where  $a \in (0, 1)$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_m$ , then it holds:*

$$v_{\max}^{-2} \sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)})] + \sum_{k=0}^{K_m-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2].$$

Two important intermediate Lemmas are needed in order to establish finite-time bounds for the vrTTEM and the fitTEM methods. We first derive an identity for the drift term of the vrTTEM :

**Lemma 3.** *Consider the vrTTEM update in (10) with  $\rho_k = \rho$ , it holds for all  $k > 0$*

$$\begin{aligned} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_{\mathbf{s}}^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2], \end{aligned}$$

where we recall that  $\ell(k)$  is the first iteration number in the epoch that iteration  $k$  is in.

The second one derives an identity for the quantity  $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2]$  using the fitTEM update:

**Lemma 4.** *Consider the fitTEM update in (11) with  $\rho_k = \rho$ . It holds for all  $k > 0$  that*

$$\begin{aligned} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_{\mathbf{s}}^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]. \end{aligned}$$

176 Recalling that  $K$  is an independent discrete r.v. drawn from  $\{1, \dots, K_m\}$  with distribution  
 177  $\{\gamma_k/P_m, 0 \leq k \leq K_m - 1\}$ , then the convergence criterion used in our study reads

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] = \frac{1}{P_m} \sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2],$$

178 where the expectation is over the stochasticity of the algorithm and  $P_m = \sum_{\ell=0}^{K_m-1} \gamma_\ell$ .

179 Denote  $\Delta V = V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_m)})$ . We now state the main result regarding the vrTTEM method:

180 **Theorem 2.** Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive  
 181 step sizes and consider the vrTTEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_m$ .  
 182 Setting  $\bar{L} = \max\{L_s, L_V\}$ ,  $\rho = \mu/(c_1 \bar{L} n^{2/3})$ ,  $m = nc_1^2/(2\mu^2 + \mu c_1^2)$ , a constant  $\mu \in (0, 1)$ ,  
 183  $\gamma_{k+1} = 1/(k^a \bar{L})$  where  $a \in (0, 1)$ , it holds:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{2n^{2/3} \bar{L}}{\mu P_m v_{\min}^2 v_{\max}^2} \left( \mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right).$$

184 Furthermore, the fitTEM method has the following convergence rate:

185 **Theorem 3.** Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive  
 186 step sizes and consider the fitTEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq$   
 187  $K_m$ . Setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\beta = 1/(\alpha n)$ ,  $\rho = 1/(\alpha c_1 \bar{L} n^{2/3})$ ,  
 188  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$  and  $\gamma_{k+1} = 1/(k^a \alpha c_1 \bar{L})$  where  $a \in (0, 1)$ , it holds:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{4\alpha \bar{L} n^{2/3}}{P_m v_{\min}^2 v_{\max}^2} \left( \mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \Gamma^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right).$$

189 Note that in those two bounds, the quantities  $\tilde{\eta}^{(k+1)}$  and  $\Xi^{(k+1)}$  depend only on the MC fluctuations  
 190  $\mathbb{E}[\|\eta_{i_k}^{(k)}\|^2]$  and some constants. While Theorem 1 suffers only from the MC noise created by the la-  
 191 tent data sampling step, Theorem 2 and Theorem 3 exhibit in their convergence bounds *two different*  
 192 *phases*. The upper bounds display a *bias term* due to the initial conditions, *i.e.*, the term  $\Delta V$ , and a  
 193 *double dynamic* burden exemplified by the term  $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]$ .

194 Indeed, the following remarks are worth doing on this quantity. (i) This term is the price we pay for  
 195 the two-timescale dynamic and corresponds to the gap between the two *asynchronous* updates (one  
 196 on  $\hat{\mathbf{s}}^{(k)}$  and the other on  $\tilde{S}^{(k)}$ ). (ii) It is readily understood that if  $\rho = 1$ , *i.e.*, there is no variance  
 197 reduction, then for any  $k > 0$

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] = \mathbb{E}[\|\mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)}\|^2] = 0 \quad \text{with} \quad \hat{\mathbf{s}}^{(0)} = \tilde{S}^{(0)} = 0,$$

198 which strengthen the fact that this quantity characterizes the impact of the variance reduction tech-  
 199 nique introduced in our class of methods. The following Lemma characterizes this gap:

200 **Lemma 5.** Considering a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho \in (0, 1)$ , we have

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \leq \frac{\rho}{1 - \rho} \sum_{\ell=0}^k (1 - \gamma_\ell)^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}),$$

201 where  $\mathcal{S}^{(k)}$  is defined either by (10) (vrTTEM) or (11) (fitTEM).



## 4 Numerical Examples

This section presents several numerical applications for our proposed class of Algorithms 1.

### 4.1 Gaussian Mixture Models

We begin by a simple and illustrative example. The authors acknowledge that the following model can be trained using deterministic EM-type of algorithms but propose to apply stochastic methods, including theirs, and to compare their performances. Given  $n$  observations  $\{y_i\}_{i=1}^n$ , we want to fit a Gaussian Mixture Model (GMM) whose distribution is modeled as a Gaussian mixture of  $M$  components, each with a unit variance. Let  $z_i \in [M]$  be the latent labels of each component, the complete log-likelihood is defined as:

$$\log f(z_i, y_i; \theta) = \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) [\log(\omega_m) - \mu_m^2/2] + \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) \mu_m y_i + \text{constant}.$$

where  $\theta := (\omega, \mu)$  with  $\omega = \{\omega_m\}_{m=1}^{M-1}$  are the mixing weights with the convention  $\omega_M = 1 - \sum_{m=1}^{M-1} \omega_m$  and  $\mu = \{\mu_m\}_{m=1}^M$  are the means. We use the penalization  $r(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \log \text{Dir}(\omega; M, \epsilon)$  where  $\delta > 0$  and  $\text{Dir}(\cdot; M, \epsilon)$  is the  $M$  dimensional symmetric Dirichlet distribution with concentration parameter  $\epsilon > 0$ . The constraint set is given by  $\Theta = \{\omega_m, m = 1, \dots, M-1 : \omega_m \geq 0, \sum_{m=1}^{M-1} \omega_m \leq 1\} \times \{\mu_m \in \mathbb{R}, m = 1, \dots, M\}$ . In the following experiments on synthetic data, we generate 30 synthetic datasets of size  $n = 10^5$  from a GMM model with  $M = 2$  components with two mixtures with means  $\mu_1 = -\mu_2 = 0.5$ . We run the EM method until convergence (to double precision) to obtain the ML estimate  $\mu^*$  averaged on 50 datasets. We compare the EM, iEM, SAEM, iSAEM, vrTTEM and fitTEM methods in terms of their precision measured by  $|\mu - \mu^*|^2$ . We set the stepsize of the SA-step of all method as  $\gamma_k = 1/k^\alpha$  with  $\alpha = 0.5$ , and the stepsizes  $\rho_k$  for vrTTEM and the fitTEM to a constant stepsize equal to  $1/n^{2/3}$ . The number of MC samples is fixed to  $M = 10$  chains. Figure 1 shows the precision  $|\mu - \mu^*|^2$  for the different methods against the epoch(s) elapsed (one epoch equals  $n$  iterations). vrTTEM and fitTEM methods outperform the other stochastic methods, supporting the benefits of our scheme.




Figure 1: Precision  $|\mu^{(k)} - \mu^*|^2$  per epoch

### 4.2 Deformable Template Model for Image Analysis

Let  $(y_i, i \in [n])$  be observed gray level images defined on a grid of pixels. Let  $u \in \mathcal{U} \subset \mathbb{R}^2$  denotes the pixel index on the image and  $x_u \in \mathcal{D} \subset \mathbb{R}^2$  its location. The model used in this experiment suggests that each image  $y_i$  is a deformation of a template, noted  $I : \mathcal{D} \rightarrow \mathbb{R}$ , common to all images of the dataset:

$$y_i(u) = I(x_u - \Phi_i(x_u, z_i)) + \varepsilon_i(u) \quad (19)$$

where  $\phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a deformation function,  $z_i$  some latent variable parameterizing this deformation and  $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$  is an observation error. The template model, given  $\{p_k\}_{k=1}^{k_p}$  landmarks on the template, a fixed known kernel  $\mathbf{K}_p$  and a vector of parameters  $\beta \in \mathbb{R}^{k_p}$  is defined as follows:

$$I_\xi = \mathbf{K}_p \beta, \quad \text{where} \quad (\mathbf{K}_p \beta)(x) = \sum_{k=1}^{k_p} \mathbf{K}_p(x, p_k) \beta_k.$$

Given a set of landmarks  $\{g_k\}_{k=1}^{k_g}$  and a fixed kernel  $\mathbf{K}_g$ , we parameterize the deformation  $\Phi_i$  as:

$$\Phi_i = \mathbf{K}_g z_i \quad \text{where} \quad (\mathbf{K}_g z_i)(x) = \sum_{k=1}^{k_g} \mathbf{K}_g(x, g_k) \left( z_i^{(1)}(k), z_i^{(2)}(k) \right),$$

where we put a Gaussian prior on the latent variables,  $z_i \sim \mathcal{N}(0, \Gamma)$  and  $z_i \in (\mathbb{R}^{k_g})^2$ . The vector of parameters we estimate is thus  $\theta = (\beta, \Gamma, \sigma)$ .

**Numerical Experiment:** We apply model (19) and our algorithms 1 to a collection of handwritten digits, called the US postal database [13], featuring  $n = 1000$  ( $16 \times 16$ )-pixel images for each class of digits from 0 to 9. The main difficulty with these data comes from the geometric dispersion

within each class of digit as shown Figure 2 for digit 5. We thus ought to use our deformable template model (19) in order to account for both sources of variability: the intrinsic template to each class of digit and the small and local deformation in each observed image.



Figure 2: Training set of the USPS database (20 images for digit 5)

Figure 3 shows the resulting synthetic images for digit 5 through several epochs, for the batch method, the online SAEM, the incremental SAEM and the various TTS methods. For all methods, the initialization of the template (20) is the mean of the gray level images. In our experiments, we have chosen Gaussian kernels for both,  $K_p$  and  $K_g$ , defined on  $\mathbb{R}^2$  and centered on the landmark points  $\{p_k\}_{k=1}^{k_p}$  and  $\{g_k\}_{k=1}^{k_g}$  with standard respective standard deviations of 0.12 and 0.3. We set  $k_p = 15$  and  $k_g = 6$  equidistributed landmarks points on the grid for the training procedure. Those hyperparameters are inspired by a relevant study in [2]. In particular, the choice of the geometric covariance, indexed by  $g$ , in such study is critical since it has a direct impact on the *sharpness* of the templates. As for the photometric hyperparameter, indexed by  $p$ , both the template and the geometry are impacted, in the sense that with a large photometric variance, the kernel centered on one landmark *spreads out* to many of its neighbors.




Figure 3: (USPS Digits) Estimation of the template. From top to bottom: batch, online, iSAEM, vrTTEM and fitTEM through 7 epochs. Note that Batch method templates are replicated in-between epochs for a fair comparison with incremental variants.

As the iterations proceed, the templates become sharper. Figure 3 displays the virtue of the vrTTEM and fitTEM methods that obtain a more *contrasted* and *accurate* template estimate. The incremental and online version are looking much better on the very first epochs compared to the batch method, which is intuitive given the high computational cost of the latter. After a few epochs, the batch SAEM estimates similar template as the incremental and online methods due to their high variance. Our variance reduced and fast incremental variants are effective in the long run and sharpen the final template estimates contrasting between the background and the regions of interest in the image.

## 5 Conclusion

This paper introduces a new class of two-timescale EM methods for learning latent variable models. In particular, the models dealt with in this paper belong to the curved exponential family and are possibly nonconvex. The nonconvexity of the problem is tackled using a Robbins-Monro type of update, which represents the *first level* of our class of methods. The scalability with the number of samples is performed through a variance reduced and incremental update, the *second* and last level of our newly introduced scheme. The various algorithms are interpreted as scaled gradient methods, in the space of the sufficient statistics, and our convergence results are *global*, in the sense of independence of the initial values, and *non-asymptotic*, *i.e.*, true for any random termination number. Numerical examples illustrate the benefits of our scheme on synthetic and real tasks.



## References

- [1] Stéphanie Allasonnière, Yali Amit, and Alain Trouvé. Towards a coherent statistical framework for dense deformable template estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):3–29, 2007.
- [2] Stéphanie Allasonnière, Estelle Kuhn, Alain Trouvé, et al. Construction of bayesian deformable models via a stochastic approximation algorithm: a convergence study. *Bernoulli*, 16(3):641–678, 2010.
- [3] Charlotte Baey, Samis Trevezas, and Paul-Henry Cournède. A non linear mixed effects model of plant growth and estimation via stochastic variants of the em algorithm. *Communications in Statistics-Theory and Methods*, 45(6):1643–1669, 2016.
- [4] Olivier Cappé. Online EM algorithm for hidden markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- [5] Olivier Cappé and Eric Moulines. On-line expectation–maximization algorithm for latent data models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):593–613, 2009.
- [6] Bradley P Carlin and Siddhartha Chib. Bayesian model choice via markov chain monte carlo methods. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(3):473–484, 1995.
- [7] Arindom Chakraborty and Kalyan Das. Inferences for joint modelling of repeated ordinal scores and time to event data. *Computational and mathematical methods in medicine*, 11(3):281–295, 2010.
- [8] Jianfei Chen, Jun Zhu, Yee Whye Teh, and Tong Zhang. Stochastic expectation maximization with variance reduction. In *Advances in Neural Information Processing Systems*, pages 7978–7988, 2018.
- [9] Bernard Delyon, Marc Lavielle, and Eric Moulines. Convergence of a stochastic approximation version of the em algorithm. *Ann. Statist.*, 27(1):94–128, 03 1999. doi: 10.1214/aos/1018031103. URL <https://doi.org/10.1214/aos/1018031103>.
- [10] Arthur P Dempster, Nan M Laird, and Donald B Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the royal statistical society. Series B (methodological)*, pages 1–38, 1977.
- [11] Bradley Efron et al. Defining the curvature of a statistical problem (with applications to second order efficiency). *The Annals of Statistics*, 3(6):1189–1242, 1975.
- [12] James P Hughes. Mixed effects models with censored data with application to hiv rna levels. *Biometrics*, 55(2):625–629, 1999.
- [13] Jonathan J. Hull. A database for handwritten text recognition research. *IEEE Transactions on pattern analysis and machine intelligence*, 16(5):550–554, 1994.
- [14] Belhal Karimi, Hoi-To Wai, Éric Moulines, and Marc Lavielle. On the global convergence of (fast) incremental expectation maximization methods. In *Advances in Neural Information Processing Systems*, pages 2833–2843, 2019.
- [15] Estelle Kuhn, Catherine Matias, and Tabea Rebafka. Properties of the stochastic approximation em algorithm with mini-batch sampling. *arXiv preprint arXiv:1907.09164*, 2019.

- 317 [16] Percy Liang and Dan Klein. Online em for unsupervised models. In *Proceedings of human*  
 318 *language technologies: The 2009 annual conference of the North American chapter of the*  
 319 *association for computational linguistics*, pages 611–619, 2009.
- 320 [17] Florian Maire, Eric Moulines, and Sidonie Lefebvre. Online em for functional data, 2016.  
 321 URL <http://arxiv.org/abs/1604.00570>. cite arxiv:1604.00570v1.pdf.
- 322 [18] Charles E McCulloch. Maximum likelihood algorithms for generalized linear mixed models.  
 323 *Journal of the American statistical Association*, 92(437):162–170, 1997.
- 324 [19] Geoffrey McLachlan and Thriyambakam Krishnan. *The EM algorithm and extensions*, volume  
 325 382. John Wiley & Sons, 2007.
- 326 [20] Sean P Meyn and Richard L Tweedie. *Markov chains and stochastic stability*. Springer Science  
 327 & Business Media, 2012.
- 328 [21] Radford M Neal and Geoffrey E Hinton. A view of the EM algorithm that justifies incremental,  
 329 sparse, and other variants. In *Learning in graphical models*, pages 355–368. Springer, 1998.
- 330 [22] Hien D Nguyen, Florence Forbes, and Geoffrey J McLachlan. Mini-batch learning of expo-  
 331 nential family finite mixture models. *Statistics and Computing*, pages 1–18, 2020.
- 332 [23] Sashank J Reddi, Suvrit Sra, Barnabás Póczos, and Alex Smola. Fast incremental method for  
 333 nonconvex optimization. *arXiv preprint arXiv:1603.06159*, 2016.
- 334 [24] Herbert Robbins and Sutton Monro. A stochastic approximation method. *The annals of math-*  
 335 *ematical statistics*, pages 400–407, 1951.
- 336 [25] Greg CG Wei and Martin A Tanner. A monte carlo implementation of the em algorithm and  
 337 the poor man’s data augmentation algorithms. *Journal of the American statistical Association*,  
 338 85(411):699–704, 1990.
- 339 [26] Rongda Zhu, Lingxiao Wang, Chengxiang Zhai, and Quanquan Gu. High-dimensional  
 340 variance-reduced stochastic gradient expectation-maximization algorithm. In *Proceedings of*  
 341 *the 34th International Conference on Machine Learning-Volume 70*, pages 4180–4188. JMLR.  
 342 org, 2017.

## 343 A Proof of Lemma 2

344 **Lemma.** Assume A3,A4. For all  $\mathbf{s} \in \mathcal{S}$ ,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (20)$$

345 **Proof** Using A3 and the fact that we can exchange integration with differentiation and the Fisher's  
346 identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left( \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left( \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (21)$$

347 Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \mathbf{s}.$$

348 Taking the gradient of the above map w.r.t.  $\mathbf{s}$  and using assumption A3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left( \nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})}|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}).$$

349 The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})))$$

350 where we recall  $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left( \mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top}$ . The proof of (20) follows directly  
351 from the assumption A4.  $\square$

## 352 B Proof of Theorem 1

353 Beforehand, We present two intermediary Lemmas important for the analysis of the incremental  
354 update of the iSAEM algorithm. The first one gives a characterization of the quantity  $\mathbb{E}[\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}]$ :  
355

356 **Lemma 6.** Assume A1. The update (9) is equivalent to the following update on the resulting statis-  
357 tics

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)})$$

358 Also:

$$\mathbb{E}[\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E}[\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E}[\eta_{i_k}^{(k+1)}]$$

359 where  $\bar{\mathbf{s}}^{(k)}$  is defined by (3) and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ .

360 **Proof** From update (9), we have:

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left( \tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_i^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned}$$

361 Since  $\tilde{S}_{i_k}^{(k+1)} = \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) + \eta_{i_k}^{(k+1)}$  we have

$$\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_i^k)} - \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)}$$

362 Taking the full expectation of both side of the equation leads to:

$$\begin{aligned}\mathbb{E}[\tilde{S}^{(k+1)} - \hat{s}^{(k)}] &= \mathbb{E}[\bar{s}^{(k)} - \hat{s}^{(k)}] + \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right] \\ &\quad - \frac{1}{n} \mathbb{E}[\mathbb{E}[\tilde{S}_i^{(\tau_i^k)} - \bar{s}_{i_k}(\theta^{(k)}) | \mathcal{F}_k]] + \frac{1}{n} \mathbb{E}[\eta_{i_k}^{(k+1)}]\end{aligned}$$

363 The following equalities:

$$\mathbb{E}[\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E}[\bar{s}_{i_k}(\theta^{(k)}) | \mathcal{F}_k] = \bar{s}^{(k)}$$

364 concludes the proof of the Lemma.  $\square$

365 And the following auxiliary Lemma setting an upper bound for the quantity  $\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2]$

366 **Lemma 7.** For any  $k \geq 0$  and consider the iSAEM update in (9), it holds that

$$\begin{aligned}\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2] &\leq 4\mathbb{E}[\|\bar{s}^{(k)} - \hat{s}^{(k)}\|^2] + \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(t_i^k)}\|^2] \\ &\quad + 2\frac{c_\eta}{M_k} + 4\mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right\|^2\right]\end{aligned}$$

367 **Proof** Applying the iSAEM update yields:

$$\begin{aligned}\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2] &= \mathbb{E}[\|\tilde{S}^{(k)} - \hat{s}^{(k)} - \frac{1}{n}(\tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(t_i^k)})\|^2] \\ &\leq 4\mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right\|^2\right] + 4\mathbb{E}[\|\bar{s}^{(k)} - \hat{s}^{(k)}\|^2] \\ &\quad + \frac{2}{n^2} \mathbb{E}[\|\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(t_i^k)}\|^2] + 2\frac{c_\eta}{M_k}\end{aligned}$$

368 The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E}[\|\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(t_i^k)}\|^2] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\bar{s}_i^{(k)} - \bar{s}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(t_i^k)}\|^2],$$

369 where (a) is due to Lemma 1 and which concludes the proof of the Lemma.  $\square$

370

371 **Theorem.** Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive  
372 step sizes and consider the iSAEM sequence  $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ .  
373 We also set  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k^a \alpha c_1 \bar{L}}$  where  
374  $a \in (0, 1)$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  $\hat{s}^{(k)} \in \mathcal{S}$  for any  $k \leq K_m$ , then it holds:

$$v_{\max}^{-2} \sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K)})] + \sum_{k=0}^{K_m-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2].$$

375 **Proof** Under the smoothness of the Lyapunov function  $V$  (cf. Lemma 1), we can write:

$$V(\hat{s}^{(k+1)}) \leq V(\hat{s}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} | \nabla V(\hat{s}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2$$

376 Taking the expectation on both sides yields:

$$\mathbb{E}[V(\hat{s}^{(k+1)})] \leq \mathbb{E}[V(\hat{s}^{(k)})] + \gamma_{k+1} \mathbb{E}[\langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} | \nabla V(\hat{s}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2]$$

377 Using Lemma 6, we obtain:

$$\begin{aligned}
& \mathbb{E} \left[ \langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&= \mathbb{E} \left[ \langle \bar{s}^{(k)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&\stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&\stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[ \left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&\stackrel{(a)}{\leq} \left( v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned}$$

378 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with  $\beta \rightarrow 1$ ). Note

379  $a_k = \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$  and

$$\begin{aligned}
a_k \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] &\leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (22)
\end{aligned}$$

380 We now give an upper bound of  $\mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right]$  using Lemma 7 and plug it into (22):

$$\begin{aligned}
(a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] &\leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] \\
&\quad + \gamma_{k+1} \left( \frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
&\quad + \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \quad (23)
\end{aligned}$$

381 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(t_i^{k+1})} \right\|^2 \right] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{n-1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \right)$$

382 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. For any  $\beta > 0$ , it holds

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(t_i^{k+1})} \right\|^2 \right] \\
&= \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 + 2 \langle \hat{s}^{(k+1)} - \hat{s}^{(k)} \mid \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \rangle \right] \\
&= \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 - 2\gamma_{k+1} \langle \hat{s}^{(k)} - \tilde{S}^{(k+1)} \mid \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \rangle \right] \\
&\leq \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 + \frac{\gamma_{k+1}}{\beta} \left\| \hat{s}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 + \gamma_{k+1} \beta \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right]
\end{aligned}$$

where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ & \leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2\right] \end{aligned}$$

Observe that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)})$ . Applying Lemma 7 yields

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ & \leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\ & \leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\ & \quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\ & \quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \end{aligned}$$

Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2]$$

From the above, we get

$$\begin{aligned} \Delta^{(k+1)} & \leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})\right) \Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ & \quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \end{aligned}$$

Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,

$c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$ ,  $\alpha \geq 8$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1}$$

which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} =$

$\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} & \leq 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_\ell}^{(\ell)}\|^2\right] \\ & \quad + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)}\right\|^2\right] \end{aligned}$$



391 Note  $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$  Summing on both sides over  $k = 0$  to  $k = K_m - 1$  yields:

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} \Delta^{(k+1)} \\
&= 4 \sum_{k=0}^{K_m-1} \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} [\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_m-1} \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[ \left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} 4 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&\leq \sum_{k=0}^{K_m-1} \frac{4 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} [\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_m-1} \frac{2 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \frac{4 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right]
\end{aligned} \tag{24}$$

392 We recall (23) where we have summed on both sides from  $k = 0$  to  $k = K_m - 1$ :

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\
&+ \sum_{k=0}^{K_m-1} \gamma_{k+1} \left( \frac{1}{2\beta} \left( 1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)}
\end{aligned} \tag{25}$$

393 Plugging (24) into (25) results in:

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \sum_{k=0}^{K_m-1} \tilde{\beta}_k \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&\leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_m-1} \tilde{\Gamma}_k \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned}$$

394 where

$$\begin{aligned}
\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\beta}_k &= \gamma_{k+1} \left( \frac{1}{2\beta} \left( 1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\Gamma}_k &= \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2 \left( \gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}}
\end{aligned}$$

395 and

$$\begin{aligned}
a_k &= \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right) \\
\Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1} \beta - \frac{2\gamma_{k+1} L_s^2}{n^2} (\gamma_{k+1} + \frac{1}{\beta}) \\
c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{L_s, L_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}
\end{aligned}$$

396 When, for any  $k > 0$ ,  $\tilde{\alpha}_k \geq 0$ , we have by Lemma 2 that:

$$\sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$$

397 which yields an upper bound of the gradient of the Lyapunov function  $V$  along the path of the  
398 iSAEM update and concludes the proof of the Theorem.  $\square$

## 399 C Proofs of Auxiliary Lemmas

### 400 C.1 Proof of Lemma 3 and Lemma 4

401 **Lemma.** For any  $k \geq 0$  and consider the vrTTEM update in (10) with  $\rho_k = \rho$ , it holds for all  $k > 0$   
402

$$\begin{aligned}
\mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} \right\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} \right\|^2] \\
&\quad + 2(1-\rho)^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)} \right\|^2] + 2\rho^2 \mathbb{E}[\left\| \eta_{i_k}^{(k+1)} \right\|^2]
\end{aligned}$$

403 where we recall that  $\ell(k)$  is the first iteration number in the epoch that iteration  $k$  is in.

404 **Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful through-  
405 out this proof:

$$\begin{aligned}
\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathcal{S}^{(k+1)}) \\
&= -\gamma_{k+1} \left( (1-\rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} \right] \right)
\end{aligned} \tag{26}$$

406 We observe, using the identity (26), that

$$\mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2] \leq 2\rho^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} \right\|^2] + 2\rho^2 \mathbb{E}[\left\| \bar{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} \right\|^2] + 2(1-\rho)^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)} \right\|^2] \tag{27}$$

407 For the latter term, we obtain its upper bound as

$$\begin{aligned}
\mathbb{E}[\left\| \bar{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} \right\|^2] &= \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{\ell(k)}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{\ell(k)}) \right\|^2 \right] \\
&\stackrel{(a)}{\leq} \mathbb{E}[\left\| \bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_{i_k}^{\ell(k)} \right\|^2] + \mathbb{E}[\left\| \eta_{i_k}^{(k+1)} \right\|^2] \stackrel{(b)}{\leq} L_s^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} \right\|^2] + \mathbb{E}[\left\| \eta_{i_k}^{(k+1)} \right\|^2]
\end{aligned}$$

408 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (27) proves the  
409 lemma.  $\square$

410 **Lemma.** For any  $k \geq 0$  and consider the fitTEM update in (11) with  $\rho_k = \rho$ , it holds for all  $k > 0$

$$\begin{aligned}
\mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} \right\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell_i^k)} \right\|^2] \\
&\quad + 2(1-\rho)^2 \mathbb{E}[\left\| \hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)} \right\|^2] + 2\rho^2 \mathbb{E}[\left\| \eta_{i_k}^{(k+1)} \right\|^2]
\end{aligned}$$

411 **Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful through-  
 412 out this proof:

$$\begin{aligned}
 \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}) \\
 &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1 - \rho)\tilde{\mathbf{S}}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\
 &= -\gamma_{k+1}\left((1 - \rho)\left[\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right] + \rho\left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\right]\right) \\
 &= -\gamma_{k+1}\left((1 - \rho)\left[\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right] + \rho\left[\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{S}}^{(k)} - (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})\right]\right)
 \end{aligned} \tag{28}$$

413 We observe, using the identity (28), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2] \leq 2\rho^2\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{S}}^{(k)}\|^2] + 2\rho^2\mathbb{E}[\|\bar{\mathbf{S}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1 - \rho)^2\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] \tag{29}$$

414 For the latter term, we obtain its upper bound as

$$\begin{aligned}
 \mathbb{E}[\|\bar{\mathbf{S}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E}\left[\left\|\frac{1}{n}\sum_{i=1}^n(\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}) - (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})\right\|^2\right] \\
 &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
 \end{aligned}$$

415 where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] = \frac{1}{n}\sum_{i=1}^n\mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_{\mathbf{S}}^2}{n}\sum_{i=1}^n\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

416 Substituting into (29) proves the lemma.  $\square$

## 417 C.2 Proof of Lemma 5

418 **Lemma.** Consider a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho$ , then the following inequality  
 419 holds:

$$\mathbb{E}\left[\left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \leq \frac{\rho}{1 - \rho}\sum_{\ell=0}^k(1 - \gamma_{\ell})^2(\mathbf{S}^{(\ell)} - \tilde{\mathbf{S}}^{(\ell)})$$

420 where  $\mathbf{S}^{(k)}$  is defined either by (11) (fiTTEM) or (10) (vrTTEM)

421 **Proof** We begin by writing the two-timescale update:

$$\begin{aligned}
 \tilde{\mathbf{S}}^{(k+1)} &= \tilde{\mathbf{S}}^{(k)} + \rho(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k)}) \\
 \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)})
 \end{aligned} \tag{30}$$

422 where  $\mathbf{S}^{(k+1)} = \frac{1}{n}\sum_{i=1}^n\tilde{\mathbf{S}}_i^{(t_i^k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})$  according to (11). Denote  $\delta^{(k+1)} = \hat{\mathbf{s}}^{(k+1)} -$   
 423  $\tilde{\mathbf{S}}^{(k+1)}$ . Then from (30), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1 - \rho}(1 - \gamma_{k+1})(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k+1)})$$

424 Using the telescoping sum and noting that  $\delta^{(0)} = 0$ , we have

$$\delta^{(k+1)} \leq \frac{\rho}{1 - \rho}\sum_{\ell=0}^k(1 - \gamma_{\ell+1})^2(\mathbf{S}^{(\ell+1)} - \tilde{\mathbf{S}}^{(\ell+1)})$$

425  $\square$

426 **C.3 Additional Intermediary Result**

427 **Lemma 8.** *At iteration  $k + 1$ , the drift term of update (11), with  $\rho_{k+1} = \rho$ , is equivalent to the*  
 428 *following :*

$$\begin{aligned} \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) \left( \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right) \end{aligned}$$

429 *where we recall that  $\eta_{i_k}^{(k+1)}$ , defined in (16), which is the gap between the MC approximation and*  
 430 *the expected statistics.*

431 **Proof** Using the fitTEM update  $\tilde{S}^{(k+1)} = (1 - \rho)\tilde{S}^{(k)} + \rho\mathcal{S}^{(k+1)}$  where  $\mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} -$   
 432  $\tilde{S}_{i_k}^{(t_{i_k}^k)})$  leads to the following decomposition:

$$\begin{aligned} &\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \\ &= (1 - \rho)\tilde{S}^{(k)} + \rho \left( \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) - \hat{\mathbf{s}}^{(k)} + \rho\bar{\mathbf{s}}^{(k)} - \rho\bar{\mathbf{s}}^{(k)} \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho(\tilde{S}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(k)}) + (1 - \rho) \left( \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) + \rho \left( \bar{\mathcal{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} + (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) \left( \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) \end{aligned}$$

433 *where we observe that  $\mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] = \bar{\mathbf{s}}^{(k)} - \bar{\mathcal{S}}^{(k)}$  and which concludes the proof.*

434 *Important Note:* Note that  $\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not equal to  $\eta_{i_k}^{(k+1)}$ , defined in (16), which is the gap  
 435 *between the MC approximation and the expected statistics. Indeed  $\tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not computed under the*  
 436 *same model as  $\bar{\mathbf{s}}_{i_k}^{(k)}$ .* □

## 437 D Proof of Theorem 2

438 **Theorem.** Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive  
 439 step sizes and consider the vrTTEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_m$ .  
 440 Setting  $\bar{L} = \max\{L_s, L_V\}$ ,  $\rho = \frac{\mu}{c_1 L n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$ , a constant  $\mu \in (0, 1)$ ,  $\gamma_{k+1} = \frac{1}{k^a \bar{L}}$  where  
 441  $a \in (0, 1)$ , it holds:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu P_m v_{\min}^2 v_{\max}^2} \left[ \mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right],$$

442 **Proof** Using the smoothness of  $V$  and update (10), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (31)$$

443 Denote  $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fitTEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 444 Taking expectations on both sides show that

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1 - \rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \rho \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}(1 - \rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad - \gamma_{k+1} \rho \mathbb{E}[\langle \eta_{i_k}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1} \rho v_{\min} + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\quad - \gamma_{k+1} \rho \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \gamma_{k+1}(1 - \rho) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \quad (32)$$

445 where we have used (26) in (a) and  $\mathbb{E}[\mathcal{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$  in (b), the growth condition in  
 446 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

447 Furthermore, for  $k+1 \leq \ell(k) + m$  (i.e.,  $k+1$  is in the same epoch as  $k$ ), we have

$$\begin{aligned} &\mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 \\ &\quad - 2\gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1 - \rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle] \\ &\leq \mathbb{E}[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1 - \rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2], \end{aligned}$$

448 where we first used (26) and the last inequality is due to the Young's inequality.

449 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2]$$

where  $b_k := \bar{b}_{k \bmod m}$  is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2) + \gamma_{k+1}^2\rho^2 L_V L_s^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0.$$

Note that  $\bar{b}_i$  is decreasing with  $i$  and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_s^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2}, \quad i = 1, 2, \dots, m.$$

For  $k+1 \leq \ell(k) + m$ , we have the following inequality

$$\begin{aligned} R_{k+1} &\leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \right] \\ &\quad + \gamma_{k+1} \mathbb{E} \left[ \rho \left\| \eta_{i_k}^{(k+1)} \right\|^2 - (1-\rho) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[ (1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[ \frac{\gamma_{k+1}\rho}{\beta} \left\| \eta_{i_k}^{(k+1)} \right\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

And using Lemma 3 we obtain:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2 - \gamma_{k+1}^2\rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2\rho^2 L_V L_s^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[ (1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left( \frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2 \right) \|\mathbf{h}_k\|^2 \right] \\ &\quad + \gamma_{k+1} \mathbb{E} \left[ (\rho + \rho^2 \gamma_{k+1} L_V) \left\| \eta_{i_k}^{(k+1)} \right\|^2 - (1-\rho - (1-\rho)^2 \gamma_{k+1} L_V) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[ \left( \frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2 \right) \left\| \eta_{i_k}^{(k+1)} \right\|^2 + \left( \frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2 \right) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

Rearranging the terms yields:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\quad + \underbrace{\left( b_{k+1}(1 + \gamma\beta + 2\gamma^2\rho^2 L_s^2) + \gamma^2\rho^2 L_V L_s^2 \right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)} \end{aligned}$$

where

$$\begin{aligned} \tilde{\eta}^{(k+1)} &= \left( \gamma_{k+1}(\rho + \rho^2 \gamma_{k+1} L_V) + b_{k+1}(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2) \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k+1)} \right\|^2 \right] \\ \chi^{(k+1)} &= \left( b_{k+1}(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2) - \gamma_{k+1}(1-\rho - (1-\rho)^2 \gamma_{k+1} L_V) \right) \\ \tilde{\chi}^{(k+1)} &= \chi^{(k+1)} \mathbb{E} \left[ \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

This leads, using Lemma 2, that for any  $\gamma_{k+1}$ ,  $\rho$  and  $\beta$  such that  $\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2) > 0$ ,

$$\begin{aligned} v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \\ &\quad + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \end{aligned}$$

We first remark that

$$\begin{aligned} &\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \\ &\geq \frac{\gamma_{k+1}\rho}{c_1} (1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1)) \end{aligned}$$



459 where  $c_1 = v_{\min}^{-1}$ . By setting  $\bar{L} = \max\{L_s, L_V\}$ ,  $\beta = \frac{c_1 \bar{L}}{n^{1/3}}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$  and  
 460  $\{\gamma_{k+1}\}$  any sequence of decreasing stepsizes in  $(0, 1)$ , it can be shown that there exists  $\mu \in (0, 1)$ ,  
 461 such that the following lower bound holds

$$\begin{aligned}
 & 1 - \gamma_{k+1} c_1 \rho L_V - b_{k+1} \left( \frac{c_1}{\beta} + 2\gamma_{k+1} \rho c_1 \right) \\
 & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0 \left( \frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
 & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V \mu^2}{c_1^2 n^{\frac{4}{3}}} \frac{(1 + \gamma\beta + 2\gamma^2 L_s^2)^m - 1}{\gamma\beta + 2\gamma^2 L_s^2} \left( \frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
 & \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2} (e - 1) \left( 1 + \frac{2\mu}{n} \right) \geq 1 - \mu - \mu(1 + 2\mu) \frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2}
 \end{aligned}$$

462 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2 L_s^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2 n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \quad \text{and} \quad (1 + \gamma\beta + 2\gamma^2 L_s^2)^m \leq e - 1.$$

463 and the required  $\mu$  in (b) can be found by solving the quadratic equation.

464 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_m})}{v_{\min} \rho} + 2 \sum_{k=0}^{K_m-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min} \rho}$$

465 Note that  $R_0 = \mathbb{E}[V(\hat{s}^{(0)})]$  and if  $K_m$  is a multiple of  $m$ , then  $R_{\max} = \mathbb{E}[V(\hat{s}^{(K_m)})]$ . Under the latter  
 466 condition, we have

$$\sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] + \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_m-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}]$$

467 This concludes our proof.

468 □

469 **E Proof of Theorem 3**

470 **Theorem.** Assume A1-A5. Let  $K_m$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive  
 471 step sizes and consider the fitTEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_m$ .  
 472 Setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\beta = \frac{1}{\alpha n}$ ,  $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq$   
 473  $c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$  and  $\gamma_{k+1} = \frac{1}{k^a \alpha c_1 \bar{L}}$  where  $a \in (0, 1)$ , it holds:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{4\alpha \bar{L} n^{2/3}}{P_m v_{\min}^2 v_{\max}^2} \left[ \mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \Gamma^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right].$$

474 **Proof** Using the smoothness of  $V$  and update (11), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (33)$$

475 Denote  $H_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fitTEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 476 Using Lemma 8 and the additional following identity:

$$\mathbb{E} \left[ \left( \bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)} \right) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] = 0 \quad (34)$$

477 we have:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \mathbb{E} \left[ \langle \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1 - \rho) \mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|H_{k+1}\|^2 \\ &\stackrel{(a)}{\leq} -v_{\min} \gamma_{k+1} \rho \mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1} \mathbb{E} \left[ \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|H_{k+1}\|^2 \\ &\stackrel{(b)}{\leq} -(v_{\min} \gamma_{k+1} \rho + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|H_{k+1}\|^2 \end{aligned}$$

478 where  $\xi^{(k+1)} = \mathbb{E}[\|\mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k]\|^2]$ . **Bounding**  $\mathbb{E}[\|H_{k+1}\|^2]$  Using Lemma 4, we obtain:

$$\begin{aligned} &\gamma_{k+1} (v_{\min} \rho + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)}) \right] + \tilde{\xi}^{(k+1)} + \left( (1 - \rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V \rho^2 L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (35)$$

479 where  $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 L_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)}$ . Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (36)$$

480 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \end{aligned}$$

481 Note that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}) = -\gamma_{k+1}\mathbf{H}_{k+1}$  and that in expectation we recall  
 482 that  $\mathbb{E}[\mathbf{H}_{k+1}|\mathcal{F}_k] = \rho\mathbf{h}_k + \rho\mathbb{E}[\eta_{i_k}^{(k+1)}|\mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)}]$  where  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ . Thus,  
 483 for any  $\beta > 0$ , it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \\ &\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\right. \\ &\quad \left.+ \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

484 where the last inequality is due to the Young's inequality. Plugging this into (36) yields:

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \\ &\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\right. \\ &\quad \left.+ \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

485 Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[\left(1 + \gamma_{k+1}\beta\right)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2\right. \\ &\quad \left.+ \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

486 We now use Lemma 4 on  $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2$  and obtain:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2\rho^2\mathbf{L}_s^2}{n} + \frac{(n-1)(1+\gamma_{k+1}\beta)}{n^2}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2\left(2\gamma_{k+1} + \frac{1}{\beta}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2\mathbf{L}_s^2}{n}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2\left(2\gamma_{k+1} + \frac{1}{\beta}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

487 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

488 From the above, we get

$$\begin{aligned} \Delta^{(k+1)} &\leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

489 Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{2, 1+2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k}$ ,  $\beta = \frac{1}{\alpha n}$ ,  $\rho = \frac{1}{\alpha c_1 L n^{2/3}}$ ,

490  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{\frac{4}{3}}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1}$$

491 which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} = \frac{1}{n} -$

492  $\gamma_{k+1}\beta - \gamma_{k+1}^2\rho^2 L_s^2$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} &\leq \sum_{\ell=0}^k \omega_{k,\ell} \left(2\gamma_{\ell+1}^2\rho^2 + \frac{\gamma_{\ell+1}^2\rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] \\ &\quad + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left(2\gamma_{\ell+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\tilde{S}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)} \end{aligned}$$

493 where  $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$  and  $\tilde{\epsilon}^{(\ell+1)} = \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]$ .

494 Summing on both sides over  $k = 0$  to  $k = K_m - 1$  yields:

$$\begin{aligned} \sum_{k=0}^{K_m-1} \Delta^{(k+1)} &\leq \sum_{k=0}^{K_m-1} \frac{2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}^2\rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ &\quad + \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)} \end{aligned}$$

495 We recall (35) where we have summed on both sides from  $k = 0$  to  $k = K_m - 1$ :

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(K_m)}) - V(\hat{\mathbf{s}}^{(0)})] \\ &\leq \sum_{k=0}^{K_m-1} \left\{ \gamma_{k+1}(-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] + \gamma^2 L_V \rho^2 L_s^2 \Delta^{(k)} \right\} \\ &\quad + \sum_{k=0}^{K_m-1} \left\{ \tilde{\xi}^{(k+1)} + \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right\} \\ &\leq \sum_{k=0}^{K_m-1} \left\{ \left[ -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_s^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right] \mathbb{E}[\|\mathbf{h}_k\|^2] \right\} \\ &\quad + \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \tag{37}$$

where

$$\Xi^{(k+1)} = \tilde{\xi}^{(k+1)} + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}$$

and

$$\Gamma^{(k+1)} = \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2 (1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}}$$

496 We now analyse the following quantity

$$\begin{aligned}
& -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \\
& = \gamma_{k+1} \left[ -(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right]
\end{aligned} \tag{38}$$

497 Furthermore, we recall that  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_S, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k}$ ,  
498  $\beta = \frac{1}{\alpha n}$ ,  $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ . Then,

$$\begin{aligned}
& \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\frac{1}{n} - \gamma_{k+1}\beta - \gamma_{k+1}^2\rho^2 L_S^2} \\
& \leq \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L}(k\alpha^2 c_1^2 n^{4/3})^{-1} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}}\right)}{\frac{1}{n} - \frac{1}{k\alpha n} - \frac{1}{k^2 \alpha^2 c_1^2 n^{4/3}}} \\
& = \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}}\right)}{(k\alpha c_1 n^{1/3})(k\alpha - 1)c_1 - 1} \\
& \stackrel{(a)}{\leq} \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k\alpha c_1^2 \bar{L} n^{1/3}} \left(\frac{2}{k\alpha n} + 1\right)}{2(\alpha c_1 n^{1/3}) - 1} \\
& \leq \frac{1}{k^2 \alpha c_1^2 \bar{L} n^{4/3}} + \frac{1}{4k\alpha^2 c_1^3 \bar{L} n^{2/3}} \\
& \leq \frac{3/4}{\alpha c_1^2 \bar{L} n^{2/3}}
\end{aligned} \tag{39}$$

where (a) is due to  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$  and  $k\alpha c_1 n^{1/3} \geq 1$ . Note also that

$$-(v_{\min}\rho + v_{\max}^2) \leq -\rho v_{\min} = -\frac{1}{\alpha c_1^2 \bar{L} n^{2/3}}$$

which yields that

$$\left[ -(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right] \leq -\frac{1/4}{\alpha c_1^2 \bar{L} n^{2/3}}$$

499 Using the Lemma 2, we know that  $v_{\max}^2 \|\nabla V(\hat{s}^{(k)})\|^2 \leq \|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2$  and using (39) on (37)  
500 yields:

$$\begin{aligned}
v_{\max}^2 \sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] \\
& \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned}$$

501 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned}
\sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] \\
& \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned}$$

502

□

## F Practical Implementations of Two-Timescale EM Methods

### F.1 Application on GMM

#### F.1.1 Explicit Updates

We first recognize that the constraint set for  $\theta$  is given by

$$\Theta = \Delta^M \times \mathbb{R}^M.$$

Using the partition of the sufficient statistics as  $S(y_i, z_i) = (S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ , the partition  $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$  and the fact that  $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$ , the complete data log-likelihood can be expressed as in (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i)y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (40)$$

and  $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$ . We also define for each  $m \in \llbracket 1, M \rrbracket$ ,  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ . Consider the following latent sample used to compute an approximation of the conditional expected value  $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}}|y = y_i]$ :

$$z_{i,m} \sim \mathbb{P}(z_i = m | y_i; \theta) \quad (41)$$

where  $m \in \llbracket 1, M \rrbracket$ ,  $i \in [n]$  and  $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$ .

In particular, given iteration  $k + 1$ , the computation of the approximated quantity  $\tilde{S}_{i_k}^{(k)}$  during Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left( \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:=\tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1})y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})y_{i_k}}_{:=\tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:=\tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (42)$$

Recall that we have used the following regularizer:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (43)$$

It can be shown that the regularized M-step evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\boldsymbol{\mu}}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (44)$$

where we have defined for all  $m \in \llbracket 1, M \rrbracket$  and  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ .

#### F.1.2 Model Assumptions (GMM example)

We use the GMM example to illustrate the required assumptions.

Many practical models can satisfy the compactness of the sets as in Assumption A1. For instance, the GMM example satisfies (14) as the sufficient statistics are composed of indicator functions and observations as defined Section F.1 Equation (40).

Assumptions A2 and A3 are standard for the curved exponential family models. For GMM, the following (strongly convex) regularization  $\mathbf{r}(\theta)$  ensures A3:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m)$$



526 since it ensures  $\theta^{(k)}$  is unique and lies in  $\text{int}(\Delta^M) \times \mathbb{R}^M$ . We remark that for A2, it is possible to  
 527 define the Lipschitz constant  $L_p$  independently for each data  $y_i$  to yield a refined characterization.

528 Again, A4 is satisfied by practical models. For GMM, it can be verified by deriving the closed form  
 529 expression for  $B(s)$  and using A1.

530 Under A1 and A3, we have  $\|\hat{s}^{(k)}\| < \infty$  since  $S$  is compact and  $\hat{\theta}^{(k)} \in \text{int}(\Theta)$  for any  $k \geq 0$  which  
 531 thus ensure that the EM methods operate in a closed set throughout the optimization process.

### 532 F.1.3 Algorithms updates

533 In the sequel, recall that, for all  $i \in [n]$  and iteration  $k$ , the computed statistic  $\tilde{S}_{i_k}^{(k)}$  is defined by (42).  
 534 At iteration  $k$ , the several E-steps defined by (9) or (10) and (11) leads to the definition of the quantity  
 535  $\hat{s}^{(k+1)}$ . For the GMM example, after the initialization of the quantity  $\hat{s}^{(0)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(0)}$ , those  
 536 E-steps break down as follows:

537 **Batch EM (EM):** for all  $i \in [n]$ , compute  $\bar{s}_i^{(k)}$  and set

$$\hat{s}^{(k+1)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(k)}.$$

538 where  $\bar{s}_i^{(k)}$  are computed using the exact conditional expected value  $\mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i]$ :

$$\tilde{\omega}_m(y_i; \theta) := \mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)},$$

539 **Incremental EM (iEM):** draw an index  $i_k$  uniformly at random on  $[n]$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}.$$

540 **batch SAEM (SAEM):** draw an index  $i_k$  uniformly at random on  $[n]$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}.$$

541 where  $= \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$  with  $\tilde{S}_i^{(k)}$  defined in (42).

542 **Incremental SAEM (iSAEM):** draw an index  $i_k$  uniformly at random on  $[n]$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_i^k)})).$$

543 **Variance Reduced Two-Timescale EM (vrTTEM):** draw an index  $i_k$  uniformly at random on  $[n]$ ,  
 544 compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}))).$$

545 **Fast Incremental Two-Timescale EM (fiTTEM):** draw an index  $i_k$  uniformly at random on  $[n]$ ,  
 546 compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}))).$$

547 Finally, the  $k$ -th update reads  $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$  where the function  $s \rightarrow \bar{\theta}(s)$  is defined by (44).

## 548 F.2 Deformable Template Model for Image Analysis

### 549 F.2.1 Model and Updates

550 The complete model belongs to the curved exponential family, see [1], which vector of sufficient  
 551 statistics  $S = (S_1(z), S_2(z), S_3(z))$  read:

$$\begin{aligned} S_1(z) &= \frac{1}{n} \sum_{i=1}^n S_1(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top y_i \\ S_2(z) &= \frac{1}{n} \sum_{i=1}^n S_2(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \frac{1}{n} \sum_{i=1}^n S_3(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (45)$$

552 where for any pixel  $u \in \mathbb{R}^2$  and  $j \in \llbracket 1, k_g \rrbracket$  we noted:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j)$$

553 Finally, the Two-Timescale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z) \hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n} \hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z) \beta(\hat{s}) - 2\beta(\hat{s}) \hat{s}_1(z) \end{pmatrix} \quad (46)$$

554 where  $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$  is the vector of statistics obtained via the SA-step (7) and using the  
 555 MC approximation of the sufficient statistics  $(S_1(z), S_2(z), S_3(z))$  defined in (45).

### 556 F.2.2 Numerical Applications

557 For the inference of the template, we use the Matlab code (online SAEM) used in [17] and implement  
 558 our own batch, incremental, Variance reduced and Fast Incremental variants. The hyperparameters  
 559 are kept the same and reads as follows  $M = 400$ ,  $\gamma_k = 1/k^{0.6}$  and  $p = 16$ . The number of  
 560 landmarks for the template is  $k_p = 15$  points and for the deformation  $k_g = 6$  points. Both have  
 561 Gaussian kernels with respectively standard deviation of 0.08 and 0.16. The standard deviation of  
 562 the measurement errors is set to 0.1.

563 For the simulation part, we use the Carlin and Chib MCMC procedure, see [6]. Refer to [17] for  
 564 more details.

## 565 G Additional Experiment: Pharmacokinetics (PK) Model with Absorption 566 Lag Time

567 This numerical example was conducted in order to characterize the pharmacokinetics (PK) of orally  
 568 administered drug to simulated patients, using a population pharmacokinetics approach.  $M = 50$   
 569 synthetic datasets were generated for  $n = 5000$  patients with 10 observations (concentration mea-  
 570 sures) per patient. The goal is to model the evolution of the concentration of the absorbed drug  
 571 using a nonlinear and latent variable model.

572 **Model and Explicit Updates:** We consider a one-compartment PK model for oral administration  
 573 with an absorption lag-time ( $T_i^{\text{lag}}$ ), assuming first-order absorption and linear elimination processes.  
 574 The final model includes the following variables:  $ka$  the absorption rate constant,  $V$  the volume of  
 575 distribution,  $k$  the elimination rate constant and  $T_i^{\text{lag}}$  the absorption lag-time. We also add several  
 576 covariates to our model such as  $D$  the dose of drug administered,  $t$  the time at which measures  
 577 are taken and the weight of the patient influencing the volume  $V$ . More precisely, the log-volume  
 578  $\log(V)$  is a linear function of the log-weight  $\log(wt/70) = \log(wt/70)$ . Let  $z_i = (T_i^{\text{lag}}, ka_i, V_i, k_i)$  be the  
 579 vector of individual PK parameters, different for each individual  $i$ . The final model reads:

$$y_{ij} = f(t_{ij}, z_i) + \varepsilon_{ij} \quad \text{where} \quad f(t_{ij}, z_i) = \frac{D ka_i}{V(ka_i - k_i)} (e^{-ka_i(t_{ij} - T_i^{\text{lag}})} - e^{-k_i(t_{ij} - T_i^{\text{lag}})}) \quad (47)$$

where  $y_{ij}$  is the  $j$ -th concentration measurement of the drug of dosage  $D$  injected at time  $t_{ij}$  for patient  $i$ . We assume in this example that the residual errors  $\varepsilon_{ij}$  are independent and normally distributed with mean 0 and variance  $\sigma^2$ . Lognormal distributions are used for the four PK parameters.

Lognormal distributions are used for the four PK parameters:

$$\log(T_i^{\text{lag}}) \sim \mathcal{N}(\log(T_{\text{pop}}^{\text{lag}}), \omega_{T^{\text{lag}}}^2), \log(ka_i) \sim \mathcal{N}(\log(ka_{\text{pop}}), \omega_{ka}^2), \quad (48)$$

$$\log(V_i) \sim \mathcal{N}(\log(V_{\text{pop}}), \omega_V^2), \log(k_i) \sim \mathcal{N}(\log(k_{\text{pop}}), \omega_k^2). \quad (49)$$

We recall that the complete model  $(y, z)$  defined by (47) belongs to the curved exponential family, which vector of sufficient statistics  $S = (S_1(z), S_2(z), S_3(z))$  read:

$$S_1(z) = \frac{1}{n} \sum_{i=1}^n z_i, \quad S_2(z) = \frac{1}{n} \sum_{i=1}^n z_i^\top z_i, \quad S_3(z) = \frac{1}{n} \sum_{i=1}^n (y_i - f(t_i, z_i))^2 \quad (50)$$

where we have noted  $y_i$  and  $t_i$  the vector of observations and time for each patient  $i$ . At iteration  $k$ , and setting the number of MC samples to 1 for the sake of clarity, the MC sampling  $z_i^{(k)} \sim p(z_i|y_i, \theta^{(k)})$  is performed using a Metropolis-Hastings procedure detailed in Algorithm 2. The quantities  $\hat{S}^{(k+1)}$  and  $\hat{s}^{(k+1)}$  are then updated according to the different methods. Finally the maximization step yields:

$$\bar{\theta}(s) = \begin{pmatrix} \hat{s}_1^{(k+1)} \\ \hat{s}_2^{(k+1)} - \hat{s}_1^{(k+1)} (\hat{s}_1^{(k+1)})^\top \\ \hat{s}_3^{(k+1)} \end{pmatrix} = \begin{pmatrix} \overline{z_{\text{pop}}}(\hat{s}^{(k+1)}) \\ \overline{\omega_z}(\hat{s}^{(k+1)}) \\ \overline{\sigma}(\hat{s}^{(k+1)}) \end{pmatrix}. \quad (51)$$

**Metropolis Hastings algorithm** During the simulation step of the MISSO method, the sampling from the target distribution  $\pi(z_i, \theta) := p(z_i|y_i, \theta)$  is performed using a Metropolis Hastings (MH) algorithm [20] with proposal distribution  $q(z_i, \delta)$  where  $\theta = (z_{\text{pop}}, \omega_z)$  and  $\delta$  is the vector of parameters of the proposal distribution. Commonly they parameterize a Gaussian proposal. The MH algorithm is summarized in 2.

---

**Algorithm 2** MH algorithm

---

```

1: Input: initialization  $z_{i,0} \sim q(z_i; \delta)$ 
2: for  $m = 1, \dots, M$  do
3:   Sample  $z_{i,m} \sim q(z_i; \delta)$ 
4:   Sample  $u \sim \mathcal{U}([0, 1])$ 
5:   Calculate the ratio  $r = \frac{\pi(z_{i,m}; \theta) / q(z_{i,m}; \delta)}{\pi(z_{i,m-1}; \theta) / q(z_{i,m-1}; \delta)}$ 
6:   if  $u < r$  then
7:     Accept  $z_{i,m}$ 
8:   else
9:      $z_{i,m} \leftarrow z_{i,m-1}$ 
10:  end if
11: end for
12: Output:  $z_{i,M}$ 

```

---

**Monte Carlo study:** We conduct a Monte Carlo study to showcase the benefits of our scheme.  $M = 50$  datasets have been simulated using the following PK parameters values:  $T_{\text{pop}}^{\text{lag}} = 1$ ,  $ka_{\text{pop}} = 1$ ,  $V_{\text{pop}} = 8$ ,  $k_{\text{pop}} = 0.1$ ,  $\omega_{T^{\text{lag}}} = 0.4$ ,  $\omega_{ka} = 0.5$ ,  $\omega_V = 0.2$ ,  $\omega_k = 0.3$  and  $\sigma^2 = 0.5$ . We define the mean square distance over the  $M$  replicates  $E_k(\ell) = \frac{1}{M} \sum_{m=1}^M \left( \theta_k^{(m)}(\ell) - \theta^* \right)^2$  and plot it against the epochs (passes over the data) Figure 4. Note that the MC-step (5) is performed using a Metropolis Hastings procedure since the posterior distribution under the model  $\theta$  noted  $p(z_i|y_i, \theta)$  is intractable due to the nonlinearity of the model (47). Figure 4 shows clear advantage of variance reduced methods (vrTTEM and fitTEM) avoiding the twists and turns displayed by the incremental and the batch methods.




Figure 4: Precision  $|ka^{(k)} - ka^*|^2$  per epoch