

363 A Proof of Lemma 2

364 **Lemma.** Assume A3,A4. For all $\mathbf{s} \in \mathcal{S}$,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (16)$$

365 **Proof** Using A3 and the fact that we can exchange integration with differentiation and the Fisher's
366 identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (17)$$

367 Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \mathbf{s}.$$

368 Taking the gradient of the above map w.r.t. \mathbf{s} and using assumption A3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left(\nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})} \Big|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}).$$

369 The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})))$$

370 where we recall $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left(\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top}$. The proof of (16) follows directly
371 from the assumption A4. \square

372 B Proof of Theorem 1

373 Beforehand, We present two intermediary Lemmas important for the analysis of the incremental
374 update of the iSAEM algorithm. The first one gives a characterization of the quantity $\mathbb{E}[\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}]$:
375

376 **Lemma.** Assume A1. The update (1) is equivalent to the following update on the resulting statistics
377

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)})$$

378 Also:

$$\mathbb{E}[\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E}[\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E}[\eta_{i_k}^{(k+1)}]$$

379 where $\bar{\mathbf{s}}^{(k)}$ is defined by (3) and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$.

380 **Proof** From update (1), we have:

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left(\tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_i^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned}$$

381 Since $\tilde{S}_{i_k}^{(k+1)} = \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k+1)}) + \eta_{i_k}^{(k+1)}$ we have

$$\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_i^k)} - \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)}$$

382 Taking the full expectation of both side of the equation leads to:

$$\begin{aligned}\mathbb{E}[\tilde{S}^{(k+1)} - \hat{s}^{(k)}] &= \mathbb{E}[\bar{s}^{(k)} - \hat{s}^{(k)}] + \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right] \\ &\quad - \frac{1}{n} \mathbb{E}[\mathbb{E}[\tilde{S}_i^{(\tau_i^k)} - \bar{s}_{i_k}(\theta^{(k)}) | \mathcal{F}_k]] + \frac{1}{n} \mathbb{E}[\eta_{i_k}^{(k+1)}]\end{aligned}$$

383 The following equalities:

$$\mathbb{E}[\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E}[\bar{s}_{i_k}(\theta^{(k)}) | \mathcal{F}_k] = \bar{s}^{(k)}$$

384 concludes the proof of the Lemma. \square

385 And the following auxiliary Lemma setting an upper bound for the quantity $\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2]$

386 **Lemma 7.** For any $k \geq 0$ and consider the iSAEM update in (1), it holds that

$$\begin{aligned}\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2] &\leq 4\mathbb{E}[\|\bar{s}^{(k)} - \hat{s}^{(k)}\|^2] + \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(t_i^k)}\|^2] \\ &\quad + 2\frac{c_\eta}{M_k} + 4\mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right\|^2\right]\end{aligned}$$

387 **Proof** Applying the iSAEM update yields:

$$\begin{aligned}\mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2] &= \mathbb{E}[\|\tilde{S}^{(k)} - \hat{s}^{(k)} - \frac{1}{n}(\tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(t_i^k)})\|^2] \\ &\leq 4\mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right\|^2\right] + 4\mathbb{E}[\|\bar{s}^{(k)} - \hat{s}^{(k)}\|^2] \\ &\quad + \frac{2}{n^2} \mathbb{E}[\|\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(t_i^k)}\|^2] + 2\frac{c_\eta}{M_k}\end{aligned}$$

388 The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E}[\|\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(t_i^k)}\|^2] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\bar{s}_i^{(k)} - \bar{s}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(t_i^k)}\|^2],$$

389 where (a) is due to Lemma 1 and which concludes the proof of the Lemma.

390 \square

391 **Theorem.** Assume A1-A5. Consider the iSAEM sequence $\{\hat{s}^{(k)}\}_{k \geq 0} \in \mathcal{S}$ obtained with $\rho_{k+1} = 1$
392 for any $k \leq K_m$ where K_m is a positive integer. Let $\{\gamma_k = 1/(k^a \alpha c_1 \bar{L})\}_{k \geq 0}$, where $a \in (0, 1)$, be a
393 sequence of stepsizes, $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\beta = c_1 \bar{L}/n$. Then:

$$v_{\max}^{-2} \sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] + \sum_{k=0}^{K_m-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2].$$

394 **Proof** Under the smoothness of the Lyapunov function V (cf. Lemma 1), we can write:

$$V(\hat{s}^{(k+1)}) \leq V(\hat{s}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} | \nabla V(\hat{s}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2$$

395 Taking the expectation on both sides yields:

$$\mathbb{E}[V(\hat{s}^{(k+1)})] \leq \mathbb{E}[V(\hat{s}^{(k)})] + \gamma_{k+1} \mathbb{E}[\langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} | \nabla V(\hat{s}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{s}^{(k)}\|^2]$$

396 Using Lemma 3, we obtain:

$$\begin{aligned}
& \mathbb{E} \left[\langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&= \mathbb{E} \left[\langle \bar{s}^{(k)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&\stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
&\stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[\left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&\stackrel{(a)}{\leq} \left(v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned}$$

397 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with $\beta \rightarrow 1$). Note

398 $a_k = \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$ and

$$\begin{aligned}
a_k \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] &\leq \mathbb{E} \left[V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (18)
\end{aligned}$$

399 We now give an upper bound of $\mathbb{E} \left[\left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right]$ using Lemma 7 and plug it into (18):

$$\begin{aligned}
(a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[\left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] &\leq \mathbb{E} \left[V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] \\
&\quad + \gamma_{k+1} \left(\frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
&\quad + \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&\quad + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \quad (19)
\end{aligned}$$

400 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{n-1}{n} \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \right)$$

401 where the equality holds as i_k and j_k are drawn independently. For any $\beta > 0$, it holds

$$\begin{aligned}
& \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \\
&= \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 + 2 \langle \hat{s}^{(k+1)} - \hat{s}^{(k)} \mid \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \rangle \right] \\
&= \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 - 2\gamma_{k+1} \langle \hat{s}^{(k)} - \tilde{S}^{(k+1)} \mid \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \rangle \right] \\
&\leq \mathbb{E} \left[\left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 + \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 + \frac{\gamma_{k+1}}{\beta} \left\| \hat{s}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 + \gamma_{k+1} \beta \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right]
\end{aligned}$$

402 where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ & \leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2\right] \end{aligned}$$

403 Observe that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)})$. Applying Lemma 7 yields

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ & \leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\ & \leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\ & \quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\ & \quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \end{aligned}$$

404 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2]$$

405 From the above, we get

$$\begin{aligned} \Delta^{(k+1)} & \leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})\right) \Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ & \quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \end{aligned}$$

406 Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$,

407 $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$, $\alpha \geq 8$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1}$$

408 which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} =$

409 $\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}\mathbf{L}_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} & \leq 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_\ell}^{(\ell)}\|^2\right] \\ & \quad + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)}\right\|^2\right] \end{aligned}$$

410 Note $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ Summing on both sides over $k = 0$ to $k = K_m - 1$ yields:

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} \Delta^{(k+1)} \\
&= 4 \sum_{k=0}^{K_m-1} \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} [\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_m-1} \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[\left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} 4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&\leq \sum_{k=0}^{K_m-1} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} [\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_m-1} \frac{2 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[\left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right]
\end{aligned} \tag{20}$$

411 We recall (19) where we have summed on both sides from $k = 0$ to $k = K_m - 1$:

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\
&+ \sum_{k=0}^{K_m-1} \gamma_{k+1} \left(\frac{1}{2\beta} \left(1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)}
\end{aligned} \tag{21}$$

412 Plugging (20) into (21) results in:

$$\begin{aligned}
& \sum_{k=0}^{K_m-1} \tilde{\alpha}_k \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \sum_{k=0}^{K_m-1} \tilde{\beta}_k \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&\leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_m-1} \tilde{\Gamma}_k \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned}$$

413 where

$$\begin{aligned}
\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\beta}_k &= \gamma_{k+1} \left(\frac{1}{2\beta} \left(1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\Gamma}_k &= \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}}
\end{aligned}$$

414 and

$$\begin{aligned}
a_k &= \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right) \\
\Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \\
c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{L_s, L_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}
\end{aligned}$$

415 When, for any $k > 0$, $\tilde{\alpha}_k \geq 0$, we have by Lemma 2 that:

$$\sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_m} \tilde{\alpha}_k \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$$

416 which yields an upper bound of the gradient of the Lyapunov function V along the path of the
417 iSAEM update and concludes the proof of the Theorem. \square

418 C Proofs of Auxiliary Lemmas

419 C.1 Proof of Lemma 4 and Lemma 5

420 **Lemma.** For any $k \geq 0$ and consider the vrTTEM update in (2) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned}
\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\
&\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned}$$

421 where we recall that $\ell(k)$ is the first iteration number in the epoch that iteration k is in.

422 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
423 out this proof:

$$\begin{aligned}
\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathcal{S}^{(k+1)}) \\
&= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} \right] \right)
\end{aligned} \tag{22}$$

424 We observe, using the identity (22), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] \tag{23}$$

425 For the latter term, we obtain its upper bound as

$$\begin{aligned}
\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)}\|^2] &= \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{\ell(k)}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{\ell(k)}) \right\|^2 \right] \\
&\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \stackrel{(b)}{\leq} L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned}$$

426 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (23) proves the
427 lemma. \square

428 **Lemma.** For any $k \geq 0$ and consider the fitTEM update in (3) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned}
\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(i_k)}\|^2] \\
&\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned}$$

429 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
 430 out this proof:

$$\begin{aligned}
 \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) \\
 &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1 - \rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\
 &= -\gamma_{k+1}\left((1 - \rho)\left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\right] + \rho\left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\right]\right) \\
 &= -\gamma_{k+1}\left((1 - \rho)\left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\right] + \rho\left[\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{S}}^{(k)} - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right]\right)
 \end{aligned} \tag{24}$$

431 We observe, using the identity (24), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{S}}^{(k)}\|^2] + 2\rho^2\mathbb{E}[\|\bar{\mathbf{S}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1 - \rho)^2\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \tag{25}$$

432 For the latter term, we obtain its upper bound as

$$\begin{aligned}
 \mathbb{E}[\|\bar{\mathbf{S}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E}\left[\left\|\frac{1}{n}\sum_{i=1}^n(\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{S}}_i^{(k)}) - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right\|^2\right] \\
 &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
 \end{aligned}$$

433 where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] = \frac{1}{n}\sum_{i=1}^n\mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_s^2}{n}\sum_{i=1}^n\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

434 Substituting into (25) proves the lemma. \square

435 C.2 Proof of Lemma 6

436 **Lemma.** Consider a decreasing stepsize $\gamma_k \in (0, 1)$ and a constant ρ , then the following inequality
 437 holds:

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \leq \frac{\rho}{1 - \rho}\sum_{\ell=0}^k(1 - \gamma_\ell)^2(\mathbf{S}^{(\ell)} - \tilde{S}^{(\ell)})$$

438 where $\mathbf{S}^{(k)}$ is defined either by (3) (fiTTEM) or (2) (vrTTEM)

439 **Proof** We begin by writing the two-timescale update:

$$\begin{aligned}
 \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho(\mathbf{S}^{(k+1)} - \tilde{S}^{(k)}) \\
 \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)})
 \end{aligned} \tag{26}$$

440 where $\mathbf{S}^{(k+1)} = \frac{1}{n}\sum_{i=1}^n\tilde{S}_i^{(t_i^k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$ according to (3). Denote $\delta^{(k+1)} = \hat{\mathbf{s}}^{(k+1)} - \tilde{S}^{(k+1)}$.
 441 Then from (26), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1 - \rho}(1 - \gamma_{k+1})(\mathbf{S}^{(k+1)} - \tilde{S}^{(k+1)})$$

442 Using the telescoping sum and noting that $\delta^{(0)} = 0$, we have

$$\delta^{(k+1)} \leq \frac{\rho}{1 - \rho}\sum_{\ell=0}^k(1 - \gamma_{\ell+1})^2(\mathbf{S}^{(\ell+1)} - \tilde{S}^{(\ell+1)})$$

443 \square

444 **C.3 Additional Intermediary Result**

445 **Lemma 8.** *At iteration $k + 1$, the drift term of update (3), with $\rho_{k+1} = \rho$, is equivalent to the*
 446 *following :*

$$\begin{aligned} \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) \left(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)} \right) \end{aligned}$$

447 *where we recall that $\eta_{i_k}^{(k+1)}$, defined in (12), which is the gap between the MC approximation and*
 448 *the expected statistics.*

449 **Proof** Using the fitTEM update $\tilde{\mathbf{S}}^{(k+1)} = (1 - \rho)\tilde{\mathbf{S}}^{(k)} + \rho\mathcal{S}^{(k+1)}$ where $\mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} -$
 450 $\tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})$ leads to the following decomposition:

$$\begin{aligned} &\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \\ &= (1 - \rho)\tilde{\mathbf{S}}^{(k)} + \rho \left(\bar{\mathcal{S}}^{(k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}) \right) - \hat{\mathbf{s}}^{(k)} + \rho\bar{\mathbf{s}}^{(k)} - \rho\bar{\mathbf{s}}^{(k)} \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho(\tilde{\mathbf{S}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(k)}) + (1 - \rho) \left(\tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) + \rho \left(\bar{\mathcal{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} + (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}) \right) \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) \left(\tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) \end{aligned}$$

451 *where we observe that $\mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}] = \bar{\mathbf{s}}^{(k)} - \bar{\mathcal{S}}^{(k)}$ and which concludes the proof.*

452 *Important Note:* Note that $\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}$ is not equal to $\eta_{i_k}^{(k+1)}$, defined in (12), which is the gap
 453 *between the MC approximation and the expected statistics. Indeed $\tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}$ is not computed under the*
 454 *same model as $\bar{\mathbf{s}}_{i_k}^{(k)}$.* □

455 D Proof of Theorem 2

456 **Theorem.** Assume A1-A5. Consider the vrTTEM sequence $\{\hat{\mathbf{s}}^{(k)}\}_{k>0} \in \mathcal{S}$ for any $k \leq K_m$ where
 457 K_m is a positive integer. Let $\{\gamma_{k+1} = 1/(k^a \bar{L})\}_{k>0}$, where $a \in (0, 1)$, be a sequence of stepsizes,
 458 $\bar{L} = \max\{L_s, L_V\}$, $\rho = \mu/(c_1 \bar{L} n^{2/3})$, $m = nc_1^2/(2\mu^2 + \mu c_1^2)$ and a constant $\mu \in (0, 1)$. Then:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu P_m v_{\min}^2 v_{\max}^2} \left(\mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right).$$

459 **Proof** Using the smoothness of V and update (2), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (27)$$

460 Denote $H_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fitTEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.
 461 Taking expectations on both sides show that

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}\rho \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|H_{k+1}\|^2] \\ &\stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\langle \eta_{i_k}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|H_{k+1}\|^2] \\ &\stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|H_{k+1}\|^2] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \gamma_{k+1}(1-\rho) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \quad (28)$$

462 where we have used (22) in (a) and $\mathbb{E}[\mathcal{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$ in (b), the growth condition in
 463 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

464 Furthermore, for $k+1 \leq \ell(k) + m$ (i.e., $k+1$ is in the same epoch as k), we have

$$\begin{aligned} &\mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|H_{k+1}\|^2 \\ &\quad - 2\gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1-\rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle] \\ &\leq \mathbb{E}[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|H_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2], \end{aligned}$$

465 where we first used (22) and the last inequality is due to the Young's inequality.

466 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2]$$

where $b_k := \bar{b}_{k \bmod m}$ is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2) + \gamma_{k+1}^2\rho^2 L_V L_s^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0.$$

Note that \bar{b}_i is decreasing with i and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_s^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2}, \quad i = 1, 2, \dots, m.$$

For $k+1 \leq \ell(k) + m$, we have the following inequality

$$\begin{aligned} R_{k+1} &\leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \right] \\ &\quad + \gamma_{k+1} \mathbb{E} \left[\rho \left\| \eta_{i_k}^{(k+1)} \right\|^2 - (1-\rho) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[\frac{\gamma_{k+1}\rho}{\beta} \left\| \eta_{i_k}^{(k+1)} \right\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

And using Lemma 4 we obtain:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2 - \gamma_{k+1}^2\rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2\rho^2 L_V L_s^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2 \right) \|\mathbf{h}_k\|^2 \right] \\ &\quad + \gamma_{k+1} \mathbb{E} \left[(\rho + \rho^2 \gamma_{k+1} L_V) \left\| \eta_{i_k}^{(k+1)} \right\|^2 - (1-\rho - (1-\rho)^2 \gamma_{k+1} L_V) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \\ &\quad + b_{k+1} \mathbb{E} \left[\left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2 \right) \left\| \eta_{i_k}^{(k+1)} \right\|^2 + \left(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2 \right) \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

Rearranging the terms yields:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\quad + \underbrace{\left(b_{k+1}(1 + \gamma\beta + 2\gamma^2\rho^2 L_s^2) + \gamma^2\rho^2 L_V L_s^2 \right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)} \end{aligned}$$

where

$$\begin{aligned} \tilde{\eta}^{(k+1)} &= \left(\gamma_{k+1}(\rho + \rho^2 \gamma_{k+1} L_V) + b_{k+1}(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2) \right) \mathbb{E} \left[\left\| \eta_{i_k}^{(k+1)} \right\|^2 \right] \\ \chi^{(k+1)} &= \left(b_{k+1}(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2) - \gamma_{k+1}(1-\rho - (1-\rho)^2 \gamma_{k+1} L_V) \right) \\ \tilde{\chi}^{(k+1)} &= \chi^{(k+1)} \mathbb{E} \left[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \end{aligned}$$

This leads, using Lemma 2, that for any γ_{k+1} , ρ and β such that $\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2) > 0$,

$$\begin{aligned} v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \\ &\quad + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \end{aligned}$$

We first remark that

$$\begin{aligned} &\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \\ &\geq \frac{\gamma_{k+1}\rho}{c_1} (1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1)) \end{aligned}$$

476 where $c_1 = v_{\min}^{-1}$. By setting $\bar{L} = \max\{L_s, L_V\}$, $\beta = \frac{c_1 \bar{L}}{n^{1/3}}$, $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$ and
 477 $\{\gamma_{k+1}\}$ any sequence of decreasing stepsizes in $(0, 1)$, it can be shown that there exists $\mu \in (0, 1)$,
 478 such that the following lower bound holds

$$\begin{aligned}
 & 1 - \gamma_{k+1} c_1 \rho L_V - b_{k+1} \left(\frac{c_1}{\beta} + 2\gamma_{k+1} \rho c_1 \right) \\
 & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0 \left(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
 & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V \mu^2}{c_1^2 n^{\frac{4}{3}}} \frac{(1 + \gamma\beta + 2\gamma^2 L_s^2)^m - 1}{\gamma\beta + 2\gamma^2 L_s^2} \left(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
 & \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2} (e - 1) \left(1 + \frac{2\mu}{n} \right) \geq 1 - \mu - \mu(1 + 2\mu) \frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2}
 \end{aligned}$$

479 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2 L_s^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2 n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \quad \text{and} \quad (1 + \gamma\beta + 2\gamma^2 L_s^2)^m \leq e - 1.$$

480 and the required μ in (b) can be found by solving the quadratic equation.

481 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_m})}{v_{\min} \rho} + 2 \sum_{k=0}^{K_m-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min} \rho}$$

482 Note that $R_0 = \mathbb{E}[V(\hat{s}^{(0)})]$ and if K_m is a multiple of m , then $R_{\max} = \mathbb{E}[V(\hat{s}^{(K_m)})]$. Under the latter
 483 condition, we have

$$\sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] + \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_m-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}]$$

484 This concludes our proof.

485

□

486 E Proof of Theorem 3

487 **Theorem.** Assume A1-A5. Consider the fitTEM sequence $\{\hat{\mathbf{s}}^{(k)}\}_{k>0} \in \mathcal{S}$ for any $k \leq K_m$ where
 488 K_m be a positive integer. Let $\{\gamma_{k+1} = 1/(k^a \alpha c_1 \bar{L})\}_{k>0}$, where $a \in (0, 1)$, be a sequence of
 489 positive stepsizes, $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\beta = 1/(\alpha n)$, $\rho = 1/(\alpha c_1 \bar{L} n^{2/3})$
 490 and $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$. Then:

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] \leq \frac{4\alpha \bar{L} n^{2/3}}{P_m v_{\min}^2 v_{\max}^2} \left(\mathbb{E}[\Delta V] + \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \Gamma^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right).$$

491 **Proof** Using the smoothness of V and update (3), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (29)$$

492 Denote $H_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fitTEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.
 493 Using Lemma 8 and the additional following identity:

$$\mathbb{E} \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] = 0 \quad (30)$$

494 we have:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \mathbb{E} \left[\langle \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1 - \rho) \mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(a)}{\leq} -v_{\min} \gamma_{k+1} \rho \mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1} \mathbb{E} \left[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(b)}{\leq} - (v_{\min} \gamma_{k+1} \rho + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \end{aligned}$$

495 where $\xi^{(k+1)} = \mathbb{E}[\|\mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k]\|^2]$. **Bounding** $\mathbb{E}[\|\mathbf{H}_{k+1}\|^2]$ Using Lemma 5, we obtain:

$$\begin{aligned} &\gamma_{k+1} (v_{\min} \rho + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)}) \right] + \tilde{\xi}^{(k+1)} + \left((1 - \rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad - \frac{\gamma_{k+1}^2 L_V \rho^2 L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (31)$$

496 where $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 L_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)}$. Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (32)$$

497 where the equality holds as i_k and j_k are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \end{aligned}$$

498 Note that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}) = -\gamma_{k+1}\mathbf{H}_{k+1}$ and that in expectation we recall
 499 that $\mathbb{E}[\mathbf{H}_{k+1}|\mathcal{F}_k] = \rho\mathbf{h}_k + \rho\mathbb{E}[\eta_{i_k}^{(k+1)}|\mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)}]$ where $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$. Thus,
 500 for any $\beta > 0$, it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \\ &\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\right. \\ &\quad \left.+ \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

501 where the last inequality is due to the Young's inequality. Plugging this into (32) yields:

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \\ &\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\right. \\ &\quad \left.+ \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

502 Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[\left(1 + \gamma_{k+1}\beta\right)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2\right. \\ &\quad \left.+ \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2]\right] \end{aligned}$$

503 We now use Lemma 5 on $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2$ and obtain:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2\rho^2\mathbf{L}_s^2}{n} + \frac{(n-1)(1+\gamma_{k+1}\beta)}{n^2}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2\left(2\gamma_{k+1} + \frac{1}{\beta}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2\mathbf{L}_s^2}{n}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2\left(2\gamma_{k+1} + \frac{1}{\beta}\right)\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

504 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

505 From the above, we get

$$\begin{aligned} \Delta^{(k+1)} &\leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

506 Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{2, 1+2v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\gamma_{k+1} = \frac{1}{k}$, $\beta = \frac{1}{\alpha n}$, $\rho = \frac{1}{\alpha c_1 L n^{2/3}}$,
507 $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{\frac{4}{3}}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1}$$

508 which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} = \frac{1}{n} -$
509 $\gamma_{k+1}\beta - \gamma_{k+1}^2\rho^2 L_s^2$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} &\leq \sum_{\ell=0}^k \omega_{k,\ell} \left(2\gamma_{\ell+1}^2\rho^2 + \frac{\gamma_{\ell+1}^2\rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] \\ &\quad + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left(2\gamma_{\ell+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\tilde{S}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)} \end{aligned}$$

510 where $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ and $\tilde{\epsilon}^{(\ell+1)} = \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]$.

511 Summing on both sides over $k = 0$ to $k = K_m - 1$ yields:

$$\begin{aligned} \sum_{k=0}^{K_m-1} \Delta^{(k+1)} &\leq \sum_{k=0}^{K_m-1} \frac{2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ &\quad + \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \sum_{k=0}^{K_m-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)} \end{aligned}$$

512 We recall (31) where we have summed on both sides from $k = 0$ to $k = K_m - 1$:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(K_m)}) - V(\hat{\mathbf{s}}^{(0)})] \\ &\leq \sum_{k=0}^{K_m-1} \left\{ \gamma_{k+1}(-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] + \gamma^2 L_V \rho^2 L_s^2 \Delta^{(k)} \right\} \\ &\quad + \sum_{k=0}^{K_m-1} \left\{ \tilde{\xi}^{(k+1)} + \left((1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right\} \\ &\leq \sum_{k=0}^{K_m-1} \left\{ \left[-\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_s^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right] \mathbb{E}[\|\mathbf{h}_k\|^2] \right\} \\ &\quad + \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \tag{33}$$

where

$$\Xi^{(k+1)} = \tilde{\xi}^{(k+1)} + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}$$

and

$$\Gamma^{(k+1)} = \left((1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2 (1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}}$$

513 We now analyse the following quantity

$$\begin{aligned}
& -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \\
& = \gamma_{k+1} \left[-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right]
\end{aligned} \tag{34}$$

514 Furthermore, we recall that $c_1 = v_{\min}^{-1}$, $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_S, L_V\}$, $\gamma_{k+1} = \frac{1}{k}$,
515 $\beta = \frac{1}{\alpha n}$, $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$. Then,

$$\begin{aligned}
& \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\frac{1}{n} - \gamma_{k+1}\beta - \gamma_{k+1}^2\rho^2 L_S^2} \\
& \leq \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L}(k\alpha^2 c_1^2 n^{4/3})^{-1} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}}\right)}{\frac{1}{n} - \frac{1}{k\alpha n} - \frac{1}{k^2 \alpha^2 c_1^2 n^{4/3}}} \\
& = \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}}\right)}{(k\alpha c_1 n^{1/3})(k\alpha - 1)c_1 - 1} \\
& \stackrel{(a)}{\leq} \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k\alpha c_1^2 \bar{L} n^{1/3}} \left(\frac{2}{k\alpha n} + 1\right)}{2(\alpha c_1 n^{1/3}) - 1} \\
& \leq \frac{1}{k^2 \alpha c_1^2 \bar{L} n^{4/3}} + \frac{1}{4k\alpha^2 c_1^3 \bar{L} n^{2/3}} \\
& \leq \frac{3/4}{\alpha c_1^2 \bar{L} n^{2/3}}
\end{aligned} \tag{35}$$

where (a) is due to $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ and $k\alpha c_1 n^{1/3} \geq 1$. Note also that

$$-(v_{\min}\rho + v_{\max}^2) \leq -\rho v_{\min} = -\frac{1}{\alpha c_1^2 \bar{L} n^{2/3}}$$

which yields that

$$\left[-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_S^2 \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \right] \leq -\frac{1/4}{\alpha c_1^2 \bar{L} n^{2/3}}$$

516 Using the Lemma 2, we know that $v_{\max}^2 \|\nabla V(\hat{s}^{(k)})\|^2 \leq \|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2$ and using (35) on (33)
517 yields:

$$\begin{aligned}
v_{\max}^2 \sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] \\
& \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned}$$

518 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned}
\sum_{k=0}^{K_m-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_m)})] \\
& \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_m-1} \Xi^{(k+1)} + \sum_{k=0}^{K_m-1} \Gamma^{(k+1)} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned}$$

519 □

520 F Practical Implementations of Two-Timescale EM Methods

521 F.1 Application on GMM

522 F.1.1 Explicit Updates

523 We first recognize that the constraint set for θ is given by

$$\Theta = \Delta^M \times \mathbb{R}^M.$$

524 Using the partition of the sufficient statistics as $S(y_i, z_i) =$
 525 $(S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$, the partition
 526 $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ and the fact that
 527 $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$, the complete data log-likelihood can be expressed as in
 528 (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i)y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (36)$$

529 and $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$. We also define for each $m \in \llbracket 1, M \rrbracket$, $j \in \llbracket 1, 3 \rrbracket$,
 530 $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$. Consider the following latent sample used to compute an approximation of
 531 the conditional expected value $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}}|y = y_i]$:

$$z_{i,m} \sim \mathbb{P}(z_i = m|y_i; \theta) \quad (37)$$

532 where $m \in \llbracket 1, M \rrbracket$, $i \in [n]$ and $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$.

533 In particular, given iteration $k + 1$, the computation of the approximated quantity $\tilde{S}_{i_k}^{(k)}$ during
 534 Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left(\underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:=\tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1})y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})y_{i_k}}_{:=\tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:=\tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (38)$$

535 Recall that we have used the following regularizer:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (39)$$

536 It can be shown that the regularized M-step evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\boldsymbol{\mu}}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (40)$$

537 where we have defined for all $m \in \llbracket 1, M \rrbracket$ and $j \in \llbracket 1, 3 \rrbracket$, $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$.

538 F.1.2 Model Assumptions (GMM example)

539 We use the GMM example to illustrate the required assumptions.

540 Many practical models can satisfy the compactness of the sets as in Assumption A1 For instance,
 541 the GMM example satisfies (11) as the sufficient statistics are composed of indicator functions and
 542 observations as defined Section F.1 Equation (36).

Assumptions A2 and A3 are standard for the curved exponential family models. For GMM, the following (strongly convex) regularization $\mathbf{r}(\theta)$ ensures A3:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m)$$

543 since it ensures $\theta^{(k)}$ is unique and lies in $\text{int}(\Delta^M) \times \mathbb{R}^M$. We remark that for A2, it is possible to
 544 define the Lipschitz constant L_p independently for each data y_i to yield a refined characterization.

545 Again, A4 is satisfied by practical models. For GMM, it can be verified by deriving the closed form
 546 expression for $B(s)$ and using A1.

547 Under A1 and A3, we have $\|\hat{s}^{(k)}\| < \infty$ since S is compact and $\hat{\theta}^{(k)} \in \text{int}(\Theta)$ for any $k \geq 0$ which
 548 thus ensure that the EM methods operate in a closed set throughout the optimization process.

549 F.1.3 Algorithms updates

550 In the sequel, recall that, for all $i \in [n]$ and iteration k , the computed statistic $\tilde{S}_{i_k}^{(k)}$ is defined by (38).
 551 At iteration k , the several E-steps defined by (1) or (2) and (3) leads to the definition of the quantity
 552 $\hat{s}^{(k+1)}$. For the GMM example, after the initialization of the quantity $\hat{s}^{(0)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(0)}$, those
 553 E-steps break down as follows:

554 **Batch EM (EM):** for all $i \in [n]$, compute $\bar{s}_i^{(k)}$ and set

$$\hat{s}^{(k+1)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(k)}.$$

555 where $\bar{s}_i^{(k)}$ are computed using the exact conditional expected value $\mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i]$:

$$\tilde{\omega}_m(y_i; \theta) := \mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)},$$

556 **Incremental EM (iEM):** draw an index i_k uniformly at random on $[n]$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}.$$

557 **batch SAEM (SAEM):** draw an index i_k uniformly at random on $[n]$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}.$$

558 where $= \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$ with $\tilde{S}_i^{(k)}$ defined in (38).

559 **Incremental SAEM (iSAEM):** draw an index i_k uniformly at random on $[n]$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \left(\tilde{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_i^k)}) \right).$$

560 **Variance Reduced Two-Timescale EM (vrTTEM):** draw an index i_k uniformly at random on $[n]$,
 561 compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}))).$$

562 **Fast Incremental Two-Timescale EM (fiTTEM):** draw an index i_k uniformly at random on $[n]$,
 563 compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}))).$$

564 Finally, the k -th update reads $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$ where the function $s \rightarrow \bar{\theta}(s)$ is defined by (40).

565 F.2 Deformable Template Model for Image Analysis

566 F.2.1 Model and Updates

567 The complete model belongs to the curved exponential family, see [1], which vector of sufficient
568 statistics $S = (S_1(z), S_2(z), S_3(z))$ read:

$$\begin{aligned} S_1(z) &= \frac{1}{n} \sum_{i=1}^n S_1(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top y_i \\ S_2(z) &= \frac{1}{n} \sum_{i=1}^n S_2(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \frac{1}{n} \sum_{i=1}^n S_3(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (41)$$

569 where for any pixel $u \in \mathbb{R}^2$ and $j \in \llbracket 1, k_g \rrbracket$ we denote:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j)$$

570 Finally, the Two-Timescale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z) \hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n} \hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z) \beta(\hat{s}) - 2\beta(\hat{s}) \hat{s}_1(z) \end{pmatrix} \quad (42)$$

571 where $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$ is the vector of statistics obtained via the SA-step (7) and using the
572 MC approximation of the sufficient statistics $(S_1(z), S_2(z), S_3(z))$ defined in (41).

573 F.2.2 Numerical Applications

574 For the inference of the template, we use the Matlab code (online SAEM) used in [19] and implement
575 our own batch, incremental, Variance reduced and Fast Incremental variants. The hyperparameters
576 are kept the same and reads as follows $M = 400$, $\gamma_k = 1/k^{0.6}$ and $p = 16$. The number of
577 landmarks for the template is $k_p = 15$ points and for the deformation $k_g = 6$ points. Both have
578 Gaussian kernels with respectively standard deviation of 0.08 and 0.16. The standard deviation of
579 the measurement errors is set to 0.1.

580 For the simulation part, we use the Carlin and Chib MCMC procedure, see [7]. Refer to [19] for
581 more details.

582 G Additional Experiment: Pharmacokinetics (PK) Model with Absorption 583 Lag Time

584 This numerical example was conducted in order to characterize the pharmacokinetics (PK) of orally
585 administered drug to simulated patients, using a population pharmacokinetics approach. $M = 50$
586 synthetic datasets were generated for $n = 5000$ patients with 10 observations (concentration mea-
587 sures) per patient. The goal is to model the evolution of the concentration of the absorbed drug
588 using a nonlinear and latent variable model.

589 **Model and Explicit Updates:** We consider a one-compartment PK model for oral administration
590 with an absorption lag-time (T^{lag}), assuming first-order absorption and linear elimination processes.
591 The final model includes the following variables: ka the absorption rate constant, V the volume of
592 distribution, k the elimination rate constant and T^{lag} the absorption lag-time. We also add several
593 covariates to our model such as D the dose of drug administered, t the time at which measures
594 are taken and the weight of the patient influencing the volume V . More precisely, the log-volume
595 $\log(V)$ is a linear function of the log-weight $lw70 = \log(wt/70)$. Let $z_i = (T_i^{\text{lag}}, ka_i, V_i, k_i)$ be the
596 vector of individual PK parameters, different for each individual i . The final model reads:

$$y_{ij} = f(t_{ij}, z_i) + \varepsilon_{ij} \quad \text{where} \quad f(t_{ij}, z_i) = \frac{D ka_i}{V(ka_i - k_i)} (e^{-ka_i(t_{ij} - T_i^{\text{lag}})} - e^{-k_i(t_{ij} - T_i^{\text{lag}})}) \quad (43)$$

where y_{ij} is the j -th concentration measurement of the drug of dosage D injected at time t_{ij} for patient i . We assume in this example that the residual errors ε_{ij} are independent and normally distributed with mean 0 and variance σ^2 . Lognormal distributions are used for the four PK parameters.

Lognormal distributions are used for the four PK parameters:

$$\log(T_i^{\text{lag}}) \sim \mathcal{N}(\log(T_{\text{pop}}^{\text{lag}}), \omega_{T^{\text{lag}}}^2), \log(ka_i) \sim \mathcal{N}(\log(ka_{\text{pop}}), \omega_{ka}^2), \quad (44)$$

$$\log(V_i) \sim \mathcal{N}(\log(V_{\text{pop}}), \omega_V^2), \log(k_i) \sim \mathcal{N}(\log(k_{\text{pop}}), \omega_k^2). \quad (45)$$

We recall that the complete model (y, z) defined by (43) belongs to the curved exponential family, which vector of sufficient statistics $S = (S_1(z), S_2(z), S_3(z))$ read:

$$S_1(z) = \frac{1}{n} \sum_{i=1}^n z_i, \quad S_2(z) = \frac{1}{n} \sum_{i=1}^n z_i^\top z_i, \quad S_3(z) = \frac{1}{n} \sum_{i=1}^n (y_i - f(t_i, z_i))^2 \quad (46)$$

where we have noted y_i and t_i the vector of observations and time for each patient i . At iteration k , and setting the number of MC samples to 1 for the sake of clarity, the MC sampling $z_i^{(k)} \sim p(z_i|y_i, \theta^{(k)})$ is performed using a Metropolis-Hastings procedure detailed in Algorithm 2. The quantities $\tilde{S}^{(k+1)}$ and $\hat{s}^{(k+1)}$ are then updated according to the different methods. Finally the maximization step yields:

$$\bar{\theta}(s) = \begin{pmatrix} \hat{s}_1^{(k+1)} \\ \hat{s}_2^{(k+1)} - \hat{s}_1^{(k+1)} (\hat{s}_1^{(k+1)})^\top \\ \hat{s}_3^{(k+1)} \end{pmatrix} = \begin{pmatrix} \overline{z_{\text{pop}}}(\hat{s}^{(k+1)}) \\ \overline{\omega_z}(\hat{s}^{(k+1)}) \\ \overline{\sigma}(\hat{s}^{(k+1)}) \end{pmatrix}. \quad (47)$$

Metropolis Hastings algorithm During the simulation step of the MISSO method, the sampling from the target distribution $\pi(z_i, \theta) := p(z_i|y_i, \theta)$ is performed using a Metropolis Hastings (MH) algorithm [22] with proposal distribution $q(z_i, \delta)$ where $\theta = (z_{\text{pop}}, \omega_z)$ and δ is the vector of parameters of the proposal distribution. Commonly they parameterize a Gaussian proposal. The MH algorithm is summarized in 2.

Algorithm 2 MH algorithm

```

1: Input: initialization  $z_{i,0} \sim q(z_i; \delta)$ 
2: for  $m = 1, \dots, M$  do
3:   Sample  $z_{i,m} \sim q(z_i; \delta)$ 
4:   Sample  $u \sim \mathcal{U}([0, 1])$ 
5:   Calculate the ratio  $r = \frac{\pi(z_{i,m}; \theta) / q(z_{i,m}; \delta)}{\pi(z_{i,m-1}; \theta) / q(z_{i,m-1}; \delta)}$ 
6:   if  $u < r$  then
7:     Accept  $z_{i,m}$ 
8:   else
9:      $z_{i,m} \leftarrow z_{i,m-1}$ 
10:  end if
11: end for
12: Output:  $z_{i,M}$ 

```

Monte Carlo study: We conduct a Monte Carlo study to showcase the benefits of our scheme. $M = 50$ datasets have been simulated using the following PK parameters values: $T_{\text{pop}}^{\text{lag}} = 1$, $ka_{\text{pop}} = 1$, $V_{\text{pop}} = 8$, $k_{\text{pop}} = 0.1$, $\omega_{T^{\text{lag}}} = 0.4$, $\omega_{ka} = 0.5$, $\omega_V = 0.2$, $\omega_k = 0.3$ and $\sigma^2 = 0.5$. We define the mean square distance over the M replicates $E_k(\ell) = \frac{1}{M} \sum_{m=1}^M (\theta_k^{(m)}(\ell) - \theta^*)^2$ and plot it against the epochs (passes over the data) Figure 4. Note that the MC-step (5) is performed using a Metropolis Hastings procedure since the posterior distribution under the model θ noted $p(z_i|y_i, \theta)$ is intractable due to the nonlinearity of the model (43). Figure 4 shows clear advantage of variance reduced methods (vrTTEM and fitTEM) avoiding the twists and turns displayed by the incremental and the batch methods.




Figure 4: Precision $|ka^{(k)} - ka^*|^2$ per epoch