

---

## On the Convergence of Decentralized Adaptive Gradient Methods (Appendix)

---

The main purpose of this appendix is to give thorough and detailed proofs for our convergence analysis described in the main paper. After having established several important Lemmas in Section A, we provide a proof for our main Theorem, namely Theorem 2, in Section B. Section C and Section D correspond to the proofs for the extension and application of Theorem 2 to the AMSGrad and AdaGrad algorithms used as prototypes of our general class of decentralized adaptive gradient methods. Section E contains additional numerical runs for more empirical insights on our scheme.

### A Proof of Auxiliary Lemmas

Similarly to [38; 8] with SGD (with momentum) and centralized adaptive gradient methods, define the following auxiliary sequence:

$$Z_t = \bar{X}_t + \frac{\beta_1}{1 - \beta_1} (\bar{X}_t - \bar{X}_{t-1}), \quad (5)$$

with  $\bar{X}_0 \triangleq \bar{X}_1$ . Such an auxiliary sequence can help us deal with the bias brought by the momentum and simplifies the convergence analysis.

**Lemma A.1.** *For the sequence defined in (5), we have*

$$Z_{t+1} - Z_t = \alpha \frac{\beta_1}{1 - \beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}}.$$

**Proof:** By update rule of Algorithm 2, we first have

$$\begin{aligned} \bar{X}_{t+1} &= \frac{1}{N} \sum_{i=1}^N x_{t+1,i} \\ &= \frac{1}{N} \sum_{i=1}^N \left( x_{t+0.5,i} - \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) \\ &= \frac{1}{N} \sum_{i=1}^N \left( \sum_{j=1}^N W_{ij} x_{t,j} - \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) \\ &\stackrel{(i)}{=} \left( \frac{1}{N} \sum_{j=1}^N x_{t,j} \right) - \frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \\ &= \bar{X}_t - \frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}}, \end{aligned}$$

where (i) is due to an interchange of summation and  $\sum_{i=1} W_{ij} = 1$ . Then, we have

$$\begin{aligned}
Z_{t+1} - Z_t &= \bar{X}_{t+1} - \bar{X}_t + \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) - \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) \\
&= \frac{1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) - \frac{\beta_1}{1 - \beta_1} (\bar{X}_{t+1} - \bar{X}_t) \\
&= \frac{1}{1 - \beta_1} \left( -\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t,i}}{\sqrt{u_{t,i}}} \right) - \frac{\beta_1}{1 - \beta_1} \left( -\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right) \\
&= \frac{1}{1 - \beta_1} \left( -\frac{1}{N} \sum_{i=1}^N \alpha \frac{\beta_1 m_{t-1,i} + (1 - \beta_1) g_{t,i}}{\sqrt{u_{t,i}}} \right) - \frac{\beta_1}{1 - \beta_1} \left( -\frac{1}{N} \sum_{i=1}^N \alpha \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right) \\
&= \alpha \frac{\beta_1}{1 - \beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}},
\end{aligned}$$

which is the desired result.  $\square$

**Lemma A.2.** Given a set of numbers  $a_1, \dots, a_n$  and denote their mean to be  $\bar{a} = \frac{1}{n} \sum_{i=1}^n a_i$ . Define  $b_i(r) \triangleq \max(a_i, r)$  and  $\bar{b}(r) = \frac{1}{n} \sum_{i=1}^n b_i(r)$ . For any  $r$  and  $r'$  with  $r' \geq r$  we have

$$\sum_{i=1}^n |b_i(r) - \bar{b}(r)| \geq \sum_{i=1}^n |b_i(r') - \bar{b}(r')| \quad (6)$$

and when  $r \leq \min_{i \in [n]} a_i$ , we have

$$\sum_{i=1}^n |b_i(r) - \bar{b}(r)| = \sum_{i=1}^n |a_i - \bar{a}|. \quad (7)$$

**Proof:** Without loss of generality, assume  $a_i \leq a_j$  when  $i < j$ , i.e.  $a_i$  is a non-decreasing sequence. Define

$$h(r) = \sum_{i=1}^n |b_i(r) - \bar{b}(r)| = \sum_{i=1}^n \left| \max(a_i, r) - \frac{1}{n} \sum_{j=1}^n \max(a_j, r) \right|.$$

We need to prove that  $h$  is a non-increasing function of  $r$ . First, it is easy to see that  $h$  is a continuous function of  $r$  with non-differentiable points  $r = a_i, i \in [n]$ , thus  $h$  is a piece-wise linear function.

Next, we will prove that  $h(r)$  is non-increasing in each piece. Define  $l(r)$  to be the largest index with  $a(l(r)) < r$ , and  $s(r)$  to be the largest index with  $a_{s(r)} < \bar{b}(r)$ . Note that we have for  $i \leq l(r)$ ,  $b_i(r) = r$  and for  $i \leq s(r)$   $b_i(r) - \bar{b}(r) \leq 0$  since  $a_i$  is a non-decreasing sequence. Therefore, we have

$$h(r) = \sum_{i=1}^{l(r)} (\bar{b}(r) - r) + \sum_{i=l(r)+1}^{s(r)} (\bar{b}(r) - a_i) + \sum_{i=s(r)+1}^n (a_i - \bar{b}(r))$$

and

$$\bar{b}(r) = \frac{1}{n} \left( l(r)r + \sum_{i=l(r)+1}^n a_i \right).$$

Taking derivative of the above form, we know the derivative of  $h(r)$  at differentiable points is

$$\begin{aligned}
h'(r) &= l(r) \left( \frac{l(r)}{n} - 1 \right) + (s(r) - l(r)) \frac{l(r)}{n} - (n - s(r)) \frac{l(r)}{n} \\
&= \frac{l(r)}{n} ((l(r) - n) + (s(r) - l(r)) - (n - s(r))).
\end{aligned}$$

Since we have  $s(r) \leq n$  we know  $(l(r) - n) + (s(r) - l(r)) - (n - s(r)) \leq 0$  and thus

$$h'(r) \leq 0,$$

which means  $h(r)$  is non-increasing in each piece. Combining with the fact that  $h(r)$  is continuous, (6) is proven. When  $r \leq a(i)$ , we have  $b(i) = \max(a_i, r) = r$ , for all  $r \in [n]$  and  $\bar{b}(r) = \frac{1}{n} \sum_{i=1}^n a_i = \bar{a}$  which proves (7).  $\square$

## 576 B Proof of Theorem 2

577 To prove convergence of the algorithm, we first define an auxiliary sequence

$$Z_t = \bar{X}_t + \frac{\beta_1}{1 - \beta_1} (\bar{X}_t - \bar{X}_{t-1}), \quad (8)$$

578 with  $\bar{X}_0 \triangleq \bar{X}_1$ . Since  $\mathbb{E}[g_{t,i}] = \nabla f(x_{t,i})$  and  $u_{t,i}$  is a function of  $G_{1:t-1}$  (which denotes  
579  $G_1, G_2, \dots, G_{t-1}$ ), we have

$$\mathbb{E}_{G_t|G_{1:t-1}} \left[ \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right] = \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}}.$$

580 Assuming smoothness (A1) we have

$$f(Z_{t+1}) \leq f(Z_t) + \langle \nabla f(Z_t), Z_{t+1} - Z_t \rangle + \frac{L}{2} \|Z_{t+1} - Z_t\|^2.$$

581 Using Lemma A.1 into the above inequality and take expectation over  $G_t$  given  $G_{1:t-1}$ , we have

$$\begin{aligned} & \mathbb{E}_{G_t|G_{1:t-1}} [f(Z_{t+1})] \\ & \leq f(Z_t) - \alpha \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle + \frac{L}{2} \mathbb{E}_{G_t|G_{1:t-1}} [\|Z_{t+1} - Z_t\|^2] \\ & \quad + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E}_{G_t|G_{1:t-1}} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]. \end{aligned}$$

582 Then take expectation over  $G_{1:t-1}$  and rearrange, we have

$$\alpha \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle \right] \quad (9)$$

$$\begin{aligned} & \leq \mathbb{E}[f(Z_t)] - \mathbb{E}[f(Z_{t+1})] + \frac{L}{2} \mathbb{E} [\|Z_{t+1} - Z_t\|^2] \\ & \quad + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]. \end{aligned} \quad (10)$$

583 In addition, we have

$$\begin{aligned} & \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\rangle \\ & = \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\rangle + \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\rangle \end{aligned} \quad (11)$$

584 and the first term on RHS of the equality can be lower bounded as

$$\begin{aligned} & \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\rangle \\ & = \frac{1}{2} \left\| \frac{\nabla f(Z_t)}{\bar{U}_t^{1/4}} \right\|^2 + \frac{1}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\nabla f(Z_t) - \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 \\ & \geq \frac{1}{4} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \frac{1}{4} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\nabla f(Z_t) - \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i})}{\bar{U}_t^{1/4}} \right\|^2 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2} \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{1}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \\
& \geq \frac{1}{2} \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{3}{2} \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 - \frac{3}{2} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2, \quad (12)
\end{aligned}$$

585 where the inequalities are all due to Cauchy-Schwartz. Substituting (12) and (11) into (9), we get

$$\begin{aligned}
\frac{1}{2} \alpha \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] & \leq \mathbb{E}[f(Z_t)] - \mathbb{E}[f(Z_{t+1})] + \frac{L}{2} \mathbb{E}[\|Z_{t+1} - Z_t\|^2] \\
& + \alpha \frac{\beta_1}{1 - \beta_1} \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\
& - \alpha \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\rangle \right] \\
& + \frac{3}{2} \alpha \mathbb{E} \left[ \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right].
\end{aligned}$$

586 Then sum over the above inequality from  $t = 1$  to  $T$  and divide both sides by  $T\alpha/2$ , we have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
& \leq \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + \frac{L}{T\alpha} \sum_{t=1}^T \mathbb{E}[\|Z_{t+1} - Z_t\|^2] \\
& + \frac{2}{T} \frac{\beta_1}{1 - \beta_1} \sum_{t=1}^T \underbrace{\mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]}_{D_1} \\
& + \frac{2}{T} \sum_{t=1}^T \underbrace{\mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{\bar{U}_t}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right]}_{D_2} \\
& + \frac{3}{T} \sum_{t=1}^T \underbrace{\mathbb{E} \left[ \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 + \left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right]}_{D_3}. \quad (13)
\end{aligned}$$

587 Now we need to upper bound all the terms on RHS of the above inequality to get the convergence  
588 rate. For the terms composing  $D_3$  in (13), we can upper bound them by

$$\begin{aligned}
\left\| \frac{\nabla f(Z_t) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 & \leq \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \|\nabla f(Z_t) - \nabla f(\bar{X}_t)\|^2 \\
& \leq L \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \underbrace{\|Z_t - \bar{X}_t\|^2}_{D_4} \quad (14)
\end{aligned}$$

589 and

$$\begin{aligned} \left\| \frac{\frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 &\leq \frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \frac{1}{N} \sum_{i=1}^N \|\nabla f_i(x_{t,i}) - \nabla f(\bar{X}_t)\|^2 \\ &\leq L \underbrace{\frac{1}{\min_{j \in [d]} [\bar{U}_t^{1/2}]_j} \frac{1}{N} \sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2}_{D_5}, \end{aligned} \quad (15)$$

590 using Jensen's inequality, Lipschitz continuity of  $f_i$ , and the fact that  $f = \frac{1}{N} \sum_{i=1}^N f_i$ . Next we need  
591 to bound  $D_4$  and  $D_5$ . Recall the update rule of  $X_t$ , we have

$$X_t = X_{t-1}W - \alpha \frac{M_{t-1}}{\sqrt{U_{t-1}}} = X_1 W^{t-1} - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} W^k, \quad (16)$$

592 where we define  $W^0 = \mathbf{I}$ . Since  $W$  is a symmetric matrix, we can decompose it as  $W = Q\Lambda Q^T$   
593 where  $Q$  is a orthonormal matrix and  $\Lambda$  is a diagonal matrix whose diagonal elements correspond  
594 to eigenvalues of  $W$  in an descending order, i.e.  $\Lambda_{ii} = \lambda_i$  with  $\lambda_i$  being  $i$ th largest eigenvalue of  
595  $W$ . In addition, because  $W$  is a doubly stochastic matrix, we know  $\lambda_1 = 1$  and  $q_1 = \frac{1}{\sqrt{N}}$ . With  
596 eigen-decomposition of  $W$ , we can rewrite  $D_5$  as

$$\sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 = \|X_t - \bar{X}_t \mathbf{1}_N^T\|_F^2 = \|X_t Q Q^T - X_t \frac{1}{N} \mathbf{1}_N \mathbf{1}_N^T\|_F^2 = \sum_{l=2}^N \|X_t q_l\|^2. \quad (17)$$

597 In addition, we can rewrite (16) as

$$X_t = X_1 W^{t-1} - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} W^k = X_1 - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} Q \Lambda^k Q^T, \quad (18)$$

598 where the last equality is because  $x_{1,i} = x_{1,j}$ , for all  $i, j$  and thus  $X_1 W = X_1$ . Then we have when  
599  $l > 1$ ,

$$X_t q_l = (X_1 - \alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} Q \Lambda^k Q^T) q_l = -\alpha \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} q_l \lambda_l^k, \quad (19)$$

600 since  $Q$  is orthonormal and  $X_1 q_l = x_{1,1} \mathbf{1}_N^T q_l = x_{1,1} \sqrt{N} q_1^T q_l = 0$ , for all  $l \neq 1$ .

601 Combining (17) and (19), we have

$$\begin{aligned} D_5 &= \sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 = \sum_{l=2}^N \|X_t q_l\|^2 \\ &= \sum_{l=2}^N \alpha^2 \left\| \sum_{k=0}^{t-2} \frac{M_{t-k-1}}{\sqrt{U_{t-k-1}}} \lambda_l^k q_l \right\|^2 \\ &\leq \alpha^2 \left( \frac{1}{1-\lambda} \right)^2 N d G_\infty^2 \frac{1}{\epsilon}, \end{aligned} \quad (20)$$

602 where the last inequality follows from the fact that  $g_{t,i} \leq G_\infty$ ,  $\|q_l\| = 1$ , and  $|\lambda_l| \leq \lambda < 1$ . Now let  
603 us turn to  $D_4$ , it can be rewritten as

$$\begin{aligned} \|Z_t - \bar{X}_t\|^2 &= \left\| \frac{\beta_1}{1-\beta_1} (\bar{X}_t - \bar{X}_{t-1}) \right\|^2 = \left( \frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{m_{t-1,i}}{\sqrt{u_{t-1,i}}} \right\|^2 \\ &\leq \left( \frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 d \frac{G_\infty^2}{\epsilon}. \end{aligned} \quad (21)$$

604 Now we know both  $D_4$  and  $D_5$  are in the order of  $\mathcal{O}(\alpha^2)$  and thus  $D_3$  is in the order of  
605  $\mathcal{O}(\alpha^2)$ . Next we will bound  $D_2$  and  $D_1$ . Define  $G_1 \triangleq \max_{t \in [T]} \max_{i \in [N]} \|\nabla f_i(x_{t,i})\|_\infty$ ,

606  $G_2 \triangleq \max_{t \in [T]} \|\nabla f(Z_t)\|_\infty$ ,  $G_3 \triangleq \max_{t \in [T]} \max_{i \in [N]} \|g_{t,i}\|_\infty$  and  $G_\infty = \max(G_1, G_2, G_3)$ .  
 607 Then we have

$$\begin{aligned}
 D_2 &= \sum_{t=1}^T \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{\bar{U}_t}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\
 &\leq \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[\bar{U}_t]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \right] \\
 &= \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[\bar{U}_t]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \frac{\sqrt{[\bar{U}_t]_j} + \sqrt{[u_{t,i}]_j}}{\sqrt{[\bar{U}_t]_j} + \sqrt{[u_{t,i}]_j}} \right] \\
 &= \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{[\bar{U}_t]_j - [u_{t,i}]_j}{[\bar{U}_t]_j \sqrt{[u_{t,i}]_j} + \sqrt{[\bar{U}_t]_j} [u_{t,i}]_j} \right| \right] \\
 &\leq \underbrace{\mathbb{E} \left[ \sum_{t=1}^T G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{[\bar{U}_t]_j - [u_{t,i}]_j}{2\epsilon^{1.5}} \right| \right]}_{D_6},
 \end{aligned} \tag{22}$$

608 where the last inequality is due to  $[u_{t,i}]_j \geq \epsilon$ , for all  $t, i, j$ . To simplify notations, define  $\|A\|_{abs} =$   
 609  $\sum_{i,j} |A_{ij}|$  to be the entry-wise  $L_1$  norm of a matrix  $A$ , then we obtain

$$\begin{aligned}
 D_6 &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\bar{U}_t \mathbf{1}^T - U_t\|_{abs} \leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\bar{U}_t \mathbf{1}^T - \tilde{U}_t\|_{abs} \\
 &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \|\tilde{U}_t \frac{1}{N} \mathbf{1}_N \mathbf{1}_N^T - \tilde{U}_t Q Q^T\|_{abs} \\
 &= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs},
 \end{aligned}$$

610 where the second inequality is due to Lemma A.2, introduced Section A, and the fact that  $U_t =$   
 611  $\max(\tilde{U}_t, \epsilon)$  (element-wise max operator). Recall from update rule of  $U_t$ , by defining  $\hat{V}_{-1} \triangleq \hat{V}_0$  and  
 612  $U_0 \triangleq U_{1/2}$ , we have for all  $t \geq 0$ ,  $\tilde{U}_{t+1} = (\tilde{U}_t - \hat{V}_{t-1} + \hat{V}_t)W$ . Thus, we obtain

$$\tilde{U}_t = \tilde{U}_0 W^t + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) W^k = \tilde{U}_0 + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) Q \Lambda^k Q^T.$$

613 Then we further obtain when  $l \neq 1$ ,

$$\tilde{U}_t q_l = (\tilde{U}_0 + \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) Q \Lambda^k Q^T) q_l = \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) q_l \lambda_l^k,$$

614 where the last equality is due to the definition  $\tilde{U}_0 \triangleq U_{1/2} = \epsilon \mathbf{1}_d \mathbf{1}_N^T = \sqrt{N} \epsilon \mathbf{1}_d \mathbf{1}_N^T$  (recall that  
 615  $q_1 = \frac{1}{\sqrt{N}} \mathbf{1}_N^T$ ) and  $q_i^T q_j = 0$  when  $i \neq j$ . Note that by definition of  $\|\cdot\|_{abs}$ , we have for all

616  $A, B, \|A + B\|_{abs} \leq \|A\|_{abs} + \|B\|_{abs}$ , then

$$\begin{aligned}
D_6 &\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs} \\
&= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \left\| - \sum_{k=1}^t (-\hat{V}_{t-1-k} + \hat{V}_{t-k}) \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_{abs} \\
&\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \left\| \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_1 \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \\
&\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \sqrt{N} \left\| \sum_{l=2}^N q_l \lambda_l^k q_l^T \right\|_2 \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \\
&\leq \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \sum_{j=1}^d \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})^T e_j\|_1 \sqrt{N} \lambda^k \\
&= \frac{G_\infty^2}{N} \sum_{t=1}^T \frac{1}{2\epsilon^{1.5}} \sum_{k=1}^t \|(-\hat{V}_{t-1-k} + \hat{V}_{t-k})\|_{abs} \sqrt{N} \lambda^k \\
&= \frac{G_\infty^2}{N} \frac{1}{2\epsilon^{1.5}} \sum_{o=0}^{T-1} \sum_{t=o+1}^T \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \sqrt{N} \lambda^{t-o} \\
&\leq \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs},
\end{aligned} \tag{23}$$

617 where  $\lambda = \max(|\lambda_2|, |\lambda_N|)$ . Combining (22) and (23), we have

$$D_2 \leq \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \frac{\lambda}{1-\lambda} \mathbb{E} \left[ \sum_{o=0}^{T-1} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right].$$

618 Now we need to bound  $D_1$ , we have

$$\begin{aligned}
D_1 &= \sum_{t=1}^T \mathbb{E} \left[ \left\langle \nabla f(Z_t), \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\rangle \right] \\
&\leq \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| \right] \\
&= \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \left( \frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right) \frac{\sqrt{[u_{t,i}]_j} + \sqrt{[u_{t-1,i}]_j}}{\sqrt{[u_{t,i}]_j} + \sqrt{[u_{t-1,i}]_j}} \right| \right] \\
&\leq \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \left| \frac{1}{2\epsilon^{1.5}} ([u_{t-1,i}]_j - [u_{t,i}]_j) \right| \right] \\
&\stackrel{(a)}{\leq} \sum_{t=1}^T \mathbb{E} \left[ G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{2\epsilon^{1.5}} |([\tilde{u}_{t-1,i}]_j - [\tilde{u}_{t,i}]_j)| \right] \\
&= G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{N} \mathbb{E} \left[ \sum_{t=1}^T \|\tilde{U}_{t-1} - \tilde{U}_t\|_{abs} \right],
\end{aligned} \tag{24}$$

where (a) is due to  $[\tilde{u}_{t-1,i}]_j = \max([u_{t-1,i}]_j, \epsilon)$  and the function  $\max(\cdot, \epsilon)$  is 1-Lipschitz. In addition, by update rule of  $U_t$ , we have

$$\begin{aligned}
& \sum_{t=1}^T \|\tilde{U}_{t-1} - \tilde{U}_t\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1} - (\tilde{U}_{t-1} - \hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1}(QQ^T - Q\Lambda Q^T) + (-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&= \sum_{t=1}^T \|\tilde{U}_{t-1}(\sum_{l=2}^N q_l(1 - \lambda_l)q_l^T) + (-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&\leq \sum_{t=1}^T \left\| \sum_{k=1}^{t-1} (-\hat{V}_{t-2-k} + \hat{V}_{t-1-k}) \sum_{l=2}^N q_l \lambda_l^k (1 - \lambda_l) q_l^T \right\|_{abs} + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})W\|_{abs} \\
&\leq \sum_{t=1}^T \left( \sum_{k=1}^{t-1} \|\hat{V}_{t-2-k} - \hat{V}_{t-1-k}\|_{abs} \sqrt{N} \lambda^k \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&= \sum_{t=1}^T \left( \sum_{o=1}^{t-1} \|\hat{V}_{t-2-o} - \hat{V}_{t-1-o}\|_{abs} \sqrt{N} \lambda^{t-o} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&= \sum_{o=1}^{T-1} \sum_{t=o+1}^T \left( \|\hat{V}_{t-2-o} - \hat{V}_{t-1-o}\|_{abs} \sqrt{N} \lambda^{t-o} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&\leq \sum_{o=1}^{T-1} \frac{\lambda}{1-\lambda} \left( \|\hat{V}_{t-2-o} - \hat{V}_{t-1-o}\|_{abs} \sqrt{N} \right) + \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \\
&\leq \frac{1}{1-\lambda} \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \sqrt{N}.
\end{aligned} \tag{25}$$

Combining (24) and (25), we have

$$D_1 \leq G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{N} \mathbb{E} \left[ \frac{1}{1-\lambda} \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \sqrt{N} \right]. \tag{26}$$

What remains is to bound  $\sum_{t=1}^T \mathbb{E} [\|Z_{t+1} - Z_t\|^2]$ . By update rule of  $Z_t$ , we have

$$\begin{aligned}
& \|Z_{t+1} - Z_t\|^2 \\
&= \left\| \alpha \frac{\beta_1}{1-\beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) - \alpha \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\
&\leq 2\alpha^2 \left\| \frac{\beta_1}{1-\beta_1} \frac{1}{N} \sum_{i=1}^N m_{t-1,i} \odot \left( \frac{1}{\sqrt{u_{t-1,i}}} - \frac{1}{\sqrt{u_{t,i}}} \right) \right\|^2 + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\
&\leq 2\alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{\sqrt{\epsilon}} \left| \frac{1}{\sqrt{[u_{t-1,i}]_j}} - \frac{1}{\sqrt{[u_{t,i}]_j}} \right| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\
&\leq 2\alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{\sqrt{\epsilon}} \left| \frac{[u_{t,i}]_j - [u_{t-1,i}]_j}{2\epsilon^{1.5}} \right| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \\
&\leq 2\alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \sum_{i=1}^N \sum_{j=1}^d \frac{1}{2\epsilon^2} |\tilde{u}_{t,i,j} - \tilde{u}_{t-1,i,j}| + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2
\end{aligned}$$



$$= 2\alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \frac{1}{2\epsilon^2} \|\tilde{U}_t - \tilde{U}_{t-1}\|_{abs} + 2\alpha^2 \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2, \quad (27)$$

where the last inequality is again due to the definition that  $[\tilde{u}_{t,i}]_j = \max([u_{t,i}]_j, \epsilon)$  and the fact that  $\max(\cdot, \epsilon)$  is 1-Lipschitz. Then, we have

$$\begin{aligned} & \sum_{t=1}^T \mathbb{E}[\|Z_{t+1} - Z_t\|^2] \\ & \leq 2\alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 \frac{1}{N} \frac{1}{2\epsilon^2} \mathbb{E} \left[ \sum_{t=1}^T \|\tilde{U}_t - \tilde{U}_{t-1}\|_{abs} \right] + 2\alpha^2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\ & \leq \alpha^2 \left( \frac{\beta_1}{1-\beta_1} \right)^2 \frac{G_\infty^2}{\sqrt{N}} \frac{1}{\epsilon^2} \frac{1}{1-\lambda} \mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] + 2\alpha^2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right], \end{aligned}$$

where the last inequality is due to (25).

We now bound the last term on RHS of the above inequality. A trivial bound can be

$$\sum_{t=1}^T \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \leq \sum_{t=1}^T d G_\infty^2 \frac{1}{\epsilon},$$

due to  $\|g_{t,i}\| \leq G_\infty$  and  $[u_{t,i}]_j \geq \epsilon$ , for all  $j$  (verified from update rule of  $u_{t,i}$  and the assumption that  $[v_{t,i}]_j \geq \epsilon$ , for all  $i$ ). However, the above bound is independent of  $N$ , to get a better bound, we need a more involved analysis to show its dependency on  $N$ . To do this, we first notice that

$$\begin{aligned} & \mathbb{E}_{G_t|G_{1:t-1}} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\ & = \mathbb{E}_{G_t|G_{1:t-1}} \left[ \frac{1}{N^2} \sum_{i=1}^N \sum_{j=1}^N \left\langle \frac{\nabla f_i(x_{t,i}) + \xi_{t,i}}{\sqrt{u_{t,i}}}, \frac{\nabla f_j(x_{t,j}) + \xi_{t,j}}{\sqrt{u_{t,j}}} \right\rangle \right] \\ & \stackrel{(a)}{=} \mathbb{E}_{G_t|G_{1:t-1}} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] + \mathbb{E}_{G_t|G_{1:t-1}} \left[ \frac{1}{N^2} \sum_{i=1}^N \left\| \frac{\xi_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \\ & \stackrel{(b)}{=} \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{1}{N^2} \sum_{i=1}^N \sum_{l=1}^d \frac{\mathbb{E}_{G_t|G_{1:t-1}}[\xi_{t,i}^2]}{[u_{t,i}]_l} \\ & \stackrel{(c)}{\leq} \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{d}{N} \frac{\sigma^2}{\epsilon}, \end{aligned}$$

where (a) is due to  $\mathbb{E}_{G_t|G_{1:t-1}}[\xi_{t,i}] = 0$  and  $\xi_{t,i}$  is independent of  $x_{t,j}, u_{t,j}$  for all  $j$ , and  $\xi_j$ , for all  $j \neq i$ , (b) comes from the fact that  $x_{t,i}, u_{t,i}$  are fixed given  $G_{1:t}$ , (c) is due to  $\mathbb{E}_{G_t|G_{1:t-1}}[\xi_{t,i}^2] \leq \sigma^2$  and  $[u_{t,i}]_l \geq \epsilon$  by definition. Then we have

$$\begin{aligned} \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] & = \mathbb{E}_{G_{1:t-1}} \left[ \mathbb{E}_{G_t|G_{1:t-1}} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] \right] \\ & \leq \mathbb{E}_{G_{1:t-1}} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 + \frac{d}{N} \frac{\sigma^2}{\epsilon} \right] \\ & = \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] + \frac{d}{N} \frac{\sigma^2}{\epsilon}. \end{aligned} \quad (28)$$

633 In traditional analysis of SGD-like distributed algorithms, the term corresponding to  
634  $\mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right]$  will be merged with the first order descent when the stepsize is cho-  
635 sen to be small enough. However, in our case, the term cannot be merged because it is different from  
636 the first order descent in our algorithm. A brute-force upper bound is possible but this will lead to a  
637 worse convergence rate in terms of  $N$ . Thus, we need a more detailed analysis for the term in the  
638 following.

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] \\
&= \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} + \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right] \\
&\leq 2\mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right] \\
&\leq 2\mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[ \frac{1}{N} \sum_{i=1}^N \left\| \nabla f_i(x_{t,i}) \odot \left( \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right) \right\|^2 \right] \\
&\leq 2\mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2\mathbb{E} \left[ \frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right].
\end{aligned}$$

639 Summing over  $T$ , we have

$$\begin{aligned}
& \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{u_{t,i}}} \right\|^2 \right] \\
&\leq 2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[ \frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right]. \quad (29)
\end{aligned}$$

640 For the last term on RHS of (29), we can bound it similarly as what we did for  $D_2$  from (22) to (23),  
641 which yields

$$\begin{aligned}
& \sum_{t=1}^T \mathbb{E} \left[ \frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \left\| \frac{1}{\sqrt{u_{t,i}}} - \frac{1}{\sqrt{\bar{U}_t}} \right\|_1 \right] \leq \sum_{t=1}^T \mathbb{E} \left[ \frac{1}{N} \sum_{i=1}^N G_\infty^2 \frac{1}{\sqrt{\epsilon}} \frac{1}{2\epsilon^{1.5}} \|u_{t,i} - \bar{U}_t\|_1 \right] \\
&= \sum_{t=1}^T \mathbb{E} \left[ \frac{1}{N} G_\infty^2 \frac{1}{2\epsilon^2} \|\bar{U}_t \mathbf{1}^T - U_t\|_{abs} \right] \\
&\leq \sum_{t=1}^T \mathbb{E} \left[ \frac{1}{N} G_\infty^2 \frac{1}{2\epsilon^2} \left\| - \sum_{l=2}^N \tilde{U}_t q_l q_l^T \right\|_{abs} \right] \\
&\leq \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[ \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right]. \quad (30)
\end{aligned}$$

642 Further, we have

$$\begin{aligned}
& \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] \\
& \leq 2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] \\
& = 2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 2 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right]
\end{aligned}$$

643 and the last term on RHS of the above inequality can be bounded following similar procedures from  
644 (15) to (20), as what we did for  $D_3$ . Completing the procedures yields

$$\begin{aligned}
\sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{\nabla f_i(\bar{X}_t) - \nabla f_i(x_{t,i})}{\sqrt{\bar{U}_t}} \right\|^2 \right] & \leq \sum_{t=1}^T \mathbb{E} \left[ L \frac{1}{\epsilon} \frac{1}{N} \sum_{i=1}^N \|x_{t,i} - \bar{X}_t\|^2 \right] \\
& \leq \sum_{t=1}^T \mathbb{E} \left[ L \frac{1}{\epsilon} \frac{1}{N} \alpha^2 \left( \frac{1}{1-\lambda} \right) N d G_\infty^2 \frac{1}{\epsilon} \right] \quad (31) \\
& = T L \frac{1}{\epsilon^2} \alpha^2 \left( \frac{1}{1-\lambda} \right) d G_\infty^2.
\end{aligned}$$

645 Finally, combining (28) to (31), we get

$$\begin{aligned}
\sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{1}{N} \sum_{i=1}^N \frac{g_{t,i}}{\sqrt{u_{t,i}}} \right\|^2 \right] & \leq 4 \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\sqrt{\bar{U}_t}} \right\|^2 \right] + 4 T L \frac{1}{\epsilon^2} \alpha^2 \left( \frac{1}{1-\lambda} \right) d G_\infty^2 \\
& \quad + 2 \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[ \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + T \frac{d}{N} \frac{\sigma^2}{\epsilon} \\
& \leq 4 \frac{1}{\sqrt{\epsilon}} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] + 4 T L \frac{1}{\epsilon^2} \alpha^2 \left( \frac{1}{1-\lambda} \right) d G_\infty^2 \\
& \quad + 2 \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[ \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + T \frac{d}{N} \frac{\sigma^2}{\epsilon}.
\end{aligned}$$

646 where the last inequality is due to each element of  $\bar{U}_t$  is lower bounded by  $\epsilon$  by definition.

647 Combining all above, we obtain

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
& \leq \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) \\
& \quad + \frac{L}{T} \alpha \left( \frac{\beta_1}{1-\beta_1} \right)^2 \frac{G_\infty^2}{\sqrt{N}} \frac{1}{\epsilon^2} \frac{1}{1-\lambda} \mathbb{E}[\mathcal{V}_T] \\
& \quad + \frac{8L}{T} \alpha \frac{1}{\sqrt{\epsilon}} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] + 8 L^2 \alpha \frac{1}{\epsilon^2} \alpha^2 \left( \frac{1}{1-\lambda} \right) d G_\infty^2 \quad (32) \\
& \quad + \frac{4L}{T} \alpha \frac{1}{\sqrt{N}} G_\infty^2 \frac{1}{2\epsilon^2} \mathbb{E} \left[ \sum_{o=0}^{T-1} \frac{\lambda}{1-\lambda} \|(-\hat{V}_{o-1} + \hat{V}_o)\|_{abs} \right] + 2 L \alpha \frac{d}{N} \frac{\sigma^2}{\epsilon}
\end{aligned}$$

$$\begin{aligned}
& + \frac{2}{T} \frac{\beta_1}{1-\beta_1} G_\infty^2 \frac{1}{2\epsilon^{1.5}} \frac{1}{\sqrt{N}} \mathbb{E} \left[ \frac{1}{1-\lambda} \mathcal{V}_T \right] \\
& + \frac{2}{T} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{2\epsilon^{1.5}} \frac{\lambda}{1-\lambda} \mathbb{E} [\mathcal{V}_T] \\
& + \frac{3}{T} \left( \sum_{t=1}^T L \left( \frac{1}{1-\lambda} \right)^2 \alpha^2 d G_\infty^2 \frac{1}{\epsilon^{1.5}} + \sum_{t=1}^T L \left( \frac{\beta_1}{1-\beta_1} \right)^2 \alpha^2 d \frac{G_\infty^2}{\epsilon^{1.5}} \right) \\
& = \frac{2}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + 2L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon} + 8L\alpha \frac{1}{\sqrt{\epsilon}} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
& + 3\alpha^2 d \left( \left( \frac{\beta_1}{1-\beta_1} \right)^2 + \left( \frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 8\alpha^3 L^2 \left( \frac{1}{1-\lambda} \right) d \frac{G_\infty^2}{\epsilon^2} \\
& + \frac{1}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left( L\alpha \left( \frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2L\alpha \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E} [\mathcal{V}_T] .
\end{aligned}$$

648 where  $\mathcal{V}_T := \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs}$ . Set  $\alpha = \frac{1}{\sqrt{dT}}$  and when  $\alpha \leq \frac{\epsilon^{0.5}}{16L}$ , we further have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] \\
& \leq \frac{4}{T\alpha} (\mathbb{E}[f(Z_1)] - \mathbb{E}[f(Z_{T+1})]) + 4L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon} \\
& \quad + 6\alpha^2 d \left( \left( \frac{\beta_1}{1-\beta_1} \right)^2 + \left( \frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 16\alpha^3 L^2 \left( \frac{1}{1-\lambda} \right) d \frac{G_\infty^2}{\epsilon^2} \\
& \quad + \frac{2}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left( L\alpha \left( \frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2L\alpha \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E} [\mathcal{V}_T] \\
& \leq \frac{4}{T\alpha} (\mathbb{E}[f(Z_1)] - \min_x f(x)) + 4L\alpha \frac{d}{N} \frac{\sigma^2}{\epsilon} \\
& \quad + 6\alpha^2 d \left( \left( \frac{\beta_1}{1-\beta_1} \right)^2 + \left( \frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}} + 16\alpha^3 d L^2 \left( \frac{1}{1-\lambda} \right) \frac{G_\infty^2}{\epsilon^2} \\
& \quad + \frac{2}{T\epsilon^{1.5}} \frac{G_\infty^2}{\sqrt{N}} \frac{1}{1-\lambda} \left( L\alpha \left( \frac{\beta_1}{1-\beta_1} \right)^2 \frac{1}{\epsilon^{0.5}} + \lambda + \frac{\beta_1}{1-\beta_1} + 2L\alpha \frac{1}{\epsilon^{0.5}} \lambda \right) \mathbb{E} [\mathcal{V}_T] \\
& \leq C_1 \left( \frac{1}{T\alpha} (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \alpha \frac{d\sigma^2}{N} \right) + C_2 \alpha^2 d + C_3 \alpha^3 d + \frac{1}{T\sqrt{N}} (C_4 + C_5 \alpha) \mathbb{E} [\mathcal{V}_T]
\end{aligned} \tag{33}$$

649 where the first inequality is obtained by moving the term  $8L\alpha \frac{1}{\sqrt{\epsilon}} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right]$  on the

650 RHS of (32) to the LHS to cancel it using the assumption  $8L\alpha \frac{1}{\sqrt{\epsilon}} \leq \frac{1}{2}$  followed by multiplying both  
651 sides by 2. The constants introduced in the last step are defined as following

$$\begin{aligned}
C_1 &= \max(4, 4L/\epsilon), \\
C_2 &= 6 \left( \left( \frac{\beta_1}{1-\beta_1} \right)^2 + \left( \frac{1}{1-\lambda} \right)^2 \right) L \frac{G_\infty^2}{\epsilon^{1.5}}, \\
C_3 &= 16L^2 \left( \frac{1}{1-\lambda} \right) \frac{G_\infty^2}{\epsilon^2}, \\
C_4 &= \frac{2}{\epsilon^{1.5}} \frac{1}{1-\lambda} \left( \lambda + \frac{\beta_1}{1-\beta_1} \right) G_\infty^2,
\end{aligned}$$

$$C_5 = \frac{2}{\epsilon^2} \frac{1}{1-\lambda} L \left( \frac{\beta_1}{1-\beta_1} \right)^2 G_\infty^2 + \frac{4}{\epsilon^2} \frac{\lambda}{1-\lambda} L G_\infty^2.$$

652 Substituting into  $Z_1 = \overline{X}_1$  completes the proof.

□

### 653 C Proof of Theorem 3

654 Under some assumptions stated in Corollary 2.1, we have that

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] &\leq C_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C_2 \frac{N}{T} + C_3 \frac{N^{1.5}}{T^{1.5}d^{0.5}} \\ &\quad + \left( C_4 \frac{1}{T\sqrt{N}} + C_5 \frac{1}{T^{1.5}d^{0.5}} \right) \mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] \end{aligned} \quad (34)$$

655 where  $\|\cdot\|_{abs}$  denotes the entry-wise  $L_1$  norm of a matrix (i.e.  $\|A\|_{abs} = \sum_{i,j} |A_{ij}|$ ) and  
656  $C_1, C_2, C_3, C_4, C_5$  are defined in Theorem 2.

657 Since Algorithm 3 is a special case of 2, building on result of Theorem 2, we just need to characterize  
658 the growth speed of  $\mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right]$  to prove convergence of Algorithm 3. By the  
659 update rule of Algorithm 3, we know  $\hat{V}_t$  is non decreasing and thus

$$\begin{aligned} \mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] &= \mathbb{E} \left[ \sum_{t=1}^T \sum_{i=1}^N \sum_{j=1}^d |-\hat{v}_{t-2,i,j} + \hat{v}_{t-1,i,j}| \right] \\ &= \mathbb{E} \left[ \sum_{t=1}^T \sum_{i=1}^N \sum_{j=1}^d (-\hat{v}_{t-2,i,j} + \hat{v}_{t-1,i,j}) \right] \\ &= \mathbb{E} \left[ \sum_{i=1}^N \sum_{j=1}^d (-\hat{v}_{-1,i,j} + \hat{v}_{T-1,i,j}) \right] \\ &= \mathbb{E} \left[ \sum_{i=1}^N \sum_{j=1}^d (-\hat{v}_{0,i,j} + \hat{v}_{T-1,i,j}) \right], \end{aligned}$$

660 where the last equality is because we defined  $\hat{V}_{-1} \triangleq \hat{V}_0$  previously.

661 Further, because  $\|g_{t,i}\|_\infty \leq G_\infty$  for all  $t, i$  and  $v_{t,i}$  is a exponential moving average of  $g_{k,i}^2, k =$   
662  $1, 2, \dots, t$ , we know  $|\hat{v}_{t,i,j}| \leq G_\infty^2$ , for all  $t, i, j$ . In addition, by update rule of  $\hat{V}_t$ , we also know  
663 each element of  $\hat{V}_t$  also cannot be greater than  $G_\infty^2$ , i.e.  $|\hat{v}_{t,i,j}| \leq G_\infty^2$ , for all  $t, i, j$ . Given the fact  
664 that  $\hat{v}_{0,i,j} \geq 0$ , we have

$$\mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] = \mathbb{E} \left[ \sum_{i=1}^N \sum_{j=1}^d (-\hat{v}_{0,i,j} + \hat{v}_{T-1,i,j}) \right] \leq \mathbb{E} \left[ \sum_{i=1}^N \sum_{j=1}^d G_\infty^2 \right] = NdG_\infty^2.$$

665 Substituting the above into (34), we have

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] &\leq C_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C_2 \frac{N}{T} + C_3 \frac{N^{1.5}}{T^{1.5}d^{0.5}} \\ &\quad + \left( C_4 \frac{1}{T\sqrt{N}} + C_5 \frac{1}{T^{1.5}d^{0.5}} \right) NdG_\infty^2 \\ &= C'_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C'_2 \frac{N}{T} + C'_3 \frac{N^{1.5}}{T^{1.5}d^{0.5}} \\ &\quad + C'_4 \frac{\sqrt{Nd}}{T} + C'_5 \frac{Nd^{0.5}}{T^{1.5}}, \end{aligned} \quad (35)$$

666 where we have

$$C'_1 = C_1 \quad C'_2 = C_2 \quad C'_3 = C_3 \quad C'_4 = C_4 G_\infty^2 \quad C'_5 = C_5 G_\infty^2. \quad (36)$$

667 and we conclude the proof.  $\square$

## 668 D Proof of Theorem 4

669 The proof follows the same flow as that of Theorem 3. Under assumptions stated in Corollary 2.1, set  
 670  $\alpha = \sqrt{N}/\sqrt{Td}$ , we have that

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] &\leq C_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C_2 \frac{N}{T} + C_3 \frac{N^{1.5}}{T^{1.5}d^{0.5}} \\ &\quad + \left( C_4 \frac{1}{T\sqrt{N}} + C_5 \frac{1}{T^{1.5}d^{0.5}} \right) \mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right], \end{aligned} \quad (37)$$

671 where  $\|\cdot\|_{abs}$  denotes the entry-wise  $L_1$  norm of a matrix (i.e.  $\|A\|_{abs} = \sum_{i,j} |A_{ij}|$ ) and  
 672  $C_1, C_2, C_3, C_4, C_5$  are defined in Theorem 2.

673 Again, Since decentralized AdaGrad is a special case of 2, we can apply Corollary 2.1 and what we  
 674 need is to upper bound  $\mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right]$  derive convergence rate. By the update rule  
 675 of decentralized AdaGrad, we have  $\hat{v}_{t,i} = \frac{1}{t} (\sum_{k=1}^t g_{k,i}^2)$  for  $t \geq 1$  and  $\hat{v}_{0,i} = \epsilon \mathbf{1}$ . Then we have for  
 676  $t \geq 3$ ,

$$\begin{aligned} &\mathbb{E} \left[ \sum_{t=1}^T \|(-\hat{V}_{t-2} + \hat{V}_{t-1})\|_{abs} \right] \\ &= \mathbb{E} \left[ \sum_{t=1}^T \sum_{i=1}^N \sum_{j=1}^d |-\hat{v}_{t-2,i} + \hat{v}_{t-1,i}| \right] \\ &\leq \mathbb{E} \left[ \sum_{t=3}^T \sum_{i=1}^N \sum_{j=1}^d \left| -\frac{1}{t-2} \left( \sum_{k=1}^{t-2} g_{k,i}^2 \right) + \frac{1}{t-1} \left( \sum_{k=1}^{t-1} g_{k,i}^2 \right) \right| \right] + Nd(G_\infty^2 - \epsilon) \\ &\leq \mathbb{E} \left[ \sum_{t=3}^T \sum_{i=1}^N \sum_{j=1}^d \left| \left( \frac{1}{t-1} - \frac{1}{t-2} \right) \left( \sum_{k=1}^{t-2} g_{k,i}^2 \right) + \frac{1}{t-1} g_{t-1,i}^2 \right| \right] + NdG_\infty^2 \\ &= \mathbb{E} \left[ \sum_{t=3}^T \sum_{i=1}^N \sum_{j=1}^d \left| \left( -\frac{1}{(t-1)(t-2)} \right) \left( \sum_{k=1}^{t-2} g_{k,i}^2 \right) + \frac{1}{t-1} g_{t-1,i}^2 \right| \right] + NdG_\infty^2 \\ &\leq \mathbb{E} \left[ \sum_{t=3}^T \sum_{i=1}^N \sum_{j=1}^d \max \left( \frac{1}{(t-1)(t-2)} \left( \sum_{k=1}^{t-2} g_{k,i}^2 \right), \frac{1}{t-1} g_{t-1,i}^2 \right) \right] + NdG_\infty^2 \\ &\leq \mathbb{E} \left[ Nd \sum_{t=3}^T \frac{G_\infty^2}{t-1} \right] + NdG_\infty^2 \\ &\leq NdG_\infty^2 \log(T) + NdG_\infty^2 \\ &= NdG_\infty^2 (\log(T) + 1) \end{aligned}$$

677 where the first equality is because we defined  $\hat{V}_{-1} \triangleq \hat{V}_0$  previously and  $\|g_{k,i}\|_\infty \leq G_\infty$  by assump-  
 678 tion.

679 Substituting the above into (37), we have

$$\begin{aligned}
\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[ \left\| \frac{\nabla f(\bar{X}_t)}{\bar{U}_t^{1/4}} \right\|^2 \right] &\leq C_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C_2 \frac{N}{T} + C_3 \frac{N^{1.5}}{T^{1.5} d^{0.5}} \\
&\quad + \left( C_4 \frac{1}{T\sqrt{N}} + C_5 \frac{1}{T^{1.5} d^{0.5}} \right) N d G_\infty^2 (\log(T) + 1) \\
&= C'_1 \frac{\sqrt{d}}{\sqrt{TN}} \left( (\mathbb{E}[f(Z_1)] - \min_x f(x)) + \sigma^2 \right) + C'_2 \frac{N}{T} + C'_3 \frac{N^{1.5}}{T^{1.5} d^{0.5}} \\
&\quad + C'_4 \frac{d\sqrt{N}(\log(T) + 1)}{T} + C'_5 \frac{(\log(T) + 1)N\sqrt{d}}{T^{1.5}},
\end{aligned}$$

680 where we have

$$C'_1 = C_1 \quad C'_2 = C_2 \quad C'_3 = C_3 \quad C'_4 = C_4 G_\infty^2 \quad C'_5 = C_5 G_\infty^2. \quad (38)$$

681 and we conclude the proof.  $\square$



## 682 E Additional Experiments and Details

683 In this section, we compare the training loss and testing accuracy of different algorithms, namely  
 684 Decentralized Stochastic Gradient Descent (D-PSGD), Decentralized Adam (DADAM) and our  
 685 proposed Decentralized AMSGrad, with different stepsizes on heterogeneous data distribution. We  
 686 use 5 nodes and the heterogeneous data distribution is created by assigning each node with data of  
 687 only two labels. Note that there are no overlapping labels between different nodes. For all algorithms,  
 688 we compare stepsizes in the grid  $[10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6}]$ .

689 Figure 3 shows the training loss and test accuracy for D-PSGD algorithm. We observe that the  
 690 stepsize  $10^{-3}$  works best for D-PSGD in terms of test accuracy and  $10^{-1}$  works best in terms of  
 691 training loss. This difference is caused by the inconsistency among the value of parameters on  
 692 different nodes when the stepsize is large. The training loss is calculated as the average of the loss  
 693 value of different local models evaluated on their local training batch. Thus, while the training loss is  
 694 small at a particular node, the test accuracy will be low when evaluating data with labels not seen by  
 695 the node (recall that each node contains data with different labels since we are in the heterogeneous  
 696 setting).




Figure 3: Performance comparison of different stepsizes for D-PSGD




Figure 4: Performance comparison of different stepsizes for decentralized AMSGrad




Figure 5: Performance comparison of different stepsizes for DADAM

697 Figure 4 shows the performance of decentralized AMSGrad with different stepsizes. We see that its  
698 best performance is better than the one of D-PSGD. Its performance is also more stable in the sense  
699 that the test performance is less sensitive to stepsize tuning according to our experiments.

700 Figure 5 displays the performance of Decentralized Adam algorithm. As expected, the performance  
701 of DADAM is not as good as D-PSGD or decentralized AMSGrad. Its divergence characteristic,  
702 highlighted Section 2.3, coupled with the heterogeneity in the data amplify its non-convergence issue  
703 in our experiments. From the experiments above, we can see the benefits of decentralized AMSGrad  
704 both in terms of performance and ease of parameter tuning, and the importance of ensuring the  
705 theoretical convergence of any newly proposed methods in the presented setting.