

---

# Fast Two-Time-Scale Noisy EM Algorithms

---

Anonymous Author(s)

Affiliation

Address

email

## Abstract

1 Training latent data models using the EM algorithm is the most common choice  
2 for current learning tasks. Variants of the EM to scale to large datasets and by-  
3 pass the impossible conditional expectation of the latent data for most nonlinear  
4 models have been initially introduced respectively by [Neal and Hinton, 1998],  
5 using incremental updates, and [Wei and Tanner, 1990, Delyon et al., 1999], using  
6 Monte-Carlo (MC) approximations. In this paper, we propose to combine those  
7 both techniques in a single class of methods called Two-Time-Scale EM Methods.  
8 We motivate the choice of a double dynamics by invoking the variance reduction  
9 virtue of each stage of the method on both noise: the incremental update and the  
10 MC approximation. We establish finite-time convergence bounds for nonconvex  
11 objective function and independent of the initialization. Numerical applications  
12 are also presented in this article to illustrate our findings.

## 1 Introduction

14 Learning latent data models is critical for modern machine learning problems, see [McLachlan and  
15 Krishnan, 2007] for references. We formulate the training of such model as the following empirical  
16 risk minimization problem:

$$\min_{\theta \in \Theta} \bar{L}(\theta) := r(\theta) + L(\theta) \quad \text{with} \quad L(\theta) = \frac{1}{n} \sum_{i=1}^n L_i(\theta) := \frac{1}{n} \sum_{i=1}^n \{ -\log g(y_i; \theta) \}, \quad (1)$$

17 We denote the observations by  $\{y_i\}_{i=1}^n$ ,  $\Theta \subset \mathbb{R}^d$  is the convex parameters space. We consider a  
18 regularized model where  $r : \Theta \rightarrow \mathbb{R}$  is a smooth convex regularization function and for  $\theta \in \Theta$ ,  
19  $g(y; \theta)$  is the (incomplete) likelihood of each individual observation. The objective function  $\bar{L}(\theta)$  is  
20 possibly *nonconvex* and is assumed to be lower bounded  $\bar{L}(\theta) > -\infty$  for all  $\theta \in \Theta$ .

21 In the latent variable model,  $g(y_i; \theta)$ , is the marginal of the complete data likelihood defined as  
22  $f(z_i, y_i; \theta)$ , i.e.  $g(y_i; \theta) = \int_{\mathcal{Z}} f(z_i, y_i; \theta) \mu(dz_i)$ , where  $\{z_i\}_{i=1}^n$  are the (unobserved) latent vari-  
23 ables. In this paper, we make the assumption of a complete model belonging to the curved expo-  
24 nential family, i.e.,

$$f(z_i, y_i; \theta) = h(z_i, y_i) \exp(\langle S(z_i, y_i) | \phi(\theta) \rangle - \psi(\theta)), \quad (2)$$

25 where  $\psi(\theta)$ ,  $h(z_i, y_i)$  are scalar functions,  $\phi(\theta) \in \mathbb{R}^k$  is a vector function, and  $S(z_i, y_i) \in \mathbb{R}^k$  is  
26 the complete data sufficient statistics.

27 Full batch EM [Dempster et al., 1977] is the method of reference for that kind of task and is a two  
28 steps procedure. The E-step amounts to computing the conditional expectation of the complete data  
29 sufficient statistics,

$$\bar{s}(\theta) = \frac{1}{n} \sum_{i=1}^n \bar{s}_i(\theta) \quad \text{where} \quad \bar{s}_i(\theta) = \int_{\mathcal{Z}} S(z_i, y_i) p(z_i | y_i; \theta) \mu(dz_i). \quad (3)$$

30 The M-step is given by

$$\text{M-step: } \hat{\theta} = \bar{\theta}(\bar{s}(\theta)) := \arg \min_{\vartheta \in \Theta} \{r(\vartheta) + \psi(\vartheta) - \langle \bar{s}(\theta) | \phi(\vartheta) \rangle\}, \quad (4)$$

31 Two caveats of this method are the following: (a) with the explosion of data, the first step of the EM  
 32 is computationally inefficient as it requires a full pass over the dataset at each iteration and (b) the  
 33 complexity of modern models makes the expectation intractable. So far, both challenges have been  
 34 addressed separately, to the best of our knowledge, and we give an overview of current solutions in  
 35 the sequel.

36 **Prior Work** Inspired by stochastic optimization procedures, [Neal and Hinton, 1998] and [Cappé  
 37 and Moulines, 2009] developed respectively an incremental and an online variant of the E-step in  
 38 models where the expectation is computable then extensively used and studied in [Nguyen et al.,  
 39 2020, Liang and Klein, 2009, Cappé, 2011]. Some improvements of that methods have been pro-  
 40 vided and analyzed, globally and in finite-time, in [Karimi et al., 2019] where variance reduction  
 41 techniques taken from the optimization literature have been efficiently applied to scale the EM algo-  
 42 rithm to large datasets.

43 Regarding the computation of the expectation under the posterior distribution, the first method was  
 44 the Monte-Carlo EM (MCEM) introduced in the seminal paper [Wei and Tanner, 1990] where a MC  
 45 approximation of this expectation is computed. A variant of that method is the Stochastic Approx-  
 46 imation of the EM (SAEM) in [Delyon et al., 1999] leveraging the power of Robbins-Monro type of  
 47 update [Robbins and Monro, 1951] to ensure pointwise convergence of the vector of estimated pa-  
 48 rameters rather using a decreasing stepsize than increasing the number of MC samples. The MCEM  
 49 and the SAEM have been successfully applied in mixed effects models [McCulloch, 1997, Hughes,  
 50 1999, Baey et al., 2016] or to do inference for joint modelling of time to event data coming from  
 51 clinical trials in [Chakraborty and Das, 2010], among other applications.

52 Recently, an incremental variant of the SAEM was proposed in [Kuhn et al., 2019] showing positive  
 53 empirical results but its analysis is limited to asymptotic consideration. Gradient-based methods  
 54 have been developed and analyzed in [Zhu et al., 2017] but they remain out of the scope of this  
 55 paper as they tackle the high-dimensionality issue.

56 **Contributions** This paper *introduces* and *analyzes* a new class of methods which purpose is to  
 57 combine both solutions proposed in the past years in a two-time-scale manner in order to optimize  
 58 (1) for current modern examples and settings. The main contributions of the paper are:

- 59 • We propose a two-time-scale method based on Stochastic Approximation (SA), to alleviate  
 60 the problem of MC computation, and on Incremental updates, to scale to large datasets.  
 61 We describe in details the edges of each level of our method based on variance reduc-  
 62 tion arguments. The derivation of such class of algorithms has two advantages. First, it  
 63 combines two powerful ideas, commonly used separately, to tackle large scale and highly  
 64 nonlinear learning tasks. Then, it gives a simple formulation as a *scaled-gradient method*,  
 65 as introduced in [Karimi et al., 2019], which makes the global analysis accessible.
- 66 • We also establish global (independent of the initialization) and finite-time (true at each  
 67 iteration) upper bounds on a classical suboptimality condition in the nonconvex literature,  
 68 *i.e.*, the second order moment of the gradient of the objective function.

69 In Section 2 we give rigorous mathematical definitions of the various updates used for both incre-  
 70 mental and Monte-Carlo EMs and we introduce the main class of new algorithms, based on two  
 71 different dynamics, we are proposing to analyze and compare to baselines algorithms. Section 3  
 72 presents the main theoretical guarantees of this newly introduced two-time-scale class of algorithms.  
 73 Results are given both in finite-time and in the nonconvex setting. Finally, we illustrate the advan-  
 74 tages of our method in Section 4 on two numerical experiments.

## 75 2 Two-Time-Scale Stochastic EM Algorithms

76 We recall and formalize in this section the different methods found in the literature that aim to solv-  
 77 ing the large scale problem and the intractable expectation. We then provide the general framework  
 78 of our method to efficiently tackle the optimization problem (1).

## 79 2.1 Monte Carlo Integration and Stochastic Approximation

80 As mentioned in the introduction, for complex and possibly nonlinear models, the expectation under  
 81 the posterior distribution defined in (3) is not tractable. In that case, the first solution involves  
 82 computing a Monte Carlo integration of that latter term. For all  $i \in \llbracket 1, n \rrbracket$ , draw for  $m \in \llbracket 1, M \rrbracket$ ,  
 83 samples  $z_{i,m} \sim p(z_i|y_i; \theta)$  and compute the MC integration  $\tilde{s}$  of the deterministic quantity  $\bar{s}(\theta)$ :

$$\text{MC-step : } \tilde{s} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}, y_i) \quad (5)$$

84 and then update the parameter  $\hat{\theta} = \bar{\theta}(\tilde{s})$ . This algorithm bypasses the intractable expectation issue  
 85 but is rather computationally expensive in order to reach point wise convergence ( $M$  needs to be  
 86 large). An alternative to that stochastic algorithm is to use a Robbins-Monro (RM) type of update.  
 87 We denote, at iteration  $k$ , the following quantity

$$\tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}^{(k)}, y_i) \quad \text{where } z_{i,m}^{(k)} \sim p(z_i|y_i; \theta^{(k)}) \quad (6)$$

88 Then, the RM updated of the sufficient statistics  $\hat{s}^{(k+1)}$  reads:

$$\text{SA-step : } \hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad (7)$$

89 where  $\{\gamma_k\}_{k \geq 1} \in (0, 1)$  is a sequence of decreasing step sizes to ensure asymptotic convergence.  
 90 This is called the Stochastic Approximation of the EM (SAEM) and has been shown theoretically to  
 91 converge to a maximum of the likelihood of the observations under very general conditions [Delyon  
 92 et al., 1999]. In the simulation step (6), since the relation between the observed data  $y_i$  and the  
 93 latent variable  $z_i$  can be non linear, sampling from the posterior distribution  $p(z_i|y_i; \theta)$ , under the  
 94 current model  $\theta$ , could require using an inference algorithm. [?] proved almost sure convergence  
 95 of the sequence of parameters obtained by this algorithm coupled with an MCMC procedure during  
 96 the simulation step. In simple scenarios, the samples  $\{z_{i,m}\}_{m=0}^{M-1}$  are conditionally independent and  
 97 identically distributed with distribution  $p(z_i, \theta)$ . Nevertheless, in most cases, sampling exactly from  
 98 this distribution is not an option and the Monte Carlo batch is sampled by Monte Carlo Markov  
 99 Chains (MCMC) algorithm. In the SA-step, the sequence of decreasing positive integers  $\{\gamma_k\}_{k \geq 1}$   
 100 controls the convergence of the algorithm. In practice,  $\gamma_k$  is set equal to 1 during the first few  
 101 iterations to let the algorithm explore the parameter space without memory and converge quickly  
 102 to a neighbourhood of the target estimate. The Stochastic Approximation is performed during the  
 103 remaining iterations where  $\gamma_k = 1/k^\alpha$ , where  $\alpha \in (0, 1)$ , ensuring the almost sure convergence of  
 104 the estimate. It is inappropriate to start with small values for step size  $\gamma_k$  and large values for the  
 105 number of simulations  $M_k$ . Rather, it is recommended that one decrease  $\gamma_k$  and keep a constant  
 106 and small number of MC samples  $M_k$  which shows a great advantage over the MC-step (5), which  
 107 requires large  $M_k$  to converge.

108 This Robbins-Monro type of update represents the *first level* of our algorithm, needed to temper  
 109 the variance and noise implied by MC integration. In the next section, we derive variants of this  
 110 algorithm to adapt to the sheer size of data of today's applications and formalize the *second level* of  
 111 our class of Two-Time-Scale EM methods.

## 112 2.2 Incremental and Bi-Level Inexact EM Methods

113 Strategies to scale to large datasets include classical incremental and variance reduced variants. We  
 114 will explicit a general update that will cover those variants and that represents the *second level* of our  
 115 algorithm, namely the incremental update of the noisy statistics  $\hat{S}^{(k)}$  inside the RM type of update.

$$\text{Incremental-step : } \tilde{S}^{(k+1)} = \tilde{S}^{(k)} + \rho_{k+1}(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}), \quad (8)$$

116 Note  $\{\rho_k\}_{k \geq 1} \in (0, 1)$  is a sequence of step sizes,  $\mathcal{S}^{(k)}$  is a proxy for  $\tilde{S}^{(k)}$ , If the stepsize is equal  
 117 to one and the proxy  $\mathcal{S}^{(k)} = \hat{S}^{(k)}$ , i.e., computed in a full batch manner as in (6), then we recover  
 118 the SAEM algorithm. Also if  $\rho_k = 1$ ,  $\gamma_k = 1$  and  $\mathcal{S}^{(k)} = \tilde{S}^{(k)}$ , then we recover the Monte Carlo  
 119 EM algorithm.

We now introduce three variants of the SAEM update depending on different definitions of the proxy  $\mathcal{S}^{(k)}$  and the choice of the stepsize  $\rho_k$ . Let  $i_k \in \llbracket 1, n \rrbracket$  be a random index drawn at iteration  $k$  and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$  be the iteration index where  $i \in \llbracket 1, n \rrbracket$  is last drawn prior to iteration  $k$ . For iteration  $k \geq 0$ , the fiTTSEM method draws *two* indices *independently* and uniformly as  $i_k, j_k \in \llbracket 1, n \rrbracket$ . In addition to  $\tau_i^k$  which was defined w.r.t.  $i_k$ , we define  $t_j^k = \{k' : j_{k'} = j, k' < k\}$  to be the iteration index where the sample  $j \in \llbracket 1, n \rrbracket$  is last drawn as  $j_k$  prior to iteration  $k$ . With the initialization  $\bar{\mathcal{S}}^{(0)} = \bar{s}^{(0)}$ , we use a slightly different update rule from SAGA inspired by [Reddi et al., 2016]. Then, we obtain:

$$(iSAEM [Karimi, 2019, Kuhn et al., 2019]) \quad \mathcal{S}^{(k+1)} = \mathcal{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)}) \quad (9)$$

$$(vrTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}) \quad (10)$$

$$(fiTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \quad (11)$$

$$\bar{\mathcal{S}}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + n^{-1} (\tilde{S}_{j_k}^{(k)} - \tilde{S}_{j_k}^{(t_{j_k}^k)}). \quad (12)$$

The stepsize is set to  $\rho_{k+1} = 1$  for the iSAEM method;  $\rho_{k+1} = \gamma$  is constant for the vrTTSEM and fiTTSEM methods. Moreover, for iSAEM we initialize with  $\mathcal{S}^{(0)} = \tilde{S}^{(0)}$ ; for vrTTSEM we set an epoch size of  $m$  and define  $\ell(k) := m \lfloor k/m \rfloor$  as the first iteration number in the epoch that iteration  $k$  is in.

### 2.3 Two-Time-Scale Noisy EM methods

We now introduce the general method derived using the two variance reduction techniques described above. Algorithm 1 leverages both levels (7) and (8) in order to output a vector of fitted parameters  $\hat{\theta}^{(K)}$  where  $K$  is some randomly chosen termination point.

The update in (14) is said to have two timescales as the step sizes satisfy  $\lim_{k \rightarrow \infty} \gamma_k / \rho_k < 1$  such that  $\tilde{S}^{(k+1)}$  is updated at a faster timescale than  $\hat{s}^{(k+1)}$ .

---

#### Algorithm 1 Two-Time-Scale Noisy EM methods.

---

- 1: **Input:** initializations  $\hat{\theta}^{(0)} \leftarrow 0, \hat{s}^{(0)} \leftarrow \hat{S}^{(0)}, K_{\max} \leftarrow \text{max. iteration number}$ .
- 2: Set the terminating iteration number,  $K \in \{0, \dots, K_{\max} - 1\}$ , as a discrete r.v. with:

$$P(K = k) = \frac{\gamma_k}{\sum_{\ell=0}^{K_{\max}-1} \gamma_{\ell}}. \quad (13)$$

- 3: **for**  $k = 0, 1, 2, \dots, K$  **do**
- 4: Draw index  $i_k \in \llbracket 1, n \rrbracket$  uniformly (and  $j_k \in \llbracket 1, n \rrbracket$  for fiTTSEM).
- 5: Compute  $\hat{S}_{i_k}^{(k)}$  using the MC-step (5), for the drawn indices.
- 6: Compute the surrogate sufficient statistics  $\mathcal{S}^{(k+1)}$  using (9) or (10) or (11).
- 7: Compute  $\hat{S}^{(k+1)}$  and  $\hat{s}^{(k+1)}$  using respectively (8) and (7):

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho_{k+1} (\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{s}^{(k+1)} &= \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \end{aligned} \quad (14)$$

- 8: Compute  $\hat{\theta}^{(k+1)}$  via the M-step (4).
  - 9: **end for**
  - 10: **Return:**  $\hat{\theta}^{(K)}$ .
- 

## 3 Global and Finite Time Analysis of the Scheme

First, we consider the following minimization problem on the statistics space:

$$\min_{\mathbf{s} \in \mathcal{S}} V(\mathbf{s}) := \bar{\mathcal{L}}(\bar{\theta}(\mathbf{s})) = \mathbf{r}(\bar{\theta}(\mathbf{s})) + \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(\bar{\theta}(\mathbf{s})) \quad (15)$$

140 It has been shown that this minimization problem is equivalent to the optimization problem (1), see  
 141 [Karimi et al., 2019, Lemma2]

142 **H1.**  $\Theta$  is an open set of  $\mathbb{R}^d$  and the sets  $Z, S$  are measurable open sets such that:

$$S \supset \left\{ n^{-1} \sum_{i=1}^n u_i, u_i \in \text{conv}(\bar{s}_i(\theta)) \right\} \quad (16)$$

143 where  $\bar{s}_i(\theta)$  is defined in (3).

144 **H2.** The conditional distribution is smooth on  $\text{int}(\Theta)$ . For any  $i \in \llbracket 1, n \rrbracket$ ,  $z \in Z$ ,  $\theta, \theta' \in \text{int}(\Theta)^2$ ,  
 145 we have  $|p(z|y_i; \theta) - p(z|y_i; \theta')| \leq L_p \|\theta - \theta'\|$ .

146 We also recall from the introduction that we consider curved exponential family models. besides:

147 **H3.** For any  $s \in S$ , the function  $\theta \mapsto L(s, \theta) := r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$  admits a unique global  
 148 minimum  $\bar{\theta}(s) \in \text{int}(\Theta)$ . In addition,  $J_\phi^\theta(\bar{\theta}(s))$  is full rank and  $\bar{\theta}(s)$  is  $L_\theta$ -Lipschitz.

149 Similar to [Karimi et al., 2019], we denote by  $H_L^\theta(s, \theta)$  the Hessian (w.r.t to  $\theta$  for a given value of  
 150  $s$ ) of the function  $\theta \mapsto L(s, \theta) = r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$ , and define

$$B(s) := J_\phi^\theta(\bar{\theta}(s)) \left( H_L^\theta(s, \bar{\theta}(s)) \right)^{-1} J_\phi^\theta(\bar{\theta}(s))^\top. \quad (17)$$

151 **H4.** It holds that  $v_{\max} := \sup_{s \in S} \|B(s)\| < \infty$  and  $0 < v_{\min} := \inf_{s \in S} \lambda_{\min}(B(s))$ . There exists  
 152 a constant  $L_B$  such that for all  $s, s' \in S^2$ , we have  $\|B(s) - B(s')\| \leq L_B \|s - s'\|$ .

153 We now formulate the main difference with the work done in [Karimi et al., 2019]. The class of  
 154 algorithms we develop in this paper are two time-scale where the first stage corresponds to the  
 155 variance reduction trick used in [Karimi et al., 2019] in order to accelerate incremental methods and  
 156 reduce the variance induced by the index sampling. The second stage is the Robbins-Monro type of  
 157 update that aims to reduce the variance induced by the MC approximations

158 Indeed the expectations (3) are never available and requires Monte Carlo approximation. Thus, at  
 159 iteration  $k + 1$ , we introduce the errors when approximating the quantity  $\bar{s}_i(\hat{\theta}(\hat{s}^{(k-1)}))$ . For all  
 160  $i \in \llbracket 1, n \rrbracket$ ,  $r > 0$  and  $\vartheta \in \Theta$ , define:

$$\eta_i^{(r)} := \tilde{S}_i^{(r)} - \bar{s}_i(\vartheta^{(r)}) \quad (18)$$

161 For instance, we consider that the MC approximation is unbiased if for all  $i \in \llbracket 1, n \rrbracket$  and  $m \in$   
 162  $\llbracket 1, M \rrbracket$ , the samples  $z_{i,m} \sim p(z_i|y_i; \theta)$  are i.i.d. under the posterior distribution, i.e.,  $\mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] = 0$   
 163 where  $\mathcal{F}_r$  is the filtration up to iteration  $r$ . The following results are derived under the assumption  
 164 of control of the fluctuations implied by the approximation stated as follows:

165 **H5.** There exist a positive sequence of MC batch size  $\{M_r\}_{r>0}$  and constants  $(C, C_\eta)$  such that for  
 166 all  $k > 0$ ,  $i \in \llbracket 1, n \rrbracket$  and  $\vartheta \in \Theta$ :

$$\mathbb{E} \left[ \left\| \eta_i^{(r)} \right\|^2 \right] \leq \frac{C_\eta}{M_r} \quad \text{and} \quad \mathbb{E} \left[ \left\| \mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] \right\|^2 \right] \leq \frac{C}{M_r} \quad (19)$$

167 In that setting, we can prove two important results on the Lyapunov function. The first one suggests  
 168 smoothness:

169 **Lemma 1.** [Karimi et al., 2019] Assume H2, H3, H4. For all  $s, s' \in S$  and  $i \in \llbracket 1, n \rrbracket$ , we have

$$\|\bar{s}_i(\bar{\theta}(s)) - \bar{s}_i(\bar{\theta}(s'))\| \leq L_s \|s - s'\|, \quad \|\nabla V(s) - \nabla V(s')\| \leq L_V \|s - s'\|, \quad (20)$$

170 where  $L_s := C_Z L_p L_\theta$  and  $L_V := v_{\max}(1 + L_s) + L_B C_S$ .

171 and the second one suggests a growth condition on the gradient of  $V$  depending on the mean field  
 172 of the algorithm:

173 **Lemma 2.** Assume H3, H4. For all  $s \in S$ ,

$$v_{\min}^{-1} \langle \nabla V(s) | s - \bar{s}(\bar{\theta}(s)) \rangle \geq \|s - \bar{s}(\bar{\theta}(s))\|^2 \geq v_{\max}^{-2} \|\nabla V(s)\|^2, \quad (21)$$

174 See proofs of this Lemma in Appendix A.

### 3.1 Global Convergence of Incremental Noisy EM Algorithms

Following the asymptotic analysis of update (9), we present a finite-time analysis of the incremental variant of the Stochastic Approximation of the EM algorithm.

The first intermediate result is the computation of the quantity  $\hat{S}^{(k+1)} - \hat{s}^{(k)}$ , which corresponds to the drift term of (7) and reads as follows:

**Lemma 3.** Assume H1. The update (9) is equivalent to the following update on the resulting statistics

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad \text{where} \quad \tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^{k+1})} \quad (22)$$

Also:

$$\mathbb{E} [\tilde{S}^{(k+1)} - \hat{s}^{(k)}] = \mathbb{E} [\bar{s}^{(k)} - \hat{s}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \quad (23)$$

where  $\bar{s}^{(k)}$  is defined by (3) and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ .

See proofs of this Lemma in Appendix B.

The following main result for the iSAEM algorithm is derived under a control of the Monte Carlo fluctuations as described by assumption H 5. Typically, the controls exhibited below are of interest when the number of MC samples  $M_k$  increase with the iteration index  $f$ .

**Theorem 1.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and consider the iSAEM sequence  $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ . We also set  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  $\hat{s}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ .

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] \leq \mathbb{E} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K)})] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (24)$$

See proof in Appendix C.

### 3.2 Global Convergence of Two-Time-Scale Noisy EM Algorithms

We now proceed by giving our main result regarding the global convergence of the fTTSEM algorithm. Two important auxiliary Lemmas, which proofs are given in Appendix D, are need in order to derive our finite-time bound. The first one derives an identity for the quantity  $\hat{S}^{(k+1)} - \hat{s}^{(k)}$  where  $\hat{S}^{(k+1)}$  is computed using the fTTSEM update:

**Lemma 4.** Assume H1. At iteration  $k+1$ , the drift term of update (11), with  $\rho_{k+1} = \rho$ , is equivalent to the following :

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{s}^{(k)} &= \rho(\bar{s}^{(k)} - \hat{s}^{(k)}) + \rho \eta_{i_k}^{(k+1)} + \rho \left[ (\bar{s}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{s}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) (\tilde{S}^{(k)} - \hat{s}^{(k)}) \end{aligned} \quad (25)$$

where we recall that  $\eta_{i_k}^{(k+1)}$ , defined in (19), is the gap between the MC approximation and the expected statistics.

The second Lemma characterizes the evolution of the quantity  $\mathbb{E} [\left\| \hat{s}^{(k)} - \tilde{S}^{(k)} \right\|^2]$ . Remark that this term is the price we pay for the two time scale dynamics and corresponds to the gap between the two asynchronous updates (one is on  $\hat{s}^{(k)}$  and the other on  $\tilde{S}^{(k)}$ ).

**Lemma 5.** Consider a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho$ , then the following inequality holds:

$$\mathbb{E} [\left\| \tilde{S}^{(k)} - \hat{s}^{(k)} \right\|^2] \leq \frac{\rho}{1 - \rho} \sum_{\ell=0}^k (1 - \gamma_{\ell})^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}) \quad (26)$$



207 where  $\mathcal{S}^{(k)}$  is defined either by (11) (fTTSEM) or (10) (vrTTSEM)

**Remark** We understand from this Lemma that, if  $\rho = 1$ , i.e., no variance reduction, then

$$\mathbb{E}[\|\tilde{S}^{(k)} - \hat{s}^{(k)}\|^2] = \mathbb{E}[\|\mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)}\|^2] = 0$$

208 which strengthen the fact that this quantity is the impact of the variance reduction technique intro-  
209 duced in our two stages class of methods.

210 We now state the main result regarding the vrTTSEM method.

211 **Theorem 2.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes  
212 and consider the vrTTSEM sequence  $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for any  $k > 0$ .

213 Assume that  $\hat{s}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\bar{L} = \max\{L_S, L_V\}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$   
214 and a constant  $\mu \in (0, 1)$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] &\leq \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3} \bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}] \end{aligned} \quad (27)$$

215 See proof in Appendix F. We now state the main result regarding the fTTSEM method.

216 **Theorem 3.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes  
217 and consider the fTTSEM sequence  $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for any  $k > 0$ .

218 Assume that  $\hat{s}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_S, L_V\}$ ,  
219  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[ \|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2 \right] \right] \end{aligned} \quad (28)$$

220 See proof in Appendix E. The authors would like to stress the fact that a very similar result can be  
221 established for the vrTTSEM method but is not presented here for the sake of space and clarity. The  
222 Numerical examples in the following section outlines those similarities.

## 223 4 Numerical Examples

### 224 4.1 Gaussian Mixture Models

225 Given  $n$  observations  $\{y_i\}_{i=1}^n$ , we want to fit a Gaussian Mixture Model (GMM) whose distribution  
226 is modeled as a Gaussian mixture of  $M$  components, each with a unit variance. Let  $z_i \in \llbracket M \rrbracket$  be  
227 the latent labels of each component, the complete log-likelihood is defined as:

$$\log f(z_i, y_i; \theta) = \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) [\log(\omega_m) - \mu_m^2/2] + \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) \mu_m y_i + \text{constant} . \quad (29)$$

228 where  $\theta := (\omega, \mu)$  with  $\omega = \{\omega_m\}_{m=1}^{M-1}$  are the mixing weights with the convention  $\omega_M =$   
229  $1 - \sum_{m=1}^{M-1} \omega_m$  and  $\mu = \{\mu_m\}_{m=1}^M$  are the means. We use the penalization  $r(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 -$   
230  $\log \text{Dir}(\omega; M, \epsilon)$  where  $\delta > 0$  and  $\text{Dir}(\cdot; M, \epsilon)$  is the  $M$  dimensional symmetric Dirichlet distribu-  
231 tion with concentration parameter  $\epsilon > 0$ . The constraint set on  $\theta$  is given by

$$\Theta = \{\omega_m, m = 1, \dots, M-1 : \omega_m \geq 0, \sum_{m=1}^{M-1} \omega_m \leq 1\} \times \{\mu_m \in \mathbb{R}, m = 1, \dots, M\}. \quad (30)$$

232 Exact two time scale updates are given in Appendix G.1.

In the following experiments on synthetic data, we generate samples from a GMM model with  $M = 2$  components with two mixtures with means  $\mu_1 = -\mu_2 = 0.5$ . We use  $n = 10^5$  synthetic samples and run the bEM method until convergence (to double precision) to obtain the ML estimate  $\mu^*$  averaged on 50 datasets. We compare the bEM, SAEM, iSAEM, vrTTSEM and fiTTSEM methods in terms of their precision measured by  $|\mu - \mu^*|^2$ . We set the step-size of the SA-step of all method as  $\gamma_k = 1/k^\alpha$  with  $\alpha = 0.5$ , and the stepsizes of the Incremental-step for vrTTSEM and the fiTTSEM to a constant stepsize equal to  $1/n^{2/3}$ .

The number of MC samples is fixed to  $M = 10$  chains. Figure 1 shows the convergence of the precision  $|\mu - \mu^*|^2$  for the different methods against the epoch(s) elapsed (one epoch equals  $n$  iterations). We observe that the vrTTSEM and fiTTSEM methods outperform the other methods, supporting our analytical results.




Figure 1: TO COMPLETE

## 4.2 Deformable Template Model for Image Analysis

Let  $(y_i, i \in \llbracket 1, n \rrbracket)$  be observed gray level images defined on a grid of pixels. Let  $u \in \mathcal{U} \subset \mathbb{R}^2$  denotes the pixel index on the image and  $x_u \in \mathcal{D} \subset \mathbb{R}^2$  its location. The model used in this experiment suggests that each image  $y_i$  is a deformation of a template, noted  $I : \mathcal{D} \rightarrow \mathbb{R}$ , common to all images of the dataset:

$$y_i(u) = I(x_u - \Phi_i(x_u, z_i)) + \varepsilon_i(u) \quad (31)$$

where  $\phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a deformation function,  $z_i$  some latent variable parametrizing this deformation and  $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$  is an observation error.

The template model, given  $(p_k, k \in \llbracket 1, k_p \rrbracket)$  landmarks on the template, a fixed known kernel  $\mathbf{K}_p$  and a vector of parameters  $\beta \in \mathbb{R}^{k_p}$  is defined as follows:

$$I_\xi = \mathbf{K}_p \beta, \quad \text{where} \quad (\mathbf{K}_p \beta)(x) = \sum_{k=1}^{k_p} \mathbf{K}_p(x, p_k) \beta_k \quad (32)$$

Besides, we parameterize the deformation model given some landmarks  $(g_k, k \in \llbracket 1, k_g \rrbracket)$  and a fixed kernel  $\mathbf{K}_g$  as:

$$\Phi_i = \mathbf{K}_g z_i \quad \text{where} \quad (\mathbf{K}_g z_i)(x) = \sum_{k=1}^{k_g} \mathbf{K}_g(x, g_k) \left( z_i^{(1)}(k), z_i^{(2)}(k) \right) \quad (33)$$

where we put a Gaussian prior on the latent variables,  $z_i \sim \mathcal{N}(0, \Gamma)$  and  $z_i \in (\mathbb{R}^{k_g})^2$ . The vector of parameters we ought to estimate is thus  $\theta = (\beta, \Gamma, \sigma)$ . The complete model belongs to the curved exponential family, see [Allasonnière et al., 2007], which vector of sufficient statistics  $S = (S_1(z), S_2(z), S_3(z))$  read:

$$\begin{aligned} S_1(z) &= \sum_{i=1}^n S_1(y_i, z_i) = \sum_{i=1}^n (\mathbf{K}_p^{z_i})^t y_i \\ S_2(z) &= \sum_{i=1}^n S_2(y_i, z_i) = \sum_{i=1}^n (\mathbf{K}_p^{z_i})^t (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \sum_{i=1}^n S_3(y_i, z_i) = \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (34)$$

where for any pixel  $u \in \mathbb{R}^2$  and  $j \in \llbracket 1, k_g \rrbracket$  we noted:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j) \quad (35)$$



265 Finally, the Two-Time-Scale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z)\hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n}\hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z)\beta(\hat{s}) - 2\beta(\hat{s})\hat{s}_1(z) \end{pmatrix} \quad (36)$$

266 where  $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$  is the vector of statistics obtained via the **SA-step (7)** and using the  
 267 MC approximation of the sufficient statistics  $(S_1(z), S_2(z), S_3(z))$  defined in **(34)**.

268 **Comparison using epochs credit**

269 **Comparison using number of training samples credit**

## 270 **5 Conclusion**

## References

- S. Allasonnière, Y. Amit, and A. Trouvé. Towards a coherent statistical framework for dense deformable template estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):3–29, 2007.
- S. Allasonnière, E. Kuhn, A. Trouvé, et al. Construction of bayesian deformable models via a stochastic approximation algorithm: a convergence study. *Bernoulli*, 16(3):641–678, 2010.
- C. Baey, S. Trevezas, and P.-H. Cournède. A non linear mixed effects model of plant growth and estimation via stochastic variants of the em algorithm. *Communications in Statistics-Theory and Methods*, 45(6):1643–1669, 2016.
- O. Cappé. Online em algorithm for hidden markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- O. Cappé and E. Moulines. On-line expectation–maximization algorithm for latent data models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):593–613, 2009.
- A. Chakraborty and K. Das. Inferences for joint modelling of repeated ordinal scores and time to event data. *Computational and mathematical methods in medicine*, 11(3):281–295, 2010.
- B. Delyon, M. Lavielle, and E. Moulines. Convergence of a stochastic approximation version of the em algorithm. *Ann. Statist.*, 27(1):94–128, 03 1999. doi: 10.1214/aos/1018031103. URL <https://doi.org/10.1214/aos/1018031103>.
- A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the royal statistical society. Series B (methodological)*, pages 1–38, 1977.
- J. P. Hughes. Mixed effects models with censored data with application to hiv rna levels. *Biometrics*, 55(2):625–629, 1999.
- B. Karimi. *Non-Convex Optimization for Latent Data Models: Algorithms, Analysis and Applications*. PhD thesis, 2019.
- B. Karimi, H.-T. Wai, É. Moulines, and M. Lavielle. On the global convergence of (fast) incremental expectation maximization methods. In *Advances in Neural Information Processing Systems*, pages 2833–2843, 2019.
- E. Kuhn, C. Matias, and T. Rebafka. Properties of the stochastic approximation em algorithm with mini-batch sampling. *arXiv preprint arXiv:1907.09164*, 2019.
- P. Liang and D. Klein. Online em for unsupervised models. In *Proceedings of human language technologies: The 2009 annual conference of the North American chapter of the association for computational linguistics*, pages 611–619, 2009.
- C. E. McCulloch. Maximum likelihood algorithms for generalized linear mixed models. *Journal of the American statistical Association*, 92(437):162–170, 1997.
- G. McLachlan and T. Krishnan. *The EM algorithm and extensions*, volume 382. John Wiley & Sons, 2007.
- R. M. Neal and G. E. Hinton. A view of the EM algorithm that justifies incremental, sparse, and other variants. In *Learning in graphical models*, pages 355–368. Springer, 1998.
- H. D. Nguyen, F. Forbes, and G. J. McLachlan. Mini-batch learning of exponential family finite mixture models. *Statistics and Computing*, pages 1–18, 2020.

- 312 S. J. Reddi, S. Sra, B. Póczos, and A. Smola. Fast incremental method for nonconvex optimization.  
313 *arXiv preprint arXiv:1603.06159*, 2016.
- 314 H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statis-*  
315 *tics*, pages 400–407, 1951.
- 316 G. C. Wei and M. A. Tanner. A monte carlo implementation of the em algorithm and the poor man’s  
317 data augmentation algorithms. *Journal of the American statistical Association*, 85(411):699–704,  
318 1990.
- 319 R. Zhu, L. Wang, C. Zhai, and Q. Gu. High-dimensional variance-reduced stochastic gradient  
320 expectation-maximization algorithm. In *Proceedings of the 34th International Conference on*  
321 *Machine Learning-Volume 70*, pages 4180–4188. JMLR. org, 2017.

## A Proof of Lemma 2

**Lemma.** Assume H3, H4. For all  $\mathbf{s} \in \mathcal{S}$ ,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (37)$$

**Proof** Using H3 and the fact that we can exchange integration with differentiation and the Fisher's identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \left( \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \left( \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (38)$$

Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top \mathbf{s}. \quad (39)$$

Taking the gradient of the above map w.r.t.  $\mathbf{s}$  and using assumption H3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left( \nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})}}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}). \quad (40)$$

The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))) \quad (41)$$

where we recall  $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left( \mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top$ . The proof of (37) follows directly from the assumption H4.  $\square$

## B Proof of Lemma 3

**Lemma.** Assume H1. The update (9) is equivalent to the following update on the resulting statistics

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}) \quad (42)$$

Also:

$$\mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \quad (43)$$

where  $\bar{\mathbf{s}}^{(k)}$  is defined by (3) and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ .

**Proof** From update (9), we have:

$$\begin{aligned} \tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left( \tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned} \quad (44)$$

Since  $\tilde{S}_{i_k}^{(k+1)} = \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) + \eta_{i_k}^{(k+1)}$  we have

$$\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)} \quad (45)$$

Taking the full expectation of both side of the equation leads to:

$$\begin{aligned} \mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] &= \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] \\ &\quad - \frac{1}{n} \mathbb{E} \left[ \mathbb{E} [\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \end{aligned} \quad (46)$$

The following equalities:

$$\mathbb{E} [\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E} [\bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] = \bar{\mathbf{s}}^{(k)} \quad (47)$$

concludes the proof of the Lemma.  $\square$

## 341 C Proof of Theorem 1

342 **Theorem.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and  
 343 consider the iSAEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ . We also set  
 344  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  
 345  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ .

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (48)$$

346 **Proof** We begin our proof by giving this auxiliary Lemma setting an upper bound for the quantity  
 347  $\mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$

348 **Lemma 6.** For any  $k \geq 0$  and consider the iSAEM update in (9), it holds that

$$\begin{aligned} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &\leq 4\mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \frac{2L_{\mathbf{s}}^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right] \\ &\quad + 2\frac{C_{\eta}}{M_k} + 4\mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \end{aligned} \quad (49)$$

349 **Proof** Applying the iSAEM update yields:

$$\begin{aligned} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &= \mathbb{E} \left[ \left\| \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} - \frac{1}{n} (\tilde{S}_{i_k}^{(\tau_i^k)} - \tilde{S}_{i_k}^{(k)}) \right\|^2 \right] \\ &\leq 4\mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + 4\mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\quad + \frac{2}{n^2} \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] + 2\frac{C_{\eta}}{M_k} \end{aligned} \quad (50)$$

350 The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)} \right\|^2 \right] \stackrel{(a)}{\leq} \frac{2L_{\mathbf{s}}^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right], \quad (51)$$

351 where (a) is due to Lemma 1 and which concludes the proof of the Lemma.

352 □

353 Under the smoothness of the Lyapunov function  $V$  (cf. Lemma 1), we can write:

$$V(\hat{\mathbf{s}}^{(k+1)}) \leq V(\hat{\mathbf{s}}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \quad (52)$$

354 Taking the expectation on both sides yields:

$$\mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k+1)}) \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) \right] + \gamma_{k+1} \mathbb{E} \left[ \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (53)$$

355 Using Lemma 3, we obtain:

$$\begin{aligned}
& \mathbb{E} \left[ \langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] = \\
& \mathbb{E} \left[ \langle \bar{s}^{(k)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] + \left( 1 - \frac{1}{n} \right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \left( 1 - \frac{1}{n} \right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[ \left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& \stackrel{(a)}{\leq} \left( v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{54}$$

356 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with  $\beta \rightarrow 1$ ). Note

357  $a_k = \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$  and

$$\begin{aligned}
a_k \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{55}$$

358 We now give an upper bound of  $\mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right]$  using Lemma 6 and plug it into (55):

$$\begin{aligned}
(a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] \\
& + \gamma_{k+1} \left( \frac{1}{2\beta} \left( 1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right]
\end{aligned} \tag{56}$$

359 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^{k+1})} \right\|^2 \right] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{n-1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \right) \tag{57}$$

360 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. For any  $\beta > 0$ , it holds

$$\begin{aligned}
& \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 - 2\gamma_{k+1}\langle \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2 + \gamma_{k+1}\beta\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right]
\end{aligned} \tag{58}$$

361 where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2\right]
\end{aligned} \tag{59}$$

362 Observe that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)})$ . Applying Lemma 6 yields

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\
&\leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\
&\quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\
&\quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right]
\end{aligned} \tag{60}$$

363 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] \tag{61}$$

364 From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} &\leq (1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}))\Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
&\quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right]
\end{aligned} \tag{62}$$

365 Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,

366  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$ ,  $\alpha \geq 8$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_{\mathbf{s}}^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1} \tag{63}$$



367 which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} =$   
 368  $\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} \leq & 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\eta_{i_\ell}^{(\ell)}\|^2] \\ & + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)}\right\|^2\right] \end{aligned} \quad (64)$$

369 Note  $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$  Summing on both sides over  $k = 0$  to  $k = K_{\max} - 1$  yields:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} \\ &= 4 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}[\|\eta_{i_\ell}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\ &\leq \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}[\|\eta_{i_\ell}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \end{aligned} \quad (65)$$

370 We recall (56) where we have summed on both sides from  $k = 0$  to  $k = K_{\max} - 1$ :

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)})] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\frac{1}{2\beta}(1 - \frac{1}{n}) + 2\gamma_{k+1} L_V\right) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n}\right) \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)} \end{aligned} \quad (66)$$

371 Plugging (65) into (66) results in:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \tilde{\alpha}_k \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \tilde{\beta}_k \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)})] \\ &+ \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2] \end{aligned} \quad (67)$$

372 where:

$$\begin{aligned}\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\beta}_k &= \gamma_{k+1} \left( \frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\Gamma}_k &= \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}}\end{aligned}$$

373 and

$$\begin{aligned}a_k &= \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1) + 1}{2n} \right) \\ \Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1} L_s^2}{n^2} (\gamma_{k+1} + \frac{1}{\beta}) \\ c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{L_s, L_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}\end{aligned}$$

374 When, for any  $k > 0$ ,  $\tilde{\alpha}_k \geq 0$ , we have by Lemma 2 that:

$$\sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (68)$$

375 which yields an upper bound of the gradient of the Lyapunov function  $V$  along the path of the  
376 iSAEM update and concludes the proof of the Theorem.  $\square$

## 377 D Proof of Lemmas 4 and Lemma 5

378 **Lemma.** Assume H1. At iteration  $k + 1$ , the drift term of update (11), with  $\rho_{k+1} = \rho$ , is equivalent  
379 to the following :

$$\begin{aligned} \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) (\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \end{aligned} \quad (69)$$

380 where we recall that  $\eta_{i_k}^{(k+1)}$ , defined in (19), which is the gap between the MC approximation and  
381 the expected statistics.

382 **Proof** Using the fiTTSEM update  $\tilde{S}^{(k+1)} = (1 - \rho)\tilde{S}^{(k)} + \rho\mathcal{S}^{(k+1)}$  where  $\mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$  leads to the following decomposition:

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= (1 - \rho)\tilde{S}^{(k)} + \rho \left( \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) - \hat{\mathbf{s}}^{(k)} + \rho\bar{\mathbf{s}}^{(k)} - \rho\bar{\mathbf{s}}^{(k)} \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho(\tilde{S}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(k)}) + (1 - \rho) (\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho \left( \bar{\mathcal{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} + (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) (\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}) \end{aligned}$$

384 where we observe that  $\mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] = \bar{\mathbf{s}}^{(k)} - \bar{\mathcal{S}}^{(k)}$  and which concludes the proof.

385 *Important Note:* Note that  $\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not equal to  $\eta_{i_k}^{(k+1)}$ , defined in (19), which is the gap  
386 between the MC approximation and the expected statistics. Indeed  $\tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not computed under the  
387 same model as  $\bar{\mathbf{s}}_{i_k}^{(k)}$ .  $\square$

388 **Lemma.** Consider a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho$ , then the following inequality  
389 holds:

$$\mathbb{E}[\|\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \leq \frac{\rho}{1 - \rho} \sum_{\ell=0}^k (1 - \gamma_\ell)^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}) \quad (70)$$

390 where  $\mathcal{S}^{(k)}$  is defined either by (11) (fiTTSEM) or (10) (vrTTSEM)

391 **Proof** We begin by writing the two-time-scale update:

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}) \end{aligned} \quad (71)$$

392 where  $\mathcal{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(t_i^k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$  according to (11). Denote  $\delta^{(k+1)} = \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k+1)}$ . Then from (71), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1 - \rho} (1 - \gamma_{k+1})(\mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)}) \quad (72)$$

394 Using the telescoping sum and noting that  $\delta^{(0)} = 0$ , we have

$$\delta^{(k+1)} \leq \frac{\rho}{1 - \rho} \sum_{\ell=0}^k (1 - \gamma_{\ell+1})^2 (\mathcal{S}^{(\ell+1)} - \tilde{S}^{(\ell+1)}) \quad (73)$$

395  $\square$

## E Proof of Theorem 3

**Theorem.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and consider the fTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for any  $k > 0$ .

Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (74)$$

**Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful throughout this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) \\ &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1 - \rho)\tilde{S}^{(k)} - \rho \mathcal{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left( (1 - \rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)} \right] \right) \\ &= -\gamma_{k+1} \left( (1 - \rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \bar{\mathcal{S}}^{(k)} - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right] \right) \end{aligned} \quad (75)$$

We begin our proof by giving this auxiliary Lemma setting an upper bound for the quantity  $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2]$ .

**Lemma 7.** For any  $k \geq 0$  and consider the fTTSEM update in (11) with  $\rho_k = \rho$ , it holds for all  $k > 0$

$$\begin{aligned} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_{\mathbf{s}}^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (76)$$

**Proof** We observe, using the identity (75), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)}\|^2] + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \quad (77)$$

For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \mathcal{S}^{(k+1)}\|^2] &= \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(k)}) - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right\|^2 \right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (78)$$

where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_{\mathbf{s}}^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \quad (79)$$

Substituting into (77) proves the lemma.  $\square$

Using the smoothness of  $V$  and update (11), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (80)$$

412 Denote  $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fTTSEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 413 Using Lemma 4 and the additional following identity:

$$\mathbb{E} \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] = 0 \quad (81)$$

414 we have:

$$\begin{aligned} & \mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ & \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho\mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}\mathbb{E} \left[ \langle \rho\mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1-\rho)\mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ & \quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ & \stackrel{(a)}{\leq} -v_{\min}\gamma_{k+1}\rho\mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1}\mathbb{E} \left[ \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] - \frac{\gamma_{k+1}\rho^2}{2}\xi^{(k+1)} - \frac{\gamma_{k+1}(1-\rho)^2}{2}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ & \quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ & \stackrel{(b)}{\leq} -(v_{\min}\gamma_{k+1}\rho + \gamma_{k+1}v_{\max}^2)\mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1}\rho^2}{2}\xi^{(k+1)} - \frac{\gamma_{k+1}(1-\rho)^2}{2}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ & \quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \end{aligned} \quad (82)$$

415 where  $\xi^{(k+1)} = \mathbb{E} \left[ \|\mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k]\|^2 \right]$ .

416 **Bounding  $\mathbb{E}[\|\mathbf{H}_{k+1}\|^2]$**  Using Lemma 7, we obtain:

$$\begin{aligned} & \gamma_{k+1}(v_{\min}\rho + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V)\mathbb{E}[\|\mathbf{h}_k\|^2] \\ & \leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)})] + \tilde{\xi}^{(k+1)} + \left( (1-\rho)^2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ & \quad + \frac{\gamma_{k+1}^2 L_V \rho^2 L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (83)$$

417 where  $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 L_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1}\rho^2}{2}\xi^{(k+1)}$ . Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (84)$$

418 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ & = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \end{aligned} \quad (85)$$

419 Note that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}\mathbf{H}_{k+1}$  and that in expectation we recall  
 420 that  $\mathbb{E}[\mathbf{H}_{k+1} | \mathcal{F}_k] = \rho\mathbf{h}_k + \rho\mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}]$  where  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ . Thus,  
 421 for any  $\beta > 0$ , it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ & = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ & \leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ & \quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (86)$$

where the last inequality is due to the Young's inequality. Plugging this into (84) yields:

$$\begin{aligned}
& \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle\right] \\
&\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\right. \\
&\quad \left.+ \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right]\right]
\end{aligned} \tag{87}$$

Subsequently, we have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\
&\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2\right. \\
&\quad \left.+ \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right]\right]
\end{aligned} \tag{88}$$

We now use Lemma 7 on  $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2 \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2$  and obtain:

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\
&\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}\left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2\right] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2\rho^2 L_s^2}{n} + \frac{(n-1)(1 + \gamma_{k+1}\beta)}{n^2}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right] \\
&\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\
&\leq \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}\left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2\right] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2}{n}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right] \\
&\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{89}$$

Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \tag{90}$$

From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} &\leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}\left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2\right] \\
&\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{91}$$

Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{\frac{4}{3}}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1} \tag{92}$$

429 which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2\rho^2 L_s^2 \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} = \frac{1}{n} -$   
 430  $\gamma_{k+1}\beta - \gamma_{k+1}^2\rho^2 L_s^2$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} &\leq \sum_{\ell=0}^k \omega_{k,\ell} \left( 2\gamma_{\ell+1}^2\rho^2 + \frac{\gamma_{\ell+1}^2\rho^2}{\beta} \right) \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)} \right\|^2 \right] \\ &\quad + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left( 2\gamma_{\ell+1} + \frac{1}{\beta} \right) \mathbb{E} \left[ \left\| \tilde{S}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)} \right\|^2 \right] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)} \end{aligned} \quad (93)$$

431 where  $\omega_{k,\ell} = \prod_{j=\ell+1}^k \left( 1 - \Lambda_{(j)} \right)$  and  $\tilde{\epsilon}^{(\ell+1)} = \left( 2\gamma_{k+1} + \frac{\rho^2}{\beta} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k+1)} \right\|^2 \right]$ .

432 Summing on both sides over  $k = 0$  to  $k = K_{\max} - 1$  yields:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} &\leq \sum_{k=0}^{K_{\max}-1} \frac{2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}^2\rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\quad + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}(1-\rho)^2 \left( 2\gamma_{k+1} + \frac{1}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)} \end{aligned} \quad (94)$$

433 We recall (83) where we have summed on both sides from  $k = 0$  to  $k = K_{\max} - 1$ :

$$\begin{aligned} &\mathbb{E} [V(\hat{\mathbf{s}}^{(K_{\max})}) - V(\hat{\mathbf{s}}^{(0)})] \\ &\leq \sum_{k=0}^{K_{\max}-1} \left\{ \gamma_{k+1}(-v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V \mathbb{E} \left[ \left\| \mathbf{h}_k \right\|^2 \right] + \gamma^2 L_V \rho^2 L_s^2 \Delta^{(k)} \right\} \\ &\quad + \sum_{k=0}^{K_{\max}-1} \left\{ \tilde{\epsilon}^{(k+1)} + \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right\} \\ &\leq \sum_{k=0}^{K_{\max}-1} \left\{ \left[ -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_s^2 \left( 2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right] \mathbb{E} \left[ \left\| \mathbf{h}_k \right\|^2 \right] \right\} \\ &\quad + \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned} \quad (95)$$

where

$$\Xi^{(k+1)} = \tilde{\epsilon}^{(k+1)} + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}$$

and

$$\Gamma_{k+1} = \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2 (1-\rho)^2 \left( 2\gamma_{k+1} + \frac{1}{\beta} \right)}{\Lambda_{(k+1)}}$$

434 We now analyse the following quantity

$$\begin{aligned} &-\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2\rho^2 L_V + \frac{\rho^2\gamma_{k+1}^2 L_V L_s^2 \left( 2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\ &= -\gamma_{k+1} \left[ (v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 L_V + \frac{\rho^2\gamma_{k+1} L_V L_s^2 \left( 2\gamma_{k+1}^2\rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right] \end{aligned} \quad (96)$$



435 Furthermore, we recall that  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{2, 1+2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  
 436  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ . Then,

$$\begin{aligned}
 & \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right)}{\frac{1}{n} - \gamma_{k+1} \beta - \gamma_{k+1}^2 \rho^2 L_s^2} \\
 & \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} (k^2 \alpha^2 c_1^2 n^{4/3})^{-1} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1 \bar{L}^2 n^{1/3}}\right)}{\frac{1}{n} - \frac{1}{k \alpha n} - \frac{1}{k^2 \alpha^2 c_1^2 n^{4/3}}} \\
 & = \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1 \bar{L}^2 n^{1/3}}\right)}{(k \alpha c_1 n^{1/3})(k \alpha - 1) c_1 - 1} \\
 & \stackrel{(a)}{\leq} \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k \alpha c_1 \bar{L} n^{1/3}} \left(\frac{2}{k \alpha n} + 1\right)}{2(\alpha c_1 n^{1/3}) - 1} \\
 & \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{1}{k \alpha^2 c_1^3 \bar{L} n^{2/3}} \\
 & \leq \frac{1}{\alpha c_1 \bar{L} n^{2/3}}
 \end{aligned} \tag{97}$$

437 where (a) is due to  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$  and  $k\alpha c_1 n^{1/3} \geq 1$ . Also, since  $-\gamma_{k+1}(v_{\min} \rho +$   
 438  $v_{\max}^2) \leq -\gamma_{k+1} \rho v_{\min} = -$

439 Using the Lemma 2, we know that  $v_{\max}^2 \|\nabla V(\hat{s}^{(k)})\|^2 \leq \|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2$  and using (97) on (95)  
 440 yields:

$$\begin{aligned}
 v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] \\
 & \quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
 \end{aligned} \tag{98}$$

441 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned}
 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] & \leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] \\
 & \quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]
 \end{aligned} \tag{99}$$

442 **Bounding**  $\mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k)}\|^2]$  Remark that this term is the price we pay for the two time scale  
 443 dynamics and corresponds to the gap between the two asynchronous updates (one is on  $\hat{s}^{(k)}$  and the  
 444 other on  $\tilde{S}^{(k)}$ ).

445 **FIND AN UPPER BOUND TO THAT GAP**

446 □

## 447 F Proof of Theorem 2

448 **Theorem.** Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and  
 449 consider the vrTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for any  $k > 0$ .

450 Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$   
 451 and a constant  $\mu \in (0, 1)$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}] \end{aligned} \quad (100)$$

452 **Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful through-  
 453 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left( (1-\rho) [\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] + \rho [\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}] \right) \end{aligned} \quad (101)$$

454 We begin our proof by giving this auxiliary Lemma setting an upper bound for the quantity  $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2]$ .  
 455

456 **Lemma 8.** For any  $k \geq 0$  and consider the vrTTSEM update in (10) with  $\rho_k = \rho$ , it holds for all  
 457  $k > 0$

$$\begin{aligned} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_{\mathbf{s}}^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (102)$$

458 where we recall that  $\ell(k)$  is the first iteration number in the epoch that iteration  $k$  is in.

459 **Proof** We observe, using the identity (101), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] \quad (103)$$

460 For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{\ell(k)}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{\ell(k)})\right\|^2\right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \stackrel{(b)}{\leq} L_{\mathbf{s}}^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (104)$$

461 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (103) proves the  
 462 lemma.  $\square$

463 Using the smoothness of  $V$  and update (10), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (105)$$

464 Denote  $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fTTSEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 465 Taking expectations on both sides show that

$$\begin{aligned}
 & \mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\
 & \stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1-\rho)\mathbb{E}\left[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle\right] - \gamma_{k+1}\rho\mathbb{E}\left[\langle \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle\right] \\
 & + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\
 & \stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho\mathbb{E}\left[\langle \mathbf{h}_k \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle\right] - \gamma_{k+1}(1-\rho)\mathbb{E}\left[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle\right] \\
 & - \gamma_{k+1}\rho\mathbb{E}\left[\langle \eta_{i_k}^{(k+1)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle\right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\
 & \stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\
 & - \gamma_{k+1}\rho\mathbb{E}\left[\left\|\eta_{i_k}^{(k+1)}\right\|^2\right] - \gamma_{k+1}(1-\rho)\mathbb{E}\left[\left\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\right\|^2\right]
 \end{aligned} \tag{106}$$

466 where we have used (101) in (a) and  $\mathbb{E}[\mathbf{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$  in (b), the growth condition in  
 467 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

468 Furthermore, for  $k+1 \leq \ell(k) + m$  (i.e.,  $k+1$  is in the same epoch as  $k$ ), we have

$$\begin{aligned}
 & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\
 & = \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} \mid \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle\right] \\
 & = \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2\right. \\
 & \quad \left. - 2\gamma_{k+1}\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} \mid \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1-\rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle\right] \\
 & \leq \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta}\|\mathbf{h}_k\|^2\right. \\
 & \quad \left.+ \frac{\gamma_{k+1}\rho}{\beta}\|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2\right],
 \end{aligned} \tag{107}$$

469 where we first used (101) and the last inequality is due to the Young's inequality.

470 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \tag{108}$$

471 where  $b_k := \bar{b}_{k \bmod m}$  is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2) + \gamma_{k+1}^2\rho^2 L_V L_s^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0. \tag{109}$$

472 Note that  $\bar{b}_i$  is decreasing with  $i$  and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_s^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_s^2}, \quad i = 1, 2, \dots, m. \tag{110}$$

473 For  $k+1 \leq \ell(k) + m$ , we have the following inequality

$$\begin{aligned}
 R_{k+1} & \leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2\right] \\
 & + \gamma_{k+1}\mathbb{E}\left[\rho\left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho)\left\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\right\|^2\right] \\
 & + b_{k+1}\mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta}\|\mathbf{h}_k\|^2\right] \\
 & + b_{k+1}\mathbb{E}\left[\frac{\gamma_{k+1}\rho}{\beta}\|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2\right]
 \end{aligned} \tag{111}$$

474 And using Lemma 8 we obtain:

$$\begin{aligned}
R_{k+1} \leq & \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1} \rho v_{\min} + \gamma_{k+1} v_{\max}^2 - \gamma_{k+1}^2 \rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2 \rho^2 L_V L_{\mathbf{s}}^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 \right] \\
& + b_{k+1} \mathbb{E} \left[ (1 + \gamma_{k+1} \beta + 2\gamma_{k+1}^2 \rho^2 L_{\mathbf{s}}^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left( \frac{\gamma_{k+1} \rho}{\beta} + 2\gamma_{k+1}^2 \rho^2 \right) \|\mathbf{h}_k\|^2 \right] \\
& + \gamma_{k+1} \mathbb{E} \left[ (\rho + \rho^2 \gamma_{k+1} L_V) \left\| \eta_{i_k}^{(k+1)} \right\|^2 - (1 - \rho - (1 - \rho)^2 \gamma_{k+1} L_V) \left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{s}}^{(k)} \right\|^2 \right] \\
& + b_{k+1} \mathbb{E} \left[ \left( \frac{\gamma_{k+1} \rho}{\beta} + 2\gamma_{k+1}^2 \rho^2 \right) \left\| \eta_{i_k}^{(k+1)} \right\|^2 + \left( \frac{\gamma_{k+1} (1 - \rho)}{\beta} + 2\gamma_{k+1}^2 (1 - \rho)^2 \right) \left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{s}}^{(k)} \right\|^2 \right]
\end{aligned} \tag{112}$$

475 Rearranging the terms yields:

$$\begin{aligned}
R_{k+1} \leq & \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} (\rho v_{\min} + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V - b_{k+1} (\frac{\rho}{\beta} + 2\gamma_{k+1} \rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\
& + \underbrace{\left( b_{k+1} (1 + \gamma\beta + 2\gamma^2 \rho^2 L_{\mathbf{s}}^2) + \gamma^2 \rho^2 L_V L_{\mathbf{s}}^2 \right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}
\end{aligned} \tag{113}$$

476 where

$$\begin{aligned}
\tilde{\eta}^{(k+1)} &= \left( \gamma_{k+1} (\rho + \rho^2 \gamma_{k+1} L_V) + b_{k+1} \left( \frac{\gamma_{k+1} \rho}{\beta} + 2\gamma_{k+1}^2 \rho^2 \right) \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k+1)} \right\|^2 \right] \\
\tilde{\chi}^{(k+1)} &= \left( b_{k+1} \left( \frac{\gamma_{k+1} (1 - \rho)}{\beta} + 2\gamma_{k+1}^2 (1 - \rho)^2 \right) - \gamma_{k+1} (1 - \rho - (1 - \rho)^2 \gamma_{k+1} L_V) \right) \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{s}}^{(k)} \right\|^2 \right]
\end{aligned} \tag{114}$$

477 This leads, using Lemma 2, that for any  $\gamma_{k+1}$ ,  $\rho$  and  $\beta$  such that  $\rho v_{\min} + v_{\max}^2 -$   
478  $\gamma_{k+1} \rho^2 L_V - b_{k+1} (\frac{\rho}{\beta} + 2\gamma_{k+1} \rho^2) > 0$ ,

$$\begin{aligned}
v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] \leq & \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1} (\rho v_{\min} + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V - b_{k+1} (\frac{\rho}{\beta} + 2\gamma_{k+1} \rho^2))} \\
& + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1} (\rho v_{\min} + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V - b_{k+1} (\frac{\rho}{\beta} + 2\gamma_{k+1} \rho^2))}
\end{aligned} \tag{115}$$

479 We first remark that

$$\begin{aligned}
& \gamma_{k+1} (\rho v_{\min} + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V - b_{k+1} (\frac{\rho}{\beta} + 2\gamma_{k+1} \rho^2)) \\
& \geq \frac{\gamma_{k+1} \rho}{c_1} (1 - \gamma_{k+1} c_1 \rho L_V - b_{k+1} (\frac{c_1}{\beta} + 2\gamma_{k+1} \rho c_1))
\end{aligned} \tag{116}$$

480 where  $c_1 = v_{\min}^{-1}$ . By setting  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\beta = \frac{c_1 \bar{L}}{n^{1/3}}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{n c_1^2}{2\mu^2 + \mu c_1^2}$  and  
481  $\{\gamma_{k+1}\}$  any sequence of decreasing stepsizes in  $(0, 1)$ , it can be shown that there exists  $\mu \in (0, 1)$ ,  
482 such that the following lower bound holds

$$\begin{aligned}
1 - \gamma_{k+1} c_1 \rho L_V - b_{k+1} (\frac{c_1}{\beta} + 2\gamma_{k+1} \rho c_1) & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0 \left( \frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
& \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V \mu^2 (1 + \gamma\beta + 2\gamma^2 L_{\mathbf{s}}^2)^m - 1}{c_1^2 n^{\frac{4}{3}} \gamma\beta + 2\gamma^2 L_{\mathbf{s}}^2} \left( \frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L} n^{\frac{2}{3}}} \right) \\
& \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2} (e - 1) \left( 1 + \frac{2\mu}{n} \right) \geq 1 - \mu - \mu(1 + 2\mu) \frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2}
\end{aligned} \tag{117}$$

483 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2 L_{\mathbf{s}}^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2 n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \text{ and } (1 + \gamma\beta + 2\gamma^2 L_{\mathbf{s}}^2)^m \leq e - 1. \tag{118}$$

484 and the required  $\mu$  in (b) can be found by solving the quadratic equation.

485 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_{\max}})}{v_{\min} \rho} + 2 \sum_{k=0}^{K_{\max}-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min} \rho} \quad (119)$$

486 Note that  $R_0 = \mathbb{E}[V(\hat{\mathbf{s}}^{(0)})]$  and if  $K_{\max}$  is a multiple of  $m$ , then  $R_{\max} = \mathbb{E}[V(\hat{\mathbf{s}}^{(K_{\max})})]$ . Under the  
 487 latter condition, we have

$$\sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}] \quad (120)$$

488 This concludes our proof.

489 □

## G Practical Implementations of Two-Time-Scale EM Methods

### G.1 Gaussian mixture models

#### G.1.1 Model assumptions

We first recognize that the constraint set for  $\theta$  is given by

$$\Theta = \Delta^M \times \mathbb{R}^M. \quad (121)$$

Using the partition of the sufficient statistics as  $S(y_i, z_i) = (S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ , the partition  $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$  and the fact that  $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$ , the complete data log-likelihood can be expressed as in (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i) y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (122)$$

and  $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$ . We also define for each  $m \in \llbracket 1, M \rrbracket$ ,  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ . Consider the following latent sample used to compute an approximation of the conditional expected value  $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}} | y = y_i]$ :

$$z_{i,m} \sim \mathbb{P}(z_i = m | y_i; \theta) \quad (123)$$

where  $m \in \llbracket 1, M \rrbracket$ ,  $i \in \llbracket 1, n \rrbracket$  and  $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$ .

In particular, given iteration  $k + 1$ , the computation of the approximated quantity  $\tilde{S}_{i_k}^{(k)}$  during Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left( \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:=\tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1})y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})y_{i_k}}_{:=\tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:=\tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (124)$$

Recall that we have used the following regularizer:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (125)$$

It can be shown that the regularized M-step in (4) evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\mu}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (126)$$

where we have defined for all  $m \in \llbracket 1, M \rrbracket$  and  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ .

#### G.1.2 Algorithms updates

In the sequel, recall that, for all  $i \in \llbracket n \rrbracket$  and iteration  $k$ , the computed statistic  $\tilde{S}_{i_k}^{(k)}$  is defined by (124). At iteration  $k$ , the several E-steps defined by (9) or (10) and (11) leads to the definition of the quantity  $\hat{\mathbf{s}}^{(k+1)}$ . For the GMM example, after the initialization of the quantity  $\hat{\mathbf{s}}^{(0)} = n^{-1} \sum_{i=1}^n \tilde{\mathbf{s}}_i^{(0)}$ , those E-steps break down as follows:

**Batch EM (EM):** for all  $i \in \llbracket 1, n \rrbracket$ , compute  $\tilde{\mathbf{s}}_i^{(k)}$  and set

$$\hat{\mathbf{s}}^{(k+1)} = n^{-1} \sum_{i=1}^n \tilde{\mathbf{s}}_i^{(k)}. \quad (127)$$

514 where  $\bar{s}_i^{(k)}$  are computed using the exact conditional expected value  $\mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}}|y=y_i]$ :

$$\tilde{\omega}_m(y_i; \theta) := \mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}}|y=y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)}, \quad (128)$$

515 **Incremental EM (iEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}. \quad (129)$$

516 **batch SAEM (SAEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}. \quad (130)$$

517 where  $= \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$  with  $\tilde{S}_i^{(k)}$  defined in (124).

518 **Incremental SAEM (iSAEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_i^k)})). \quad (131)$$

520 **Variance Reduced Two-Time-Scale EM (vrTTSEM):** draw an index  $i_k$  uniformly at random on  
521  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}))). \quad (132)$$

522 **Fast Incremental Two-Time-Scale EM (fiTTSEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ ,  
523 compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)}(1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)}(1 - \rho) + \rho(\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}))). \quad (133)$$

524 Finally, the  $k$ -th update reads  $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$  where the function  $s \rightarrow \bar{\theta}(s)$  is defined by (126).