

---

# Fast Two-Time-Scale Noisy EM Algorithms

---

Anonymous Author(s)

Affiliation

Address

email

## Abstract

Training latent data models using the EM algorithm is the most popular choice for current learning tasks. For today's modern and complex tasks, variants of the EM have been initially introduced by [Neal and Hinton, 1998], using incremental updates to scale to large dataset, and by [Wei and Tanner, 1990, Delyon et al., 1999], using Monte-Carlo (MC) approximations to bypass the impossible conditional expectation of the latent data for most nonconvex models. In this paper, we propose to combine both techniques in a single class of methods called Two-Time-Scale EM Methods. We motivate the choice of a double dynamics by invoking the variance reduction virtue of each stage of the method on both noises: the incremental update and the MC approximation. We establish finite-time and independent of the initialization convergence bounds for nonconvex objective function. Numerical applications are also presented in this article to illustrate our findings.

## 1 Introduction

Learning latent data models is critical for modern machine learning problems, see [McLachlan and Krishnan, 2007] for references. We formulate the training of such model as the following empirical risk minimization problem:

$$\min_{\theta \in \Theta} \bar{L}(\theta) := r(\theta) + L(\theta) \quad \text{with} \quad L(\theta) = \frac{1}{n} \sum_{i=1}^n L_i(\theta) := \frac{1}{n} \sum_{i=1}^n \{ -\log g(y_i; \theta) \}, \quad (1)$$

We denote the observations by  $\{y_i\}_{i=1}^n$ ,  $\Theta \subset \mathbb{R}^d$  is the convex parameters space. We consider a regularized model where  $r : \Theta \rightarrow \mathbb{R}$  is a smooth convex regularization function and for  $\theta \in \Theta$ ,  $g(y; \theta)$  is the (incomplete) likelihood of each individual observation. The objective function  $\bar{L}(\theta)$  is possibly *nonconvex* and is assumed to be lower bounded  $\bar{L}(\theta) > -\infty$  for all  $\theta \in \Theta$ .

In the latent variable model,  $g(y_i; \theta)$ , is the marginal of the complete data likelihood defined as  $f(z_i, y_i; \theta)$ , i.e.  $g(y_i; \theta) = \int_{\mathcal{Z}} f(z_i, y_i; \theta) \mu(dz_i)$ , where  $\{z_i\}_{i=1}^n$  are the (unobserved) latent variables. In this paper, we make the assumption of a complete model belonging to the curved exponential family, i.e.,

$$f(z_i, y_i; \theta) = h(z_i, y_i) \exp(\langle S(z_i, y_i) | \phi(\theta) \rangle - \psi(\theta)), \quad (2)$$

where  $\psi(\theta)$ ,  $h(z_i, y_i)$  are scalar functions,  $\phi(\theta) \in \mathbb{R}^k$  is a vector function, and  $S(z_i, y_i) \in \mathbb{R}^k$  is the complete data sufficient statistics.

Full batch EM [Dempster et al., 1977] is the method of reference for that kind of task and is a two steps procedure. The E-step amounts to computing the conditional expectation of the complete data sufficient statistics,

$$\bar{s}(\theta) = \frac{1}{n} \sum_{i=1}^n \bar{s}_i(\theta) \quad \text{where} \quad \bar{s}_i(\theta) = \int_{\mathcal{Z}} S(z_i, y_i) p(z_i | y_i; \theta) \mu(dz_i). \quad (3)$$

30 The M-step is given by

$$\text{M-step: } \hat{\theta} = \bar{\theta}(\bar{s}(\theta)) := \arg \min_{\vartheta \in \Theta} \{ r(\vartheta) + \psi(\vartheta) - \langle \bar{s}(\theta) | \phi(\vartheta) \rangle \}, \quad (4)$$

31 Two caveats of this method are the following: (a) with the explosion of data, the first step of the EM  
 32 is computationally inefficient as it requires a full pass over the dataset at each iteration and (b) the  
 33 complexity of modern models makes the expectation intractable. So far, both challenges have been  
 34 addressed separately, to the best of our knowledge, and we give an overview of current solutions in  
 35 the sequel.

36 **Prior Work** Inspired by stochastic optimization procedures, [Neal and Hinton, 1998] and [Cappé  
 37 and Moulines, 2009] developed respectively an incremental and an online variant of the E-step in  
 38 models where the expectation is computable then extensively used and studied in [Nguyen et al.,  
 39 2020, Liang and Klein, 2009, Cappé, 2011]. Some improvements of that methods have been pro-  
 40 vided and analyzed, globally and in finite-time, in [Karimi et al., 2019] where variance reduction  
 41 techniques taken from the optimization literature have been efficiently applied to scale the EM algo-  
 42 rithm to large datasets.

43 Regarding the computation of the expectation under the posterior distribution, the first method was  
 44 the Monte-Carlo EM (MCEM) introduced in the seminal paper [Wei and Tanner, 1990] where a MC  
 45 approximation of this expectation is computed. A variant of that method is the Stochastic Approx-  
 46 imation of the EM (SAEM) in [Delyon et al., 1999] leveraging the power of Robbins-Monro type of  
 47 update [Robbins and Monro, 1951] to ensure pointwise convergence of the vector of estimated pa-  
 48 rameters rather using a decreasing stepsize than increasing the number of MC samples. The MCEM  
 49 and the SAEM have been successfully applied in mixed effects models [McCulloch, 1997, Hughes,  
 50 1999, Baey et al., 2016] or to do inference for joint modelling of time to event data coming from  
 51 clinical trials in [Chakraborty and Das, 2010], among other applications.

52 Recently, an incremental variant of the SAEM was proposed in [Kuhn et al., 2019] showing positive  
 53 empirical results but its analysis is limited to asymptotic consideration. Gradient-based methods  
 54 have been developed and analyzed in [Zhu et al., 2017] but they remain out of the scope of this  
 55 paper as they tackle the high-dimensionality issue.

56 **Contributions** This paper *introduces* and *analyzes* a new class of methods which purpose is to  
 57 combine both solutions proposed in the past years in a two-time-scale manner in order to optimize  
 58 (1) for current modern examples and settings. The main contributions of the paper are:

- 59 • We propose a two-time-scale method based on Stochastic Approximation (SA), to alleviate  
 60 the problem of MC computation, and on Incremental updates, to scale to large datasets. We  
 61 describe in details the edges of each level of our method based on variance reduction argu-  
 62 ments. The derivation of such class of algorithms has two advantages. First, it combines  
 63 two powerful ideas, commonly used separately, to tackle large scale and highly nonlin-  
 64 ear learning tasks. Then, it gives a simple formulation as a *scaled-gradient method* which  
 65 makes the global analysis accessible.
- 66 • We also establish global (independent of the initialization) and finite-time (true at each  
 67 iteration) upper bounds on a classical suboptimality condition in the nonconvex literature,  
 68 *i.e.*, the second order moment of the gradient of the objective function.

69 In Section 2 we give rigorous mathematical definitions of the various updates used for both incre-  
 70 mental and Monte-Carlo EMs and we introduce the main class of new algorithms, based on two  
 71 different dynamics, we are proposing to analyze and compare to baselines algorithms. Section 3  
 72 presents the main theoretical guarantees of this newly introduced two-time-scale class of algorithms.  
 73 Results are given both in finite-time and in the nonconvex setting. Finally, we illustrate the advan-  
 74 tages of our method in Section 4 on two numerical experiments.

## 75 2 Two-Time-Scale Stochastic EM Algorithms

76 We recall and formalize in this section the different methods found in the literature that aim to solv-  
 77 ing the large scale problem and the intractable expectation. We then provide the general framework  
 78 of our method to efficiently tackle the optimization problem (1).

## 79 2.1 Monte Carlo Integration and Stochastic Approximation

80 As mentioned in the introduction, for complex and possibly nonlinear models, the expectation under  
 81 the posterior distribution defined in (3) is not tractable. In that case, the first solution involves  
 82 computing a Monte Carlo integration of that latter term. For all  $i \in \llbracket 1, n \rrbracket$ , draw for  $m \in \llbracket 1, M \rrbracket$ ,  
 83 samples  $z_{i,m} \sim p(z_i|y_i; \theta)$  and compute the MC integration  $\tilde{s}$  of the deterministic quantity  $\bar{s}(\theta)$ :

$$\text{MC-step : } \tilde{s} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}, y_i) \quad (5)$$

84 and then update the parameter  $\hat{\theta} = \bar{\theta}(\tilde{s})$ . This algorithm bypasses the intractable expectation issue  
 85 but is rather computationally expensive in order to reach point wise convergence ( $M$  needs to be  
 86 large). An alternative to that stochastic algorithm is to use a Robbins-Monro (RM) type of update.  
 87 We denote, at iteration  $k$ , the following quantity

$$\tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}^{(k)}, y_i) \quad \text{where } z_{i,m}^{(k)} \sim p(z_i|y_i; \theta^{(k)}) \quad (6)$$

88 Then, the RM updated of the sufficient statistics  $\hat{s}^{(k+1)}$  reads:

$$\text{SA-step : } \hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad (7)$$

89 where  $\{\gamma_k\}_{k \geq 1} \in (0, 1)$  is a sequence of decreasing step sizes to ensure asymptotic convergence.  
 90 This is called the Stochastic Approximation of the EM (SAEM) and has been shown theoretically to  
 91 converge to a maximum of the likelihood of the observations under very general conditions [Delyon  
 92 et al., 1999]. In the simulation step (6), since the relation between the observed data  $y_i$  and the  
 93 latent variable  $z_i$  can be non linear, sampling from the posterior distribution  $p(z_i|y_i; \theta)$ , under the  
 94 current model  $\theta$ , could require using an inference algorithm. [?] proved almost sure convergence  
 95 of the sequence of parameters obtained by this algorithm coupled with an MCMC procedure during  
 96 the simulation step. In simple scenarios, the samples  $\{z_{i,m}\}_{m=0}^{M-1}$  are conditionally independent and  
 97 identically distributed with distribution  $p(z_i, \theta)$ . Nevertheless, in most cases, sampling exactly from  
 98 this distribution is not an option and the Monte Carlo batch is sampled by Monte Carlo Markov  
 99 Chains (MCMC) algorithm. In the SA-step, the sequence of decreasing positive integers  $\{\gamma_k\}_{k \geq 1}$   
 100 controls the convergence of the algorithm. In practice,  $\gamma_k$  is set equal to 1 during the first few  
 101 iterations to let the algorithm explore the parameter space without memory and converge quickly  
 102 to a neighbourhood of the target estimate. The Stochastic Approximation is performed during the  
 103 remaining iterations where  $\gamma_k = 1/k^\alpha$ , where  $\alpha \in (0, 1)$ , ensuring the almost sure convergence of  
 104 the estimate. It is inappropriate to start with small values for step size  $\gamma_k$  and large values for the  
 105 number of simulations  $M_k$ . Rather, it is recommended that one decrease  $\gamma_k$  and keep a constant  
 106 and small number of MC samples  $M_k$  which shows a great advantage over the MC-step (5), which  
 107 requires large  $M_k$  to converge.

108 This Robbins-Monro type of update represents the *first level* of our algorithm, needed to temper  
 109 the variance and noise implied by MC integration. In the next section, we derive variants of this  
 110 algorithm to adapt to the sheer size of data of today's applications and formalize the *second level* of  
 111 our class of Two-Time-Scale EM methods.

## 112 2.2 Incremental and Bi-Level Inexact EM Methods

113 Strategies to scale to large datasets include classical incremental and variance reduced variants. We  
 114 will explicit a general update that will cover those variants and that represents the *second level* of our  
 115 algorithm, namely the incremental update of the noisy statistics  $\hat{S}^{(k)}$  inside the RM type of update.

$$\text{Incremental-step : } \tilde{S}^{(k+1)} = \tilde{S}^{(k)} + \rho_{k+1}(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}), \quad (8)$$

116 Note  $\{\rho_k\}_{k \geq 1} \in (0, 1)$  is a sequence of step sizes,  $\mathcal{S}^{(k)}$  is a proxy for  $\tilde{S}^{(k)}$ , If the stepsize is equal  
 117 to one and the proxy  $\mathcal{S}^{(k)} = \hat{S}^{(k)}$ , i.e., computed in a full batch manner as in (6), then we recover  
 118 the SAEM algorithm. Also if  $\rho_k = 1$ ,  $\gamma_k = 1$  and  $\mathcal{S}^{(k)} = \tilde{S}^{(k)}$ , then we recover the Monte Carlo  
 119 EM algorithm.

120 We now introduce three variants of the SAEM update depending on different definitions of the  
 121 proxy  $\mathcal{S}^{(k)}$  and the choice of the stepsize  $\rho_k$ . Let  $i_k \in \llbracket 1, n \rrbracket$  be a random index drawn at iteration  
 122  $k$  and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$  be the iteration index where  $i \in \llbracket 1, n \rrbracket$  is last drawn  
 123 prior to iteration  $k$ . For iteration  $k \geq 0$ , the fiTTSEM method draws *two* indices *independently* and  
 124 uniformly as  $i_k, j_k \in \llbracket 1, n \rrbracket$ . In addition to  $\tau_i^k$  which was defined w.r.t.  $i_k$ , we define  $t_j^k = \{k' : j_{k'} = j, k' < k\}$  to be the iteration index where the sample  $j \in \llbracket 1, n \rrbracket$  is last drawn as  $j_k$  prior to  
 125 iteration  $k$ . With the initialization  $\bar{\mathcal{S}}^{(0)} = \bar{s}^{(0)}$ , we use a slightly different update rule from SAGA  
 126 inspired by [Reddi et al., 2016]. Then, we obtain:

$$(iSAEM [Kuhn et al., 2019]) \quad \mathcal{S}^{(k+1)} = \mathcal{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)}) \quad (9)$$

$$(vrTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}) \quad (10)$$

$$(fiTTSEM This paper) \quad \mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \quad (11)$$

$$\bar{\mathcal{S}}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + n^{-1} (\tilde{S}_{j_k}^{(k)} - \tilde{S}_{j_k}^{(t_{j_k}^k)}). \quad (12)$$

128 The stepsize is set to  $\rho_{k+1} = 1$  for the iSAEM method;  $\rho_{k+1} = \gamma$  is constant for the vrTTSEM and  
 129 fiTTSEM methods. Moreover, for iSAEM we initialize with  $\mathcal{S}^{(0)} = \tilde{S}^{(0)}$ ; for vrTTSEM we set an  
 130 epoch size of  $m$  and define  $\ell(k) := m \lfloor k/m \rfloor$  as the first iteration number in the epoch that iteration  
 131  $k$  is in.

### 132 2.3 Two-Time-Scale Noisy EM methods

133 We now introduce the general method derived using the two variance reduction techniques described  
 134 above. Algorithm 1 leverages both levels (7) and (8) in order to output a vector of fitted parameters  
 135  $\hat{\theta}^{(K)}$  where  $K$  is some randomly chosen termination point.

136 The update in (14) is said to have two timescales as the step sizes satisfy  $\lim_{k \rightarrow \infty} \gamma_k / \rho_k < 1$  such that  
 137  $\tilde{S}^{(k+1)}$  is updated at a faster timescale than  $\hat{s}^{(k+1)}$ .

---

#### Algorithm 1 Two-Time-Scale Noisy EM methods.

---

- 1: **Input:** initializations  $\hat{\theta}^{(0)} \leftarrow 0, \hat{s}^{(0)} \leftarrow \tilde{S}^{(0)}, K_{\max} \leftarrow \text{max. iteration number}$ .
- 2: Set the terminating iteration number,  $K \in \{0, \dots, K_{\max} - 1\}$ , as a discrete r.v. with:

$$P(K = k) = \frac{\gamma_k}{\sum_{\ell=0}^{K_{\max}-1} \gamma_\ell}. \quad (13)$$

- 3: **for**  $k = 0, 1, 2, \dots, K$  **do**
- 4:   Draw index  $i_k \in \llbracket 1, n \rrbracket$  uniformly (and  $j_k \in \llbracket 1, n \rrbracket$  for fiTTSEM).
- 5:   Compute  $\hat{S}_{i_k}^{(k)}$  using the MC-step (5), for the drawn indices.
- 6:   Compute the surrogate sufficient statistics  $\mathcal{S}^{(k+1)}$  using (9) or (10) or (11).
- 7:   Compute  $\hat{S}^{(k+1)}$  and  $\hat{s}^{(k+1)}$  using respectively (8) and (7):

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho_{k+1} (\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{s}^{(k+1)} &= \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \end{aligned} \quad (14)$$

- 8:   Compute  $\hat{\theta}^{(k+1)}$  via the M-step (4).
  - 9: **end for**
  - 10: **Return:**  $\hat{\theta}^{(K)}$ .
- 

138 The next section presents the main results of this paper and establishes global and finite-time bounds  
 139 for the three different updates of our two-time-scale scheme..

### 3 Finite Time Analysis of the Two-Time-Scale Scheme

Following [Cappé and Moulines, 2009], it can be shown that stationary points of the objective function (1) corresponds to the stationary points of the following *nonconvex* Lyapunov function:

$$\min_{\mathbf{s} \in \mathcal{S}} V(\mathbf{s}) := \bar{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) = r(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(\bar{\boldsymbol{\theta}}(\mathbf{s})) \quad (15)$$

We thus propose to study the latter minimization problem in the sequel.

An important assumption in order to derive convergence guarantees reads as follows:

**H1.** *The sets  $\mathcal{Z}, \mathcal{S}$  are compact. There exists constants  $C_S, C_Z$  such that:*

$$C_S := \max_{\mathbf{s}, \mathbf{s}' \in \mathcal{S}} \|\mathbf{s} - \mathbf{s}'\| < \infty, \quad C_Z := \max_{i \in \llbracket 1, n \rrbracket} \int_{\mathcal{Z}} |S(z, y_i)| \mu(dz) < \infty. \quad (16)$$

**H2.** *The conditional distribution is smooth on  $\text{int}(\Theta)$ . For any  $i \in \llbracket 1, n \rrbracket$ ,  $z \in \mathcal{Z}$ ,  $\boldsymbol{\theta}, \boldsymbol{\theta}' \in \text{int}(\Theta)^2$ , we have  $|p(z|y_i; \boldsymbol{\theta}) - p(z|y_i; \boldsymbol{\theta}')| \leq L_p \|\boldsymbol{\theta} - \boldsymbol{\theta}'\|$ .*

We also recall from the introduction that we consider curved exponential family models. besides:

**H3.** *For any  $\mathbf{s} \in \mathcal{S}$ , the function  $\boldsymbol{\theta} \mapsto L(\mathbf{s}, \boldsymbol{\theta}) := r(\boldsymbol{\theta}) + \psi(\boldsymbol{\theta}) - \langle \mathbf{s} | \phi(\boldsymbol{\theta}) \rangle$  admits a unique global minimum  $\bar{\boldsymbol{\theta}}(\mathbf{s}) \in \text{int}(\Theta)$ . In addition,  $J_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))$  is full rank,  $L_{\phi}$ -Lipschitz and  $\bar{\boldsymbol{\theta}}(\mathbf{s})$  is  $L_{\theta}$ -Lipschitz.*

We denote by  $H_L^{\boldsymbol{\theta}}(\mathbf{s}, \boldsymbol{\theta})$  the Hessian (w.r.t to  $\boldsymbol{\theta}$  for a given value of  $\mathbf{s}$ ) of the function  $\boldsymbol{\theta} \mapsto L(\mathbf{s}, \boldsymbol{\theta}) = r(\boldsymbol{\theta}) + \psi(\boldsymbol{\theta}) - \langle \mathbf{s} | \phi(\boldsymbol{\theta}) \rangle$ , and define

$$\mathbf{B}(\mathbf{s}) := J_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left( H_L^{\boldsymbol{\theta}}(\mathbf{s}, \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} J_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top}. \quad (17)$$

**H4.** *It holds that  $v_{\max} := \sup_{\mathbf{s} \in \mathcal{S}} \|\mathbf{B}(\mathbf{s})\| < \infty$  and  $0 < v_{\min} := \inf_{\mathbf{s} \in \mathcal{S}} \lambda_{\min}(\mathbf{B}(\mathbf{s}))$ . There exists a constant  $L_B$  such that for all  $\mathbf{s}, \mathbf{s}' \in \mathcal{S}^2$ , we have  $\|\mathbf{B}(\mathbf{s}) - \mathbf{B}(\mathbf{s}')\| \leq L_B \|\mathbf{s} - \mathbf{s}'\|$ .*

The class of algorithms we develop in this paper are two time-scale where the first stage corresponds to the variance reduction trick used in [Karimi et al., 2019] in order to accelerate incremental methods and reduce the variance induced by the index sampling. The second stage is the Robbins-Monro type of update that aims to reduce the variance induced by the MC approximations

Indeed the expectations (3) are never available and requires Monte Carlo approximation. Thus, at iteration  $k + 1$ , we introduce the errors when approximating the quantity  $\bar{s}_i(\hat{\boldsymbol{\theta}}(\hat{\mathbf{s}}^{(k-1)}))$ . For all  $i \in \llbracket 1, n \rrbracket$ ,  $r > 0$  and  $\vartheta \in \Theta$ , define:

$$\eta_i^{(r)} := \tilde{S}_i^{(r)} - \bar{s}_i(\vartheta^{(r)}) \quad (18)$$

For instance, we consider that the MC approximation is unbiased if for all  $i \in \llbracket 1, n \rrbracket$  and  $m \in \llbracket 1, M \rrbracket$ , the samples  $z_{i,m} \sim p(z_i|y_i; \boldsymbol{\theta})$  are i.i.d. under the posterior distribution, i.e.,  $\mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] = 0$  where  $\mathcal{F}_r$  is the filtration up to iteration  $r$ . The following results are derived under the assumption of control of the fluctuations implied by the approximation stated as follows:

**H5.** *There exist a positive sequence of MC batch size  $\{M_r\}_{r>0}$  and constants  $(C, C_{\eta})$  such that for all  $k > 0$ ,  $i \in \llbracket 1, n \rrbracket$  and  $\vartheta \in \Theta$ :*

$$\mathbb{E} \left[ \left\| \eta_i^{(r)} \right\|^2 \right] \leq \frac{C_{\eta}}{M_r} \quad \text{and} \quad \mathbb{E} \left[ \left\| \mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] \right\|^2 \right] \leq \frac{C}{M_r} \quad (19)$$

In that setting, we can prove two important results on the Lyapunov function. The first one suggests smoothness:

**Lemma 1.** [Karimi et al., 2019] *Assume H1-H4. For all  $\mathbf{s}, \mathbf{s}' \in \mathcal{S}$  and  $i \in \llbracket 1, n \rrbracket$ , we have*

$$\|\bar{s}_i(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \bar{s}_i(\bar{\boldsymbol{\theta}}(\mathbf{s}'))\| \leq L_S \|\mathbf{s} - \mathbf{s}'\|, \quad \|\nabla V(\mathbf{s}) - \nabla V(\mathbf{s}')\| \leq L_V \|\mathbf{s} - \mathbf{s}'\|, \quad (20)$$

where  $L_S := C_Z L_p L_{\theta}$  and  $L_V := v_{\max}(1 + L_S) + L_B C_S$ .

and the second one suggests a growth condition on the gradient of  $V$  depending on the mean field of the algorithm:

**Lemma 2.** *Assume H3, H4. For all  $\mathbf{s} \in \mathcal{S}$ ,*

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (21)$$

Proof of this Lemma can be found in Appendix A.

### 3.1 Global Convergence of Incremental Noisy EM Algorithms

We present in this section a finite-time analysis of the incremental variant of the Stochastic Approximation of the EM algorithm. We want to draw the attention of the readers that the word "global" here does not mean for a global optimum of the nonconvex function, but of the independence of our analysis on the initialization and the iteration  $k$  (finite time).

The first intermediate result is the computation of the quantity  $\hat{S}^{(k+1)} - \hat{s}^{(k)}$ , which corresponds to the drift term of (7) and reads as follows:

**Lemma 3.** *The update (9) is equivalent to the following update on the resulting statistics*

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \quad \text{where} \quad \tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^{k+1})} \quad (22)$$

Also:

$$\mathbb{E}[\tilde{S}^{(k+1)} - \hat{s}^{(k)}] = \mathbb{E}[\bar{s}^{(k)} - \hat{s}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)}\right] + \frac{1}{n} \mathbb{E}[\eta_{i_k}^{(k+1)}] \quad (23)$$

where  $\bar{s}^{(k)}$  is defined by (3) and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ .

Proof of this Lemma can be found in Appendix B.

The following main result for the iSAEM algorithm is derived under a control of the Monte Carlo fluctuations as described by assumption H 5. Typically, the controls exhibited below are of interest when the number of MC samples  $M_k$  increase with the iteration index  $f$ .

**Theorem 1.** *Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and consider the iSAEM sequence  $\{\hat{s}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ . We also set  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  $\hat{s}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ .*

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K)})] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E}[\|\eta_{i_k}^{(k)}\|^2] \quad (24)$$

Proof of this Theorem can be found in Appendix C.

### 3.2 Global Convergence of Two-Time-Scale Noisy EM Algorithms

We now proceed by giving our main result regarding the global convergence of the fiTTSEM algorithm. Two important auxiliary Lemmas, which proofs are given in Appendix D.1, are need in order to derive our finite-time bound. The first one derives an identity for the quantity  $\mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2]$  using the vrTTSEM update:

**Lemma 4.** *For any  $k \geq 0$  and consider the vrTTSEM update in (10) with  $\rho_k = \rho$ , it holds for all  $k > 0$*

$$\begin{aligned} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E}[\|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(\ell(k))}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{s}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (25)$$

where we recall that  $\ell(k)$  is the first iteration number in the epoch that iteration  $k$  is in.

The second one derives an identity for the quantity  $\mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2]$  using the fiTTSEM update:

**Lemma 5.** *For any  $k \geq 0$  and consider the fiTTSEM update in (11) with  $\rho_k = \rho$ , it holds for all  $k > 0$*

$$\begin{aligned} \mathbb{E}[\|\hat{s}^{(k)} - \tilde{S}^{(k+1)}\|^2] &\leq 2\rho^2 \mathbb{E}[\|\hat{s}^{(k)} - \bar{s}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{s}^{(k)} - \hat{s}^{(\tau_i^k)}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{s}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (26)$$



206 We now state the main result regarding the vrTTSEM method.

207 **Theorem 2.** Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of  
 208 positive step sizes and consider the vrTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for  
 209 any  $k > 0$ .

210 Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$   
 211 and a constant  $\mu \in (0, 1)$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (27)$$

212 Proof of this Theorem can be found in Appendix E. We now state the main result regarding the  
 213 fiTTSEM method.

214 **Theorem 3.** Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of  
 215 positive step sizes and consider the fiTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for  
 216 any  $k > 0$ .

217 Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  
 218  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (28)$$

219 Proof of this Theorem can be found in Appendix F. Note that in those two bounds, the quantities  
 220  $\tilde{\eta}^{(k+1)}$  and  $\Xi^{(k+1)}$  are abstraction that depends only on the MC fluctuations  $\mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]$  and some  
 221 constants.

222 **Remarks:** The following remarks are worth noting on the quantity  $\mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right]$ :

- 223 • This term is the price we pay for the two time scale dynamics and corresponds to the gap
- 224 between the two asynchronous updates (one is on  $\hat{\mathbf{s}}^{(k)}$  and the other on  $\tilde{S}^{(k)}$ ).
- It is trivial to see that if  $\rho = 1$ , i.e., there is no variance reduction, then

$$\mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] = \mathbb{E} \left[ \left\| \mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)} \right\|^2 \right] = 0 \quad \text{with} \quad \hat{\mathbf{s}}^{(0)} = \tilde{S}^{(0)} = 0$$

225 which strengthen the fact that this quantity characterizes the impact of the variance reduc-  
 226 tion technique introduced in our two stages class of methods.

227 The following lemma, which proof can be found in Appendix D.2, can be derived to characterize  
 228 this gap:

229 **Lemma 6.** Consider a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho \in (0, 1)$ , then the following  
 230 inequality holds:

$$\mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_{\ell})^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}) \quad (29)$$

231 where  $\mathcal{S}^{(k)}$  is defined either by (10) (vrTTSEM) or (11) (fiTTSEM).

232 In the next section, we illustrate the benefits of our two-time-scale class of methods on several  
 233 numerical applications.

## 4 Numerical Examples

### 4.1 Gaussian Mixture Models

We begin by a simple and illustrative example. The authors acknowledge that the following model can be trained using deterministic EM-type of algorithms but propose to apply stochastic methods, including theirs, and to compare their performances. Given  $n$  observations  $\{y_i\}_{i=1}^n$ , we want to fit a Gaussian Mixture Model (GMM) whose distribution is modeled as a Gaussian mixture of  $M$  components, each with a unit variance. Let  $z_i \in \llbracket M \rrbracket$  be the latent labels of each component, the complete log-likelihood is defined as:

$$\log f(z_i, y_i; \theta) = \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) [\log(\omega_m) - \mu_m^2/2] + \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) \mu_m y_i + \text{constant} . \quad (30)$$

where  $\theta := (\omega, \mu)$  with  $\omega = \{\omega_m\}_{m=1}^{M-1}$  are the mixing weights with the convention  $\omega_M = 1 - \sum_{m=1}^{M-1} \omega_m$  and  $\mu = \{\mu_m\}_{m=1}^M$  are the means. We use the penalization  $r(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \log \text{Dir}(\omega; M, \epsilon)$  where  $\delta > 0$  and  $\text{Dir}(\cdot; M, \epsilon)$  is the  $M$  dimensional symmetric Dirichlet distribution with concentration parameter  $\epsilon > 0$ . The constraint set on  $\theta$  is given by

$$\Theta = \{\omega_m, m = 1, \dots, M-1 : \omega_m \geq 0, \sum_{m=1}^{M-1} \omega_m \leq 1\} \times \{\mu_m \in \mathbb{R}, m = 1, \dots, M\}. \quad (31)$$

Exact two time scale updates are given in Appendix G.1.

In the following experiments on synthetic data, we generate samples from a GMM model with  $M = 2$  components with two mixtures with means  $\mu_1 = -\mu_2 = 0.5$ . We use  $n = 10^5$  synthetic samples and run the bEM method until convergence (to double precision) to obtain the ML estimate  $\mu^*$  averaged on 50 datasets. We compare the bEM, iEM (incremental EM), SAEM, iSAEM, vrTTSEM and fiTTSEM methods in terms of their precision measured by  $|\mu - \mu^*|^2$ . We set the stepsize of the SA-step of all method as  $\gamma_k = 1/k^\alpha$  with  $\alpha = 0.5$ , and the stepsizes of the Incremental-step for vrTTSEM and the fiTTSEM to a constant stepsize equal to  $1/n^{2/3}$ .

The number of MC samples is fixed to  $M = 10$  chains. Figure 1 shows the convergence of the precision  $|\mu - \mu^*|^2$  for the different methods against the epoch(s) elapsed (one epoch equals  $n$  iterations). We observe that the vrTTSEM and fiTTSEM methods outperform the other stochastic methods, supporting the benefits of our newly introduced scheme.




Figure 1: TO COMPLETE

### 4.2 Deformable Template Model for Image Analysis

Let  $(y_i, i \in \llbracket 1, n \rrbracket)$  be observed gray level images defined on a grid of pixels. Let  $u \in \mathcal{U} \subset \mathbb{R}^2$  denotes the pixel index on the image and  $x_u \in \mathcal{D} \subset \mathbb{R}^2$  its location. The model used in this experiment suggests that each image  $y_i$  is a deformation of a template, noted  $I : \mathcal{D} \rightarrow \mathbb{R}$ , common to all images of the dataset:

$$y_i(u) = I(x_u - \Phi_i(x_u, z_i)) + \varepsilon_i(u) \quad (32)$$

where  $\phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a deformation function,  $z_i$  some latent variable parametrizing this deformation and  $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$  is an observation error.

The template model, given  $(p_k, k \in \llbracket 1, k_p \rrbracket)$  landmarks on the template, a fixed known kernel  $\mathbf{K}_p$  and a vector of parameters  $\beta \in \mathbb{R}^{k_p}$  is defined as follows:

$$I_\xi = \mathbf{K}_p \beta, \quad \text{where} \quad (\mathbf{K}_p \beta)(x) = \sum_{k=1}^{k_p} \mathbf{K}_p(x, p_k) \beta_k \quad (33)$$

Besides, we parameterize the deformation model given some landmarks  $(g_k, k \in \llbracket 1, k_g \rrbracket)$  and a fixed kernel  $\mathbf{K}_g$  as:

$$\Phi_i = \mathbf{K}_g z_i \quad \text{where} \quad (\mathbf{K}_g z_i)(x) = \sum_{k=1}^{k_g} \mathbf{K}_g(x, g_k) \left( z_i^{(1)}(k), z_i^{(2)}(k) \right) \quad (34)$$



where we put a Gaussian prior on the latent variables,  $z_i \sim \mathcal{N}(0, \Gamma)$  and  $z_i \in (\mathbb{R}^{k_g})^2$ . The vector of parameters we ought to estimate is thus  $\theta = (\beta, \Gamma, \sigma)$ . The complete model belongs to the curved exponential family, see [Allasonnière et al., 2007], which vector of sufficient statistics  $S = (S_1(z), S_2(z), S_3(z))$  read:

$$\begin{aligned} S_1(z) &= \frac{1}{n} \sum_{i=1}^n S_1(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top y_i \\ S_2(z) &= \frac{1}{n} \sum_{i=1}^n S_2(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \frac{1}{n} \sum_{i=1}^n S_3(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (35)$$

where for any pixel  $u \in \mathbb{R}^2$  and  $j \in \llbracket 1, k_g \rrbracket$  we noted:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j) \quad (36)$$

Finally, the Two-Time-Scale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z) \hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n} \hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z) \beta(\hat{s}) - 2\beta(\hat{s}) \hat{s}_1(z) \end{pmatrix} \quad (37)$$

where  $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$  is the vector of statistics obtained via the SA-step (7) and using the MC approximation of the sufficient statistics  $(S_1(z), S_2(z), S_3(z))$  defined in (42).

### Comparison using epochs credit

### Comparison using number of training samples credit

## 4.3 PK Model with Absorption Lag Time

This numerical example was conducted in order to characterize the pharmacokinetics of orally administered drug to simulated patients, using a population pharmacokinetic approach.  $M = 50$  synthetic datasets were generated for  $n = 500$  patients with 10 observations (concentration measures) per patient.

**The model:** We consider a one-compartment pharmacokinetics (PK) model for oral administration with an absorption lag-time ( $T^{\text{lag}}$ ), assuming first-order absorption and linear elimination processes. The final model includes  $ka$  is the absorption rate constant,  $V$  the volume of distribution,  $k$  the elimination rate constant. We also add several covariates to our model such as  $D$  the dose of drug administered,  $t$  the time at which measures are taken and the weight such as  $V$  is function of it. More precisely, the log-volume  $\log(V)$  is a linear function of the log-weight  $lw70 = \log(wt/70)$ . The final reads:

$$f(t, ka, V, k) = \frac{D ka}{V(ka - k)} (e^{-ka(t-T^{\text{lag}})} - e^{-k(t-T^{\text{lag}})}) , \quad (38)$$

Here,  $T^{\text{lag}}$ ,  $ka$ ,  $V$  and  $k$  are PK parameters that can change from one individual to another.

Let  $z_i = (T_i^{\text{lag}}, ka_i, V_i, k_i)$  be the vector of individual PK parameters for individual  $i$ . The model for the  $j$ -th measured concentration, noted  $y_{ij}$ , for individual  $i$  writes:

$$y_{ij} = f(t_{ij}, z_i) + \varepsilon_{ij} . \quad (39)$$

We assume in this example that the residual errors are independent and normally distributed with mean 0 and variance  $\sigma^2$ . Lognormal distributions are used for the three PK parameters:

$$\log(T_i^{\text{lag}}) \sim \mathcal{N}(\log(T_{\text{pop}}^{\text{lag}}), \omega_{T^{\text{lag}}}^2) \sim \mathcal{N}(\log(ka_{\text{pop}}), \omega_{ka}^2) , \quad (40)$$

$$\log(V_i) \sim \mathcal{N}(\log(V_{\text{pop}}), \omega_V^2) , \log(k_i) \sim \mathcal{N}(\log(k_{\text{pop}}), \omega_k^2) . \quad (41)$$

302 The complete model belongs to the curved exponential family, which vector of sufficient statistics  
 303  $S = (S_1(z), S_2(z), S_3(z))$  read:

$$S_1(z) = \frac{1}{n} \sum_{i=1}^n z_i, \quad S_2(z) = \frac{1}{n} \sum_{i=1}^n z_i^\top z_i, \quad S_3(z) = \frac{1}{n} \sum_{i=1}^n (y_i - f(t_i, z_i))^2 \quad (42)$$

304 where we have noted  $y_i$  and  $t_i$  the vector of observations and time for each patient  $i$ .

305 **Monte Carlo study:** We conduct a Monte  
 306 Carlo study to showcase the benefits of our  
 307 scheme.

308  $M = 50$  datasets have been simulated using  
 309 the following PK parameters values:  $T_{\text{pop}}^{\text{lag}} =$   
 310  $1$ ,  $ka_{\text{pop}} = 1$ ,  $V_{\text{pop}} = 8$ ,  $k_{\text{pop}} = 0.1$ ,  
 311  $\omega_{T^{\text{lag}}} = 0.4$ ,  $\omega_{ka} = 0.5$ ,  $\omega_V = 0.2$ ,  $\omega_k =$   
 312  $0.3$  and  $\sigma^2 = 0.5$ . We define the mean  
 313 square distance over the  $M$  replicates  $E_k(\ell) =$

314  $\frac{1}{M} \sum_{m=1}^M \left( \theta_k^{(m)}(\ell) - \theta^* \right)^2$  and plot it against

315 the epochs (passes over the data) Figure 2. Note that the MC-step (5) is performed using a Metropo-  
 316 lis Hastings procedure since the posterior distribution under the model  $\theta$  noted  $p(z_i|y_i, \theta)$  is in-  
 317 tractable due to the nonlinearity of the model (38). Figure 2 shows




Figure 2: TO COMPLETE

## 318 5 Conclusion

## References

- S. Allasonnière, Y. Amit, and A. Trouvé. Towards a coherent statistical framework for dense deformable template estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):3–29, 2007.
- S. Allasonnière, E. Kuhn, A. Trouvé, et al. Construction of bayesian deformable models via a stochastic approximation algorithm: a convergence study. *Bernoulli*, 16(3):641–678, 2010.
- C. Baey, S. Trevezas, and P.-H. Cournède. A non linear mixed effects model of plant growth and estimation via stochastic variants of the em algorithm. *Communications in Statistics-Theory and Methods*, 45(6):1643–1669, 2016.
- O. Cappé. Online em algorithm for hidden markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- O. Cappé and E. Moulines. On-line expectation–maximization algorithm for latent data models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):593–613, 2009.
- A. Chakraborty and K. Das. Inferences for joint modelling of repeated ordinal scores and time to event data. *Computational and mathematical methods in medicine*, 11(3):281–295, 2010.
- B. Delyon, M. Lavielle, and E. Moulines. Convergence of a stochastic approximation version of the em algorithm. *Ann. Statist.*, 27(1):94–128, 03 1999. doi: 10.1214/aos/1018031103. URL <https://doi.org/10.1214/aos/1018031103>.
- A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the royal statistical society. Series B (methodological)*, pages 1–38, 1977.
- J. P. Hughes. Mixed effects models with censored data with application to hiv rna levels. *Biometrics*, 55(2):625–629, 1999.
- B. Karimi. *Non-Convex Optimization for Latent Data Models: Algorithms, Analysis and Applications*. PhD thesis, 2019.
- B. Karimi, H.-T. Wai, É. Moulines, and M. Lavielle. On the global convergence of (fast) incremental expectation maximization methods. In *Advances in Neural Information Processing Systems*, pages 2833–2843, 2019.
- E. Kuhn, C. Matias, and T. Rebafka. Properties of the stochastic approximation em algorithm with mini-batch sampling. *arXiv preprint arXiv:1907.09164*, 2019.
- P. Liang and D. Klein. Online em for unsupervised models. In *Proceedings of human language technologies: The 2009 annual conference of the North American chapter of the association for computational linguistics*, pages 611–619, 2009.
- C. E. McCulloch. Maximum likelihood algorithms for generalized linear mixed models. *Journal of the American statistical Association*, 92(437):162–170, 1997.
- G. McLachlan and T. Krishnan. *The EM algorithm and extensions*, volume 382. John Wiley & Sons, 2007.
- R. M. Neal and G. E. Hinton. A view of the EM algorithm that justifies incremental, sparse, and other variants. In *Learning in graphical models*, pages 355–368. Springer, 1998.
- H. D. Nguyen, F. Forbes, and G. J. McLachlan. Mini-batch learning of exponential family finite mixture models. *Statistics and Computing*, pages 1–18, 2020.

- 360 S. J. Reddi, S. Sra, B. Póczos, and A. Smola. Fast incremental method for nonconvex optimization.  
361 *arXiv preprint arXiv:1603.06159*, 2016.
- 362 H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical statis-*  
363 *tics*, pages 400–407, 1951.
- 364 G. C. Wei and M. A. Tanner. A monte carlo implementation of the em algorithm and the poor man’s  
365 data augmentation algorithms. *Journal of the American statistical Association*, 85(411):699–704,  
366 1990.
- 367 R. Zhu, L. Wang, C. Zhai, and Q. Gu. High-dimensional variance-reduced stochastic gradient  
368 expectation-maximization algorithm. In *Proceedings of the 34th International Conference on*  
369 *Machine Learning-Volume 70*, pages 4180–4188. JMLR. org, 2017.

## A Proof of Lemma 2

**Lemma.** Assume H3, H4. For all  $\mathbf{s} \in \mathcal{S}$ ,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (43)$$

**Proof** Using H3 and the fact that we can exchange integration with differentiation and the Fisher's identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \left( \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \left( \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^\top \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (44)$$

Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top \mathbf{s}. \quad (45)$$

Taking the gradient of the above map w.r.t.  $\mathbf{s}$  and using assumption H3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left( \nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})}}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}). \quad (46)$$

The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))) \quad (47)$$

where we recall  $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left( \mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^\top$ . The proof of (43) follows directly from the assumption H4.  $\square$

## B Proof of Lemma 3

**Lemma.** Assume H??. The update (9) is equivalent to the following update on the resulting statistics

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}) \quad (48)$$

Also:

$$\mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n}\right) \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \quad (49)$$

where  $\bar{\mathbf{s}}^{(k)}$  is defined by (3) and  $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ .

**Proof** From update (9), we have:

$$\begin{aligned} \tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{\mathbf{S}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left( \tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned} \quad (50)$$

Since  $\tilde{S}_{i_k}^{(k+1)} = \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) + \eta_{i_k}^{(k+1)}$  we have

$$\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{\mathbf{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left( \tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)} \quad (51)$$

Taking the full expectation of both side of the equation leads to:

$$\begin{aligned} \mathbb{E} [\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] &= \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] \\ &\quad - \frac{1}{n} \mathbb{E} \left[ \mathbb{E} [\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}] \end{aligned} \quad (52)$$

The following equalities:

$$\mathbb{E} [\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E} [\bar{s}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k] = \bar{\mathbf{s}}^{(k)} \quad (53)$$

concludes the proof of the Lemma.  $\square$

## C Proof of Theorem 1

**Theorem.** Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of positive step sizes and consider the iSAEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = 1$  for any  $k > 0$ . We also set  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ . Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ .

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \quad (54)$$

**Proof** We begin our proof by giving this auxiliary Lemma setting an upper bound for the quantity  $\mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$

**Lemma 7.** For any  $k \geq 0$  and consider the iSAEM update in (9), it holds that

$$\begin{aligned} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &\leq 4\mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right] \\ &\quad + 2\frac{C_\eta}{M_k} + 4\mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \end{aligned} \quad (55)$$

**Proof** Applying the iSAEM update yields:

$$\begin{aligned} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] &= \mathbb{E} \left[ \left\| \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} - \frac{1}{n} (\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \tilde{S}_{i_k}^{(k)}) \right\|^2 \right] \\ &\leq 4\mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + 4\mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\quad + \frac{2}{n^2} \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] + 2\frac{C_\eta}{M_k} \end{aligned} \quad (56)$$

The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)} \right\|^2 \right] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)} \right\|^2 \right] \stackrel{(a)}{\leq} \frac{2L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \right\|^2 \right], \quad (57)$$

where (a) is due to Lemma 1 and which concludes the proof of the Lemma.  $\square$

Under the smoothness of the Lyapunov function  $V$  (cf. Lemma 1), we can write:

$$V(\hat{\mathbf{s}}^{(k+1)}) \leq V(\hat{\mathbf{s}}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \quad (58)$$

Taking the expectation on both sides yields:

$$\mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k+1)}) \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(k)}) \right] + \gamma_{k+1} \mathbb{E} \left[ \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (59)$$



403 Using Lemma 3, we obtain:

$$\begin{aligned}
& \mathbb{E} \left[ \langle \tilde{S}^{(k+1)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] = \\
& \mathbb{E} \left[ \langle \bar{s}^{(k)} - \hat{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] + \left( 1 - \frac{1}{n} \right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \left( 1 - \frac{1}{n} \right) \mathbb{E} \left[ \left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[ \langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{s}^{(k)}) \rangle \right] \\
& \stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[ \left\| \nabla V(\hat{s}^{(k)}) \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& \stackrel{(a)}{\leq} \left( v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{60}$$

404 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with  $\beta \rightarrow 1$ ). Note

405  $a_k = \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$  and

$$\begin{aligned}
a_k \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned} \tag{61}$$

406 We now give an upper bound of  $\mathbb{E} \left[ \left\| \tilde{S}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right]$  using Lemma 7 and plug it into (61):

$$\begin{aligned}
(a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[ \left\| \bar{s}^{(k)} - \hat{s}^{(k)} \right\|^2 \right] & \leq \mathbb{E} \left[ V(\hat{s}^{(k)}) - V(\hat{s}^{(k+1)}) \right] \\
& + \gamma_{k+1} \left( \frac{1}{2\beta} \left( 1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{s}^{(k)} \right\|^2 \right] \\
& + \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
& + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right]
\end{aligned} \tag{62}$$

407 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^{k+1})} \right\|^2 \right] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(k)} \right\|^2 \right] + \frac{n-1}{n} \mathbb{E} \left[ \left\| \hat{s}^{(k+1)} - \hat{s}^{(\tau_i^k)} \right\|^2 \right] \right) \tag{63}$$

408 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. For any  $\beta > 0$ , it holds

$$\begin{aligned}
& \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 - 2\gamma_{k+1}\langle \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\
&\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2 + \gamma_{k+1}\beta\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right]
\end{aligned} \tag{64}$$

409 where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2\right]
\end{aligned} \tag{65}$$

410 Observe that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)})$ . Applying Lemma 7 yields

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\
&\leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\
&\leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\
&\quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\
&\quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right]
\end{aligned} \tag{66}$$

411 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] \tag{67}$$

412 From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} &\leq (1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}))\Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
&\quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right]
\end{aligned} \tag{68}$$

413 Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{8, 1 + 6v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,

414  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$ ,  $\alpha \geq 8$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1} \tag{69}$$

415 which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} =$   
 416  $\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} \leq & 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E} \left[ \left\| \eta_{i_\ell}^{(\ell)} \right\|^2 \right] \\ & + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)} \right\|^2 \right] \end{aligned} \quad (70)$$

417 Note  $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$  Summing on both sides over  $k = 0$  to  $k = K_{\max} - 1$  yields:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} \\ &= 4 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_{\max}-1} (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E} \left[ \left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\ &+ \sum_{k=0}^{K_{\max}-1} 4 (\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \omega_{k,1} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &\leq \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \end{aligned} \quad (71)$$

418 We recall (62) where we have summed on both sides from  $k = 0$  to  $k = K_{\max} - 1$ :

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left( \frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)} \end{aligned} \quad (72)$$

419 Plugging (71) into (72) results in:

$$\begin{aligned} & \sum_{k=0}^{K_{\max}-1} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\beta}_k \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[ V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\ &+ \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[ \left\| \eta_{i_k}^{(k)} \right\|^2 \right] \end{aligned} \quad (73)$$

420 where:

$$\begin{aligned}\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\beta}_k &= \gamma_{k+1} \left( \frac{1}{2\beta} (1 - \frac{1}{n}) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}} \\ \tilde{\Gamma}_k &= \gamma_{k+1} \left( \gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta})}{\Lambda_{(k+1)}}\end{aligned}$$

421 and

$$\begin{aligned}a_k &= \gamma_{k+1} \left( v_{\min} - v_{\max}^2 \frac{\beta(n-1) + 1}{2n} \right) \\ \Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1} L_s^2}{n^2} (\gamma_{k+1} + \frac{1}{\beta}) \\ c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{L_s, L_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}\end{aligned}$$

422 When, for any  $k > 0$ ,  $\tilde{\alpha}_k \geq 0$ , we have by Lemma 2 that:

$$\sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[ \left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \quad (74)$$

423 which yields an upper bound of the gradient of the Lyapunov function  $V$  along the path of the  
424 iSAEM update and concludes the proof of the Theorem.  $\square$

## 425 D Proofs of Auxiliary Lemmas

### 426 D.1 Proof of Lemma 4 and Lemma 5

427 **Lemma.** For any  $k \geq 0$  and consider the vrTTSEM update in (10) with  $\rho_k = \rho$ , it holds for all  
428  $k > 0$

$$\begin{aligned} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (75)$$

429 where we recall that  $\ell(k)$  is the first iteration number in the epoch that iteration  $k$  is in.

430 **Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful through-  
431 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left( (1-\rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \end{aligned} \quad (76)$$

432 We observe, using the identity (76), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] \quad (77)$$

433 For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{\ell(k)}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{\ell(k)}) \right\|^2 \right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \stackrel{(b)}{\leq} L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{\ell(k)}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (78)$$

434 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (77) proves the  
435 lemma.  $\square$

436 **Lemma.** For any  $k \geq 0$  and consider the fiTTSEM update in (11) with  $\rho_k = \rho$ , it holds for all  $k > 0$   
437

$$\begin{aligned} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_k)}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned} \quad (79)$$

438 **Proof** Beforehand, we provide a rewriting of the quantity  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$  that will be useful through-  
439 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) \\ &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left( (1-\rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \\ &= -\gamma_{k+1} \left( (1-\rho) \left[ \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[ \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_k)}) \right] \right) \end{aligned} \quad (80)$$

440 We observe, using the identity (80), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{((k))} - \tilde{S}^{(k)}\|^2] \quad (81)$$

441 For the latter term, we obtain its upper bound as

$$\begin{aligned}\mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{S}}_i^{(k)}) - (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})\right\|^2\right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]\end{aligned}\quad (82)$$

442 where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \quad (83)$$

443 Substituting into (81) proves the lemma.  $\square$

## 444 D.2 Proof of Lemma 6

445 **Lemma.** Consider a decreasing stepsize  $\gamma_k \in (0, 1)$  and a constant  $\rho$ , then the following inequality  
446 holds:

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_\ell)^2 (\mathbf{S}^{(\ell)} - \tilde{\mathbf{S}}^{(\ell)}) \quad (84)$$

447 where  $\mathbf{S}^{(k)}$  is defined either by (11) (fTTSEM) or (10) (vrTTSEM)

448 **Proof** We begin by writing the two-time-scale update:

$$\begin{aligned}\tilde{\mathbf{S}}^{(k+1)} &= \tilde{\mathbf{S}}^{(k)} + \rho(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k)}) \\ \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)})\end{aligned}\quad (85)$$

449 where  $\mathbf{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(t_i^k)} + (\tilde{\mathbf{S}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)})$  according to (11). Denote  $\delta^{(k+1)} = \hat{\mathbf{s}}^{(k+1)} -$   
450  $\tilde{\mathbf{S}}^{(k+1)}$ . Then from (85), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1-\rho} (1 - \gamma_{k+1})(\mathbf{S}^{(k+1)} - \tilde{\mathbf{S}}^{(k+1)}) \quad (86)$$

451 Using the telescoping sum and noting that  $\delta^{(0)} = 0$ , we have

$$\delta^{(k+1)} \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1 - \gamma_{\ell+1})^2 (\mathbf{S}^{(\ell+1)} - \tilde{\mathbf{S}}^{(\ell+1)}) \quad (87)$$

452  $\square$

## 453 D.3 Additional Intermediary Result

454 **Lemma 8.** At iteration  $k + 1$ , the drift term of update (11), with  $\rho_{k+1} = \rho$ , is equivalent to the  
455 following :

$$\begin{aligned}\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{\mathbf{S}}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1 - \rho) (\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)})\end{aligned}\quad (88)$$

456 where we recall that  $\eta_{i_k}^{(k+1)}$ , defined in (19), which is the gap between the MC approximation and  
457 the expected statistics.



458 **Proof** Using the fTTSEM update  $\tilde{S}^{(k+1)} = (1-\rho)\tilde{S}^{(k)} + \rho\mathcal{S}^{(k+1)}$  where  $\mathcal{S}^{(k+1)} = \overline{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$  leads to the following decomposition:

$$\begin{aligned}
& \tilde{S}^{(k+1)} - \hat{s}^{(k)} \\
&= (1-\rho)\tilde{S}^{(k)} + \rho\left(\overline{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right) - \hat{s}^{(k)} + \rho\overline{\mathcal{S}}^{(k)} - \rho\overline{\mathcal{S}}^{(k)} \\
&= \rho(\overline{\mathcal{S}}^{(k)} - \hat{s}^{(k)}) + \rho(\tilde{S}_{i_k}^{(k)} - \overline{\mathcal{S}}_{i_k}^{(k)}) + (1-\rho)\left(\tilde{S}^{(k)} - \hat{s}^{(k)}\right) + \rho\left(\overline{\mathcal{S}}^{(k)} - \overline{\mathcal{S}}^{(k)} + (\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})\right) \\
&= \rho(\overline{\mathcal{S}}^{(k)} - \hat{s}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho\left[(\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}]\right] \\
&+ (1-\rho)\left(\tilde{S}^{(k)} - \hat{s}^{(k)}\right)
\end{aligned}$$

460 where we observe that  $\mathbb{E}[\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] = \overline{\mathcal{S}}^{(k)} - \overline{\mathcal{S}}^{(k)}$  and which concludes the proof.

461 *Important Note:* Note that  $\overline{\mathcal{S}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not equal to  $\eta_{i_k}^{(k+1)}$ , defined in (19), which is the gap  
462 between the MC approximation and the expected statistics. Indeed  $\tilde{S}_{i_k}^{(t_{i_k}^k)}$  is not computed under the  
463 same model as  $\overline{\mathcal{S}}_{i_k}^{(k)}$ . □

## 464 E Proof of Theorem 2

465 **Theorem.** Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of  
 466 positive step sizes and consider the vrTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for  
 467 any  $k > 0$ .

468 Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\bar{L} = \max\{L_S, L_V\}$ ,  $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$   
 469 and a constant  $\mu \in (0, 1)$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (89)$$

470 **Proof** Using the smoothness of  $V$  and update (10), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (90)$$

471 Denote  $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fiTTSEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 472 Taking expectations on both sides show that

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}\rho \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\langle \eta_{i_k}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \gamma_{k+1}(1-\rho) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \quad (91)$$

473 where we have used (76) in (a) and  $\mathbb{E}[\mathbf{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$  in (b), the growth condition in  
 474 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

475 Furthermore, for  $k+1 \leq \ell(k) + m$  (i.e.,  $k+1$  is in the same epoch as  $k$ ), we have

$$\begin{aligned} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 \\ &\quad - 2\gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1-\rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle] \\ &\leq \mathbb{E}[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2], \end{aligned} \quad (92)$$

476 where we first used (76) and the last inequality is due to the Young's inequality.

477 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \quad (93)$$

478 where  $b_k := \bar{b}_{k \bmod m}$  is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0. \quad (94)$$

479 Note that  $\bar{b}_i$  is decreasing with  $i$  and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2}, \quad i = 1, 2, \dots, m. \quad (95)$$

480 For  $k+1 \leq \ell(k) + m$ , we have the following inequality

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[\rho \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\frac{\gamma_{k+1}\rho}{\beta} \left\|\eta_{i_k}^{(k+1)}\right\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned} \quad (96)$$

481 And using Lemma 4 we obtain:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2 - \gamma_{k+1}^2\rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \|\mathbf{h}_k\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[(\rho + \rho^2\gamma_{k+1} L_V) \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho - (1-\rho)^2\gamma_{k+1} L_V) \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \left\|\eta_{i_k}^{(k+1)}\right\|^2 + \left(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2\right) \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned} \quad (97)$$

482 Rearranging the terms yields:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\quad + \underbrace{\left(b_{k+1}(1 + \gamma\beta + 2\gamma^2\rho^2 L_{\mathbf{s}}^2) + \gamma^2\rho^2 L_V L_{\mathbf{s}}^2\right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)} \end{aligned} \quad (98)$$

483 where

$$\begin{aligned} \tilde{\eta}^{(k+1)} &= \left(\gamma_{k+1}(\rho + \rho^2\gamma_{k+1} L_V) + b_{k+1}(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2)\right) \mathbb{E}\left[\left\|\eta_{i_k}^{(k+1)}\right\|^2\right] \\ \chi^{(k+1)} &= \left(b_{k+1}(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2) - \gamma_{k+1}(1-\rho - (1-\rho)^2\gamma_{k+1} L_V)\right) \\ \tilde{\chi}^{(k+1)} &= \chi^{(k+1)} \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned} \quad (99)$$

484 This leads, using Lemma 2, that for any  $\gamma_{k+1}$ ,  $\rho$  and  $\beta$  such that  $\rho v_{\min} + v_{\max}^2 -$   
485  $\gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2) > 0$ ,

$$\begin{aligned} v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \\ &\quad + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \end{aligned} \quad (100)$$

486 We first remark that

$$\begin{aligned} & \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \\ & \geq \frac{\gamma_{k+1}\rho}{c_1}(1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1)) \end{aligned} \quad (101)$$

487 where  $c_1 = v_{\min}^{-1}$ . By setting  $\bar{L} = \max\{L_s, L_V\}$ ,  $\beta = \frac{c_1\bar{L}}{n^{1/3}}$ ,  $\rho = \frac{\mu}{c_1\bar{L}n^{2/3}}$ ,  $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$  and  
 488  $\{\gamma_{k+1}\}$  any sequence of decreasing stepsizes in  $(0, 1)$ , it can be shown that there exists  $\mu \in (0, 1)$ ,  
 489 such that the following lower bound holds

$$\begin{aligned} & 1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1) \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V\mu^2}{c_1^2n^{\frac{4}{3}}}\frac{(1 + \gamma\beta + 2\gamma^2L_s^2)^m - 1}{\gamma\beta + 2\gamma^2L_s^2}(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2}(e - 1)(1 + \frac{2\mu}{n}) \geq 1 - \mu - \mu(1 + 2\mu)\frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2} \end{aligned} \quad (102)$$

490 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2L_s^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \text{ and } (1 + \gamma\beta + 2\gamma^2L_s^2)^m \leq e - 1. \quad (103)$$

491 and the required  $\mu$  in (b) can be found by solving the quadratic equation.

492 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_{\max}})}{v_{\min}\rho} + 2 \sum_{k=0}^{K_{\max}-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min}\rho} \quad (104)$$

493 Note that  $R_0 = \mathbb{E}[V(\hat{s}^{(0)})]$  and if  $K_{\max}$  is a multiple of  $m$ , then  $R_{\max} = \mathbb{E}[V(\hat{s}^{(K_{\max})})]$ . Under the  
 494 latter condition, we have

$$\sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}] \quad (105)$$

495 This concludes our proof.

496 □

497 **F Proof of Theorem 3**

498 **Theorem.** Assume H1-H5. Let  $K_{\max}$  be a positive integer. Let  $\{\gamma_k, k \in \mathbb{N}\}$  be a sequence of  
 499 positive step sizes and consider the fiTTSEM sequence  $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$  obtained with  $\rho_{k+1} = \rho$  for  
 500 any  $k > 0$ .

501 Assume that  $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$  for any  $k \leq K_{\max}$ . By setting  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{L_s, L_V\}$ ,  
 502  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[ \Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned} \quad (106)$$

503 **Proof** Using the smoothness of  $V$  and update (11), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (107)$$

504 Denote  $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$  the drift term of the fiTTSEM update in (7) and  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ .  
 505 Using Lemma 8 and the additional following identity:

$$\mathbb{E} \left[ (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(k)}] \right] = 0 \quad (108)$$

506 we have:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \mathbb{E} \left[ \langle \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1 - \rho) \mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(a)}{\leq} -v_{\min} \gamma_{k+1} \rho \mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1} \mathbb{E} \left[ \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(b)}{\leq} -(v_{\min} \gamma_{k+1} \rho + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \end{aligned} \quad (109)$$

507 where  $\xi^{(k+1)} = \mathbb{E} \left[ \left\| \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] \right\|^2 \right]$ .

508 **Bounding**  $\mathbb{E}[\|\mathbf{H}_{k+1}\|^2]$  Using Lemma 5, we obtain:

$$\begin{aligned} &\gamma_{k+1} (v_{\min} \rho + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\leq \mathbb{E} [V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)})] + \tilde{\xi}^{(k+1)} + \left( (1 - \rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V \rho^2 L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (110)$$

509 where  $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 \mathbb{L}_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1}\rho^2}{2} \xi^{(k+1)}$ . Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (111)$$

510 where the equality holds as  $i_k$  and  $j_k$  are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \end{aligned} \quad (112)$$

511 Note that  $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}\mathbf{H}_{k+1}$  and that in expectation we recall  
 512 that  $\mathbb{E}[\mathbf{H}_{k+1}|\mathcal{F}_k] = \rho\mathbf{h}_k + \rho\mathbb{E}[\eta_{i_k}^{(k+1)}|\mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}]$  where  $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$ . Thus,  
 513 for any  $\beta > 0$ , it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (113)$$

514 where the last inequality is due to the Young's inequality. Plugging this into (111) yields:

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (114)$$

515 Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2] \\ &\quad + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned} \quad (115)$$



516 We now use Lemma 5 on  $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2 \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2$  and obtain:

$$\begin{aligned}
& \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\
& \leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n} + \frac{(n-1)(1+\gamma_{k+1}\beta)}{n^2}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\
& \leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{116}$$

517 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \tag{117}$$

518 From the above, we get

$$\begin{aligned}
\Delta^{(k+1)} & \leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
& + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]
\end{aligned} \tag{118}$$

519 Setting  $c_1 = v_{\min}^{-1}$ ,  $\alpha = \max\{2, 1 + 2v_{\min}\}$ ,  $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,  $\beta = \frac{c_1 \bar{L}}{n}$ ,  
520  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ , we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{4/3}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1} \tag{119}$$

521 which shows that  $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \in (0, 1)$  for any  $k > 0$ . Denote  $\Lambda_{(k+1)} = \frac{1}{n} -$   
522  $\gamma_{k+1}\beta - \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2$  and note that  $\Delta^{(0)} = 0$ , thus the telescoping sum yields:

$$\begin{aligned}
\Delta^{(k+1)} & \leq \sum_{\ell=0}^k \omega_{k,\ell} \left(2\gamma_{\ell+1}^2 \rho^2 + \frac{\gamma_{\ell+1} \rho^2}{\beta}\right) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] \\
& + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left(2\gamma_{\ell+1} + \frac{1}{\beta}\right) \mathbb{E}[\|\tilde{\mathbf{S}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)}
\end{aligned} \tag{120}$$

523 where  $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$  and  $\tilde{\epsilon}^{(\ell+1)} = \left(2\gamma_{\ell+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(\ell+1)}\|^2]$ .

524 Summing on both sides over  $k = 0$  to  $k = K_{\max} - 1$  yields:

$$\begin{aligned}
\sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} & \leq \sum_{k=0}^{K_{\max}-1} \frac{2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1} \rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\
& + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}
\end{aligned} \tag{121}$$

525 We recall (110) where we have summed on both sides from  $k = 0$  to  $k = K_{\max} - 1$ :

$$\begin{aligned}
& \mathbb{E}[V(\hat{\mathbf{s}}^{(K_{\max})}) - V(\hat{\mathbf{s}}^{(0)})] \\
& \leq \sum_{k=0}^{K_{\max}-1} \left\{ \gamma_{k+1}(-(\nu_{\min}\rho + \nu_{\max}^2) + \gamma_{k+1}\rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] + \gamma^2 L_V \rho^2 L_{\mathbf{s}}^2 \Delta^{(k)} \right\} \\
& + \sum_{k=0}^{K_{\max}-1} \left\{ \tilde{\xi}^{(k+1)} + \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right\} \\
& \leq \sum_{k=0}^{K_{\max}-1} \left\{ -\gamma_{k+1}(\nu_{\min}\rho + \nu_{\max}^2) + \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left( 2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right\} \mathbb{E}[\|\mathbf{h}_k\|^2] \\
& + \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]
\end{aligned} \tag{122}$$

where

$$\Xi^{(k+1)} = \tilde{\xi}^{(k+1)} + \frac{\gamma_{k+1}^3 L_V \rho^2 L_{\mathbf{s}}^2}{\Lambda_{(k+1)}} \epsilon^{(k+1)}$$

and

$$\Gamma_{k+1} = \left( (1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_{\mathbf{s}}^2 (1-\rho)^2 \left( 2\gamma_{k+1} + \frac{1}{\beta} \right)}{\Lambda_{(k+1)}}$$

526 We now analyse the following quantity

$$\begin{aligned}
& -\gamma_{k+1}(\nu_{\min}\rho + \nu_{\max}^2) + \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left( 2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\
& = -\gamma_{k+1} \left[ (\nu_{\min}\rho + \nu_{\max}^2) + \gamma_{k+1} \rho^2 L_V + \frac{\rho^2 \gamma_{k+1} L_V L_{\mathbf{s}}^2 \left( 2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right]
\end{aligned} \tag{123}$$

527 Furthermore, we recall that  $c_1 = \nu_{\min}^{-1}$ ,  $\alpha = \max\{2, 1+2\nu_{\min}\}$ ,  $\bar{L} = \max\{L_{\mathbf{s}}, L_V\}$ ,  $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$ ,

528  $\beta = \frac{c_1 \bar{L}}{n}$ ,  $\rho = \frac{1}{n^{2/3}}$ ,  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ ,  $\alpha \geq 2$ . Then,

$$\begin{aligned}
& \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_{\mathbf{s}}^2 \left( 2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\frac{1}{n} - \gamma_{k+1}\beta - \gamma_{k+1}^2 \rho^2 L_{\mathbf{s}}^2} \\
& \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} (k^2 \alpha^2 c_1^2 n^{4/3})^{-1} \left( \frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1^2 \bar{L}^2 n^{1/3}} \right)}{\frac{1}{n} - \frac{1}{k \alpha n} - \frac{1}{k^2 \alpha^2 c_1^2 n^{4/3}}} \\
& = \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left( \frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k \alpha c_1^2 \bar{L}^2 n^{1/3}} \right)}{(k \alpha c_1 n^{1/3})(k \alpha - 1) c_1 - 1} \\
& \stackrel{(a)}{\leq} \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k \alpha c_1^2 \bar{L} n^{1/3}} \left( \frac{2}{k \alpha n} + 1 \right)}{2(\alpha c_1 n^{1/3}) - 1} \\
& \leq \frac{1}{k^2 \alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{1}{k \alpha^2 c_1^3 \bar{L} n^{2/3}} \\
& \leq \frac{1}{\alpha c_1 \bar{L} n^{2/3}}
\end{aligned} \tag{124}$$

529 where (a) is due to  $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$  and  $k \alpha c_1 n^{1/3} \geq 1$ . Also, since  $-\gamma_{k+1}(\nu_{\min}\rho +$   
530  $\nu_{\max}^2) \leq -\gamma_{k+1}\rho \nu_{\min} = -$

531 Using the Lemma 2, we know that  $v_{\max}^2 \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \leq \|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2$  and using (124) on (122)  
 532 yields:

$$\begin{aligned} v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min}} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned} \quad (125)$$

533 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[ \left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned} \quad (126)$$

534

□

## G Practical Implementations of Two-Time-Scale EM Methods

### G.1 Application on GMM

We first recognize that the constraint set for  $\theta$  is given by

$$\Theta = \Delta^M \times \mathbb{R}^M. \quad (127)$$

Using the partition of the sufficient statistics as  $S(y_i, z_i) = (S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ , the partition  $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$  and the fact that  $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$ , the complete data log-likelihood can be expressed as in (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i) y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (128)$$

and  $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$ . We also define for each  $m \in \llbracket 1, M \rrbracket$ ,  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ . Consider the following latent sample used to compute an approximation of the conditional expected value  $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}} | y = y_i]$ :

$$z_{i,m} \sim \mathbb{P}(z_i = m | y_i; \theta) \quad (129)$$

where  $m \in \llbracket 1, M \rrbracket$ ,  $i \in \llbracket 1, n \rrbracket$  and  $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$ .

In particular, given iteration  $k + 1$ , the computation of the approximated quantity  $\tilde{S}_{i_k}^{(k)}$  during Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left( \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:= \tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}) y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1}) y_{i_k}}_{:= \tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:= \tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (130)$$

Recall that we have used the following regularizer:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (131)$$

It can be shown that the regularized M-step in (4) evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\boldsymbol{\mu}}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (132)$$

where we have defined for all  $m \in \llbracket 1, M \rrbracket$  and  $j \in \llbracket 1, 3 \rrbracket$ ,  $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$ .

### G.2 Model Assumptions (GMM example)

We use the GMM example to illustrate the required assumptions.

Many practical models can satisfy the compactness of the sets as in Assumption H1. For instance, the GMM example satisfies (16) as the sufficient statistics are composed of indicator functions and observations as defined Section G.1 Equation (128).

Assumptions H2 and H3 are standard for the curved exponential family models. For GMM, the following (strongly convex) regularization  $\mathbf{r}(\theta)$  ensures H3:

$$\mathbf{r}(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m)$$

557 since it ensures  $\theta^{(k)}$  is unique and lies in  $\text{int}(\Delta^M) \times \mathbb{R}^M$ . We remark that for H2, it is possible to  
 558 define the Lipschitz constant  $L_p$  independently for each data  $y_i$  to yield a refined characterization.

559 Again, H4 is satisfied by practical models. For GMM, it can be verified by deriving the closed form  
 560 expression for  $B(s)$  and using H1.

561 Under H1 and H3, we have  $\|\hat{s}^{(k)}\| < \infty$  since  $S$  is compact and  $\hat{\theta}^{(k)} \in \text{int}(\Theta)$  for any  $k \geq 0$  which  
 562 thus ensure that the EM methods operate in a closed set throughout the optimization process.

### 563 G.3 Algorithms updates

564 In the sequel, recall that, for all  $i \in \llbracket n \rrbracket$  and iteration  $k$ , the computed statistic  $\tilde{S}_{i_k}^{(k)}$  is defined by  
 565 (130). At iteration  $k$ , the several E-steps defined by (9) or (10) and (11) leads to the definition of the  
 566 quantity  $\hat{s}^{(k+1)}$ . For the GMM example, after the initialization of the quantity  $\hat{s}^{(0)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(0)}$ ,  
 567 those E-steps break down as follows:

568 **Batch EM (EM):** for all  $i \in \llbracket 1, n \rrbracket$ , compute  $\bar{s}_i^{(k)}$  and set

$$\hat{s}^{(k+1)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(k)}. \quad (133)$$

569 where  $\bar{s}_i^{(k)}$  are computed using the exact conditional expected value  $\mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i]$ :

$$\tilde{\omega}_m(y_i; \theta) := \mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)}, \quad (134)$$

570 **Incremental EM (iEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}. \quad (135)$$

571 **batch SAEM (SAEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}. \quad (136)$$

572 where  $\tilde{S}^{(k)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$  with  $\tilde{S}_i^{(k)}$  defined in (130).

573 **Incremental SAEM (iSAEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

574

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_i^k)})). \quad (137)$$

575 **Variance Reduced Two-Time-Scale EM (vrTTSEM):** draw an index  $i_k$  uniformly at random on

576  $\llbracket n \rrbracket$ , compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} (1 - \rho) + \rho (\tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}))). \quad (138)$$

577 **Fast Incremental Two-Time-Scale EM (fiTTSEM):** draw an index  $i_k$  uniformly at random on  $\llbracket n \rrbracket$ ,

578 compute  $\bar{s}_{i_k}^{(k)}$  and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} (1 - \rho) + \rho (\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}))). \quad (139)$$

579 Finally, the  $k$ -th update reads  $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$  where the function  $s \rightarrow \bar{\theta}(s)$  is defined by (132).