
Fast Two-Timescale Stochastic EM Algorithms

Anonymous Author(s)

Affiliation

Address

email

Abstract

Using the Expectation-Maximization (EM) algorithm is the most popular choice for current latent data model learning tasks. For today's modern and complex models, variants of the EM have been initially introduced by [20], using incremental updates to scale to large datasets, and by [24, 8], using Monte-Carlo (MC) approximations to bypass the impossible conditional expectation of the latent data for most nonconvex models. In this paper, we propose a general class of methods called Two-Timescale EM Methods based on double stages of stochastic updates to tackle an essential large and nonconvex optimization task for latent data models. We motivate the choice of a double dynamics by invoking the variance reduction virtue of each stage of the method on both sources of noise: the incremental update and the MC approximation. We establish finite-time and global convergence bounds for nonconvex objective functions. Numerical applications are also presented in this article to illustrate our findings.

1 Introduction

Learning latent data models is critical for modern machine learning problems, see [18] for references. We formulate the training of such model as an empirical risk minimization problem:

$$\min_{\theta \in \Theta} \bar{L}(\theta) := r(\theta) + L(\theta) \quad \text{with} \quad L(\theta) = \frac{1}{n} \sum_{i=1}^n L_i(\theta) := \frac{1}{n} \sum_{i=1}^n \{ -\log g(y_i; \theta) \}, \quad (1)$$

We denote the observations by $\{y_i\}_{i=1}^n$, $\Theta \subset \mathbb{R}^d$ is the convex parameters space. We consider a smooth convex regularization noted $r : \Theta \rightarrow \mathbb{R}$ and $g(y; \theta)$ is the (incomplete) likelihood of each observation. The objective function $\bar{L}(\theta)$ is possibly *nonconvex* and is assumed to be lower bounded.

In the latent variable model, $g(y_i; \theta)$, is the marginal of the complete data likelihood defined as $f(z_i, y_i; \theta)$, i.e. $g(y_i; \theta) = \int_{\mathcal{Z}} f(z_i, y_i; \theta) \mu(dz_i)$, where $\{z_i\}_{i=1}^n$ are the latent variables. In this paper, we make the assumption of a complete model belonging to the curved exponential family:

$$f(z_i, y_i; \theta) = h(z_i, y_i) \exp(\langle S(z_i, y_i) | \phi(\theta) \rangle - \psi(\theta)), \quad (2)$$

where $\psi(\theta)$, $h(z_i, y_i)$ are scalar functions, $\phi(\theta) \in \mathbb{R}^k$ is a vector function, and $S(z_i, y_i) \in \mathbb{R}^k$ is the complete data sufficient statistics. Full batch EM [9] is the method of reference for that kind of task and is a two steps procedure. The **E-step** amounts to computing the conditional expectation of the complete data sufficient statistics,

$$\bar{s}(\theta) = \frac{1}{n} \sum_{i=1}^n \bar{s}_i(\theta) \quad \text{where} \quad \bar{s}_i(\theta) = \int_{\mathcal{Z}} S(z_i, y_i) p(z_i | y_i; \theta) \mu(dz_i). \quad (3)$$

The **M-step** is given by

$$\text{M-step: } \hat{\theta} = \bar{\theta}(\bar{s}(\theta)) := \arg \min_{\vartheta \in \Theta} \{ r(\vartheta) + \psi(\vartheta) - \langle \bar{s}(\theta) | \phi(\vartheta) \rangle \}, \quad (4)$$

Two caveats of this method are the following: (a) with the explosion of data, the first step of the EM is computationally inefficient as it requires a full pass over the dataset at each iteration and (b) the complexity of modern models makes the expectation in (3) intractable. So far, both challenges have been addressed separately, to the best of our knowledge, as detailed the sequel.

Prior Work Inspired by stochastic optimization procedures, [20] and [5] developed respectively an incremental and an online variant of the E-step in models where the expectation is computable then extensively used and studied in [21, 15, 4]. Some improvements of that methods have been provided and analyzed, globally and in finite-time, in [12] where variance reduction techniques taken from the optimization literature have been efficiently applied to scale the EM algorithm to large datasets.

Regarding the computation of the expectation under the posterior distribution, the first method was the Monte-Carlo EM (MCEM) introduced in the seminal paper [24] where a MC approximation for this expectation is computed. A variant of that method is the Stochastic Approximation of the EM (SAEM) in [8] leveraging the power of Robbins-Monro type of update [23] to ensure pointwise convergence of the vector of estimated parameters rather using a decreasing stepsize than increasing the number of MC samples. The MCEM and the SAEM have been successfully applied in mixed effects models [17, 10, 3] or to do inference for joint modeling of time to event data coming from clinical trials in [7], among other applications. Recently, an incremental variant of the SAEM was proposed in [14] showing positive empirical results but its analysis is limited to asymptotic consideration. Gradient-based methods have been developed and analyzed in [25] but they remain out of the scope of this paper as they tackle the high-dimensionality issue.

Contributions This paper *introduces* and *analyzes* a new class of methods which purpose is to update two proxies for target expected quantities in a two-timescale manner. Those approximated quantities are then used to optimize (1) for modern examples and settings using EM Maximization step. The main contributions of the paper are:

- We propose a two-timescale method based on Stochastic Approximation (SA), to alleviate the problem of MC computation, and on Incremental updates, to scale to large datasets. We describe in details the edges of each level of our method based on variance reduction arguments. The derivation of such class of algorithms has two advantages. First, it naturally leverages variance reduction and Robbins-Monro type of updates to tackle large-scale and highly nonlinear learning tasks. Then, it gives a simple formulation as a *scaled-gradient method* which makes the global analysis and the implementation accessible.
- We also establish global (independent of the initialization) and finite-time (true at each iteration) upper bounds on a classical suboptimality condition in the nonconvex literature, *i.e.*, the second order moment of the gradient of the objective function.

In Section 2 we formalize both incremental and Monte-Carlo variants of the EM. Then, we introduce our two-timescale class of EM algorithms for which we derive several global statistical guarantees in Section 3 for possibly *nonconvex* functions. Section 4 is devoted to numerical illustrations.

2 Two-Timescale Stochastic EM Algorithms

We recall and formalize in this section the different methods found in the literature that aim to solving the large-scale problem and the intractable expectation. We then provide the general framework of our method that efficiently tackles the optimization problem (1).

2.1 Monte Carlo Integration and Stochastic Approximation

As mentioned in the introduction, for complex and possibly nonlinear models, the expectation under the posterior distribution defined in (3) is not tractable. In that case, the first solution involves computing a Monte Carlo integration of that latter term. For all $i \in \llbracket 1, n \rrbracket$, draw for $m \in \llbracket 1, M \rrbracket$, samples $z_{i,m} \sim p(z_i | y_i; \theta)$ and compute the MC integration \tilde{s} of the deterministic quantity $\bar{s}(\theta)$:

$$\text{MC-step : } \tilde{s} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}, y_i), \quad (5)$$

and then update the parameter $\hat{\theta} = \bar{\theta}(\tilde{s})$. This algorithm bypasses the intractable expectation issue but is rather computationally expensive in order to reach point wise convergence (M needs to be

large). An alternative to that stochastic algorithm is to use a Robbins-Monro (RM) type of update. We denote, at iteration k , the following quantity

$$\tilde{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \frac{1}{M} \sum_{m=1}^M S(z_{i,m}^{(k)}, y_i) \quad \text{where} \quad z_{i,m}^{(k)} \sim p(z_i | y_i; \theta^{(k)}) . \quad (6)$$

Then, the RM updated of the sufficient statistics $\hat{s}^{(k+1)}$ reads:

$$\text{SA-step : } \hat{s}^{(k+1)} = \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) , \quad (7)$$

where $\{\gamma_k\}_{k \geq 1} \in (0, 1)$ is a sequence of decreasing step sizes to ensure asymptotic convergence. This is called the Stochastic Approximation of the EM (SAEM) and has been shown to converge to a maximum likelihood of the observations under very general conditions [8]. In the simulation step (6), since the loss function between the observed data y_i and the latent variable z_i can be nonconvex, sampling from the posterior distribution $p(z_i | y_i; \theta)$, under the current model θ , requires using an inference algorithm. [13] proved almost sure convergence of the sequence of parameters obtained by this algorithm coupled with an MCMC procedure during the simulation step. In simple scenarios, the samples $\{z_{i,m}\}_{m=0}^{M-1}$ are conditionally independent and identically distributed with distribution $p(z_i, \theta)$. Nevertheless, in most cases, sampling exactly from this distribution is not an option and the Monte Carlo batch is sampled by Monte Carlo Markov Chains (MCMC) algorithm. In the SA-step, the sequence of decreasing positive integers $\{\gamma_k\}_{k \geq 1}$ controls the convergence of the algorithm. In practice, γ_k is set equal to 1 during the first few iterations to let the algorithm explore the parameter space without memory and converge quickly to a neighborhood of the target estimate. The Stochastic Approximation is performed during the remaining iterations where $\gamma_k = 1/k^\alpha$, where $\alpha \in (0, 1)$, ensuring the almost sure convergence of the estimate. It is inappropriate to start with small values for step size γ_k and large values for the number of simulations M_k . Rather, it is recommended that one decrease γ_k and keep a constant and small number of MC samples M_k which shows a great advantage over the MC-step (5), which requires large M_k to converge.

This Robbins-Monro type of update represents the *first level* of our algorithm, needed to temper the variance and noise implied by MC integration. In the next section, we derive variants of this algorithm to adapt to the sheer size of data of today's applications and formalize the *second level* of our class of two-timescale EM methods.

2.2 Incremental and Bi-Level Stochastic EM Methods

Strategies to scale to large datasets include classical incremental and variance reduced variants. We will explicit a general update that will cover those variants and that represents the *second level* of our algorithm, namely the incremental update of the approximated statistics $\tilde{S}^{(k)}$ inside the RM type of update.

$$\text{Incremental-step : } \tilde{S}^{(k+1)} = \tilde{S}^{(k)} + \rho_{k+1} (\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) . \quad (8)$$

Note $\{\rho_k\}_{k \geq 1} \in (0, 1)$ is a sequence of step sizes, $\mathcal{S}^{(k)}$ is a proxy for $\tilde{S}^{(k)}$, If the stepsize is equal to one and the proxy $\mathcal{S}^{(k)} = \tilde{S}^{(k)}$, i.e., computed in a full batch manner as in (6), then we recover the SAEM algorithm. Also if $\rho_k = 1$, $\gamma_k = 1$ and $\mathcal{S}^{(k)} = \tilde{S}^{(k)}$, then we recover the MCEM [24].

We now introduce three variants of the SAEM update depending on different definitions of the proxy $\mathcal{S}^{(k)}$ and the choice of the stepsize ρ_k . Let $i_k \in \llbracket 1, n \rrbracket$ be a random index drawn at iteration k and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$ be the iteration index where $i \in \llbracket 1, n \rrbracket$ is last drawn prior to iteration k . For iteration $k \geq 0$, the fitTEM method draws *two* indices *independently* and uniformly as $i_k, j_k \in \llbracket 1, n \rrbracket$. In addition to τ_i^k which was defined w.r.t. i_k , we define $t_j^k = \{k' : j_{k'} = j, k' < k\}$ to be the iteration index where the sample $j \in \llbracket 1, n \rrbracket$ is last drawn as j_k prior to iteration k . With the initialization $\overline{\mathcal{S}}^{(0)} = \overline{s}^{(0)}$, we use a slightly different update rule from SAGA inspired by [22].

116 Then, we obtain:

$$(iSAEM [14]) \quad \mathcal{S}^{(k+1)} = \mathcal{S}^{(k)} + \frac{1}{n} (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)}) \quad (9)$$

$$(vrTTEM) \quad \mathcal{S}^{(k+1)} = \tilde{S}^{(\ell(k))} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}) \quad (10)$$

$$(fitTEM) \quad \mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}), \quad \bar{\mathcal{S}}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + n^{-1} (\tilde{S}_{j_k}^{(k)} - \tilde{S}_{j_k}^{(t_{j_k}^k)}) \quad (11)$$

117 where $\tilde{S}_{i_k}^{(k)}$ is the MC approximation of the expectation $\bar{S}_{i_k}(\theta^{(k)})$:

$$\tilde{S}_{i_k}^{(k)} = \frac{1}{M_k} \sum_{m=1}^{M_k} S(z_{i_k, m}^{(k)}, y_{i_k}) \quad \text{with} \quad z_{i_k, m}^{(k)} \sim p(z_{i_k} | y_{i_k}; \theta^{(k)}) .$$

118 The stepsize is set to $\rho_{k+1} = 1$ for the iSAEM method and we initialize with $\mathcal{S}^{(0)} = \tilde{S}^{(0)}$; $\rho_{k+1} = \gamma$
 119 is constant for the vrTTEM and fitTEM methods. Moreover, for vrTTEM we set an epoch size of
 120 m and define $\ell(k) := m \lfloor k/m \rfloor$ as the first iteration number in the epoch that iteration k is in.

121 **Two-Timescale Stochastic EM methods:** We now introduce the general method derived using the
 122 two variance reduction techniques described above. Algorithm 1 leverages both levels (7) and (8) in
 123 order to output a vector of fitted parameters $\hat{\theta}^{(K)}$ where K is a randomly chosen termination point.

Algorithm 1 Two-Timescale Stochastic EM methods.

- 1: **Input:** initializations $\hat{\theta}^{(0)} \leftarrow 0, \hat{s}^{(0)} \leftarrow \hat{S}^{(0)}, K_{\max} \leftarrow \text{max. iteration number}$.
- 2: Set the terminating iteration number, $K \in \{0, \dots, K_{\max} - 1\}$, as a discrete r.v. with:

$$P(K = k) = \frac{\gamma_k}{\sum_{\ell=0}^{K_{\max}-1} \gamma_\ell} = \frac{\gamma_k}{P_{\max}} . \quad (12)$$

- 3: **for** $k = 0, 1, 2, \dots, K$ **do**
- 4: Draw index $i_k \in \llbracket 1, n \rrbracket$ uniformly (and $j_k \in \llbracket 1, n \rrbracket$ for fitTEM).
- 5: Compute $\hat{S}_{i_k}^{(k)}$ using the MC-step (5), for the drawn indices.
- 6: Compute the surrogate sufficient statistics $\mathcal{S}^{(k+1)}$ using (9) or (10) or (11).
- 7: Compute $\hat{S}^{(k+1)}$ and $\hat{s}^{(k+1)}$ using respectively (8) and (7):

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho_{k+1} (\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{s}^{(k+1)} &= \hat{s}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{s}^{(k)}) \end{aligned} \quad (13)$$

- 8: Compute $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$ via the M-step (4).
 - 9: **end for**
 - 10: **Return:** $\hat{\theta}^{(K)}$.
-

124 The update in (13) is said to have two-timescale as the step sizes satisfy $\lim_{k \rightarrow \infty} \gamma_k / \rho_k < 1$ such that
 125 $\tilde{S}^{(k+1)}$ is updated at a faster time-scale, determined by ρ_k , than $\hat{s}^{(k+1)}$, determined by γ_k . The next
 126 section presents the main results of this paper and establishes global and finite-time bounds for the
 127 three different updates of our two-timescale scheme.

128 3 Finite Time Analysis of the Two-Timescale Scheme

129 Following [5], it can be shown that stationary points of the objective function (1) corresponds to the
 130 stationary points of the following *nonconvex* Lyapunov function:

$$\min_{s \in S} V(s) := \bar{L}(\bar{\theta}(s)) = r(\bar{\theta}(s)) + \frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(\bar{\theta}(s)) , \quad (14)$$

131 that we propose to study in this article. Several critical assumptions required to derive convergence
 132 guarantees read as follows:

133 **H1.** The sets Z, S are compact. There exists constants C_S, C_Z such that:

$$C_S := \max_{s, s' \in S} \|s - s'\| < \infty, \quad C_Z := \max_{i \in \llbracket 1, n \rrbracket} \int_Z |S(z, y_i)| \mu(dz) < \infty. \quad (15)$$

134 **H2.** The conditional distribution is smooth on $\text{int}(\Theta)$. For any $i \in \llbracket 1, n \rrbracket$, $z \in \mathcal{Z}$, $\theta, \theta' \in \text{int}(\Theta)^2$,
 135 we have $|p(z|y_i; \theta) - p(z|y_i; \theta')| \leq L_p \|\theta - \theta'\|$.

136 We also recall from the introduction that we consider curved exponential family models. besides:

137 **H3.** For any $s \in \mathcal{S}$, the function $\theta \mapsto L(s, \theta) := r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$ admits a unique global
 138 minimum $\bar{\theta}(s) \in \text{int}(\Theta)$. In addition, $J_\phi^\theta(\bar{\theta}(s))$ is full rank, L_ϕ -Lipschitz and $\bar{\theta}(s)$ is L_θ -Lipschitz.

139 We denote by $H_L^\theta(s, \theta)$ the Hessian (w.r.t to θ for a given value of s) of the function $\theta \mapsto L(s, \theta) =$
 140 $r(\theta) + \psi(\theta) - \langle s | \phi(\theta) \rangle$, and define

$$B(s) := J_\phi^\theta(\bar{\theta}(s)) \left(H_L^\theta(s, \bar{\theta}(s)) \right)^{-1} J_\phi^\theta(\bar{\theta}(s))^\top. \quad (16)$$

141 **H4.** It holds that $v_{\max} := \sup_{s \in \mathcal{S}} \|B(s)\| < \infty$ and $0 < v_{\min} := \inf_{s \in \mathcal{S}} \lambda_{\min}(B(s))$. There exists
 142 a constant L_B such that for all $s, s' \in \mathcal{S}$, we have $\|B(s) - B(s')\| \leq L_B \|s - s'\|$.

143 The class of algorithms we develop in this paper are two-timescale where the first stage corresponds
 144 to the variance reduction trick used in [12] in order to accelerate incremental methods and reduce the
 145 variance induced by the index sampling. The second stage is the Robbins-Monro type of update that
 146 aims to reduce the variance induced by the MC approximations As the expectations (3) are never
 147 available, we introduce the errors when approximating the quantity $\bar{s}_i(\hat{\theta}(\hat{s}^{(k-1)}))$ at iteration $k + 1$:

$$\eta_i^{(r)} := \tilde{S}_i^{(r)} - \bar{s}_i(\vartheta^{(r)}) \quad \text{for all } i \in \llbracket 1, n \rrbracket, r > 0 \quad \text{and} \quad \vartheta \in \Theta. \quad (17)$$

149 For instance, we consider that the MC approximation is unbiased if for all $i \in \llbracket 1, n \rrbracket$ and $m \in$
 150 $\llbracket 1, M \rrbracket$, the samples $z_{i,m} \sim p(z_i|y_i; \theta)$ are i.i.d. under the posterior distribution, i.e., $\mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] = 0$
 151 where \mathcal{F}_r is the filtration up to iteration r . The following results are derived under the assumption
 152 of control of the fluctuations implied by the approximation stated as follows:

153 **H5.** There exist a positive sequence of MC batch size $\{M_r\}_{r>0}$ and constants (C, C_η) such that for
 154 all $k > 0$, $i \in \llbracket 1, n \rrbracket$ and $\vartheta \in \Theta$:

$$\mathbb{E} \left[\left\| \eta_i^{(r)} \right\|^2 \right] \leq \frac{C_\eta}{M_r} \quad \text{and} \quad \mathbb{E} \left[\left\| \mathbb{E}[\eta_i^{(r)} | \mathcal{F}_r] \right\|^2 \right] \leq \frac{C}{M_r}. \quad (18)$$

155 We can prove two important results on the Lyapunov function. The first one suggests smoothness:

156 **Lemma 1.** [12] Assume H1-H4. For all $s, s' \in \mathcal{S}$ and $i \in \llbracket 1, n \rrbracket$, we have

$$\|\bar{s}_i(\bar{\theta}(s)) - \bar{s}_i(\bar{\theta}(s'))\| \leq L_s \|s - s'\|, \quad \|\nabla V(s) - \nabla V(s')\| \leq L_V \|s - s'\|, \quad (19)$$

157 where $L_s := C_Z L_p L_\theta$ and $L_V := v_{\max}(1 + L_s) + L_B C_S$.

158 and the second one suggests a growth condition on the gradient of V depending on the mean field
 159 of the algorithm:

160 **Lemma 2.** Assume H3, H4. For all $s \in \mathcal{S}$,

$$v_{\min}^{-1} \langle \nabla V(s) | s - \bar{s}(\bar{\theta}(s)) \rangle \geq \|s - \bar{s}(\bar{\theta}(s))\|^2 \geq v_{\max}^{-2} \|\nabla V(s)\|^2, \quad (20)$$

161 Proof of this Lemma can be found in Appendix A.

162 3.1 Global Convergence of Incremental Stochastic EM Algorithms

163 We present in this section a finite-time analysis of the incremental variant of the Stochastic Approx-
 164 imation of the EM algorithm. We want to draw the attention of the readers that the word "global"
 165 here does not mean for a global optimum of the nonconvex function, but of the independence of our
 166 analysis on the initialization and the iteration k (finite time).

167 The following main result for the iSAEM algorithm, which proof can be found in Appendix B, is
 168 derived under a control of the Monte Carlo fluctuations as described by assumption H 5 and is built
 169 upon an intermediary Lemma, detailed in in Appendix B, characterizing the quantity of interest
 170 $\hat{S}^{(k+1)} - \hat{s}^{(k)}$. Typically, the controls exhibited above are of interest when the number of MC
 171 samples M_k increase with k .

Theorem 1. Assume *H1-H5*. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and consider the iSAEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = 1$ for any $k > 0$. We also set $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\gamma_{k+1} = \frac{1}{k^\alpha \alpha c_1 \bar{L}}$ where $a \in (0, 1)$, $\beta = \frac{c_1 \bar{L}}{n}$. Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$.

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right].$$

3.2 Global Convergence of Two-Timescale Stochastic EM Algorithms

We now proceed by giving our main result regarding the global convergence of the fitTEM algorithm. Two important auxiliary Lemmas, which proofs are given in Appendix C.1, are need in order to derive our finite-time bound. The first one derives an identity for the quantity $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2]$ using the vrTTEM update:

Lemma 3. For any $k \geq 0$ and consider the vrTTEM update in (10) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{\mathbf{S}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2], \end{aligned}$$

where we recall that $\ell(k)$ is the first iteration number in the epoch that iteration k is in.

The second one derives an identity for the quantity $\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2]$ using the fitTEM update:

Lemma 4. For any $k \geq 0$ and consider the fitTEM update in (11) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + 2(1 - \rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{\mathbf{S}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]. \end{aligned}$$

Recalling that K is an independent discrete r.v. drawn from $\{1, \dots, K_{\max}\}$ with distribution $\{\gamma_k / P_{\max}, 0 \leq k \leq K_{\max} - 1\}$, as in (12), we have

$$\mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] = \frac{1}{P_{\max}} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2].$$

We now state the main result regarding the vrTTEM method, see proof in Appendix D:

Theorem 2. Assume *H1-H5*. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and consider the vrTTEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$. Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. Setting $\bar{L} = \max\{L_s, L_V\}$, $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$, a constant $\mu \in (0, 1)$, $\gamma_{k+1} = \frac{1}{k^\alpha \bar{L}}$ where $a \in (0, 1)$, we have the following bound:

$$\begin{aligned} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] &\leq \frac{2n^{2/3} \bar{L}}{\mu P_{\max} v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3} \bar{L}}{\mu P_{\max} v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)} \right\|^2 \right] \right]. \end{aligned}$$

We now state the main result regarding the fitTEM method.

Theorem 3. Assume *H1-H5*. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of positive step sizes and consider the fitTEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any $k > 0$. Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\alpha = \max\{2, 1 + 2v_{\min}\}$,

199 $\bar{L} = \max\{L_S, L_V\}$, $\beta = \frac{1}{\alpha n}$, $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$ and $\gamma_{k+1} = \frac{1}{k^a \alpha c_1 \bar{L}}$
 200 where $a \in (0, 1)$, we have the following bound:

$$\begin{aligned} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(K)})\|^2] &\leq \frac{4\alpha \bar{L} n^{2/3}}{P_{\max} v_{\min}^2 v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{4\alpha \bar{L} n^{2/3}}{P_{\max} v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right]. \end{aligned}$$

201 Proof of this Theorem can be found in Appendix E. Note that in those two bounds, the quantities
 202 $\tilde{\eta}^{(k+1)}$ and $\Xi^{(k+1)}$ depend only on the MC fluctuations $\mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]$ and some constants.

203 While Theorem 1 suffers only from the MC noise induced by the latent data sampling step, Theo-
 204 rem 2 and Theorem 3 exhibit in their convergence bounds two different phases. The upper bounds
 205 display a bias term due to the initial conditions, *i.e.*, the term $V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})$, and a double
 206 dynamics burden exemplified by the term $\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right]$.

207 Indeed, the following remarks are worth noting on the quantity $\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right]$:

- 208 • This term is the price we pay for the two-timescale dynamics and corresponds to the gap
 209 between the two asynchronous updates (one is on $\hat{\mathbf{s}}^{(k)}$ and the other on $\tilde{S}^{(k)}$).
- It is trivial to see that if $\rho = 1$, *i.e.*, there is no variance reduction, then for any $k > 0$

$$\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] = \mathbb{E} \left[\left\| \mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)} \right\|^2 \right] = 0 \quad \text{with} \quad \hat{\mathbf{s}}^{(0)} = \tilde{S}^{(0)} = 0$$

210 which strengthen the fact that this quantity characterizes the impact of the variance reduc-
 211 tion technique introduced in our two stages class of methods.

212 The following lemma, which proof can be found in Appendix C.2, characterizes this gap:

213 **Lemma 5.** Consider a decreasing stepsize $\gamma_k \in (0, 1)$ and a constant $\rho \in (0, 1)$, then the following
 214 inequality holds:

$$\mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_{\ell})^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)}),$$

215 where $\mathcal{S}^{(k)}$ is defined either by (10) (vrTTEM) or (11) (fiTTEM).

216 In the next section, we illustrate the benefits of our two-timescale class of algorithms on several
 217 numerical applications.

218 4 Numerical Examples

219 For the sake of space, we provide details on the experiments in Appendix F.

220 4.1 Gaussian Mixture Models

221 We begin by a simple and illustrative example. The authors acknowledge that the following model
 222 can be trained using deterministic EM-type of algorithms but propose to apply stochastic methods,
 223 including theirs, and to compare their performances. Given n observations $\{y_i\}_{i=1}^n$, we want to
 224 fit a Gaussian Mixture Model (GMM) whose distribution is modeled as a Gaussian mixture of M
 225 components, each with a unit variance. Let $z_i \in \llbracket M \rrbracket$ be the latent labels of each component, the
 226 complete log-likelihood is defined as:

$$\log f(z_i, y_i; \boldsymbol{\theta}) = \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) [\log(\omega_m) - \mu_m^2/2] + \sum_{m=1}^M \mathbb{1}_{\{m\}}(z_i) \mu_m y_i + \text{constant}. \quad (21)$$

227 where $\boldsymbol{\theta} := (\boldsymbol{\omega}, \boldsymbol{\mu})$ with $\boldsymbol{\omega} = \{\omega_m\}_{m=1}^{M-1}$ are the mixing weights with the convention $\omega_M =$
 228 $1 - \sum_{m=1}^{M-1} \omega_m$ and $\boldsymbol{\mu} = \{\mu_m\}_{m=1}^M$ are the means. We use the penalization $r(\boldsymbol{\theta}) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 -$

229 $\log \text{Dir}(\omega; M, \epsilon)$ where $\delta > 0$ and $\text{Dir}(\cdot; M, \epsilon)$ is the M dimensional symmetric Dirichlet distribu-
 230 tion with concentration parameter $\epsilon > 0$. The constraint set on θ is given by

$$\Theta = \{\omega_m, m = 1, \dots, M-1 : \omega_m \geq 0, \sum_{m=1}^{M-1} \omega_m \leq 1\} \times \{\mu_m \in \mathbb{R}, m = 1, \dots, M\}.$$

231 In the following experiments on synthetic data, we generate 30 synthetic datasets of size $n = 10^5$
 232 from a GMM model with $M = 2$ components with two mixtures with means $\mu_1 = -\mu_2 = 0.5$.
 233 We run the bEM method until convergence (to
 234 double precision) to obtain the ML estimate μ^*
 235 averaged on 50 datasets. We compare the bEM,
 236 iEM (incremental EM), SAEM, iSAEM, vrT-
 237 TEM and fitTEM methods in terms of their
 238 precision measured by $|\mu - \mu^*|^2$. We set
 239 the stepsize of the SA-step of all method as
 240 $\gamma_k = 1/k^\alpha$ with $\alpha = 0.5$, and the stepsizes
 241 of the Incremental-step for vrTTEM and the
 242 fitTEM to a constant stepsize equal to $1/n^{2/3}$.
 243 The number of MC samples is fixed to $M = 10$
 244 chains. Figure 1 shows the convergence of the
 245 precision $|\mu - \mu^*|^2$ for the different methods
 246 against the epoch(s) elapsed (one epoch equals
 247 n iterations). Besides, vrTTEM and fitTEM methods outperform the other stochastic methods,
 248 supporting the benefits of our scheme.




Figure 1: Precision $|\mu^{(k)} - \mu^*|^2$ per epoch

249 4.2 Deformable Template Model for Image Analysis

250 Let $(y_i, i \in \llbracket 1, n \rrbracket)$ be observed gray level images defined on a grid of pixels. Let $u \in \mathcal{U} \subset \mathbb{R}^2$
 251 denotes the pixel index on the image and $x_u \in \mathcal{D} \subset \mathbb{R}^2$ its location. The model used in this
 252 experiment suggests that each image y_i is a deformation of a template, noted $I : \mathcal{D} \rightarrow \mathbb{R}$, common
 253 to all images of the dataset:

$$y_i(u) = I(x_u - \Phi_i(x_u, z_i)) + \varepsilon_i(u) \quad (22)$$

254 where $\phi_i : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a deformation function, z_i some latent variable parameterizing this deforma-
 255 tion and $\varepsilon_i \sim \mathcal{N}(0, \sigma^2)$ is an observation error.

256 The template model, given $\{p_k\}_{k=1}^{k_p}$ landmarks on the template, a fixed known kernel \mathbf{K}_p and a
 257 vector of parameters $\beta \in \mathbb{R}^{k_p}$ is defined as follows:

$$I_\xi = \mathbf{K}_p \beta, \quad \text{where} \quad (\mathbf{K}_p \beta)(x) = \sum_{k=1}^{k_p} \mathbf{K}_p(x, p_k) \beta_k.$$

258 Besides, we parameterize the deformation model given some landmarks $\{g_k\}_{k=1}^{k_g}$ and a fixed kernel
 259 \mathbf{K}_g as:

$$\Phi_i = \mathbf{K}_g z_i \quad \text{where} \quad (\mathbf{K}_g z_i)(x) = \sum_{k=1}^{k_g} \mathbf{K}_g(x, g_k) \begin{pmatrix} z_i^{(1)}(k) \\ z_i^{(2)}(k) \end{pmatrix},$$

260 where we put a Gaussian prior on the latent variables, $z_i \sim \mathcal{N}(0, \Gamma)$ and $z_i \in (\mathbb{R}^{k_g})^2$. The vector
 261 of parameters we estimate is thus $\theta = (\beta, \Gamma, \sigma)$.

262 **Numerical Experiment:** We apply model (22) and our algorithms to a collection of handwritten
 263 digits, called the US postal database [11], featuring $n = 1000$ (16×16)-pixel images for each
 264 class of digits from 0 to 9. The main difficulty with these data comes from the geometric dispersion
 265 within each class of digit as shown Figure 2 for digit 5. We thus ought to use our deformable
 266 template model (22) in order to account for both sources of variability: the intrinsic template to each
 267 class of digit and the small and local deformation in each observed image.



Figure 2: Training set of the USPS database (20 images for figit 5)

Figure 4 shows the resulting synthetic images for digit 5 through several epochs and for the batch method, the online SAEM, the incremental SAEM and the various TTS methods. We choose Gaussian kernels for both, \mathbf{K}_p and \mathbf{K}_g , defined on \mathbb{R}^2 and centered on the landmark points $\{p_k\}_{k=1}^{k_p}$ and $\{g_k\}_{k=1}^{k_g}$ with standard respective standard deviations of 0.12 and 0.3. $k_p = 15$ and $k_g = 6$ equidistributed landmarks points are chosen on the grid for the training. Those hyperparameters are inspired by a relevant study in [2]. The kernel covariance matrices are important hyperparameters in such study since they have a direct impact on the sharpness of the templates. Intuitively, if those variances are large, the kernels centered around the equidistributed landmark spread out on too many of its neighbors. Bad choices of such hyperparameters can lead to thicker shapes.



Figure 3: Estimation of the template: from top to bottom: batch, online, iSAEM ,vrTTEM and fitTEM . Columns represent 1 to 7 epochs.



Figure 4: Estimation of the template: from top to bottom: batch, online, iSAEM ,vrTTEM and fitTEM through 7 epochs. Note that Batch method templates are replicated in-between epochs for a fair comparison with incremental variants.

Figure 4 displays the virtue of the vrTTEM and fitTEM methods that obtain a more *contrasted* and *accurate* template estimate. The incremental and online version are looking much better on the very first epochs compared to the batch method, which is intuitive given the high computational cost of the batch method. After a few epochs, the batch SAEM seem to estimate similar template as the incremental an online methods due to the high variance of the latter procedures. Our variance

reduced and fast incremental come into play in the long run and sharpen the final template estimates contrasting between the background and the regions of interest in the image.

4.3 PK Model with Absorption Lag Time

This numerical example was conducted in order to characterize the pharmacokinetics (PK) of orally administered drug to simulated patients, using a population pharmacokinetic approach. $M = 50$ synthetic datasets were generated for $n = 5000$ patients with 10 observations (concentration measures) per patient. The goal is to model the evolution of the concentration of the absorbed drug using a nonlinear and latent data model.

The model: We consider a one-compartment PK model for oral administration with an absorption lag-time (T^{lag}), assuming first-order absorption and linear elimination processes. The final model includes the following variables: ka the absorption rate constant, V the volume of distribution, k the elimination rate constant and T^{lag} the absorption lag-time. We also add several covariates to our model such as D the dose of drug administered, t the time at which measures are taken and the weight of the patient influencing the volume V . More precisely, the log-volume $\log(V)$ is a linear function of the log-weight $lw70 = \log(wt/70)$. Let $z_i = (T_i^{\text{lag}}, ka_i, V_i, k_i)$ be the vector of individual PK parameters, different for each individual i . The final model reads:

$$y_{ij} = f(t_{ij}, z_i) + \varepsilon_{ij} \quad \text{where} \quad f(t_{ij}, z_i) = \frac{D ka_i}{V(ka_i - k_i)} (e^{-ka_i(t_{ij} - T_i^{\text{lag}})} - e^{-k_i(t_{ij} - T_i^{\text{lag}})}), \quad (23)$$

where y_{ij} is the j -th concentration measurement of the drug of dosage D injected at time t_{ij} for patient i . We assume in this example that the residual errors ε_{ij} are independent and normally distributed with mean 0 and variance σ^2 . Lognormal distributions are used for the four PK parameters.

Monte Carlo study: We conduct a Monte Carlo study to showcase the benefits of our scheme. $M = 50$ datasets have been simulated using the following PK parameters values: $T_{\text{pop}}^{\text{lag}} = 1$, $ka_{\text{pop}} = 1$, $V_{\text{pop}} = 8$, $k_{\text{pop}} = 0.1$, $\omega_{T^{\text{lag}}} = 0.4$, $\omega_{ka} = 0.5$, $\omega_V = 0.2$, $\omega_k = 0.3$ and $\sigma^2 = 0.5$. We define the mean square distance over the M replicates $E_k(\ell) = \frac{1}{M} \sum_{m=1}^M (\theta_k^{(m)}(\ell) - \theta^*)^2$ and plot it against the epochs (passes over the data) Figure 5. Note that the MC-step (5) is performed using a Metropolis Hastings procedure since the posterior distribution under the model θ noted $p(z_i|y_i, \theta)$ is intractable due to the nonlinearity of the model (23). Figure 5 shows clear advantage of variance reduced methods (vrTTEM and fiTTEM) avoiding the twists and turns displayed by the incremental and the batch methods.




Figure 5: Precision $|ka^{(k)} - ka^*|^2$ per epoch

5 Conclusion

This paper introduces a new class of two-timescale EM methods for learning latent data models. In particular, the models dealt with in this paper belong to the curved exponential family and are possibly nonconvex. The nonconvexity of the problem is tackled using a Robbins-Monro type of update, which represent the *first* level of our class of methods and the scalability with the number of samples is performed through a variance reduced and incremental type of update, the *second* and last level of our newly introduced scheme. The various methods are interpreted as scaled gradient methods, in the space of the sufficient statistics, and our convergence results are *global*, in the sense of independent of the initial values, and *non-asymptotic*, true for any random termination number.

References

- [1] S. Allasonnière, Y. Amit, and A. Trouvé. Towards a coherent statistical framework for dense deformable template estimation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(1):3–29, 2007.
- [2] S. Allasonnière, E. Kuhn, A. Trouvé, et al. Construction of bayesian deformable models via a stochastic approximation algorithm: a convergence study. *Bernoulli*, 16(3):641–678, 2010.
- [3] C. Baey, S. Trevezas, and P.-H. Cournède. A non linear mixed effects model of plant growth and estimation via stochastic variants of the em algorithm. *Communications in Statistics-Theory and Methods*, 45(6):1643–1669, 2016.
- [4] O. Cappé. Online em algorithm for hidden markov models. *Journal of Computational and Graphical Statistics*, 20(3):728–749, 2011.
- [5] O. Cappé and E. Moulines. On-line expectation–maximization algorithm for latent data models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):593–613, 2009.
- [6] B. P. Carlin and S. Chib. Bayesian model choice via markov chain monte carlo methods. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(3):473–484, 1995.
- [7] A. Chakraborty and K. Das. Inferences for joint modelling of repeated ordinal scores and time to event data. *Computational and mathematical methods in medicine*, 11(3):281–295, 2010.
- [8] B. Delyon, M. Lavielle, and E. Moulines. Convergence of a stochastic approximation version of the em algorithm. *Ann. Statist.*, 27(1):94–128, 03 1999. doi: 10.1214/aos/1018031103. URL <https://doi.org/10.1214/aos/1018031103>.
- [9] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *Journal of the royal statistical society. Series B (methodological)*, pages 1–38, 1977.
- [10] J. P. Hughes. Mixed effects models with censored data with application to hiv rna levels. *Biometrics*, 55(2):625–629, 1999.
- [11] J. J. Hull. A database for handwritten text recognition research. *IEEE Transactions on pattern analysis and machine intelligence*, 16(5):550–554, 1994.
- [12] B. Karimi, H.-T. Wai, É. Moulines, and M. Lavielle. On the global convergence of (fast) incremental expectation maximization methods. In *Advances in Neural Information Processing Systems*, pages 2833–2843, 2019.
- [13] E. Kuhn and M. Lavielle. Coupling a stochastic approximation version of em with an mcmc procedure. *ESAIM: Probability and Statistics*, 8:115–131, 2004.
- [14] E. Kuhn, C. Matias, and T. Rebafka. Properties of the stochastic approximation em algorithm with mini-batch sampling. *arXiv preprint arXiv:1907.09164*, 2019.
- [15] P. Liang and D. Klein. Online em for unsupervised models. In *Proceedings of human language technologies: The 2009 annual conference of the North American chapter of the association for computational linguistics*, pages 611–619, 2009.
- [16] F. Maire, E. Moulines, and S. Lefebvre. Online em for functional data, 2016. URL <http://arxiv.org/abs/1604.00570>. cite arxiv:1604.00570v1.pdf.
- [17] C. E. McCulloch. Maximum likelihood algorithms for generalized linear mixed models. *Journal of the American statistical Association*, 92(437):162–170, 1997.

- 368 [18] G. McLachlan and T. Krishnan. *The EM algorithm and extensions*, volume 382. John Wiley
369 & Sons, 2007.
- 370 [19] S. P. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Springer Science &
371 Business Media, 2012.
- 372 [20] R. M. Neal and G. E. Hinton. A view of the EM algorithm that justifies incremental, sparse,
373 and other variants. In *Learning in graphical models*, pages 355–368. Springer, 1998.
- 374 [21] H. D. Nguyen, F. Forbes, and G. J. McLachlan. Mini-batch learning of exponential family
375 finite mixture models. *Statistics and Computing*, pages 1–18, 2020.
- 376 [22] S. J. Reddi, S. Sra, B. Póczos, and A. Smola. Fast incremental method for nonconvex opti-
377 mization. *arXiv preprint arXiv:1603.06159*, 2016.
- 378 [23] H. Robbins and S. Monro. A stochastic approximation method. *The annals of mathematical*
379 *statistics*, pages 400–407, 1951.
- 380 [24] G. C. Wei and M. A. Tanner. A monte carlo implementation of the em algorithm and the poor
381 man’s data augmentation algorithms. *Journal of the American statistical Association*, 85(411):
382 699–704, 1990.
- 383 [25] R. Zhu, L. Wang, C. Zhai, and Q. Gu. High-dimensional variance-reduced stochastic gradient
384 expectation-maximization algorithm. In *Proceedings of the 34th International Conference on*
385 *Machine Learning-Volume 70*, pages 4180–4188. JMLR. org, 2017.

386 A Proof of Lemma 2

387 **Lemma.** Assume H3, H4. For all $\mathbf{s} \in S$,

$$v_{\min}^{-1} \langle \nabla V(\mathbf{s}) | \mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \rangle \geq \|\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))\|^2 \geq v_{\max}^{-2} \|\nabla V(\mathbf{s})\|^2, \quad (24)$$

388 **Proof** Using H3 and the fact that we can exchange integration with differentiation and the Fisher's
389 identity, we obtain

$$\begin{aligned} \nabla_{\mathbf{s}} V(\mathbf{s}) &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{L}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \left(\nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \right) \\ &= \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s})^{\top} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} (\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))), \end{aligned} \quad (25)$$

390 Consider the following vector map:

$$\mathbf{s} \rightarrow \nabla_{\boldsymbol{\theta}} L(\mathbf{s}, \boldsymbol{\theta})|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})} = \nabla_{\boldsymbol{\theta}} \psi(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \nabla_{\boldsymbol{\theta}} \mathbf{r}(\bar{\boldsymbol{\theta}}(\mathbf{s})) - \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top} \mathbf{s}.$$

391 Taking the gradient of the above map w.r.t. \mathbf{s} and using assumption H3, we show that:

$$\mathbf{0} = -\mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) + \underbrace{\left(\nabla_{\boldsymbol{\theta}}^2 (\psi(\boldsymbol{\theta}) + \mathbf{r}(\boldsymbol{\theta}) - \langle \phi(\boldsymbol{\theta}) | \mathbf{s} \rangle) \right)|_{\boldsymbol{\theta}=\bar{\boldsymbol{\theta}}(\mathbf{s})}}_{=\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \boldsymbol{\theta})} \mathbf{J}_{\bar{\boldsymbol{\theta}}}^{\mathbf{s}}(\mathbf{s}).$$

392 The above yields

$$\nabla_{\mathbf{s}} V(\mathbf{s}) = \mathbf{B}(\mathbf{s})(\mathbf{s} - \bar{\mathbf{s}}(\bar{\boldsymbol{\theta}}(\mathbf{s})))$$

393 where we recall $\mathbf{B}(\mathbf{s}) = \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s})) \left(\mathbf{H}_L^{\boldsymbol{\theta}}(\mathbf{s}; \bar{\boldsymbol{\theta}}(\mathbf{s})) \right)^{-1} \mathbf{J}_{\phi}^{\boldsymbol{\theta}}(\bar{\boldsymbol{\theta}}(\mathbf{s}))^{\top}$. The proof of (24) follows directly
394 from the assumption H4. \square

395 B Proof of Theorem 1

396 Beforehand, We present two intermediary Lemmas important for the analysis of the incremen-
397 tal update of the iSAEM algorithm. The first one gives a characterization of the quantity
398 $\mathbb{E} [\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}]$:

399 **Lemma 6.** Assume H1. The update (9) is equivalent to the following update on the resulting statis-
400 tics

$$\hat{\mathbf{s}}^{(k+1)} = \hat{\mathbf{s}}^{(k)} + \gamma_{k+1} (\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)})$$

401 Also:

$$\mathbb{E} [\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}] = \mathbb{E} [\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] + \frac{1}{n} \mathbb{E} [\eta_{i_k}^{(k+1)}]$$

402 where $\bar{\mathbf{s}}^{(k)}$ is defined by (3) and $\tau_i^k = \max\{k' : i_{k'} = i, k' < k\}$.

403 **Proof** From update (9), we have:

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} + \frac{1}{n} \left(\tilde{S}_{i_k}^{(k+1)} - \tilde{S}_{i_k}^{(\tau_{i_k}^k)} \right) \\ &= \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \tilde{S}_{i_k}^{(k+1)} \right) \end{aligned}$$

404 Since $\tilde{S}_{i_k}^{(k+1)} = \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) + \eta_{i_k}^{(k+1)}$ we have

$$\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} + \tilde{S}^{(k)} - \bar{\mathbf{s}}^{(k)} - \frac{1}{n} \left(\tilde{S}_{i_k}^{(\tau_{i_k}^k)} - \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) \right) + \frac{1}{n} \eta_{i_k}^{(k+1)}$$

405 Taking the full expectation of both side of the equation leads to:

$$\begin{aligned}\mathbb{E} \left[\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \right] &= \mathbb{E} \left[\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right] + \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right] \\ &\quad - \frac{1}{n} \mathbb{E} \left[\mathbb{E} \left[\tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k \right] \right] + \frac{1}{n} \mathbb{E} \left[\eta_{i_k}^{(k+1)} \right]\end{aligned}$$

406 The following equalities:

$$\mathbb{E} \left[\tilde{S}_i^{(\tau_i^k)} | \mathcal{F}_k \right] = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} \quad \text{and} \quad \mathbb{E} \left[\bar{\mathbf{s}}_{i_k}(\boldsymbol{\theta}^{(k)}) | \mathcal{F}_k \right] = \bar{\mathbf{s}}^{(k)}$$

407 concludes the proof of the Lemma. \square

408 And the following auxiliary Lemma setting an upper bound for the quantity $\mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right]$

409 **Lemma 7.** For any $k \geq 0$ and consider the iSAEM update in (9), it holds that

$$\begin{aligned}\mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] &\leq 4\mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] + \frac{2\mathbf{L}_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 \right] \\ &\quad + 2\frac{C_\eta}{M_k} + 4\mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right]\end{aligned}$$

410 **Proof** Applying the iSAEM update yields:

$$\begin{aligned}\mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] &= \mathbb{E} \left[\left\| \tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} - \frac{1}{n} (\tilde{S}_i^{(\tau_i^k)} - \tilde{S}_{i_k}^{(k)}) \right\|^2 \right] \\ &\leq 4\mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + 4\mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] \\ &\quad + \frac{2}{n^2} \mathbb{E} \left[\left\| \bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_i^k)} \right\|^2 \right] + 2\frac{C_\eta}{M_k}\end{aligned}$$

411 The last expectation can be further bounded by

$$\frac{2}{n^2} \mathbb{E} \left[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_i^k)}\|^2 \right] = \frac{2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2 \right] \stackrel{(a)}{\leq} \frac{2\mathbf{L}_s^2}{n^3} \sum_{i=1}^n \mathbb{E} \left[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 \right],$$

412 where (a) is due to Lemma 1 and which concludes the proof of the Lemma. \square

413

414 **Theorem.** Assume H1-H5. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of
415 positive step sizes and consider the iSAEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = 1$ for any
416 $k > 0$. We also set $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$, $\gamma_{k+1} = \frac{1}{k^\alpha \alpha c_1 \bar{L}}$ where
417 $a \in (0, 1)$, $\beta = \frac{c_1 \bar{L}}{n}$. Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$.

$$v_{\max}^{-2} \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]$$

418 **Proof** Under the smoothness of the Lyapunov function V (cf. Lemma 1), we can write:

$$V(\hat{\mathbf{s}}^{(k+1)}) \leq V(\hat{\mathbf{s}}^{(k)}) + \gamma_{k+1} \langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 \mathbf{L}_V}{2} \|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2$$

419 Taking the expectation on both sides yields:

$$\mathbb{E} \left[V(\hat{\mathbf{s}}^{(k+1)}) \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) \right] + \gamma_{k+1} \mathbb{E} \left[\langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right]$$

420 Using Lemma 6, we obtain:

$$\begin{aligned} & \mathbb{E} \left[\langle \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] = \\ & \mathbb{E} \left[\langle \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ & \stackrel{(a)}{\leq} -v_{\min} \mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] + \left(1 - \frac{1}{n} \right) \mathbb{E} \left[\left\langle \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \right\rangle \right] + \frac{1}{n} \mathbb{E} \left[\langle \eta_{i_k}^{(k)} \mid \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ & \stackrel{(b)}{\leq} -v_{\min} \mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ & + \frac{\beta(n-1)+1}{2n} \mathbb{E} \left[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\|\eta_{i_k}^{(k)}\|^2 \right] \\ & \stackrel{(a)}{\leq} \left(v_{\max}^2 \frac{\beta(n-1)+1}{2n} - v_{\min} \right) \mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] + \frac{1 - \frac{1}{n}}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + \frac{1}{2n} \mathbb{E} \left[\|\eta_{i_k}^{(k)}\|^2 \right] \end{aligned}$$

421 where (a) is due to the growth condition (2) and (b) is due to Young's inequality (with $\beta \rightarrow 1$). Note

422 $a_k = \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1)+1}{2n} \right)$ and

$$\begin{aligned} a_k \mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] & \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)}) \right] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] \\ & + \frac{\gamma_{k+1}(1 - \frac{1}{n})}{2\beta} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] + \frac{\gamma_{k+1}}{2n} \mathbb{E} \left[\|\eta_{i_k}^{(k)}\|^2 \right] \end{aligned} \quad (26)$$

423 We now give an upper bound of $\mathbb{E} \left[\|\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \right]$ using Lemma 7 and plug it into (26):

$$\begin{aligned} (a_k - 2\gamma_{k+1}^2 L_V) \mathbb{E} \left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2 \right] & \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)}) \right] \\ & + \gamma_{k+1} \left(\frac{1}{2\beta} \left(1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\ & + \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[\|\eta_{i_k}^{(k)}\|^2 \right] \\ & + \frac{\gamma_{k+1}^2 L_V L_{\mathbf{s}}^2}{n^3} \sum_{i=1}^n \mathbb{E} [\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] \end{aligned} \quad (27)$$

424 Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E} [\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E} [\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E} [\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2] \right)$$

where the equality holds as i_k and j_k are drawn independently. For any $\beta > 0$, it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\ &= \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 - 2\gamma_{k+1}\langle \hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)} \mid \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)} \rangle\right] \\ &\leq \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2 + \gamma_{k+1}\beta\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \end{aligned}$$

where the last inequality is due to the Young's inequality. Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}\left[\left(1 + \gamma_{k+1}\beta\right)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2 + \frac{\gamma_{k+1}}{\beta}\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)}\|^2\right] \end{aligned}$$

Observe that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k+1)})$. Applying Lemma 7 yields

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\tau_i^{k+1})}\|^2] \\ &\leq (\gamma_{k+1}^2 + \frac{n-1}{n} \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\tilde{\mathbf{S}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2\right] \\ &\leq 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] \\ &\quad + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \\ &\quad + \sum_{i=1}^n \mathbb{E}\left[\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})}{n} \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2\right] \end{aligned}$$

Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\tau_i^k)}\|^2]$$

From the above, we get

$$\begin{aligned} \Delta^{(k+1)} &\leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})\right) \Delta^{(k)} + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] \\ &\quad + 2(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_k}^{(k)}\|^2\right] + 4(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)}\right\|^2\right] \end{aligned}$$

Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{8, 1 + 6v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}$, $\beta = \frac{c_1 \bar{L}}{n}$,

$c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 6$, $\alpha \geq 8$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \leq 1 - \frac{c_1(k\alpha - 1) - 4}{k\alpha n c_1} \leq 1 - \frac{2}{k\alpha n c_1}$$

which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta}) \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} =$

$\frac{1}{n} - \gamma_{k+1}\beta - \frac{2\gamma_{k+1}L_s^2}{n^2}(\gamma_{k+1} + \frac{1}{\beta})$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} &\leq 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2] + 2 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\|\eta_{i_\ell}^{(\ell)}\|^2\right] \\ &\quad + 4 \sum_{\ell=0}^k \prod_{j=\ell+1}^k \left(1 - \Lambda_{(j)}\right) (\gamma_{\ell+1}^2 + \frac{\gamma_{\ell+1}}{\beta}) \mathbb{E}\left[\left\|\frac{1}{n} \sum_{i=1}^n \tilde{\mathbf{S}}_i^{(\tau_i^\ell)} - \bar{\mathbf{s}}^{(\ell)}\right\|^2\right] \end{aligned}$$

434 Note $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ Summing on both sides over $k = 0$ to $k = K_{\max} - 1$ yields:

$$\begin{aligned}
& \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} \\
&= 4 \sum_{k=0}^{K_{\max}-1} \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + 2 \sum_{k=0}^{K_{\max}-1} \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[\left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_{\max}-1} 4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right) \omega_{k,1} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&\leq \sum_{k=0}^{K_{\max}-1} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{k=0}^{K_{\max}-1} \frac{2 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[\left\| \eta_{i_\ell}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_{\max}-1} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right]
\end{aligned} \tag{28}$$

435 We recall (27) where we have summed on both sides from $k = 0$ to $k = K_{\max} - 1$:

$$\begin{aligned}
& \sum_{k=0}^{K_{\max}-1} \left(a_k - 2\gamma_{k+1}^2 L_V \right) \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\
&+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\frac{1}{2\beta} \left(1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right] \\
&+ \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \Delta^{(k)}
\end{aligned} \tag{29}$$

436 Plugging (28) into (29) results in:

$$\begin{aligned}
& \sum_{k=0}^{K_{\max}-1} \tilde{\alpha}_k \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right] + \sum_{k=0}^{K_{\max}-1} \tilde{\beta}_k \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(\tau_i^k)} - \bar{\mathbf{s}}^{(k)} \right\|^2 \right] \leq \mathbb{E} \left[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K)}) \right] \\
&+ \sum_{k=0}^{K_{\max}-1} \tilde{\Gamma}_k \mathbb{E} \left[\left\| \eta_{i_k}^{(k)} \right\|^2 \right]
\end{aligned}$$

437 where:

$$\begin{aligned}
\tilde{\alpha}_k &= a_k - 2\gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\beta}_k &= \gamma_{k+1} \left(\frac{1}{2\beta} \left(1 - \frac{1}{n} \right) + 2\gamma_{k+1} L_V \right) - \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{4 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}} \\
\tilde{\Gamma}_k &= \gamma_{k+1} \left(\gamma_{k+1} L_V + \frac{1}{2n} \right) + \frac{\gamma_{k+1}^2 L_V L_s^2}{n^2} \frac{2 \left(\gamma_{k+1}^2 + \frac{\gamma_{k+1}}{\beta} \right)}{\Lambda_{(k+1)}}
\end{aligned}$$

438 and

$$\begin{aligned}
a_k &= \gamma_{k+1} \left(v_{\min} - v_{\max}^2 \frac{\beta(n-1) + 1}{2n} \right) \\
\Lambda_{(k+1)} &= \frac{1}{n} - \gamma_{k+1} \beta - \frac{2\gamma_{k+1} \mathbf{L}_{\mathbf{s}}^2}{n^2} \left(\gamma_{k+1} + \frac{1}{\beta} \right) \\
c_1 &= v_{\min}^{-1}, \alpha = \max\{8, 1 + 6v_{\min}\}, \bar{L} = \max\{\mathbf{L}_{\mathbf{s}}, \mathbf{L}_V\}, \gamma_{k+1} = \frac{1}{k\alpha c_1 \bar{L}}, \beta = \frac{c_1 \bar{L}}{n}
\end{aligned}$$

439 When, for any $k > 0$, $\tilde{\alpha}_k \geq 0$, we have by Lemma 2 that:

$$\sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \nabla V(\hat{\mathbf{s}}^{(k)}) \right\|^2 \right] \leq v_{\max}^2 \sum_{k=0}^{K_{\max}} \tilde{\alpha}_k \mathbb{E} \left[\left\| \bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)} \right\|^2 \right]$$

440 which yields an upper bound of the gradient of the Lyapunov function V along the path of the
441 iSAEM update and concludes the proof of the Theorem. \square

442 C Proofs of Auxiliary Lemmas

443 C.1 Proof of Lemma 3 and Lemma 4

444 **Lemma.** For any $k \geq 0$ and consider the vrTTEM update in (10) with $\rho_k = \rho$, it holds for all $k > 0$
 445

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

446 where we recall that $\ell(k)$ is the first iteration number in the epoch that iteration k is in.

447 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
 448 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \end{aligned} \quad (30)$$

449 We observe, using the identity (30), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] \quad (31)$$

450 For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \tilde{S}_i^{(\ell(k))}) - (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(\ell(k))}) \right\|^2 \right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \stackrel{(b)}{\leq} L_s^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

451 where (a) uses the variance inequality and (b) uses Lemma 1. Substituting into (31) proves the
 452 lemma. \square

453 **Lemma.** For any $k \geq 0$ and consider the fitTEM update in (11) with $\rho_k = \rho$, it holds for all $k > 0$

$$\begin{aligned} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} \right\|^2 \right] &\leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \frac{L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

454 **Proof** Beforehand, we provide a rewriting of the quantity $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}$ that will be useful through-
 455 out this proof:

$$\begin{aligned} \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) \\ &= -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - (1-\rho)\tilde{S}^{(k)} - \rho\mathbf{S}^{(k+1)}) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} \right] \right) \\ &= -\gamma_{k+1} \left((1-\rho) \left[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right] + \rho \left[\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)} - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right] \right) \end{aligned} \quad (32)$$

456 We observe, using the identity (32), that

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2] \leq 2\rho^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] + 2\rho^2 \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] + 2(1-\rho)^2 \mathbb{E}[\|\hat{\mathbf{s}}^{(\ell(k))} - \tilde{S}^{(k)}\|^2] \quad (33)$$

457 For the latter term, we obtain its upper bound as

$$\begin{aligned} \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)}\|^2] &= \mathbb{E} \left[\left\| \frac{1}{n} \sum_{i=1}^n (\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(k)}) - (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right\|^2 \right] \\ &\stackrel{(a)}{\leq} \mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(\ell(k))}\|^2] + \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

where (a) uses the variance inequality. We can further bound the last expectation using Lemma 1:

$$\mathbb{E}[\|\bar{\mathbf{s}}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(t_{i_k}^k)}\|^2] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\bar{\mathbf{s}}_i^{(k)} - \bar{\mathbf{s}}_i^{(t_i^k)}\|^2] \stackrel{(a)}{\leq} \frac{L_{\mathbf{s}}^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

Substituting into (33) proves the lemma. \square

C.2 Proof of Lemma 5

Lemma. Consider a decreasing stepsize $\gamma_k \in (0, 1)$ and a constant ρ , then the following inequality holds:

$$\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1-\gamma_{\ell})^2 (\mathcal{S}^{(\ell)} - \tilde{S}^{(\ell)})$$

where $\mathcal{S}^{(k)}$ is defined either by (11) (fiTTEM) or (10) (vrTTEM)

Proof We begin by writing the two-timescale update:

$$\begin{aligned} \tilde{S}^{(k+1)} &= \tilde{S}^{(k)} + \rho(\mathcal{S}^{(k+1)} - \tilde{S}^{(k)}) \\ \hat{\mathbf{s}}^{(k+1)} &= \hat{\mathbf{s}}^{(k)} + \gamma_{k+1}(\tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)}) \end{aligned} \quad (34)$$

where $\mathcal{S}^{(k+1)} = \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(t_i^k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$ according to (11). Denote $\delta^{(k+1)} = \hat{\mathbf{s}}^{(k+1)} - \tilde{S}^{(k+1)}$. Then from (34), doing the subtraction of both equations yields:

$$\delta^{(k+1)} = (1 - \gamma_{k+1})\delta^{(k)} + \frac{\rho}{1-\rho}(1 - \gamma_{k+1})(\mathcal{S}^{(k+1)} - \tilde{S}^{(k+1)})$$

Using the telescoping sum and noting that $\delta^{(0)} = 0$, we have

$$\delta^{(k+1)} \leq \frac{\rho}{1-\rho} \sum_{\ell=0}^k (1 - \gamma_{\ell+1})^2 (\mathcal{S}^{(\ell+1)} - \tilde{S}^{(\ell+1)})$$

\square

C.3 Additional Intermediary Result

Lemma 8. At iteration $k+1$, the drift term of update (11), with $\rho_{k+1} = \rho$, is equivalent to the following:

$$\begin{aligned} \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} &= \rho(\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} + \rho \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1-\rho) \left(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right) \end{aligned}$$

where we recall that $\eta_{i_k}^{(k+1)}$, defined in (18), which is the gap between the MC approximation and the expected statistics.

Proof Using the fiTTEM update $\tilde{S}^{(k+1)} = (1-\rho)\tilde{S}^{(k)} + \rho\mathcal{S}^{(k+1)}$ where $\mathcal{S}^{(k+1)} = \bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)})$ leads to the following decomposition:

$$\begin{aligned} \tilde{S}^{(k+1)} - \hat{\mathbf{s}}^{(k)} &= (1-\rho)\tilde{S}^{(k)} + \rho \left(\bar{\mathcal{S}}^{(k)} + (\tilde{S}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) - \hat{\mathbf{s}}^{(k)} + \rho\bar{\mathbf{s}}^{(k)} - \rho\bar{\mathbf{s}}^{(k)} \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho(\tilde{S}_{i_k}^{(k)} - \bar{\mathbf{s}}_{i_k}^{(k)}) + (1-\rho) \left(\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) + \rho \left(\bar{\mathcal{S}}^{(k)} - \bar{\mathbf{s}}^{(k)} + (\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) \right) \\ &= \rho(\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}) + \rho\eta_{i_k}^{(k+1)} - \rho \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] \\ &\quad + (1-\rho) \left(\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)} \right) \end{aligned}$$

476 where we observe that $\mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] = \bar{\mathbf{s}}^{(k)} - \bar{\boldsymbol{\mathcal{S}}}^{(k)}$ and which concludes the proof.

477 *Important Note:* Note that $\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}$ is not equal to $\eta_{i_k}^{(k+1)}$, defined in (18), which is the gap

478 between the MC approximation and the expected statistics. Indeed $\tilde{S}_{i_k}^{(t_{i_k}^k)}$ is not computed under the

479 same model as $\bar{\mathbf{s}}_{i_k}^{(k)}$. □

480 D Proof of Theorem 2

481 **Theorem.** Assume H1-H5. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of
 482 positive step sizes and consider the vrTTEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for
 483 any $k > 0$.

484 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\bar{L} = \max\{L_S, L_V\}$, $\rho = \frac{\mu}{c_1 \bar{L} n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$
 485 and a constant $\mu \in (0, 1)$ and $\gamma_{k+1} = \frac{1}{k^a \bar{L}}$ where $a \in (0, 1)$, we have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\tilde{\eta}^{(k+1)} + \chi^{(k+1)} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned}$$

486 **Proof** Using the smoothness of V and update (10), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (35)$$

487 Denote $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fitTEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.
 488 Taking expectations on both sides show that

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\stackrel{(a)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}\rho \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \mathbf{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(b)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}\rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1}(1-\rho) \mathbb{E}[\langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\langle \eta_{i_k}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\stackrel{(c)}{\leq} \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] + \frac{\gamma_{k+1}^2 L_V}{2} \mathbb{E}[\|\mathbf{H}_{k+1}\|^2] \\ &\quad - \gamma_{k+1}\rho \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \gamma_{k+1}(1-\rho) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \end{aligned} \quad (36)$$

489 where we have used (30) in (a) and $\mathbb{E}[\mathbf{S}^{(k+1)}] = \bar{\mathbf{s}}^{(k)} + \mathbb{E}[\eta_{i_k}^{(k+1)}]$ in (b), the growth condition in
 490 Lemma 2 and the Young's inequality with the constant equal to 1 in (c).

491 Furthermore, for $k+1 \leq \ell(k) + m$ (i.e., $k+1$ is in the same epoch as k), we have

$$\begin{aligned} &\mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] = \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} + \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} \rangle] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 \\ &\quad - 2\gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))} | \rho(\mathbf{h}_k - \eta_{i_k}^{(k+1)}) + (1-\rho)(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}) \rangle] \\ &\leq \mathbb{E}[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2], \end{aligned}$$

492 where we first used (30) and the last inequality is due to the Young's inequality.

493 Consider the following sequence

$$R_k := \mathbb{E}[V(\hat{\mathbf{s}}^{(k)}) + b_k \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2]$$

494 where $b_k := \bar{b}_{k \bmod m}$ is a periodic sequence where:

$$\bar{b}_i = \bar{b}_{i+1}(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2, \quad i = 0, 1, \dots, m-1 \quad \text{with } \bar{b}_m = 0.$$

495 Note that \bar{b}_i is decreasing with i and this implies

$$\bar{b}_i \leq \bar{b}_0 = \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \frac{(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2)^m - 1}{\gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2}, \quad i = 1, 2, \dots, m.$$

496 For $k+1 \leq \ell(k) + m$, we have the following inequality

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2) \|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[\rho \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \gamma_{k+1}^2 \|\mathbf{H}_{k+1}\|^2 + \frac{\gamma_{k+1}\rho}{\beta} \|\mathbf{h}_k\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\frac{\gamma_{k+1}\rho}{\beta} \|\eta_{i_k}^{(k+1)}\|^2 + \frac{\gamma_{k+1}(1-\rho)}{\beta} \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned}$$

497 And using Lemma 3 we obtain:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}\left[V(\hat{\mathbf{s}}^{(k)}) - (\gamma_{k+1}\rho v_{\min} + \gamma_{k+1}v_{\max}^2 - \gamma_{k+1}^2\rho^2 L_V) \|\mathbf{h}_k\|^2 + \gamma_{k+1}^2\rho^2 L_V L_{\mathbf{s}}^2 \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[(1 + \gamma_{k+1}\beta + 2\gamma_{k+1}^2\rho^2 L_{\mathbf{s}}^2) \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2 + \left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \|\mathbf{h}_k\|^2\right] \\ &\quad + \gamma_{k+1} \mathbb{E}\left[(\rho + \rho^2\gamma_{k+1} L_V) \left\|\eta_{i_k}^{(k+1)}\right\|^2 - (1-\rho - (1-\rho)^2\gamma_{k+1} L_V) \left\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\right\|^2\right] \\ &\quad + b_{k+1} \mathbb{E}\left[\left(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2\right) \|\eta_{i_k}^{(k+1)}\|^2 + \left(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2\right) \|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned}$$

498 Rearranging the terms yields:

$$\begin{aligned} R_{k+1} &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\quad + \underbrace{\left(b_{k+1}(1 + \gamma\beta + 2\gamma^2\rho^2 L_{\mathbf{s}}^2) + \gamma^2\rho^2 L_V L_{\mathbf{s}}^2\right)}_{=b_k \text{ since } k+1 \leq \ell(k) + m} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(\ell(k))}\|^2] + \tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)} \end{aligned}$$

499 where

$$\begin{aligned} \tilde{\eta}^{(k+1)} &= \left(\gamma_{k+1}(\rho + \rho^2\gamma_{k+1} L_V) + b_{k+1}(\frac{\gamma_{k+1}\rho}{\beta} + 2\gamma_{k+1}^2\rho^2)\right) \mathbb{E}\left[\left\|\eta_{i_k}^{(k+1)}\right\|^2\right] \\ \chi^{(k+1)} &= \left(b_{k+1}(\frac{\gamma_{k+1}(1-\rho)}{\beta} + 2\gamma_{k+1}^2(1-\rho)^2) - \gamma_{k+1}(1-\rho - (1-\rho)^2\gamma_{k+1} L_V)\right) \\ \tilde{\chi}^{(k+1)} &= \chi^{(k+1)} \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{\mathbf{S}}^{(k)}\|^2\right] \end{aligned}$$

500 This leads, using Lemma 2, that for any γ_{k+1} , ρ and β such that $\rho v_{\min} + v_{\max}^2 -$
 501 $\gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2) > 0$,

$$\begin{aligned} v_{\max}^2 \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2] \leq \frac{R_k - R_{k+1}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \\ &\quad + \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{\gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2))} \end{aligned}$$

502 We first remark that

$$\begin{aligned} & \gamma_{k+1}(\rho v_{\min} + v_{\max}^2 - \gamma_{k+1}\rho^2 L_V - b_{k+1}(\frac{\rho}{\beta} + 2\gamma_{k+1}\rho^2)) \\ & \geq \frac{\gamma_{k+1}\rho}{c_1}(1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1)) \end{aligned}$$

503 where $c_1 = v_{\min}^{-1}$. By setting $\bar{L} = \max\{L_s, L_V\}$, $\beta = \frac{c_1\bar{L}}{n^{1/3}}$, $\rho = \frac{\mu}{c_1\bar{L}n^{2/3}}$, $m = \frac{nc_1^2}{2\mu^2 + \mu c_1^2}$ and
 504 $\{\gamma_{k+1}\}$ any sequence of decreasing stepsizes in $(0, 1)$, it can be shown that there exists $\mu \in (0, 1)$,
 505 such that the following lower bound holds

$$\begin{aligned} & 1 - \gamma_{k+1}c_1\rho L_V - b_{k+1}(\frac{c_1}{\beta} + 2\gamma_{k+1}\rho c_1) \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \bar{b}_0(\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \geq 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{L_V\mu^2}{c_1^2n^{\frac{4}{3}}} \frac{(1 + \gamma\beta + 2\gamma^2 L_s^2)^m - 1}{\gamma\beta + 2\gamma^2 L_s^2} (\frac{n^{\frac{1}{3}}}{\bar{L}} + \frac{2\mu}{\bar{L}n^{\frac{2}{3}}}) \\ & \stackrel{(a)}{\geq} 1 - \frac{\mu}{n^{\frac{2}{3}}} - \frac{\mu}{c_1^2} (e - 1)(1 + \frac{2\mu}{n}) \geq 1 - \mu - \mu(1 + 2\mu) \frac{e - 1}{c_1^2} \stackrel{(b)}{\geq} \frac{1}{2} \end{aligned}$$

506 where the simplification in (a) is due to

$$\frac{\mu}{n} \leq \gamma\beta + 2\gamma^2 L_s^2 \leq \frac{\mu}{n} + \frac{2\mu^2}{c_1^2 n^{\frac{4}{3}}} \leq \frac{\mu c_1^2 + 2\mu^2}{c_1^2} \frac{1}{n} \text{ and } (1 + \gamma\beta + 2\gamma^2 L_s^2)^m \leq e - 1.$$

507 and the required μ in (b) can be found by solving the quadratic equation.

508 Finally, these results yield:

$$v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2(R_0 - R_{K_{\max}})}{v_{\min}\rho} + 2 \sum_{k=0}^{K_{\max}-1} \frac{\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}}{v_{\min}\rho}$$

509 Note that $R_0 = \mathbb{E}[V(\hat{s}^{(0)})]$ and if K_{\max} is a multiple of m , then $R_{\max} = \mathbb{E}[V(\hat{s}^{(K_{\max})})]$. Under the
 510 latter condition, we have

$$\sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{s}^{(k)})\|^2] \leq \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \mathbb{E}[V(\hat{s}^{(0)}) - V(\hat{s}^{(K_{\max})})] + \frac{2n^{2/3}\bar{L}}{\mu v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} [\tilde{\eta}^{(k+1)} + \tilde{\chi}^{(k+1)}]$$

511 This concludes our proof.

512

□

513 **E Proof of Theorem 3**

514 **Theorem.** Assume H1-H5. Let K_{\max} be a positive integer. Let $\{\gamma_k, k \in \mathbb{N}\}$ be a sequence of
 515 positive step sizes and consider the fitTEM sequence $\{\hat{\mathbf{s}}^{(k)}, k \in \mathbb{N}\}$ obtained with $\rho_{k+1} = \rho$ for any
 516 $k > 0$.

517 Assume that $\hat{\mathbf{s}}^{(k)} \in \mathcal{S}$ for any $k \leq K_{\max}$. By setting $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$,
 518 $\beta = \frac{c_1 \bar{L}}{n}$, $\rho = \frac{1}{n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$ and $\gamma_{k+1} = \frac{1}{k^a \alpha c_1 \bar{L}}$ where $a \in (0, 1)$, we
 519 have the following bound:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] &\leq \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ &\quad + \frac{\alpha \bar{L} n^{2/3}}{v_{\min} v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \left[\Xi^{(k+1)} + \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \right] \end{aligned}$$

520 **Proof** Using the smoothness of V and update (11), we obtain:

$$\begin{aligned} V(\hat{\mathbf{s}}^{(k+1)}) &\leq V(\hat{\mathbf{s}}^{(k)}) + \langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{L_V}{2} \|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 \\ &\leq V(\hat{\mathbf{s}}^{(k)}) - \gamma_{k+1} \langle \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)} | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle + \frac{\gamma_{k+1}^2 L_V}{2} \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2 \end{aligned} \quad (37)$$

521 Denote $\mathbf{H}_{k+1} := \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}$ the drift term of the fitTEM update in (7) and $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$.
 522 Using Lemma 8 and the additional following identity:

$$\mathbb{E} \left[(\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}) - \mathbb{E}[\bar{\mathbf{s}}_{i_k}^{(k)} - \tilde{S}_{i_k}^{(t_{i_k}^k)}] \right] = 0 \quad (38)$$

523 we have:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(k+1)})] \\ &\leq \mathbb{E}[V(\hat{\mathbf{s}}^{(k)})] - \gamma_{k+1} \rho \mathbb{E}[\langle \mathbf{h}_k | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle] - \gamma_{k+1} \mathbb{E} \left[\langle \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1 - \rho) \mathbb{E}[\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}] | \nabla V(\hat{\mathbf{s}}^{(k)}) \rangle \right] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(a)}{\leq} -v_{\min} \gamma_{k+1} \rho \mathbb{E}[\|\mathbf{h}_k\|^2] - \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \\ &\stackrel{(b)}{\leq} -(v_{\min} \gamma_{k+1} \rho + \gamma_{k+1} v_{\max}^2) \mathbb{E}[\|\mathbf{h}_k\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)} - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V}{2} \|\mathbf{H}_{k+1}\|^2 \end{aligned}$$

524 where $\xi^{(k+1)} = \mathbb{E} \left[\left\| \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] \right\|^2 \right]$. **Bounding $\mathbb{E}[\|\mathbf{H}_{k+1}\|^2]$** Using Lemma 4, we obtain:

$$\begin{aligned} &\gamma_{k+1} (v_{\min} \rho + v_{\max}^2 - \gamma_{k+1} \rho^2 L_V) \mathbb{E}[\|\mathbf{h}_k\|^2] \\ &\leq \mathbb{E} [V(\hat{\mathbf{s}}^{(k)}) - V(\hat{\mathbf{s}}^{(k+1)})] + \tilde{\xi}^{(k+1)} + \left((1 - \rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1} (1 - \rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \\ &\quad + \frac{\gamma_{k+1}^2 L_V \rho^2 L_s^2}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \end{aligned} \quad (39)$$

525 where $\tilde{\xi}^{(k+1)} = \gamma_{k+1}^2 \rho^2 \mathbf{L}_V \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] - \frac{\gamma_{k+1} \rho^2}{2} \xi^{(k+1)}$. Next, we observe that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n} \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \right) \quad (40)$$

526 where the equality holds as i_k and j_k are drawn independently. Next,

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \end{aligned}$$

527 Note that $\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} = -\gamma_{k+1}(\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}) = -\gamma_{k+1} \mathbf{H}_{k+1}$ and that in expectation we recall
528 that $\mathbb{E}[\mathbf{H}_{k+1} | \mathcal{F}_k] = \rho \mathbf{h}_k + \rho \mathbb{E}[\eta_{i_k}^{(k+1)} | \mathcal{F}_k] + (1-\rho)\mathbb{E}[\tilde{S}^{(k)} - \hat{\mathbf{s}}^{(k)}]$ where $\mathbf{h}_k = \hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}$. Thus,
529 for any $\beta > 0$, it holds

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned}$$

530 where the last inequality is due to the Young's inequality. Plugging this into (40) yields:

$$\begin{aligned} & \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &= \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + \|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + 2\langle \hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)} | \hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)} \rangle] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 + (1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\quad + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned}$$

531 Subsequently, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2] + \frac{n-1}{n^2} \sum_{i=1}^n \mathbb{E}[(1 + \gamma_{k+1}\beta)\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\|\mathbf{h}_k\|^2 \\ &\quad + \frac{\gamma_{k+1}\rho^2}{\beta}\mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] + \frac{\gamma_{k+1}(1-\rho)^2}{\beta}\mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2]] \end{aligned}$$

532 We now use Lemma 4 on $\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(k)}\|^2 = \gamma_{k+1}^2 \|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k+1)}\|^2$ and obtain:

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k+1)} - \hat{\mathbf{s}}^{(t_i^{k+1})}\|^2] \\ &\leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{\gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n} + \frac{(n-1)(1 + \gamma_{k+1}\beta)}{n^2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \\ &\leq \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right) \mathbb{E}[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2] + \sum_{i=1}^n \left(\frac{1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2}{n} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2] \\ &\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] + \left(2\gamma_{k+1}^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

533 Let us define

$$\Delta^{(k)} := \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(t_i^k)}\|^2]$$

534 From the above, we get

$$\begin{aligned} \Delta^{(k+1)} &\leq \left(1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2\right) \Delta^{(k)} + \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right) \mathbb{E}\left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2\right] \\ &\quad + \gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right) \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2\right] + \gamma_{k+1} \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2] \end{aligned}$$

535 Setting $c_1 = v_{\min}^{-1}$, $\alpha = \max\{2, 1+2v_{\min}\}$, $\bar{L} = \max\{\mathbf{L}_s, \mathbf{L}_V\}$, $\gamma_{k+1} = \frac{1}{k}$, $\beta = \frac{1}{\alpha n}$, $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$,

536 $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$, we observe that

$$1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \leq 1 - \frac{1}{n} + \frac{1}{\alpha k n} + \frac{1}{\alpha^2 c_1^2 k^2 n^{4/3}} \leq 1 - \frac{c_1(k\alpha - 1) - 1}{k\alpha n c_1} \leq 1 - \frac{1}{k\alpha n c_1}$$

537 which shows that $1 - \frac{1}{n} + \gamma_{k+1}\beta + \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2 \in (0, 1)$ for any $k > 0$. Denote $\Lambda_{(k+1)} = \frac{1}{n} -$

538 $\gamma_{k+1}\beta - \gamma_{k+1}^2 \rho^2 \mathbf{L}_s^2$ and note that $\Delta^{(0)} = 0$, thus the telescoping sum yields:

$$\begin{aligned} \Delta^{(k+1)} &\leq \sum_{\ell=0}^k \omega_{k,\ell} \left(2\gamma_{\ell+1}^2 \rho^2 + \frac{\gamma_{\ell+1}^2 \rho^2}{\beta}\right) \mathbb{E}\left[\|\bar{\mathbf{s}}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2\right] \\ &\quad + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} (1-\rho)^2 \left(2\gamma_{\ell+1} + \frac{1}{\beta}\right) \mathbb{E}\left[\|\tilde{S}^{(\ell)} - \hat{\mathbf{s}}^{(\ell)}\|^2\right] + \sum_{\ell=0}^k \omega_{k,\ell} \gamma_{\ell+1} \tilde{\epsilon}^{(\ell+1)} \end{aligned}$$

539 where $\omega_{k,\ell} = \prod_{j=\ell+1}^k (1 - \Lambda_{(j)})$ and $\tilde{\epsilon}^{(\ell+1)} = \left(2\gamma_{k+1} + \frac{\rho^2}{\beta}\right) \mathbb{E}[\|\eta_{i_k}^{(k+1)}\|^2]$.

540 Summing on both sides over $k = 0$ to $k = K_{\max} - 1$ yields:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \Delta^{(k+1)} &\leq \sum_{k=0}^{K_{\max}-1} \frac{2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}^2 \rho^2}{\beta}}{\Lambda_{(k+1)}} \mathbb{E}\left[\|\bar{\mathbf{s}}^{(k)} - \hat{\mathbf{s}}^{(k)}\|^2\right] \\ &\quad + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}(1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2\right] + \sum_{k=0}^{K_{\max}-1} \frac{\gamma_{k+1}}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)} \end{aligned}$$

541 We recall (39) where we have summed on both sides from $k = 0$ to $k = K_{\max} - 1$:

$$\begin{aligned} &\mathbb{E}[V(\hat{\mathbf{s}}^{(K_{\max})}) - V(\hat{\mathbf{s}}^{(0)})] \\ &\leq \sum_{k=0}^{K_{\max}-1} \left\{ \gamma_{k+1}(-v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}\rho^2 \mathbf{L}_V \mathbb{E}[\|\mathbf{h}_k\|^2] + \gamma^2 \mathbf{L}_V \rho^2 \mathbf{L}_s^2 \Delta^{(k)} \right\} \\ &\quad + \sum_{k=0}^{K_{\max}-1} \left\{ \tilde{\xi}^{(k+1)} + \left((1-\rho)^2 \gamma_{k+1}^2 \mathbf{L}_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) \mathbb{E}[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2] \right\} \\ &\leq \sum_{k=0}^{K_{\max}-1} \left\{ -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2 \rho^2 \mathbf{L}_V + \frac{\rho^2 \gamma_{k+1}^2 \mathbf{L}_V \mathbf{L}_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta}\right)}{\Lambda_{(k+1)}} \mathbb{E}[\|\mathbf{h}_k\|^2] \right\} \\ &\quad + \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E}\left[\|\hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)}\|^2\right] \end{aligned} \tag{41}$$

where

$$\Xi^{(k+1)} = \tilde{\xi}^{(k+1)} + \frac{\gamma_{k+1}^3 \mathbf{L}_V \rho^2 \mathbf{L}_s^2}{\Lambda_{(k+1)}} \tilde{\epsilon}^{(k+1)}$$

and

$$\Gamma_{k+1} = \left((1-\rho)^2 \gamma_{k+1}^2 L_V - \frac{\gamma_{k+1}(1-\rho)^2}{2} \right) + \frac{\gamma_{k+1}^3 L_V \rho^2 L_s^2 (1-\rho)^2 \left(2\gamma_{k+1} + \frac{1}{\beta} \right)}{\Lambda_{(k+1)}}$$

542 We now analyse the following quantity

$$\begin{aligned} & -\gamma_{k+1}(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1}^2 \rho^2 L_V + \frac{\rho^2 \gamma_{k+1}^2 L_V L_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\ & = \gamma_{k+1} \left[-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1} \rho^2 L_V + \frac{\rho^2 \gamma_{k+1} L_V L_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right] \end{aligned} \quad (42)$$

543 Furthermore, we recall that $c_1 = v_{\min}^{-1}$, $\alpha = \max\{2, 1 + 2v_{\min}\}$, $\bar{L} = \max\{L_s, L_V\}$, $\gamma_{k+1} = \frac{1}{k}$,

544 $\beta = \frac{1}{\alpha n}$, $\rho = \frac{1}{\alpha c_1 \bar{L} n^{2/3}}$, $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$, $\alpha \geq 2$. Then,

$$\begin{aligned} & \gamma_{k+1} \rho^2 L_V + \frac{\rho^2 \gamma_{k+1} L_V L_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \\ & \leq \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{n} - \gamma_{k+1}\beta - \gamma_{k+1}^2 \rho^2 L_s^2}{\bar{L}(k\alpha^2 c_1^2 n^{4/3})^{-1} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}} \right)} \\ & = \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\bar{L} \left(\frac{2}{k^2 \alpha^2 c_1^2 \bar{L}^2 n^{4/3}} + \frac{1}{k\alpha c_1^2 \bar{L}^2 n^{1/3}} \right)}{(k\alpha c_1 n^{1/3})(k\alpha - 1)c_1 - 1} \\ & \stackrel{(a)}{\leq} \frac{1}{k\alpha^2 c_1^2 \bar{L} n^{4/3}} + \frac{\frac{1}{k\alpha c_1^2 \bar{L} n^{1/3}} \left(\frac{2}{k\alpha n} + 1 \right)}{2(\alpha c_1 n^{1/3}) - 1} \\ & \leq \frac{1}{k^2 \alpha c_1^2 \bar{L} n^{4/3}} + \frac{1}{4k\alpha^2 c_1^3 \bar{L} n^{2/3}} \\ & \leq \frac{3/4}{\alpha c_1^2 \bar{L} n^{2/3}} \end{aligned} \quad (43)$$

where (a) is due to $c_1(k\alpha - 1) \geq c_1(\alpha - 1) \geq 2$ and $k\alpha c_1 n^{1/3} \geq 1$. Note also that

$$-(v_{\min}\rho + v_{\max}^2) \leq -\rho v_{\min} = -\frac{1}{\alpha c_1^2 \bar{L} n^{2/3}}$$

which yields that

$$\left[-(v_{\min}\rho + v_{\max}^2) + \gamma_{k+1} \rho^2 L_V + \frac{\rho^2 \gamma_{k+1} L_V L_s^2 \left(2\gamma_{k+1}^2 \rho^2 + \frac{\gamma_{k+1}\rho^2}{\beta} \right)}{\Lambda_{(k+1)}} \right] \leq -\frac{1/4}{\alpha c_1^2 \bar{L} n^{2/3}}$$

545 Using the Lemma 2, we know that $v_{\max}^2 \|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2 \leq \|\hat{\mathbf{s}}^{(k)} - \bar{\mathbf{s}}^{(k)}\|^2$ and using (43) on (41)
546 yields:

$$\begin{aligned} v_{\max}^2 \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ & \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned}$$

547 proving the final bound on the gradient of the Lyapunov function:

$$\begin{aligned} \sum_{k=0}^{K_{\max}-1} \gamma_{k+1} \mathbb{E}[\|\nabla V(\hat{\mathbf{s}}^{(k)})\|^2] & \leq \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} [V(\hat{\mathbf{s}}^{(0)}) - V(\hat{\mathbf{s}}^{(K_{\max})})] \\ & \quad + \frac{4\alpha \bar{L} n^{2/3}}{v_{\min}^2 v_{\max}^2} \sum_{k=0}^{K_{\max}-1} \Xi^{(k+1)} + \sum_{k=0}^{K_{\max}-1} \Gamma_{k+1} \mathbb{E} \left[\left\| \hat{\mathbf{s}}^{(k)} - \tilde{S}^{(k)} \right\|^2 \right] \end{aligned}$$

F Practical Implementations of Two-Timescale EM Methods

F.1 Application on GMM

F.1.1 Explicit Updates

We first recognize that the constraint set for θ is given by

$$\Theta = \Delta^M \times \mathbb{R}^M.$$

Using the partition of the sufficient statistics as $S(y_i, z_i) = (S^{(1)}(y_i, z_i)^\top, S^{(2)}(y_i, z_i)^\top, S^{(3)}(y_i, z_i)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$, the partition $\phi(\theta) = (\phi^{(1)}(\theta)^\top, \phi^{(2)}(\theta)^\top, \phi^{(3)}(\theta)^\top)^\top \in \mathbb{R}^{M-1} \times \mathbb{R}^{M-1} \times \mathbb{R}$ and the fact that $\mathbb{1}_{\{M\}}(z_i) = 1 - \sum_{m=1}^{M-1} \mathbb{1}_{\{m\}}(z_i)$, the complete data log-likelihood can be expressed as in (2) with

$$\begin{aligned} s_{i,m}^{(1)} &= \mathbb{1}_{\{m\}}(z_i), \quad \phi_m^{(1)}(\theta) = \left\{ \log(\omega_m) - \frac{\mu_m^2}{2} \right\} - \left\{ \log(1 - \sum_{j=1}^{M-1} \omega_j) - \frac{\mu_M^2}{2} \right\}, \\ s_{i,m}^{(2)} &= \mathbb{1}_{\{m\}}(z_i)y_i, \quad \phi_m^{(2)}(\theta) = \mu_m, \quad s_i^{(3)} = y_i, \quad \phi^{(3)}(\theta) = \mu_M, \end{aligned} \quad (44)$$

and $\psi(\theta) = -\left\{ \log(1 - \sum_{m=1}^{M-1} \omega_m) - \frac{\mu_M^2}{2\sigma^2} \right\}$. We also define for each $m \in \llbracket 1, M \rrbracket$, $j \in \llbracket 1, 3 \rrbracket$, $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$. Consider the following latent sample used to compute an approximation of the conditional expected value $\mathbb{E}_\theta[\mathbb{1}_{\{z_i=m\}}|y = y_i]$:

$$z_{i,m} \sim \mathbb{P}(z_i = m | y_i; \theta) \quad (45)$$

where $m \in \llbracket 1, M \rrbracket$, $i \in \llbracket 1, n \rrbracket$ and $\theta = (\mathbf{w}, \boldsymbol{\mu}) \in \Theta$.

In particular, given iteration $k + 1$, the computation of the approximated quantity $\tilde{S}_{i_k}^{(k)}$ during Incremental-step updates, see (8) can be written as

$$\tilde{S}_{i_k}^{(k)} = \left(\underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1}), \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})}_{:=\tilde{s}_{i_k}^{(1)}}, \underbrace{\mathbb{1}_{\{1\}}(z_{i_k,1})y_{i_k}, \dots, \mathbb{1}_{\{M-1\}}(z_{i_k,M-1})y_{i_k}}_{:=\tilde{s}_{i_k}^{(2)}}, \underbrace{y_{i_k}}_{:=\tilde{s}_{i_k}^{(3)}(\theta^{(k)})} \right)^\top. \quad (46)$$

Recall that we have used the following regularizer:

$$r(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m), \quad (47)$$

It can be shown that the regularized M-step in (4) evaluates to

$$\bar{\theta}(\mathbf{s}) = \begin{pmatrix} (1 + \epsilon M)^{-1} (s_1^{(1)} + \epsilon, \dots, s_{M-1}^{(1)} + \epsilon)^\top \\ ((s_1^{(1)} + \delta)^{-1} s_1^{(2)}, \dots, (s_{M-1}^{(1)} + \delta)^{-1} s_{M-1}^{(2)})^\top \\ (1 - \sum_{m=1}^{M-1} s_m^{(1)} + \delta)^{-1} (s^{(3)} - \sum_{m=1}^{M-1} s_m^{(2)}) \end{pmatrix} = \begin{pmatrix} \bar{\omega}(\mathbf{s}) \\ \bar{\boldsymbol{\mu}}(\mathbf{s}) \\ \bar{\mu}_M(\mathbf{s}) \end{pmatrix}. \quad (48)$$

where we have defined for all $m \in \llbracket 1, M \rrbracket$ and $j \in \llbracket 1, 3 \rrbracket$, $s_m^{(j)} = n^{-1} \sum_{i=1}^n s_{i,m}^{(j)}$.

F.1.2 Model Assumptions (GMM example)

We use the GMM example to illustrate the required assumptions.

Many practical models can satisfy the compactness of the sets as in Assumption H1. For instance, the GMM example satisfies (15) as the sufficient statistics are composed of indicator functions and observations as defined Section F.1 Equation (44).

Assumptions H2 and H3 are standard for the curved exponential family models. For GMM, the following (strongly convex) regularization $r(\theta)$ ensures H3:

$$r(\theta) = \frac{\delta}{2} \sum_{m=1}^M \mu_m^2 - \epsilon \sum_{m=1}^M \log(\omega_m) - \epsilon \log(1 - \sum_{m=1}^{M-1} \omega_m)$$

since it ensures $\theta^{(k)}$ is unique and lies in $\text{int}(\Delta^M) \times \mathbb{R}^M$. We remark that for H2, it is possible to define the Lipschitz constant L_p independently for each data y_i to yield a refined characterization.

Again, H4 is satisfied by practical models. For GMM, it can be verified by deriving the closed form expression for $B(s)$ and using H1.

Under H1 and H3, we have $\|\hat{s}^{(k)}\| < \infty$ since S is compact and $\hat{\theta}^{(k)} \in \text{int}(\Theta)$ for any $k \geq 0$ which thus ensure that the EM methods operate in a closed set throughout the optimization process.

F.1.3 Algorithms updates

In the sequel, recall that, for all $i \in \llbracket n \rrbracket$ and iteration k , the computed statistic $\tilde{S}_{i_k}^{(k)}$ is defined by (46). At iteration k , the several E-steps defined by (9) or (10) and (11) leads to the definition of the quantity $\hat{s}^{(k+1)}$. For the GMM example, after the initialization of the quantity $\hat{s}^{(0)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(0)}$, those E-steps break down as follows:

Batch EM (EM): for all $i \in \llbracket 1, n \rrbracket$, compute $\bar{s}_i^{(k)}$ and set

$$\hat{s}^{(k+1)} = n^{-1} \sum_{i=1}^n \bar{s}_i^{(k)}.$$

where $\bar{s}_i^{(k)}$ are computed using the exact conditional expected value $\mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i]$:

$$\tilde{\omega}_m(y_i; \theta) := \mathbb{E}_{\theta}[\mathbb{1}_{\{z_i=m\}} | y = y_i] = \frac{\omega_m \exp(-\frac{1}{2}(y_i - \mu_i)^2)}{\sum_{j=1}^M \omega_j \exp(-\frac{1}{2}(y_i - \mu_j)^2)},$$

Incremental EM (iEM): draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)}) = n^{-1} \sum_{i=1}^n \bar{s}_i^{(\tau_i^k)}.$$

batch SAEM (SAEM): draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} \tilde{S}^{(k)}.$$

where $= \frac{1}{n} \sum_{i=1}^n \tilde{S}_i^{(k)}$ with $\tilde{S}_i^{(k)}$ defined in (46).

Incremental SAEM (iSAEM): draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} + \frac{1}{n} (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\tau_i^k)})).$$

Variance Reduced Two-Timescale EM (vrTTEM): draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} (1 - \rho) + \rho (\tilde{S}^{(\ell(k))} + (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(\ell(k))}))).$$

Fast Incremental Two-Timescale EM (fiTTEM): draw an index i_k uniformly at random on $\llbracket n \rrbracket$, compute $\bar{s}_{i_k}^{(k)}$ and set

$$\hat{s}^{(k+1)} = \hat{s}^{(k)} (1 - \gamma_{k+1}) + \gamma_{k+1} (\tilde{S}^{(k)} (1 - \rho) + \rho (\bar{\mathcal{S}}^{(k)} + (\bar{s}_{i_k}^{(k)} - \bar{s}_{i_k}^{(t_{i_k}^k)}))).$$

Finally, the k -th update reads $\hat{\theta}^{(k+1)} = \bar{\theta}(\hat{s}^{(k+1)})$ where the function $s \rightarrow \bar{\theta}(s)$ is defined by (48).

595 F.2 Deformable Template Model for Image Analysis

596 F.2.1 Model and Updates

597 The complete model belongs to the curved exponential family, see [1], which vector of sufficient
598 statistics $S = (S_1(z), S_2(z), S_3(z))$ read:

$$\begin{aligned} S_1(z) &= \frac{1}{n} \sum_{i=1}^n S_1(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top y_i \\ S_2(z) &= \frac{1}{n} \sum_{i=1}^n S_2(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n (\mathbf{K}_p^{z_i})^\top (\mathbf{K}_p^{z_i}) \\ S_3(z) &= \frac{1}{n} \sum_{i=1}^n S_3(y_i, z_i) = \frac{1}{n} \sum_{i=1}^n z_i^t z_i \end{aligned} \quad (49)$$

599 where for any pixel $u \in \mathbb{R}^2$ and $j \in \llbracket 1, k_g \rrbracket$ we noted:

$$\mathbf{K}_p^{z_i}(x_u, j) = \mathbf{K}_p^{z_i}(x_u - \phi_i(x_u, z_i), p_j)$$

600 Finally, the Two-Timescale M-step yields the following parameter updates:

$$\bar{\theta}(\hat{s}) = \begin{pmatrix} \beta(\hat{s}) = \hat{s}_2^{-1}(z) \hat{s}_1(z) \\ \Gamma(\hat{s}) = \frac{1}{n} \hat{s}_3(z) \\ \sigma(\hat{s}) = \beta(\hat{s})^\top \hat{s}_2(z) \beta(\hat{s}) - 2\beta(\hat{s}) \hat{s}_1(z) \end{pmatrix} \quad (50)$$

601 where $\hat{s} = (\hat{s}_1(z), \hat{s}_2(z), \hat{s}_3(z))$ is the vector of statistics obtained via the SA-step (7) and using the
602 MC approximation of the sufficient statistics $(S_1(z), S_2(z), S_3(z))$ defined in (53).

603 F.2.2 Numerical Applications

604 For the inference of the template, we use the Matlab code (online SAEM) used in [16] and implement
605 our own batch, incremental, Variance reduced and Fast Incremental variants. The hyperparameters
606 are kept the same and reads as follows $M = 400$, $\gamma_k = 1/k^{0.6}$ and $p = 16$. The number of
607 landmarks for the template is $k_p = 15$ points and for the deformation $k_g = 6$ points. Both have
608 Gaussian kernels with respectively standard deviation of 0.08 and 0.16. The standard deviation of
609 the measurement errors is set to 0.1.

610 For the simulation part, we use the Carlin and Chib MCMC procedure, see [6]. Refer to [16] for
611 more details.

612 F.3 Application on PK Model

613 F.3.1 Model and Explicit Updates

614 Lognormal distributions are used for the four PK parameters:

$$\log(T_i^{\text{lag}}) \sim \mathcal{N}(\log(T_{\text{pop}}^{\text{lag}}), \omega_{T^{\text{lag}}}^2), \log(ka_i) \sim \mathcal{N}(\log(ka_{\text{pop}}), \omega_{ka}^2), \quad (51)$$

$$\log(V_i) \sim \mathcal{N}(\log(V_{\text{pop}}), \omega_V^2), \log(k_i) \sim \mathcal{N}(\log(k_{\text{pop}}), \omega_k^2). \quad (52)$$

615 We recall that the complete model (y, z) defined by (23) belongs to the curved exponential family,
616 which vector of sufficient statistics $S = (S_1(z), S_2(z), S_3(z))$ read:

$$S_1(z) = \frac{1}{n} \sum_{i=1}^n z_i, \quad S_2(z) = \frac{1}{n} \sum_{i=1}^n z_i^\top z_i, \quad S_3(z) = \frac{1}{n} \sum_{i=1}^n (y_i - f(t_i, z_i))^2 \quad (53)$$

617 where we have noted y_i and t_i the vector of observations and time for each patient i . At iter-
618 ation k , and setting the number of MC samples to 1 for the sake of clarity, the MC sampling
619 $z_i^{(k)} \sim p(z_i | y_i, \theta^{(k)})$ is performed using a Metropolis-Hastings procedure detailed in algorithm 2.

620 The quantities $\tilde{S}^{(k+1)}$ and $\hat{\mathbf{s}}^{(k+1)}$ are then updated according to the different methods. Finally the
 621 maximization step yields:

$$\bar{\boldsymbol{\theta}}(\mathbf{s}) = \begin{pmatrix} \hat{\mathbf{s}}_1^{(k+1)} \\ \hat{\mathbf{s}}_2^{(k+1)} - \hat{\mathbf{s}}_1^{(k+1)} \left(\hat{\mathbf{s}}_1^{(k+1)} \right)^\top \\ \hat{\mathbf{s}}_3^{(k+1)} \end{pmatrix} = \begin{pmatrix} \overline{\mathbf{z}_{\text{pop}}}(\hat{\mathbf{s}}^{(k+1)}) \\ \overline{\boldsymbol{\omega}_z}(\hat{\mathbf{s}}^{(k+1)}) \\ \overline{\boldsymbol{\sigma}}(\hat{\mathbf{s}}^{(k+1)}) \end{pmatrix}. \quad (54)$$

622 F.3.2 Metropolis Hastings algorithm

623 During the simulation step of the MISSO method, the sampling from the target distribution
 624 $\pi(z_i, \boldsymbol{\theta}) := p(z_i | y_i, \boldsymbol{\theta})$ is performed using a Metropolis Hastings (MH) algorithm [19] with pro-
 625 posal distribution $q(z_i, \boldsymbol{\delta})$ where $\boldsymbol{\theta} = (z_{\text{pop}}, \omega_z)$ and $\boldsymbol{\delta}$ is the vector of parameters of the proposal
 626 distribution. Commonly they parameterize a Gaussian proposal. The MH algorithm is summarized
 627 in 2.

Algorithm 2 MH algorithm

```

1: Input: initialization  $z_{i,0} \sim q(z_i; \boldsymbol{\delta})$ 
2: for  $m = 1, \dots, M$  do
3:   Sample  $z_{i,m} \sim q(z_i; \boldsymbol{\delta})$ 
4:   Sample  $u \sim \mathcal{U}([0, 1])$ 
5:   Calculate the ratio  $r = \frac{\pi(z_{i,m}; \boldsymbol{\theta}) / q(z_{i,m}; \boldsymbol{\delta})}{\pi(z_{i,m-1}; \boldsymbol{\theta}) / q(z_{i,m-1}; \boldsymbol{\delta})}$ 
6:   if  $u < r$  then
7:     Accept  $z_{i,m}$ 
8:   else
9:      $z_{i,m} \leftarrow z_{i,m-1}$ 
10:  end if
11: end for
12: Output:  $z_{i,M}$ 

```
