

MATH1860J Recitation Class 02

Foundations: Chapters 06-08

Shuhan Yu

SJTU Global College

Fall Term 2026

Outline

1. Recap for Last Week
2. Complex Numbers
3. Functions and Maps
4. Sequences
5. Exercise Solutions

1. Recap for Last Week

- Logic
- Set
- Numbers
- Countability for $\mathbb{Q}$ and $\mathbb{R}$
- Subset of Real Numbers

Limit Inferior and Limit Superior

Definition

Let (a_n) be a bounded real sequence. Define

$$b_n := \inf\{a_k : k \geq n\} \quad \text{and} \quad c_n := \sup\{a_k : k \geq n\}.$$

The **limit inferior** of (a_n) is defined as

$$\liminf_{n \rightarrow \infty} a_n := \lim_{n \rightarrow \infty} b_n = \sup_{n \in \mathbb{N}} \inf\{a_k : k \geq n\}.$$

The **limit superior** of (a_n) is defined as

$$\limsup_{n \rightarrow \infty} a_n := \lim_{n \rightarrow \infty} c_n = \inf_{n \in \mathbb{N}} \sup\{a_k : k \geq n\}.$$

Equivalently: $\overline{\lim} U$ is the maximum accumulation point/maximum limit of convergent subsequences in U , $\underline{\lim} U$ is the minimum such limit.

Limit Inferior and Limit Superior

Properties:

- (b_n) is monotonically increasing and (c_n) is monotonically decreasing.
- $\liminf_{n \rightarrow \infty} a_n \leq \limsup_{n \rightarrow \infty} a_n$.
- (a_n) converges if and only if $\liminf_{n \rightarrow \infty} a_n = \limsup_{n \rightarrow \infty} a_n$.

Exercise (1)

Consider the set $U \subset \mathbb{R}$, where $U = A \cup B \cup C$ with

$$A = \{x \in \mathbb{R} : 0 < x \leq 1\},$$

$$B = \{x \in \mathbb{R} : x = 2 - 1/n, n \in \mathbb{N} \setminus \{0\}\},$$

$$C = \{x \in \mathbb{R} : x = -1/n, n \in \mathbb{N} \setminus \{0\}\}.$$

Find $\overline{\lim} U, \underline{\lim} U$. State if one or more of them don't exist.

2. Complex Numbers

Complex Numbers

Problem: The equation $x^2 + 1 = 0$ has no solution in $\mathbb{R}$. $\mathbb{R}$ is algebraically incomplete.



Solution: Introduce a new number i such that $i^2 = -1$. Then the solutions are $x = i$ and $x = -i$. We define addition in $\mathbb{C}$ by

$$(a_1, b_1) + (a_2, b_2) := (a_1 + a_2, b_1 + b_2),$$

so in particular the sum of two real numbers is again a real number. Multiplication is defined as

$$(a_1, b_1) \cdot (a_2, b_2) := (a_1 a_2 - b_1 b_2, a_1 b_2 + a_2 b_1).$$

Neutral element: $(1, 0)$

Definition

The **modulus** or **absolute value** of z is defined by

$$|z| := \sqrt{a^2 + b^2} = \sqrt{z \cdot \bar{z}},$$

Definition

The **open ball** of radius R centered at z_0 is

$$B_R(z_0) := \{z \in \mathbb{C} : |z - z_0| < R\}.$$

A set $S \subset \mathbb{C}$ is called **bounded** if there exists some $R > 0$ such that

$$S \subset B_R(0).$$

3. Functions and Maps

Definition

Let X and Y be sets, and let P be a predicate with domain $X \times Y$. Define

$$f := \{(x, y) \in X \times Y : P(x, y)\}.$$

Assume f satisfies the property:

$$(x_1, y_1), (x_2, y_2) \in f, \quad x_1 = x_2 \implies y_1 = y_2.$$

Define the **domain** of f by

$$\text{dom } f := \{x \in X : \exists y \in Y, (x, y) \in f\}.$$

Then f is called a **function** from $\text{dom } f$ to Y . The set Y is called the **codomain** of f , and the **range** of f is defined by

$$\text{ran } f := \{y \in Y : \exists x \in X, (x, y) \in f\}.$$

Definition

Let $f : X \rightarrow \mathbb{R}$ (or $f : X \rightarrow \mathbb{C}$) be a function. We say that f is **bounded** if there exists a constant $M > 0$ such that

$$|f(x)| \leq M \quad \text{for all } x \in X.$$

- A function is **bounded above** if there exists $M \in \mathbb{R}$ such that $f(x) \leq M$ for all $x \in X$.
- A function is **bounded below** if there exists $m \in \mathbb{R}$ such that $f(x) \geq m$ for all $x \in X$.
- A function is **bounded** if and only if it is both bounded above and bounded below.

Injective, Surjective, and Bijective Functions

Definition

We say that a function $f : X \rightarrow Y$ is

- **injective** if

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

for all $x_1, x_2 \in X$.

- **surjective** if $\text{ran } f = Y$.
- **bijective** if f is both injective and surjective.

Exercise (2)

Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be functions, with $A \neq \emptyset$. Show that:

- a If $g \circ f$ is **injective**, then f is **injective**.
- b If $g \circ f$ is **surjective**, then g is **surjective**.

Definition

Let X be any set. The set X is said to be

- **finite** if there exists a bijective function $f : X \rightarrow \{1, 2, \dots, n\}$ for some $n \in \mathbb{N}$.
- **infinite** if it is not finite.
- **countably infinite** if there exists a bijective function $f : X \rightarrow \mathbb{N}$.
- **countable** if there exists an injective function $f : X \rightarrow \mathbb{N}$, and **uncountable** otherwise.

4. Sequences

Definition

Let $\mathcal{I} \subset \mathbb{N}$ be an infinite set. A map $\mathcal{I} \rightarrow \mathbb{R}$ is called a **real sequence**, a map $\mathcal{I} \rightarrow \mathbb{C}$ is called a **complex sequence**, or simply a **sequence**.

- We denote the values of a sequence by a_n , i.e., a sequence maps

$$n \mapsto a_n, \quad n \in \mathcal{I} \subset \mathbb{N}.$$

- For simplicity, we often consider the case $\mathcal{I} = \mathbb{N}$. Generalizations are straightforward.
- Formally, a sequence is the set

$$\{(n, a_n) : n \in \mathcal{I}\}.$$

For simplicity, we write

$$(a_n)_{n \in \mathbb{N}} = (a_n) = (a_0, a_1, a_2, \dots) = a_0; a_1; a_2; \dots$$

The values a_n are also called the **terms** of the sequence.

Definition

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence. A **subsequence** of (a_n) is a sequence of the form $(a_{n_k})_{k \in \mathbb{N}}$, where $(n_k)_{k \in \mathbb{N}}$ is a strictly increasing sequence of natural numbers, i.e.,

$$n_0 < n_1 < n_2 < \cdots$$

- Informally, a subsequence is obtained by selecting infinitely many terms from the original sequence while preserving their order.
- Example: If $a_n = \frac{1}{n}$, then $a_{2k} = \frac{1}{2k}$ is a subsequence.

Convergence

Definition

A complex sequence $(a_n) : \mathcal{I} \rightarrow \mathbb{C}$, $\mathcal{I} \subset \mathbb{N}$, is said to **converge** with limit $a \in \mathbb{C}$ if

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ such that } n > N \implies a_n \in B_\varepsilon(a),$$

where

$$B_\varepsilon(a) := \{z \in \mathbb{C} : |z - a| < \varepsilon\}$$

is called the ε -neighborhood of a .

Equivalently, we write

$$\lim_{n \rightarrow \infty} a_n = a \quad \text{or} \quad a_n \rightarrow a \text{ as } n \rightarrow \infty.$$

A sequence that does not converge (to any limit) is said to **diverge**.

Definition

A real sequence (a_n) is said to **diverge to infinity** if

$$\forall C > 0, \exists N \in \mathbb{N} \text{ such that } n > N \implies a_n > C.$$

We then write

$$\lim_{n \rightarrow \infty} a_n = \infty \quad \text{or} \quad a_n \rightarrow \infty \text{ as } n \rightarrow \infty.$$

A real sequence (a_n) is said to **diverge to minus infinity** if

$$\forall C < 0, \exists N \in \mathbb{N} \text{ such that } n > N \implies a_n < C.$$

In this case, we write

$$\lim_{n \rightarrow \infty} a_n = -\infty \quad \text{or} \quad a_n \rightarrow -\infty \text{ as } n \rightarrow \infty.$$

Convergence

Lemma

A convergent sequence is bounded.

Lemma

For a sequence (a_n) and $a \in \mathbb{C}$:

$$a_n \rightarrow a \iff a_n - a \rightarrow 0, \quad a_n \rightarrow 0 \iff |a_n| \rightarrow 0.$$

Lemma

A convergent sequence has exactly one limit.

Convergence

Definition

Let (a_n) and (b_n) be convergent sequences with

$$a_n \rightarrow a \quad \text{and} \quad b_n \rightarrow b, \quad a, b \in \mathbb{C}.$$

Then the following hold:

- ① $\lim_{n \rightarrow \infty} (a_n + b_n) = a + b$
- ② $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = ab$
- ③ $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$ if $b \neq 0$

Exercise (3)

Prove the third lemma.

Theorem (Squeeze Theorem)

Let (a_n) , (b_n) , and (c_n) be real sequences such that

$$a_n \leq c_n \leq b_n$$

for all sufficiently large n . Suppose that

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = a.$$

Then the sequence (c_n) converges, and

$$\lim_{n \rightarrow \infty} c_n = a.$$

Exercise (4)

- 1 Prove that $\lim_{n \rightarrow \infty} n^{1/n} = 1$.
- 2 Prove that $\lim_{n \rightarrow \infty} nq^n = 0$, $|q| < 1$.
- 3 Let (a_n) be a sequence converging to a . Prove that the sequence $b_n = \left(\frac{1}{n} \sum_{j=1}^n a_j \right)$ is convergent with limit a .
- 4 Consider the sequence defined by $a_1 = 2$, $a_{n+1} = 1 + \frac{1}{a_n}$. Show that (a_n) is convergent and find its limit.

Convergence

- ① (2.2.24) Every monotonic and bounded sequence (a_n) converges to its supremum or infimum.
- ② (2.2.30) Let (a_n) be a convergent sequence with $\lim a_n = a$. Then any subsequence of (a_n) is convergent with the same limit.
- ③ (2.2.31) Every real sequence has a monotonic subsequence.
- ④ (2.2.33) If a sequence converges, then the limit is the only accumulation point.
- ⑤ (2.2.35) A number a is an accumulation point of a sequence (a_n) iff there exists a subsequence of (a_n) that converges to a .
- ⑥ (2.2.36) **Bolzano-Weierstrass Theorem:** Every bounded real sequence has an accumulation point.

Reminders

- 1 Get used to $B_\epsilon(a)$.
- 2 When you're dealing with limits, the finite elements don't matter. Things beyond N matter.
- 3 When dealing with some complex sequences, you could consider the squeeze theorem to transform the problem into investigating some easier sequences.
- 4 The terms of sequences are discrete points.
For the properties of continuous functions, you can't take it for granted when dealing with sequences.

5. Exercise Solutions

Solution to Exercise 1

Let

$$U = A \cup B \cup C,$$

where

$$A = (0, 1], \quad B = \left\{ 2 - \frac{1}{n} \mid n \in \mathbb{N}^+ \right\}, \quad C = \left\{ -\frac{1}{n} \mid n \in \mathbb{N}^+ \right\}.$$

- ① $\lim U$ (limit inferior): The accumulation points of U are $[0, 1] \cup \{2\}$. The smallest accumulation point is 0. Hence $\lim U = 0$.
- ② $\overline{\lim} U$ (limit superior): The largest accumulation point is 2. Hence $\overline{\lim} U = 2$.

Equivalently: $\overline{\lim} U$ is the maximum limit of convergent subsequences in U , $\lim U$ is the minimum such limit.

Solution to Exercise 2

Proof.

(a) Suppose $g \circ f$ is injective. To show f is injective, assume $f(x_1) = f(x_2)$ for some $x_1, x_2 \in A$. Then

$$(g \circ f)(x_1) = g(f(x_1)) = g(f(x_2)) = (g \circ f)(x_2).$$

Since $g \circ f$ is injective, we have $x_1 = x_2$. Therefore, f is injective.

(b) Suppose $g \circ f$ is surjective. To show g is surjective, let $c \in C$ be arbitrary. Since $g \circ f$ is surjective, there exists $a \in A$ such that $(g \circ f)(a) = c$. Let $b = f(a) \in B$. Then

$$g(b) = g(f(a)) = (g \circ f)(a) = c.$$

Therefore, g is surjective. □

Solution to Exercise 3

Proof.

We need to prove: If $a_n \rightarrow a$ and $b_n \rightarrow b$ with $b \neq 0$, then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{a}{b}$.

Step 1: Show that $\frac{1}{b_n} \rightarrow \frac{1}{b}$.

Since $b \neq 0$, there exists N_1 such that for all $n > N_1$, we have $|b_n - b| < \frac{|b|}{2}$. Thus

$$|b_n| \geq |b| - |b_n - b| > |b| - \frac{|b|}{2} = \frac{|b|}{2}.$$

For any $\varepsilon > 0$, there exists N_2 such that for all $n > N_2$,
 $|b_n - b| < \frac{|b|^2 \varepsilon}{2}$.



Solution to Exercise 3 (continued)

Proof (continued).

For $n > \max\{N_1, N_2\}$:

$$\left| \frac{1}{b_n} - \frac{1}{b} \right| = \frac{|b - b_n|}{|b_n| \cdot |b|} < \frac{|b|^2 \varepsilon / 2}{(|b|/2) \cdot |b|} = \varepsilon.$$

Step 2: Apply the product rule.

Since $a_n \rightarrow a$ and $\frac{1}{b_n} \rightarrow \frac{1}{b}$, by the product rule (Lemma 2 from the convergence properties):

$$\frac{a_n}{b_n} = a_n \cdot \frac{1}{b_n} \rightarrow a \cdot \frac{1}{b} = \frac{a}{b}.$$



Solution to Exercise 4(1)

Proof.

Let $a_n = n^{1/n}$. For $n \geq 2$, we have $a_n > 1$. Write $a_n = 1 + h_n$ where $h_n > 0$.

By the binomial theorem:

$$n = a_n^n = (1 + h_n)^n \geq 1 + nh_n + \frac{n(n-1)}{2}h_n^2.$$

Therefore:

$$\frac{n(n-1)}{2}h_n^2 \leq n-1 \implies h_n^2 \leq \frac{2(n-1)}{n(n-1)} = \frac{2}{n}.$$

Thus $0 < h_n \leq \sqrt{\frac{2}{n}}$. Since $\sqrt{\frac{2}{n}} \rightarrow 0$ as $n \rightarrow \infty$, by the squeeze theorem, $h_n \rightarrow 0$.

Therefore, $n^{1/n} = 1 + h_n \rightarrow 1$. □

Solution to Exercise 4(2)

Proof.

Since $|q| < 1$, we can write $|q| = \frac{1}{1+r}$ for some $r > 0$.

By the binomial theorem, for $n \geq 2$:

$$(1+r)^n = \sum_{k=0}^n \binom{n}{k} r^k \geq \binom{n}{2} r^2 = \frac{n(n-1)}{2} r^2.$$

Therefore:

$$|nq^n| = n|q|^n = \frac{n}{(1+r)^n} \leq \frac{n}{\frac{n(n-1)}{2} r^2} = \frac{2}{(n-1)r^2}.$$

Since $\frac{2}{(n-1)r^2} \rightarrow 0$ as $n \rightarrow \infty$, by the squeeze theorem:

$$\lim_{n \rightarrow \infty} nq^n = 0.$$



Solution to Exercise 4(3)

Proof.

Let $a_n \rightarrow a$. We prove that $\left(\frac{1}{n} \sum_{j=1}^n a_j\right) \rightarrow a$.

Let $\varepsilon > 0$. Since $a_n \rightarrow a$, there exists N such that for all $j > N$, $|a_j - a| < \frac{\varepsilon}{2}$.

For $n > N$, write:

$$\left| \frac{1}{n} \sum_{j=1}^n a_j - a \right| = \left| \frac{1}{n} \sum_{j=1}^n (a_j - a) \right| \leq \frac{1}{n} \sum_{j=1}^n |a_j - a|.$$



Solution to Exercise 4(3) (continued)

Proof (continued).

Split the sum:

$$\frac{1}{n} \sum_{j=1}^n |a_j - a| = \frac{1}{n} \sum_{j=1}^N |a_j - a| + \frac{1}{n} \sum_{j=N+1}^n |a_j - a|.$$

The first term: $\frac{1}{n} \sum_{j=1}^N |a_j - a| \leq \frac{NM}{n} < \frac{\varepsilon}{2}$ for sufficiently large n , where

$$M = \max_{1 \leq j \leq N} |a_j - a|.$$

The second term: $\frac{1}{n} \sum_{j=N+1}^n |a_j - a| < \frac{n-N}{n} \cdot \frac{\varepsilon}{2} < \frac{\varepsilon}{2}.$



Solution to Exercise 4(4)

Proof.

Consider $a_1 = 2$, $a_{n+1} = 1 + \frac{1}{a_n}$.

Step 1: Show that (a_n) is bounded. We claim $1 < a_n < 2$ for all $n \geq 2$ (by induction).

Base: $a_2 = 1 + \frac{1}{2} = \frac{3}{2} \in (1, 2)$.

Step: If $1 < a_n < 2$, then $a_{n+1} = 1 + \frac{1}{a_n} \in (1 + \frac{1}{2}, 1 + 1) = (\frac{3}{2}, 2)$.

Step 2: Suppose (a_n) converges to L . Then taking limits in $a_{n+1} = 1 + \frac{1}{a_n}$:

$$L = 1 + \frac{1}{L} \implies L^2 - L - 1 = 0 \implies L = \frac{1 + \sqrt{5}}{2} \text{ (golden ratio).}$$

Step 3: To show convergence, observe that odd/even subsequences are monotonic and bounded, hence converge (by Theorem 2.2.24). Both must converge to the same limit by the recurrence relation. □

Thanks for Listening!

Any questions? Please feel free to ask!

References:

- Horst Hohberger, *math186_all_lecture_slides (2026)*
- Wenxin Chen, *MATH1860J RC*, 25 Fall
- Leqi Tang, *MATH1860J RC*, 25 Fall
- Zhun Wang, *MATH1860J RC*, 25 Fall

To be continued...

- Metric Spaces
- Cauchy Sequences

Enjoy the National Holiday!