

The following is a list of approved medical abbreviations. In general, the use of abbreviations should be limited to this list.

A&O x 3 - alert and oriented to person, place and time

A&O x 4 - alert and oriented to person, place, time and event

A-FIB - atrial fibrillation

AAA - abdominal aortic aneurysm
ABC - airway, breathing, circulation

ABD - abdomen (abdominal)

ACLS - advanced cardiac life support
AKA - above the knee amputation
ALS - advanced life support
- against medical advice

AMS - altered mental status

AMT - amount

APPROX - approximately

ASA - aspirin - associated

BG - blood glucose BILAT - bilateral

BKA - below the knee amputation

BLS - basic life support

BM - bowel movement

BP - blood pressure

BS - breath sounds

BVM - bag-valve-mask

C-SECTION - caesarean section
C-SPINE - cervical spine

C/O - complaint of (complains of)

CA - cancer

CABG - coronary artery bypass graft - coronary artery disease

CATH - catheter

CC - chief complaint

CEPH - cephalic

CHF - congestive heart failure
CNS - central nervous system

COPD - chronic obstructive pulmonary disease

CP - chest pain

CPR - cardiopulmonary resuscitation

CSF - cerebrospinal fluid

CT - cat scan

CVA - cerebrovascular accident (stroke)



D5W - 5% dextrose in water
DKA - diabetic ketoacidosis
DNR - do not resuscitate
DOA - dead on arrival
DT - delirium tremens

Dx - diagnosis

ECG - electrocardiogram - electroencephelogram

ET - endotracheal
ETOH - ethanol (alcohol)
ETT - endotracheal tube
EXT - external (extension)

FB - foreign body
FLEX - flexion
Fx - fracture

g - gram(s)

GI - gastrointestinal - gunshot wound

gtts - drops

GU - gastrourinary

GYN - gynecology (gynecological)

H/A - headache

HEENT - head, eyes, ears, nose, throat

HR - heart rate (hour)
HTN - hypertension

Hx - history

ICP - intracranial pressure
ICU - intensive care unit
IM - intramuscular
IV - intravenous

JVD - jugular vein distension

kg - kilogram

KVO - keep vein open



L-SPINE - lumbar spine

L/S-SPINE - lumbarsacral spine L&D - labor and delivery

LAT - lateral lb - pound

LLQ - left lower quadrant LMP - last mestrual period

LOC - level of consciousness (loss of consciousness)

LR - lactated ringers LUQ - left upper quadrant

MAST - military anti-shock trousers

mcg - microgram(s)
MED - medicine
mg - milligram(s)

MI - myocardial infarction (heart attack)

min - minimum / minute MS - mental status

MS - mental status change

MSO4 - morphine

MVC - motor vehicle crash

N/V - nausea/vomiting

N/V/D - nausea/vomiting/diarrheaNAD - no apparant distressNC - nasal cannula

NEB - nebulizer

NKDA - no known drug allergies

NRB - non-rebreather
NS - normal saline

NSR - normal sinus rhythm

OB/GYN - obstetrics/gynecology

PALP - palpation

PAC - premature atrial contraction

PE - pulmonary embolus

PEARL - pupils equal and reactive to light

PMHx - past medical history

PO - orally

PRB - partial rebreather

PRN - as needed PT - patient

PVC - premature ventricular contraction



RLQ - right lower quadrant RUQ - right upper quadrant

Rx - medicine RXN - reaction

S/P - status post

SOB - shortness of breath
SQ - subcutaneous
ST - sinus tachycardia

SVT - supraventricular tachycardia

Sx - symptom SZ - seizure

T-SPINE - thoracic spine - temperature

TIA - transient ischemic attack

TKO - to keep open (refers to IV's - same as KVO)

Tx - treatment

UOA - upon our arrival

URI - upper respiratory infection
UTI - urinary tract infection

VF - ventricular fibrillation

VS - vital signs

VT - ventricular tachycardia

WAP - wandering atrial pacemaker

WNL - within normal limits

YO (YOA) - years old (years of age)

M or ♂ - male
F or ♀ - female
+ - positive
- negative
? - questionable

Ψ - psychiatric
- approximately
- greater than
- less than
- equal



↑	upper (increased)beforeafterwithwithout
Δ L R ↓ 1° 2°	changeleftrightlower (decreased)primarysecondary