



# Asthma Attack!

by

Hollie L. Leavitt  
Department of Biology  
College of Western Idaho, Nampa, ID

## Part I – Background Information

Jaden Hunter, 11 years old, stumbled into his parents' bedroom. It was 4 a.m. and completely dark. Unable to talk, he simply grabbed his mother's foot to get her attention. His mom awoke immediately and shouted "Brandon, get up!" Without explanation, she knew what the problem was. She helped Jaden onto the bed next to her husband, then raced down the stairs and grabbed the backpack full of medical equipment they used when things got this bad.

When she returned Jaden was sitting on the bed being comforted by his father. While slipping a mask over his face to deliver nebulized albuterol, his mother recalled the events earlier in the day. Jaden had enjoyed a field trip with his STEM club where they attended the local race track to learn about the engineering of race cars. At one point during the event, one of the drivers offered to start up the car for the group of excited school kids. He warned "Before I do this, if you have asthma you'll want to step way back." A happy and excited Jaden didn't want to leave, but his mom recognized the danger and led him away from the car. Just a few moments later the driver revved the engine. The deafening sound was followed by a plume of green smoke—nitromethane exhaust. Jaden immediately needed his quick-relief rescue inhaler, and his mother took him home.

Asthma is a condition caused by chronic inflammation of the small airways in the lungs. This leads to swelling and increased mucus production within conducting zone passageways. Due to the chronic inflammation, an asthmatic's airways are already more narrow than the airways of an individual without this disease (Figure 1). Situations that may cause the airways to constrict or spasm are common. Exposure to dry and/or cold air, contact with pollen or other allergens, illnesses such as a cold or the flu, certain medications and foods, or even just stress can cause this bronchoconstriction to occur (AAAAI, *n.d.*). This usually does not cause a problem for a non-asthmatic, but for someone with asthma, bronchoconstriction can severely decrease the diameter of the already swollen, mucus-producing airways, making it very difficult to move air into and out of the lungs. The decrease in air flow can range from mild to life-threatening, and may cause a great deal of anxiety for someone actively suffering from an acute asthma exacerbation, or asthma attack.




Figure 1. The respiratory passageways in a non-asthmatic (L) and asthmatic (R) lung. Credit: BruceBlaus, CC BY 3.0.

## Questions

You may need to use the internet or another source to help answer some of the questions in this case study.

1. According to the passage above, what are the three factors involving the airways that lead to an asthma attack?
2. What is contained within a quick-relief rescue inhaler?
3. How did the quick-relief rescue inhaler help Jaden to breathe easier at the race track?
4. What is nebulized albuterol, and how does it help with an asthma exacerbation?

Answers:

1)

3 factors that lead to an asthma attack:

- swelling of airways
- increased mucus production
- bronchoconstriction

In asthmatics, airways are chronically inflamed, meaning swelling and increased mucus production are constant. When they have bronchoconstriction on top of this, it leads to acute asthma exacerbation.

2)

most quick relief rescue inhalers contain a medication called albuterol, which is a bronchodilator

3)

The the bronchodilating medication albuterol opens up the airways so there is less resistance to airflow in and out of the lungs

4)

Nebulized albuterol is liquid albuterol that is broken up into a fine mist (or nebulized) by a machine. This fine mist is delivered to the lungs through an oxygen mask for an extended period of time (usually several minutes). In the case of more severe asthma exacerbations, nebulized albuterol can be preferable to a quick relief rescue inhaler.

## Part II – The Oxygen-Hemoglobin Dissociation Curve

Following his nebulizer treatment, Jaden was breathing a little better and able to talk.

“I used my rescue inhaler all night . . . it didn’t help!” He paused for a breath and then said, “We used my nebulizer . . . right before bed . . . I still couldn’t sleep . . . My chest feels tight.”

“Do you feel like we need to go to the hospital?” his mom gently asked. Jaden looked down and simply nodded his head yes. Having suffered from asthma since the age of four, he understood that his symptoms had reached a point where they couldn’t be adequately managed at home.

Luckily, the hospital was just a short drive away. Brandon stayed home with Jaden’s little sister, Chloe, and Jaden and his mom headed to the car. It wasn’t the first time they had made this trip in the middle of the night. In fact, it seemed like every time Jaden had breathing problems severe enough to warrant an emergency room trip it was in the middle of the night.

As they arrived at the hospital Jaden’s chest tightness was once again beginning to increase and he was unable to take deep breaths. The panic this caused him led to tears, which only made his breathing problems worse. His mom knew that the best thing she could do to help him was to stay calm herself and gently guide him through breathing exercises, so she set aside her own fears and panic, and started to do just that as she checked him into the ER.

Once back in the examining room a pulse oximeter was immediately placed on Jaden’s finger and vital signs monitored. Jaden’s oxygen saturation ( $SO_2$ ), or the percentage of his hemoglobin binding sites bound to oxygen, was 82%.

Use Figure 2 of the oxygen-hemoglobin dissociation curve to answer the questions below.




Figure 2. The oxygen-hemoglobin dissociation curve.  
Credit: OpenStax College, CC BY 3.0.

### Questions

1. Based on his  $SO_2$ , what is the partial pressure of oxygen ( $PO_2$ ) dissolved in Jaden’s blood plasma?
2. How does Jaden’s  $PO_2$  compare to the normal, expected  $PO_2$  for arterial blood?
3. Provide a reason for the change from normal arterial  $PO_2$ , and explain how this change affects oxygen delivery to Jaden’s tissues.

## Answers

1)

The  $\text{PO}_2$  of Jaden's blood is  $\approx 50\text{mm Hg}$  (for how this value was obtained, see the annotations on the graph found on the page above)

2)

The expected  $\text{PO}_2$  for arterial blood is  $80\text{--}100\text{mm Hg}$ ; Jaden's  $\text{PO}_2$  is lower than normal

3)

The airway obstruction caused by bronchoconstriction, inflammation, and mucus build-up is making it difficult for Jaden to fill his lungs during inhalation. With less  $\text{O}_2$  in the lungs, less  $\text{O}_2$  diffuses into the blood. His blood is not being effectively oxygenated.

### Part III – The Effects of Air Flow on $P_{O_2}$ and $P_{CO_2}$

Because asthma is an obstructive respiratory disease caused by a decreased diameter in the respiratory passageways, Jaden was having problems both with breathing in and with breathing out. Air flow through the respiratory passageways can be calculated by using the following formula:

$$F = \Delta P / R$$

Where:

$F$  = air flow

$\Delta P$  = the difference between atmospheric and intrapulmonary pressure

$R$  = resistance

#### Questions

1. How does the body create a difference between atmospheric and intrapulmonary pressures to cause air flow to and from the lungs?
2. In Jaden's case, which factor in the above equation changed, causing air flow to and from his lungs to decrease?
3. Without medication or treatment, how could Jaden compensate to maintain airflow to and from his lungs despite his narrowed airways? In your answer, be sure to reference the formula for air flow given above.
4. Explain why asthma exacerbations and other obstructive lung diseases that make it difficult for air to move into and out of the lungs can be exhausting for the sufferer. (*Note:* this goes beyond the fact that the sufferer is oxygen deficient and that often the problem happens at night. You need to discuss the mechanical strain that this places on the body due to the requirements to maintain airflow to and from the lungs.)
5. The  $P_{CO_2}$  of venous blood is normally 45 mmHg. How would you expect Jaden's current  $P_{CO_2}$  level to compare to the normal level? Explain your answer.
6. How would Jaden's  $P_{CO_2}$  level affect the rate at which his oxygen is dissociating from hemoglobin? (Remember the Bohr effect.)

1)

movement of the diaphragm and other respiratory muscles influences the volume of the thoracic cavity. As the thoracic cavity volume changes, so does the pressure inside of it. These changes in pressure within the thoracic cavity (and within the lungs) drive airflow to and from the lungs.

2)

The resistance (R) in Jaden's airways had really increased due to bronchoconstriction, inflammation, and mucus build up.

3)

In order to overcome the increased resistance and maintain airflow, Jaden would have to increase his  $\Delta P$ , because  $F = \Delta P/R$ . To do this, Jaden would need to increase the volume of the thoracic cavity more than normal (to drop the pressure in the lungs more than normal) during inhalation, and decrease the volume of the thoracic cavity (to increase the pressure in the lungs more than normal) during exhalation.

4)

To increase (and decrease) the volume of the thoracic cavity more than normal to allow to airflow during an asthma attack, an asthmatic will need to contract additional muscles with every breath. On inspiration, they may be activating the sternocleidomastoid, scalenes, serratus anterior, pectoralis minor, and upper trapezius in addition to the muscles normally activated (external intercostals and diaphragm). Under quiet breathing situations in people with healthy lung function, expiration is entirely passive requiring no energy. For an asthmatic suffering from an exacerbation, exhalation may require activation muscles. Using additional muscles during inhalational and exhalation requires a lot of extra energy and can cause breathing to be tiring.

5)

because airway obstruction associated with an asthma attack also makes it difficult to breathe out, Jaden is having a hard time clearing the  $\text{CO}_2$  that is diffusing into his lungs and it is instead building up in the lungs. With increased alveolar  $\text{CO}_2$ , it becomes more difficult for  $\text{CO}_2$  to diffuse from the blood into the lungs, so  $\text{Pco}_2$  begins to increase as well. You would expect Jaden's  $\text{Pco}_2$  to be above normal

6)

the high carbon dioxide levels in Jaden's blood are causing oxygen to dissociate from his hemoglobin at a rate faster than normal

## Part IV – Spirometry

Spirometry is a test that is done as part of a normal, routine check-up for an asthmatic. In addition, it is also often performed during asthma exacerbations to assess lung function. During the test a patient is required to breathe through a tube that measures air flow into and out of their lungs. Usually the patient is asked to breathe in and out normally, before being required to breathe in as deeply as possible and then breathe out as deeply and quickly as possible.

Familiarize yourself with Figure 3 and then answer the questions below.




Figure 3. Respiratory volumes. Credit: LungVolume.jpg, CC BY-SA 3.0, <<https://commons.wikimedia.org/wiki/File:Lungvolumes.svg>>.

### Questions

1. Provide definitions for the following: *tidal volume*, *inspiratory reserve volume*, and *expiratory reserve volume*.
2. How do you think that each of these lung volumes (tidal, inspiratory reserve, and expiratory reserve) would change for someone experiencing an asthma attack?
3. Explain why spirometry (the measuring of these lung volumes) can be used to get an idea of basic lung function and determine whether or not someone is currently suffering from asthma or another respiratory disorder.

1)

- tidal volume:
  - the volume of air inhaled or exhaled during normal quiet breathing
- inspiratory reserve volume:
  - the amount of air that can be inhaled after normal tidal volume inhalation
- expiratory volume:
  - the amount of air that can be exhaled after a normal tidal volume exhalation

2)

Because asthma is an obstruction to air flow, both breathing in and breathing out are affected and become more difficult. For that reason, ALL of the lung volumes given here will decrease in someone with an asthma attack.

3)

Spirometry measures lung volumes which we know decrease with an asthma attack due to resistance to airflow in the airways. Therefore, if a person has a lower than expected lung volume during a spirometry test, it would be indicative that there is some obstruction to airflow.

study:

- respiratory case studies will likely show up on the final



## Part V – Treatment

Jaden's asthma specialist, Dr. Palmer, was on call at the hospital and popped into the ER to see how Jaden was doing. Jaden had been placed on a continuous nebulizer, given 20 mg of prednisone, and was continuing to have his vitals monitored. His  $\text{SO}_2$  had increased to 95% and had remained there for approximately an hour.

"Well, it looks like we've had some improvement," the doctor said. "Blood oxygen levels have been above 90% for a while now. Let's discontinue the nebulizer. If we can keep his  $\text{SO}_2$  above 90% without it, then we'll let you guys out of here soon."

It was now 7 a.m. and Jaden was dozing on the table. Dr. Palmer put his hand on Jaden's shoulder to gently wake him and said, "It seems like the last time I saw you we put you on a long-term asthma preventative—Advair was it? Have you been taking your Advair twice a day like we talked about?"

"Well, I take it sometimes. But I forgot to take it for a while because I couldn't find the inhaler."

"Jaden, if you don't want to come to the hospital in the middle of the night, or at any other time for that matter, you've got to take your Advair like we talked about. We're going to keep you on prednisone for the next five days. Please take 20 mg twice a day and continue to use your nebulizer every 4–6 hours until you feel like you are breathing better and your chest is no longer tight."

An hour later Jaden and his mom headed home after a very long night.

### Questions

1. Use the oxygen-hemoglobin dissociation curve (Figure 2 in Part II) to determine what Jaden's  $\text{PO}_2$  is if  $\text{SO}_2$  is 95%.
2. Why is Dr. Palmer concerned that Jaden's  $\text{SO}_2$  stay above 90%?
3. How do long-term asthma control medications such as Advair work to prevent asthma attacks?
4. How did prednisone help to reverse Jaden's symptoms?

### Internet Resources

AAAAI (American Academy of Allergy, Asthma and Immunology). *n.d.* Asthma triggers and management. <<https://www.aaaai.org/conditions-and-treatments/library/asthma-library/asthma-triggers-and-management>>.

Mayo Clinic. *n.d.* Hypoxemia. <<http://www.mayoclinic.org/symptoms/hypoxemia/basics/definition/sym-20050930>>.

Bottrell, J. 2015. Understanding oxygen and oxygen levels with COPD. <<http://www.healthcentral.com/asthma/c/52325/175572/understanding-oxygen-levels/>>.

National Center for Biotechnology Information. PubChem Compound Database. Prednisone, *CID=5865*. <<https://pubchem.ncbi.nlm.nih.gov/compound/5865>>.