

# Lecture 7b: Non-Bayesian Information Processing

EC 404: Behavioral Economics  
Professor: Ben Bushong

December 9, 2021

# An Alternative Model: The Law of Small Numbers

## An Alternative Model: The Law of Small Numbers

Based on Rabin (*QJE* 2002)

Assumes one particular misconception of randomness: People act as if they believe recent draws of one outcome increase the odds of the next draw being a different outcome.

Specifically, an *N-Freddy* acts as if he believes that probabilistic events are drawn from an “urn” with  $N$  balls without replacement — specifically, he believes that the ball just drawn is not in the “urn” for the current draw.

# An Alternative Model: The Law of Small Numbers

## An Alternative Model: The Law of Small Numbers

Based on Rabin (*QJE* 2002)

Assumes one particular misconception of randomness: People act as if they believe recent draws of one outcome increase the odds of the next draw being a different outcome.

Specifically, an *N-Freddy* acts as if he believes that probabilistic events are drawn from an “urn” with  $N$  balls without replacement — specifically, he believes that the ball just drawn is not in the “urn” for the current draw.

# An Alternative Model: The Law of Small Numbers

## An Alternative Model: The Law of Small Numbers

Based on Rabin (*QJE* 2002)

Assumes one particular misconception of randomness: People act as if they believe recent draws of one outcome increase the odds of the next draw being a different outcome.

Specifically, an *N-Freddy* acts as if he believes that probabilistic events are drawn from an “urn” with  $N$  balls without replacement — specifically, he believes that the ball just drawn is not in the “urn” for the current draw.

# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = ?$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = ?$

# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = ?$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = ?$

# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = ?$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = ?$

# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = ?$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = ?$



# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = \text{ ? }$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = \text{ ? }$

# An Example

Suppose ( $N = 12$ )-Freddy faces a die with 4  $H$ 's and 2  $M$ 's.

On the first roll:  $\Pr(H) = 2/3$

On the second roll:

- ▶ If first roll is  $H$ , then  $\Pr(H) = 7/11 < 2/3$
- ▶ If first roll is  $M$ , then  $\Pr(H) = 8/11 > 2/3$

More generally:

- ▶ If previous roll is  $H$ , then  $\Pr(H) = ?$
- ▶ If previous roll is  $M$ , then  $\Pr(H) = ?$

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).



# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Formal Assumptions

Suppose the objective probability of outcome  $x$  is  $p$ .

On the first draw:

- ▶  $N$ -Freddy believes  $\Pr(x) = p$ .

On the second draw:

- ▶ If first draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If first draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

More generally, on  $n^{\text{th}}$  draw (where  $n \geq 2$ ):

- ▶ If  $(n-1)^{\text{st}}$  draw is  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN-1}{N-1}$ .
- ▶ If  $(n-1)^{\text{st}}$  draw NOT  $x$ ,  $N$ -Freddy believes  $\Pr(x) = \frac{pN}{N-1}$ .

A nice feature: The larger is  $N$ , the smaller is the bias  
(and as  $N \rightarrow \infty$  an  $N$ -Freddy becomes a Bayesian).

# Implication 1

Implication 1: An  $N$ -Freddy will exhibit the gambler's fallacy.

- ▶ If there is a known objective probability, then an  $N$ -Freddy will act as if he believes a recent draw of one outcome increases the odds of the next draw being a different outcome (nearly tautological).

# Implication 1

Implication 1: An  $N$ -Freddy will exhibit the gambler's fallacy.

- If there is a known objective probability, then an  $N$ -Freddy will act as if he believes a recent draw of one outcome increases the odds of the next draw being a different outcome (nearly tautological).

## Implication 2

Implication 2: An  $N$ -Freddy will exhibit over-inference from small samples.

Recall Example C: Suppose there are two dice, a red one and a blue one. On the red die, there are 4  $H$ 's and 2  $M$ 's, whereas on the blue die, there are 2  $H$ 's and 4  $M$ 's. I flip a fair coin, and if it comes up heads I use the red die, and if it comes up tails I use the blue die (and you cannot observe the coin flip). I then roll that die twice. If the die comes up  $HH$ , what is the likelihood that I'm rolling the red die?

For a Bayesian:  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$ .

$\gamma(HH|\text{red}) = 4/9$  and  $\gamma(HH|\text{blue}) = 1/9$ .

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 4/5.$$

## Implication 2

Implication 2: An  $N$ -Freddy will exhibit over-inference from small samples.

Recall Example C: Suppose there are two dice, a red one and a blue one. On the red die, there are 4  $H$ 's and 2  $M$ 's, whereas on the blue die, there are 2  $H$ 's and 4  $M$ 's. I flip a fair coin, and if it comes up heads I use the red die, and if it comes up tails I use the blue die (and you cannot observe the coin flip). I then roll that die twice. If the die comes up  $HH$ , what is the likelihood that I'm rolling the red die?

For a Bayesian:  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$ .

$\gamma(HH|\text{red}) = 4/9$  and  $\gamma(HH|\text{blue}) = 1/9$ .

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 4/5.$$



## Implication 2

Implication 2: An  $N$ -Freddy will exhibit over-inference from small samples.

Recall Example C: Suppose there are two dice, a red one and a blue one. On the red die, there are 4  $H$ 's and 2  $M$ 's, whereas on the blue die, there are 2  $H$ 's and 4  $M$ 's. I flip a fair coin, and if it comes up heads I use the red die, and if it comes up tails I use the blue die (and you cannot observe the coin flip). I then roll that die twice. If the die comes up  $HH$ , what is the likelihood that I'm rolling the red die?

For a Bayesian:  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$ .

$$\gamma(HH|\text{red}) = 4/9 \text{ and } \gamma(HH|\text{blue}) = 1/9.$$

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 4/5.$$

## Implication 2

Implication 2: An  $N$ -Freddy will exhibit over-inference from small samples.

Recall Example C: Suppose there are two dice, a red one and a blue one. On the red die, there are 4  $H$ 's and 2  $M$ 's, whereas on the blue die, there are 2  $H$ 's and 4  $M$ 's. I flip a fair coin, and if it comes up heads I use the red die, and if it comes up tails I use the blue die (and you cannot observe the coin flip). I then roll that die twice. If the die comes up  $HH$ , what is the likelihood that I'm rolling the red die?

For a Bayesian:  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$ .

$$\gamma(HH|\text{red}) = 4/9 \text{ and } \gamma(HH|\text{blue}) = 1/9.$$

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 4/5.$$

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).

## Implication 2

For an ( $N = 12$ )-Freddy:

- ▶  $q_0(\text{red}) = q_0(\text{blue}) = 1/2$  (no change).
- ▶  $\gamma(HH|\text{red}) = 2/3 * 7/11 = 14/33$
- ▶  $\gamma(HH|\text{blue}) = 1/3 * 3/11 = 1/11$
- ▶ Now apply Bayes' Rule:

$$q_1(\text{red}|HH) = \frac{\gamma(HH|\text{red})q_0(\text{red})}{\gamma(HH|\text{red})q_0(\text{red}) + \gamma(HH|\text{blue})q_0(\text{blue})} = 14/17$$

Intuition: The law of small numbers makes Freddy think that  $HH$  is less likely to occur than it actually is; but since this reduction is larger for the die less likely to roll  $HH$  (i.e., for the blue die), Freddy will update too much toward the die more likely to roll  $HH$  (i.e., toward the red die).



## Implication 3

Implication 3: An  $N$ -Freddy will exhibit the hot-hand fallacy.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.

# Implication 3

## A Simplified Example:

- ▶ Suppose that in each game a player takes two shots. In reality, the player's hit rate is always 50%.

Consider an ( $N = 8$ )-Freddy who entertains the notion that the player might have hot and cold games. Specifically, for any given game, he believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Initially, Freddy doesn't know  $q$ , but Freddy attempts to infer  $q$  from watching the player's performance across a large number of games. What will Freddy infer?

Note: Let's assume that Freddy's processing reflects that there is a new urn at the start of every game.



## Implication 3

What does Freddy actually see?

Given that the true hit rate is always 50%, after observing a “large” sample, Freddy will see pairs in the following proportions:

- ▶  $(H, H) = 1/4$
- ▶  $(M, M) = 1/4$
- ▶  $(H, M) \text{ or } (M, H) = 1/2$

## Implication 3

What does Freddy actually see?

Given that the true hit rate is always 50%, after observing a “large” sample, Freddy will see pairs in the following proportions:

- ▶  $(H, H) = 1/4$
- ▶  $(M, M) = 1/4$
- ▶  $(H, M) \text{ or } (M, H) = 1/2$

## Implication 3

What does Freddy actually see?

Given that the true hit rate is always 50%, after observing a “large” sample, Freddy will see pairs in the following proportions:

- ▶  $(H, H) = 1/4$
- ▶  $(M, M) = 1/4$
- ▶  $(H, M) \text{ or } (M, H) = 1/2$

## Implication 3

What does Freddy actually see?

Given that the true hit rate is always 50%, after observing a “large” sample, Freddy will see pairs in the following proportions:

- ▶  $(H, H) = 1/4$
- ▶  $(M, M) = 1/4$
- ▶  $(H, M) \text{ or } (M, H) = 1/2$

## Implication 3

What does Freddy actually see?

Given that the true hit rate is always 50%, after observing a “large” sample, Freddy will see pairs in the following proportions:

- ▶  $(H, H) = 1/4$
- ▶  $(M, M) = 1/4$
- ▶  $(H, M) \text{ or } (M, H) = 1/2$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$



# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M)$  or  $(M, H) = 8/14 - 2q/7$

## Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M)$  or  $(M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy expect to see?

Recall that Freddy believes:

- ▶ With probability  $q$ , the player's hit rate is 75%.
- ▶ With probability  $q$ , the player's hit rate is 25%.
- ▶ With probability  $1 - 2q$ , the player's hit rate is 50%.

Hence, as a function of  $q$ , our  $(N = 8)$ -Freddy expects to see pairs in the following proportions:

- ▶  $(H, H) = 3/14 + q/7$
- ▶  $(M, M) = 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) = 8/14 - 2q/7$

# Implication 3

What does Freddy infer about  $q$ ?

Recall: Freddy actually sees:

- ▶  $(H, H) \rightarrow 1/4$
- ▶  $(M, M) \rightarrow 1/4$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 1/2$

Recall: Freddy expects to see:

- ▶  $(H, H) \rightarrow 3/14 + q/7$
- ▶  $(M, M) \rightarrow 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 8/14 - 2q/7$

Hence, an  $N = 8$  Freddy's estimate  $q$  will be the  $q$  such that ...



# Implication 3

What does Freddy infer about  $q$ ?

Recall: Freddy actually sees:

- ▶  $(H, H) \rightarrow 1/4$
- ▶  $(M, M) \rightarrow 1/4$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 1/2$

Recall: Freddy expects to see:

- ▶  $(H, H) \rightarrow 3/14 + q/7$
- ▶  $(M, M) \rightarrow 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 8/14 - 2q/7$

Hence, an  $N = 8$  Freddy's estimate  $q$  will be the  $q$  such that ...

# Implication 3

What does Freddy infer about  $q$ ?

Recall: Freddy actually sees:

- ▶  $(H, H) \rightarrow 1/4$
- ▶  $(M, M) \rightarrow 1/4$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 1/2$

Recall: Freddy expects to see:

- ▶  $(H, H) \rightarrow 3/14 + q/7$
- ▶  $(M, M) \rightarrow 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 8/14 - 2q/7$

Hence, an  $N = 8$  Freddy's estimate  $q$  will be the  $q$  such that ...

# Implication 3

What does Freddy infer about  $q$ ?

Recall: Freddy actually sees:

- ▶  $(H, H) \rightarrow 1/4$
- ▶  $(M, M) \rightarrow 1/4$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 1/2$

Recall: Freddy expects to see:

- ▶  $(H, H) \rightarrow 3/14 + q/7$
- ▶  $(M, M) \rightarrow 3/14 + q/7$
- ▶  $(H, M) \text{ or } (M, H) \rightarrow 8/14 - 2q/7$

Hence, an  $N = 8$  Freddy's estimate  $q$  will be the  $q$  such that ...

## Implication 3

Even though the player does NOT have a hot hand, Freddy will come to believe that the player is “hot” in 25% of games, “cold” in 25% of games, and “average” in 50% of games.

Intuition: The law of small numbers makes Freddy think that a 50% shooter is less likely to have *HH* or *MM* games than the player actually is. Hence, when Freddy sees a lot of *HH* and *MM* games, Freddy is forced to conclude that the player must be having hot and cold games.

## Implication 3

Even though the player does NOT have a hot hand, Freddy will come to believe that the player is “hot” in 25% of games, “cold” in 25% of games, and “average” in 50% of games.

Intuition: The law of small numbers makes Freddy think that a 50% shooter is less likely to have *HH* or *MM* games than the player actually is. Hence, when Freddy sees a lot of *HH* and *MM* games, Freddy is forced to conclude that the player must be having hot and cold games.

# Class Conclusion Slide

I don't care if you struggled with the calculus in this course or if you found the algebra tedious. (So did I, when I was coming up with all this stuff)

I want you to take these lessons:

- ▶ People are dumb (duh)
- ▶ People are systematically dumb
- ▶ Formal models can help us understand the implications of their dumbness.
- ▶ Formal thinking can help us understand what we don't understand.

# Class Conclusion Slide

I don't care if you struggled with the calculus in this course or if you found the algebra tedious. (So did I, when I was coming up with all this stuff)

I want you to take these lessons:

- ▶ People are dumb (duh)
- ▶ People are systematically dumb
- ▶ Formal models can help us understand the implications of their dumbness.
- ▶ Formal thinking can help us understand what we don't understand.

# Class Conclusion Slide

I don't care if you struggled with the calculus in this course or if you found the algebra tedious. (So did I, when I was coming up with all this stuff)

I want you to take these lessons:

- ▶ People are dumb (duh)
- ▶ People are systematically dumb
- ▶ Formal models can help us understand the implications of their dumbness.
- ▶ Formal thinking can help us understand what we don't understand.



# Class Conclusion Slide

I don't care if you struggled with the calculus in this course or if you found the algebra tedious. (So did I, when I was coming up with all this stuff)

I want you to take these lessons:

- ▶ People are dumb (duh)
- ▶ People are systematically dumb
- ▶ Formal models can help us understand the implications of their dumbness.
- ▶ Formal thinking can help us understand what we don't understand.

# Class Conclusion Slide

I don't care if you struggled with the calculus in this course or if you found the algebra tedious. (So did I, when I was coming up with all this stuff)

I want you to take these lessons:

- ▶ People are dumb (duh)
- ▶ People are systematically dumb
- ▶ Formal models can help us understand the implications of their dumbness.
- ▶ Formal thinking can help us understand what we don't understand.