Fefferman Statement (B) on $\mathbb{T}^3$: A Precise Lean Formalization, a Kernel-Checked Reduction, and the Boundaries of the Frohmanian Symplectic Tether

Benjamin Stanley Frohman
ORCID: 0009-0006-7068-3718 | GitHub: BenFrohman/Clay-
Concept DOI (all versions): 10.5281/zenodo.23653012
Previously archived version DOI: 10.5281/zenodo.23940667

Earliest Commit Provenance: 2026-10-07 06:16:45 AM CDT (UTC-5)
Revised edition: November 6, 2026; smoothness-index correction November 9, 2026 (America/Chicago, UTC−6). Not peer reviewed.

Abstract

This monograph records, in Lean 4 against Mathlib, a precise formal statement of Fefferman's Statement (B) on global smoothness of the unforced incompressible Navier–Stokes equations on $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$, together with an exact ledger of the analytic inputs that the Frohmanian symplectic tether does *not* supply. Statement (B) is *not* proved here. What is proved is a reduction: Statement (B) follows from the classical local theory with the Beale–Kato–Majda continuation criterion and from the open hypothesis that the Beale–Kato–Majda integral is finite on every finite interval of existence. We also record the two-dimensional clock test: constant vortex direction ($\sin\varphi \equiv 0$) coexists with linear growth $\|\phi(t)\|_{L^\infty} = M_0^2\, t$ on steady Euler equilibria.

1 Scope and non-claim boundary

This repository is a boundary and gap ledger. It is **not** a resolution of the Clay Millennium Problem. Every open input is a named proposition in Lean; none is assumed as an axiom, and no proof uses `sorry`. The build and axiom audit (`lean/AxiomReport.lean`, output in `AXIOMS.md`) reports only `propext`, `Classical.choice` and `Quot.sound`. The active continuation of this programme, in which the remaining gap is decomposed into finer named statements (all open or conditional, apart from individually proved lemmas such as the planar `sinShear` obstruction; the open statement is the Frohmanian Tether Hypothesis (A)/(B), see `NAMING.md`), is BenFrohman/periodic-ns-classical-continuation.

- **Repository:** https://github.com/BenFrohman/Clay-
- **Concept DOI:** 10.5281/zenodo.23653012
- **Previously archived version DOI:** 10.5281/zenodo.23940667
- **Earliest commit:** `9e61806681bfce3f0edee28293d22a6a711f0780` (2026-10-07, 06:16:45 CDT, UTC−5)
- **License:** Copyright © 2026 Benjamin Stanley Frohman. Apache License, Version 2.0.

2 Statement (B)

Following the official problem description of Fefferman [1], fix $\nu > 0$ and let u° be a smooth, divergence-free vector field on $\mathbb{R}^3$ with $u^\circ(x + e_j) = u^\circ(x)$ for $j = 1, 2, 3$. Take $f \equiv 0$.

> **Statement (B).** There exist smooth functions $p(x,t)$, $u_i(x,t)$ on $\mathbb{R}^3 \times [0,\infty)$ such that
>
> $$\partial_t u_i + \sum_{j=1}^{3} u_j\, \partial_j u_i = \nu\, \Delta u_i - \partial_i p, \qquad \operatorname{div} u = 0, \qquad u(x,0) = u^\circ(x),$$
>
> and u, p are 1-periodic in each spatial coordinate.

In Lean this is the proposition `Clay.StatementB.FeffermanStatementB` (file `lean/Clay/StatementB.lean`). Space is $\mathbb{R}^3$ realised as Euclidean space on $\mathrm{Fin}\,3$; partial derivatives are Fréchet derivatives along the standard basis; the time derivative is taken within $[0,\infty)$; smoothness is C^∞ on $[0,\infty) \times \mathbb{R}^3$, written with the Mathlib index $\infty = ((\top : \mathbb{N}_\infty) : \texttt{WithTop}\ \mathbb{N}_\infty)$. No component of the statement is opaque or trivial.

Correction (November 9, 2026). Earlier versions of `lean/Clay/StatementB.lean` (up to commit `1849c26`), and earlier editions of this monograph, including those archived as Zenodo versions 1.1.0–1.3.0, described the formal smoothness as C^∞, but the Lean source wrote a bare $\top$ in `SmoothPeriodicData` and `NSOn`. At the pinned Mathlib $\top : \texttt{WithTop}\ \mathbb{N}_\infty$ is the index ω, which means real-analytic. Those earlier formal statements were therefore analytic-regularity statements, not Fefferman's C^∞ Statement (B); by an informal shear-flow argument (spatial-bridge paper, Remark 2.1) that earlier analytic version is false if classical uniqueness of smooth solutions is granted. The source now uses ∞ in `SmoothPeriodicData`, `NSOn`, `reversedHolderClaim` and `spikeObstruction`. The full library rebuilds with the same proofs (one proof needed the bound $2 \le \infty$ in place of $2 \le \omega$) and the same axioms; `lean/Clay/Notes/SmoothnessIndex.lean` kernel-checks that the indices differ and that a C^∞ function on $\mathbb{R}$ need not be analytic.

3 The proved reduction

Write $\omega = \operatorname{curl} u$. Consider the propositions

- `PeriodicMaximalSolutionWithBKM` (classical; Kato, Fujita–Kato, Beale–Kato–Majda [2]; not yet formalized): for all admissible data there are $T^* \in (0,\infty]$ and a smooth solution on $[0,T^*)$ such that $T^* < \infty$ implies $\int_0^{T^*} \|\omega(t)\|_{L^\infty}\, dt = \infty$;
- `PeriodicBKMIntegrability` (**open**; the Clay core): every smooth solution on a finite interval $[0,T)$ satisfies $\int_0^{T} \|\omega(t)\|_{L^\infty}\, dt < \infty$.

Theorem 1 (`feffermanStatementB_of_bkm`)**.** `PeriodicMaximalSolutionWithBKM` *and* `PeriodicBKMIntegrability` *together imply Statement (B).* [Proved as an implication; Conditional on both hypotheses]

```
theorem feffermanStatementB_of_bkm (hloc : PeriodicMaximalSolutionWithBKM)
    (hbkm : PeriodicBKMIntegrability) : FeffermanStatementB
```

Theorem 2 (`bkmIntegralFinite_of_holographic`). *If* $\|\omega(t)\|_{L^\infty}$ *is bounded on every finite interval of existence, then* `PeriodicBKMIntegrability` *holds.* [Proved as an implication; the bound itself is open]

The second hypothesis of the first theorem is open and is equivalent in strength to Statement (B) itself. Consequently the reduction does not prove Statement (B).

4 Ledger of missing inputs

The file `lean/Clay/Gaps/MissingInputs.lean` states each former placeholder as a precise proposition:

- `metriplecticControlsEnstrophyL2`: $\sup_{[0,T)} \int_{\mathbb{T}^3} |\omega|^2 < \infty$ for every solution on a finite interval (open);
- `czGivesBMONotLinf`: no constant C gives $\|\nabla v\|_{L^\infty} \le C\|\operatorname{curl} v\|_{L^\infty}$ for smooth periodic divergence-free v (classical, not formalized);
- `spikeObstruction`: no constant C gives $\|f\|_{L^\infty} \le C\|f\|_{L^6(\mathbb{T}^3)}$ for smooth periodic f (classical, not formalized);
- `holographicNotAZFCBound`: the uniform ceiling $\sup_{[0,T)} \|\omega\|_{L^\infty} < \infty$ (open);
- `depletedStretching`: $\omega \cdot S\omega \le C|\omega|^2$ uniformly on finite intervals (open).

The former trivial definitions have been removed; the repository history retains them. For planar fields $v = (v_1(x_1, x_2), v_2(x_1, x_2), 0)$ the vorticity is proved parallel to e_3. Kernel-checked elementary results include the identity $\Pi_u u = 0$ for the pointwise tether projection, the unboundedness of the history integral $m^2 t$ for $m > 0$, and the conditional Riccati rearrangement.

5 Build and axiom status

Proved (kernel-checked). After the smoothness-index correction a full `lake build` of the Clay library, the regression file and the axiom report pass on Lean v4.16.0 against the pinned Mathlib build cache (Mathlib itself is not rebuilt from source). The full-library axiom report in `AXIOMS.md` lists 133 theorem declarations (including generated auxiliaries), 244 audited declarations, and no nonstandard axiom; before the correction it listed 125 and 236, and those 125 rows are unchanged. The substantive kernel-checked results are `feffermanStatementB_of_bkm`, `statementB_of_bkmIntegralFinite`, `bkmIntegralFinite_of_holographic`, `bkmIntegralFinite_of_vortSup_le`, the two-dimensional clock results `steady_euler_clock`, `alignment_does_not_delete_clock`, `clock_fails_on_aligned_2d_solutions`, the planar-vorticity lemma `curl_planar`, the tether identities `tether_degeneracy`, `tether_bracket_preservation`, and the integer rearrangement `conditional_riccati`. This is a machine check of the Lean statements, not peer review.

Conditional on. The global-existence reduction takes `PeriodicMaximalSolutionWithBKM` (classical local theory and continuation, not proved here) and `PeriodicBKMIntegrability` (the still-undischarged BKM integrability assumption) as arguments. Neither is supplied by the reduction.

The separate declaration `conditional_riccati` is an algebraic implication over integers. For $\text{rate}, C, \kappa, M \in \mathbb{Z}$, `CorrectedMaxRate` is exactly

$$\text{rate} \le M(CM - \kappa M^2),$$

and the theorem rewrites its right-hand side as $CM^2 - \kappa M^3$. Here `rate` is an integer variable, not a time derivative. No real-valued trajectory, differential inequality, vorticity maximum principle, or connection of this predicate to Navier–Stokes is proved. An analytical max-rate law and its PDE bridge remain separate missing inputs; this rearrangement does not supply them. Finer decompositions of the open BKM input belong to the continuation repository and are not formalized here.

Open. `PeriodicBKMIntegrability`; every item of the ledger above marked open; and therefore Statement (B) itself. The Clay Millennium Problem is not solved by this work.

An abstract criterion is not an additional analytic input. The new semantic regression `abstract_dhy_criterion_iff_periodicBKMIntegrability` proves that the current `dhyImpliesRegularity` proposition is equivalent to `PeriodicBKMIntegrability`: its existential packet has no connection to the solution and can have all four functions identically zero. This is a diagnosis of the abstract interface, not a proof of a Deng–Hou–Yu geometric criterion or of BKM integrability. An actual geometric criterion must bind its packet to vortex lines of the solution and justify the required quantitative estimates.

A solution-linked, nondegenerate vortex-line implementation

The module `Clay.Depletion.SolutionLinkedVortexLines` now supplies a `PacketuT` tied to the actual `curl(ut)`. It records moving endpoints, parameterized curves joining them, actual normalized-vorticity tangents, nonvanishing along active segments, measurable observed endpoint magnitudes, and an actual bound on the integral of the direction divergence. At nonzero peak vorticity this implementation additionally requires positive segment length: it is an explicitly nondegenerate specialization of the spatial setting of Deng–Hou–Yu Theorem 1 [4], not a transcription of every degenerate case. The paper's moving-segment Theorem 2 is for Euler and is not transferred to viscous Navier–Stokes here.

Proved. The tangent has norm one wherever vorticity is nonzero. A zero-length packet forces the actual vorticity to vanish everywhere at that time; any nonzero actual vorticity forces positive packet length. The actual zero solution satisfies `NSOn`, admits a packet, and passes the endpoint and BKM regressions. The actual spatial comparison is now derived from existing NSOn smoothness and the packet. Given separately the finite observed endpoint integral, the new theorem proves

$$\Omega(t) \le K|\omega(t,x(t))| \le Ke^{C}|\omega(t,y(t))|, \qquad \int_0^T \Omega(t)\,dt \le Ke^{C}\int_0^T |\omega(t,y(t))|\,dt < \infty.$$

This is a conditional geometric-to-BKM criterion, not a proof of global regularity. Neither endpoint-integral finiteness nor BKM finiteness is hidden in the packet fields.

Spatial bridge proved. The new module `Clay.Depletion.ActualSpatialBridge` supplies `spatialBridgeObligation_proved`, an inhabitant of the exact original `SpatialBridgeObligation`. Spatial C2 regularity is derived by restricting NSOn's existing joint smoothness to a time slice. Symmetry of actual mixed partials gives $\operatorname{div}(\operatorname{curl} u) = 0$; product and chain rules give the log-vorticity derivative along the actual unit tangent. FTC and exponentiation then establish the endpoint comparison, without assuming that comparison or adding a calculus regularity hypothesis.

Still unproved. Constructing suitable measurable packets with uniform geometric bounds and proving their actual endpoint time integrals finite for arbitrary solutions remain open. The theorem `bkmIntegralFinite_of_NSOn_and_endpoint` retains this last integral as an explicit argument. Thus Statement (B) is not proved. The old free-packet equivalence remains a diagnostic, not the interpretation of the linked data.

6 Provenance hashes

Source files are fingerprinted with SHA-256 (`provenance/PROVENANCE_HASHES.json`) and a Merkle root (`provenance/PROVENANCE_MERKLE_ROOT.json`), signed with Ed25519 key fingerprint `b428fb72676f1c52`. These are file-integrity fingerprints of an earlier snapshot only; they are not a certificate of correctness or review. The authoritative correctness check is the build and axiom report above, rerun by CI on every push.

References

[1] C. L. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, in: The Millennium Prize Problems, Clay Mathematics Institute, 2006, pp. 57–67.

[2] J. T. Beale, T. Kato, A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Comm. Math. Phys. **94** (1984), 61–66.

[3] P. Constantin, C. Fefferman, *Direction of vorticity and the problem of global regularity for the Navier–Stokes equations*, Indiana Univ. Math. J. **42** (1993), 775–789.

[4] J. Deng, T. Y. Hou, X. Yu, *Geometric Properties and Non-blowup of 3-D Incompressible Euler Flow*, https://arxiv.org/abs/math-ph/0402032, Theorem 1 and Lemma 1; Theorem 2 concerns Euler flow.

Attribution and licensing. Copyright © 2026 Benjamin Stanley Frohman (ORCID 0009-0006-7068-3718). Licensed under the Apache License, Version 2.0.