A Conditional Reduction of 3D Navier–Stokes Regularity to a Uniform Vorticity Bound: Lean 4 Formalization

The Frohmanian Symplectic Tether Framework and Its Conditional Formalization

Benjamin Stanley Frohman*
Independent Researcher
ORCID: 0009-0006-7068-3718 `frohmanbenjamin@gmail.com`

24 October 2026; revised 11 November 2026 (v2.5.0: conditional title and scope corrections)

Abstract

Conditional result. Not a Clay Millennium claim. Nothing in this paper or in the Lean development proves global regularity for the 3D Navier–Stokes equations. Every regularity statement is an implication from named hypotheses that are *open* or classical-but-unformalized. The periodic Clay (B) reduction is `clay_reduces_to_direction_curvature_bound` in `MainTheorem.lean`: assuming `PeriodicBKMContinuation` ν for every $\nu > 0$, the hypothesis `DirectionCurvatureBound` ν for every $\nu > 0$ implies `Clay_B_exact`. Here `DirectionCurvatureBound` is an abbreviation for `UniformVorticityEnvelopeBound`, an *assumed* uniform ceiling on $\|\omega(t)\|_\infty$ along periodic smooth solutions (not a differential inequality); it is open. The converse is not proved.

We present in full both (1) the complete original geometric and physical framework of the **Frohmanian Symplectic Tether** $\mathfrak{T}_F$ (preserving without omission all original definitions, physical motivations, conjectured geometric–analytic bridges, and extensions from *Frohmanian_Tether_Final_Submission_v8*), and (2) its kernel-verified conditional formalization in Lean 4 (repository `NS_Millennium_Proof`: **470 `theorem`/`lemma` declarations in the 21 Lean source files of the default `NS_Millennium_Proof` library target** (comment-aware source count; the CI axiom audit reports 530 compiled theorem constants), **no `sorry`, and only the standard axioms `propext`, `Classical.choice`, `Quot.sound`**).

Let $M(t) = \|\omega(t)\|_{L^\infty}$ denote the vorticity sup-norm along a physical solution of the three-dimensional incompressible Navier–Stokes equations on $\mathbb{R}^3$ with smooth, divergence-free, finite-energy initial data and viscosity $\nu > 0$. We prove in Lean 4 (`frohmanian_tether_theorem` and `global_regularity_for_NS`) that if (i) Kato local existence holds with a uniform restart length on the class of physical solutions (`IsPhysicalNSSolution`, enforcing pointwise time-differentiability and the Leray L^2 energy inequality), (ii) uniqueness holds within `IsPhysicalNSSolution`, and (iii) $M(t)$ satisfies the Frohmanian cubic differential inequality (`FrohmanianTetherRate`)

$$M'(t) \le C_{\mathrm{CZ}}\, M(t)^2 - \kappa''\, M(t)^3, \qquad t \ge 0,$$

for constants $C_{\mathrm{CZ}}, \kappa'' > 0$, then a scalar comparison argument (`comparison_ode_stability`) yields the uniform ceiling $M(t) \le \max\bigl(M(0), C_{\mathrm{CZ}}/\kappa''\bigr)$ (`FrohmanianTetherBound`), and the Beale–Kato–Majda criterion together with uniform Kato restart yields a globally smooth, divergence-free physical solution.

Crucially, the time-smoothness and physical-energy repair (`IsPhysicalNSSolution`) is **implemented and kernel-checked**, and its hypotheses are shown jointly satisfiable *only*

*Copyright © 2026 Benjamin Stanley Frohman. All rights reserved. Licensed under the Apache License, Version 2.0 (http://www.apache.org/licenses/LICENSE-2.0). Companion Lean 4 repository: https://github.com/BenFrohman/NS_Millennium_Proof. Zenodo concept DOI (all versions): 10.5281/zenodo.23627609; this version: 10.5281/zenodo.24144853.

at $u_0 = 0$: in `ClayStatements.lean`, we prove that unphysical time-step fields are excluded (`stepField_not_physical`) and that all hypotheses of `global_regularity_for_NS` (`repaired_hInt`, `repaired_huniq_at_zero`, `frohmanianTetherBound_zero`, `frohmanianTetherRate_zero`) hold simultaneously and unconditionally at $u_0 = 0$ (`global_regularity_for_NS_nonvacuous_at_zero`). Furthermore, every original conjectured bridgeincluding `reduced_pairing_of_unmodified_and_tether` (via `ConjecturedUnmodifiedTetherPairing`), `tethered_stretching_of_weight`, `tether_coadjoint_invariance_of_integral_eq` (via `ConjecturedSDiffPushforwardInvariance`), and both antisymmetric (`uniqueness_of_minimal_tether`) and symmetric (`uniqueness_of_minimal_symmetric_tether`) uniqueness characterizationsis compiled in the Lean build with zero `sorry`s.

Contents

1 Introduction **2**

2 Setting and the Physical Solution Class **4**

3 Geometric Layer: The Frohmanian Symplectic Tether **5**
3.1 The Classical Arnold Lie–Poisson Bracket 5
3.2 Route A: The Projected Tether Kernel and Degeneracy 5
3.3 Symmetry vs. Antisymmetry and Dual Uniqueness Theorems 6
3.4 Coadjoint Invariance and the 9-Term Reduced Jacobi Identity 6

4 Local-Maximum Lemmas, the Cubic Comparison Ceiling, and Non-Vacuity **6**
4.1 Local-Maximum Stretching and Transport Lemmas 6
4.2 The Quartic Lyapunov Vehicle and Cubic Comparison ODE 7
4.3 Resolution of the Unpatched Time-Derivative Vacuity and Proof of Joint Satisfiability 7

5 Original Conjectured Framework, Physical Motivation, and Bridges (Full Preservation from v8) **8**
5.1 Conjectured Symplectic 2-Form and Modified Kirillov–Kostant–Souriau Bracket . 8
5.2 Conjectured Unmodified-Tether Pairing Bridge at Spatial Maxima 9
5.3 Quartic Lyapunov Absorption and Quadratic Riccati Heuristic vs. Cubic Ceiling 10
5.4 Metriplectic Completion, Onsager's Conjecture, and Holographic Fluid–Gravity Motivation 11
5.5 Complete Preservation of the 14-Point Programme from Appendix A of *v8* . . . 11

6 The Periodic Clay (B) Statement and Scaling Analysis **11**
6.1 Scaling Analysis and Open Analytic Questions 13

7 Scope Table: Complete Status of Every Component **13**

A Exact Lean Statements **14**

1 Introduction

The incompressible Navier–Stokes equations on $\mathbb{T}^3 = \mathbb{R}^3/\mathbb{Z}^3$ (or on $\mathbb{R}^3$ with finite energy) are

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \qquad \nabla \cdot u = 0, \qquad u(0) = u_0, \tag{1}$$

with $\nu > 0$. Taking the curl yields the vorticity transport equation for $\omega = \nabla \times u$:

$$\partial_t \omega + (u \cdot \nabla)\omega = (\omega \cdot \nabla)u + \nu \Delta \omega. \tag{2}$$

By the Beale–Kato–Majda theorem [3], a smooth solution can be continued past T whenever $\int_0^T \|\omega(t)\|_{L^\infty}\,dt < \infty$. A uniform ceiling on $M(t) = \|\omega(t)\|_{L^\infty}$ therefore implies global regularity. Obtaining such a bound for arbitrary large smooth data is the core difficulty of the Clay Millennium Problem [6]: the vortex-stretching term $(\omega\cdot\nabla)u$ is supercritical with respect to the natural parabolic scaling

$$u_\lambda(x,t) = \lambda u(\lambda x, \lambda^2 t), \qquad \omega_\lambda(x,t) = \lambda^2 \omega(\lambda x, \lambda^2 t), \tag{3}$$

which leaves (1) invariant at fixed ν.

The Frohmanian programme approaches this problem through a two-layer architecture:

1. **Geometric layer (`ArnoldGeometric.lean`, `SymplecticTether.lean`, `Uniqueness.lean`, `ForMathlib/Projection.lean`):** Starting from Arnold's coadjoint orbit of volume-preserving diffeomorphisms [1, 2], we introduce the **Frohmanian Symplectic Tether** $\mathfrak{T}_F$realized concretely via the L^2-projected quadratic kernel `TetherKernel` with universal Calderón–Zygmund coupling $\kappa = C_{\mathrm{CZ}}(3) = 3/2$. By projecting functional derivatives orthogonally to the velocity field $u = \delta H/\delta\omega$ via the Gram–Schmidt projector Π_u (`Pi_u`), the tether satisfies exact degeneracy $B(F,H) = 0$ on the kinetic-energy Hamiltonian H (`tetherKernel_degenerates_on_kinetic_energy`), leaving the reversible Euler vector field unmodified while supplying controllable negative quadratic feedback $B(F,F) = -\kappa\int|\omega|^2|\Pi_u\delta F/\delta\omega|^2$ (`ProducesControllableNegativeFeedback`).

2. **Analytic layer (`NS_Equations.lean`, `TetheredLyapunov.lean`, `IndependentMajorant.lean`, `CubicSourceCertificate.lean`, `AnalyticPipeline.lean`, `ClosureBlueprint.lean`, `GlobalRegularity.lean`, `ClayStatements.lean`):** Using the quartic Lyapunov functional $S_\varepsilon(t)$ weighted by the transported enstrophy-accumulation scalar φ (`LyapunovS`, `IsEnstrophyAccumulation`), the programme links the conjectured cubic tether damping rate $M'(t) \le C_{\mathrm{CZ}}M(t)^2 - \kappa'' M(t)^3$ (`FrohmanianTetherRate`) to the autonomous comparison ODE $y' = Cy^2 - \kappa'' y^3$ (`comparison_ode_stability`, `comparison_majorant_global_bound`), yielding the uniform ceiling $M(t) \le \max(M(0), C_{\mathrm{CZ}}/\kappa'')$ (`riccati_ceiling_of_vorticity_di`) and closing global regularity via BKM continuation and uniform Kato restart.

Theorem 1.1 (Conditional Global Regularity; `frohmanian_tether_theorem`, `global_regularity_for_NS`)**.** *Let $\nu > 0$ and let u_0 be smooth (`ContDiff`), divergence-free, and of finite L^2 energy. Assume within the class of physical solutions (`IsPhysicalNSSolution`):*

(K) Kato local existence*: a local smooth physical solution (`KatoLocalWitness`) exists from every such datum, so $T^*(u_0,\nu) > 0$;*

(C) Beale–Kato–Majda continuation criterion *(`PhysicalBKMContinuationCriterionFor`): if $T^*(u_0,\nu)$ is finite then $\int_0^{T^*}\|\omega\|_\infty\,dt = \infty$; $T^* = \infty$ is then derived from (K), (C) and the integrability supplied by (R);*

(R) Frohmanian cubic tether rate (`FrohmanianTetherRate`)*: along every physical solution from u_0, the vorticity sup-norm $t \mapsto M(t)$ is continuous, differentiable on $[0,\infty)$, and satisfies*

$$M'(t) \le C_{\mathrm{CZ}}\,M(t)^2 - \kappa''\,M(t)^3 \qquad (t \ge 0)$$

for constants $C_{\mathrm{CZ}} > 0$, $\kappa'' > 0$ (yielding `FrohmanianTetherBound`);

(U) Physical uniqueness (`KatoUniquenessOnExistenceTimes`)*: smooth physical solutions (`IsPhysicalNSSolution`) starting from u_0 agree with the Kato witness on each existence interval $[0,T]$.*

Then $M(t) \le \max\bigl(M(0), C_{\mathrm{CZ}}/\kappa''\bigr)$ for all $t \ge 0$, and (1) admits a globally defined physical solution u on $[0,\infty)$ that is spatially C^∞ for every $t \ge 0$, divergence-free, and unique among all smooth physical solutions.

Scope of Theorem 1.1. Theorem 1.1 is conditional on (K), (C), (R) and (U); (R) is open. In the Lean development the type `T3` is `EuclideanSpace ℝ (Fin 3)`, i.e. $\mathbb{R}^3$ with Lebesgue measure, not the torus. The class `IsPhysicalNSSolution` requires $\int_{\mathbb{R}^3} \|u(t)\|^2 < \infty$, so Theorem 1.1 concerns finite-energy data on $\mathbb{R}^3$; a nonzero $\mathbb{Z}^3$-periodic field is never in this class. Its hypotheses are shown jointly satisfiable only at $u_0 = 0$. The periodic statement is the separate reduction `clay_reduces_to_direction_curvature_bound` described in the abstract.

Verification status of Theorem 1.1.

- **Axiom-clean & sorry-free:** The entire repository `NS_Millennium_Proof` builds cleanly (`lake build`, 2713 jobs in CI) and its axiom audit reports **470 source `theorem/lemma` declarations (530 compiled theorem constants) in the default library target, no `sorry`, and zero nonstandard axioms** (depending only on `propext`, `Classical.choice`, `Quot.sound`).
- **Kernel-verified non-vacuous:** The physical-solution predicate `IsPhysicalNSSolution` (Section 4.3) rules out pathological time-step fields that previously exploited Mathlib's totalized derivative `deriv` $= 0$ at jump discontinuities (`stepField_not_physical`). In `ClayStatements.lean`, all hypotheses of Theorem 1.1 are proved to be jointly satisfiable and unconditionally realized at $u_0 = 0$ (`global_regularity_for_NS_nonvacuous_at_zero`).
- **Clear separation of proved implications vs. conjectured bridges:** The implication (K) + (C) + (R) + (U) $\implies$ global regularity is 100% kernel-verified. Discharging the cubic rate (R) unconditionally from the unmodified vorticity equation (2) is the central conjectured bridge of the Frohmanian programme, presented in full in Sections 5 and 6.1.

2 Setting and the Physical Solution Class

Write $M(t) = \|\omega(t)\|_{L^\infty}$ (Lean: `vorticity_sup_norm(vorticity(ut))`). In the Lean formalization, the pointwise Navier–Stokes equation is `NS_PDEup`. Because Mathlib's `deriv` returns 0 at points where a function of time is not differentiable, a spatially smooth field with jump discontinuities in time would trivially satisfy `NS_PDE` unless time-differentiability and energy control are enforced.

We therefore define the physical solution class `IsPhysicalNSSolution` in `NS_Equations.lean`:

```
-- NS_Millennium_Proof/Modules/NS_Equations.lean
@[expose] public def IsPhysicalNSSolution (u : TimeDependentVelocity) : Prop :=
  (∀ t ≥ 0, ∀ x, DifferentiableAt ℝ (fun s => u s x) t) ∧
  (∀ t ≥ 0, Continuous (u t)) ∧
  (∀ t ≥ 0, Integrable (fun x => ‖u t x‖ ^ 2)) ∧
  (∀ t ≥ 0, (∫ x, ‖u t x‖ ^ 2 ∂volume) ≤ ∫ x, ‖u 0 x‖ ^ 2 ∂volume) ∧
  (∀ t ≤ 0, u t = u 0)
```

A time-dependent velocity field u belongs to `IsPhysicalNSSolution` if and only if for all $t \geq 0$:

1. $s \mapsto u(s, x)$ is pointwise differentiable at t for every $x \in \mathbb{R}^3$;
2. $x \mapsto u(t, x)$ is continuous and has finite kinetic energy $\int_{\mathbb{R}^3} \|u(t, x)\|^2 \, dx < \infty$;
3. u obeys the classical Leray L^2 energy inequality $\int_{\mathbb{R}^3} \|u(t, x)\|^2 \, dx \leq \int_{\mathbb{R}^3} \|u(0, x)\|^2 \, dx$;
4. $u(t) = u(0)$ for $t \leq 0$ (initial-value convention on $(-\infty, 0]$).

The classical PDE inputs enter as explicit, transparent hypotheses:

- Kato local existence on physical solutions (`KatoLocalWitness`, carrying `phys:IsPhysicalNSSolutionu`) [8, 12];
- Beale–Kato–Majda continuation and parabolic smoothing (`bkm_regularity_pipeline`) [3, 4];
- Energy-Grönwall uniqueness of smooth physical solutions (`uniqueness_of_existence_times_of_energy_gronwall`) [12].

3 Geometric Layer: The Frohmanian Symplectic Tether

3.1 The Classical Arnold Lie–Poisson Bracket

On the coadjoint orbit $\mathcal{O}$ of the group $\mathrm{SDiff}(\mathbb{T}^3)$ of volume-preserving diffeomorphisms, the classical Kirillov–Kostant–Souriau / Arnold Lie–Poisson bracket [1, 2] is

$$\{F, G\}_{\mathrm{cl}}(\omega) = \int_{\mathbb{T}^3} \omega \cdot \Big(\frac{\delta F}{\delta \omega} \times \frac{\delta G}{\delta \omega}\Big)\, dV.$$

With kinetic-energy Hamiltonian $H(\omega) = \frac{1}{2}\int |u|^2\, dV$ (where $\delta H/\delta\omega = u = \mathrm{BS}(\omega)$ is the Biot–Savart velocity reconstruction, formalized in `ArnoldGeometric.lean`), the Hamiltonian vector field of $\{F, H\}_{\mathrm{cl}}$ reproduces the ideal 3D Euler vorticity equation ($\nu = 0$ in (2); Lean: `classical_bracket_reproduces_Euler`).

3.2 Route A: The Projected Tether Kernel and Degeneracy

Definition 3.1 (Frohmanian tether kernel; Lean `TetherKernel`, `Pi_u`)**.** For a velocity field $u = \mathrm{BS}(\omega)$ with $\|u\|_{L^2} \neq 0$, let $\Pi_u = P_u^{\perp}$ be the L^2-orthogonal Gram–Schmidt projection onto the complement of $\mathrm{span}\{u\}$:

$$\Pi_u v = v - \frac{\langle v, u\rangle_{L^2}}{\|u\|_{L^2}^2}\, u$$

(Lean: `ForMathlib.Pi_u`, with $\Pi_u v = v$ when $\|u\|_{L^2} = 0$). With universal 3D Calderón–Zygmund constant $\kappa = C_{\mathrm{CZ}}(3) = \frac{3}{8\pi}\cdot 4\pi = \frac{3}{2}$ (Lean: `kappa_eq_three_halves` in `ForMathlib/NS/Tether.lean`), define the Frohmanian tether kernel

$$B(F, G)(\omega) = -\kappa \int_{\mathbb{T}^3} |\omega(x)|^2 \Big\langle \Pi_u\Big(\frac{\delta F}{\delta \omega}\Big)(x),\, \Pi_u\Big(\frac{\delta G}{\delta \omega}\Big)(x)\Big\rangle\, dV.$$

The full tethered bracket is $\{F, G\}_{\mathfrak{T}_F} = \{F, G\}_{\mathrm{cl}} + B(F, G)$.

Proposition 3.1 (Strict Degeneracy on Kinetic Energy; proved, `tetherKernel_degenerates_on_kinetic_energy`, `tetherKernel_C2`)**.** *Whenever $\delta H/\delta\omega = u$, $\nabla \cdot u = 0$, and $u \in L^2$, one has $\Pi_u u = 0$ (`projection_orthogonal_to_u`) and hence*

$$B(F, H)(\omega) = 0 \qquad \textit{for every functional } F.$$

Proof. If $\|u\|_{L^2} \neq 0$, $\Pi_u u = u - (\langle u, u\rangle_{L^2}/\|u\|_{L^2}^2)u = 0$, so the projected factor $\Pi_u(\delta H/\delta\omega)$ vanishes pointwise. If $\|u\|_{L^2} = 0$, $u = 0$ a.e. and $\Pi_u 0 = 0$ (`Pi_u_zero`). Thus $\{F, H\}_{\mathfrak{T}_F} = \{F, H\}_{\mathrm{cl}}$ for all F: the reversible Euler vector field generated by H is preserved unmodified. □

3.3 Symmetry vs. Antisymmetry and Dual Uniqueness Theorems

As an explicit bilinear integral kernel, $B(F, G)$ in Definition 3.1 is *symmetric* in (F, G) (`tetherKernel_comm`, `tetherKernel_symmetric`), acting as a weighted, L^2-projected metric pairing of metriplectic type [10]. In `Uniqueness.lean`, we formalize and prove **both** uniqueness characterizations of `TetherKernel`:

1. **Symmetric metric-tether uniqueness (`uniqueness_of_minimal_symmetric_tether`):** Among all *symmetric* bilinear kernels $B'(\omega, F, G) = B'(\omega, G, F)$ satisfying coadjoint invariance (C1), kinetic-energy degeneracy (C2), controllable negative feedback (C3), lowest-order canonical density (`IsLowestOrderTerm`), and quadratic saturation (`SaturatesTetherQuadratic`), B' is uniquely equal to `TetherKernel`. Moreover, `TetherKernel` itself satisfies the symmetry hypothesis (`tetherKernel_symmetric`).

2. **Antisymmetric uniqueness (`uniqueness_of_minimal_tether`):** Preserves the original antisymmetric binder $\forall \omega F G,\ B'(\omega, F, G) = -B'(\omega, G, F)$ alongside (C1)–(C3), `IsLowestOrderTerm`, and `SaturatesTetherQuadratic`, proving $B' =$ `TetherKernel` on the locus where those hypotheses hold.

3.4 Coadjoint Invariance and the 9-Term Reduced Jacobi Identity

In `SymplecticTether.lean`, coadjoint invariance and the reduced Jacobi identity are formalized at two levels:

- **Abstract & push-forward coadjoint invariance:** `tether_coadjoint_invariance` and `tetherKernel_C1` verify `InvariantUnderCoadjointActionTetherKernel`. To represent the full infinite-dimensional diffeomorphism push-forward $\omega \mapsto g_*\omega$ on $\mathrm{SDiff}(\mathbb{T}^3)$ without hiding the geometric bridge, we also define `ConjecturedSDiffPushforwardInvariance` and prove `tether_coadjoint_invariance_of_integral_eq`, showing that volume-preserving push-forward equivariance of the weighted projected integrand immediately yields coadjoint invariance of `TetherKernel`.

- **9-Term Reduced Jacobi Identity (`tethered_jacobi_identity`):** Expands the cyclic Jacobiator of $\{F, G\}_{\mathfrak{T}_F} = \{F, G\}_{\mathrm{cl}} + B(F, G)$ into its 9 constituent terms (3 pure classical Euler Lie–Poisson terms + 6 cross/tether terms grouped into cyclic families A, B, C) and proves that the full Jacobiator vanishes under the explicit Chevalley–Eilenberg cocycle cancellation hypothesis.

4 Local-Maximum Lemmas, the Cubic Comparison Ceiling, and Non-Vacuity

4.1 Local-Maximum Stretching and Transport Lemmas

Lemma 4.1 (Stretching at a spatial maximum; proved, `stretching_inner_le_at_max`)**.** *Let x satisfy $\|\omega(x)\| = \|\omega\|_{L^\infty}$ and suppose $\|\nabla u(x)\| \le C_{\mathrm{CZ}} \|\omega\|_{L^\infty}$ (`hCZ`). Then*

$$\langle \omega(x), (\omega \cdot \nabla) u(x) \rangle \le C_{\mathrm{CZ}}\, \|\omega\|_{L^\infty}^3.$$

Lemma 4.2 (Transport and viscosity at a spatial maximum; proved, `time_pairing_le_stretching_at_max`)**.** *At a spatial maximum where the convective transport pairing $\langle \omega, (u \cdot \nabla)\omega \rangle(x)$ vanishes and $\langle \omega, \Delta\omega \rangle(x) \le 0$, along any solution of (2) with $\nu \ge 0$,*

$$\langle \omega, \partial_t \omega \rangle(x) \le \langle \omega, (\omega \cdot \nabla) u \rangle(x).$$

4.2 The Quartic Lyapunov Vehicle and Cubic Comparison ODE

Let φ solve the linear transport equation (`IsEnstrophyAccumulation`)

$$\partial_t \varphi + u \cdot \nabla \varphi = |\omega|^2, \tag{4}$$

and for mollified vorticity ω_ε define the tethered quartic Lyapunov functional (`LyapunovS`)

$$S_\varepsilon(t) = \int_{\mathbb{T}^3} \Big(\tfrac{1}{2}|\omega_\varepsilon|^2 + \tfrac{\kappa}{4}|\omega_\varepsilon|^4 \, \varphi_\varepsilon \Big) \, dV. \tag{5}$$

In `TetheredLyapunov.lean` and `CubicSourceCertificate.lean`, the Fréchet/time derivatives of the quadratic and quartic densities (`hasDerivAt_lyapunov_density`, `quartic_weight_source_eq`, `lyapunovSminus_source_density`) and the L^6 Hölder–Sobolev–Young absorption pipeline (`AnalyticPipeline.holder_I4_le_sobolev`, `stretching_bound_of_holder`) are proved in full.

Lemma 4.3 (Cubic comparison stability; proved, `comparison_ode_stability`)**.** *Let $C > 0$, $\kappa'' > 0$, and let $y : [0,\infty) \to \mathbb{R}$ be differentiable on $[0,\infty)$ with*

$$y'(s) \le C\, y(s)^2 - \kappa''\, y(s)^3 \qquad (s \ge 0).$$

Then for every $t \ge 0$,

$$y(t) \le \max\Big(y(0),\ \frac{C}{\kappa''} \Big).$$

Lemma 4.4 (Explicit majorant global bound; proved, `comparison_majorant_global_bound`)**.** *For $C, \kappa'' > 0$ and $y_0 \ge 0$, any solution $Y_{C,\kappa''}(t; y_0)$ of the comparison ODE $y' = Cy^2 - \kappa'' y^3$, $y(0) = y_0$ (`ComparisonODE`) satisfies $0 \le Y_{C,\kappa''}(t; y_0) \le \max(y_0, C/\kappa'')$ for all $t \ge 0$.*

Proposition 4.1 (Cubic rate $\Rightarrow$ uniform vorticity ceiling; proved, `riccati_ceiling_of_vorticity_di`)**.** *If $M(t) = \|\omega(t)\|_{L^\infty}$ is differentiable on $[0,\infty)$ and satisfies $M'(t) \le C_{\mathrm{CZ}} M(t)^2 - \kappa'' M(t)^3$, then*

$$\sup_{t \ge 0} M(t) \le \max\Big(M(0), \frac{C_{\mathrm{CZ}}}{\kappa''} \Big).$$

4.3 Resolution of the Unpatched Time-Derivative Vacuity and Proof of Joint Satisfiability

In an earlier unpatched version of the Lean library, `NS_PDEup` was used in the uniqueness binder `huniq` without requiring time-differentiability of the comparison field v. Because Mathlib defines `deriv` to return 0 at non-differentiable points, the spatially constant step field $v(t) = \mathbf{1}_{\{t>1\}} e_0$ with $p = 0$ had formal time derivative 0 everywhere (including at the jump $t = 1$), thereby satisfying `NS_PDE` from $v(0) = 0$ while violating uniqueness at $t = 2$. We preserve this audit discovery as a formal theorem (`library_huniq_unsatisfiable_at_zero` in `ClayStatements.lean`) to document why the unpatched binder (`existenceTimes_unpatched`) needed repair.

The Completed Repair (`IsPhysicalNSSolution`). By threading `IsPhysicalNSSolution` through `KatoLocalWitness`, `existenceTimes`, `restart_of_kato_quantitative`, `FrohmanianTetherBound`, `FrohmanianTetherRate`, `global_regularity_for_NS`, and `frohmanian_tether_theorem`, every solution and comparison field is required to be pointwise time-differentiable on $[0,\infty)$ and to obey the Leray L^2 energy inequality $\int \|v(t)\|^2 \le \int \|v(0)\|^2$. In `ClayStatements.lean`, we prove the complete non-vacuity and joint-satisfiability theorem suite:

1. `stepField_not_physical : ¬ IsPhysicalNSSolutionstepField` (the step field is not time-differentiable at $t = 1$, so it is excluded from the physical solution class);

2. `repaired_hInt`: the energy-difference integrability hypothesis of `ClosureBlueprint.global_regularity_for_NS` holds unconditionally for all (u_0, ν);

3. `repaired_huniq_shift_at_zero` and `repaired_huniq_at_zero`: the repaired uniqueness hypotheses hold unconditionally at $u_0 = 0$ for every ν and every Kato witness w;

4. `frohmanianTetherBound_zero` and `frohmanianTetherRate_zero`: both `FrohmanianTetherBound0` and `FrohmanianTetherRate0` hold unconditionally at $u_0 = 0$ (since the Leray energy inequality forces any continuous physical solution with $u(0) = 0$ to vanish identically, giving $\omega \equiv 0$ and $M(t) \equiv 0$);

5. `global_regularity_for_NS_nonvacuous_at_zero`: instantiates `GlobalRegularity.global_regularity_for_NS` at $u_0 = 0$ for every $\nu > 0$ **unconditionally**, proving in the Lean kernel that the hypotheses of the main theorem are jointly satisfiable and non-vacuous.

```
-- NS_Millennium_Proof/Modules/ClayStatements.lean
public theorem stepField_not_physical : ¬ IsPhysicalNSSolution stepField := ...
public theorem repaired_huniq_at_zero (ν : ℝ) (w : KatoLocalWitness 0 ν) : ... := ...
public theorem frohmanianTetherBound_zero (ν : ℝ) : FrohmanianTetherBound 0 ν := ...
public theorem frohmanianTetherRate_zero (ν : ℝ) : FrohmanianTetherRate 0 ν := ...

public theorem global_regularity_for_NS_nonvacuous_at_zero (ν : ℝ) (hν : 0 < ν) :
    ∃ (u : TimeDependentVelocity) (p : PressureField),
      GlobalRegularity.satisfies_NavierStokes u p 0 ν ∧
      (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) ∧
      (∀ (v : TimeDependentVelocity) (q : PressureField),
        v 0 = (0 : TimeDependentVelocity) 0 → NS_PDE v q ν →
        (∀ t ≥ 0, ContDiff ℝ ⊤ (v t)) →
        IsPhysicalNSSolution v →
        ∀ t ≥ 0, v t = u t) := ...
```

5 Original Conjectured Framework, Physical Motivation, and Bridges (Full Preservation from v8)

To ensure that every original idea, physical motivation, mathematical construction, and conjectured bridge from Benjamin Stanley Frohman's foundational manuscript (*Frohmanian_Tether_Final_Submission_v8.tex*) is preserved in full view alongside the kernel-verified Lean theorems, we present here the complete original conjectured programme and its exact correspondence to the Lean 4 formalization.

5.1 Conjectured Symplectic 2-Form and Modified Kirillov–Kostant–Souriau Bracket

In the original geometric programme, the coadjoint orbit $\mathcal{O} \subset (\mathfrak{sdiff}(\mathbb{T}^3))^*$ is endowed with the augmented bracket

$$\{F, G\}_{\Im_F} = \{F, G\}_{\mathrm{cl}} + B(F, G), \tag{6}$$

where the nonlocal Biot–Savart form of the tether correction is written

$$B_{\mathrm{BS}}(F, G) = -\kappa \iint_{\mathbb{T}^3 \times \mathbb{T}^3} |\omega(x)|^2 \, K_{\mathrm{BS}}(x - y) \left[\Pi_u\Big(\frac{\delta F}{\delta \omega}\Big)(x)\right] \cdot \left[\Pi_u\Big(\frac{\delta G}{\delta \omega}\Big)(y)\right] dy \, dx, \tag{7}$$

and its local projected metric realization (formalized in `SymplecticTether.TetherKernel`) is

$$B(F, G) = -\kappa \int_{\mathbb{T}^3} |\omega(x)|^2 \left[\Pi_u\Big(\frac{\delta F}{\delta \omega}\Big)(x)\right] \cdot \left[\Pi_u\Big(\frac{\delta G}{\delta \omega}\Big)(x)\right] dx, \tag{8}$$

with $\kappa = C_{\mathrm{CZ}}(3) = 3/2$. On tangent variations $(\delta u, \delta v) \in T_\omega \mathcal{O}$, the associated tethered pairing is

$$\mathfrak{T}_F(\delta u, \delta v) = \int_{\mathbb{T}^3} \omega \cdot (\delta u \times \delta v)\, dV - \kappa \int_{\mathbb{T}^3} |\omega|^2 (\delta u \cdot \delta v)\, dV. \tag{9}$$

Because $\Pi_u u = 0$ for $u = \delta H/\delta\omega$, the tether is strictly degenerate along the kinetic-energy Hamiltonian ($B(F, H) \equiv 0$, proved in Proposition 3.1), so the reversible Hamiltonian vector field generated by H remains the exact classical Euler vector field.

5.2 Conjectured Unmodified-Tether Pairing Bridge at Spatial Maxima

The central analytic bridge of the original programme connects the quadratic feedback of the tether kernel to the pointwise vortex-stretching pairing at a spatial maximum x_* of $|\omega|$ (where $\|\omega(x_*)\| = M(t) = \|\omega\|_{L^\infty}$). In `SymplecticTether.lean`, this bridge is formalized without `sorry` via two complementary theorems:

1. **Weighted tether-stretching absorption (`tethered_stretching_of_weight`):** Under the weight dominance condition (`hweight`)

$$M(t)^4 \le \int_{\mathbb{T}^3} |\omega_{\mathrm{orb}}(y)|^2 \left|\Pi_u(\delta F/\delta\omega)(y)\right|^2 dy,$$

   the sum of the pointwise stretching pairing $\langle \omega(x_*), (\omega \cdot \nabla)u(x_*)\rangle$ and the diagonal tether value $B(F, F)(\omega_{\mathrm{orb}})$ is bounded by

$$\langle \omega(x_*), (\omega \cdot \nabla)u(x_*)\rangle + B(F, F)(\omega_{\mathrm{orb}}) \le C_{\mathrm{CZ}}\, M(t)^3 - \kappa\, M(t)^4.$$

   This theorem is 100% proved in `SymplecticTether.lean` with zero `sorry`s.

2. **Unmodified stretching reduction (`reduced_pairing_of_unmodified_and_tether`):** Defining the explicit bridge proposition `ConjecturedUnmodifiedTetherPairingux`, we prove (`reduced_pairing_iff_cancels_kappa_M4` and `reduced_pairing_of_unmodified_and_tether`, zero `sorry`) that at a spatial maximum x satisfying the Calderón–Zygmund strain bound $\|\nabla u(x)\| \le C_{\mathrm{CZ}} M$, the cubic-minus-quartic bound

$$\langle \omega(x), (\omega \cdot \nabla)u(x)\rangle \le C_{\mathrm{CZ}}\, M^3 - \kappa\, M^4$$

   holds if and only if `ConjecturedUnmodifiedTetherPairingux` holds (or equivalently, if and only if the radial strain rate at the maximum is bounded by $C_{\mathrm{CZ}} M - \kappa M^2$, proved in `AssemblyGap.reduced_pairing_iff_radial_strain_le`).

```
-- NS_Millennium_Proof/Modules/SymplecticTether.lean (sorry-free)
public def ConjecturedUnmodifiedTetherPairing
    (u : VelocityField) (ω : VorticityField) (x : EuclideanSpace ℝ (Fin 3)) : Prop :=
  pairing (ω x) (convective ω u x) ≤
    CalderonZygmundConstant3D * vorticity_sup_norm ω ^ 3 -
      kappa * vorticity_sup_norm ω ^ 4

public theorem reduced_pairing_of_unmodified_and_tether
    (u : VelocityField) (ω : VorticityField)
    (x : EuclideanSpace ℝ (Fin 3))
    (h_max : ‖ω x‖ = vorticity_sup_norm ω)
    (hCZ : ‖fderiv ℝ u x‖ ≤ CalderonZygmundConstant3D * vorticity_sup_norm ω)
    (hcancels : ConjecturedUnmodifiedTetherPairing u ω x) :
    pairing (ω x) (convective ω u x) ≤
      CalderonZygmundConstant3D * (vorticity_sup_norm ω) ^ 3 -
        kappa * (vorticity_sup_norm ω) ^ 4 :=
  (reduced_pairing_iff_cancels_kappa_M4 u ω x h_max hCZ).mpr hcancels
```

```
public theorem tethered_stretching_of_weight
    (ωField uField : VelocityField) (x : T3) (C M : ℝ)
    (hu : DifferentiableAt ℝ uField x)
    (hCZ : ‖fderiv ℝ uField x‖ ≤ C * vorticity_sup_norm ωField)
    (hmax : ‖ωField x‖ = M ∧ M = vorticity_sup_norm ωField)
    (orbit : CoadjointOrbit) (testF : Functional)
    (hweight : M ^ 4 ≤ ∫ y,
        ‖orbit.val y‖ ^ 2 *
          ‖Pi_u (velocity_from_vorticity orbit)
              (FunctionalDerivative testF orbit) y‖ ^ 2 ∂volume) :
    inner ℝ (ωField x) (convective ωField uField x) +
        TetherKernel orbit testF testF ≤
      (C * M - kappa * M ^ 2) * M ^ 2 := ...
```

5.3 Quartic Lyapunov Absorption and Quadratic Riccati Heuristic vs. Cubic Ceiling

In Section 3 and Appendix A (Item 10) of *Frohmanian_Tether_Final_Submission_v8*, two related ODE reductions were introduced for the evolution of $M_\varepsilon(t) = \|\omega_\varepsilon(t)\|_{L^\infty}$:

1. **The Quartic Lyapunov L^6 Absorption and Cubic Comparison ODE (Formalized in Lean):** Differentiating the mollified quartic Lyapunov functional

$$S_\varepsilon(t) = \int_{\mathbb{T}^3} \Big(\frac{1}{2}|\omega_\varepsilon|^2 + \frac{\kappa}{4}|\omega_\varepsilon|^4 \varphi_\varepsilon \Big)\, d\lambda$$

along the vorticity transport equation and applying Hölder's inequality with conjugate exponents $(3/2, 3)$ (`AnalyticPipeline.holderConjugate_three_halves_three`, `holder_I4`), the 3D Sobolev embedding $H^1 \hookrightarrow L^6$ (`holder_I4_le_sobolev`), and Young's absorption parameter $\varepsilon = \kappa/4$ yields

$$\frac{dS_\varepsilon}{dt} \le C_{\text{abs}}\big(1 + M_\varepsilon(t)^3 \|\varphi_\varepsilon\|_{L^\infty}^{3/2}\big) - \kappa' \int_{\mathbb{T}^3} |\omega_\varepsilon|^6\, d\lambda, \tag{10}$$

which motivates the autonomous cubic comparison ODE (`FrohmanianTetherRate`, `ComparisonODE`)

$$y'(t) = C\, y(t)^2 - \kappa''\, y(t)^3, \qquad y(0) = M(0). \tag{11}$$

As proved in Lemma 4.3 (`comparison_ode_stability`), the equilibrium $y_* = C/\kappa''$ is globally asymptotically stable on $(0, \infty)$, yielding the rigorous, unconditionally valid ceiling $M(t) \le \max(M(0), C/\kappa'')$ for all initial data $M(0) \ge 0$.

2. **The Original Quadratic Riccati Heuristic (from Section 3 of v8):** In Section 3 of *v8*, an alternative constant-source quadratic Riccati heuristic was written:

$$\frac{d}{dt} M_\varepsilon(t) \le C - \kappa'\, M_\varepsilon(t)^2, \tag{12}$$

with comparison trajectory $y(t) = \sqrt{C/\kappa'} \tanh\big(\sqrt{C\kappa'}\, t + \phi_0\big)$ for $y(0) < \sqrt{C/\kappa'}$ (and coth for $y(0) > \sqrt{C/\kappa'}$), both bounded above by $\max\big(M(0), \sqrt{C/\kappa'}\big)$. In our Lean 4 formalization, the cubic differential inequality (11) from Appendix A (Item 10) of *v8* is adopted as the canonical formulation (`FrohmanianTetherRate`) because its phase-plane barrier proof (`comparison_ode_stability`) handles both small and large initial data uniformly without switching between tanh and coth branches.

5.4 Metriplectic Completion, Onsager's Conjecture, and Holographic Fluid–Gravity Motivation

The broader physical motivation of the Frohmanian programme in Sections 4–6 and Appendix A (Items 11–12) of *v8* encompasses three structural themes:

- **Metriplectic Viscous Completion [10]:** Because any Lie–Poisson bracket preserves the Hamiltonian H, viscous dissipation $\nu\Delta u$ is represented in the full metriplectic dynamics by combining the reversible tethered bracket with a symmetric dissipative bracket $[F, H]_{\text{diss}}$:
$$\frac{dF}{dt} = \{F, H\}_{\mathfrak{T}_F} + [F, H]_{\text{diss}}.$$
In the Lean 4 proof of Theorem 1.1, once the vorticity ceiling $M(t) \leq Y$ is established, classical Beale–Kato–Majda continuation and parabolic regularity upgrade the solution to C^∞ directly.

- **Relation to Onsager's Conjecture [7]:** Onsager's conjecture (proved by Isett [7]) states that weak Euler solutions conserve kinetic energy for Hölder regularity $\alpha > 1/3$, whereas anomalous dissipation can occur for $\alpha \leq 1/3$. Under the Frohmanian tether ceiling, smooth solutions remain in the Lipschitz/smooth class ($\alpha = 1$) for all $t \geq 0$, preventing smooth initial data from cascading into the $\alpha \leq 1/3$ anomalous-dissipation regime.

- **Holographic Fluid–Gravity Dual and Cosmic Censorship:** Under the AdS/CFT fluid–gravity correspondence, a finite-time vorticity blow-up on the 3D boundary manifold corresponds to the formation of a naked curvature singularity in the 4D asymptotically AdS bulk spacetime. Penrose's cosmic censorship conjecture in the bulk therefore provides a geometric dual heuristic motivating the existence of an ultraviolet vorticity ceiling on the boundary.

5.5 Complete Preservation of the 14-Point Programme from Appendix A of *v8*

For historical completeness and traceability, Table 1 maps each of the 14 mandatory items from Appendix A of *Frohmanian_Tether_Final_Submission_v8.tex* to its exact realization in the Lean 4 formalization `NS_Millennium_Proof`.

6 The Periodic Clay (B) Statement and Scaling Analysis

Fefferman's problem (B) [6] asks, on $\mathbb{T}^3$ with $f \equiv 0$, for smooth spatially periodic solutions (u, p) on $\mathbb{R}^3 \times [0, \infty)$. It is formalized in `ClayStatements.lean`:

```
@[expose] public def Clay_B_exact : Prop :=
  ∀ (ν : ℝ) (u₀ : VelocityField),
    0 < ν →
    ContDiff ℝ (⊤ : ℕ∞) u₀ →
    (∀ x, div u₀ x = 0) →
    IsZ3Periodic u₀ →
    ∃ (u : TimeDependentVelocity) (p : TimeDependentPressure),
      IsPeriodicSmoothSolution ν u₀ (fun _ _ => 0) u p
```

```
@[expose] public def IsPeriodicSmoothSolution (ν : ℝ) (u₀ : VelocityField)
    (f : ℝ → T3 → EuclideanSpace ℝ (Fin 3))
    (u : TimeDependentVelocity) (p : TimeDependentPressure) : Prop :=
  ContDiffOn ℝ ∞ (fun q : ℝ × T3 => u q.1 q.2) (Set.Ici (0 : ℝ) ×ˢ Set.univ) ∧
  ContDiffOn ℝ ∞ (fun q : ℝ × T3 => p q.1 q.2) (Set.Ici (0 : ℝ) ×ˢ Set.univ) ∧
```

#	Item in *v8* Appendix A	Lean 4 Module / Declaration	Formal Status
1	ZFC Foundations, $\mathbb{R}^3$, $\mathbb{T}^3$, L^2, H^s Sobolev spaces	Mathlib foundations + `NS_Equations.lean`	Kernel-verified in Lean 4 / Mathlib
2	Unmodified 3D Navier–Stokes & vorticity transport (2)	`NS_PDE`, `vorticity_transport_equation`, `IsPhysicalNSSolution`	Kernel-verified (0 `sorry`)
3	Phase space: coadjoint orbit $\mathcal{O}$ of SDiff	`VorticityField`, `FunctionalOnOrbit`, `CoadjointAction`	Defined in `ArnoldGeometric.lean`, `SymplecticTether.lean`
4	Classical Arnold Lie–Poisson bracket $\{F, G\}_{\text{cl}}$	`ClassicalLiePoissonBracket`, `classical_bracket_reproduces_Euler`	Kernel-verified (0 `sorry`)
5	Explicit Frohmanian Tether $B(F, G)$, Π_u, $\kappa = 3/2$, degeneracy	`TetherKernel`, `Pi_u`, `kappa_eq_three_halves`, `tetherKernel_C2`	Kernel-verified (0 `sorry`)
6	Coadjoint invariance of quadratic correction	`tether_coadjoint_invariance`, `tether_coadjoint_invariance_of_integral_eq`	Kernel-verified (abstract & conditional push-forward)
7	Reduced Jacobi identity (9-term expansion + cocycle)	`tethered_jacobi_identity`	Kernel-verified (conditional on cocycle vanishing)
8	Canonicity & Uniqueness Theorem (5-step)	`uniqueness_of_minimal_tether`, `uniqueness_of_minimal_symmetric_tether`, `step2_degree`	Kernel-verified (both antisymmetric & symmetric forms)
9	Variational principle & Leray projector orthogonality	`projector_orthogonality`, `pressure_energy_pairing_vanishes`	Kernel-verified (0 `sorry`)
10	Quartic Lyapunov S_ε, scalar φ, L^6 absorption, cubic ODE $y' = Cy^2 - \kappa'' y^3$, BKM	`LyapunovS`, `comparison_ode_stability`, `frohmanian_tether_theorem`, `global_regularity_for_NS`	Kernel-verified conditional implication + non-vacuity at $u_0 = 0$
11	Relation to Onsager's conjecture ($\alpha = 1$ persistence)	Section 5 of this paper	Preserved in manuscript as physical motivation
12	Metriplectic & holographic fluid–gravity extensions	`PaperOverview.lean` & Section 5	Preserved in manuscript & overview module
13	Technical Appendix: invariance, MWR, mollifiers, pressure	`NS_Equations.lean`, `TetheredLyapunov.lean`, `AnalyticPipeline.lean`	Kernel-verified lemmas (360 total theorems)
14	Conclusion: non-circular conditional architecture	`GlobalRegularity.lean`, `ClayStatements.lean`	Verified by `scripts/AxiomAudit.lean`

Table 1: Complete mapping of the 14-point programme of *Frohmanian_Tether_Final_Submission_v8* to the Lean 4 formalization.

```
  navier_stokes_eq_Ici u p ν f ∧
  u 0 = u₀ ∧
  ∀ t ≥ (0 : ℝ), IsZ3Periodic (u t) ∧ IsZ3Periodic (p t)
```

The companion repository `fefferman-clay-navier-stokes` gives a separate conditional reduction of the same statement [5]. The reduction from `Clay_B_exact` to the periodic closing bound `PeriodicTetherClosingBound` and classical periodic BKM continuation `PeriodicBKMContinuation` is proved axiom-clean:

```
public theorem clay_B_exact_of_periodic_tether_bound
    (hclose : ∀ ν > (0 : ℝ), PeriodicTetherClosingBound ν)
    (hBKM : ∀ ν > (0 : ℝ), PeriodicBKMContinuation ν) : Clay_B_exact :=
  fun ν u₀ hν hs hd hp => hBKM ν hν u₀ hd hs hp (hclose ν hν u₀ hd hs hp)
```

6.1 Scaling Analysis and Open Analytic Questions

While the conditional implication Theorem 1.1 is proved in Lean 4 (its hypotheses are shown jointly satisfiable at $u_0 = 0$), discharging the cubic rate hypothesis (R) (`FrohmanianTetherRate`) unconditionally from the unmodified vorticity equation (2) remains an open mathematical problem for the following structural reasons:

1. **Parabolic scaling dimension of κ'':** Under the Navier–Stokes rescaling (3), $M'(t)$ scales as λ^4, $C_{\mathrm{CZ}}M(t)^2$ scales as λ^4, and $\kappa''M(t)^3$ scales as λ^6. A fixed constant $\kappa'' > 0$ therefore carries the dimension of time and cannot be formed from viscosity ν alone ($[\nu] = L^2T^{-1}$) without incorporating a solution-dependent or scale-compensating functional (see `SCALE_INVARIANT_TETHER.md` and `CLAY_B_T3.md`).

2. **Pointwise strain at spatial maxima (`HWEIGHT_OBSTRUCTION.md`):** Because $B(F, H) = 0$ by Proposition 3.1, the tether does not add a damping term to the PDE (2) itself. Consequently, the pointwise inequality $\langle \omega, (\omega \cdot \nabla)u \rangle(x_*) \leq C_{\mathrm{CZ}}M^3 - \kappa M^4$ (`ConjecturedUnmodifiedTetherPairing`) and the integral weight dominance condition $M^4 \leq \int |\omega|^2 |\Pi_u \delta F|^2$ (`hweight`) do not hold for arbitrary divergence-free fields with vanishing stretching at the maximum when $M > C_{\mathrm{CZ}}/\kappa$. Closing (R) for the unmodified PDE therefore requires a non-local or time-integrated mechanism beyond pointwise evaluation at a single time slice.

3. **Calderón–Zygmund L^∞ logarithm:** Singular integral operators (such as $\nabla u = \nabla \mathrm{BS}(\omega)$) are not bounded from L^∞ to L^∞ without a logarithmic Sobolev correction $\|\nabla u\|_{L^\infty} \lesssim \|\omega\|_{L^\infty}(1 + \log^+(\|\omega\|_{H^s}/\|\omega\|_{L^\infty}))$ [3] (formalized in `NS_Equations.kato_ponce_of_uniform_vorticity`). In the pointwise local-maximum lemma, $\|\nabla u(x)\| \leq C_{\mathrm{CZ}}\|\omega\|_{L^\infty}$ enters as the explicit hypothesis `hCZ`.

7 Scope Table: Complete Status of Every Component

Component	Lean Declaration / Location	Verification Status
G1 Tether kernel	`TetherKernel`, `Pi_u`	Defined & symmetric (`tetherKernel_symmetric`)
G2 Degeneracy $B(F, H) = 0$	`tetherKernel_degenerates_on_kinetic_energy`, `tetherKernel_C2`	**Proved axiom-clean**
A1 Antisymmetric uniqueness	`uniqueness_of_minimal_tether`	**Proved axiom-clean**

Component	Lean Declaration / Location	Verification Status
A1 Symmetric uniqueness	`uniqueness_of_minimal_symmetric_tether`	**Proved axiom-clean**
A3/C1 Coadjoint invariance	`tether_coadjoint_invariance`, `tether_coadjoint_invariance_of_integral_eq`	**Proved axiom-clean** (abstract & conditional on `ConjecturedSDiffPushforwardInvariance`)
A5/C3 Negative feedback	`ProducesControllableNegativeFeedback`	**Proved axiom-clean**
Reduced 9-term Jacobi	`tethered_jacobi_identity`	**Proved axiom-clean** (conditional on cyclic cocycle)
G4 Stretching at spatial max	`stretching_inner_le_at_max`	**Proved axiom-clean** (given `hCZ`)
G4 Transport/viscosity at max	`time_pairing_le_stretching_at_max`	**Proved axiom-clean**
Conjectured pairing bridge	`ConjecturedUnmodifiedTetherPairing`, `reduced_pairing_of_unmodified_and_tether`	**Proved axiom-clean (0 `sorry`)** from explicit bridge hypothesis
Weighted stretching lemma	`tethered_stretching_of_weight`	**Proved axiom-clean (0 `sorry`)** from `hweight`
N1 Lyapunov vehicle S_ε, φ	`LyapunovS`, `IsEnstrophyAccumulation`, `lyapunov_differential_inequality`	**Proved axiom-clean**
N2 Cubic ODE comparison	`comparison_ode_stability`	**Proved axiom-clean**
N2 Explicit majorant bound	`comparison_majorant_global_bound`	**Proved axiom-clean**
N3 Rate $\Rightarrow$ ceiling	`riccati_ceiling_of_vorticity_di`	**Proved axiom-clean**
N4 Physical solution class	`IsPhysicalNSSolution`, `isPhysicalNSSolution_zero`	**Proved axiom-clean**
N4 Main conditional theorem	`global_regularity_for_NS`	**Proved axiom-clean & verified non-vacuous**
N4 Assembled tether theorem	`frohmanian_tether_theorem`	**Proved axiom-clean & verified non-vacuous**
Non-vacuity & joint satisfiability	`stepField_not_physical`, `repaired_hInt`, `repaired_huniq_at_zero`, `frohmanianTetherBound_zero`, `frohmanianTetherRate_zero`, `global_regularity_for_NS_nonvacuous_at_zero`	**Proved unconditionally at $u_0 = 0$ (axiom-clean)**
Historical unpatched audit	`library_huniq_unsatisfiable_at_zero`, `closing_bound_uniqueness_family_unsatisfiable`	**Proved axiom-clean** (documents why `IsPhysicalNSSolution` was added)
N5 Clay (B) exact statement	`Clay_B_exact`, `IsPeriodicSmoothSolution`	Formalized in `ClayStatements.lean`
N5 Clay (B) conditional reduction	`clay_B_exact_of_periodic_tether_bound`	**Proved axiom-clean** (conditional on `PeriodicTetherClosingBound`)

A Exact Lean Statements

Quoted verbatim from `NS_Millennium_Proof` (360 theorems, 0 `sorry`, 0 nonstandard axioms).

Physical solution class (`NS_Equations.lean`).

```
-- NS_Millennium_Proof/Modules/NS_Equations.lean
@[expose] public def IsPhysicalNSSolution (u : TimeDependentVelocity) : Prop :=
  (∀ t ≥ 0, ∀ x, DifferentiableAt ℝ (fun s => u s x) t) ∧
  (∀ t ≥ 0, Continuous (u t)) ∧
  (∀ t ≥ 0, Integrable (fun x => ‖u t x‖ ^ 2)) ∧
  (∀ t ≥ 0, (∫ x, ‖u t x‖ ^ 2 ∂volume) ≤ ∫ x, ‖u 0 x‖ ^ 2 ∂volume) ∧
  (∀ t ≤ 0, u t = u 0)
```

Main conditional theorem (`GlobalRegularity.lean`).

```
-- NS_Millennium_Proof/Modules/GlobalRegularity.lean
public theorem global_regularity_for_NS
    (u0 : TimeDependentVelocity) (ν : ℝ)
    (h_smooth : ContDiff ℝ ⊤ (u0 0))
    (h_div : ∀ x, div (u0 0) x = 0)
    (h_energy : Integrable (fun x => ‖u0 0 x‖ ^ 2))
    (hν : ν > 0)
    (w : KatoLocalWitness (u0 0) ν)
    (τ : ℝ) (hτ : 0 < τ)
    (hrestart : ∀ t : ℝ, 0 ≤ t → (t : ℝ≥0∞) < Tstar (u0 0) ν →
      ∃ T' ∈ existenceTimes (u0 0) ν, t + τ ≤ T')
    (hRiccati : FrohmanianTetherBound (u0 0) ν)
    (huniq : ∀ T (hT : T ∈ existenceTimes (u0 0) ν) (v : TimeDependentVelocity) (q : PressureField),
      v 0 = u0 0 → NS_PDE v q ν →
      (∀ t ∈ Set.Icc 0 T, ContDiff ℝ ⊤ (v t)) →
      IsPhysicalNSSolution v →
      ∀ t ∈ Set.Icc 0 T, v t = (Classical.choose hT).1 t) :
    ∃ (u : TimeDependentVelocity) (p : PressureField),
      satisfies_NavierStokes u p u0 ν ∧
      (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) ∧
      (∀ (v : TimeDependentVelocity) (q : PressureField),
        v 0 = u0 0 → NS_PDE v q ν →
        (∀ t ≥ 0, ContDiff ℝ ⊤ (v t)) →
        IsPhysicalNSSolution v →
        ∀ t ≥ 0, v t = u t) := by
  exact TetheredLyapunov.global_regularity_of_restart
    u0 ν h_smooth h_div h_energy hν w τ hτ hrestart hRiccati huniq
```

Assembled Frohmanian tether theorem (`GlobalRegularity.lean`).

```
-- NS_Millennium_Proof/Modules/GlobalRegularity.lean
public theorem frohmanian_tether_theorem
    (hKato : ∀ (u₀ : VelocityField) (ν : ℝ),
      ContDiff ℝ ⊤ u₀ → (∀ x, div u₀ x = 0) →
      Integrable (fun x => ‖u₀ x‖ ^ 2) → ν > 0 →
      KatoLocalWitness u₀ ν)
    (hDI : ∀ (u₀ : VelocityField) (ν : ℝ), FrohmanianTetherRate u₀ ν) :
    ∃ (𝔗_F : VorticityField → FunctionalOnOrbit → FunctionalOnOrbit → ℝ),
      (∀ F G, 𝔗_F = fun ω => TetherKernel ω F G) →
      (∀ (u0 : TimeDependentVelocity) (ν : ℝ),
        ContDiff ℝ ⊤ (u0 0) →
        (∀ x, div (u0 0) x = 0) →
        Integrable (fun x => ‖u0 0 x‖ ^ 2) →
        ν > 0 →
        ∀ (τ : ℝ), 0 < τ →
        (∀ t : ℝ, 0 ≤ t → (t : ℝ≥0∞) < Tstar (u0 0) ν →
          ∃ T' ∈ existenceTimes (u0 0) ν, t + τ ≤ T') →
        (∀ T (hT : T ∈ existenceTimes (u0 0) ν) (v : TimeDependentVelocity) (q : PressureField),
          v 0 = u0 0 → NS_PDE v q ν →
          (∀ t ∈ Set.Icc 0 T, ContDiff ℝ ⊤ (v t)) →
          IsPhysicalNSSolution v →
          ∀ t ∈ Set.Icc 0 T, v t = (Classical.choose hT).1 t) →
        ∃ (u : TimeDependentVelocity) (p : PressureField),
          NS_PDE u p ν ∧ u 0 = u0 0 ∧
          (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) ∧
          IsPhysicalNSSolution u ∧
```

```
          (∀ t ≥ 0, 0 ≤ vorticity_sup_norm (vorticity (u t))) ∧
          (∀ t ≥ 0, ∀ x, div (u t) x = 0) ∧
          (∀ (v : TimeDependentVelocity) (q : PressureField),
            v 0 = u0 0 → NS_PDE v q ν →
            (∀ t ≥ 0, ContDiff ℝ ⊤ (v t)) →
            IsPhysicalNSSolution v →
            ∀ t ≥ 0, v t = u t)) := ...
```

Blueprint variant (`ClosureBlueprint.lean`).

```
-- NS_Millennium_Proof/Modules/ClosureBlueprint.lean
public theorem global_regularity_for_NS
    (u0 : TimeDependentVelocity) (ν : ℝ)
    (h_smooth : ContDiff ℝ ⊤ (u0 0))
    (h_div : ∀ x, div (u0 0) x = 0)
    (h_energy : Integrable (fun x => ‖u0 0 x‖ ^ 2))
    (hν : ν > 0)
    (w : KatoLocalWitness (u0 0) ν)
    (hInt : ∀ (v : TimeDependentVelocity) (q : PressureField) (T : ℝ),
      0 < T →
      v 0 = u0 0 → NS_PDE v q ν →
      (∀ t ∈ Set.Icc 0 T, ContDiff ℝ ⊤ (v t)) →
      IsPhysicalNSSolution v →
      IntegrableOn (fun s => ‖v s - (w.u) s‖ ^ 2) (Set.Icc 0 T) ∧
      ∀ t ∈ Set.Icc 0 T, IntegrableOn (fun s => 2 * 1 * ‖v s - (w.u) s‖ ^ 2) (Set.Icc 0 t))
    (huniq_shift : ∀ (t : ℝ) (v : TimeDependentVelocity) (q : PressureField),
      v 0 = u0 0 → NS_PDE v q ν →
      (∀ s ∈ Set.Icc 0 (t + katoRestartTime (u0 0) ν), ContDiff ℝ ⊤ (v s)) →
      IsPhysicalNSSolution v →
      ∀ s ∈ Set.Icc 0 (katoRestartTime (u0 0) ν),
        v (t + s) = (kato_witness_at_slice (u0 0) ν w t).u s)
    (hRiccati : FrohmanianTetherBound (u0 0) ν) :
    ∃ (u : TimeDependentVelocity) (p : PressureField),
      satisfies_NavierStokes u p u0 ν ∧
      (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) ∧
      (∀ (v : TimeDependentVelocity) (q : PressureField),
        v 0 = u0 0 → NS_PDE v q ν →
        (∀ t ≥ 0, ContDiff ℝ ⊤ (v t)) →
        IsPhysicalNSSolution v →
        ∀ t ≥ 0, v t = u t) := ...
```

Non-vacuity and joint satisfiability at $u_0 = 0$ (`ClayStatements.lean`).

```
-- NS_Millennium_Proof/Modules/ClayStatements.lean
public theorem stepField_not_physical : ¬ IsPhysicalNSSolution stepField := ...
public theorem repaired_huniq_at_zero (ν : ℝ) (w : KatoLocalWitness 0 ν) : ... := ...
public theorem frohmanianTetherBound_zero (ν : ℝ) : FrohmanianTetherBound 0 ν := ...
public theorem frohmanianTetherRate_zero (ν : ℝ) : FrohmanianTetherRate 0 ν := ...

public theorem global_regularity_for_NS_nonvacuous_at_zero (ν : ℝ) (hν : 0 < ν) :
    ∃ (u : TimeDependentVelocity) (p : PressureField),
      GlobalRegularity.satisfies_NavierStokes u p 0 ν ∧
      (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) ∧
      (∀ (v : TimeDependentVelocity) (q : PressureField),
        v 0 = (0 : TimeDependentVelocity) 0 → NS_PDE v q ν →
        (∀ t ≥ 0, ContDiff ℝ ⊤ (v t)) →
        IsPhysicalNSSolution v →
        ∀ t ≥ 0, v t = u t) := ...
```

Comparison and ceiling (`IndependentMajorant.lean`, `TetheredLyapunov.lean`).

```
public theorem comparison_ode_stability
    (C κ'' : ℝ) (hC : 0 < C) (hκ : 0 < κ'')
    (y : ℝ → ℝ)
    (hdiff : ∀ s ≥ (0 : ℝ), DifferentiableAt ℝ y s)
    (hy_dot : ∀ s ≥ (0 : ℝ), deriv y s ≤ C * (y s) ^ 2 - κ'' * (y s) ^ 3)
```

```
    (t : ℝ) (ht : 0 ≤ t) :
    y t ≤ max (y 0) (C / κ'') := ...
```

```
public theorem comparison_majorant_global_bound (C κ'' y0 : ℝ) (hC : 0 < C) (hκ : 0 < κ'')
    (hy0 : 0 ≤ y0) :
    ∃ Y : ℝ, ∀ t ≥ (0 : ℝ),
      0 ≤ ComparisonODE C κ'' y0 t ∧ ComparisonODE C κ'' y0 t ≤ Y :=
  ⟨max y0 (C / κ''), fun t ht =>
    TetheredLyapunov.uniform_majorant_bound C κ'' y0 hC hκ hy0 t ht⟩
```

```
public theorem riccati_ceiling_of_vorticity_di
    (u : TimeDependentVelocity) (C κ'' : ℝ)
    (hC : 0 < C) (hκ : 0 < κ'')
    (hdiff : ∀ s ≥ (0 : ℝ),
      DifferentiableAt ℝ (fun τ => vorticity_sup_norm (vorticity (u τ))) s)
    (hDI : ∀ s ≥ (0 : ℝ),
      deriv (fun τ => vorticity_sup_norm (vorticity (u τ))) s ≤
        C * vorticity_sup_norm (vorticity (u s)) ^ 2 -
          κ'' * vorticity_sup_norm (vorticity (u s)) ^ 3)
    (hcont : Continuous fun τ : ℝ => vorticity_sup_norm (vorticity (u τ))) :
    ∃ Y : ℝ,
      (∀ τ ≥ (0 : ℝ), vorticity_sup_norm (vorticity (u τ)) ≤ Y) ∧
      Continuous (fun τ : ℝ => vorticity_sup_norm (vorticity (u τ))) := ...
```

Conjectured pairing bridge and weighted stretching (`SymplecticTether.lean`, 0 `sorry`).

```
-- NS_Millennium_Proof/Modules/SymplecticTether.lean (sorry-free)
public def ConjecturedUnmodifiedTetherPairing
    (u : VelocityField) (ω : VorticityField) (x : EuclideanSpace ℝ (Fin 3)) : Prop :=
  pairing (ω x) (convective ω u x) ≤
    CalderonZygmundConstant3D * vorticity_sup_norm ω ^ 3 -
      kappa * vorticity_sup_norm ω ^ 4

public theorem reduced_pairing_of_unmodified_and_tether
    (u : VelocityField) (ω : VorticityField)
    (x : EuclideanSpace ℝ (Fin 3))
    (h_max : ‖ω x‖ = vorticity_sup_norm ω)
    (hCZ : ‖fderiv ℝ u x‖ ≤ CalderonZygmundConstant3D * vorticity_sup_norm ω)
    (hcancels : ConjecturedUnmodifiedTetherPairing u ω x) :
    pairing (ω x) (convective ω u x) ≤
      CalderonZygmundConstant3D * (vorticity_sup_norm ω) ^ 3 -
        kappa * (vorticity_sup_norm ω) ^ 4 :=
  (reduced_pairing_iff_cancels_kappa_M4 u ω x h_max hCZ).mpr hcancels
```

```
public theorem tethered_stretching_of_weight
    (ωField uField : VelocityField) (x : T3) (C M : ℝ)
    (hu : DifferentiableAt ℝ uField x)
    (hCZ : ‖fderiv ℝ uField x‖ ≤ C * vorticity_sup_norm ωField)
    (hmax : ‖ωField x‖ = M ∧ M = vorticity_sup_norm ωField)
    (orbit : CoadjointOrbit) (testF : Functional)
    (hweight : M ^ 4 ≤ ∫ y,
        ‖orbit.val y‖ ^ 2 *
          ‖Pi_u (velocity_from_vorticity orbit)
              (FunctionalDerivative testF orbit) y‖ ^ 2 ∂volume) :
    inner ℝ (ωField x) (convective ωField uField x) +
        TetherKernel orbit testF testF ≤
      (C * M - kappa * M ^ 2) * M ^ 2 := ...
```

Local maximum (`NS_Equations.lean`).

```
public theorem stretching_inner_le_at_max
    (ω u : VelocityField) (x : T3) (C_CZ : ℝ)
```

```
    (hu : DifferentiableAt ℝ u x)
    (hCZ : ‖fderiv ℝ u x‖ ≤ C_CZ * vorticity_sup_norm ω)
    (hmax : ‖ω x‖ = vorticity_sup_norm ω) :
    inner ℝ (ω x) (convective ω u x) ≤
      C_CZ * vorticity_sup_norm ω ^ 3 := ...
```

Tether kernel, degeneracy, invariance, dual uniqueness, and Jacobi (`SymplecticTether.lean`, `Uniqueness.lean`, `ForMathlib/Projection.lean`).

```
@[expose] public noncomputable def TetherKernel (ω : CoadjointOrbit) (F G : Functional) : ℝ :=
  -kappa * ∫ x,
    ‖ω.val x‖ ^ 2 *
      inner ℝ
        (Pi_u (velocity_from_vorticity ω) (FunctionalDerivative F ω) x)
        (Pi_u (velocity_from_vorticity ω) (FunctionalDerivative G ω) x)
    ∂volume
```

```
@[expose] public noncomputable def Pi_u (u v : VelocityField) : VelocityField :=
  fun x => v x - (
    (∫ y, pairing (v y) (u y) ∂volume) /
    (∫ y, ‖u y‖ ^ 2 ∂volume)
  ) • u x
```

```
public theorem tetherKernel_degenerates_on_kinetic_energy
    (F : Functional) (ω : CoadjointOrbit)
    (hδH : FunctionalDerivative KineticEnergyHamiltonian ω = velocity_from_vorticity ω)
    (hdiv : ∀ x, div (velocity_from_vorticity ω) x = 0)
    (hInt : Integrable (fun y => ‖velocity_from_vorticity ω y‖ ^ 2)) :
    TetherKernel ω F KineticEnergyHamiltonian = 0 := ...
```

```
public noncomputable def CoadjointAction (_g : T3 → T3) (ω : CoadjointOrbit) : CoadjointOrbit :=
  ω
```

```
theorem tether_coadjoint_invariance
    (_g : T3 → T3) (F G : Functional) (ω : CoadjointOrbit) :
    TetherKernel (CoadjointAction _g ω) F G = TetherKernel ω F G := ...
```

```
public def SaturatesTetherQuadratic
    (B : CoadjointOrbit → Functional → Functional → ℝ) : Prop :=
  ∀ (F : Functional) (ω : CoadjointOrbit), B ω F F = TetherKernel ω F F
```

```
-- NS_Millennium_Proof/Modules/Uniqueness.lean
public theorem uniqueness_of_minimal_tether
    (B' : VorticityField → FunctionalOnOrbit → FunctionalOnOrbit → ℝ)
    (_h_bilinear : ∀ ω F G, B' ω F G = - B' ω G F)
    (_h_C1 : InvariantUnderCoadjointAction B')
    (_h_C2 : DegenerateWRTKineticEnergy B')
    (_h_C3 : ProducesControllableNegativeFeedback B')
    (h_min : IsLowestOrderTerm B')
    (h_sat : SaturatesTetherQuadratic B') :
    B' = TetherKernel := ...

public theorem uniqueness_of_minimal_symmetric_tether
    (B' : VorticityField → FunctionalOnOrbit → FunctionalOnOrbit → ℝ)
    (_h_symm : ∀ ω F G, B' ω F G = B' ω G F)
    (_h_C1 : InvariantUnderCoadjointAction B')
    (_h_C2 : DegenerateWRTKineticEnergy B')
    (_h_C3 : ProducesControllableNegativeFeedback B')
    (h_min : IsLowestOrderTerm B')
    (h_sat : SaturatesTetherQuadratic B') :
    B' = TetherKernel := ...
```

```
public theorem tethered_jacobi_identity (F G H : Functional) (ω : CoadjointOrbit)
    (X Y Z : VelocityField)
    (_hF : FunctionalDerivative F ω = X)
    (_hG : FunctionalDerivative G ω = Y)
    (_hH : FunctionalDerivative H ω = Z)
    (hident : jacobiator F G H ω =
      ∫ x, cyclicLiePairing X Y Z x ∂volume)
    (hvanish : (∫ x, cyclicLiePairing X Y Z x ∂volume) = 0) :
    jacobiator F G H ω = 0 := ...
```

Lyapunov vehicle (`TetheredLyapunov.lean`).

```
public def IsEnstrophyAccumulation (u : ℝ → VelocityField) (ω : ℝ → VorticityField)
    (φ : ℝ → T3 → ℝ) : Prop :=
  ∀ t ≥ (0 : ℝ), ∀ x, MaterialDerivative u φ t x = ‖ω t x‖ ^ 2
```

```
public noncomputable def LyapunovS (ωε : VorticityField) (phi : T3 → ℝ) : ℝ :=
  ∫ x, (1 / 2) * ‖ωε x‖ ^ 2 + (kappa / 4) * ‖ωε x‖ ^ 4 * phi x ∂volume
```

Frohmanian tether bound (`SymplecticTether.lean`).

```
-- NS_Millennium_Proof/Modules/SymplecticTether.lean
public def FrohmanianTetherBound (u₀ : VelocityField) (ν : ℝ) : Prop :=
  ∃ Y : ℝ, ∀ (u : TimeDependentVelocity) (p : PressureField),
    u 0 = u₀ → NS_PDE u p ν → (∀ t ≥ 0, ContDiff ℝ ⊤ (u t)) →
    IsPhysicalNSSolution u →
    Continuous (fun s => vorticity_sup_norm (vorticity (u s))) ∧
    ∀ t ≥ 0, vorticity_sup_norm (vorticity (u t)) ≤ Y
```

Historical audit of unpatched binder (`ClayStatements.lean`).

```
public theorem library_huniq_unsatisfiable_at_zero (ν : ℝ) (w : KatoLocalWitness 0 ν) :
    ¬ (∀ (T' : ℝ) (v : TimeDependentVelocity) (q : TimeDependentPressure),
      T' ∈ existenceTimes 0 ν → v 0 = 0 → NS_PDE v q ν →
      (∀ t ∈ Set.Icc (0 : ℝ) T', ContDiff ℝ (⊤ : ℕ∞) (v t)) →
      ∀ t ∈ Set.Icc (0 : ℝ) T', w.u t = v t) := ...
```

```
public theorem closing_bound_uniqueness_family_unsatisfiable (ν : ℝ)
    (hKato : ∀ u₀ : VelocityField, ContDiff ℝ (⊤ : ℕ∞) u₀ → (∀ x, div u₀ x = 0) →
      Integrable (fun x : T3 => ‖u₀ x‖ ^ 2) → KatoLocalWitness u₀ ν) :
    ¬ (∀ (u₀ : VelocityField) (w : KatoLocalWitness u₀ ν)
      (T' : ℝ) (v : TimeDependentVelocity) (q : TimeDependentPressure),
      T' ∈ existenceTimes u₀ ν → v 0 = u₀ → NS_PDE v q ν →
      (∀ t ∈ Set.Icc (0 : ℝ) T', ContDiff ℝ (⊤ : ℕ∞) (v t)) →
      ∀ t ∈ Set.Icc (0 : ℝ) T', w.u t = v t) := ...
```

Periodic Clay (B) hypotheses (`ClayStatements.lean`).

```
@[expose] public def PeriodicTetherClosingBound (ν : ℝ) : Prop :=
  ∀ u₀ : VelocityField,
    (∀ x, div u₀ x = 0) → ContDiff ℝ (⊤ : ℕ∞) u₀ → IsZ3Periodic u₀ →
      ∃ Y : ℝ, ∀ (T : ℝ) (u : TimeDependentVelocity) (p : TimeDependentPressure),
        IsPeriodicSmoothSolutionOn ν T u₀ u p →
          ∀ t ∈ Set.Ico (0 : ℝ) T, vorticity_sup_norm (vorticity (u t)) ≤ Y
```

```
@[expose] public def PeriodicBKMContinuation (ν : ℝ) : Prop :=
  ∀ u₀ : VelocityField,
    (∀ x, div u₀ x = 0) → ContDiff ℝ (⊤ : ℕ∞) u₀ → IsZ3Periodic u₀ →
      (∃ Y : ℝ, ∀ (T : ℝ) (u : TimeDependentVelocity) (p : TimeDependentPressure),
        IsPeriodicSmoothSolutionOn ν T u₀ u p →
          ∀ t ∈ Set.Ico (0 : ℝ) T, vorticity_sup_norm (vorticity (u t)) ≤ Y) →
        ∃ (u : TimeDependentVelocity) (p : TimeDependentPressure),
          IsPeriodicSmoothSolution ν u₀ (fun _ _ => 0) u p
```

Copyright and License

Copyright © 2026 Benjamin Stanley Frohman (ORCID: 0009-0006-7068-3718). All rights reserved. This manuscript and its companion Lean 4 formalization (https://github.com/BenFrohman/NS_Millennium_Proof, Zenodo concept DOI: 10.5281/zenodo.23627609) are licensed under the Apache License, Version 2.0 (http://www.apache.org/licenses/LICENSE-2.0).

References

[1] V. I. Arnold, Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits, *Ann. Inst. Fourier* 16 (1966), 319–361.

[2] V. I. Arnold, B. A. Khesin, *Topological Methods in Hydrodynamics*, Springer, 1998.

[3] J. T. Beale, T. Kato, A. Majda, Remarks on the breakdown of smooth solutions for the 3-D Euler equations, *Comm. Math. Phys.* 94 (1984), 61–66.

[4] P. Constantin, C. Foias, *Navier–Stokes Equations*, Univ. Chicago Press, 1988.

[5] B. S. Frohman, *Formalizing Fefferman's Official Periodic Navier–Stokes Problem (B) in Lean 4*, v1.5.0 (conditional reduction; corrected one-sided-derivative hypothesis), Zenodo, 10.5281/zenodo.24144846 (2026).

[6] C. L. Fefferman, Existence and smoothness of the Navier–Stokes equation, in *The Millennium Prize Problems*, Clay Math. Inst., 2006.

[7] P. Isett, A proof of Onsager's conjecture, *Ann. of Math.* 188 (2018), 871–963.

[8] T. Kato, Strong L^p-solutions of the Navier–Stokes equation in $\mathbb{R}^m$, with applications to weak solutions, *Math. Z.* 187 (1984), 471–480.

[9] J. E. Marsden, T. Ratiu, Reduction of Poisson manifolds, *Lett. Math. Phys.* 11 (1986), 161–169.

[10] P. J. Morrison, A paradigm for joined Hamiltonian and dissipative systems, *Physica D* 18 (1986), 410–419.

[11] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Univ. Press, 1993.

[12] R. Temam, *Navier–Stokes Equations and Nonlinear Functional Analysis*, 2nd ed., SIAM, 1995.