Work in progress — draft, not peer reviewed

A Lean 4 Formalization of the Classical Continuation Chain for Periodic Navier–Stokes on $\mathbb{T}^3$: Energy Identity, H^3 Commutator Reduction and the BKM Route

Companion paper to `BenFrohman/periodic-ns-classical-continuation`

Benjamin Stanley Frohman
ORCID 0009-0006-7068-3718 · frohmanbenjamin@gmail.com

Draft of 31 October 2026 (Lean development at commit `f9a26c7`)
DOI 10.5281/zenodo.23842555 · Companion reduction paper: 10.5281/zenodo.23842295

How to cite / What is claimed. In Lean 4 with Mathlib, we give a machine-checked reduction of Clay Millennium Statement (B) (periodic Navier–Stokes on $\mathbb{T}^3$) to the single hypothesis `PeriodicBKMIntegrability`. We prove this hypothesis is equivalent to global existence of smooth solutions for every smooth periodic initial field. The BKM criterion and H^3 control are proved unconditionally, using only the standard axioms. `PeriodicBKMIntegrability` remains open. The equivalence of vorticity integrability with continuation of smooth solutions is classical (Beale–Kato–Majda, 1984 [1]); the novelty here is the complete, `sorry`-free formalization. Cite as: B. S. Frohman, reduction paper, DOI 10.5281/zenodo.23842295; main paper, DOI 10.5281/zenodo.23842555.

Draft — tracks build in progress. This document reports exactly what the Lean development proves at the stated commit. It does *not* solve the Clay Millennium problem, and does not prove global regularity of the 3D Navier–Stokes equations. The step `PeriodicBKMIntegrability` is the open Clay core and remains an explicit hypothesis.

Abstract

We describe a Lean 4/Mathlib [4, 15] formalization of the classical continuation steps for smooth divergence-free solutions of the incompressible Navier–Stokes equations on the unit torus $\mathbb{T}^3$, along the Statement (B) route [5] of the companion repository `fefferman-clay-navier-stokes`. Proved with no hypothesis Props: local existence for $\nu \geq 0$, uniqueness via an L^2 energy-difference estimate, the Fourier coefficient ODE, coefficient decay, the H^s energy identity, Parseval's identity on the unit cube, and the Kato–Ponce toolkit [7] at $s = 3$ (integration by parts, Leibniz rule, a Gagliardo–Nirenberg inequality, a bound on every lower-order term, and vanishing of the top-order term from $\operatorname{div} u = 0$). Also proved: the $s = 3$ physical-space commutator bound `periodicPhysicalCommutatorBound`. Also proved: the low-frequency vorticity bound `periodicLowFreqVorticityBound`, and hence, with no hypothesis Props, H^3 Beale–Kato–Majda control (`periodicBKMHsControl_three`) and the periodic BKM criterion (`periodicBKMCriterion`). Statement (B) is proven conditionally on `PeriodicBKMIntegrability`, which is open (`clay_B_exact_of_integrability`). Small zero-mean H^1 data give $T^* = \infty$ conditionally on `PeriodicEnstrophyRateBound` and `PeriodicH1BlowupCriterion`; at $\kappa = 0$ the tether headline is equivalent to the open `PeriodicDirectionRegularity`; `periodicBKMIntegrability_of_bending` gives integrability from `PeriodicDirectionEnstrophyGrowthL1`, `PeriodicH1BlowupCriterion` and the open `PeriodicBendingRateHiddenProperty`.

2020 Mathematics Subject Classification. 35Q30, 76D05, 68V15; 35B65, 03B35.

Keywords. Navier–Stokes equations; periodic domain; Beale–Kato–Majda criterion; continuation; Lean 4; Mathlib; formal verification; work in progress.

© 2026 B. S. Frohman · Apache-2.0

1 Introduction

The incompressible Navier–Stokes equations on the three-dimensional torus $\mathbb{T}^3$ are the setting of Statement (B) of the Clay Millennium Problem [5]. Their classical continuation theory—local existence, uniqueness, the H^s energy identities and the Beale–Kato–Majda (BKM) criterion [1]—is well established on paper. The purpose of this work is to formalize that theory in Lean 4 with Mathlib [4, 15], and to record precisely which steps are machine-checked without hypotheses, which are machine-checked relative to explicitly named hypotheses, and which remain open.

Section 4 collects the results proved with no hypothesis Props. Section 5 records the conditional results, culminating in the reduction of Statement (B) to the single open hypothesis `PeriodicBKMIntegrability`. Section 6 documents the further reductions of that hypothesis obtained so far, and Section 7 summarizes the status of every named Prop. No result in this paper resolves the Clay problem.

2 Conventions and naming

Every hypothesis is a named `Prop` passed as an explicit theorem argument; no `axiom`, `sorry` or `admit` appears. The axiom audit (`lake env lean AxiomAudit.lean`) reports only `propext`, `Classical.choice` and `Quot.sound`. Lean identifiers defined in this repository are *coined in this build* (marked †), not names from the author's May/June 2026 manuscript. The identifiers `Clay_B_exact`, `PeriodicLocalExistence`, `PeriodicBKMCriterion` and `PeriodicBKMIntegrability` come from the dependency `fefferman-clay-navier-stokes` (pinned at `ebcf109`).

3 Relation to the May/June 2026 manuscript

The author's manuscript frames regularity through a tethered Lyapunov functional S_ε and a comparison ODE

$$y' = Cy^2 - \kappa'' y^3, \tag{3.1}$$

with transport cancellation, viscous dissipation, a Hölder–Sobolev gate and Young absorption at $\varepsilon = \kappa/4$ (Proposition 2). For *unmodified* Navier–Stokes ($\kappa = 0$) the cubic damping is absent, and the remaining gap is hypothesis (H5): a non-circular majorant for the vorticity. In this repository that gap appears exactly as `PeriodicBKMIntegrability`, i.e.

$$\int_0^T \|\omega(t)\|_{L^\infty}\,\mathrm{d}t < \infty. \tag{3.2}$$

Nothing here closes it. The transport cancellation of the manuscript corresponds to `top_term_zero` below.

4 Proved results (no hypothesis Props)

Theorem 4.1 (Local existence)**.** `periodicLocalExistence_of_nonneg`†*: for $\nu \geq 0$,* `PeriodicLocalExistence` *ν holds (Galerkin method at $s = 3$).*

Theorem 4.2 (Energy difference / uniqueness)**.** `periodicEnergyDifferenceEstimate`†*: for $\nu \geq 0$ and two classical solutions u, v with equal data,*

$$E(t) = \int_{[0,1]^3} |u - v|^2\,\mathrm{d}x \quad \textit{satisfies} \quad E'(t) \leq 2C\,E(t).$$

© 2026 B. S. Frohman · Apache-2.0

With Grönwall (`eq_zero_of_deriv_le_mul`†*,* `periodicVelocityUniqueness_of_energy`†*) this gives uniqueness.*

Theorem 4.3 (Fourier coefficient ODE, decay, H^s energy identity)**.** `periodicFourierCoeffODE`†*,* `periodicFourierCoeffDecay`† *and* `periodicFourierHsEnergyIdentity`† *(file* `CoeffODE.lean`*) prove the Props* `PeriodicFourierCoeffODE`*,* `PeriodicFourierCoeffDecay`*,* `PeriodicFourierHsEnergyIdentity` *for every ν (and every s): each Fourier coefficient $\hat{u}_k(t)$ obeys the projected ODE, coefficients decay faster than any power, and*

$$\frac{\mathrm{d}}{\mathrm{d}t}\|u\|_{H^s}^2 = -2\nu\sum_k (1+|k|^2)^s\, 4\pi^2|k|^2|\hat{u}_k|^2 + \big(\textit{nonlinear rate}\big). \tag{4.1}$$

The bridge is `periodicFourierHsEnergyIdentity_of_coeff`† *(*`HsEnergyIdentity.lean`*).*

Theorem 4.4 (Parseval on the unit cube)**.** `hasSum_prod_cubeCoeff`† *(*`Parseval.lean`*): for continuous periodic f, g,*

$$\sum_{k\in\mathbb{Z}^3} \overline{\hat{f}_k}\,\hat{g}_k = \int_{[0,1]^3} \bar{f}\, g\,\mathrm{d}x,$$

via `torusLift`† *and Mathlib's torus Parseval.*

Theorem 4.5 (Physical form of the H^3 rate)**.** `hasSum_rate_three`†*: the Fourier-side H^3 nonlinear rate is summable and equals*

$$-2\sum_i \int \Big(1-\frac{\Delta}{4\pi^2}\Big)^3 v_i\,\big((v\cdot\nabla)v\big)_i\,\mathrm{d}x.$$

Theorem 4.6 (Kato–Ponce toolkit, $s=3$; `KatoPonce.lean`)**.** *Proved: unit-cell integration by parts* `cellI_ibp`†*,* `cellI_ibp'`†*; Leibniz rule* `DL_mul`†*; Gagliardo–Nirenberg on the cell* `gagliardoNirenberg_cell`† *and* `cellI_sq_mul_abs_le`†*; and*

- `lower_term_bound`†*: every lower-order Leibniz term (first factor of order ≥ 1) satisfies*

  $$\Big|\int D^L v_i \cdot D^{p_1} v_l\, D^{p_2}\partial_l v_i\,\mathrm{d}x\Big| \leq 5\|\nabla v\|_{L^\infty}\, Q(v),$$

  with $Q =$ `physQ`†*;*
- `top_term_zero`†*:* $\displaystyle\sum_{i,l}\int D^L v_i\, v_l\, D^L\partial_l v_i\,\mathrm{d}x = 0$*, because* $\operatorname{div} v = 0$ *(transport cancellation).*

Theorem 4.7 (Continuation alternative)**.** `periodicBKMCriterion_of_hsControl`†*: for $\nu \geq 0$, $s > 5/2$,* `PeriodicBKMHsControl` *ν s implies* `PeriodicBKMCriterion` *ν (the H^s blow-up alternative is proved).*

5 Conditional results

Named hypothesis 5.1 (`PeriodicPhysicalCommutatorBound` $C^\dagger$, $s=3$)**.** For smooth periodic divergence-free v,

$$-2\sum_i \int \Big(1-\frac{\Delta}{4\pi^2}\Big)^3 v_i\,\big((v\cdot\nabla)v\big)_i\,\mathrm{d}x \leq C\|\nabla v\|_{L^\infty}\|v\|_{H^3}^2. \tag{5.1}$$

Status: **proved** in `CommutatorAssembly.lean`: `periodicPhysicalCommutatorBound`† gives it with $C = 2\cdot 64\cdot 315\cdot 64\,(4\pi^2)^3$, from `lower_term_bound`, `top_term_zero`, `word_bound`† and `physQ_le_hsSq`†. No longer a hypothesis.

© 2026 B. S. Frohman · Apache-2.0

Named hypothesis 5.2 (`PeriodicLowFreqVorticityBound`†)**.** There is $K \geq 0$ such that for all $R \geq 2$,

$$\|V_R \nabla v\|_{L^\infty} \leq K(1 + \log R)\|\omega\|_{L^\infty}. \tag{5.2}$$

Status: **proved** in `LowFreqAssembly.lean` (`periodicLowFreqVorticityBound`†). The proof writes the Biot–Savart law in Riesz form,

$$\widehat{\partial_j v_i} = q_{j,i+2}\,\hat{\omega}_{i+1} - q_{j,i+1}\,\hat{\omega}_{i+2}, \qquad q_{jb} = \frac{k_j k_b}{|k|^2}.$$

It splits q into a low piece and $S = \lceil \log_2 R \rceil + 3$ smooth dyadic blocks, each with a uniformly L^1-bounded kernel (`DyadicBlock.lean`). Every piece is followed by the uniformly L^1-bounded de la Vallée Poussin kernel (`VPKernel.lean`). No longer a hypothesis.

Named hypothesis 5.3 (`PeriodicBKMIntegrability` ν)**.** From the dependency `fefferman-clay-navier-stokes`: $\int_0^T \|\omega\|_{L^\infty}\,\mathrm{d}t < \infty$ on every finite interval of existence. *Status:* **open**. This is the Clay core and the manuscript's (H5) gap.

Proposition 5.4. `periodicFourierCommutatorBound_of_physical`†*:* `PeriodicPhysicalCommutatorBound` $C \Rightarrow$ `PeriodicFourierCommutatorBound` 3 C*.* `periodicBKMHsControl_of_physical_lowFreq`†*:* $(\exists C \geq 0,$ `PeriodicPhysicalCommutatorBound` $C)$ *and* `PeriodicLowFreqVorticityBound` *give* `PeriodicBKMHsControl` ν *3. Combined with the proved bound:* `periodicBKMHsControl_three_of_lowFreq`† *gives* `PeriodicBKMHsControl` ν *3 from* `PeriodicLowFreqVorticityBound` *alone.*

Theorem 5.5 (Unconditional corollaries, `BKMFinal.lean`)**.** `periodicBKMHsControl_three`†*:* `PeriodicBKMHsControl` ν *3 for every* ν*, with no hypotheses.* `periodicBKMCriterion`†*:* `PeriodicBKMCriterion` ν *for every* ν*, with no hypotheses.*

Proposition 5.6 (Statement (B), conditional)**.** `clay_B_exact_of_hsControl`†*: for* $s > 5/2$*,* `PeriodicBKMHsControl` ν s *and* `PeriodicBKMIntegrability` ν *give* `Clay_B_exact` ν*. Also* `clay_B_exact_of_corrected_route'`† *and* `clay_B_exact_of_bkm`†*. The final form,* `clay_B_exact_of_integrability`†*, is conditional* only *on the open* `PeriodicBKMIntegrability` ν*.*

Remark 5.7 (A failed route). A Wiener-algebra/Fourier commutator estimate was tried and abandoned: its low-frequency part grows like $R^{3/2}$, not $\log R$. The $s = 3$ physical-space estimate replaces it.

6 Further reductions of the open hypothesis (work in progress)

Files `IntegrabilityAttack.lean`, `SmallDataAttack.lean`, `TetherClaimAttack.lean`, `EnstrophyIdentity.lean`, `TetherKappaZero.lean`, `DirectionCriterion.lean` (commits `f981e71`, `052cef1`, `ea5133d`, `0cd3551`, `a35d13a`); the file names are historical. All identifiers in this section are coined. **This does not prove `PeriodicBKMIntegrability`.**

- `integrableOn_vorticity_of_lt_tstar` (proved): if $T < T^*$, then $\int_0^T \|\omega\|_{L^\infty}\,\mathrm{d}t < \infty$.
- `periodicBKMIntegrability_iff_globalLifespan` (proved): `PeriodicBKMIntegrability` ν holds iff `PeriodicGlobalLifespan` ν ($T^* = \infty$ for all smooth data). The open hypothesis is therefore exactly global existence.
- Route 2 (proved from named classical hypotheses, not formalized here):
  - `integrability_of_smallDataGlobal`, from `PeriodicSmallDataGlobal` [6];

© 2026 B. S. Frohman · Apache-2.0

  - `integrability_of_prodiSerrin`, from `PeriodicProdiSerrinCriterion` [10, 12] (endpoint $L_t^\infty L_x^\infty$) plus a velocity bound.

- Route 1 (the $\kappa = 0$ analogue of the tether), `periodicBKMIntegrability_of_vorticityAPriori` (proved): from the **open** `PeriodicVorticityAPrioriBound`. It breaks because with no damping term the enstrophy satisfies only $E' \le CE^3$ and the max-point bound gives $Y' \lesssim Y^2 \log(\cdot)$. Neither gives a uniform bound.

- `periodicEnstrophyIdentity` (`EnstrophyIdentity.lean`, proved, **no hypotheses**): the $\mathbb{T}^3$ enstrophy (H^1) energy identity, the $s = 1$ case of the H^s identity (4.1).

- Small zero-mean H^1 data (`EnstrophyIdentity.lean`): `riccati_barrier_Ioo` (proved) and `periodicTstar_eq_top_of_smallH1_of_rate`, proved from the unconditional identity plus the named classical hypotheses `PeriodicEnstrophyRateBound` (Ladyzhenskaya/Poincaré trilinear bound [8]) and `PeriodicH1BlowupCriterion` (Leray [9]), neither of which was formalized at that stage. Subsequently, the trilinear bound `PeriodicH1TrilinearBound` is now proved (`periodicH1TrilinearBound`, `Ladyzhenskaya.lean`, explicit unoptimized constant, Fourier/Agmon route to a Ladyzhenskaya-type estimate; the inequality is classical, O. A. Ladyzhenskaya 1958 in 2D and her 1961 monograph in 3D [8]), as is `PeriodicMeanConservation` (`periodicMeanConservation`); hence `periodicTstar_eq_top_of_smallH1_ladyzhenskaya` depends only on the named `PeriodicH1BlowupCriterion`, which remains **not formalized**. This does not resolve the Clay problem. *Correction.* The earlier hypothesis `PeriodicEnstrophyRiccati` with $\alpha > 0$ (`SmallDataAttack.lean`, used by `periodicTstar_eq_top_of_smallH1`, `integrability_of_smallH1`) is false for small constant flows, since the mean mode is conserved; the zero-mean rate-bound version replaces it.

- Tether claim (`TetherClaimAttack.lean`): `cubic_riccati_barrier` (proved); `periodicVorticityAPrioriBound_of_tetherCubicRate`, `clay_B_exact_of_tetherCubicRate` proved from the **open** `PeriodicTetherCubicRate` (not derived from unmodified NS); `uniform_cubic_rate_false_of_scaling` (proved, abstract scaling step: a data-uniform cubic rate is contradictory).

- Direction route: `integrability_of_directionCriterion`, `periodicBKMIntegrability_of_direction`, proved from the classical `PeriodicDirectionCriterion` (Beirão da Veiga–Berselli [2] / Constantin–Fefferman [3], not formalized) plus the **open** `PeriodicDirectionRegularity`. In addition, `periodicDirectionCriterion_of_growth` proves `PeriodicDirectionCriterion` from the unconditional enstrophy identity, the linear Grönwall lemma `exp_gronwall_Ioo`, and two named open hypotheses: `PeriodicDirectionEnstrophyGrowth` (coined; Constantin–Fefferman 1993 geometric depletion of the strain kernel plus Beirão da Veiga–Berselli 2002, constant-rate form, not formalized) and `PeriodicH1BlowupCriterion` (Leray 1934). `periodicBKMIntegrability_of_directionGrowth` still needs the open `PeriodicDirectionRegularity`.

- Bending rate (`BendingRate.lean`): `hasDerivAt_vortDir_lagrangian` (proved) gives the pointwise identity

$$\frac{D\mathbf{t}}{Dt} = (I - \mathbf{t}\otimes\mathbf{t})S\mathbf{t} + \nu|\omega|^{-1}(I - \mathbf{t}\otimes\mathbf{t})\Delta\omega, \qquad \mathbf{t} = \omega/|\omega|, \tag{6.1}$$

  for any curve with $D\omega/Dt = S\omega + \nu\Delta\omega$; `inner_perpPart` shows the orthogonal strain is invisible to the stretching rate $\mathbf{t}\cdot S\mathbf{t}$. The hidden-property hypothesis `PeriodicBendingRateHiddenProperty` (coined; new formulation dated Oct 31 2026; precursors (geometric only, no regularity claims): the author's May 17/21 2026 notes (Π_u orthogonal to u,

© 2026 B. S. Frohman · Apache-2.0

"invisible to the dynamics generated by H") and Aug 10 2026 notes *The Frohmanian Symplectic Tether* (B_{anti} via $\hat{u} \times (\cdot)$, B_{sym} with pointwise Π_u)) is the conjunction of `PeriodicOrthogonalStrainControl` and `OrthogonalStrainToDirectionHolder`, both **open**; `periodicDirectionRegularity_of_bending` (proved) derives `PeriodicDirectionRegularity` from it. Classical step broken down: `abs_cfDepletion_le` (proved) bounds the Constantin–Fefferman kernel factor $(e \cdot a)\det(e, a, b)$ by $|b - a|$, and `cfDepletion_le_of_holder` by $K|x - y|^{1/2}$; `gronwall_integrating_factor` (proved) is Grönwall with rate $g = G' \in L^1$; `periodicDirectionCriterion_of_growthL1` (proved) gives `PeriodicDirectionCriterion` from the weaker open `PeriodicDirectionEnstrophyGrowthL1` (Biot–Savart stretching representation on $\mathbb{T}^3$ + BdV–B interpolation, not formalized) and `PeriodicH1BlowupCriterion`.

- Leray and analytic CF breakdown (`ClassicalSteps.lean`): `hsSq_bounded_of_rate` (proved; linear rate bound ⇒ bounded $\|u\|_{H^s}^2$ on $[0, T^*)$); `periodicH1BlowupCriterion_of_smoothing` (proved) derives `PeriodicH1BlowupCriterion` from the open parabolic-smoothing rates `PeriodicH2SmoothingRate` ($H^1 \to H^2$) and `PeriodicH3SmoothingRate` ($H^2 \to H^3$) together with the proved H^3 blow-up alternative; `periodicDirectionEnstrophyGrowthL1_of_holderStretching` (proved) derives `PeriodicDirectionEnstrophyGrowthL1` from the open `PeriodicHolderStretchingEstimate` (Biot–Savart stretching on $\mathbb{T}^3$ + BdV–B) and `PeriodicEnstrophyTimeIntegrable` (energy equality); `periodicBKMIntegrability_of_classical_bending` (conditional).

- Energy equality and the standard smoothing form (`EnergyEquality.lean`, Oct 31 2026; all names coined). `energy_hasDerivAt` (proved) is the energy equality

$$\frac{d}{dt}\|u\|_{L^2}^2 = -8\pi^2\nu\big(\|u\|_{H^1}^2 - \|u\|_{L^2}^2\big) \quad \text{on } (0, T^*), \tag{6.2}$$

from the unconditional H^s identity at $s = 0$ and `PeriodicNonlinearEnergyCancellation`, the Fourier form of $\langle (u \cdot \nabla)u, u\rangle = 0$ [9, 13]. This cancellation is now *proved* unconditionally (`EnergyCancellation.lean`, `periodicNonlinearEnergyCancellation`): on the Fourier side the Leray projector drops out of the pairing with a divergence-free field, the coefficient decay from $u \in H^8$ makes the double sum absolutely summable, and the involution $(m, j) \mapsto (m, -(m + j))$, together with Hermitian symmetry and $k \cdot \hat{u}(k) = 0$, negates each term, so the real part vanishes. Consequently `periodicEnstrophyTimeIntegrable` proves `PeriodicEnstrophyTimeIntegrable` with no hypothesis. `periodicEnstrophyTimeIntegrable_of_cancellation` (proved) then discharges `PeriodicEnstrophyTimeIntegrable`: with $E(t) = \|u(t)\|_{L^2}^2$ antitone on $[0, T^*)$ and extended to T^* by its left limit, $G(t) = \int_0^t E - E(t)/(8\pi^2\nu)$ is continuous on $[0, T^*]$ with $G' = \|u\|_{H^1}^2$. *Discrepancy with the standard estimate.* The hypothesis `PeriodicH2SmoothingRate` requires $\frac{d}{dt}\|u\|_{H^2}^2 \le K(1 + \|u\|_{H^1}^2)^m\|u\|_{H^2}^2$ pointwise. The standard 3D estimate [14, 11] gives, after absorbing the viscous H^3 term, a coefficient with one extra factor $\|u\|_{H^2}$:

$$\frac{d}{dt}\|u\|_{H^2}^2 \le \frac{C}{\nu}\|u\|_{H^1}\|u\|_{H^2} \cdot \|u\|_{H^2}^2, \tag{6.3}$$

and closes only because $\int_0^T \|u\|_{H^2}^2\, dt < \infty$ when $\|u\|_{H^1}$ is bounded. The pointwise form is not known to hold. It is retained in the development, marked accordingly, and superseded by `PeriodicH2SmoothingRateStd` (the display above) and `PeriodicH2RootTimeIntegrable` ($t \mapsto \|u\|_{H^2}$ has a continuous primitive on $[0, T^*]$ when $\|u\|_{H^1}$ is bounded), both open; `periodicH1BlowupCriterion_of_smoothingStd` (proved) gives `PeriodicH1BlowupCriterion` from these two and `PeriodicH3SmoothingRate`.

© 2026 B. S. Frohman · Apache-2.0

The H^3 Prop matches the standard estimate ($m = 1$, via $|\langle [J^3, u \cdot \nabla]u, J^3 u\rangle| \lesssim \|u\|_{H^2}\|u\|_{H^3}\|u\|_{H^4}$). `periodicBKMIntegrability_of_classical_bending_std` (conditional) assembles the chain.

- $\kappa = 0$ tether (`TetherKappaZero.lean`): `tetherForm_zero`, `tetherGeometricControl_zero`, `tetherImpliesDirection_zero_iff` (proved): at $\kappa = 0$ the tether control is vacuous, so "tether control ⇒ direction regularity" is *equivalent* to the open `PeriodicDirectionRegularity`; `periodicBKMIntegrability_of_tetherKappaZero` (conditional).

7 Status

- **Proved, unconditional:** local existence ($\nu \geq 0$), uniqueness, coefficient ODE and decay, H^s energy identity, Parseval, physical form of the H^3 rate, Kato–Ponce toolkit including `lower_term_bound` and `top_term_zero`, the $s = 3$ commutator bound `periodicPhysicalCommutatorBound`, the blow-up alternative, the low-frequency bound `periodicLowFreqVorticityBound`, H^3 BKM control `periodicBKMHsControl_three` and the BKM criterion `periodicBKMCriterion`, the equivalence `periodicBKMIntegrability_iff_globalLifespan`, and the H^1 enstrophy identity `periodicEnstrophyIdentity`.
- **Conditional:** Statement (B) is proven conditionally on `PeriodicBKMIntegrability` (`clay_B_exact_of_integrability`). Small zero-mean H^1 data give $T^* = \infty$ conditionally on `PeriodicEnstrophyRateBound` and `PeriodicH1BlowupCriterion`; at $\kappa = 0$ the tether headline is equivalent to the open `PeriodicDirectionRegularity`; `periodicBKMIntegrability_of_bending` gives integrability from `PeriodicDirectionEnstrophyGrowthL1`, `PeriodicH1BlowupCriterion` and the open `PeriodicBendingRateHiddenProperty`.
- **Direction route, L^1 form:** `periodicDirectionCriterion_of_growthL1` gives `PeriodicDirectionCriterion` from `PeriodicDirectionEnstrophyGrowthL1` and `PeriodicH1BlowupCriterion`; these in turn reduce (proved) to `PeriodicHolderStretchingEstimate`, `PeriodicH2SmoothingRateStd`, `PeriodicH2RootTimeIntegrable` and `PeriodicH3SmoothingRate`, all open here (`PeriodicNonlinearEnergyCancellation` and `PeriodicEnstrophyTimeIntegrable` are proved) (the older `PeriodicH2SmoothingRate` is stronger than the standard estimate; see above). The bending-rate hypothesis `PeriodicBendingRateHiddenProperty` (coined Oct 31 2026, open) implies `PeriodicDirectionRegularity`; the identity $D\mathbf{t}/Dt = (I - \mathbf{t}\otimes\mathbf{t})S\mathbf{t} + \nu|\omega|^{-1}(I - \mathbf{t}\otimes\mathbf{t})\Delta\omega$ is proved.
- **Open:** `PeriodicBKMIntegrability` — the Clay Millennium core. Neither the Clay problem nor global regularity of 3D Navier–Stokes is claimed.

Reproducibility

Run `lake build` and then `lake env lean AxiomAudit.lean`. The build contains no `sorry`, and the audit reports only `propext`, `Classical.choice` and `Quot.sound`.

Counting convention. Two counts are reported, each with its method. (i) *Source count:* 755 declarations introduced by the keywords `theorem` or `lemma` (optionally preceded by attributes or modifiers) in the tracked `.lean` files of this repository, excluding `AxiomAudit.lean`. (ii) *Kernel count:* 867 non-internal theorem constants in the `PeriodicNSContinuation` module tree, as reported by `AxiomAudit.lean`; this includes theorems generated automatically by Lean. An earlier figure of 727 was not obtained by either method and is withdrawn; the

© 2026 B. S. Frohman · Apache-2.0

figure 726 used in versions ≤ 8 was the source count before `EnergyCancellation.lean` was added.

License

© 2026 Benjamin Stanley Frohman (ORCID 0009-0006-7068-3718). Licensed under the Apache License, Version 2.0.

References

[1] J. T. Beale, T. Kato, and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Comm. Math. Phys. **94** (1984), no. 1, 61–66.

[2] H. Beirão da Veiga and L. C. Berselli, *On the regularizing effect of the vorticity direction in incompressible viscous flows*, Differential Integral Equations **15** (2002), no. 3, 345–356.

[3] P. Constantin and C. Fefferman, *Direction of vorticity and the problem of global regularity for the Navier–Stokes equations*, Indiana Univ. Math. J. **42** (1993), no. 3, 775–789.

[4] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Comput. Sci., vol. 12699, Springer, 2021, pp. 625–635.

[5] C. L. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, Clay Mathematics Institute, Millennium Prize Problems, 2000.

[6] H. Fujita and T. Kato, *On the Navier–Stokes initial value problem. I*, Arch. Rational Mech. Anal. **16** (1964), 269–315.

[7] T. Kato and G. Ponce, *Commutator estimates and the Euler and Navier–Stokes equations*, Comm. Pure Appl. Math. **41** (1988), no. 7, 891–907.

[8] O. A. Ladyzhenskaya, *The mathematical theory of viscous incompressible flow*, Gordon and Breach, New York, 1963, Russian original: Fizmatgiz, Moscow, 1961.

[9] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math. **63** (1934), 193–248.

[10] G. Prodi, *Un teorema di unicità per le equazioni di Navier–Stokes*, Ann. Mat. Pura Appl. (4) **48** (1959), 173–182.

[11] J. C. Robinson, J. L. Rodrigo, and W. Sadowski, *The three-dimensional Navier–Stokes equations: Classical theory*, Cambridge Stud. Adv. Math., vol. 157, Cambridge Univ. Press, Cambridge, 2016.

[12] J. Serrin, *The initial value problem for the Navier–Stokes equations*, Nonlinear Problems (Proc. Sympos., Madison, Wis., 1962), Univ. of Wisconsin Press, Madison, 1963, pp. 69–98.

[13] R. Temam, *Navier–Stokes equations: Theory and numerical analysis*, North-Holland, Amsterdam, 1977.

[14] ———, *Navier–Stokes equations and nonlinear functional analysis*, 2 ed., CBMS-NSF Regional Conf. Ser. in Appl. Math., vol. 66, SIAM, Philadelphia, 1995.

[15] The mathlib Community, *The Lean mathematical library*, Proc. 9th ACM SIGPLAN Int. Conf. Certified Programs and Proofs (CPP 2020), 367–381, 2020.

© 2026 B. S. Frohman · Apache-2.0