Dean's Office, Faculty of Science 2004 TEST AND ANSWERS

ANSWER SHEET

FORM CC1

SECTION C : ANSWERS

USE 2B PENCILS ONLY

INSTRUCTIONS

Suggested answers to each question are given in the question paper. Choose an answer and shade the corresponding circle.

EXAMPLES OF SHADING

CORRECT

INCORRECT

8 0

1

1

SECTION A STUDENT'S NAME

MA 1505

MODULE:

YEAR/SEMESTER: 2004

DATE:

SECTION B MATRICULATION NUMBER

000000000000 G 🗷 🛢 🖯 🗷 🗷 🗷 🖯 🖂 🗎 🖂 🗷 0000000000000

> 2. Now SHADE the the grid for each digit or corresponding circle in

Si 82828282828282

8

® 2 ® 2 ® 2 ® 2 ® 2 ® 2 ® 2 ® 2

1 2 73 (A) (B) (B) 2 74 (A) (B) (B) 2 75 (A) (B) (B) 2 76 (A) (B) (B) 2 77 (A) (B) (B) 2 78 (A) (B) (B) 2 80 (A) (B) (B) (B)

4 0 4 0 4 0 4 0 4 0 4 0 4 0 m 5 m 5 m 5 m 5 m 5 m 5 m 5 m 5 m 5 m

65

67

8

1. Write your matriculation

<u>5</u> 7	င္သာ		52	5			10		9		00		7		on.		C)		4		ω		2		_	
3 -	(B)	_	(A) -					_	(4	(-	(_	(_	(_	(2)	_		_	(_	(_
® №													•													
(i) a													0													
4 📵	6111111111111		100000000000000000000000000000000000000	12 12 12 12 12									ŏ													
(m) 4													<u>—</u>													
		-			٠.						-		0			Til		77	-	Min	0	Till.	0	77777	0	-
cm ·	con .		6	6			N		_		_		_		-		_		_		-		_		_	
64							0		9 (8		17 A		6		5		4		3	EU.	2	uun.	0	
(2)							(A)		0		0		0	7	0		0		0		0		-			_
® №	(00)	2	(B) N	(B)	100	- 1							B													
(C) 64													0													
4 🖯																										
(m) cs	•	C)	(m) cr	•	OT .		m	cn	m	C)	m	cn	m	CT	•	CI		CII		CTI	(m)	C)		CIT	(m)	បា
74	73		72	71			30		29		28		27		26		25		24		23		22		2	
> -	(D)	_	D -	((2)	-	(3)	-	(3)	-	3	_	(2)	-	(_	(2)	-	(-	9	-	(_
@ N													®													
(i) w													0													
4 📵													0													
(m) u													<u></u>													
		7		0			0		0		0		0		0		0		0		0		0		0	
																						*				
20	co		00	00			4		ယ		డు		37		ట		ယ		cu		ట	ħ	ట		ယ	
																										-
(E) -1	0												(2)													
® №													®													
@ w	111111111111111111111111111111111111111		11.7000177	17 F (50 75 C									0													
a													0													
m •	(m)	CI	(m) a		CT			CI		CII		CI	m	CII		GI	(m)	OI .	(17)	CI		CT		CII		CI
22							50										5									
> -													(2)													
® N		N	® №		N								•													
@ ca	0	w	(F) 64	0	ယ		0	ω	0	ω	0	ω	0	ω	0	ယ	0	çu	0	cu	0	cu	0	ယ	0	ω
4 😑																										
(m) u													(m)													
									_																	

1 2 3 4 A B C D	2 3 4 B C D	1 2 3 4 A B C D	1 2 3 4 A B C D	1 2 3 4 5 86 A B C D E	1 2 3 4 A B C D	1 2 3 4 A B C D	B C D	
B 2 3	B 2 3	B 2 3	B 0	1 2 3 4 96 (A) (B) (C) (D)	B 2 3	B 2 3	B 2 3	

```
1. hy=xhx \Rightarrow y'(y)=1+hx \Rightarrow y'=x^{2}(1+hx)
                           y'=0 \iff 1+\ln x=0, i.e. \ln x=-1 or x=e^{-1}
      2. 3x^2 + 3y^2y' = 6(y + xy'). at (3,3), 3(9) + 27y' = 6(3+3y')

9y' = -9 \Rightarrow y' = -1, tangent line: \frac{y-3}{x-3} = -1 \Rightarrow y-3+x-3=0
       3. dx(sin'x cosnx) = nsin'-1x cos x cos nx +(sin'x)(-nsinnx)
                  = n sinn-1 x (cos nx cos x - sin nx sinx) = n sinn-1x cos (n+1)x
     4. f(x) = 1 + \frac{9}{x^2 q} \implies f'(x) = -18(x^2 - 9)^2 x
                f''(x) = -18 \left[ (x^2 - 9)^2 + x(-2)(x^2 - 9)^3 (2x) \right]
               f''(9) = -18(81-9)^{-3} [72 - 4(81)] = \frac{252(18)}{72^{2}(72)} = \frac{63}{72(72)} = \frac{7}{576}
   5. let y^{\pm} = 3 + x, \lim_{y \to 3^{\pm}} \frac{y - 2y^{\pm} - 3^{\pm} + b}{y^{\pm} - 3} = \lim_{y \to 3^{\pm}} \frac{1 - 8y^{3}}{4y^{3}} = \lim_{y \to 3^{\pm}} \frac{1}{4y^{3}} = 0
6. f'(x) = 6(x-2)(x+1) = 0 at x = -1, 2, f(-2) = 1, f(-1) = 12
-15 = f(2) \le f(-2) \le f(-1) \le f(4) = 37
    7. 6\int \frac{6x}{\sqrt{3}x^2+5} dx = 6\int (3x^2+5)^{\frac{1}{2}} d(3x^2+5) = 6(2)(3x^2+5)^{\frac{1}{2}} + C
      8. d = (\sin \sqrt{x^4}) \int_0^{x^4} \sin \sqrt{t} dt \frac{d(x^4)}{dx} = (\sin \sqrt{x^4}) 4x^3 = (\sin |x^2|) 4x^3
      9. Since Jsin x JI-sin x = Jsin x / 45 x = Jsh(J-x) / Gr (J-x) on EO, T.
                \int_{0}^{\pi} \sqrt{\sin x} |u_{n} x| dx = 2 \int_{0}^{\frac{\pi}{2}} (\sin x)^{\frac{1}{2}} d(\sin x) = \frac{4}{3} (\sin x)^{\frac{3}{2}} \Big|_{0}^{\frac{\pi}{2}} = \frac{4}{3}
   10. In(wxx) d(tanx) = In(wxx) tonx- 5 tonx (-sinx) dx
                 = \tan x \ln w x + \int sec^2 x - 1 dx = (\tan x)(\ln w x + 1) - x + c
    11. y' = 4x(x^2-2) = 0 at x = 0, \pm \sqrt{2}, y'(\pm \sqrt{2}) -2 - \sqrt{2}

Required Area = 2\int_0^{\sqrt{2}} |y_1 - y_2| dx

= 2\int_0^{\sqrt{2}} x^4 - 4x^2 + 4 dx = 2\left[\frac{x^5}{5} - \frac{4x^3}{3} + 4x\right]_0^{\sqrt{2}}
     2. y'=xe^{x}+e^{x}>0 for x>0

Volume of typical disk=\pi(y-o)^{2}\Delta x

Required Volume=\int_{0}^{1}\pi x^{2}e^{2x}dx=\frac{\pi}{2}\int_{0}^{1}x^{2}d(e^{2x})

=\frac{\pi}{2}[x^{2}e^{2x}]'-[\frac{1}{2}e^{2x}](x^{2})
= 2\sqrt{2} \left[ \frac{4}{5} - \frac{8}{3} + 4 \right] = 2\sqrt{2} \left( \frac{32}{15} \right)
12. y' = xe^{x} + e^{x} > 0 for x \ge 0

Volume of typical disk = \pi(y-0)^{2}\Delta x
       = \underline{\mathcal{I}} \left[ x^{2} e^{2x} \Big|_{o}^{1} - \int_{o}^{1} e^{2x} (2x) dx \right] = \underline{\mathcal{I}} e^{2} - \underline{\mathcal{I}} \int_{o}^{1} x d(e^{2x}) dx = \underline{\mathcal{I}} e^{2} - \underline{\mathcal{I}} \int_{o}^{1} x e^{2x} \Big|_{o}^{1} - \int_{o}^{1} e^{2x} dx = \underline{\mathcal{I}} e^{2} - \underline{\mathcal{I}} e^{2} + \underline{\mathcal{I}} (e^{2x}) \Big|_{o}^{1} = \underline{\mathcal{I}} (e^{2} - e^{2}) dx = \underline{\mathcal{I}} e^{2} - \underline{\mathcal{I}} (e^{2x}) \Big|_{o}^{1} = \underline{\mathcal{I}} (e^{2x}) e^{2x} dx = \underline{\mathcal{I}} (e^{2x}) e^{2x} dx
```

- 1. Let $f(x) = x^x$, x > 0. Then f'(x) = 0 at
 - (A) x = e
 - **(B)** $x = e^{-1}$
 - (C) x=1
 - $(\mathbf{D}) \quad x = e^{-2}$
 - $(\mathbf{E}) \quad x = e^2$
- 2. Find the equation of the tangent to the curve $x^3 + y^3 = 6xy$ at the point (3,3).
 - $(\mathbf{A}) \quad x y = 0$
 - **(B)** x 2y + 3 = 0
 - (C) 2x y + 3 = 0
 - **(D)** x + y 6 = 0
 - **(E)** x + 2y 9 = 0
- $3. \ \frac{d}{dx} \left(\sin^n x \cos nx \right) =$
 - **(A)** $n \sin^{n+1} x \cos(n+1) x$
 - (B) $n\sin^{n-1}x\cos(n-1)x$
 - (C) $n\sin^{n-1}x\sin(n+1)x$
 - **(D)** $n\sin^{n-1}x\sin(n-1)x$
 - (E) $n \sin^{n-1} x \sin nx$

4. Let f(x) be a function defined by

$$f\left(x\right) = \frac{x^2}{x^2 - 9}$$

where $x \in [5, 10]$. Then f''(9) =

- (A) $\frac{3}{864}$
- **(B)** $\frac{7}{864}$
- (C) $\frac{3}{576}$
- **(D)** $\frac{7}{576}$
- (\mathbf{E}) ∞

5. Evaluate $\lim_{x\to 0} \frac{(3+x)^{1/4}-2(3+x)-(3)^{1/4}+6}{x}$.

- (A) The limit does not exist
- **(B)** $\frac{1}{4(27)^{1/4}}$
- (C) $\frac{1}{4(3)^{1/4}} 2$
- (D) $\frac{1}{4(3)^{1/4}}$
- (E) $\frac{1}{4(27)^{1/4}} 2$

6. Let $f(x) = 2x^3 - 3x^2 - 12x + 5$, $x \in [-2, 4]$. Let M and m denote the absolute maximum value and absolute minimum value of f respectively. Then

3

- (A) M = 12, m = 1
- **(B)** M = 37, m = -15
- (C) M = 15, m = 0
- **(D)** M = 40, m = -12
- **(E)** M = 45, m = 5

$$7. \int \frac{x}{\sqrt{3x^2+5}} \mathrm{d}x =$$

- (A) $\frac{1}{9} (3x^2 + 5)^{3/2} + C$
- **(B)** $\frac{1}{4} (3x^2 + 5)^{3/2} + C$
- (C) $\frac{1}{12} (3x^2 + 5)^{1/2} + C$
- **(D)** $\frac{1}{3} (3x^2 + 5)^{1/2} + C$
- **(E)** $\frac{3}{2} (3x^2 + 5)^{1/2} + C$

$$8. \ \frac{d}{dx} \int_0^{x^4} \sin \sqrt{t} dt =$$

- (A) $\sin x^4$
- (B) $-\cos x^2$
- (C) $x^4 \sin x^2$
- **(D)** $1 \cos x^2$
- (E) $4x^3 \sin x^2$

9.
$$\int_0^{\pi} \sqrt{\sin x - \sin^3 x} dx =$$

- (A) $\frac{3}{2}$
- **(B)** $\frac{2}{3}$
- **(C)** 0
- (D) $\frac{3}{4}$
- (E) $\frac{4}{3}$

10.
$$\int \frac{\ln(\cos x)}{\cos^2 x} dx =$$

- (A) $\tan x (1 + \ln(\cos x)) + \sin x + C$
- (B) $\tan x (1 + \ln(\cos x)) \sin x + C$
- (C) $\tan x (1 + \ln(\cos x)) + x + C$
- **(D)** $\tan x (1 + \ln(\cos x)) x + C$
- **(E)** $\tan x (1 + \ln(\cos x)) + C$
- 11. Find the area of the region bounded by the curves $y = x^4 4x^2$ and y = -4.
 - **(A)** $\frac{8}{3}\sqrt{2}$
 - **(B)** $\frac{16}{5}\sqrt{2}$
 - (C) $\frac{64}{15}\sqrt{2}$
 - **(D)** $\frac{96}{25}\sqrt{2}$
 - **(E)** $\frac{128}{75}\sqrt{2}$
- 12. Let R denote the region in the first quadrant bounded by $y = xe^x$, y = 0 and x = 1. Find the volume generated by revolving the region R about the x-axis.
 - **(A)** $\frac{\pi}{4} \left(e^2 1 \right)$
 - **(B)** $\frac{\pi}{4}e^2$
 - (C) $\frac{\pi}{4} \left(1 \frac{1}{e^2}\right)$
 - **(D)** $\frac{\pi}{4} \left(e^2 2 \right)$
 - (E) $\frac{\pi}{4} \left(e^2 \frac{1}{2} \right)$

END OF PAPER