

Tuple Relational Calculus

1. Find the names of pizzas that come in a 10 inch size

$$\{T \mid \exists T1 \\ (T1 \in \text{pizza} \wedge T1.\text{size} = 10 \\ \wedge T1.\text{name} = T.\text{name})\}$$

2. Find the names of pizzas that come in a 10 inch or a 12 inch size

$$\{T \mid \exists T1 \\ (T1 \in \text{pizza} \wedge (T1.\text{size} = 10 \vee T1.\text{size} = 12) \\ \wedge T1.\text{name} = T.\text{name})\}$$

3. Find the names of pizzas that come in both a 10 inch and a 12 inch size

$$\{T \mid \exists T1 \exists T2 \\ (T1 \in \text{pizza} \wedge T2 \in \text{pizza} \wedge T1.\text{name} = T2.\text{name} \wedge T1.\text{size} = 10 \wedge T2.\text{size} = 12 \\ \wedge T1.\text{name} = T.\text{name})\}$$

4. Find the pairs of different codes of pizzas with the same name and the same size (is there any?)

$$\{T \mid \exists T1 \exists T2 \\ (T1 \in \text{pizza} \wedge T2 \in \text{pizza} \wedge T1.\text{code} \neq T2.\text{code} \wedge T1.\text{name} = T2.\text{name} \wedge T1.\text{size} = \\ T2.\text{size} \\ \wedge T.\text{code1} = T1.\text{code} \wedge T.\text{code2} = T2.\text{code})\}$$

Yes there some possibly {code} is the key, not {name, size}

5. Find the names and phone numbers of the stores in "College Park" or "Greenbelt" that sell a 10 inch pizza named "pepperoni" for less than \$8

$$\{T \mid \exists T1 \exists T2 \exists T3 \\ (T1 \in \text{pizza} \wedge T2 \in \text{store} \wedge T3 \in \text{sells} \wedge T1.\text{code} = T3.\text{code} \wedge T2.\text{name} = T3.\text{store_name} \wedge \\ (T2.\text{area} = \text{« College Park »} \vee T2.\text{area} = \text{« Greenbelt »}) \wedge T1.\text{name} = \text{“pepperoni”} \wedge T1.\text{size} = 10 \\ \wedge T3.\text{price} < 8 \\ \wedge T2.\text{name} = T.\text{name} \wedge T2.\text{phone} = T.\text{phone})\}$$

6. Find the codes of the most expensive pizzas – assume the scheme of the database is reduced to a relation pizza(code, price) to simplify –

$$\{T \mid \exists T1 \forall T2 \\ (T1 \in \text{pizza} \wedge (T2 \in \text{pizza} \Rightarrow T1.\text{price} \geq T2.\text{price}) \\ \wedge T1.\text{code} = T.\text{code})\}$$

7. Find the names of the stores that sell all the pizzas

$$\{T \mid \exists T1 \forall T2 \exists T3 \\ (T1 \in \text{store} \wedge (T2 \in \text{pizza} \Rightarrow (T3 \in \text{sells} \wedge T2.\text{code} = T3.\text{code} \wedge T1.\text{name} = T3.\text{store_name})) \\ \wedge T1.\text{name} = T.\text{name})\}$$