

EE3206/EE3206E INTRODUCTION TO COMPUTER VISION AND IMAGE PROCESSING

Tutorial Set F – Solution



Question 1

Part (a)

Consider a rectangle with sides a, a, ka, ka . The compactness is

$$\begin{aligned}
 \gamma &= \frac{P^2}{4\pi A} \\
 &= \frac{(2ka + 2a)^2}{4\pi \times ka^2} \\
 &= \frac{4(k+1)^2}{4\pi k} \\
 &= \frac{(k+1)^2}{\pi k} \quad (1)
 \end{aligned}$$


The minimum value of γ is $4/\pi$ when $k = 1$, i.e., a square.



Part (b)

$$\begin{aligned}
 \gamma &= \frac{\pi^2 \times 2(a^2 + b^2)}{4\pi \times \pi ab} \\
 &= \frac{a^2 + b^2}{2ab} \\
 &= \frac{1 + \epsilon^2}{2\epsilon}, \quad \text{where } \epsilon = \frac{a}{b} \geq 1
 \end{aligned} \tag{2}$$

Minimum value of γ is 1, when $\epsilon = 1$, i.e., a circle.



Question 2

Procedure

1. Find a suitable threshold, T , by determining the valley between the two modes, or use the inter-means algorithm.
2. Threshold image at gray level T :


$$g(x, y) = \begin{cases} 1 & \text{if } f(x, y) > T \\ 0 & \text{if } f(x, y) \leq T. \end{cases} \quad (3)$$

3. Apply component labelling algorithm to $g(x, y)$ with $V = 1$.
4. This will result in 5 connected components labelled A to E . (As an additional step, small regions due to noise may be discarded by imposing a minimum-area requirement, or by using morphology.)
5. For each R_i , compute m_{00} , m_{01} and m_{10} to obtain the centroid (\bar{x}, \bar{y}) :

$$\bar{x} = \frac{m_{10}}{m_{00}} \quad \bar{y} = \frac{m_{01}}{m_{00}}$$


For example, for Region A :

$$m_{pq} = \sum_{(x,y) \in A} x^p y^q f(x, y) \quad (4)$$



Question 3

Part (a)



$$m_{pq} = \iint x^p y^q f(x, y) dx dy = \iint_{R_1, R_2} x^p y^q dx dy = \iint_{R_1} x^p y^q dx dy + \iint_{R_2} x^p y^q dx dy$$

$$f(x, y) = \begin{cases} 1 & (x, y) \in R_1, R_2 \\ 0 & \text{elsewhere} \end{cases}$$

Take the origin to be the lower left corner.

$$m_{00} = \text{area of object} = 16$$

$$\begin{aligned} m_{10} &= \iint x dx dy = \int_0^6 \left\{ \int_0^2 x dx \right\} dy + \int_0^2 \left\{ \int_2^4 x dx \right\} dy \\ &= \int_0^6 [x^2/2]_0^2 dy + \int_0^2 [x^2/2]_2^4 dy \\ &= \int_0^6 2 dy + \int_0^2 6 dy \\ &= 12 + 12 \\ &= 24 \end{aligned}$$

$$\begin{aligned} m_{01} &= \iint y dx dy = \int_0^6 \left\{ \int_0^2 y dx \right\} dy + \int_0^2 \left\{ \int_2^4 y dx \right\} dy \\ &= \int_0^6 2y dy + \int_0^2 2y dy \\ &= [y^2]_0^6 + [y^2]_0^2 \\ &= 36 + 4 \\ &= 40 \end{aligned}$$

$$\bar{x} = m_{10}/m_{00} = 1.5$$

$$\bar{y} = m_{01}/m_{00} = 2.5$$

$$\begin{aligned}
\mu_{11} &= \iint (x - 1.5)(y - 2.5) \, dx \, dy \\
&= \int_0^6 \left\{ \int_0^2 (x - 1.5)(y - 2.5) \, dx \right\} dy + \int_0^2 \left\{ \int_2^4 (x - 1.5)(y - 2.5) \, dx \right\} dy \\
&= -3 - 9 \\
&= -12
\end{aligned}$$

$$\begin{aligned}
\eta_{11} &= \mu_{11}/\mu_{00}^2 \\
&= -12/16^2 \\
&= -0.0469
\end{aligned}$$

Alternatively, use

$$\mu_{11} = m_{11} - \bar{y}m_{10}$$

We have

$$\begin{aligned}
m_{11} &= \iint xy \, dx \, dy = \int_0^6 \left\{ \int_0^2 xy \, dx \right\} dy + \int_0^2 \left\{ \int_2^4 xy \, dx \right\} dy \\
&= \int_0^6 [x^2/2]_0^2 y \, dy + \int_0^2 [x^2/2]_2^4 y \, dy \\
&= \int_0^6 2y \, dy + \int_0^2 6y \, dy \\
&= [y^2]_0^6 + [3y^2]_0^2 \\
&= 36 + 12 \\
&= 48
\end{aligned}$$


$$\begin{aligned}
\mu_{11} &= 48 - 2.5 \times 24 \\
&= -12
\end{aligned}$$

$$\begin{aligned}
\eta_{11} &= \mu_{11}/\mu_{00}^2 \\
&= -12/16^2 \\
&= -0.0469
\end{aligned}$$

$$\begin{aligned}
m_{20} &= \iint x^2 dx dy = \int_0^6 \left\{ \int_0^2 x^2 dx \right\} dy + \int_0^2 \left\{ \int_2^4 x^2 dx \right\} dy \\
&= 16 + \frac{112}{3} \\
&= 53.3333 \\
m_{02} &= \iint y^2 dx dy = \int_0^6 \left\{ \int_0^2 y^2 dx \right\} dy + \int_0^2 \left\{ \int_2^4 y^2 dx \right\} dy \\
&= \frac{432}{3} + \frac{16}{3} \\
&= 149.3333
\end{aligned}$$

$$\begin{aligned}
\mu_{20} &= m_{20} - \bar{x}m_{10} \\
&= 53.3333 - 1.5 \times 24 = 17.3333 \\
\mu_{02} &= m_{02} - \bar{y}m_{01} \\
&= 149.3333 - 2.5 \times 40 = 49.3333 \\
\eta_{20} &= \mu_{20}/\mu_{00}^2 = 0.0677 \\
\eta_{02} &= \mu_{02}/\mu_{00}^2 = 0.1927 \\
\phi_1 &= \eta_{20} + \eta_{02} \\
&= 0.26
\end{aligned}$$

Part (b)



Choose origin at bottom left pixel.

$$m_{00} = 16$$

$$\begin{aligned} m_{10} &= \sum \sum x^1 y^0 f(x, y) = \sum \sum x f(x, y) = \sum_{x=0}^3 x \left\{ \sum_{y=0}^5 f(x, y) \right\} \\ &= 0 \times \sum_{y=0}^5 f(x, y) + 1 \times \sum_{y=0}^5 f(x, y) + 2 \times \sum_{y=0}^5 f(x, y) + 3 \times \sum_{y=0}^5 f(x, y) \\ &= (0 \times 6) + (1 \times 6) + (2 \times 2) + (3 \times 2) \\ &= 16 \end{aligned}$$

$$\begin{aligned} m_{01} &= \sum \sum x^0 y^1 f(x, y) = \sum \sum y f(x, y) = \sum_{y=0}^5 y \left\{ \sum_{x=0}^3 f(x, y) \right\} \\ &= (0 \times 4) + (1 \times 4) + (2 \times 2) + (3 \times 2) + (4 \times 2) + (5 \times 2) \\ &= 32 \end{aligned}$$

$$\bar{x} = m_{10}/m_{00} = 1$$

$$\bar{y} = m_{01}/m_{00} = 2$$

$$\begin{aligned} \mu_{11} &= \sum \sum (x - 1)(y - 2) f(x, y) = \sum_y (y - 2) \left\{ \sum_x (x - 1) f(x, y) \right\} \\ &= (-2)(-1 + 0 + 1 + 2) + (-1)(-1 + 0 + 1 + 2) + (0)(-1 + 0) \\ &\quad + (1)(-1 + 0) + (2)(-1 + 0) + (3)(-1 + 0) \\ &= -12 \end{aligned}$$

$$\eta_{11} = \mu_{11}/\mu_{00}^2 = -0.0469$$

$$\begin{aligned}
m_{20} &= \sum \sum x^2 f(x, y) = \sum_x x^2 \left\{ \sum_y f(x, y) \right\} \\
&= (0^2 \times 6) + (1^2 \times 6) + (2^2 \times 2) + (3^2 \times 2) \\
&= 32
\end{aligned}$$

$$\begin{aligned}
m_{02} &= \sum \sum y^2 f(x, y) = \sum_y y^2 \left\{ \sum_x f(x, y) \right\} \\
&= (0^2 \times 4) + (1^2 \times 4) + (2^2 \times 2) + (3^2 \times 2) + (4^2 \times 2) + (5^2 \times 2) \\
&= 112
\end{aligned}$$

$$\begin{aligned}
\mu_{20} &= m_{20} - \bar{x}m_{10} \\
&= 32 - 1 \times 16 = 16
\end{aligned}$$

$$\begin{aligned}
\mu_{02} &= m_{02} - \bar{y}m_{01} \\
&= 112 - 2 \times 32 = 48
\end{aligned}$$

$$\begin{aligned}
\eta_{20} &= \mu_{20}/\mu_{00}^2 \\
&= 16/16^2 = 0.0625
\end{aligned}$$


$$\begin{aligned}
\eta_{02} &= \mu_{02}/\mu_{00}^2 \\
&= 48/16^2 = 0.1875
\end{aligned}$$

$$\begin{aligned}
\phi_1 &= \eta_{20} + \eta_{02} \\
&= 0.25
\end{aligned}$$

Question 4

Part (a)

$$\begin{array}{cccccc}
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 \dots & 0 & 0 & 1 & 1 & 2 & 2 & \dots \\
 \dots & 0 & 0 & 1 & 1 & 2 & 2 & \dots \\
 \dots & 2 & 2 & 0 & 0 & 1 & 1 & \dots \\
 \dots & 2 & 2 & 0 & 0 & 1 & 1 & \dots \\
 \dots & 1 & 1 & 2 & 2 & 0 & 0 & \dots \\
 \dots & 1 & 1 & 2 & 2 & 0 & 0 & \dots \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots
 \end{array}$$



Texture P

$$\begin{array}{cc}
 \uparrow & \uparrow \\
 \mathbf{A}_{0,1} = \begin{bmatrix} 6 & 0 & 6 \\ 6 & 6 & 0 \\ 0 & 6 & 6 \end{bmatrix} & \mathbf{A}_{0,2} = \begin{bmatrix} 0 & 0 & 12 \\ 12 & 0 & 0 \\ 0 & 12 & 0 \end{bmatrix} \\
 \mathbf{C}_{0,1} = \begin{bmatrix} \frac{1}{6} & 0 & \frac{1}{6} \\ \frac{1}{6} & \frac{1}{6} & 0 \\ 0 & \frac{1}{6} & \frac{1}{6} \end{bmatrix} & \mathbf{C}_{0,2} = \begin{bmatrix} 0 & 0 & \frac{1}{3} \\ \frac{1}{3} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \end{bmatrix}
 \end{array}$$


$$\begin{aligned}
 D &= \sum_{i=1}^3 \sum_{j=1}^3 (i-j)^2 c_{ij} \\
 &= (1-1)^2 c_{11} + (1-2)^2 c_{12} + (1-3)^2 c_{13} \\
 &\quad + (2-1)^2 c_{21} + (2-2)^2 c_{22} + (2-3)^2 c_{23} \\
 &\quad + (3-1)^2 c_{31} + (3-2)^2 c_{32} + (3-3)^2 c_{33}
 \end{aligned}$$

$$\text{For } \mathbf{C}_{0,1} : \quad D = 2^2 \times \frac{1}{6} + 1^2 \times \frac{1}{6} + 1^2 \times \frac{1}{6} = 1.0$$

$$\text{For } \mathbf{C}_{0,2} : \quad D = 2^2 \times \frac{1}{3} + 1^2 \times \frac{1}{3} + 1^2 \times \frac{1}{3} = 2.0$$

$$\begin{array}{cccccc}
& \vdots & \vdots & \vdots & \vdots & \vdots \\
\dots & 0 & 1 & 2 & 0 & 1 & 2 & \dots \\
\dots & 1 & 2 & 0 & 1 & 2 & 0 & \dots \\
\dots & 2 & 0 & 1 & 2 & 0 & 1 & \dots \\
\dots & 0 & 1 & 2 & 0 & 1 & 2 & \dots \\
\dots & 1 & 2 & 0 & 1 & 2 & 0 & \dots \\
\dots & 2 & 0 & 1 & 2 & 0 & 1 & \dots \\
& \vdots & \vdots & \vdots & \vdots & \vdots & \vdots
\end{array}$$

Texture Q



$$\swarrow \mathbf{A}_{-1,1} = \begin{bmatrix} 0 & 0 & 12 \\ 12 & 0 & 0 \\ 0 & 12 & 0 \end{bmatrix}$$

$$\mathbf{C}_{-1,1} = \begin{bmatrix} 0 & 0 & \frac{1}{3} \\ \frac{1}{3} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \end{bmatrix}$$

$$\northeast \mathbf{A}_{1,1} = \begin{bmatrix} 12 & 0 & 0 \\ 0 & 12 & 0 \\ 0 & 0 & 12 \end{bmatrix}$$


$$\mathbf{C}_{1,1} = \begin{bmatrix} \frac{1}{3} & 0 & 0 \\ 0 & \frac{1}{3} & 0 \\ 0 & 0 & \frac{1}{3} \end{bmatrix}$$

$$\begin{aligned}
D &= \sum_{i=1}^3 \sum_{j=1}^3 (i-j)^2 c_{ij} \\
&= (1-1)^2 c_{11} + (1-2)^2 c_{12} + (1-3)^2 c_{13} \\
&+ (2-1)^2 c_{21} + (2-2)^2 c_{22} + (2-3)^2 c_{23} \\
&+ (3-1)^2 c_{31} + (3-2)^2 c_{32} + (3-3)^2 c_{33}
\end{aligned}$$

$$\text{For } \mathbf{C}_{-1,1} : \quad D = 2^2 \times \frac{1}{3} + 1^2 \times \frac{1}{3} + 1^2 \times \frac{1}{3} = 2.0$$

$$\text{For } \mathbf{C}_{1,1} : \quad D = 0$$

Part (b)



The n th moment of a about its mean, m is

$$\mu_n(a) = \sum_{i=1}^K (a_i - m)^n p(a_i) \quad \text{where} \quad m = \sum_{i=1}^K a_i p(a_i)$$

For the resulting histograms $h_{g,P}$ and $h_{g,Q}$, we compute m and μ_2 (the variance).

	m	μ_2
Texture P: $h_{g,P}$	0	1
Texture Q: $h_{g,Q}$	0	2

Hence, the textures can be differentiated by using the μ_2 values.