

EE3206/EE3206E ICVIP**Tutorial Set B – Solutions**

Question 1**Part (a)**

The FT of $\delta(x, y)$ is

$$\begin{aligned}\mathcal{F}\{\delta(x, y)\} &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \delta(x, y) \exp[-j2\pi(ux + vy)] dx dy \quad (\text{by definition}) \\ &= \exp[-j2\pi(ux + vy)]|_{x,y=0} \\ &= 1\end{aligned}$$

Part (b)

The inverse FT of $\delta(u, v)$ is


$$\begin{aligned}\mathcal{F}^{-1}\{\delta(u, v)\} &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \delta(u, v) \exp[j2\pi(ux + vy)] du dv \quad (\text{by definition}) \\ &= \exp[j2\pi(ux + vy)]|_{u,v=0} \\ &= 1\end{aligned}$$

Taking the FT of both sides,

$$\delta(u, v) = \mathcal{F}\{1\}$$

i.e., the FT of $f(x, y) = 1$ is $\delta(u, v)$.

Question 2



The image function $f(x, y)$ can be regarded as the sum of two component functions $f_1(x, y)$ and $f_2(x, y)$:

$$f(x, y) = f_1(x, y) + f_2(x, y)$$

where

$$\begin{aligned} f_1(x, y) &= 10 \quad \text{for all } x, y \\ f_2(x, y) &= \begin{cases} 40 & -1 \leq x \leq +1, -2.5 \leq y \leq +0.5 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

Method I - (not recommended)

$$\begin{aligned} F(u, v) &= \mathcal{F}\{f(x, y)\} \\ &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} 10 \exp[-j2\pi(ux + vy)] dx dy \\ &\quad + \int_{-2.5}^{0.5} \int_{-1}^{1} 40 \exp[-j2\pi(ux + vy)] dx dy \\ &= \dots \end{aligned}$$

$$\text{Fourier spectrum} = |F(u, v)|$$

Method II

Use $|F(u, v)| = |\mathcal{F}\{f(x, y)\}| = |\mathcal{F}\{f(x, y - 1)\}|$

We have


$$f(x, y) = f_1(x, y) + f_2(x, y)$$

Hence,

$$\begin{aligned} f(x, y - 1) &= f_1(x, y) + f_2(x, y - 1) \\ \mathcal{F}\{f(x, y - 1)\} &= \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} 10 \exp[-j2\pi(ux + vy)] dx dy \\ &\quad + 40 \int_{-1.5}^{+1.5} \int_{-1}^{+1} \exp[-j2\pi(ux + vy)] dx dy \\ &= 10\delta(u, v) + 40 \left[\frac{\exp(-j2\pi ux)}{-j2\pi u} \right]_{-1}^{+1} \left[\frac{\exp(-j2\pi vy)}{-j2\pi v} \right]_{-1.5}^{+1.5} \\ &= 10\delta(u, v) + 40[2\text{sinc}(2u)] [3\text{sinc}(3v)] \\ &= 10\delta(u, v) + 240\text{sinc}(2u)\text{sinc}(3v) \end{aligned}$$

The Fourier spectrum of $f(x, y)$ is

$$\begin{aligned} |F(u, v)| &= |\mathcal{F}\{f(x, y)\}| \\ &= |\mathcal{F}\{f(x, y - 1)\}| \\ &= 10\delta(u, v) + 240|\text{sinc}(2u)||\text{sinc}(3v)| \end{aligned}$$



Question 3

$$\begin{aligned}
 \mathcal{F}^{-1}\{\delta(u-a, v-b)\} &= \int \int \delta(u-a, v-b) \exp[j2\pi(ux+vy)] du dv \\
 &= \exp[j2\pi(ux+vy)]|_{u=a, v=b} \\
 &= \exp[j2\pi(ax+by)]
 \end{aligned}$$

Similarly,

$$\mathcal{F}^{-1}\{\delta(u+a, v+b)\} = \exp[-j2\pi(ax+by)]$$

Hence,

$$\begin{aligned}
 \mathcal{F}^{-1}\{\delta(u-a, v-b) + \delta(u+a, v+b)\} &= \exp[j2\pi(ax+by)] + \exp[-j2\pi(ax+by)] \\
 &= 2 \cos 2\pi(ax+by)
 \end{aligned}$$

Thus,

$$\cos 2\pi(ax+by) \leftrightarrow \frac{1}{2}\delta(u-a, v-b) + \frac{1}{2}\delta(u+a, v+b) \quad (1)$$


Part (a)

From Eq. (1), we let $a = 10$ and $b = 0$:

$$f_1(x, y) = \cos(20\pi x) \leftrightarrow \frac{1}{2}\delta(u-10, v) + \frac{1}{2}\delta(u+10, v)$$

Hence, the Fourier spectrum of $f_1(x, y)$ is

$$|F_1(u, v)| = \frac{1}{2}\delta(u-10, v) + \frac{1}{2}\delta(u+10, v)$$



Part (b)

We first obtain the Fourier transform of $\sin 2\pi(ax + by)$. Similar to Eq. (1),

$$\begin{aligned}\mathcal{F}^{-1}\{\delta(u - a, v - b) - \delta(u + a, v + b)\} &= \exp[j2\pi(ax + by)] - \exp[-j2\pi(ax + by)] \\ &= 2j \sin 2\pi(ax + by)\end{aligned}$$

Hence,

$$\sin 2\pi(ax + by) \leftrightarrow \frac{1}{2j}\delta(u - a, v - b) - \frac{1}{2j}\delta(u + a, v + b) \quad (2)$$


$$= -\frac{j}{2}\delta(u - a, v - b) + \frac{j}{2}\delta(u + a, v + b) \quad (3)$$

From Eq. (3), we let $a = 0$ and $b = 20$:

$$f_2(x, y) = \sin(40\pi y) \leftrightarrow \frac{j}{2}\delta(u, v + 20) - \frac{j}{2}\delta(u, v - 20)$$

Hence, the Fourier spectrum of $f_2(x, y)$ is

$$|F_2(u, v)| = \frac{1}{2}\delta(u, v + 20) + \frac{1}{2}\delta(u, v - 20)$$



Part (c)

$$\begin{aligned}
 f_3(x, y) &= \sin(30x + 40y) \\
 &= \sin 2\pi \left(\frac{15}{\pi}x + \frac{20}{\pi}y \right) \\
 F_3(u, v) &= \frac{j}{2} \delta \left(u + \frac{15}{\pi}, v + \frac{20}{\pi} \right) - \frac{j}{2} \delta \left(u - \frac{15}{\pi}, v - \frac{20}{\pi} \right)
 \end{aligned}$$


Hence, the Fourier spectrum of $f_3(x, y)$ is

$$|F_3(u, v)| = \frac{1}{2} \delta \left(u + \frac{15}{\pi}, v + \frac{20}{\pi} \right) + \frac{1}{2} \delta \left(u - \frac{15}{\pi}, v - \frac{20}{\pi} \right) \quad (4)$$

Part (d)

$f_4(x, y) = \sin(30x + 40y + 30)$ is a translated version of $f_3(x, y)$.
Hence

$$|F_4(u, v)| = |F_3(u, v)|$$



Question 4

Consider the transform $F(u)$ shifted by u_0 , i.e.,

$$F_t(u) = F(u - u_0)$$

The inverse DFT of $F_t(u)$ is

$$\begin{aligned} f_t(x) &= \mathcal{F}^{-1}\{F(u - u_0)\} \\ &= f(x) \exp(j2\pi u_0 x/N) \end{aligned}$$

from the translation property of the DFT (Ch. 3B).

We wish to shift the transform $F(u)$ to the right by $N/2$:

$$F(u) \rightarrow F(u - N/2)$$

i.e., $u_0 = N/2$. Shifting $F(u)$ by u_0 is obtained by multiplying $f(x)$ by $\exp(j2\pi u_0 x/N)$. In this case, the exponential term is

$$\begin{aligned} \exp(j2\pi u_0 x/N) &= \exp\left(j2\pi \frac{N}{2} \frac{x}{N}\right) \\ &= \exp(j\pi x) \\ &= (e^{j\pi})^x \\ &= (-1)^x \end{aligned}$$

Thus,

$$f_t(x) = f(x)(-1)^x$$

and

$$\mathcal{F}\{f(x)(-1)^x\} = F(u - N/2)$$

$$\begin{aligned} F(u) &= \frac{1}{N} \sum_{x=0}^{N-1} f(x) e^{-j2\pi u x/N} \\ F(u - u_0) &= \frac{1}{N} \sum_{x=0}^{N-1} f(x) e^{-j2\pi (u - u_0) x/N} \\ &= \frac{1}{N} \sum_{x=0}^{N-1} \{f(x) e^{j2\pi u_0 x/N}\} e^{-j2\pi u x/N} \\ &= \mathcal{F}\{f(x) e^{j2\pi u_0 x/N}\} \end{aligned}$$

Question 5

Given

$$f(x) = 1, 1, 1, 1, 0, 0, 0, 0$$

Part (a)

$$F(u) = \frac{1}{N} \sum_{x=0}^{N-1} f(x) \exp(-j2\pi ux/N) = \frac{1}{8} \sum_{x=0}^3 \exp(-j2\pi ux/8)$$

$$F(0) = \frac{1}{8} \sum_{x=0}^3 \exp(0) = 0.5$$

$$\begin{aligned} F(1) &= \frac{1}{8} \sum_{x=0}^3 \exp(-j2\pi(1)x/8) \\ &= \frac{1}{8} \exp(0) + \frac{1}{8} \exp(-j2\pi/8) + \frac{1}{8} \exp(-j2\pi \times 2/8) \\ &\quad + \frac{1}{8} \exp(-j2\pi \times 3/8) \\ &= 0.327\angle -1.18 \end{aligned}$$

$$\begin{aligned} F(2) &= \frac{1}{8} \sum_{x=0}^3 \exp(-j2\pi(2)x/8) \\ &= \frac{1}{8} \exp(0) + \frac{1}{8} \exp(-j2\pi \times 2/8) + \frac{1}{8} \exp(-j2\pi \times 4/8) \\ &\quad + \frac{1}{8} \exp(-j2\pi \times 6/8) \\ &= 0 \end{aligned}$$

$$\begin{aligned} F(3) &= \frac{1}{8} \sum_{x=0}^3 \exp(-j2\pi(3)x/8) \\ &= \frac{1}{8} \exp(0) + \frac{1}{8} \exp(-j2\pi \times 3/8) + \frac{1}{8} \exp(-j2\pi \times 6/8) \\ &\quad + \frac{1}{8} \exp(-j2\pi \times 9/8) \\ &= 0.135\angle -0.393 \end{aligned}$$

$$\begin{aligned}
F(4) &= \frac{1}{8} \sum_{x=0}^3 \exp(-j2\pi(4)x/8) \\
&= \frac{1}{8} \exp(0) + \frac{1}{8} \exp(-j2\pi \times 4/8) + \frac{1}{8} \exp(-j2\pi \times 8/8) \\
&\quad + \frac{1}{8} \exp(-j2\pi \times 12/8) \\
&= 0
\end{aligned}$$

$$\begin{aligned}
F(5) &= F^*(-5) \quad (\text{conjugate symmetry}) \\
&= F^*(8-5) \quad (\text{periodicity}) \\
&= F^*(3) \\
&= 0.135\angle +0.393
\end{aligned}$$

$$\begin{aligned}
F(6) &= F^*(2) \\
&= 0
\end{aligned}$$


$$\begin{aligned}
F(7) &= F^*(1) \\
&= 0.327\angle 1.18
\end{aligned}$$

Part (b)

$$f(x) = 1, \quad 1, \quad 1, \quad 1, \quad 0, \quad 0, \quad 0, \quad 0$$


$$F(u) = 0.5 \quad 0.327\angle -1.18, \quad 0, \quad 0.135\angle -0.393$$

$$0, \quad 0.135\angle +0.393, \quad 0, \quad 0.327\angle 1.18$$



Part (c)

$$\begin{aligned}
 g(x) &= (-1)^x f(x) = +1, \quad -1, \quad +1, \quad -1 \quad 0, \quad 0, \quad ,0, \quad 0 \\
 G(u) &= 0, \quad 0.135 \angle 0.393, \quad 0, \quad 0.327 \angle 1.18, \\
 &\quad 0.5, \quad 0.327 \angle -1.18, \quad 0, \quad 0.135 \angle -0.393
 \end{aligned}$$



Question 6

Part (a)

$$f_1(x) = 1, 1, 1, 1$$

$$F_1(u) = \frac{1}{N} \sum_{x=0}^{N-1} f_1(x) \exp(-j2\pi ux/N) = \frac{1}{4} \sum_{x=0}^3 \exp(-j2\pi ux/4)$$

$$F_1(0) = \frac{1}{4} \sum_{x=0}^3 \exp(0) = \frac{1}{4} \times 4 = 1$$

$$F_1(1) = \frac{1}{4} \sum_{x=0}^3 \exp(-j2\pi(1)x/4) = \frac{1}{4} \times 0 = 0$$

$$F_1(2) = \frac{1}{4} \sum_{x=0}^3 \exp(-j2\pi(2)x/4) = \frac{1}{4} \times 0 = 0$$

$$F_1(3) = \frac{1}{4} \sum_{x=0}^3 \exp(-j2\pi(3)x/4) = \frac{1}{4} \times 0 = 0$$

Hence,

$$F_1(u) = 1, 0, 0, 0$$

Part(b)

$$f_2(x) = 1, 0, 0, 0$$

$$\begin{aligned} F_2(u) &= \frac{1}{N} \sum_{x=0}^{N-1} f_2(x) \exp(-j2\pi ux/N) \\ &= \frac{1}{4} \sum_{x=0}^0 1 \\ &= \frac{1}{4} \end{aligned}$$

$$F_2(0) = \frac{1}{4}, \quad F_2(1) = \frac{1}{4}, \quad F_2(2) = \frac{1}{4}, \quad F_2(3) = \frac{1}{4}$$

Hence,

$$F(u) = \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$$

Part (c)

From the periodicity property, we note that

$$f_3(x) = f_2(x - 1)$$

Hence,

$$\begin{aligned} F_3(u) &= F_2(u) \exp(-j2\pi ua/4) \quad (\text{translation property}) \\ &= F_2(u) \exp(-j\pi u/2) \end{aligned}$$

$$\begin{aligned} F_3(0) &= F_2(0) \exp(0) = F_2(0) \\ F_3(1) &= F_2(1) \exp(-j\pi/2) = -jF_2(1) \\ F_3(2) &= F_2(2) \exp(-j\pi) = -F_2(2) \\ F_3(3) &= F_2(3) \exp(-j3\pi/2) = jF_2(3) \end{aligned}$$

$$F_3(u) = \frac{1}{4}, -\frac{j}{4}, -\frac{1}{4}, \frac{j}{4}$$

Part (d)

$$f_4(x, y) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$F_4(x, v) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$F_4(u, v) = \frac{1}{4} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

Part (e)

$$f_5(x, y) = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

$$F_5(x, v) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

$$F_5(u, v) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Note that $F_5(u, v) = \delta(u, v)$.