

NATIONAL UNIVERSITY OF SINGAPORE  
Department of Mathematics

MA1506 Laboratory 3 (MATLAB)  
Comments and Suggested Solutions  
Semester II 2010/2011

*Exercise 3*

1. 716.25

2. 70. From the information available,  $M = \begin{bmatrix} 0.7 & 0.2 & 0.3 \\ 0.2 & 0.5 & 0.3 \\ 0.1 & 0.3 & 0.4 \end{bmatrix}$ . The ‘brute force’ way

to compute the long run behaviour is to just calculate very large powers of  $M$  and compare the values of the entries.

```
>> M^(200) - M^(100)
>> M^(100)*[100; 100; 100]
```

In the first command, we can see that the difference between the entries in  $M^{200}$  and  $M^{100}$  are less than  $10^{-14}$ . So we are sufficiently confident that  $M^{100}$  is close enough to the long run value. Alternatively, we can do things properly by diagonalizing  $M$  and taking limits.

```
>> [P D] = eig(M)
>> D1= diag( [ 1 0 0 ])
>> P*D1*inv(P)*[100; 100; 100]
```

The **diag** command creates a diagonal matrix with the entries specified. It is clear that D1 is the limit of  $D^n$  as  $n$  gets large.

3. 104 million

4. Eigenvalues are -4, 3 and 3.

5. In this case we have complex eigenvalues  $\pm i$  and note that  $P$  is also a matrix with complex entries.

```
6. >> C=[ 5 0 0; 1 5 0; 0 1 5]
>> [P D] = eig(C)
>> det(P)
```

In this case, we have the eigenvalue 5 repeated 3 times, and the determinant of  $P$  is approximately zero. Actually, matrix  $A$  is not diagonalizable and the determinant of  $P$  should be exactly zero. This slight error is due to the algorithm used by the **eig** function.

```
7. >> A=[5 6 2; 0 -1 -8; 1 0 -2]
>> poly(A)

ans =

    1.0000   -2.0000  -15.0000   36.0000

>> 1*A^3 -2*A^2 -15*A+36*eye(3)
>> B= [ -2 -1 ; 5 2]
>> poly(B)

ans =

    1.0000    0.0000    1.0000

>> 1*B^2 + eye(2)
>> C=[ 5 0 0; 1 5 0; 0 1 5]
>> poly(C)

ans =

    1   -15    75  -125

>> 1*C^3 -15*C^2 + 75*C -125*eye(3)
```

—The End—