Inter-ecosystem specifications of energy transfer in trophic interactions

Benjamin Mercier 1,2,‡
 Université de Sherbrooke; ² Québec Centre for Biodiversity Sciences † These authors contributed equally to the work
Correspondance to: Benjamin Mercier — benjamin.b.mercier@usherbrooke.ca
This is a very short abstract
1
Introduction
Blabla this is an introduction. This is a citation test Brose <i>et al.</i> (2019), and this is also another citation test (Brose <i>et al.</i> 2019).
2
The data
The Ecopath data were obtain from Jacquet <i>et al.</i> (2016) on request to the corresponding author. Before manipulation, the data initially represented 116 Ecopath trophic networks. Ecopath is a modeling software which aims to quantify species interactions statically and is mass-balanced (Christensen n.d.). One weakness, if I may, is that a lot of these Ecopath networks are not taxonomically resolved to the species but encompass trophic groups or guilds. The first step here was then, for each of these networks, match the data to the original article from which they originated. This matching was done to
3
Figures
4
Analyses
Conclusion

the poisot lab

©

Keywords: trophic interactions energy fluxes ecosystem

Brose, U., Archambault, P., Barnes, A.D., Bersier, L.-F., Boy, T., Canning-Clode, J., *et al.* (2019). Predator traits determine food-web architecture across ecosystems. *Nature Ecology & Evolution*, 3, 919–927.




Figure 1 This is a prelim figure about the fluxes.




Figure 2 This is the map of network

Christensen, V. (n.d.). Ecopath with Ecosim: A User's Guide, 155.

Jacquet, C., Moritz, C., Morissette, L., Legagneux, P., Massol, F., Archambault, P., *et al.* (2016). No complexity stability relationship in empirical ecosystems. *Nature Communications*, 7, 12573.