# Review Session 1: Homework 3 & OLS Regression

Jiahui Shui jishui@ucsd.edu

Rady School of Management, UCSD

November 8, 2024

#### Hints for HW3

- ▶ Period: January 1926 until December 2018
- CRSP\_vw and CRSP\_vwx: with / without dividends
- Don't spend to much time on Question 8 and 9

### Econometric issues in return predictability

Consider the following system

$$r_{t+1} = \alpha + \beta dp_t + \varepsilon_{t+1}$$

$$dp_{t+1} = \mu + \phi dp_t + u_{t+1}$$
(1)

- ▶ In this system,  $dp_t$  is highly persistent  $(\phi \approx 1)$ ,  $\beta > 0$ .
- Now we suppose that  $\mathbb{E}[dp_{t-1}\varepsilon_t] = 0$  by construction.
- $ightharpoonup \hat{\phi}$  tends to be downward biased. This is standard issue in OLS.

$$\mathbb{E}[\hat{\phi}] = \phi - \frac{1+3\phi}{T} + O(1/T^2) \tag{2}$$

And  $\hat{\beta}$  is upward biased, which means that we reject the null of no predictability too often.

$$\mathbb{E}[\hat{\beta} - \beta] = \frac{\sigma_{\varepsilon u}}{\sigma_{u}^{2}} \mathbb{E}[\hat{\phi} - \phi] = -\frac{\sigma_{\varepsilon u}}{\sigma_{u}^{2}} \frac{1 + 3\phi}{T}$$
(3)

▶ Q: Why  $\sigma_{\varepsilon u}$  < 0? A positive dp shock usually has no news about dividends, so it means a negative p shock and a negative r shock.

1. Consider the following regression equation

$$y_t = \alpha + \beta x_t + \varepsilon_t \tag{4}$$

Assume that  $\mathbb{E}[\varepsilon_t|x_t] = 0$  for all t.

(a) Prove that  $\mathbb{E}[arepsilon_t|x_t]=0$  implies  $\mathbb{E}[arepsilon_tx_t]=0$ 

Solution: By Law of Iterated Expection, we have

$$\mathbb{E}[\varepsilon_t x_t] = \mathbb{E}[\mathbb{E}[\varepsilon_t x_t | x_t]] = \mathbb{E}[x_t \mathbb{E}[\varepsilon_t | x_t]] = 0$$
 (5)

(b) Find the OLS estimator  $\hat{\alpha}$  and  $\hat{\beta}$ 

Solution: Define

$$Q(\alpha, \beta) = \sum_{t=1}^{T} (y_t - \alpha - \beta x_t)^2$$
 (6)

The FOCs are

$$\frac{\partial Q}{\partial \alpha} = -2 \sum_{t=1}^{T} (y_t - \alpha - \beta x_t) = 0$$

$$\frac{\partial Q}{\partial \beta} = -2 \sum_{t=1}^{T} (y_t - \alpha - \beta x_t) x_t = 0$$
(7)

The first equation can be simplified as  $\bar{y} = \hat{\alpha} + \hat{\beta}\bar{x}$ . Substitute this into the second equation, we get

$$\hat{\beta} = \frac{\sum_{t=1}^{I} x_t y_t - T \bar{x} \bar{y}}{\sum_{t=1}^{T} x_t^2 - T \bar{x}^2}$$
 (8)

(c) Let  $\hat{y}_t := \hat{\alpha} + \hat{\beta} x_t$ . Define  $e_t = y_t - \hat{y}_t$ . Show that

$$\sum_{t=1}^T e_t = 0$$

Solution: We have

$$\sum_{t=1}^{T} e_t = \sum_{t=1}^{T} (y_t - \hat{\alpha} - \hat{\beta} x_t)$$
 (9)

Note that this is just the first FOC.

(d) Show that 
$$\hat{\beta}$$
 is unbiased, i.e.  $\mathbb{E}[\hat{\beta}|X] = \beta$ , where  $X = (x_1, \dots, x_T)'$ 

Solution: Note that

$$\mathbb{E}[\hat{\beta}|X] = \mathbb{E}\left[\frac{\sum_{t=1}^{T} x_t y_t - T\bar{x}\bar{y}}{\sum_{t=1}^{T} x_t^2 - T\bar{x}^2} \middle| X\right] = \frac{\sum_{t=1}^{T} x_t \mathbb{E}[y_t|X] - T\bar{x}\mathbb{E}[\bar{y}|X]}{\sum_{t=1}^{T} - T\bar{x}^2}$$
(10)

and

$$\mathbb{E}[y_t|X] = \mathbb{E}[\alpha + \beta x_t + \varepsilon_t|X] = \alpha + \beta x_t \tag{11}$$

Then

$$\mathbb{E}[\bar{y}|X] = \frac{1}{T}\mathbb{E}[\sum_{t=1}^{T} y_t|X] = \alpha + \beta\bar{x}$$
 (12)

Substituting those two equations into equation (10) yields

$$\mathbb{E}[\hat{\beta}|X] = \beta \tag{13}$$

Notes: Unbiased property requires  $\mathbb{E}[\varepsilon_t|X] = 0$ .

(e) Now, consider the following regression equation

$$y_t = \beta x_t + \varepsilon_t \tag{14}$$

Find the OLS estimator  $\hat{\beta}$ . Calculate  $\sum_{t=1}^{T} e_t$  again.

**Solution**: Now

$$Q(\beta) = \sum_{t=1}^{T} (y_t - \beta x_t)^2$$
 (15)

The FOC gives us

$$\frac{\partial Q(\beta)}{\beta} = -2\sum_{t=1}^{T} (y_t - \beta x_t) x_t = 0 \Rightarrow \hat{\beta} = \frac{\sum_{t=1}^{T} x_t y_t}{\sum_{t=1}^{T} x_t^2}$$
(16)

Now  $\sum_{t=1}^{T} e_t$  is not necessarily to be 0.

$$\sum_{t=1}^{T} e_t = \sum_{t=1}^{T} y_t - \frac{\sum_{t=1}^{T} x_t y_t}{\sum_{t=1}^{T} x_t^2} \sum_{t=1}^{T} x_t$$
 (17)

2. Consider the following AR(1) process

$$x_t = \rho x_{t-1} + \varepsilon_t \tag{18}$$

where  $|\rho| < 1$ ,  $\{\varepsilon_t\}_{t=0}^{\infty}$  is white noise with variance  $\sigma^2$ . Suppose that  $\mathbb{E}[x_{t-1}\varepsilon_t] = 0$ .

(a) Is the OLS estimator  $\hat{\rho}$  unbiased? Is  $\hat{\rho}$  consistent?

**Solution**: No,  $\hat{\rho}$  is biased. Since unbiased estimation requires that  $\mathbb{E}[\varepsilon_t|X]=0$ , meaning  $\varepsilon_t$  is uncorrelated with all  $x_t$ , both past and future. Obviously here  $\varepsilon_t$  is correlated to future  $x_t$ . But  $\hat{\rho}$  is consistent. Consistency only requires predetermined explanatory variables, i.e.  $\mathbb{E}[x_{t-1}\varepsilon_t]=0$ .

(b) Find  $\mathbb{E}[x_t]$  and  $\operatorname{Var}(x_t)$ 

**Solution**: There are some tricks here. If we impose  $x_t$  is stationary here, then  $\mathbb{E}[x_t] = \mathbb{E}[x_{t-1}], \mathrm{Var}(x_t) = \mathrm{Var}(x_{t-1})$ . Hence

$$\mathbb{E}[x_t] = \rho \mathbb{E}[x_{t-1}] \Rightarrow \mathbb{E}[x_t] = 0 \tag{19}$$

$$\operatorname{Var}(x_t) = \rho^2 \operatorname{Var}(x_{t-1}) + \sigma^2 \Rightarrow \operatorname{Var}(x_t) = \frac{\sigma^2}{1 - \rho^2}$$
 (20)