

Proof: A 2x2 Matrix is a Vector Space

Bill Cruise (UNCC Student Id: 800932170)

Wednesday, March 23, 2016

$\mathbb{M}_{2 \times 2}(\mathbb{R}) = \{\mathbb{M}_{2 \times 2} : a_{ij} \in \mathbb{R}\}$ with

$$A + B = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} + \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix}, \text{ and}$$

$$cA = c \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} ca_{11} & ca_{12} \\ ca_{21} & ca_{22} \end{bmatrix}$$

Closure:

1. Let $A, B \in \mathbb{M}_{2 \times 2}(\mathbb{R})$.

$$A + B = C = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix}.$$

$a_{11}, b_{11}, a_{12}, b_{12}, a_{21}, b_{21}, a_{22}, b_{22} \in \mathbb{R} \implies a_{11} + b_{11}, a_{12} + b_{12}, a_{21} + b_{21}, a_{22} + b_{22} \in \mathbb{R}$,

and $C \in \mathbb{M}_{2 \times 2}$, so $A + B \in \mathbb{M}_{2 \times 2}(\mathbb{R})$

That is, $\mathbb{M}_{2 \times 2}(\mathbb{R})$ is closed under vector addition.

2. Let $A \in \mathbb{M}_{2 \times 2}(\mathbb{R}), c \in \mathbb{R}$.

$$cA = \begin{bmatrix} ca_{11} & ca_{12} \\ ca_{21} & ca_{22} \end{bmatrix}.$$

$c, a_{11}, a_{12}, a_{21}, a_{22} \in \mathbb{R} \implies ca_{11}, ca_{12}, ca_{21}, ca_{22} \in \mathbb{R}$, so $cA \in \mathbb{M}_{2 \times 2}(\mathbb{R})$.

That is, $\mathbb{M}_{2 \times 2}(\mathbb{R})$ is closed under scalar multiplication.

Addition:

3. Commutativity - $A + B = B + A$ for every $A, B \in \mathbb{M}_{2 \times 2}(\mathbb{R})$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix} = \begin{bmatrix} b_{11} + a_{11} & b_{12} + a_{12} \\ b_{21} + a_{21} & b_{22} + a_{22} \end{bmatrix} \text{ by commutativity of addition in } \mathbb{R}.$$

$$\begin{bmatrix} b_{11} + a_{11} & b_{12} + a_{12} \\ b_{21} + a_{21} & b_{22} + a_{22} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} + \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = B + A.$$

4. Associativity - $(A + B) + C = A + (B + C)$ for every $A, B, C \in \mathbb{M}_{2 \times 2}(\mathbb{R})$.

$$(A + B) + C = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix} + \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} = \begin{bmatrix} (a_{11} + b_{11}) + c_{11} & (a_{12} + b_{12}) + c_{12} \\ (a_{21} + b_{21}) + c_{21} & (a_{22} + b_{22}) + c_{22} \end{bmatrix} =$$
$$\begin{bmatrix} a_{11} + (b_{11} + c_{11}) & a_{12} + (b_{12} + c_{12}) \\ a_{21} + (b_{21} + c_{21}) & a_{22} + (b_{22} + c_{22}) \end{bmatrix} \text{ by associativity of addition in } \mathbb{R}.$$

$$\begin{bmatrix} a_{11} + (b_{11} + c_{11}) & a_{12} + (b_{12} + c_{12}) \\ a_{21} + (b_{21} + c_{21}) & a_{22} + (b_{22} + c_{22}) \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} + \begin{bmatrix} b_{11} + c_{11} & b_{12} + c_{12} \\ b_{21} + c_{21} & b_{22} + c_{22} \end{bmatrix} = A + (B + C).$$

5. Existence of an additive identity ($\vec{0}$), i.e., there must exist a $\vec{0}$ such that $A + \vec{0} = \vec{0} + A = A$ for every $A \in \mathbb{M}_{2 \times 2}(\mathbb{R})$.

$$A + \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} = \begin{bmatrix} a_{11} + c_{11} & a_{12} + c_{12} \\ a_{21} + c_{21} & a_{22} + c_{22} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \implies \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \text{ i.e., } \vec{0} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \text{ is the additive identity of } \mathbb{M}_{2 \times 2}(\mathbb{R}).$$

6. Existence of additive inverses, i.e., for every $A \in \mathbb{M}_{2 \times 2}(\mathbb{R})$ there must exist some B such that $A + B = B + A = \vec{0}$.

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \implies b_{11} = -a_{11}, b_{12} = -a_{12}, b_{21} = -a_{21}, \text{ and } b_{22} = -a_{22}, \text{ so } B = \begin{bmatrix} -a_{11} & -a_{12} \\ -a_{21} & -a_{22} \end{bmatrix}.$$

Multiplication:

7. Distribution of a scalar onto the sum of two vectors, i.e., for every $c \in \mathbb{R}$ and every $A, B \in \mathbb{M}_{2 \times 2}(\mathbb{R})$, $c(A + B) = cA + cB$.

$$c(A + B) = c \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{bmatrix} = \begin{bmatrix} c(a_{11} + b_{11}) & c(a_{12} + b_{12}) \\ c(a_{21} + b_{21}) & c(a_{22} + b_{22}) \end{bmatrix} = \begin{bmatrix} ca_{11} + cb_{11} & ca_{12} + cb_{12} \\ ca_{21} + cb_{21} & ca_{22} + cb_{22} \end{bmatrix} = \begin{bmatrix} ca_{11} & ca_{12} \\ ca_{21} & ca_{22} \end{bmatrix} + \begin{bmatrix} cb_{11} & cb_{12} \\ cb_{21} & cb_{22} \end{bmatrix} = c \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} + c \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = cA + cB.$$

8. Distribution of a vector onto the sum of two scalars, i.e., for every $c, d \in \mathbb{R}$ and every $A \in \mathbb{M}_{2 \times 2}(\mathbb{R})$, $(c + d)A = cA + dA$.

$$(c + d)A = \begin{bmatrix} (c + d)a_{11} & (c + d)a_{12} \\ (c + d)a_{21} & (c + d)a_{22} \end{bmatrix} = \begin{bmatrix} ca_{11} + da_{11} & ca_{12} + da_{12} \\ ca_{21} + da_{21} & ca_{22} + da_{22} \end{bmatrix} = \begin{bmatrix} ca_{11} & ca_{12} \\ ca_{21} & ca_{22} \end{bmatrix} + \begin{bmatrix} da_{11} & da_{12} \\ da_{21} & da_{22} \end{bmatrix} = cA + dA.$$

9. Scalar multiplication of a scalar multiplication, i.e., for every $c, d \in \mathbb{R}$ and $A \in \mathbb{M}_{2 \times 2}(\mathbb{R})$, $c(dA) = (cd)A$.

$$c(dA) = c \begin{bmatrix} da_{11} & da_{12} \\ da_{21} & da_{22} \end{bmatrix} = \begin{bmatrix} cda_{11} & cda_{12} \\ cda_{21} & cda_{22} \end{bmatrix} = \begin{bmatrix} (cd)a_{11} & (cd)a_{12} \\ (cd)a_{21} & (cd)a_{22} \end{bmatrix} = (cd) \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = (cd)A.$$

10. Scalar multiplication with the multiplicative identity of the field (\mathbb{R}), i.e., for every $A \in \mathbb{M}_{2 \times 2}(\mathbb{R})$, $1A = A$.

$$1A = 1 \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 1a_{11} & 1a_{12} \\ 1a_{21} & 1a_{22} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = A.$$