

# Machine Intelligence

## 4. Reasoning under Uncertainty, Part I: Basics

(Our Machinery for) Thinking About What is Likely to be True

Álvaro Torralba



AALBORG UNIVERSITET


Fall 2022

Thanks to Thomas D. Nielsen and Jörg Hoffmann for slide sources


# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

# The Wumpus World




# The Wumpus World



- **Actions:** *GoForward*, *TurnRight* (by  $90^\circ$ ), *TurnLeft* (by  $90^\circ$ ), *Grab* object in current cell, *Shoot* arrow in direction you're facing (you got exactly one arrow), *Leave cave* if you're in cell [1,1].

→ Fall down *Pit*, meet live *Wumpus*: *Game Over*.

# The Wumpus World




- **Actions:** *GoForward*, *TurnRight* (by  $90^\circ$ ), *TurnLeft* (by  $90^\circ$ ), *Grab* object in current cell, *Shoot* arrow in direction you're facing (you got exactly one arrow), *Leave* cave if you're in cell [1,1].

→ Fall down *Pit*, meet live *Wumpus*: **Game Over.**

- **Initial knowledge:** You're in cell [1,1] facing east. There's a Wumpus, and there's gold.
- **Goal:** Have the gold and be outside the cave.

# The Wumpus World




- **Actions:** *GoForward*, *TurnRight* (by  $90^\circ$ ), *TurnLeft* (by  $90^\circ$ ), *Grab* object in current cell, *Shoot* arrow in direction you're facing (you got exactly one arrow), *Leave* cave if you're in cell [1,1].

→ Fall down *Pit*, meet live *Wumpus*: *Game Over*.

- **Initial knowledge:** You're in cell [1,1] facing east. There's a Wumpus, and there's gold.
- **Goal:** Have the gold and be outside the cave.

**Percepts:** [*Stench*, *Breeze*, *Glitter*, *Bump*, *Scream*]

# The Wumpus World



- **Actions:** *GoForward*, *TurnRight* (by  $90^\circ$ ), *TurnLeft* (by  $90^\circ$ ), *Grab* object in current cell, *Shoot* arrow in direction you're facing (you got exactly one arrow), *Leave* cave if you're in cell [1,1].

→ Fall down *Pit*, meet live *Wumpus*: *Game Over*.

- **Initial knowledge:** You're in cell [1,1] facing east. There's a Wumpus, and there's gold.
- **Goal:** Have the gold and be outside the cave.

**Percepts:** [*Stench*, *Breeze*, *Glitter*, *Bump*, *Scream*]

- Cell adjacent (i.e. north, south, west, east) to Wumpus: *Stench* (else: *None*).
- Cell adjacent to Pit: *Breeze* (else: *None*).
- Cell that contains gold: *Glitter* (else: *None*).
- You walk into a wall: *Bump* (else: *None*).
- Wumpus shot by arrow: *Scream* (else: *None*).

# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2 OK	2,2	3,2	4,2
1,1 <div>A</div> OK	2,1 OK	3,1	4,1

(1) Initial state



# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1 A OK	2,1 OK	3,1	4,1

(1) Initial state

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2 P?	3,2	4,2
OK			
1,1 V OK	2,1 A B OK	3,1 P?	4,1

(2) One step to right

# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1 A OK	2,1 OK	3,1	4,1

(1) Initial state

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2 P?	3,2	4,2
OK			
1,1 V OK	2,1 A B OK	3,1 P?	4,1

(2) One step to right

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

(3) Back, and up to [1,2]

→ The Wumpus is in [1,3]! How do we know?

# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1 A OK	2,1 OK	3,1	4,1

(1) Initial state

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2 P?	3,2	4,2
OK			
1,1 V OK	2,1 A B OK	3,1 P?	4,1

(2) One step to right

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

(3) Back, and up to [1,2]

→ The Wumpus is in [1,3]! How do we know? Because in [2,1] we perceived no Stench, the Stench in [1,2] can only come from [1,3].

# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
1,1 A OK	2,1 OK	3,1	4,1

(1) Initial state

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2 P?	3,2	4,2
1,1 V OK	2,1 A B OK	3,1 P?	4,1

(2) One step to right

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

(3) Back, and up to [1,2]

→ The Wumpus is in [1,3]! How do we know? Because in [2,1] we perceived no Stench, the Stench in [1,2] can only come from [1,3].

→ There's a Pit in [3,1]! How do we know?

# Reasoning in the Wumpus World

**A:** Agent, **V:** Visited, **OK:** Safe, **P:** Pit, **W:** Wumpus, **B:** Breeze, **S:** Stench, **G:** Gold

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1 A OK	2,1 OK	3,1	4,1

(1) Initial state

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2 P?	3,2	4,2
OK			
1,1 V OK	2,1 A B OK	3,1 P?	4,1

(2) One step to right

1,4	2,4	3,4	4,4
1,3 W!	2,3	3,3	4,3
1,2 A S OK	2,2 OK	3,2	4,2
1,1 V OK	2,1 B V OK	3,1 P!	4,1

(3) Back, and up to [1,2]

→ The Wumpus is in [1,3]! How do we know? Because in [2,1] we perceived no Stench, the Stench in [1,2] can only come from [1,3].

→ There's a Pit in [3,1]! How do we know? Because in [1,2] we perceived no Breeze, the Breeze in [2,1] can only come from [3,1].

# Agents that Think Rationally

## Think Before You Act!

```

function KB-AGENT(percept) returns an action
  persistent: KB, a knowledge base
               t, a counter, initially 0, indicating time

  TELL(KB, MAKE-PERCEPT-SENTENCE(percept, t))
  action ← ASK(KB, MAKE-ACTION-QUERY(t))
  TELL(KB, MAKE-ACTION-SENTENCE(action, t))
  t ← t + 1
  return action

```

# Agents that Think Rationally

## Think Before You Act!

```
function KB-AGENT(percept) returns an action  
  persistent: KB, a knowledge base  
               t, a counter, initially 0, indicating time  
  
  TELL(KB, MAKE-PERCEPT-SENTENCE(percept, t))  
  action ← ASK(KB, MAKE-ACTION-QUERY(t))  
  TELL(KB, MAKE-ACTION-SENTENCE(action, t))  
  t ← t + 1  
  return action
```

→ "Thinking" = Reasoning about knowledge represented using logic.

## Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.



## Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.
- **Quantifying Uncertainty:** What to do when all we know is we don't know.  
→ A bit of motivation for what comes next.

## Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.
- **Quantifying Uncertainty:** What to do when all we know is we don't know.  
→ A bit of motivation for what comes next.
- **Unconditional Probabilities and Conditional Probabilities:** Which concepts and properties of probabilities will be used?  
→ Mostly a recap of things you're familiar with from school.

## Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.
- **Quantifying Uncertainty:** What to do when all we know is we don't know.  
→ A bit of motivation for what comes next.
- **Unconditional Probabilities and Conditional Probabilities:** Which concepts and properties of probabilities will be used?  
→ Mostly a recap of things you're familiar with from school.
- **Independence and Basic Probabilistic Reasoning Methods:** What simple methods are there to avoid enumeration and to deduce probabilities from other probabilities?  
→ A basic tool set we'll need. (Still familiar from school?)

## Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.
- **Quantifying Uncertainty:** What to do when all we know is we don't know.  
→ A bit of motivation for what comes next.
- **Unconditional Probabilities and Conditional Probabilities:** Which concepts and properties of probabilities will be used?  
→ Mostly a recap of things you're familiar with from school.
- **Independence and Basic Probabilistic Reasoning Methods:** What simple methods are there to avoid enumeration and to deduce probabilities from other probabilities?  
→ A basic tool set we'll need. (Still familiar from school?)
- **Bayes' Rule:** What's that "Bayes"? How is it used and why is it important?  
→ The basic insight about how to invert the "direction" of conditional probabilities.

# Our Agenda for This Chapter

- **Propositional Logic:** What's the syntax and semantics? How can we capture deduction?  
→ A brief introduction to logical reasoning.
- **Quantifying Uncertainty:** What to do when all we know is we don't know.  
→ A bit of motivation for what comes next.
- **Unconditional Probabilities and Conditional Probabilities:** Which concepts and properties of probabilities will be used?  
→ Mostly a recap of things you're familiar with from school.
- **Independence and Basic Probabilistic Reasoning Methods:** What simple methods are there to avoid enumeration and to deduce probabilities from other probabilities?  
→ A basic tool set we'll need. (Still familiar from school?)
- **Bayes' Rule:** What's that "Bayes"? How is it used and why is it important?  
→ The basic insight about how to invert the "direction" of conditional probabilities.
- **Conditional Independence:** How to capture and exploit complex relations between random variables?  
→ Explains the difficulties arising when using Bayes' rule on multiple evidences. Conditional independence is used to ameliorate these difficulties.

# Agenda

- 1 Introduction
- 2 **Propositional Logic**
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

# Propositional Logic: Syntax

## Atomic Propositions

*Boolean variables* are now seen as **atomic propositions**. Convention: start with lowercase letter.

Constraints	Logic
$A = \text{true}$	$a$
$A = \text{false}$	$\neg a$

# Propositional Logic: Syntax

## Atomic Propositions

*Boolean variables* are now seen as **atomic propositions**. Convention: start with lowercase letter.

Constraints	Logic
$A = \text{true}$	$a$
$A = \text{false}$	$\neg a$

## Propositions (Formulas)

Using **logical connectives** more complex propositions are constructed:

$\neg p$	<b>not</b> $p$
$(p \wedge q)$	$p$ <b>and</b> $q$
$(p \vee q)$	$p$ <b>or</b> $q$
$(p \rightarrow q)$	$p$ <b>implies</b> $q$

**Example:** "If it rains I'll take my umbrella, or I'll stay home"



# Propositional Logic: Syntax

## Atomic Propositions

*Boolean variables* are now seen as **atomic propositions**. Convention: start with lowercase letter.

Constraints	Logic
$A = \text{true}$	$a$
$A = \text{false}$	$\neg a$

## Propositions (Formulas)

Using **logical connectives** more complex propositions are constructed:

$\neg p$	<b>not</b> $p$
$(p \wedge q)$	$p$ <b>and</b> $q$
$(p \vee q)$	$p$ <b>or</b> $q$
$(p \rightarrow q)$	$p$ <b>implies</b> $q$

**Example:** "If it rains I'll take my umbrella, or I'll stay home"

$$\text{rains} \rightarrow (\text{umbrella} \vee \text{home})$$

## Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

# Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

An interpretation defines a truth value for all propositions:

$\pi(p)$	$\pi(\neg p)$
<i>true</i>	<i>false</i>
<i>false</i>	<i>true</i>

$\pi(p)$	$\pi(q)$	$\pi(p \wedge q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>false</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \vee q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>true</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \rightarrow q)$
<i>true</i>	<i>true</i>	
<i>true</i>	<i>false</i>	
<i>false</i>	<i>true</i>	
<i>false</i>	<i>false</i>	

# Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

An interpretation defines a truth value for all propositions:

$\pi(p)$	$\pi(\neg p)$
<i>true</i>	<i>false</i>
<i>false</i>	<i>true</i>

$\pi(p)$	$\pi(q)$	$\pi(p \wedge q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>false</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \vee q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>true</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \rightarrow q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	
<i>false</i>	<i>true</i>	
<i>false</i>	<i>false</i>	

# Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

An interpretation defines a truth value for all propositions:

$\pi(p)$	$\pi(\neg p)$
<i>true</i>	<i>false</i>
<i>false</i>	<i>true</i>

$\pi(p)$	$\pi(q)$	$\pi(p \wedge q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>false</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \vee q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>true</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \rightarrow q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	
<i>false</i>	<i>false</i>	

# Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

An interpretation defines a truth value for all propositions:

$\pi(p)$	$\pi(\neg p)$
<i>true</i>	<i>false</i>
<i>false</i>	<i>true</i>

$\pi(p)$	$\pi(q)$	$\pi(p \wedge q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>false</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \vee q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>true</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \rightarrow q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>true</i>

# Propositional Logic: Semantics I

An **interpretation**  $\pi$  for a set of atomic propositions  $a_1, a_2, \dots, a_n$  is an assignment of a truth value to each proposition (= possible world when atomic propositions seen as boolean variables):

$$\pi(a_i) \in \{true, false\}$$

An interpretation defines a truth value for all propositions:

$\pi(p)$	$\pi(\neg p)$
<i>true</i>	<i>false</i>
<i>false</i>	<i>true</i>

$\pi(p)$	$\pi(q)$	$\pi(p \wedge q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>false</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \vee q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>true</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>false</i>

$\pi(p)$	$\pi(q)$	$\pi(p \rightarrow q)$
<i>true</i>	<i>true</i>	<i>true</i>
<i>true</i>	<i>false</i>	<i>false</i>
<i>false</i>	<i>true</i>	<i>true</i>
<i>false</i>	<i>false</i>	<i>true</i>

# Knowledge as Propositional Formulas

## Satisfiability

A formula  $\varphi$  is:

- **satisfiable** if there exists  $I$  that satisfies  $\varphi$ .
- **unsatisfiable** if  $\varphi$  is not satisfiable.
- **falsifiable** if there exists  $I$  that doesn't satisfy  $\varphi$ .
- **valid** if  $I \models \varphi$  holds for all  $I$ . We also call  $\varphi$  a **tautology**.



# Knowledge as Propositional Formulas

## Satisfiability

A formula  $\varphi$  is:

- **satisfiable** if there exists  $I$  that satisfies  $\varphi$ .
- **unsatisfiable** if  $\varphi$  is not satisfiable.
- **falsifiable** if there exists  $I$  that doesn't satisfy  $\varphi$ .
- **valid** if  $I \models \varphi$  holds for all  $I$ . We also call  $\varphi$  a **tautology**.

## Knowledge Base

A **Knowledge Base (KB)** is a set of formulas that describe the agent's knowledge.

→ Knowledge Base = set of formulas, interpreted as a conjunction.

# Knowledge as Propositional Formulas

## Satisfiability

A formula  $\varphi$  is:

- **satisfiable** if there exists  $I$  that satisfies  $\varphi$ .
- **unsatisfiable** if  $\varphi$  is not satisfiable.
- **falsifiable** if there exists  $I$  that doesn't satisfy  $\varphi$ .
- **valid** if  $I \models \varphi$  holds for all  $I$ . We also call  $\varphi$  a **tautology**.

## Knowledge Base

A **Knowledge Base (KB)** is a set of formulas that describe the agent's knowledge.

→ **Knowledge Base = set of formulas, interpreted as a conjunction.**

**Definition (Model).** A **model** of a knowledge base KB is an interpretation  $I$  in which all the formulas in the knowledge base are true:  $I \models \varphi$  for all  $\varphi \in \text{KB}$ .

→ a model is a possible world that satisfies the constraint.

We denote by  $M(\varphi)$  the set of all models of  $\varphi$  (i.e., the set of possible worlds where the formula is true).

# Deduction

## Deduction

deriving of a conclusion by reasoning

**Remember (slide 4)?** Does our knowledge of the cave entail a definite Wumpus position?

→ We don't know everything; what can we conclude from the things we *do* know?

# Deduction

## Deduction

deriving of a conclusion by reasoning

**Remember (slide 4)?** Does our knowledge of the cave entail a definite Wumpus position?

→ We don't know everything; what can we conclude from the things we *do* know?

## Logical consequence (entailment)

**Definition (Entailment).** Let  $\Sigma$  be a set of atomic propositions. We say that a set of formulas  $KB$  *entails* a formula  $\varphi$ , written  $KB \models \varphi$ , if  $\varphi$  is true in all models of  $KB$ , i.e.,  $M(\bigwedge_{\psi \in KB}) \subseteq M(\varphi)$ . In this case, we also say that  $\varphi$  *follows* from  $KB$ .

# Deduction

## Deduction

deriving of a conclusion by reasoning

**Remember (slide 4)?** Does our knowledge of the cave entail a definite Wumpus position?

→ We don't know everything; what can we conclude from the things we *do* know?

## Logical consequence (entailment)

**Definition (Entailment).** Let  $\Sigma$  be a set of atomic propositions. We say that a set of formulas  $KB$  *entails* a formula  $\varphi$ , written  $KB \models \varphi$ , if  $\varphi$  is true in all models of  $KB$ , i.e.,  $M(\bigwedge_{\psi \in KB}) \subseteq M(\varphi)$ . In this case, we also say that  $\varphi$  *follows* from  $KB$ .

A formula  $\varphi$  is a **logical consequence** of a knowledge base  $KB$ , if every model of  $KB$  is a model of  $\varphi$ . Written:

$$KB \models \varphi$$

(whenever  $KB$  is true, then  $\varphi$  also is true).

Example:  $KB = \{man \rightarrow mortal, man\}$ . Then

# Deduction

## Deduction

deriving of a conclusion by reasoning

**Remember (slide 4)?** Does our knowledge of the cave entail a definite Wumpus position?

→ We don't know everything; what can we conclude from the things we *do* know?

## Logical consequence (entailment)

**Definition (Entailment).** Let  $\Sigma$  be a set of atomic propositions. We say that a set of formulas  $KB$  *entails* a formula  $\varphi$ , written  $KB \models \varphi$ , if  $\varphi$  is true in all models of  $KB$ , i.e.,  $M(\bigwedge_{\psi \in KB}) \subseteq M(\varphi)$ . In this case, we also say that  $\varphi$  *follows* from  $KB$ .

A formula  $\varphi$  is a **logical consequence** of a knowledge base  $KB$ , if every model of  $KB$  is a model of  $\varphi$ . Written:

$$KB \models \varphi$$

(whenever  $KB$  is true, then  $\varphi$  also is true).

Example:  $KB = \{man \rightarrow mortal, man\}$ . Then  $KB \models mortal$

# Simple Example

$$KB = \begin{cases} p \leftarrow q. \\ q. \\ r \leftarrow s. \end{cases}$$

	<i>p</i>	<i>q</i>	<i>r</i>	<i>s</i>	Model?
<i>I</i> <sub>1</sub>	<i>true</i>	<i>true</i>	<i>true</i>	<i>true</i>	
<i>I</i> <sub>2</sub>	<i>false</i>	<i>false</i>	<i>false</i>	<i>false</i>	
<i>I</i> <sub>3</sub>	<i>true</i>	<i>true</i>	<i>false</i>	<i>false</i>	
<i>I</i> <sub>4</sub>	<i>true</i>	<i>true</i>	<i>true</i>	<i>false</i>	
<i>I</i> <sub>5</sub>	<i>true</i>	<i>true</i>	<i>false</i>	<i>true</i>	
...					

# Simple Example

$$KB = \begin{cases} p \leftarrow q. \\ q. \\ r \leftarrow s. \end{cases}$$

	$p$	$q$	$r$	$s$	Model?
$I_1$	<i>true</i>	<i>true</i>	<i>true</i>	<i>true</i>	is a model of $KB$
$I_2$	<i>false</i>	<i>false</i>	<i>false</i>	<i>false</i>	not a model of $KB$
$I_3$	<i>true</i>	<i>true</i>	<i>false</i>	<i>false</i>	is a model of $KB$
$I_4$	<i>true</i>	<i>true</i>	<i>true</i>	<i>false</i>	is a model of $KB$
$I_5$	<i>true</i>	<i>true</i>	<i>false</i>	<i>true</i>	not a model of $KB$
...					



## Simple Example

$$KB = \begin{cases} p \leftarrow q. \\ q. \\ r \leftarrow s. \end{cases}$$

	$p$	$q$	$r$	$s$	Model?
$I_1$	<i>true</i>	<i>true</i>	<i>true</i>	<i>true</i>	is a model of $KB$
$I_2$	<i>false</i>	<i>false</i>	<i>false</i>	<i>false</i>	not a model of $KB$
$I_3$	<i>true</i>	<i>true</i>	<i>false</i>	<i>false</i>	is a model of $KB$
$I_4$	<i>true</i>	<i>true</i>	<i>true</i>	<i>false</i>	is a model of $KB$
$I_5$	<i>true</i>	<i>true</i>	<i>false</i>	<i>true</i>	not a model of $KB$
...					

Which of  $p, q, r, s$  logically follow from  $KB$ ?

## Simple Example

$$KB = \begin{cases} p \leftarrow q. \\ q. \\ r \leftarrow s. \end{cases}$$

	$p$	$q$	$r$	$s$	Model?
$I_1$	<i>true</i>	<i>true</i>	<i>true</i>	<i>true</i>	is a model of $KB$
$I_2$	<i>false</i>	<i>false</i>	<i>false</i>	<i>false</i>	not a model of $KB$
$I_3$	<i>true</i>	<i>true</i>	<i>false</i>	<i>false</i>	is a model of $KB$
$I_4$	<i>true</i>	<i>true</i>	<i>true</i>	<i>false</i>	is a model of $KB$
$I_5$	<i>true</i>	<i>true</i>	<i>false</i>	<i>true</i>	not a model of $KB$
...					

Which of  $p, q, r, s$  logically follow from  $KB$ ?

$$KB \models p, KB \models q, KB \not\models r, KB \not\models s$$

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell  $(2,2)$ . Can we conclude the cell  $(2,2)$  is free?**

(A): Yes

(B): No

## Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

### Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell  $(2,2)$ . Can we conclude the cell  $(2,2)$  is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell  $(2,2)$ . Can we conclude the cell  $(2,2)$  is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell  $(2,2)$ . Can we conclude the cell  $(2,2)$  is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ Entailment can be tested via satisfiability.

To deduce “Cell  $(2,2)$  is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell (2,2). Can we conclude the cell (2,2) is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ Entailment can be tested via satisfiability.

To deduce “Cell (2,2) is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

Does there exist an interpretation that agrees with all formulas in our KB?

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell (2,2). Can we conclude the cell (2,2) is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ Entailment can be tested via satisfiability.

To deduce “Cell (2,2) is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

Does there exist an interpretation that agrees with all formulas in our KB?

Any deja vu?



# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell (2,2). Can we conclude the cell (2,2) is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ Entailment can be tested via satisfiability.

To deduce “Cell (2,2) is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

Does there exist an interpretation that agrees with all formulas in our KB?

Any deja vu?

Does there exist a variable assignment that agrees with all constraints?

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<input type="checkbox"/>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell (2,2). Can we conclude the cell (2,2) is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ Entailment can be tested via satisfiability.

To deduce “Cell (2,2) is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

Does there exist an interpretation that agrees with all formulas in our KB?

Any deja vu?

Does there exist a variable assignment that agrees with all constraints? CSP!

# Proof by Contradiction

1,4	2,4	3,4	4,4
1,3	2,3	3,3	4,3
1,2	2,2	3,2	4,2
OK			
1,1	2,1	3,1	4,1
<div>A</div>			
OK	OK		

## Question!

**Suppose that there exists an interpretation  $I$  in  $M(KB)$  where the Wumpus is not at cell  $(2,2)$ . Can we conclude the cell  $(2,2)$  is free?**

(A): Yes

(B): No

No, it could not be there, but that does not mean that it is not there!

**Contradiction Theorem.**  $KB \models \varphi$  if and only if  $KB \cup \{\neg\varphi\}$  is unsatisfiable.

→ **Entailment can be tested via satisfiability.**

To deduce “Cell  $(2,2)$  is free”, assume the opposite “ $\neg(\text{cell}(2,2) \text{ is free})$ ” look for an interpretation. If none exists, you can conclude the opposite of your assumption.

Does there exist an interpretation that agrees with all formulas in our KB?

Any deja vu?

Does there exist a variable assignment that agrees with all constraints? **CSP!**

In this course, we don't cover algorithms for testing satisfiability (this is part of the Algorithms and Satisfiability course on DAT6). But the principles are similar to what we covered in **Chapter 3**.

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty**
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

## Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

## Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about: *toothache* → *cavity*

→ Is this rule correct?

## Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about: *toothache* → *cavity*

→ Is this rule correct? No, toothaches may have different causes.

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about: *toothache* → *cavity*

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this: *toothache* → *cavity* ∨ *gum\_disease* ∨ ...



## Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about: *toothache* → *cavity*

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this: *toothache* → *cavity* ∨ *gum\_disease* ∨ ...

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about:  $toothache \rightarrow cavity$

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this:  $toothache \rightarrow cavity \vee gum\_disease \vee \dots$

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

**Attempt 3:** Perhaps a *causal* rule is better?  $cavity \rightarrow toothache$

- Is this rule correct?

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about:  $toothache \rightarrow cavity$

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this:  $toothache \rightarrow cavity \vee gum\_disease \vee \dots$

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

**Attempt 3:** Perhaps a *causal* rule is better?  $cavity \rightarrow toothache$

- Is this rule correct? No, not all cavities cause toothaches.
- Does this rule allow to deduce a cause from a symptom?

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about:  $toothache \rightarrow cavity$

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this:  $toothache \rightarrow cavity \vee gum\_disease \vee \dots$

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

**Attempt 3:** Perhaps a *causal* rule is better?  $cavity \rightarrow toothache$

- Is this rule correct? No, not all cavities cause toothaches.
- Does this rule allow to deduce a cause from a symptom? No, setting *toothache* to true here has no consequence on the truth of *cavity*.

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about:  $toothache \rightarrow cavity$

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this:  $toothache \rightarrow cavity \vee gum\_disease \vee \dots$

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

**Attempt 3:** Perhaps a *causal* rule is better?  $cavity \rightarrow toothache$

- Is this rule correct? No, not all cavities cause toothaches.
- Does this rule allow to deduce a cause from a symptom? No, setting *toothache* to true here has no consequence on the truth of *cavity*. [Note: If *toothache* is *false*, we would conclude  $\neg cavity$  ... which would be incorrect, cf. previous question.]

# Uncertainty and Logic

**Diagnosis:** We want to build an expert dental diagnosis system, that deduces the cause (the disease) from the symptoms.

→ Can we base this on logic?

**Attempt 1:** Say we have a toothache. How's about:  $toothache \rightarrow cavity$

→ Is this rule correct? No, toothaches may have different causes.

**Attempt 2:** So what about this:  $toothache \rightarrow cavity \vee gum\_disease \vee \dots$

→ We don't know all possible causes. And we'd like to be able to deduce which causes are more plausible!

**Attempt 3:** Perhaps a *causal* rule is better?  $cavity \rightarrow toothache$

- Is this rule correct? No, not all cavities cause toothaches.
- Does this rule allow to deduce a cause from a symptom? No, setting *toothache* to true here has no consequence on the truth of *cavity*. [Note: If *toothache* is *false*, we would conclude  $\neg cavity \dots$  which would be incorrect, cf. previous question.]
- Anyway, this still doesn't allow to compare the plausibility of different causes.

→ Logic does not allow to weigh different alternatives, and it does not allow to express incomplete knowledge ("cavity does not always come with a toothache, nor vice versa").

## Beliefs and Probabilities

**What do we model with probabilities?**

## Beliefs and Probabilities

**What do we model with probabilities?** Incomplete knowledge! We are not 100% sure, but we *believe to a certain degree* that something is true.

→ Probability  $\approx$  Our degree of belief, given our current knowledge.



## Beliefs and Probabilities

**What do we model with probabilities?** Incomplete knowledge! We are not 100% sure, but we *believe to a certain degree* that something is true.

→ Probability  $\approx$  Our degree of belief, given our current knowledge.

### Example (Diagnosis)

- *toothache*  $\rightarrow$  *cavity* with 80% probability.

## Beliefs and Probabilities

**What do we model with probabilities?** Incomplete knowledge! We are not 100% sure, but we *believe to a certain degree* that something is true.

→ Probability  $\approx$  Our degree of belief, given our current knowledge.

### Example (Diagnosis)

- *toothache*  $\rightarrow$  *cavity* with 80% probability.
- But, for any given  $p$ , in reality we do, or do not, have cavity: 1 or 0!

## Beliefs and Probabilities

**What do we model with probabilities?** Incomplete knowledge! We are not 100% sure, but we *believe to a certain degree* that something is true.

→ Probability  $\approx$  Our degree of belief, given our current knowledge.

### Example (Diagnosis)

- *toothache* → *cavity* with 80% probability.
- But, for any given  $p$ , in reality we do, or do not, have cavity: 1 or 0!  
 → The “probability” depends on our knowledge! The “80%” refers to **the fraction of cavity, within the set of all persons that are indistinguishable from the one being evaluated based on our knowledge.**

## Beliefs and Probabilities

**What do we model with probabilities?** Incomplete knowledge! We are not 100% sure, but we *believe to a certain degree* that something is true.

→ Probability  $\approx$  Our degree of belief, given our current knowledge.

### Example (Diagnosis)

- *toothache* → *cavity* with 80% probability.
- But, for any given  $p$ , in reality we do, or do not, have cavity: 1 or 0!  
→ The “probability” depends on our knowledge! The “80%” refers to **the fraction of cavity, within the set of all persons that are indistinguishable from the one being evaluated based on our knowledge.**
- If we receive new knowledge (e.g., if we know *gum\_disease*) is true, the probability of *cavity* changes!

→ Probabilities represent and measure the uncertainty that stems from lack of knowledge.

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus**
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion


# Probability Measures

Probability theory is built on the foundation of variables and worlds.

Worlds described by the variables:

- Filled: {true, false}
- Shape: {circle, triangle, star}

as well as position.



## Probability measures

$\Omega$ : set of all possible worlds (for a given, fixed set of variables). A **probability measure over  $\Omega$** , is a function  $P$ , that assigns **probability values**

$$P(\Omega') \in [0, 1]$$

to subsets  $\Omega' \subseteq \Omega$ , such that

**Axiom 1:**  $P(\Omega) = 1$ .

**Axiom 2:** if  $\Omega_1 \cap \Omega_2 = \emptyset$ , then  $P(\Omega_1 \cup \Omega_2) = P(\Omega_1) + P(\Omega_2)$ .

## Simplification for finite $\Omega$

If all variables have a finite domain, then

- $\Omega$  is finite, and
- a probability distribution is defined by assigning a probability value  $P(\omega)$  to each individual possible world  $\omega \in \Omega$ .


## Simplification for finite $\Omega$

If all variables have a finite domain, then

- $\Omega$  is finite, and
- a probability distribution is defined by assigning a probability value  $P(\omega)$  to each individual possible world  $\omega \in \Omega$ .

For any  $\Omega' \subseteq \Omega$  then

$$P(\Omega') = \sum_{\omega \in \Omega'} P(\omega)$$



$$P(\Omega') = 0.08 + 0.13 + 0.03 + 0.21 = 0.45$$




## Simplification for finite $\Omega$

If all variables have a finite domain, then

- $\Omega$  is finite, and
- a probability distribution is defined by assigning a probability value  $P(\omega)$  to each individual possible world  $\omega \in \Omega$ .

For any  $\Omega' \subseteq \Omega$  then

$$P(\Omega') = \sum_{\omega \in \Omega'} P(\omega)$$



$$P(\Omega') = 0.08 + 0.13 + 0.03 + 0.21 = 0.45$$

**Note:** In general, random variables can have arbitrary domains. Here, we consider **finite-domain** random variables only, and **Boolean** random variables most of the time.

# Random Variables and Distributions

**Definition (Random Variables).** Variables defining possible worlds on which probabilities are defined are called **random variables**.

## Distributions

For a random variable  $A$ , and  $a \in D_A$  we have the probability

$$P(A = a) = P(\{\omega \in \Omega \mid A = a \text{ in } \omega\})$$

# Random Variables and Distributions

**Definition (Random Variables).** Variables defining possible worlds on which probabilities are defined are called **random variables**.

## Distributions

For a random variable  $A$ , and  $a \in D_A$  we have the probability

$$P(A = a) = P(\{\omega \in \Omega \mid A = a \text{ in } \omega\})$$

The **probability distribution of**  $A$  (written  $P(A)$ ) is the function on  $D_A$  that maps each  $a$  to its probability  $P(A = a)$ .

**Example:**

$$\mathbf{P}(\textit{Headache}) = \langle F \mapsto 0.1, T \mapsto 0.9 \rangle$$

$$\mathbf{P}(\textit{Weather}) = \langle \textit{sunny} \mapsto 0.7, \textit{rain} \mapsto 0.2, \textit{cloudy} \mapsto 0.08, \textit{snow} \mapsto 0.02 \rangle$$

# Joint Probability Distributions

Extension to several random variables:

$$P(A_1, \dots, A_k)$$

is the **joint distribution** of  $A_1, \dots, A_k$ . The joint distribution maps tuples  $(a_1, \dots, a_k)$  with  $a_i \in D_{A_i}$  to the probability

$$P(A_1 = a_1, \dots, A_k = a_k)$$

**Example:**  $\mathbf{P}(\text{Headache}, \text{Weather}) =$

	<i>Headache = true</i>	<i>Headache = false</i>
<i>Weather = sunny</i>	$P(W = \text{sunny} \wedge \text{headache})$	$P(W = \text{sunny} \wedge \neg \text{headache})$
<i>Weather = rain</i>		
<i>Weather = cloudy</i>		
<i>Weather = snow</i>		

## Terminology:

- Given random variables  $\{X_1, \dots, X_n\}$ , an **atomic event (world)** is an assignment of values to all variables.
- Given random variables  $\{X_1, \dots, X_n\}$ , the **full joint probability distribution**, denoted  $\mathbf{P}(X_1, \dots, X_n)$ , lists the probabilities of all atomic events.

→ All worlds are disjoint (their pairwise conjunctions all are  $\perp$ ); the sum of all fields is 1 (corresponds to their disjunction  $\top$ ).

## Probabilities of propositions

A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .

### Example



Assume probability for each world is 0.1:

•  $P(\text{Shape} = \text{circle}) =$

## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .

### Example



Assume probability for each world is 0.1:

- $P(\text{Shape} = \text{circle}) = 0.5$
- $P(\text{Filled} = \text{false}) =$

## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .

### Example



Assume probability for each world is 0.1:

- $P(\text{Shape} = \text{circle}) = 0.5$
- $P(\text{Filled} = \text{false}) = 0.4$
- $P(\text{Shape} = c \wedge \text{Filled} = f) =$



## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .

### Example



Assume probability for each world is 0.1:

- $P(\text{Shape} = \text{circle}) = 0.5$
- $P(\text{Filled} = \text{false}) = 0.4$
- $P(\text{Shape} = c \wedge \text{Filled} = f) = 0.1$

## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .


### Example



Assume probability for each world is 0.1:

- $P(\text{Shape} = \text{circle}) = 0.5$
- $P(\text{Filled} = \text{false}) = 0.4$
- $P(\text{Shape} = c \wedge \text{Filled} = f) = 0.1$

### Another example



## Probabilities of propositions


A probability distribution over possible worlds defines probabilities for formulas  $\varphi$ :

$$P(\alpha) = \sum_{\omega \in \Omega: \omega \in M(\varphi)} P(\omega)$$

→ Propositions represent sets of atomic events: the interpretations satisfying the formula.

**Notation:** Instead of  $P(a \wedge b)$ , we often write  $P(a, b)$ .


### Example



Assume probability for each world is 0.1:

- $P(\text{Shape} = \text{circle}) = 0.5$
- $P(\text{Filled} = \text{false}) = 0.4$
- $P(\text{Shape} = c \wedge \text{Filled} = f) = 0.1$

### Another example



$$\begin{aligned} P(\text{Color} = \text{red}) &= 0.08 + 0.13 + 0.03 + 0.21 \\ &= 0.45 \end{aligned}$$

## Basic probability axioms

### Axiom

If  $\mathcal{A}$  and  $\mathcal{B}$  are disjoint, then  $P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B})$ .

### Example

Consider a deck with 52 cards. If  $\mathcal{A} = \{2, 3, 4, 5\}$  and  $\mathcal{B} = \{7, 8\}$ , then

$$P(\mathcal{A} \cup \mathcal{B}) =$$

## Basic probability axioms

### Axiom

If  $\mathcal{A}$  and  $\mathcal{B}$  are disjoint, then  $P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B})$ .

### Example

Consider a deck with 52 cards. If  $\mathcal{A} = \{2, 3, 4, 5\}$  and  $\mathcal{B} = \{7, 8\}$ , then

$$P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B}) = \frac{4}{13} + \frac{2}{13} = \frac{6}{13}.$$

# Basic probability axioms

## Axiom

If  $\mathcal{A}$  and  $\mathcal{B}$  are disjoint, then  $P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B})$ .

## Example

Consider a deck with 52 cards. If  $\mathcal{A} = \{2, 3, 4, 5\}$  and  $\mathcal{B} = \{7, 8\}$ , then

$$P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B}) = \frac{4}{13} + \frac{2}{13} = \frac{6}{13}.$$


## More generally

If  $\mathcal{C}$  and  $\mathcal{D}$  are not disjoint, then  $P(\mathcal{C} \cup \mathcal{D}) = P(\mathcal{C}) + P(\mathcal{D}) - P(\mathcal{C} \cap \mathcal{D})$ .

## Example

If  $\mathcal{C} = \{2, 3, 4, 5\}$  and  $\mathcal{D} = \{\spadesuit\}$ , then

$$P(\mathcal{C} \cup \mathcal{D}) =$$



# Basic probability axioms

## Axiom

If  $\mathcal{A}$  and  $\mathcal{B}$  are disjoint, then  $P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B})$ .

## Example

Consider a deck with 52 cards. If  $\mathcal{A} = \{2, 3, 4, 5\}$  and  $\mathcal{B} = \{7, 8\}$ , then

$$P(\mathcal{A} \cup \mathcal{B}) = P(\mathcal{A}) + P(\mathcal{B}) = \frac{4}{13} + \frac{2}{13} = \frac{6}{13}.$$


## More generally

If  $\mathcal{C}$  and  $\mathcal{D}$  are not disjoint, then  $P(\mathcal{C} \cup \mathcal{D}) = P(\mathcal{C}) + P(\mathcal{D}) - P(\mathcal{C} \cap \mathcal{D})$ .

## Example

If  $\mathcal{C} = \{2, 3, 4, 5\}$  and  $\mathcal{D} = \{\spadesuit\}$ , then

$$P(\mathcal{C} \cup \mathcal{D}) = \frac{4}{13} + \frac{1}{4} - \frac{4}{52} = \frac{25}{52}.$$



# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities**
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion



## Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?

## Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?

Yes! Probabilities model our *belief*, thus they depend on our knowledge. Given new information (**evidence**), degrees of belief change.

**Example:** Your “probability of missing the connection train” increases when you are informed that your current train has 30 minutes delay.

## Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?

Yes! Probabilities model our *belief*, thus they depend on our knowledge. Given new information (**evidence**), degrees of belief change.

**Example:** Your “probability of missing the connection train” increases when you are informed that your current train has 30 minutes delay.

Evidence can consist of:

- learning that a certain proposition  $p$  is true (“*switch\_up*”)
- obtaining partial information on the value of some variable (“ $\text{room\_ai} \neq 0.2.90$ ”)
- ...

# Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?


Yes! Probabilities model our *belief*, thus they depend on our knowledge. Given new information (**evidence**), degrees of belief change.

**Example:** Your “probability of missing the connection train” increases when you are informed that your current train has 30 minutes delay.

Evidence can consist of:

- learning that a certain proposition  $p$  is true (“*switch\_up*”)
- obtaining partial information on the value of some variable (“*room\_ai*  $\neq$  0.2.90”)
- ...

→ In all cases: **evidence can be represented as the set of possible world  $\Omega'$  not ruled out by the observation.** How should the probabilities be updated, when I observe  $\Omega'$ ?:



# Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?


Yes! Probabilities model our *belief*, thus they depend on our knowledge. Given new information (**evidence**), degrees of belief change.

**Example:** Your “probability of missing the connection train” increases when you are informed that your current train has 30 minutes delay.

Evidence can consist of:

- learning that a certain proposition  $p$  is true (“*switch\_up*”)
- obtaining partial information on the value of some variable (“*room\_ai*  $\neq$  0.2.90”)
- ...

→ In all cases: **evidence can be represented as the set of possible world  $\Omega'$  not ruled out by the observation.** How should the probabilities be updated, when I observe  $\Omega'$ ?:



- worlds that are not consistent with evidence have probability 0

# Updating Your Beliefs

→ Do probabilities change as we gather new knowledge?


Yes! Probabilities model our *belief*, thus they depend on our knowledge. Given new information (**evidence**), degrees of belief change.

**Example:** Your “probability of missing the connection train” increases when you are informed that your current train has 30 minutes delay.

Evidence can consist of:

- learning that a certain proposition  $p$  is true (“*switch\_up*”)
- obtaining partial information on the value of some variable (“*room\_ai*  $\neq$  0.2.90”)
- ...

→ In all cases: **evidence can be represented as the set of possible world  $\Omega'$  not ruled out by the observation.** How should the probabilities be updated, when I observe  $\Omega'$ ?:



- worlds that are not consistent with evidence have probability 0
- the probabilities of worlds consistent with evidence are proportional to their probability before observation, and they must sum to 1

## Conditional Probabilities

**Definition.** Given propositions  $p$  and  $e$  where  $P(e) \neq 0$ , the *conditional probability*, or *posterior probability*, of  $p$  given  $e$ , written  $P(p \mid e)$ , is defined as:

$$P(p \mid e) = \frac{P(p \wedge e)}{P(e)}$$

→ The likelihood of having  $p$  **and**  $e$ , **within** the set of outcomes where we have  $e$ .


## Conditional Probabilities

**Definition.** Given propositions  $p$  and  $e$  where  $P(e) \neq 0$ , the *conditional probability*, or *posterior probability*, of  $p$  given  $e$ , written  $P(p \mid e)$ , is defined as:

$$P(p \mid e) = \frac{P(p \wedge e)}{P(e)}$$

→ The likelihood of having  $p$  **and**  $e$ , **within** the set of outcomes where we have  $e$ .

### Example



(probability for each world is 0.1)

$$\begin{aligned}
 P(S=\text{circ.} \mid \text{Fill} = f) &= \frac{P(S=\text{circ.} \wedge \text{Fill} = f)}{P(\text{Fill} = f)} \\
 &= \frac{0.1}{0.4} = 0.25
 \end{aligned}$$




## Conditional Probabilities

**Definition.** Given propositions  $p$  and  $e$  where  $P(e) \neq 0$ , the *conditional probability*, or *posterior probability*, of  $p$  given  $e$ , written  $P(p \mid e)$ , is defined as:

$$P(p \mid e) = \frac{P(p \wedge e)}{P(e)}$$

→ The likelihood of having  $p$  and  $e$ , within the set of outcomes where we have  $e$ .

### Example



(probability for each world is 0.1)

$$\begin{aligned} P(S=circ. \mid Fill = f) &= \frac{P(S=circ. \wedge Fill = f)}{P(Fill = f)} \\ &= \frac{0.1}{0.4} = 0.25 \end{aligned}$$

What is the probability of  $P(S=star \mid Fill = f)$ ?


# Conditional Probabilities

**Definition.** Given propositions  $p$  and  $e$  where  $P(e) \neq 0$ , the *conditional probability*, or *posterior probability*, of  $p$  given  $e$ , written  $P(p \mid e)$ , is defined as:

$$P(p \mid e) = \frac{P(p \wedge e)}{P(e)}$$

→ The likelihood of having  $p$  and  $e$ , within the set of outcomes where we have  $e$ .

## Example




(probability for each world is 0.1)

$$\begin{aligned} P(S=circ. \mid Fill = f) &= \frac{P(S=circ. \wedge Fill = f)}{P(Fill = f)} \\ &= \frac{0.1}{0.4} = 0.25 \end{aligned}$$

What is the probability of  $P(S=star \mid Fill = f)$ ?

## Another example



- $e$  and  $p$  are represented by possible worlds  $\Omega_1$  and  $\Omega_2$
- division by  $P(\Omega_1)$  already in green numbers

$$P(\Omega_2 \mid \Omega_1) = 0.18 + 0.29$$

## Conditional Probability Distributions

**Definition.** Given random variables  $X$  and  $Y$ , the *conditional probability distribution* of  $X$  given  $Y$ , written  $\mathbf{P}(X \mid Y)$ , is the table of all conditional probabilities of values of  $X$  given values of  $Y$ .

→ For sets of variables:  $\mathbf{P}(X_1, \dots, X_n \mid Y_1, \dots, Y_m)$ .

# Conditional Probability Distributions

**Definition.** Given random variables  $X$  and  $Y$ , the *conditional probability distribution* of  $X$  given  $Y$ , written  $\mathbf{P}(X \mid Y)$ , is the table of all conditional probabilities of values of  $X$  given values of  $Y$ .

→ For sets of variables:  $\mathbf{P}(X_1, \dots, X_n \mid Y_1, \dots, Y_m)$ .

**Example:**  $\mathbf{P}(\text{Weather} \mid \text{Headache}) =$

	<i>Headache = true</i>	<i>Headache = false</i>
<i>Weather = sunny</i>	$P(W = \text{sunny} \mid \text{headache})$	$P(W = \text{sunny} \mid \neg \text{headache})$
<i>Weather = rain</i>		
<i>Weather = cloudy</i>		
<i>Weather = snow</i>		

# Conditional Probability Distributions

**Definition.** Given random variables  $X$  and  $Y$ , the *conditional probability distribution* of  $X$  given  $Y$ , written  $\mathbf{P}(X \mid Y)$ , is the table of all conditional probabilities of values of  $X$  given values of  $Y$ .

→ For sets of variables:  $\mathbf{P}(X_1, \dots, X_n \mid Y_1, \dots, Y_m)$ .

**Example:**  $\mathbf{P}(\text{Weather} \mid \text{Headache}) =$

	<i>Headache = true</i>	<i>Headache = false</i>
<i>Weather = sunny</i>	$P(W = \text{sunny} \mid \text{headache})$	$P(W = \text{sunny} \mid \neg \text{headache})$
<i>Weather = rain</i>		
<i>Weather = cloudy</i>		
<i>Weather = snow</i>		

→ "The probability of sunshine given that I have a headache?"

# Conditional Probability Distributions

**Definition.** Given random variables  $X$  and  $Y$ , the *conditional probability distribution* of  $X$  given  $Y$ , written  $\mathbf{P}(X \mid Y)$ , is the table of all conditional probabilities of values of  $X$  given values of  $Y$ .

→ For sets of variables:  $\mathbf{P}(X_1, \dots, X_n \mid Y_1, \dots, Y_m)$ .

**Example:**  $\mathbf{P}(\text{Weather} \mid \text{Headache}) =$

	<i>Headache = true</i>	<i>Headache = false</i>
<i>Weather = sunny</i>	$P(W = \text{sunny} \mid \text{headache})$	$P(W = \text{sunny} \mid \neg \text{headache})$
<i>Weather = rain</i>		
<i>Weather = cloudy</i>		
<i>Weather = snow</i>		

→ "The probability of sunshine given that I have a headache?" If you're susceptible to headaches depending on weather conditions, this makes sense. Otherwise, the two variables are *independent* (we'll get to this later).

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods**
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

# The Product Rule

**Proposition (Product Rule).** *Given propositions  $A$  and  $B$ ,  $P(a \wedge b) = P(a \mid b)P(b)$ . (Direct from definition.)*

**Example:**  $P(\text{cavity} \wedge \text{toothache}) = P(\text{toothache} \mid \text{cavity})P(\text{cavity})$ .



# The Product Rule

**Proposition (Product Rule).** *Given propositions  $A$  and  $B$ ,  $P(a \wedge b) = P(a \mid b)P(b)$ . (Direct from definition.)*

**Example:**  $P(\text{cavity} \wedge \text{toothache}) = P(\text{toothache} \mid \text{cavity})P(\text{cavity})$ .

→ If we know the values of  $P(a \mid b)$  and  $P(b)$ , then we can compute  $P(a \wedge b)$ .

→ Similarly,  $P(a \wedge b) = P(b \mid a)P(a)$ .

# The Product Rule

**Proposition (Product Rule).** Given propositions  $A$  and  $B$ ,  $P(a \wedge b) = P(a | b)P(b)$ .  
(Direct from definition.)

**Example:**  $P(\text{cavity} \wedge \text{toothache}) = P(\text{toothache} | \text{cavity})P(\text{cavity})$ .

→ If we know the values of  $P(a | b)$  and  $P(b)$ , then we can compute  $P(a \wedge b)$ .

→ Similarly,  $P(a \wedge b) = P(b | a)P(a)$ .

**Notation:**  $\mathbf{P}(X, Y) = \mathbf{P}(X | Y)\mathbf{P}(Y)$  is a **system of equations**:

$$\begin{aligned}
 P(W = \text{sunny} \wedge \text{headache}) &= P(W = \text{sunny} | \text{headache})P(\text{headache}) \\
 P(W = \text{rain} \wedge \text{headache}) &= P(W = \text{rain} | \text{headache})P(\text{headache}) \\
 \dots &= \dots \\
 P(W = \text{snow} \wedge \neg \text{headache}) &= P(W = \text{snow} | \neg \text{headache})P(\neg \text{headache})
 \end{aligned}$$

→ Similar for unconditional distributions,  $\mathbf{P}(X, Y) = \mathbf{P}(X)\mathbf{P}(Y)$ .

# The Chain Rule

**Proposition (Chain Rule).** *Given random variables  $X_1, \dots, X_n$ , we have*

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_2 \mid X_1) * \mathbf{P}(X_1).$$

**Example:**  $P(\neg brush \wedge cavity \wedge toothache)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity, \neg brush)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity \mid \neg brush)P(\neg brush).$

# The Chain Rule

**Proposition (Chain Rule).** *Given random variables  $X_1, \dots, X_n$ , we have*

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_2 \mid X_1) * \mathbf{P}(X_1).$$

**Example:**  $P(\neg brush \wedge cavity \wedge toothache)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity, \neg brush)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity \mid \neg brush)P(\neg brush).$

**Proof.** Iterated application of Product Rule.

# The Chain Rule

**Proposition (Chain Rule).** *Given random variables  $X_1, \dots, X_n$ , we have*

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_2 \mid X_1) * \mathbf{P}(X_1).$$

**Example:**  $P(\neg brush \wedge cavity \wedge toothache)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity, \neg brush)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity \mid \neg brush)P(\neg brush).$

**Proof.** Iterated application of Product Rule.  $\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1}, \dots, X_1)$  by Product Rule. In turn,  $\mathbf{P}(X_{n-1}, \dots, X_1) = \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \mathbf{P}(X_{n-2}, \dots, X_1)$ , etc.

# The Chain Rule

**Proposition (Chain Rule).** Given random variables  $X_1, \dots, X_n$ , we have

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_2 \mid X_1) * \mathbf{P}(X_1).$$

**Example:**  $P(\neg brush \wedge cavity \wedge toothache)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity, \neg brush)$   
 $= P(toothache \mid cavity, \neg brush)P(cavity \mid \neg brush)P(\neg brush).$

**Proof.** Iterated application of Product Rule.  $\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1}, \dots, X_1)$  by Product Rule. In turn,  $\mathbf{P}(X_{n-1}, \dots, X_1) = \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \mathbf{P}(X_{n-2}, \dots, X_1)$ , etc.

**Note:** This works *for any ordering* of the variables.

→ We can recover the probability of atomic events from sequenced conditional probabilities for any ordering of the variables.

→ First of the four basic techniques in Bayesian networks.

# Marginalization

→ Extracting a sub-distribution from a larger joint distribution:

**Proposition (Marginalization).** *Given sets  $\mathbf{X}$  and  $\mathbf{Y}$  of random variables, we have:*

$$P(\mathbf{X}) = \sum_{\mathbf{y} \in \mathbf{Y}} P(\mathbf{X}, \mathbf{y})$$

where  $\sum_{\mathbf{y} \in \mathbf{Y}}$  sums over all possible value combinations of  $\mathbf{Y}$ .

# Marginalization

→ Extracting a sub-distribution from a larger joint distribution:

**Proposition (Marginalization).** *Given sets  $\mathbf{X}$  and  $\mathbf{Y}$  of random variables, we have:*

$$\mathbf{P}(\mathbf{X}) = \sum_{\mathbf{y} \in \mathbf{Y}} \mathbf{P}(\mathbf{X}, \mathbf{y})$$

where  $\sum_{\mathbf{y} \in \mathbf{Y}}$  sums over all possible value combinations of  $\mathbf{Y}$ .

**Example:** (Note: Equation system!)

$$\mathbf{P}(\text{Cavity}) = \sum_{y \in \text{Toothache}} \mathbf{P}(\text{Cavity}, y)$$

$$P(\text{cavity}) = P(\text{cavity}, \text{toothache}) + P(\text{cavity}, \neg \text{toothache})$$

$$P(\neg \text{cavity}) = P(\neg \text{cavity}, \text{toothache}) + P(\neg \text{cavity}, \neg \text{toothache})$$



## Questionnaire

### Question!

**Say**  $P(dog) = 0.4$ ,  $\neg dog \leftrightarrow cat$ , **and**  $P(likeslasagna \mid cat) = 0.5$ . **Then**  
 $P(likeslasagna \wedge cat) =$

(A): 0.2

(B): 0.5

(C): 0.475

(D): 0.3

### Question!

**Can we compute the value of**  $P(likeslasagna)$ , **given the above informations?**

(A): Yes.

(B): No.

## Questionnaire

### Question!

**Say**  $P(dog) = 0.4$ ,  $\neg dog \leftrightarrow cat$ , **and**  $P(likeslasagna \mid cat) = 0.5$ . **Then**  
 $P(likeslasagna \wedge cat) =$

(A): 0.2

(B): 0.5

(C): 0.475

(D): 0.3

→ We have  $P(cat) = 0.6$  and  $P(likeslasagna \mid cat) = 0.5$ , hence (D) by the product rule.

### Question!

**Can we compute the value of**  $P(likeslasagna)$ , **given the above informations?**

(A): Yes.

(B): No.

# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $\neg dog \leftrightarrow cat$ , **and**  $P(likeslasagna \mid cat) = 0.5$ . **Then**  
 $P(likeslasagna \wedge cat) =$

(A): 0.2

(B): 0.5

(C): 0.475

(D): 0.3

→ We have  $P(cat) = 0.6$  and  $P(likeslasagna \mid cat) = 0.5$ , hence (D) by the product rule.

## Question!

**Can we compute the value of**  $P(likeslasagna)$ , **given the above informations?**

(A): Yes.

(B): No.

→ No. We don't know the probability that *dogs* like lasagna, i.e.,  $P(likeslasagna \mid dog)$ .

## Normalization: Idea

**Problem:** We know  $P(\text{cavity} \wedge \text{toothache})$  but don't know  $P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.12}{P(\text{toothache})}$$

# Normalization: Idea

**Problem:** We know  $P(\text{cavity} \wedge \text{toothache})$  but don't know  $P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.12}{P(\text{toothache})}$$

**Step 1:** Case distinction over the values of  $\text{Cavity}$ :

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.08}{P(\text{toothache})}$$

# Normalization: Idea

**Problem:** We know  $P(\text{cavity} \wedge \text{toothache})$  but don't know  $P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.12}{P(\text{toothache})}$$

**Step 1:** Case distinction over the values of  $\text{Cavity}$ :

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.08}{P(\text{toothache})}$$

**Step 2:** Assuming placeholder  $\alpha := 1/P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \alpha P(\text{cavity} \wedge \text{toothache}) = \alpha 0.12$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \alpha P(\neg \text{cavity} \wedge \text{toothache}) = \alpha 0.08$$

# Normalization: Idea

**Problem:** We know  $P(\text{cavity} \wedge \text{toothache})$  but don't know  $P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.12}{P(\text{toothache})}$$

**Step 1:** Case distinction over the values of  $\text{Cavity}$ :

$$P(\neg \text{cavity} \mid \text{toothache}) = \frac{P(\neg \text{cavity} \wedge \text{toothache})}{P(\text{toothache})} = \frac{0.08}{P(\text{toothache})}$$

**Step 2:** Assuming placeholder  $\alpha := 1/P(\text{toothache})$ :

$$P(\text{cavity} \mid \text{toothache}) = \alpha P(\text{cavity} \wedge \text{toothache}) = \alpha 0.12$$

$$P(\neg \text{cavity} \mid \text{toothache}) = \alpha P(\neg \text{cavity} \wedge \text{toothache}) = \alpha 0.08$$

**Step 3:** Fixing  $\text{toothache}$  to be true, view  $P(\text{cavity} \wedge \text{toothache})$  vs.  $P(\neg \text{cavity} \wedge \text{toothache})$  as the **relative weights of  $P(\text{cavity})$  vs.  $P(\neg \text{cavity})$  within  $\text{toothache}$** . Then normalize their summed-up weight to 1:

$$1 = \alpha(0.12 + 0.08) \Rightarrow \alpha = 1/(0.12 + 0.08) = 1/0.2 = 5$$

→  $\alpha$  is the **normalization constant** scaling the sum of relative weights to 1.

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .



## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X \mid e) = \alpha P(X, e)$ .

**Proof.**

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X | e) = \alpha P(X, e)$ .

**Proof.** For each value  $x$  of  $X$ ,  $P(X = x | e) = P(X = x \wedge e) / P(e)$ . So all we need to prove is that  $\alpha = 1 / P(e)$ .

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X | e) = \alpha P(X, e)$ .

**Proof.** For each value  $x$  of  $X$ ,  $P(X = x | e) = P(X = x \wedge e) / P(e)$ . So all we need to prove is that  $\alpha = 1 / P(e)$ . By definition,  $\alpha = 1 / \sum_x P(X = x \wedge e)$ ,

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X | e) = \alpha P(X, e)$ .

**Proof.** For each value  $x$  of  $X$ ,  $P(X = x | e) = P(X = x \wedge e) / P(e)$ . So all we need to prove is that  $\alpha = 1 / P(e)$ . By definition,  $\alpha = 1 / \sum_x P(X = x \wedge e)$ , so we need to prove  $P(e) = \sum_x P(X = x \wedge e)$  which holds by Marginalization.

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X | e) = \alpha P(X, e)$ .

**Proof.** For each value  $x$  of  $X$ ,  $P(X = x | e) = P(X = x \wedge e) / P(e)$ . So all we need to prove is that  $\alpha = 1 / P(e)$ . By definition,  $\alpha = 1 / \sum_x P(X = x \wedge e)$ , so we need to prove  $P(e) = \sum_x P(X = x \wedge e)$  which holds by Marginalization.

**Example:**  $\alpha \langle P(\text{cavity} \wedge \text{toothache}), P(\neg \text{cavity} \wedge \text{toothache}) \rangle = \alpha \langle 0.12, 0.08 \rangle$ , so  $P(\text{cavity} | \text{toothache}) = 0.6$ , and  $P(\neg \text{cavity} | \text{toothache}) = 0.4$ .

## Normalization: Formal

**Definition.** Given a vector  $\langle w_1, \dots, w_k \rangle$  of numbers in  $[0, 1]$  where  $\sum_{i=1}^k w_i \leq 1$ , the *normalization constant*  $\alpha$  is  $\alpha \langle w_1, \dots, w_k \rangle := 1 / \sum_{i=1}^k w_i$ .

**Example:**  $\alpha \langle 0.12, 0.08 \rangle = 5 \langle 0.12, 0.08 \rangle = \langle 0.6, 0.4 \rangle$ .

**Proposition (Normalization).** Given a random variable  $X$  and an event  $e$ , we have  $P(X | e) = \alpha P(X, e)$ .

**Proof.** For each value  $x$  of  $X$ ,  $P(X = x | e) = P(X = x \wedge e) / P(e)$ . So all we need to prove is that  $\alpha = 1 / P(e)$ . By definition,  $\alpha = 1 / \sum_x P(X = x \wedge e)$ , so we need to prove  $P(e) = \sum_x P(X = x \wedge e)$  which holds by Marginalization.

**Example:**  $\alpha \langle P(\text{cavity} \wedge \text{toothache}), P(\neg \text{cavity} \wedge \text{toothache}) \rangle = \alpha \langle 0.12, 0.08 \rangle$ , so  $P(\text{cavity} | \text{toothache}) = 0.6$ , and  $P(\neg \text{cavity} | \text{toothache}) = 0.4$ .

**Normalization+Marginalization:** Given “query variable”  $X$ , “observed event”  $e$ , and “hidden variables” set  $\mathbf{Y}$ :  $P(X | e) = \alpha P(X, e) = \alpha \sum_{\mathbf{y} \in \mathbf{Y}} P(X, e, \mathbf{y})$ .

→ Second of the four basic techniques in Bayesian networks.

## Questionnaire

### Question!

**Say we know  $P(\text{likeschappi} \wedge \text{dog}) = 0.32$  and  $P(\neg \text{likeschappi} \wedge \text{dog}) = 0.08$ . Can we compute  $P(\text{likeschappi} \mid \text{dog})$ ?**

(A): Yes.

(B): No.

# Questionnaire

## Question!

**Say we know  $P(\text{likeschappi} \wedge \text{dog}) = 0.32$  and  $P(\neg \text{likeschappi} \wedge \text{dog}) = 0.08$ . Can we compute  $P(\text{likeschappi} \mid \text{dog})$ ?**

(A): Yes.

(B): No.

→ Yes, because we can compute  $P(\text{dog}) = P(\text{likeschappi} \wedge \text{dog}) + P(\neg \text{likeschappi} \wedge \text{dog})$ , and thus  $P(\text{likeschappi} \mid \text{dog}) = \frac{P(\text{likeschappi} \wedge \text{dog})}{P(\text{dog})}$ .



# Questionnaire

## Question!

**Say we know  $P(\text{likeschappi} \wedge \text{dog}) = 0.32$  and  $P(\neg \text{likeschappi} \wedge \text{dog}) = 0.08$ . Can we compute  $P(\text{likeschappi} \mid \text{dog})$ ?**

(A): Yes.

(B): No.

→ Yes, because we can compute  $P(\text{dog}) = P(\text{likeschappi} \wedge \text{dog}) + P(\neg \text{likeschappi} \wedge \text{dog})$ , and thus  $P(\text{likeschappi} \mid \text{dog}) = \frac{P(\text{likeschappi} \wedge \text{dog})}{P(\text{dog})}$ .

→ In other words, we can use Normalization:  $\mathbf{P}(X \mid \mathbf{e}) = \alpha \mathbf{P}(X, \mathbf{e})$ .

Inserting *LikesChappi* for  $X$  and *dog* for  $\mathbf{e}$ , we get  $\mathbf{P}(\text{LikesChappi} \mid \text{dog}) = \alpha \mathbf{P}(\text{LikesChappi}, \text{dog}) = \alpha \langle P(\text{likeschappi} \wedge \text{dog}), P(\neg \text{likeschappi} \wedge \text{dog}) \rangle = \alpha \langle 0.32, 0.08 \rangle$ .

# Questionnaire

## Question!

**Say we know  $P(\text{likeschappi} \wedge \text{dog}) = 0.32$  and  $P(\neg \text{likeschappi} \wedge \text{dog}) = 0.08$ . Can we compute  $P(\text{likeschappi} \mid \text{dog})$ ?**

(A): Yes.

(B): No.

→ Yes, because we can compute  $P(\text{dog}) = P(\text{likeschappi} \wedge \text{dog}) + P(\neg \text{likeschappi} \wedge \text{dog})$ , and thus  $P(\text{likeschappi} \mid \text{dog}) = \frac{P(\text{likeschappi} \wedge \text{dog})}{P(\text{dog})}$ .

→ In other words, we can use Normalization:  $\mathbf{P}(X \mid \mathbf{e}) = \alpha \mathbf{P}(X, \mathbf{e})$ .

Inserting *LikesChappi* for  $X$  and *dog* for  $\mathbf{e}$ , we get  $\mathbf{P}(\text{LikesChappi} \mid \text{dog}) = \alpha \mathbf{P}(\text{LikesChappi}, \text{dog}) = \alpha \langle P(\text{likeschappi} \wedge \text{dog}), P(\neg \text{likeschappi} \wedge \text{dog}) \rangle = \alpha \langle 0.32, 0.08 \rangle$ .

→ So what is  $P(\text{likeschappi} \mid \text{dog})$ ?

# Questionnaire

## Question!

**Say we know  $P(\text{likeschappi} \wedge \text{dog}) = 0.32$  and  $P(\neg \text{likeschappi} \wedge \text{dog}) = 0.08$ . Can we compute  $P(\text{likeschappi} \mid \text{dog})$ ?**

(A): Yes.

(B): No.

→ Yes, because we can compute  $P(\text{dog}) = P(\text{likeschappi} \wedge \text{dog}) + P(\neg \text{likeschappi} \wedge \text{dog})$ , and thus  $P(\text{likeschappi} \mid \text{dog}) = \frac{P(\text{likeschappi} \wedge \text{dog})}{P(\text{dog})}$ .

→ In other words, we can use Normalization:  $\mathbf{P}(X \mid \mathbf{e}) = \alpha \mathbf{P}(X, \mathbf{e})$ .

Inserting *LikesChappi* for  $X$  and *dog* for  $\mathbf{e}$ , we get  $\mathbf{P}(\text{LikesChappi} \mid \text{dog}) = \alpha \mathbf{P}(\text{LikesChappi}, \text{dog}) = \alpha \langle P(\text{likeschappi} \wedge \text{dog}), P(\neg \text{likeschappi} \wedge \text{dog}) \rangle = \alpha \langle 0.32, 0.08 \rangle$ .

→ So what is  $P(\text{likeschappi} \mid \text{dog})$ ? 0.8, because  $\alpha = 1/P(\text{dog}) = 1/(0.32 + 0.08) = 2.5$ .

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule**
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

# Bayes' Rule

**Proposition (Bayes' Rule).** *Given propositions  $A$  and  $B$  where  $P(a) \neq 0$  and  $P(b) \neq 0$ , we have:*

$$P(a | b) = \frac{P(b | a)P(a)}{P(b)}$$

# Bayes' Rule

**Proposition (Bayes' Rule).** *Given propositions  $A$  and  $B$  where  $P(a) \neq 0$  and  $P(b) \neq 0$ , we have:*

$$P(a | b) = \frac{P(b | a)P(a)}{P(b)}$$

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$

# Bayes' Rule

**Proposition (Bayes' Rule).** *Given propositions  $A$  and  $B$  where  $P(a) \neq 0$  and  $P(b) \neq 0$ , we have:*

$$P(a | b) = \frac{P(b | a)P(a)}{P(b)}$$

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$  which by product rule  $P(a \wedge b) = P(b | a)P(a)$  is equal to the claim.

# Bayes' Rule

**Proposition (Bayes' Rule).** *Given propositions  $A$  and  $B$  where  $P(a) \neq 0$  and  $P(b) \neq 0$ , we have:*

$$P(a | b) = \frac{P(b | a)P(a)}{P(b)}$$

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$  which by product rule  $P(a \wedge b) = P(b | a)P(a)$  is equal to the claim.

**Notation:** (System of equations)

$$\mathbf{P}(X | Y) = \frac{\mathbf{P}(Y | X)\mathbf{P}(X)}{\mathbf{P}(Y)}$$



## Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ :

## Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ : By Bayes' rule,

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{toothache})} = \frac{0.6 * 0.2}{0.2} = 0.6.$$

## Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ : By Bayes' rule,

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{toothache})} = \frac{0.6 \cdot 0.2}{0.2} = 0.6.$$

**Ok, but:** Why don't we simply assess  $P(\text{cavity} \mid \text{toothache})$  directly?

- $P(\text{toothache} \mid \text{cavity})$  is **causal**,  $P(\text{cavity} \mid \text{toothache})$  is **diagnostic**.

# Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ : By Bayes' rule,

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{toothache})} = \frac{0.6 * 0.2}{0.2} = 0.6.$$

**Ok, but:** Why don't we simply assess  $P(\text{cavity} \mid \text{toothache})$  directly?

- $P(\text{toothache} \mid \text{cavity})$  is **causal**,  $P(\text{cavity} \mid \text{toothache})$  is **diagnostic**.
- **Causal dependencies are robust over frequency of the causes.**  
→ Example: If there is a cavity epidemic then  $P(\text{cavity} \mid \text{toothache})$  increases, but  $P(\text{toothache} \mid \text{cavity})$  remains the same.

# Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ : By Bayes' rule,

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{toothache})} = \frac{0.6 \cdot 0.2}{0.2} = 0.6.$$

**Ok, but:** Why don't we simply assess  $P(\text{cavity} \mid \text{toothache})$  directly?

- $P(\text{toothache} \mid \text{cavity})$  is **causal**,  $P(\text{cavity} \mid \text{toothache})$  is **diagnostic**.
- **Causal dependencies are robust over frequency of the causes.**  
→ Example: If there is a cavity epidemic then  $P(\text{cavity} \mid \text{toothache})$  increases, but  $P(\text{toothache} \mid \text{cavity})$  remains the same.
- Also, causal dependencies are often easier to assess.

# Applying Bayes' Rule

**Example:** Say we know that  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{cavity}) = 0.2$ , and  $P(\text{toothache}) = 0.2$ .

→ We can compute  $P(\text{cavity} \mid \text{toothache})$ : By Bayes' rule,

$$P(\text{cavity} \mid \text{toothache}) = \frac{P(\text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{toothache})} = \frac{0.6 \cdot 0.2}{0.2} = 0.6.$$

**Ok, but:** Why don't we simply assess  $P(\text{cavity} \mid \text{toothache})$  directly?

- $P(\text{toothache} \mid \text{cavity})$  is **causal**,  $P(\text{cavity} \mid \text{toothache})$  is **diagnostic**.
- **Causal dependencies are robust over frequency of the causes.**  
→ Example: If there is a cavity epidemic then  $P(\text{cavity} \mid \text{toothache})$  increases, but  $P(\text{toothache} \mid \text{cavity})$  remains the same.
- Also, causal dependencies are often easier to assess.

→ Bayes' rule allows to perform diagnosis (observing a symptom, what is the cause?) based on prior probabilities and causal dependencies.

# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $P(likeschappi \mid dog) = 0.8$ , **and**  $P(likeschappi) = 0.5$ . **What is**  $P(dog \mid likeschappi)$ ?

(A): 0.8

(B): 0.64

(C): 0.9

(D): 0.32

# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $P(likeschappi \mid dog) = 0.8$ , and  $P(likeschappi) = 0.5$ . **What is**  $P(dog \mid likeschappi)$ ?

(A): 0.8

(B): 0.64

(C): 0.9

(D): 0.32

→ By Bayes' rule,  $P(dog \mid likeschappi) = \frac{P(likeschappi \mid dog)P(dog)}{P(likeschappi)} = \frac{0.8 \cdot 0.4}{0.5} = 0.64$  so (B).



# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $P(likeschappi \mid dog) = 0.8$ , and  $P(likeschappi) = 0.5$ . **What is**  $P(dog \mid likeschappi)$ ?

(A): 0.8

(B): 0.64

(C): 0.9

(D): 0.32

→ By Bayes' rule,  $P(dog \mid likeschappi) = \frac{P(likeschappi \mid dog)P(dog)}{P(likeschappi)} = \frac{0.8 \cdot 0.4}{0.5} = 0.64$  so (B).

→ Is  $P(likeschappi \mid dog)$  causal or diagnostic?

# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $P(likeschappi \mid dog) = 0.8$ , and  $P(likeschappi) = 0.5$ . **What is**  $P(dog \mid likeschappi)$ ?

(A): 0.8

(B): 0.64

(C): 0.9

(D): 0.32

→ By Bayes' rule,  $P(dog \mid likeschappi) = \frac{P(likeschappi \mid dog)P(dog)}{P(likeschappi)} = \frac{0.8 \cdot 0.4}{0.5} = 0.64$  so (B).

→ Is  $P(likeschappi \mid dog)$  causal or diagnostic? Causal; liking or not liking dog food may be caused by being or not being a dog.

→ Is  $P(dog \mid likeschappi)$  causal or diagnostic?

# Questionnaire

## Question!

**Say**  $P(dog) = 0.4$ ,  $P(likeschappi \mid dog) = 0.8$ , and  $P(likeschappi) = 0.5$ . **What is**  $P(dog \mid likeschappi)$ ?

(A): 0.8

(B): 0.64

(C): 0.9

(D): 0.32

→ By Bayes' rule,  $P(dog \mid likeschappi) = \frac{P(likeschappi \mid dog)P(dog)}{P(likeschappi)} = \frac{0.8 \cdot 0.4}{0.5} = 0.64$  so (B).

→ Is  $P(likeschappi \mid dog)$  causal or diagnostic? Causal; liking or not liking dog food may be caused by being or not being a dog.

→ Is  $P(dog \mid likeschappi)$  causal or diagnostic? Diagnostic; liking Chappi does not cause anybody to be a dog.

## Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

## Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)}$$

## Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

# Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

Consider the variables

- Temp :  $\text{sp}(\text{Temp}) = \{l, m, h\}$
- Sensor :  $\text{sp}(\text{Sensor}) = \{l, m, h\}$

$$P(\text{Temp}) = (0.1, 0.6, 0.3)$$

$$P(\text{Sensor}|\text{Temp}) =$$

		Temp		
		l	m	h
Sensor	l	0.8	0.1	0.05
	m	0.15	0.8	0.1
	h	0.05	0.1	0.85

## Question!

Assume we observe  $S = \text{low}$ , what is our updated belief in the temperature?

(A):  $P(T = \text{low} \mid S = \text{low}) = 0.1$

(B):  $P(T = \text{low} \mid S = \text{low}) = 0.52$

(C):  $P(T = \text{low} \mid S = \text{low}) = 0.8$

(D):  $P(T = \text{low} \mid S = \text{low}) = 1$

# Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

Consider the variables

- Temp :  $\text{sp}(\text{Temp}) = \{l, m, h\}$
- Sensor :  $\text{sp}(\text{Sensor}) = \{l, m, h\}$

$$P(\text{Temp}) = (0.1, 0.6, 0.3)$$

$$P(\text{Sensor}|\text{Temp}) =$$

		Temp		
		l	m	h
Sensor	l	0.8	0.1	0.05
	m	<del>0.15</del>	<del>0.8</del>	<del>0.1</del>
	h	<del>0.05</del>	<del>0.1</del>	<del>0.85</del>

## Question!

Assume we observe  $S = \text{low}$ , what is our updated belief in the temperature?

- (A):  $P(T = \text{low} \mid S = \text{low}) = 0.1$  (B):  $P(T = \text{low} \mid S = \text{low}) = 0.52$   
 (C):  $P(T = \text{low} \mid S = \text{low}) = 0.8$  (D):  $P(T = \text{low} \mid S = \text{low}) = 1$

$$P(T|\text{low}) = \frac{P(\text{low}|T)P(T)}{\sum_T P(\text{low}|T)P(T)} = \frac{S = \text{low}}{\text{Temp}=l \quad \text{Temp}=m \quad \text{Temp}=h}$$



# Bayes' rule for variables

Bayes' rule can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

Consider the variables

- Temp :  $\text{sp}(\text{Temp}) = \{l, m, h\}$
- Sensor :  $\text{sp}(\text{Sensor}) = \{l, m, h\}$

$$P(\text{Temp}) = (0.1, 0.6, 0.3)$$

$$P(\text{Sensor}|\text{Temp}) =$$

		Temp		
		l	m	h
Sensor	l	0.8	0.1	0.05
	m	0.15	0.8	0.1
	h	0.05	0.1	0.85

## Question!

Assume we observe  $S = \text{low}$ , what is our updated belief in the temperature?

(A):  $P(T = \text{low} \mid S = \text{low}) = 0.1$

(B):  $P(T = \text{low} \mid S = \text{low}) = 0.52$

(C):  $P(T = \text{low} \mid S = \text{low}) = 0.8$

(D):  $P(T = \text{low} \mid S = \text{low}) = 1$

$$P(T|\text{low}) = \frac{P(\text{low}|T)P(T)}{\sum_T P(\text{low}|T)P(T)} = \frac{S = \text{low}}{\text{Temp}=l \quad \text{Temp}=m \quad \text{Temp}=h}$$

$$P(\text{low}|T)P(T) = \frac{S = \text{low}}{\begin{array}{|c|c|c|} \hline l & m & h \\ \hline 0.08 & 0.06 & 0.015 \\ \hline \end{array}} \rightarrow \frac{P(S = \text{low})}{0.155}$$

# Bayes' rule for variables

**Bayes' rule** can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

Consider the variables

- Temp :  $\text{sp}(\text{Temp}) = \{l, m, h\}$
- Sensor :  $\text{sp}(\text{Sensor}) = \{l, m, h\}$

$$P(\text{Temp}) = (0.1, 0.6, 0.3)$$

$P(\text{Sensor}|\text{Temp}) =$

		Temp		
		l	m	h
Sensor	l	0.8	0.1	0.05
	m	0.15	0.8	0.1
	h	0.05	0.1	0.85

## Question!

Assume we observe  $S = \text{low}$ , what is our updated belief in the temperature?

- (A):**  $P(T = \text{low} \mid S = \text{low}) = 0.1$
- (B):**  $P(T = \text{low} \mid S = \text{low}) = 0.52$
- (C):**  $P(T = \text{low} \mid S = \text{low}) = 0.8$
- (D):**  $P(T = \text{low} \mid S = \text{low}) = 1$

$$P(T|\text{low}) = \frac{P(\text{low}|T)P(T)}{\sum_T P(\text{low}|T)P(T)} = \frac{S = \text{low}}{\begin{array}{|c|c|c|} \hline \text{Temp}=l & \text{Temp}=m & \text{Temp}=h \\ \hline \text{0.08} / \text{0.155} & \text{0.06} / \text{0.155} & \text{0.015} / \text{0.155} \\ \hline \end{array}}$$

$$P(\text{low}|T)P(T) = \frac{S = \text{low}}{\begin{array}{|c|c|c|} \hline l & m & h \\ \hline \text{0.08} & \text{0.06} & \text{0.015} \\ \hline \end{array}} \rightarrow \frac{P(S = \text{low})}{\text{0.155}}$$

# Bayes' rule for variables

**Bayes' rule** can also be written for variables:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)} = \frac{P(B|A)P(A)}{\sum_A P(A, B)} = \frac{P(B|A)P(A)}{\sum_A P(B|A)P(A)}$$

Consider the variables

- Temp :  $\text{sp}(\text{Temp}) = \{l, m, h\}$
- Sensor :  $\text{sp}(\text{Sensor}) = \{l, m, h\}$

$$P(\text{Temp}) = (0.1, 0.6, 0.3)$$

$$P(\text{Sensor}|\text{Temp}) =$$

		Temp		
		l	m	h
Sensor	l	0.8	0.1	0.05
	m	<del>0.15</del>	<del>0.8</del>	<del>0.1</del>
	h	<del>0.05</del>	<del>0.1</del>	<del>0.85</del>

## Question!

**Assume we observe  $S = \text{low}$ , what is our updated belief in the temperature?**

- (A):  $P(T = \text{low} \mid S = \text{low}) = 0.1$ 
 (B):  $P(T = \text{low} \mid S = \text{low}) = 0.52$
- (C):  $P(T = \text{low} \mid S = \text{low}) = 0.8$ 
 (D):  $P(T = \text{low} \mid S = \text{low}) = 1$

$$P(T|\text{low}) = \frac{P(\text{low}|T)P(T)}{\sum_T P(\text{low}|T)P(T)} = \frac{S = \text{low}}{\begin{array}{|c|c|c|} \hline \text{Temp}=l & \text{Temp}=m & \text{Temp}=h \\ \hline 0.52 & 0.39 & 0.09 \\ \hline \end{array}}$$

$$P(\text{low}|T)P(T) = \frac{S = \text{low}}{\begin{array}{|c|c|c|} \hline l & m & h \\ \hline 0.08 & 0.06 & 0.015 \\ \hline \end{array}} \rightarrow \frac{P(S = \text{low})}{0.155}$$

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence**
- 9 Conditional Independence
- 10 Conclusion

# Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

# Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ?

## Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

## Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \vee \text{toothache})$ ?



## Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \vee \text{toothache})$ ? Sum across atomic events:

$$P(\text{cavity} \wedge \text{toothache}) + P(\neg \text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

## Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \vee \text{toothache})$ ? Sum across atomic events:

$$P(\text{cavity} \wedge \text{toothache}) + P(\neg \text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \mid \text{toothache})$ ?

# Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \vee \text{toothache})$ ? Sum across atomic events:

$$P(\text{cavity} \wedge \text{toothache}) + P(\neg \text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \mid \text{toothache})$ ?  $\frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$

## Working with the Full Joint Probability Distribution

**Example:**

	<i>toothache</i>	$\neg$ <i>toothache</i>
<i>cavity</i>	0.12	0.08
$\neg$ <i>cavity</i>	0.08	0.72

→ How to compute  $P(\text{cavity})$ ? Sum across the row:

$$P(\text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \vee \text{toothache})$ ? Sum across atomic events:

$$P(\text{cavity} \wedge \text{toothache}) + P(\neg \text{cavity} \wedge \text{toothache}) + P(\text{cavity} \wedge \neg \text{toothache})$$

→ How to compute  $P(\text{cavity} \mid \text{toothache})$ ?  $\frac{P(\text{cavity} \wedge \text{toothache})}{P(\text{toothache})}$

→ All relevant probabilities can be computed using the full joint probability distribution, by expressing propositions as disjunctions of atomic events.

## Working with the Full Joint Probability Distribution??

→ So, is it a good idea to use the full joint probability distribution?

## Working with the Full Joint Probability Distribution??

→ So, is it a good idea to use the full joint probability distribution? No:

- Given  $n$  random variables with  $k$  values each, the joint probability distribution contains  $k^n$  probabilities.
- Computational cost of dealing with this size.
- *Practically impossible to assess all these probabilities.*

## Working with the Full Joint Probability Distribution??

→ So, is it a good idea to use the full joint probability distribution? No:

- Given  $n$  random variables with  $k$  values each, the joint probability distribution contains  $k^n$  probabilities.
- Computational cost of dealing with this size.
- *Practically impossible to assess all these probabilities.*

→ So, is there a compact way to represent the full joint probability distribution? Is there an efficient method to work with that representation?

## Working with the Full Joint Probability Distribution??

→ So, is it a good idea to use the full joint probability distribution? No:

- Given  $n$  random variables with  $k$  values each, the joint probability distribution contains  $k^n$  probabilities.
- Computational cost of dealing with this size.
- *Practically impossible to assess all these probabilities.*

→ So, is there a compact way to represent the full joint probability distribution? Is there an efficient method to work with that representation?

→ Not in general, but it works in many cases. We can work directly with conditional probabilities, and exploit (conditional) independence.

→ Bayesian networks.



## Working with the Full Joint Probability Distribution??

→ So, is it a good idea to use the full joint probability distribution? No:

- Given  $n$  random variables with  $k$  values each, the joint probability distribution contains  $k^n$  probabilities.
- Computational cost of dealing with this size.
- *Practically impossible to assess all these probabilities.*

→ So, is there a compact way to represent the full joint probability distribution? Is there an efficient method to work with that representation?

→ Not in general, but it works in many cases. We can work directly with conditional probabilities, and exploit **(conditional) independence**.

→ **Bayesian networks**. (First, we do the simple case.)

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a \mid b) = P(a)$ .

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a \mid b) = P(a)$ .

**Proof.** By definition,  $P(a \mid b) = \frac{P(a \wedge b)}{P(b)}$ ,

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a \mid b) = P(a)$ .

**Proof.** By definition,  $P(a \mid b) = \frac{P(a \wedge b)}{P(b)}$ , which by independence is equal to  $\frac{P(a)P(b)}{P(b)} = P(a)$ .  
 (→ Similarly, if  $P(a) \neq 0$ , we have  $P(b \mid a) = P(b)$ .)

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a | b) = P(a)$ .

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ , which by independence is equal to  $\frac{P(a)P(b)}{P(b)} = P(a)$ .  
 (→ Similarly, if  $P(a) \neq 0$ , we have  $P(b | a) = P(b)$ .)

**Examples:**

- $P(\text{Dice1} = 6 \wedge \text{Dice2} = 6) = 1/36$ .
- $P(W = \text{sunny} | \text{headache}) = P(W = \text{sunny})$  unless you're weather-sensitive (cf. slide 26).

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a | b) = P(a)$ .

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ , which by independence is equal to  $\frac{P(a)P(b)}{P(b)} = P(a)$ .  
 (→ Similarly, if  $P(a) \neq 0$ , we have  $P(b | a) = P(b)$ .)

## Examples:

- $P(\text{Dice1} = 6 \wedge \text{Dice2} = 6) = 1/36$ .
- $P(W = \text{sunny} | \text{headache}) = P(W = \text{sunny})$  unless you're weather-sensitive (cf. slide 26).
- **But toothache and cavity are NOT independent.** The fraction of "cavity" is higher within "toothache" than within " $\neg$ toothache".  $P(\text{toothache}) = 0.2$  and  $P(\text{cavity}) = 0.2$ , but  $P(\text{toothache} \wedge \text{cavity}) = 0.12 > 0.04$ .

# Independence

**Definition.** Events  $a$  and  $b$  are *independent* if  $P(a \wedge b) = P(a)P(b)$ .

**Proposition.** Given independent events  $a$  and  $b$  where  $P(b) \neq 0$ , we have  $P(a | b) = P(a)$ .

**Proof.** By definition,  $P(a | b) = \frac{P(a \wedge b)}{P(b)}$ , which by independence is equal to  $\frac{P(a)P(b)}{P(b)} = P(a)$ .  
 (→ Similarly, if  $P(a) \neq 0$ , we have  $P(b | a) = P(b)$ .)

## Examples:

- $P(\text{Dice1} = 6 \wedge \text{Dice2} = 6) = 1/36$ .
- $P(W = \text{sunny} | \text{headache}) = P(W = \text{sunny})$  unless you're weather-sensitive (cf. slide 26).
- **But toothache and cavity are NOT independent.** The fraction of "cavity" is higher within "toothache" than within " $\neg$ toothache".  $P(\text{toothache}) = 0.2$  and  $P(\text{cavity}) = 0.2$ , but  $P(\text{toothache} \wedge \text{cavity}) = 0.12 > 0.04$ .

**Definition.** Random variables  $X$  and  $Y$  are independent if  $\mathbf{P}(X, Y) = \mathbf{P}(X)\mathbf{P}(Y)$ . (System of equations!)



## Example: Football statistics

Results for Bayern Munich and SC Freiburg in seasons 2001/02 and 2003/04. (Not counting the matches Munich vs. Freiburg):

$$D_{\text{Munich}} = D_{\text{Freiburg}} = \{\text{Win}, \text{Draw}, \text{Loss}\}$$

2001/02

Munich: LWDWWWWWWLDDLDDLWLDWWWDWDDWWWW

Freiburg: WLLDDWLDWDWLLLLDDLWDDLLDLLLWLW

2003/04

Munich: WDWWLDWWDWLWDDWDWLWWDDWWLWLL

Freiburg: LDDWDWLWLLLWVLWLWLLDWLDDWDLWLWLD

Summary:

Munich	Freiburg			
	W	D	L	
W	12	9	15	36
D	3	4	9	16
L	6	4	2	12
	21	17	26	

# Independence of Outcomes

The joint distribution of *Munich* and *Freiburg*:

$P(\text{Munich}, \text{Freiburg})$ :

<i>Munich</i>	<i>Freiburg</i>			$P(\text{Munich})$
	W	D	L	
W	.1875	.1406	.2344	.5625
D	.0468	.0625	.1406	.25
L	.0937	.0625	.0312	.1875
$P(\text{Freiburg})$	.3281	.2656	.4062	

## Independence of Outcomes

The joint distribution of *Munich* and *Freiburg*:

$P(\text{Munich}, \text{Freiburg})$ :

<i>Munich</i>	<i>Freiburg</i>			$P(\text{Munich})$
	W	D	L	
W	.1875 .571	.1406 .529	.2344 .577	.5625
D	.0468 .143	.0625 .235	.1406 .346	
L	.0937 .285	.0625 .235	.0312 .077	.1875
$P(\text{Freiburg})$	.3281	.2656	.4062	

Conditional distribution:  $P(\text{Munich} \mid \text{Freiburg})$

## Independence of Outcomes

The joint distribution of *Munich* and *Freiburg*:

$P(\text{Munich}, \text{Freiburg})$ :

<i>Munich</i>	<i>Freiburg</i>			$P(\text{Munich})$
	W	D	L	
W	.1875 .571	.1406 .529	.2344 .577	.5625
D	.0468 .143	.0625 .235	.1406 .346	
L	.0937 .285	.0625 .235	.0312 .077	.1875
$P(\text{Freiburg})$	.3281	.2656	.4062	

Conditional distribution:  $P(\text{Munich} \mid \text{Freiburg})$

We have (almost):

$$P(\text{Munich} \mid \text{Freiburg}) = P(\text{Munich})$$

The variables *Munich* and *Freiburg* are **independent**.

# Independent Variables

## Definition of Independence

The variables  $A_1, \dots, A_k$  and  $B_1, \dots, B_m$  are **independent** if

$$P(A_1, \dots, A_k \mid B_1, \dots, B_m) = P(A_1, \dots, A_k)$$

This is equivalent to:

$$P(B_1, \dots, B_m \mid A_1, \dots, A_k) = P(B_1, \dots, B_m)$$

and also to:

$$P(A_1, \dots, A_k, B_1, \dots, B_m) = P(A_1, \dots, A_k) \cdot P(B_1, \dots, B_m)$$

## Compact Specifications by Independence

Independence properties can greatly simplify the specification of a joint distribution:

$M =$	$F =$			$P(M)$
	W	D	L	
W	<i>M and F are independent</i>			.5625
D				.25
L				.1875
$P(F)$	.3281	.2656	.4062	

The probability for each possible world then is defined, e.g.

$$P(M = D, F = L) = 0.25 \cdot 0.4062 = 0.10155$$

→ Independence can be exploited to represent the full joint probability distribution more compactly.

→ Usually, random variables are independent only under particular conditions: **conditional independence**, see next section.

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence**
- 10 Conclusion

## Questionnaire

*Hair length* :  $D_{\text{Hair length}} = \{\text{long, short}\}$

*Height* :  $D_{\text{Height}} = \{\text{tall, medium}\}$

### Question!

**Are Hair length and Height independent?**

(A): Yes

(B): No



## Questionnaire

Hair length :  $D_{\text{Hair length}} = \{\text{long, short}\}$

Height :  $D_{\text{Height}} = \{\text{tall, medium}\}$

Sex :  $D_{\text{Sex}} = \{\text{male, female}\}$

### Question!

**Are Hair length and Height independent?**

(A): Yes

(B): No

Joint Distribution:

	Sex			
	male		female	
Height	Hair length		Hair length	
	long	short	long	short
tall	0.06	0.24	0.07	0.03
medium	0.04	0.16	0.28	0.12

## Questionnaire

Hair length :  $D_{\text{Hair length}} = \{\text{long}, \text{short}\}$

Height :  $D_{\text{Height}} = \{\text{tall}, \text{medium}\}$

Sex :  $D_{\text{Sex}} = \{\text{male}, \text{female}\}$

### Question!

**Are Hair length and Height independent?**

(A): Yes

(B): No

Joint Distribution:

	Sex			
	male		female	
Height	Hair length		Hair length	
	long	short	long	short
tall	0.06	0.24	0.07	0.03
medium	0.04	0.16	0.28	0.12

$P(\text{Hair length}, \text{Height})$   $P(\text{Height})$ ,  
 $P(\text{Height} \mid \text{Hair length})$ :

Height	Hair length		
	long	short	
tall	0.13	0.27	0.4
	0.289	0.49	
medium	0.32	0.28	0.6
	0.711	0.51	

↪ No, *Hair length* and *Height* are not independent. Tall people have shorter hair!

Does that sound right to you?

## Conditional Independence

**Definition.** Given sets of random variables  $\mathbf{Z}_1$ ,  $\mathbf{Z}_2$ ,  $\mathbf{Z}$ , we say that  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are *conditionally independent given  $\mathbf{Z}$*  if:

$$\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z})\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z})$$

We alternatively say that  $\mathbf{Z}_1$  is *conditionally independent of  $\mathbf{Z}_2$  given  $\mathbf{Z}$* .

## Conditional Independence

**Definition.** Given sets of random variables  $\mathbf{Z}_1$ ,  $\mathbf{Z}_2$ ,  $\mathbf{Z}$ , we say that  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are *conditionally independent given  $\mathbf{Z}$*  if:

$$P(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) = P(\mathbf{Z}_1 \mid \mathbf{Z})P(\mathbf{Z}_2 \mid \mathbf{Z})$$

We alternatively say that  $\mathbf{Z}_1$  is *conditionally independent of  $\mathbf{Z}_2$  given  $\mathbf{Z}$* .

$P(\text{Hair length, Height} \mid \text{Sex} = \text{female})$ ,  $P(\text{Height} \mid \text{Sex} = \text{female})$ ,  
 $P(\text{Height} \mid \text{Hair length, Sex} = \text{female})$ :

Height	Hair length		
	long	short	
tall	0.14	0.06	0.2
	0.2	0.2	
medium	0.56	0.24	0.8
	0.8	0.8	

$\leadsto$  Hair length and Height are independent given Sex=female.

Also: Hair length and Height are independent given Sex=male.

$\leadsto$  Hair length and Height are independent given Sex.

# Conditional Independence

**Definition.** Given sets of random variables  $\mathbf{Z}_1$ ,  $\mathbf{Z}_2$ ,  $\mathbf{Z}$ , we say that  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are *conditionally independent given  $\mathbf{Z}$*  if:

$$P(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) = P(\mathbf{Z}_1 \mid \mathbf{Z})P(\mathbf{Z}_2 \mid \mathbf{Z})$$

We alternatively say that  $\mathbf{Z}_1$  is *conditionally independent of  $\mathbf{Z}_2$  given  $\mathbf{Z}$* .

$P(\text{Hair length, Height} \mid \text{Sex} = \text{female})$ ,  $P(\text{Height} \mid \text{Sex} = \text{female})$ ,  
 $P(\text{Height} \mid \text{Hair length, Sex} = \text{female})$ :

Height	Hair length		
	long	short	
tall	0.14	0.06	0.2
	0.2	0.2	
medium	0.56	0.24	0.8
	0.8	0.8	

↪ Hair length and Height are independent given Sex=female.

Also: Hair length and Height are independent given Sex=male.

↪ Hair length and Height are independent given Sex.

**Note:** The definition is symmetric regarding the roles of  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$ : *hairlength* is conditionally independent of *height*, and vice versa.

## Bayes' Rule with Multiple Evidence

**Example:** Say we know from medicinal studies that  $P(\text{cavity}) = 0.2$ ,  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{toothache} \mid \neg \text{cavity}) = 0.1$ ,  $P(\text{catch} \mid \text{cavity}) = 0.9$ , and  $P(\text{catch} \mid \neg \text{cavity}) = 0.2$ . Now, in case we did observe the symptoms toothache and catch (the dentist's probe catches in the aching tooth), what would be the likelihood of having a cavity? What is  $P(\text{cavity} \mid \text{catch}, \text{toothache})$ ?

# Bayes' Rule with Multiple Evidence

**Example:** Say we know from medicinal studies that  $P(\text{cavity}) = 0.2$ ,  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{toothache} \mid \neg \text{cavity}) = 0.1$ ,  $P(\text{catch} \mid \text{cavity}) = 0.9$ , and  $P(\text{catch} \mid \neg \text{cavity}) = 0.2$ . Now, in case we did observe the symptoms toothache and catch (the dentist's probe catches in the aching tooth), what would be the likelihood of having a cavity? What is  $P(\text{cavity} \mid \text{catch}, \text{toothache})$ ?

By Bayes' rule we get:

$$P(\text{cavity} \mid \text{catch}, \text{toothache}) = \frac{P(\text{catch}, \text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{catch}, \text{toothache})}$$

## Bayes' Rule with Multiple Evidence

**Example:** Say we know from medicinal studies that  $P(\text{cavity}) = 0.2$ ,  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{toothache} \mid \neg \text{cavity}) = 0.1$ ,  $P(\text{catch} \mid \text{cavity}) = 0.9$ , and  $P(\text{catch} \mid \neg \text{cavity}) = 0.2$ . Now, in case we did observe the symptoms toothache and catch (the dentist's probe catches in the aching tooth), what would be the likelihood of having a cavity? What is  $P(\text{cavity} \mid \text{catch}, \text{toothache})$ ?

By Bayes' rule we get:

$$P(\text{cavity} \mid \text{catch}, \text{toothache}) = \frac{P(\text{catch}, \text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{catch}, \text{toothache})}$$

Question!

So, is everything fine? Do we just need some more medicinal studies?

(A): Yes.

(B): No.



# Bayes' Rule with Multiple Evidence

**Example:** Say we know from medicinal studies that  $P(\text{cavity}) = 0.2$ ,  $P(\text{toothache} \mid \text{cavity}) = 0.6$ ,  $P(\text{toothache} \mid \neg \text{cavity}) = 0.1$ ,  $P(\text{catch} \mid \text{cavity}) = 0.9$ , and  $P(\text{catch} \mid \neg \text{cavity}) = 0.2$ . Now, in case we did observe the symptoms toothache and catch (the dentist's probe catches in the aching tooth), what would be the likelihood of having a cavity? What is  $P(\text{cavity} \mid \text{catch}, \text{toothache})$ ?

By Bayes' rule we get:

$$P(\text{cavity} \mid \text{catch}, \text{toothache}) = \frac{P(\text{catch}, \text{toothache} \mid \text{cavity})P(\text{cavity})}{P(\text{catch}, \text{toothache})}$$

Question!

So, is everything fine? Do we just need some more medicinal studies?

(A): Yes.

(B): No.

→ No! We would need  $P(\text{toothache} \wedge \text{catch} \mid \text{Cavity})$ , i.e., causal dependencies for all combinations of symptoms! ( $\gg 2$ , in general)

## Bayes' Rule with Multiple Evidence, ctd.

**Second attempt:** First Normalization (slide 34), then Chain Rule (slide 30) using ordering  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Catch}$ ,  $X_3 = \text{Toothache}$ :

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} \mid \text{catch}, \text{toothache}) = \\
 & \alpha \mathbf{P}(\text{Cavity}, \text{catch}, \text{toothache}) = \\
 & \alpha \mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity}) \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$

## Bayes' Rule with Multiple Evidence, ctd.

**Second attempt:** First Normalization (slide 34), then Chain Rule (slide 30) using ordering  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Catch}$ ,  $X_3 = \text{Toothache}$ :

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} \mid \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{Cavity}, \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity}) \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$

**Close, but no Banana:** Less red (i.e.unknown) probabilities, but still  $\mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity})$ .

## Bayes' Rule with Multiple Evidence, ctd.

**Second attempt:** First Normalization (slide 34), then Chain Rule (slide 30) using ordering  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Catch}$ ,  $X_3 = \text{Toothache}$ :

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} \mid \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{Cavity}, \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity}) \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$

**Close, but no Banana:** Less red (i.e. unknown) probabilities, but still  $\mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity})$ .

**But:** *Are Toothache and Catch independent?*

## Bayes' Rule with Multiple Evidence, ctd.

**Second attempt:** First Normalization (slide 34), then Chain Rule (slide 30) using ordering  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Catch}$ ,  $X_3 = \text{Toothache}$ :

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} \mid \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{Cavity}, \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity}) \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$

**Close, but no Banana:** Less red (i.e. unknown) probabilities, but still  $\mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity})$ .

**But:** *Are Toothache and Catch independent?*

→ No. If a probe catches, we probably have a cavity which probably causes toothache.

## Bayes' Rule with Multiple Evidence, ctd.

**Second attempt:** First Normalization (slide 34), then Chain Rule (slide 30) using ordering  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Catch}$ ,  $X_3 = \text{Toothache}$ :

$$\begin{aligned}
 & \mathbf{P}(\text{Cavity} \mid \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{Cavity}, \text{catch}, \text{toothache}) = \\
 & \propto \mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity}) \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{Cavity})
 \end{aligned}$$

**Close, but no Banana:** Less red (i.e. unknown) probabilities, but still  $\mathbf{P}(\text{toothache} \mid \text{catch}, \text{Cavity})$ .

**But:** *Are Toothache and Catch independent?*

→ No. If a probe catches, we probably have a cavity which probably causes toothache.

**But:** *They are independent given the presence or absence of cavity!*

→ See next slide.

## Conditional Independence, ctd.

**Proposition.** *If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then*  

$$\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}).$$

## Conditional Independence, ctd.

**Proposition.** *If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z})$ .*

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$



## Conditional Independence, ctd.

**Proposition.** If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then

$$\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}).$$

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by product rule is equal to

$$\frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$$

## Conditional Independence, ctd.

**Proposition.** If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then

$$\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}).$$

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by product rule is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by prerequisite is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$ .

## Conditional Independence, ctd.

**Proposition.** *If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then*

$$\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}).$$

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by product rule is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by prerequisite is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$ . Since  $\frac{\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})} = 1$  this proves the claim.

## Conditional Independence, ctd.

**Proposition.** If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z})$ .

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by product rule is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by prerequisite is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$ . Since  $\frac{\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})} = 1$  this proves the claim.

**Example:** Using  $\{\textit{Toothache}\}$  as  $\mathbf{Z}_1$ ,  $\{\textit{Catch}\}$  as  $\mathbf{Z}_2$ , and  $\{\textit{Cavity}\}$  as  $\mathbf{Z}$ :  
 $\mathbf{P}(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) = \mathbf{P}(\textit{Toothache} \mid \textit{Cavity})$ .

→ In the presence of conditional independence, we can drop variables from the right-hand side of conditional probabilities.

→ Third of the four basic techniques in Bayesian networks.

## Conditional Independence, ctd.

**Proposition.** If  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are conditionally independent given  $\mathbf{Z}$ , then  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z})$ .

**Proof.** By definition,  $\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}_2, \mathbf{Z}) = \frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2, \mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})}$  which by product rule is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1, \mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}$  which by prerequisite is equal to  $\frac{\mathbf{P}(\mathbf{Z}_1 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}$ . Since  $\frac{\mathbf{P}(\mathbf{Z}_2 \mid \mathbf{Z}) \mathbf{P}(\mathbf{Z})}{\mathbf{P}(\mathbf{Z}_2, \mathbf{Z})} = 1$  this proves the claim.

**Example:** Using  $\{\textit{Toothache}\}$  as  $\mathbf{Z}_1$ ,  $\{\textit{Catch}\}$  as  $\mathbf{Z}_2$ , and  $\{\textit{Cavity}\}$  as  $\mathbf{Z}$ :  
 $\mathbf{P}(\textit{Toothache} \mid \textit{Catch}, \textit{Cavity}) = \mathbf{P}(\textit{Toothache} \mid \textit{Cavity})$ .

→ In the presence of conditional independence, we can drop variables from the right-hand side of conditional probabilities.

→ Third of the four basic techniques in Bayesian networks. Last missing technique: "Capture variable dependencies in a graph"; illustration see Conclusions, details see **Next Chapter**.

# Agenda

- 1 Introduction
- 2 Propositional Logic
- 3 Quantifying Uncertainty
- 4 Basic Probability Calculus
- 5 Conditional Probabilities
- 6 Basic Probabilistic Reasoning Methods
- 7 Bayes' Rule
- 8 Independence
- 9 Conditional Independence
- 10 Conclusion

## Summary

- Reasoning can be attained by a combination of logic and probability.
- Deduction is about deriving conclusions that follow logically from our knowledge base.
- Uncertainty is unavoidable in many environments, namely whenever agents do not have perfect knowledge.
- Probabilities express the degree of belief of an agent, given its knowledge, into an event.
- Conditional probabilities express the likelihood of an event given observed evidence.
- Assessing a probability means to use statistics to approximate the likelihood of an event.
- Bayes' rule allows us to derive, from probabilities that are easy to assess, probabilities that aren't easy to assess.
- Given multiple evidence, we can exploit conditional independence.


# Summary

- Reasoning can be attained by a combination of logic and probability.
- Deduction is about deriving conclusions that follow logically from our knowledge base.
- Uncertainty is unavoidable in many environments, namely whenever agents do not have perfect knowledge.
- Probabilities express the degree of belief of an agent, given its knowledge, into an event.
- Conditional probabilities express the likelihood of an event given observed evidence.
- Assessing a probability means to use statistics to approximate the likelihood of an event.
- Bayes' rule allows us to derive, from probabilities that are easy to assess, probabilities that aren't easy to assess.
- Given multiple evidence, we can exploit conditional independence.
  - Bayesian networks (up next) do this, in a comprehensive manner (see next slides for some spoilers of where are we headed).



# Exploiting Conditional Independence: Overview

## 1. Graph captures variable dependencies: (Variables $X_1, \dots, X_n$ )



→ Given evidence  $e$ , want to know  $\mathbf{P}(X \mid e)$ . Remaining vars:  $\mathbf{Y}$ .

# Exploiting Conditional Independence: Overview

## 1. Graph captures variable dependencies: (Variables $X_1, \dots, X_n$ )



→ Given evidence  $\mathbf{e}$ , want to know  $\mathbf{P}(X \mid \mathbf{e})$ . Remaining vars:  $\mathbf{Y}$ .

## 2. Normalization+Marginalization:

$$\mathbf{P}(X \mid \mathbf{e}) = \alpha \mathbf{P}(X, \mathbf{e}); \text{ if } \mathbf{Y} \neq \emptyset \text{ then } \mathbf{P}(X \mid \mathbf{e}) = \alpha \sum_{\mathbf{y} \in \mathbf{Y}} \mathbf{P}(X, \mathbf{e}, \mathbf{y})$$

→ A sum over atomic events!

# Exploiting Conditional Independence: Overview

## 1. Graph captures variable dependencies: (Variables $X_1, \dots, X_n$ )



→ Given evidence  $e$ , want to know  $\mathbf{P}(X \mid e)$ . Remaining vars:  $\mathbf{Y}$ .

## 2. Normalization+Marginalization:

$$\mathbf{P}(X \mid e) = \alpha \mathbf{P}(X, e); \text{ if } \mathbf{Y} \neq \emptyset \text{ then } \mathbf{P}(X \mid e) = \alpha \sum_{\mathbf{y} \in \mathbf{Y}} \mathbf{P}(X, e, \mathbf{y})$$

→ A sum over atomic events!

## 3. Chain rule: Order $X_1, \dots, X_n$ consistently with dependency graph.

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_1)$$

# Exploiting Conditional Independence: Overview

## 1. Graph captures variable dependencies: (Variables $X_1, \dots, X_n$ )



→ Given evidence  $e$ , want to know  $\mathbf{P}(X \mid e)$ . Remaining vars:  $\mathbf{Y}$ .

## 2. Normalization+Marginalization:

$$\mathbf{P}(X \mid e) = \alpha \mathbf{P}(X, e); \text{ if } \mathbf{Y} \neq \emptyset \text{ then } \mathbf{P}(X \mid e) = \alpha \sum_{\mathbf{y} \in \mathbf{Y}} \mathbf{P}(X, e, \mathbf{y})$$

→ A sum over atomic events!

## 3. Chain rule: Order $X_1, \dots, X_n$ consistently with dependency graph.

$$\mathbf{P}(X_1, \dots, X_n) = \mathbf{P}(X_n \mid X_{n-1}, \dots, X_1) * \mathbf{P}(X_{n-1} \mid X_{n-2}, \dots, X_1) * \dots * \mathbf{P}(X_1)$$

4. **Exploit conditional independence:** Instead of  $\mathbf{P}(X_i \mid X_{i-1}, \dots, X_1)$ , with previous slide we can use  $\mathbf{P}(X_i \mid \text{Parents}(X_i))$ .

→ Bayesian networks!

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ Given *toothache, catch*, want  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch})$ . Remaining vars:  $\emptyset$ .

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ **Given** *toothache, catch*, want  $\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch}) = \alpha \mathbf{P}(\textit{Cavity}, \textit{toothache}, \textit{catch})$$

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ Given *toothache*, *catch*, want  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch}) = \alpha \mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch})$$

3. **Chain rule:** Order  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Toothache}$ ,  $X_3 = \text{Catch}$ .

$$\begin{aligned} &\mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch}) = \\ &\mathbf{P}(\text{catch} \mid \text{toothache}, \text{Cavity}) \mathbf{P}(\text{toothache} \mid \text{Cavity}) \mathbf{P}(\text{Cavity}) \end{aligned}$$

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ Given *toothache*, *catch*, want  $\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch}) = \alpha \mathbf{P}(\textit{Cavity}, \textit{toothache}, \textit{catch})$$

3. **Chain rule:** Order  $X_1 = \textit{Cavity}$ ,  $X_2 = \textit{Toothache}$ ,  $X_3 = \textit{Catch}$ .

$$\begin{aligned} &\mathbf{P}(\textit{Cavity}, \textit{toothache}, \textit{catch}) = \\ &\mathbf{P}(\textit{catch} \mid \textit{toothache}, \textit{Cavity}) \mathbf{P}(\textit{toothache} \mid \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \end{aligned}$$

4. **Exploit conditional independence:**

Instead of  $\mathbf{P}(\textit{catch} \mid \textit{toothache}, \textit{Cavity})$  use  $\mathbf{P}(\textit{catch} \mid \textit{Cavity})$ .



## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ **Given** *toothache*, *catch*, **want**  $\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch}) = \alpha \mathbf{P}(\textit{Cavity}, \textit{toothache}, \textit{catch})$$

3. **Chain rule:** Order  $X_1 = \textit{Cavity}$ ,  $X_2 = \textit{Toothache}$ ,  $X_3 = \textit{Catch}$ .

$$\begin{aligned} &\mathbf{P}(\textit{Cavity}, \textit{toothache}, \textit{catch}) = \\ &\mathbf{P}(\textit{catch} \mid \textit{toothache}, \textit{Cavity}) \mathbf{P}(\textit{toothache} \mid \textit{Cavity}) \mathbf{P}(\textit{Cavity}) \end{aligned}$$

4. **Exploit conditional independence:**

Instead of  $\mathbf{P}(\textit{catch} \mid \textit{toothache}, \textit{Cavity})$  use  $\mathbf{P}(\textit{catch} \mid \textit{Cavity})$ .

**Thus:**  $\mathbf{P}(\textit{Cavity} \mid \textit{toothache}, \textit{catch}) =$

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ **Given** *toothache*, *catch*, **want**  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch}) = \alpha \mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch})$$

3. **Chain rule:** Order  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Toothache}$ ,  $X_3 = \text{Catch}$ .

$$\begin{aligned} &\mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch}) = \\ &\mathbf{P}(\text{catch} \mid \text{toothache}, \text{Cavity}) \mathbf{P}(\text{toothache} \mid \text{Cavity}) \mathbf{P}(\text{Cavity}) \end{aligned}$$

4. **Exploit conditional independence:**

Instead of  $\mathbf{P}(\text{catch} \mid \text{toothache}, \text{Cavity})$  use  $\mathbf{P}(\text{catch} \mid \text{Cavity})$ .

**Thus:**  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch}) =$   
 $\alpha \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{toothache} \mid \text{Cavity}) \mathbf{P}(\text{Cavity}) =$   
 $\alpha \langle 0.9 * 0.6 * 0.2, 0.2 * 0.1 * 0.8 \rangle = \alpha \langle 0.108, 0.016 \rangle.$

## Exploiting Conditional Independence: Example

1. **Graph captures variable dependencies:** (See previous slide.)

→ **Given** *toothache*, *catch*, **want**  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch})$ . Remaining vars:  $\emptyset$ .

2. **Normalization+Marginalization:**

$$\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch}) = \alpha \mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch})$$

3. **Chain rule:** Order  $X_1 = \text{Cavity}$ ,  $X_2 = \text{Toothache}$ ,  $X_3 = \text{Catch}$ .

$$\begin{aligned} &\mathbf{P}(\text{Cavity}, \text{toothache}, \text{catch}) = \\ &\mathbf{P}(\text{catch} \mid \text{toothache}, \text{Cavity}) \mathbf{P}(\text{toothache} \mid \text{Cavity}) \mathbf{P}(\text{Cavity}) \end{aligned}$$

4. **Exploit conditional independence:**

Instead of  $\mathbf{P}(\text{catch} \mid \text{toothache}, \text{Cavity})$  use  $\mathbf{P}(\text{catch} \mid \text{Cavity})$ .

**Thus:**  $\mathbf{P}(\text{Cavity} \mid \text{toothache}, \text{catch}) =$

$$\alpha \mathbf{P}(\text{catch} \mid \text{Cavity}) \mathbf{P}(\text{toothache} \mid \text{Cavity}) \mathbf{P}(\text{Cavity}) =$$

$$\alpha \langle 0.9 * 0.6 * 0.2, 0.2 * 0.1 * 0.8 \rangle = \alpha \langle 0.108, 0.016 \rangle. \text{ So } \alpha \approx 8.06 \text{ and}$$

$$\mathbf{P}(\text{cavity} \mid \text{toothache} \wedge \text{catch}) \approx 0.87.$$