

Multilevel Analysis of School Examination Data

Benjamin Chun Ho Chan*

November 27, 2018

Abstract

A real dataset called EXAM is downloaded from Centre for Multilevel Modelling at University of Bristol, consists of 4,059 students from 65 schools in Inner London. The objective is to investigate the relationship between the exam score and the student-level and/or school-level variables.

Contents

1	Introduction	2
1.1	Motivation for Multilevel Models	2
1.2	Data Set Description and Notation	2
1.3	Exploratory Data Analysis	3
1.4	Methodology in Bayesian Estimation	4
2	Full Models	5
2.1	Full Model with Random and Fixed Intercept	5
2.1.1	Bayesian Setting	5
2.2	Full Model with Random but no Fixed Intercept	6
2.3	Full Model with Random Intercept and LRT Effect	6
2.4	Full Model without Intercept, with Random LRT Effect	7
3	Submodels	7
3.1	Submodel with Intercepts, without School Effect Predictor	7
3.2	Submodel with Intercepts, without Student VR Band	8
3.3	Submodel with Intercepts, without Student VR, Intake Band	8
3.4	Submodel with Intercepts, without Student Gender and VR	9
3.5	Submodel with Intercepts, without Student VR, School Gender	9
3.6	Submodel with Intercepts, without Student VR, School Intake	10
3.7	Submodel with Random Intercept, without School Effect Predictor	10
4	Recommended Model	11
4.1	Result	11
4.2	Intuitive Interpretation	11
5	Appendix	12
5.1	Running Mean Plots of Recommended Model	12

*For enquiry, please email to 1155049861@link.cuhk.edu.hk.

1 Introduction

The dataset EXAM is downloaded from Centre for Multilevel Modelling at University of Bristol: <http://www.bristol.ac.uk/cmm/learning/mmsoftware/data-rev.html>. It consists of 4,059 students from 65 schools in Inner London. The objective is to investigate the relationship between the exam score and the student-level and/or school-level variables. For implementation in R, please refer to my GitHub repository: <https://github.com/BenjaminChanChunHo/Advanced-Modeling-and-Data-Analysis>.

1.1 Motivation for Multilevel Models

Multilevel models are appropriate for hierarchically arranged data with nested sources of variability. In EXAM, the sources of variability include different student abilities, i.e. subject-specific random errors, and differences between schools, i.e. random cluster effects. It is cross sectional in the sense that students at the lower level cluster at the higher level, i.e. schools. To address the similarity between students in the same school, multilevel models are used.

1.2 Data Set Description and Notation

The attribute name, attribute type and description are given on the website as follows:

Name	Data Type	Description
School ID	Integer	Unique to School
Student ID	Integer	Non-unique across School
Exam Score	Continuous	Standardized Response
London Reading Test Score	Continuous	Standardized Predictor
Student Gender	Binary	0 = Boy; 1 = Girl
School Gender	Trinary	1 = Mixed; 2 = Boys; 3 = Girls
School Intake Score	Continuous	Unique to School
Student Verbal Reasoning Band	Trinary	1 = Bot 25%; 2 = Mid 50%; 3 = Top 25%
Student Intake Band	Trinary	1 = Bot 25%; 2 = Mid 50%; 3 = Top 25%

Table 1: Attribute Information

Let y_{ij} be exam score for school $j = 1, \dots, 65$ and student $i = 1, \dots, n_j$, where $\sum_{j=1}^{65} n_j = 4059$. Full Model is a model using all variables while Submodel is one using some variables.

Variable Name	Range	Notation	Original Variable
School ID	$\{1, 2, \dots, 65\}$	j	Same
Exam Score	-3.66 to 3.67	y	Same
LR Test Score	-2.93 to 3.02	x_1	Same
Student Gender	$\{0, 1\}$	x_2	Same
Student VR Mid	$\{0, 1\}$	x_3	VR Score = 2
Student VR Top	$\{0, 1\}$	x_4	VR Score = 3
Student Band Mid	$\{0, 1\}$	x_5	Intake Band = 2
Student Band Top	$\{0, 1\}$	x_6	Intake Band = 3
School Intake Score	-0.76 to 0.64	w_1	Same
School Boys	$\{0, 1\}$	w_2	School Gender = 2
School Girls	$\{0, 1\}$	w_3	School Gender = 3

Table 2: Notation

1.3 Exploratory Data Analysis

Univariate Plots at Student Level

Note that y and x_1 follow a standard normal distribution approximately:




Figure 1: Histogram of y




Figure 2: Histogram of x_1

Univariate Plots at School Level

Heterogeneity among schools is supported by the histograms:




Figure 3: Histogram of w_1




Figure 4: Histogram of \bar{y}_j

where \bar{y}_j is the sample mean of y under School ID j .

Bivariate Relationship at Student Level

Exam Score y and LR Test Score x_1 are positively correlated:




Figure 5: Scatter Plot of x_1 vs y

The higher the Student VR Band is, the higher the Mean Exam Score is

Student VR Band	Mean Exam Score
1	-0.412
2	-0.062
3	0.349

Table 3: Mean Exam Score under each Student VR Band

The lower the Student Intake Band is, the higher the Mean Exam Score is

Student Intake Band	Mean Exam Score
1	0.736
2	-0.142
3	-0.990

Table 4: Mean Exam Score under each Student Intake Band

Bivariate Relationship at School Level

Sample mean of exam scores for each school \bar{y}_j and School Intake Score w_1 are positively correlated:




Figure 6: Scatter Plot of w_1 vs \bar{y}_j

1.4 Methodology in Bayesian Estimation

Markov Chain Monte Carlo (MCMC) algorithms are used to simulate a large number of observations from the joint posterior distribution. The Bayesian estimates of unknown parameters can be obtained from the corresponding sample means of simulated observations.

The software WinBUGS is used via the R package R2WinBUGS to do the Bayesian inference. For simplicity, the conjugate prior distributions (e.g. normal and gamma distributions) are used and hyperparameters are specified. Two sequences are generated with different initial values. The burn-in iterations required for achieving convergence of the MCMC algorithms are set to be 8,000. In fact, it can be checked by running mean plots of parameters. The number of simulated observations collected after the burn-in iterations is set to be 6,000.

The posterior means and standard error estimates of parameters are reported. Treatment is similar for most models and hence details are omitted after introducing the first model. Please refer to GitHub for code and running mean plots. DIC and parsimony principle are used to compare models.

2 Full Models

2.1 Full Model with Random and Fixed Intercept

Let the full model with random and fixed intercept be

$$y_{ij} = \mathbf{x}_{ij}^T \boldsymbol{\beta} + u_{0j} + \varepsilon_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \cdots + \beta_6 x_{6ij} + \varepsilon_{ij}$$

where $j = 1, \dots, 65$ and $i = 1, \dots, n_j$.

Note that \mathbf{x}_{ij} is 7×1 vector of covariates (including constant 1) and u_{0j} is level 2 random intercepts (school effects). Assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$. Using school level predictors $\mathbf{w}_j^T = (1, w_{1j}, w_{2j}, w_{3j})$,

$$\mu_{0j} = \mathbf{w}_j^T \boldsymbol{\alpha}_0 = \alpha_{00} + \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}.$$

2.1.1 Bayesian Setting

The conjugate prior distributions of σ_0^2 , σ_y^2 , $\boldsymbol{\beta}$ and $\boldsymbol{\alpha}_0$ are

$$\sigma_0^{-2} \stackrel{D}{=} \text{Gamma}[a_0, b_0], \quad \sigma_y^{-2} \stackrel{D}{=} \text{Gamma}[a_y, b_y], \quad \boldsymbol{\beta} \stackrel{D}{=} N(\boldsymbol{\mu}_\beta, \boldsymbol{\Sigma}_\beta) \text{ and } \boldsymbol{\alpha}_0 \stackrel{D}{=} N(\boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0),$$

where a_0 , b_0 , a_y , b_y , and elements in $\boldsymbol{\mu}_\beta$, $\boldsymbol{\mu}_0$, positive definite $\boldsymbol{\Sigma}_\beta$ and $\boldsymbol{\Sigma}_0$ are hyperparameters. Since there is no available prior knowledge, the hyperparameters are specified for convenience:

$$a_0 = 6, \quad b_0 = 10, \quad a_y = 6, \quad b_y = 10, \quad \boldsymbol{\mu}_\beta = \mathbf{0}, \quad \boldsymbol{\Sigma}_\beta = \mathbf{I}, \quad \boldsymbol{\mu}_0 = \mathbf{0} \quad \text{and} \quad \boldsymbol{\Sigma}_0 = \mathbf{I}.$$

There are two sets of initial values of parameters (denoted by *). The first set is:

$$\boldsymbol{\beta}^* = \mathbf{0}, \quad \boldsymbol{\alpha}_0^* = \mathbf{0}, \quad \sigma_0^{2*} = 1, \quad \sigma_y^{2*} = 1.$$

The second set is

$$\begin{aligned} \beta_0^* &= 0, \quad \beta_1^* = \beta_3^* = 1, \quad \beta_2^* = \beta_4^* = \beta_5^* = \beta_6^* = -1, \\ \alpha_{00}^* &= 0, \quad \alpha_{01}^* = \alpha_{03}^* = 1, \quad \alpha_{02}^* = -1, \quad \sigma_0^{2*} = 2, \quad \sigma_y^{2*} = 2. \end{aligned}$$

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	0.42	0.72	-0.80	1.68	Y	
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_3	Student.VR.Mid	0.00	0.25	-0.49	0.48	Y	
β_4	Student.VR.Top	0.08	0.39	-0.72	0.84	Y	
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
α_{00}	u_Int	-0.26	0.67	-1.42	0.77	Y	
α_{01}	School.Intake.Score	0.21	0.41	-0.58	1.03	Y	
α_{02}	School.Boys	0.19	0.22	-0.24	0.61	Y	
α_{03}	School.Girls	0.14	0.17	-0.20	0.48	Y	
σ_0^2	var_u0	0.37	0.06	0.26	0.51	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 5: Full Model with Random and Fixed Intercept; DIC = 9033.81

The fixed intercepts β_0 and α_{00} are not significant in the sense that each of the Bayesian credible intervals contains 0. It is intuitive since all the variables are standardized, i.e. mean 0.

2.2 Full Model with Random but no Fixed Intercept

Let the full model with random but no fixed intercept be

$$y_{ij} = u_{0j} + \beta_1 x_{1ij} + \cdots + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_3	Student.VR.Mid	0.13	0.13	-0.13	0.38	Y	
β_4	Student.VR.Top	0.28	0.21	-0.13	0.74	Y	
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
α_{01}	School.Intake.Score	0.04	0.29	-0.55	0.59	Y	
α_{02}	School.Boys	0.23	0.21	-0.19	0.64	Y	
α_{03}	School.Girls	0.16	0.17	-0.18	0.49	Y	
σ_0^2	var_u0	0.37	0.06	0.26	0.51	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 6: Full Model with Random but no Fixed Intercept; DIC = 9033.82

2.3 Full Model with Random Intercept and LRT Effect

Consider random intercept and random slope associated with the continuous LR Test Score x_1 . Let the full model with random intercept and LRT effect be

$$y_{ij} = u_{0j} + (\beta_1 + u_{1j})x_{1ij} + \beta_2 x_{2ij} + \cdots + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}, \quad \mu_{1j} = \alpha_{11} w_{1j} + \alpha_{12} w_{2j} + \alpha_{13} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$, $u_{1j} \sim N(\mu_{1j}, \sigma_1^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_1	LR.Test.Score	0.38	0.09	0.22	0.57	N	*
β_2	Student.Gender	0.17	0.03	0.10	0.23	N	*
β_3	Student.VR.Mid	0.14	0.13	-0.12	0.40	Y	
β_4	Student.VR.Top	0.21	0.22	-0.17	0.67	Y	
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.34	N	*
β_6	Student.Band.Top	-0.77	0.05	-0.87	-0.66	N	*
α_{01}	School.Intake.Score	0.14	0.30	-0.46	0.71	Y	
α_{02}	School.Boys	0.22	0.21	-0.20	0.64	Y	
α_{03}	School.Girls	0.16	0.17	-0.18	0.49	Y	
α_{11}	School.Intake.Score	0.12	0.21	-0.30	0.53	Y	
α_{12}	School.Boys	-0.01	0.20	-0.40	0.38	Y	
α_{13}	School.Girls	-0.03	0.16	-0.35	0.28	Y	
σ_0^2	var_u0	0.37	0.06	0.26	0.51	N	*
σ_1^2	var_u1	0.32	0.05	0.23	0.44	N	*
σ_y^2	var_y	0.52	0.01	0.50	0.55	N	*

Table 7: Full Model with Random Intercept and LRT Effect; DIC = 8990.33

2.4 Full Model without Intercept, with Random LRT Effect

Consider removing the random intercept and leaving the random LRT effect. Let the full model without intercept, with random LRT effect be

$$y_{ij} = (\beta_1 + u_{1j})x_{1ij} + \beta_2x_{2ij} + \cdots + \beta_6x_{6ij} + \varepsilon_{ij}$$

$$\mu_{1j} = \alpha_{11}w_{1j} + \alpha_{12}w_{2j} + \alpha_{13}w_{3j}$$

and assume that $u_{1j} \sim N(\mu_{1j}, \sigma_1^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_1	LR.Test.Score	0.39	0.10	0.21	0.57	N	*
β_2	Student.Gender	0.17	0.02	0.12	0.22	N	*
β_3	Student.VR.Mid	0.17	0.03	0.12	0.23	N	*
β_4	Student.VR.Top	0.30	0.03	0.23	0.36	N	*
β_5	Student.Band.Mid	-0.37	0.03	-0.43	-0.32	N	*
β_6	Student.Band.Top	-0.71	0.05	-0.81	-0.62	N	*
α_{11}	School.Intake.Score	0.14	0.21	-0.27	0.56	Y	
α_{12}	School.Boys	0.00	0.21	-0.41	0.40	Y	
α_{13}	School.Girls	-0.02	0.16	-0.34	0.30	Y	
σ_1^2	var_u1	0.33	0.06	0.24	0.46	N	*
σ_y^2	var_y	0.59	0.01	0.56	0.61	N	*

Table 8: Full Model without Intercept, with Random LRT Effect; DIC = 9391.79

The DIC rises dramatically, indicating that random intercept may be needed.

3 Submodels

3.1 Submodel with Intercepts, without School Effect Predictor

Let the submodel with random and fixed intercept, without school effect predictor be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1x_{1ij} + \cdots + \beta_6x_{6ij} + \varepsilon_{ij},$$

and assume that $u_{0j} \sim N(0, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	0.15	0.16	-0.18	0.45	Y	
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_3	Student.VR.Mid	0.08	0.19	-0.28	0.44	Y	
β_4	Student.VR.Top	0.24	0.21	-0.16	0.65	Y	
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 9: Submodel with Intercepts, without School Effect Predictor; DIC = 9033.65

3.2 Submodel with Intercepts, without Student VR Band

Parameter estimates of Student VR Band have been shown to be insignificant in many models. Consider removing it as a predictor. Let the submodel with random and fixed intercept, without student VR band be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_5 x_{5ij} + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{00} + \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	0.54	0.55	-0.43	1.69	Y	
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.22	N	*
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
α_{00}	u_Int	-0.36	0.55	-1.51	0.64	Y	
α_{01}	School.Intake.Score	0.29	0.22	-0.14	0.71	Y	
α_{02}	School.Boys	0.19	0.21	-0.23	0.61	Y	
α_{03}	School.Girls	0.15	0.17	-0.18	0.48	Y	
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 10: Submodel with Intercepts, without Student VR Band; DIC = 9033.75

3.3 Submodel with Intercepts, without Student VR, Intake Band

For experimental purpose, consider removing Student Intake Band as a predictor. Let the submodel with random and fixed intercept, without student VR and intake band be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{00} + \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	0.02	0.48	-0.87	0.89	Y	
β_1	LR.Test.Score	0.56	0.01	0.53	0.58	N	*
β_2	Student.Gender	0.17	0.03	0.10	0.24	N	*
α_{00}	u_Int	-0.18	0.49	-1.07	0.73	Y	
α_{01}	School.Intake.Score	0.31	0.22	-0.12	0.74	Y	
α_{02}	School.Boys	0.19	0.22	-0.24	0.62	Y	
α_{03}	School.Girls	0.14	0.17	-0.19	0.48	Y	
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.57	0.01	0.54	0.59	N	*

Table 11: Submodel with Intercepts, without Student VR and Intake Band; DIC = 9248.36

The DIC rises dramatically, indicating that Student Intake Band may be needed.

3.4 Submodel with Intercepts, without Student Gender and VR

For experimental purpose, consider removing Student Gender and VR Band as predictors. Let the submodel with random and fixed intercept, without student gender and VR band be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \beta_5 x_{5ij} + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{00} + \alpha_{01} w_{1j} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	1.07	0.45	-0.16	1.76	Y	
β_1	LR.Test.Score	0.39	0.02	0.36	0.42	N	*
β_5	Student.Band.Mid	-0.41	0.03	-0.48	-0.35	N	*
β_6	Student.Band.Top	-0.76	0.05	-0.86	-0.65	N	*
α_{00}	u_Int	-0.80	0.46	-1.51	0.42	Y	
α_{01}	School.Intake.Score	0.29	0.22	-0.13	0.72	Y	
α_{02}	School.Boys	0.10	0.22	-0.33	0.52	Y	
α_{03}	School.Girls	0.22	0.17	-0.11	0.55	Y	
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.54	0.01	0.52	0.56	N	*

Table 12: Submodel with Intercepts, without Student Gender and VR Band; DIC = 9055.27

3.5 Submodel with Intercepts, without Student VR, School Gender

For experimental purpose, consider removing predictors Student VR Band and School Gender. Let the submodel with random, fixed intercept, without student VR band, school gender be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_5 x_{5ij} + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{00} + \alpha_{01} w_{1j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	-0.09	0.52	-1.20	0.67	Y	
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
α_{00}	u_Int	0.35	0.52	-0.43	1.45	Y	
α_{01}	School.Intake.Score	0.28	0.22	-0.15	0.70	Y	
σ_0^2	var_u0	0.36	0.06	0.26	0.49	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 13: Submodel with Intercepts, without Student VR, School Gender; DIC = 9033.53

3.6 Submodel with Intercepts, without Student VR, School Intake

For experimental purpose, consider removing Student VR Band and School Intake Score. Let the submodel with random, fixed intercept, without student VR, school intake score be

$$y_{ij} = (\beta_0 + u_{0j}) + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_5 x_{5ij} + \beta_6 x_{6ij} + \varepsilon_{ij}$$

$$\mu_{0j} = \alpha_{00} + \alpha_{02} w_{2j} + \alpha_{03} w_{3j}$$

and assume that $u_{0j} \sim N(\mu_{0j}, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_0	y_Int	0.84	0.36	0.05	1.45	N	*
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.86	-0.65	N	*
α_{00}	u_Int	-0.66	0.36	-1.30	0.15	Y	
α_{02}	School.Boys	0.16	0.21	-0.26	0.58	Y	
α_{03}	School.Girls	0.14	0.17	-0.19	0.48	Y	
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 14: Submodel with Intercepts, without Student VR, School Intake; DIC = 9033.55

3.7 Submodel with Random Intercept, without School Effect Predictor

Parameter estimates of school effect predictors have been shown to be insignificant in many models. Consider removing all of them while keeping all student level predictors.

Let the submodel with random intercept and without school effect predictor be

$$y_{ij} = u_{0j} + \beta_1 x_{1ij} + \dots + \beta_6 x_{6ij} + \varepsilon_{ij}$$

and assume that $u_{0j} \sim N(0, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

Result

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_1	LR.Test.Score	0.39	0.02	0.35	0.42	N	*
β_2	Student.Gender	0.16	0.03	0.10	0.23	N	*
β_3	Student.VR.Mid	0.21	0.11	0.00	0.42	N	*
β_4	Student.VR.Top	0.37	0.15	0.08	0.66	N	*
β_5	Student.Band.Mid	-0.41	0.03	-0.47	-0.35	N	*
β_6	Student.Band.Top	-0.75	0.05	-0.85	-0.64	N	*
σ_0^2	var_u0	0.36	0.06	0.26	0.50	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 15: Submodel with Random Intercept, without School Level Predictor; DIC = 9033.34

The effect of Student VR Band is rather weak so consider removing it as a predictor.

4 Recommended Model

Let the model with random intercept, without Student VR band and school effect predictor be

$$y_{ij} = u_{0j} + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_5 x_{5ij} + \beta_6 x_{6ij} + \varepsilon_{ij}$$

and assume that $u_{0j} \sim N(0, \sigma_0^2)$ and $\varepsilon_{ij} \sim N(0, \sigma_y^2)$.

4.1 Result

It is recommended because DIC value is 9034.77, close to the best value obtained before. Moreover, it is a parsimonious model that aims to explain phenomena using fewer parameters.

	Variable	Estimate		95% CI		Significance	
		Mean	SE	2.5%	97.5%	Contain 0	Significant
β_1	LR.Test.Score	0.40	0.02	0.36	0.43	N	*
β_2	Student.Gender	0.18	0.03	0.12	0.25	N	*
β_5	Student.Band.Mid	-0.38	0.03	-0.44	-0.33	N	*
β_6	Student.Band.Top	-0.71	0.05	-0.82	-0.61	N	*
σ_0^2	var_u0	0.40	0.07	0.29	0.54	N	*
σ_y^2	var_y	0.54	0.01	0.51	0.56	N	*

Table 16: Model with Random Intercept, without Student VR Band and School Effect Predictor

4.2 Intuitive Interpretation

It is worth mentioning that the unconditional variance of y is around 1. After fitting the model, $\sigma_0^2 = 0.40$ (0.07) means that multilevel model is useful since there are random school/cluster effects while $\sigma_y^2 = 0.54$ (0.01) means there are student-specific random errors.

In addition, $\beta_1 = 0.40$ (0.02) indicates that the higher the LR Test Score is, the higher the Exam Score is. It is consistent with Figure 5.

Girls perform better than boys do in general as indicated by $\beta_2 = 0.18$ (0.03), the coefficient estimate corresponding to indicator variable of girl. It is consistent with the data that the mean exam score of girls is 0.09 while that of boys is -0.14 .

Note that $\beta_5 = -0.38$ (0.03) and $\beta_6 = -0.71$ (0.05), which are the coefficient estimates corresponding to indicator variables of student intake band 2 and 3 respectively. It means that the lower the Student Intake Band is, the higher the mean exam score is. It is consistent with Table 4.

Student Verbal Reasoning Score Band is not important to Exam Score, in the presence of other variables. Possible explanation is that other student-level variables (say LR Test Score) have incorporated the information. School Intake Score and School Gender are not important to random school effects, in the presence of other variables. In case of further study, other school-level variables may be needed, e.g. school teaching quality measure.

5 Appendix

5.1 Running Mean Plots of Recommended Model




Figure 7: Running Mean Plots of $\beta_1, \beta_2, \beta_5, \beta_6$




Figure 8: Running Mean Plots of σ_0^2




Figure 9: Running Mean Plots of σ_y^2